Evaluation of a Real-time Plasmid Transmission Detection Pipeline

Natalie Scherff[a,b]#, Jörg Rothgänger[b], Thomas Weniger[b], Alexander Mellmann[a], Dag

Harmsen[c]

[a]Institute of Hygiene, University Hospital Münster, Münster, Germany;

[b]Ridom GmbH, Münster, Germany;

[c]Department of Periodontology and Operative Dentistry, University Hospital Münster,

Münster, Germany

Running Head: Plasmid Transmission Detection Pipeline

#Address correspondence to Natalie Scherff, Natalie.Scherff@ukmuenster.de.

https://orcid.org/0000-0002-8414-0259

Abstract

The spread of antimicrobial resistance among bacteria by horizontal plasmid

transmissions poses a major challenge for clinical microbiology. Here, we evaluate a new

real-time plasmid transmission detection pipeline implemented in the SeqSphere[+] (Ridom

GmbH, Münster, Germany) software.

Within the pipeline, a local Mash plasmid database is created and Mash searches with a

distance threshold of 0.001 are used to trigger plasmid transmission early warning alerts

(EWA). Clonal transmissions are detected using cgMLST allelic differences. The

integrated tools pyGenomeViz and MUMmer were used for visual pairwise plasmid

comparisons. We evaluated the pipeline using published hybrid assemblies (Oxford

Nanopore Technology R9.4.1/Illumina) of 81 carbapenem-resistant isolates from

Addenbrooke's Hospital. To emulate prospective usage, samples were imported in

chronological order of sampling date. Different combinations of the user-adjustable

parameters sketch size (1,000 vs 10,000) and size correction were tested and

discrepancies between resulting clusters were analyzed with Quast.

The SeqSphere[+] pipeline agreed with the published data and produced the same clonal

(n=7) and carbapenemase-carrying plasmid (n=9) clusters. EWAs also were in the correct

chronological order. A reduced sketch size of 1,000 led to one missing cluster but could

be compensated using size correction. Among 44 potential transmission events, 12 were

clonal, 18 single plasmid and 14 co-transmissions of two or more plasmids to the same

recipient.

In summary, the developed pipeline presented here is suitable for integration into clinical

microbiology settings with limited bioinformatics knowledge due to its automated

analyses and alert system, which are integrated into the GUI-based SeqSphere^{+} platform.

Importance

Plasmid-mediated spread of antimicrobial resistance (AMR) is a major challenge for

clinical microbiology and monitoring of potential plasmid transmissions is essential to

combat further dissemination. Whole-genome sequencing (WGS) is often used to surveil

nosocomial transmissions but usually limited to the detection of clonal transmissions

(based on chromosomal markers). Recent advances in long-read sequencing technologies

enable full reconstruction of plasmids and the detection of very similar plasmids but so

far easy-to-use tools for this purpose were missing. Here we present an evaluation of an

innovative real-time plasmid transmission detection pipeline. It is integrated into the

GUI-based SeqSphere^{+} software, which already offers cgMLST based pathogen outbreak

detection. It requires very limited bioinformatics knowledge and its automated analyses

and alert system make it well suited for clinical application.

Introduction

Antimicrobial resistance (AMR) is a major challenge in clinical microbiology as it is

limiting the therapeutic options to manage infectious diseases (1). This is largely

mediated by the rapid spread of resistance genes among bacterial populations via

horizontal gene transfer (HGT) facilitated by mobile genetic elements (MGE) such as

plasmids (2). Plasmids are extrachromosomal DNA elements that can serve as a vehicle

to exchange genes between different bacterial strains and even species (3).

Recently, multi-species hospital outbreaks of carbapenem-resistant bacteria have been

reported (4, 5). Whole-genome sequencing (WGS) based surveillance has become an

important tool to monitor nosocomial transmissions in many hospitals (6), but these

activities are usually limited to the detection of clonal transmissions, i.e., based on

chromosomal markers, of single species. Thus, tools are needed that enable the detection

of similar plasmids from WGS data. A variety of software tools has been developed for

plasmid reconstruction and characterization (reviewed in (7)). Among the most popular

are currently CGE PlasmidFinder (8), COPLA (9), PLACNET (10), and MOB-suite (11),

which relies on Mash (12) for comparing k-mers of plasmids. All these tools require

some form of bioinformatic expertise, which is often lacking in routine microbiology

laboratories. Another major limitation is that plasmids are generally difficult to fully

reconstruct from short-reads due to their high number of repetitive regions (7). However,

with the recent advances in long-read sequencing, e.g., Oxford Nanopore Technology

(ONT), accurate long-read data will become more available to clinical microbiology

laboratories (13).

The GUI-based Ridom SeqSphere[+] software (14) is widely used in clinical microbiology

laboratories. One of its main purposes is the detection of clonal transmissions using core-

genome multi-locus sequence typing (cgMLST) based similarity of bacterial isolates.

Here, we evaluated the newly implemented real-time plasmid detection pipeline to enable

users to detect potential plasmid transmissions within their own WGS datasets

complementary to clonal transmissions. We used an already published dataset of hybrid

assemblies of carbapenem-resistant bacteria from a United Kingdom (UK) hospital (15).

Materials and Methods

SeqSphere[+] real-time plasmid transmission detection pipeline

We developed the pipeline as an extra module for the SeqSphere[+] (Ridom GmbH,

Münster, Germany) software. The first steps involve the reconstruction and

characterization of plasmids from assembled data. These steps are based on the MOB-

suite (v3.18) software package (11) and work for both short- and long-read data. Short-

read and non-circular long read data go into the MOB-recon module first to identify and

reconstruct plasmids. The reconstructed plasmids are then forwarded to MOB-typer,

which assigns a primary and secondary cluster ID (Mash distance cluster threshold =

0.006 and 0.025, respectively), rep and relaxase type(s), and mobility prediction to each

plasmid. For long-read data, the FASTA headers are checked for circularity terms

([topology = circular]) and only non-circular contigs a forwarded to MOB-recon. Circular

contigs that are above 500kb in size are considered to be chromosomes, smaller contigs

are considered to be plasmids and directly go into MOB-typer. Optionally, the MOB-

recon module can also be skipped using a [non-recon] term in the FASTA header. AMR

targets are determined using the NCBI AMRFinderPlus (v3.11.26) (16). Integrated

mobile genetic elements (iMGEs) are detected with CGE MobileElementFinder (v1.1.2)

(17). The typing results from MOB-typer, AMRFinderPlus, and MobileElementFinder

are then summarized in a tabular "Chromosome & Plasmid" overview. From here,

plasmid contigs can be exported as FASTA files for downstream analyses.

AMRFinderPlus results are grouped into two columns, one of them only showing priority

AMR genes, which are defined as targets that might confer resistance to carbapenems,

colistin, vancomycin, or methicillin, and genes that contain ESBL or AmpC in their

name. The detected iMGE(s) are also shown in a separate column if they enclose a

priority AMR gene.

<u>Long-read data plasmid transmission analysis module</u>

With the recently introduced version 10 of the SeqSphere$^+$ software with the long-read

data plasmid transmission analysis module, potential plasmid transmissions can be

detected using Mash (v2.1) comparisons (12). The identified plasmids can be used to

build a local Mash plasmid database from the users' own data. Pairwise Mash distances

can then be utilized to detect very similar plasmids. This information can be used in

several ways. First, if new plasmids are imported via the pipeline mode, they are

compared with all existing plasmids in the database and if there are matches below a

certain threshold (default = 0.001), an early warning alert (EWA) is triggered and shown

on the SeqSphere$^+$ start screen. No EWA is generated if the respective samples contain a

cgMLST scheme and have a clonal (i.e., cgMLST allelic distance below the species-

specific threshold) relationship. Second, the complete plasmid database can be used to

perform an all-against-all comparison to create an exportable distance matrix and/or a

single-linkage clustering. Third, the database can be searched for similar plasmids from a

single plasmid of interest. This plasmid search can be triggered from both the

"Chromosome & Plasmid" overview and EWA reports. Within the resulting table,

sample metadata and plasmid typing results for each plasmid are included. Finally,

partially annotated plasmids (origin(s) of replication, priority AMR gene(s), other AMR

gene(s), and iMGE(s)) can also be pairwise aligned and visualized using MUMmer v3.23

(18) and pyGenomeViz (v0.4.4; https://moshi4.github.io/pyGenomeViz/). To create a

better visualization, there is an optional fixstart function to change the start and

orientation of contigs. It uses *dnaA* for chromosomes and for plasmids the first

occurrence of a *rep* gene as the start of the reoriented contig. Figure 1 shows an overview

of the complete pipeline.

SeqSphere^{+} allows for configuration of multiple parameters including sketch size,

sequence length compensation, and Mash distance threshold for matches (default:0.001).

For the sketch size the user has the choice between 1,000 and 10,000. The k-mer size of

21 is fixed. Due to the large variability in plasmid length and as Mash distances favor

contigs of similar size we have implemented a scaling factor for comparing plasmids with

difference in unique content where the match threshold is lowered for each 1% of size

difference. The default is a lowering of 0.0003 per 1% size difference up to a limit of

40% size difference but both the lowering value and the upper limit can be adjusted by

the user.

<u>Evaluation of the pipeline using a published dataset</u>

To test the pipeline functionality and evaluate parameter settings, the pipeline was

applied to a dataset of 85 isolates of phenotypically carbapenem-resistant Gram-negative

bacteria from a single hospital in the UK (Addenbrooke's Hospital, Cambridge) that were

collected over a period of six years (15). We excluded four samples without a detected

carbapenemase gene leading to a total number of 81 samples in our test dataset. These

isolates were sequenced using a combination of Illumina HiSeq and Oxford Nanopore

Technologies (ONT) MinION R9.4.1 technologies. Hybrid assemblies were created using

Flye (v2.8) (19), Unicycler (v0.4.8) (20) and Canu (v2.1.1) (21). These hybrid assemblies

were downloaded in FASTA format from the European Nucleotide Archive (ENA),

where they were deposited under the study accession PRJEB30134. From the information

in the original publication, we did assume that each contig, except the chromosome, was

a single plasmid. However, since we did not have circularity information, we used a

[non-recon] term in the FASTA headers to skip the MOB-recon part as additional

plasmid reconstruction was not done in the original publication. The study was done

retrospectively but to test the EWAs, we emulated a prospective analysis by importing

the files in the chronological order of sampling date.

Plasmid clusters were defined as plasmids with a Mash distance ≤ 0.001. In addition to

clusters, the number of potential transmissions was counted. If a sample carried multiple

plasmids that belonged to different plasmid clusters, it was assessed whether all of these

plasmids were transferred to the same or different receiving samples. If all plasmids were

shared with the same sample, this was counted as one co-transmission event. If plasmids

of one sample were connected to different samples, each connection was counted as a

single plasmid transmission event. If two samples had a clonal relation, the respective

plasmids were not counted as a plasmid but as a clonal transmission event.

To assess clonal relations, we used cgMLST allelic distances. For species with public

cgMLST schemes hosted on the Ridom nomenclature server (www.cgMLST.org), we

used the species-specific default threshold. For species without a public scheme, we

created *ad hoc* schemes, using the NCBI Genomes reference sequence of each species as

seed genome and a threshold of 15 alleles to determine clonal clusters. To check for

additional transmissions of only iMGEs, a user-defined task template was created from

an allele database that contained all iMGE with a carbapenemase gene that were found

using the CGE MobileElementFinder with thresholds of ≥90% identity and ≥95%

alignment.

For parameter evaluation, we tested four combinations: a sketch size of 1,000 and 10,000,

each with and without size compensation applied to the Mash distance. A cluster

threshold of ≤ 0.001 was used for all four approaches. Plasmid clusters with

discrepancies between these different runs were analyzed with Quast (v5.2) (22). Here,

the longest plasmid of each cluster was set as reference. Mismatches, InDels, and larger

gaps were analyzed, and each counted as single evolutionary events. Finally, the pipeline

was applied to non carbapenemase-carrying plasmids as well.

Results

<u>Analysis of clonal clusters</u>

Clonal clusters were determined based on cgMLST allelic distances with species-specific

thresholds (15 for species without public cgMLST schemes). In total, seven clonal

clusters were detected, three for *Klebsiella pneumoniae,* two for *Escherichia coli,* one for

Pseudomonas putida, and one for *Enterobacter hormaechei* (Table 1). Except one *K.*

pneumoniae cluster comprising eleven isolates (kleb_2), all clusters contained two

isolates. The allelic distances were between 0 and 11. In addition, we found two *E.*

roggenkampii isolates with a distance of one allele but since these isolates were from the

same patient, they were denoted as intra-host variability.

<u>Analysis of carbapenemase-carrying plasmids</u>

First, we only analyzed the plasmids carrying a carbapenemase gene. Out of the 81

isolates, eleven isolates contained chromosomally encoded carbapenemase genes. The

other 70 isolates had carbapenemase-carrying plasmids. One isolate comprised two

carbapenemase-carrying plasmids leading to a total number of 71 analyzed plasmids.

Using a sketch size of 10,000, a Mash distance threshold of 0.001, and size

compensation, we found nine plasmid clusters (Table 1). One additional plasmid cluster

consisted of only clonal isolates (kleb_2). Three plasmid clusters (Clust_148, Clust_9,

Clust_72) partly contained clonal isolates. The clusters comprised between two and eight

isolates from one to three different species. The relatedness of the 71 carbapenemase-

carrying plasmids is visualized with a population snapshot using the exported clustering

distance matrix (Figure 2).

To evaluate the parameters, we repeated this analysis with a lower sketch size of 1,000

and tested both sketch sizes with and without size compensation and compared the

resulting clusters. Out of the nine plasmid clusters, four showed discrepancies between

the different methods (Table 2). One plasmid pair (Clust_54) was missed with a sketch

size of 1,000 but only when size compensation was turned off. In Clust_9, one additional

sample was included in the cluster for both sketch sizes when size compensation was

applied. Clust_148 also contained an additional plasmid with size compensation, but only

with a sketch size of 10,000. Finally, Clust_98/Clust_217 was lumped together and

contained nine samples with size compensation but without size compensation it was

divided in two clusters with six and three samples, respectively. We counted the number

of evolutionary events between the disputable plasmid and the reference plasmid of each

cluster. For Clust_54 there were three, for Clust_148 five, and for Clust_9 13

evolutionary events, respectively. For Clust_98/Clust_217 we compared all samples of

Clust_217 with one reference of Clust_98, which resulted in two evolutionary events.

In one cluster (Clust_72), we detected a case, where one plasmid (size = 12,282 bp)

contained a duplicated sequence, which was otherwise completely identical with another

plasmid (size = 6,141 bp) that only contained the single sequence, i.e. their uncorrected

Mash distance was 0. We checked for additional transmissions of iMGEs by retrieving

the sequences of all iMGEs containing a carbapenemase gene and searching for these

sequences in all isolates utilizing a user-defined task template. No additional

transmissions of iMGEs were found using this approach.

<u>Analysis of all plasmids</u>

After analysis of the carbapenemase-carrying plasmids, we also had a look at the overall

plasmid landscape of these samples. In total, 324 plasmids were found in 78 of the 81

samples (three samples did not contain plasmids). Each sample carried 1-10 plasmids

with an average of four plasmids per sample. Of these, 175 plasmids from 58 samples

belonged to a cluster (Mash distance 0.001). On average, three plasmids per isolate

belonged to a cluster. A total of 51 clusters (including clonal and carbapenemase-carrying

plasmid clusters) was detected with 2-11 isolates per cluster (median = 2, mean = 3.4).

In addition to clonal and plasmid clusters, we counted the number of potential

transmission events. In total, 44 single transmission events were counted. Of these,

twelve were clonal transmissions, 18 were single plasmid transmissions, and 14 were co-

transmissions of two or more plasmids to the same sample. Three of the single and five of

the co-plasmid-transmissions were intra-host transmissions. Fifteen isolates transferred

more than one plasmid; for nine of them, all plasmids were transferred to the same and

for six of them to different receiving samples. Furthermore, we detected two cases where

an isolate received plasmids from a clonal and an additional plasmid transmission. One of

these involved two different species, *K. pneumoniae* and *E. coli* (Figure 3). Here, the two

K. pneumoniae samples had identical MLST sequence types (STs) and cgMLST allelic

profiles. However, the sample that was isolated later carried additional plasmids. Two of

these plasmids (size = 1.8 kb & 5.2 kb) clustered with plasmids of an *E. coli* isolate that

also shared a 145 kb NDM1-carrying plasmid with both *K. pneumoniae* samples.

Discussion

In this study, we used a published dataset of plasmids from carbapenem-resistant bacteria

to evaluate a new plasmid detection pipeline implemented in the SeqSphere$^+$ software.

The results are in complete agreement with the findings from the original study, in

particular, the same clonal and plasmid clusters were found. Moreover, the same

resistance genes and plasmid rep-types were detected, despite using different databases.

In the original study, Abricate (https://github.com/tseemann/abricate) was used with the

CARD (23) and PlasmidFinder (8) databases, while our pipeline is based on NCBI

AMRFinderPlus (16) and MOB-suite (11). Different approaches were also used for

clonal cluster detection. While Roberts *et al.* used a combination of single nucleotide

polymorphisms (SNPs) and split k-mer analysis (SKA), the SeqSphere$^+$ pipeline is based

on cgMLST allelic distances.

In both the Roberts *et al.* and our study, Mash with a sketch size of 10,000 and a distance

threshold of 0.001 was used to detect plasmid results, leading not surprisingly to the same

cluster results. However, in contrast to a file-based Mash approach, our automated,

database-based pipeline enables a prospective real-time surveillance of plasmid

transmissions.

We compared the clustering results of Mash databases constructed with a sketch size of

1,000 and 10,000. Both are values used in the literature (15, 24). While the higher sketch

size increases the sensitivity, it also requires more computational memory and power and

enlarges the database size. We found that a sketch size of 1,000 was generally enough to

detect the same clusters at a threshold of 0.001, with one exception. The sensitivity can

be increased without a significant increase in computational power requirements by

applying a size compensation. The size compensation allows to detect potential

transmissions of plasmids that share the same backbone but may have lost or gained

larger unique segments. However, size compensation must have an upper limit to prevent

too many false positives. An interesting finding was a plasmid pair, where one plasmid

consisted of two exact copies of another plasmid. This is likely a result of multimer

formation. To account for such phenomena, we decided to limit the size compensation to

a maximum of 40% plasmid size differences but keep the uncorrected Mash distance

value for plasmids with a larger size difference. Thus, plasmids with large size

differences but very similar sequences, as for example multimers, can still be detected.

When analyzing the whole plasmid landscape of the dataset, we found that the

reconstruction of potential transmission scenarios can be complicated due to complex co-

transmission patterns. Based on genetic clustering, the number of plasmid transmission

events within the analyzed dataset is estimated to be twice as common as clonal

transmission of pathogens but detailed epidemiological information is needed to confirm

these results. In particular, it is still largely unknown how often very similar plasmids

occur without a transmission, i.e. as "core plasmids" of certain lineages or species. As

other plasmid comparison tools, the pipeline only works with well reconstructed, ideally

circular plasmids and thus requires long-read sequences. Moreover, Mash as a k-mer-

based approach does not take synteny into account. To check for rearrangements between

plasmid sequences, pyGenomeViz can be utilized as a visual approach. Alternatively,

new tools such as Pling (25), which is alignment-based, could be used to zoom into

specific plasmid clusters.

The new SeqSphere[+] plasmid transmission detection pipeline allows the detection of

plasmid clusters in both retrospective (single linkage clustering of plasmid databases) and

prospective (Early Warning Alerts) studies. The automated, database-based approach

enables real-time surveillance and allows for rapid actions.

References

1. Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Aguilar GR, Gray A, Han C, Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G, Hamadani BHK, Kumaran EAP, McManigal B, Agarwal R, Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalheiro C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Chiurchiù S, Chowdhury F, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe S, Day NPJ, Luca MD, Dokova K, Dramowski A, Dunachie SJ, Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garrett D, Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Khorana M, Kissoon N, Kobeissi E, Kostyanev T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Limmathurotsakul D, Loftus MJ, Lunn M, Ma J, Mturi N, Munera-Huertas T, Musicha P, Mussi-Pinhata MM, Nakamura T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Peleg AY, Perrone C, Plakkal N, Ponce-de-Leon A, Raad M, Ramdin T, Riddell A, Roberts T, Robotham JV, Roca A, Rudd KE, Russell N, Schnall J, Scott JAG, Shivamallappa M, Sifuentes-Osornio J, Steenkeste N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Turner C, Turner P, Doorn HR van, Velaphi S, Vongpradith A, Vu H, Walsh T, Waner S, Wangrangsimakul T, Wozniak T, Zheng P, Sartorius B, Lopez AD, Stergachis A,

Moore C, Dolecek C, Naghavi M. 2022. Global burden of bacterial antimicrobial

resistance in 2019: a systematic analysis. The Lancet 399:629–655.

2. Sheppard AE, Stoesser N, Wilson DJ, Sebra R, Kasarskis A, Anson LW, Giess A,

Pankhurst LJ, Vaughan A, Grim CJ, Cox HL, Yeh AJ, Modernising Medical

Microbiology (MMM) Informatics Group, Sifri CD, Walker AS, Peto TE, Crook

DW, Mathers AJ. 2016. Nested Russian doll-like genetic mobility drives rapid

dissemination of the carbapenem resistance gene *bla*KPC. Antimicrob Agents

Chemother 60:3767–3778.

3. Orlek A, Stoesser N, Anjum MF, Doumith M, Ellington MJ, Peto T, Crook D,

Woodford N, Sarah Walker A, Phan H, Sheppard AE. 2017. Plasmid classification in

an era of whole-genome sequencing: Application in studies of antibiotic resistance

epidemiology. Frontiers in Microbiology 8:1–10.

4. Weber RE, Pietsch M, Frühauf A, Pfeifer Y, Martin M, Luft D, Gatermann S,

Pfennigwerth N, Kaase M, Werner G, Fuchs S. 2019. IS26-mediated transfer of

*bla*NDM–1 as the main route of resistance transmission during a polyclonal,

multispecies outbreak in a German hospital. Frontiers in Microbiology 10:1–14.

5. Marimuthu K, Venkatachalam I, Koh V, Harbarth S, Perencevich E, Cherng BPZ,

Fong RKC, Pada SK, Ooi ST, Smitasin N, Thoon KC, Tambyah PA, Hsu LY, Koh

TH, De PP, Tan TY, Chan D, Deepak RN, Tee NWS, Kwa A, Cai Y, Teo Y-Y,

Thevasagayam NM, Prakki SRS, Xu W, Khong WX, Henderson D, Stoesser N, Eyre

DW, Crook D, Ang M, Lin RTP, Chow A, Cook AR, Teo J, Ng OT, Carbapenemase-

Producing Enterobacteriaceae in Singapore (CaPES) Study Group, Marimuthu K,

Venkatachalam I, Cherng BPZ, Fong RKC, Pada SK, Ooi ST, Smitasin N, Thoon

KC, Hsu LY, Koh TH, De PP, Tan TY, Chan D, Deepak RN, Tee NWS, Ang M, Lin

RTP, Teo J, Ng OT. 2022. Whole genome sequencing reveals hidden transmission of

carbapenemase-producing Enterobacterales. Nat Commun 13:3052.

6. Mellmann A, Bletz S, Böking T, Kipp F, Becker K, Schultes A, Prior K, Harmsen D.

2016. Real-time genome sequencing of resistant bacteria provides precision infection

control in an institutional setting. Journal of Clinical Microbiology 54:2874–2881.

7. Paganini JA, Plantinga NL, Arredondo-Alonso S, Willems RJL, Schürch AC. 2021.

Recovering *Escherichia coli* plasmids in the absence of long-read sequencing data.

Microorganisms 9:1613.

8. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L,

Moller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids

using PlasmidFinder and plasmid multilocus sequence typing. Antimicrobial agents

and chemotherapy 58:3895–3903.

9. Redondo-Salvo S, Bartomeus-Peñalver R, Vielva L, Tagg KA, Webb HE, Fernández-

López R, de la Cruz F. 2021. COPLA, a taxonomic classifier of plasmids. BMC

Bioinformatics 22:390.

10. Vielva L, de Toro M, Lanza VF, de la Cruz F. 2017. PLACNETw: a web-based tool

for plasmid reconstruction from bacterial genomes. Bioinformatics 33:3796–3798.

11. Robertson J, Nash JHE. 2018. MOB-suite: software tools for clustering,

reconstruction and typing of plasmids from draft assemblies. Microbial Genomics 4.

12. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy

AM. 2016. Mash: Fast genome and metagenome distance estimation using MinHash.

Genome Biology 17:1–14.

13. Sereika M, Kirkegaard RH, Karst SM, Michaelsen TY, Sørensen EA, Wollenberg

RD, Albertsen M. 2022. Oxford Nanopore R10.4 long-read sequencing enables the

generation of near-finished bacterial genomes from pure cultures and metagenomes

without short-read or reference polishing. Nat Methods 19:823–826.

14. Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, Mellmann

A, Goesmann A, von Haeseler A, Stoye J, Harmsen D. 2013. Updating benchtop

sequencing performance comparison. Nature biotechnology 31:294–296.

15. Roberts LW, Enoch DA, Khokhar F, Blackwell GA, Wilson H, Warne B, Gouliouris

T, Iqbal Z, Török ME. 2023. Long-read sequencing reveals genomic diversity and

associated plasmid movement of carbapenemase-producing bacteria in a UK hospital

over 6 years. Microbial Genomics 9:001048.

16. Feldgarden M, Brover V, Gonzalez-Escalona N, Frye JG, Haendiges J, Haft DH,

Hoffmann M, Pettengill JB, Prasad AB, Tillman GE, Tyson GH, Klimke W. 2021.

AMRFinderPlus and the Reference Gene Catalog facilitate examination of the

genomic links among antimicrobial resistance, stress response, and virulence.

Scientific Reports 11:1–9.

17. Johansson MHK, Bortolaia V, Tansirichaiya S, Aarestrup FM, Roberts AP, Petersen

TN. 2021. Detection of mobile genetic elements associated with antibiotic resistance

in *Salmonella enterica* using a newly developed web tool: MobileElementFinder. J

Antimicrob Chemother 76:101–109.

18. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg

SL. 2004. Versatile and open software for comparing large genomes. Genome Biol

5:R12.

19. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone

reads using repeat graphs. Nat Biotechnol 37:540–546.

20. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. PLoS Comput Biol

13:e1005595.

21. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu:

scalable and accurate long-read assembly via adaptive k-mer weighting and repeat

separation. Genome Res 27:722–736.

22. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile

genome assembly evaluation with QUAST-LG. Bioinformatics 34:i142–i150.

23. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave

BM, Pereira S, Sharma AN, Doshi S, Courtot MM, Lo R, Williams LE, Frye JG,

Elsayegh T, Sardar D, Westman EL, Pawlowski AC, Johnson TA, Brinkman FSLL,

Wright GD, McArthur AG. 2017. CARD 2017: Expansion and model-centric

curation of the comprehensive antibiotic resistance database. Nucleic Acids Research

45:D566–D573.

24. Robertson J, Bessonov K, Schonfeld J, Nash JHE. 2020. Universal whole-sequence-

based plasmid typing and its utility to prediction of host range and epidemiological

surveillance. Microbial Genomics 6:1–12.

25. Frolova D, Lima L, Roberts L, Bohnenkämper L, Wittler R, Stoye J, Iqbal Z. 2024.

Applying rearrangement distances to enable plasmid epidemiology with pling.

bioRxiv https://doi.org/10.1101/2024.06.12.598623.

Acknowledgments

We thank Leah Roberts (CIIC, Queensland University of Technology) for providing help with sequence and epidemiological data. Further, we thank James Robertson and Kyrylo Bessonov (both National Microbiology Laboratory, Public Health Agency of Canada) for critical reading and improvement of the manuscript.

Disclaimers

NS, JR, and TW are (part-time) employees of Ridom GmbH. JR, TW, and DH are shareholders of Ridom GmbH. AM declares no conflict of interest.

428 **Tables**

429 **Table 1:** Overview of detected clonal and carbapenemase-carrying plasmid transmission
430 clusters. Cluster names correspond to the original publication.
431

Cluster	No. of isolates	Species	Carbapenemase gene	Rep types
Clonal clusters				
pseu_1	2	*Pseudomonas putida*	IMP-70	
ecoli_1	2	*Escherichia coli*	OXA-232	
ecoli_2	2	*Escherichia coli*	NDM-5	
ehor_1	2	*Enterobacter hormaechei*	IMP-70	
kleb_1	2	*Klebsiella pneumoniae*	NDM-1	
kleb_2	11	*Klebsiella pneumoniae*	NDM-1	
kleb_3	2	*Klebsiella pneumoniae*	OXA-232	
Intra-host variability				
erog_1	2	*Enterobacter roggenkampii*	OXA-181	
Plasmid clusters				
Clust_33	2	*Escherichia coli, Klebsiella pneumoniae*	NDM-6	IncFIB
Clust_133	2	*Acinetobacter baumannii*	OXA-72	rep_1127
Clust_17	4	*Escherichia coli, Klebsiella aerogenes, Klebsiella pneumonia*	OXA-48	IncI
Clust_148*	4	*Escherichia coli, Klebsiella pneumoniae*	NDM-1	IncC
Clust_54	2	*Escherichia coli*	NDM-5	IncFIB/IncFIA
Clust_9*	3	*Enterobacter hormaechei*	IMP-70	IncC
Clust_72*	8	*Escherichia coli, Klebsiella pneumoniae*	OXA-232	Col
Clust_98	6	*Escherichia coli, Enterobacter hormaechei, Klebsiella pneumonia*	OXA-181	IncX3
Clust_217	4	*Enterobacter roggenkampii, Klebsiella grimontii, Raoultella ornithinolytica*	OXA-181	IncX3

* includes clonal isolates

Plasmid clusters			Mash distance to reference				Quast results			
Cluster	Plasmid size (kb)	Size difference to reference %	1k	1k + sc	10k	10k + sc	Mismatches	Indels	Large gaps	Sum events
Clust_9	127,2	0%								
	127,2	0%	0,0000	0,0000	0,0000	0,0000	0	0	0	0
	126,2	1%	0,0002	0,0000	0,0000	0,0000	0	1	2	3
	96,3	24%	0,0068	0,0000	0,0074	0,0002	1	10	2	13
Clust_148	145,8	0%								
	145,8	0%	0,0000	0,0000	0,0000	0,0000	0	0	0	0
	145,8	0%	0,0000	0,0000	0,0000	0,0000	0	0	0	0
	140,1	4%	0,0022	0,00103	0,0019	0,0007	4	0	1	5
Clust_54	153,2	0%								
	148,4	3%	0,00101	0,0001	0,00099	0,00004	1	1	1	3
Clust_98	51,5	0%	0,0000	0,0000	0,0000	0,0000				
	51,5	0%	0,0000	0,0000	0,0000	0,0000				
	51,5	0%	0,0000	0,0000	0,0000	0,0000				
	51,5	0%	0,0000	0,0000	0,0000	0,0000				
	51,5	0%	0,0000	0,0000	0,0000	0,0000				
	51,5	**0%**								
Clust_217	47,1	9%	0,0021	0,0000	0,0022	0	1	0	1	2
	47,1	9%	0,0021	0,0000	0,0022	0	1	0	1	2
	48,5	6%	0,0022	0,0004	0,0022	0,0005	1	0	1	2

441   **Figures**

443   **Figure 1:** Overview of plasmid transmission detection pipeline.

Figure 2: Plasmid population snapshot based on Mash distances between 71 carbapenemase-carrying plasmids. Each circle represents one plasmid, plasmids with a

Mash distance of 0 are lumped together in single circles with a larger size. Connection

lines are drawn for plasmids with a Mash distance <0.06. Fill color represents the species

of the respective isolates and colored, dotted circles mark clonal clusters among these

isolates. Plasmid clusters (Mash distance threshold <0.001) are illustrated by grey shaded

areas around the circles. Clonal (in color) and plasmid (in grey) cluster names are stated

next to the clusters.

Figure 3: Visualization of a combination of clonal and horizontal plasmid transmissions.

ST = MLST sequence type.