Developmental, regenerative, and behavioral dynamics in acoel reproduction

Vikram Chandra[#*], Samantha E. Tseng[#†], Allison P. Kann, D. Marcela Bolaños, Mansi

Srivastava*

Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology, Harvard

University, 52 Oxford Street, Cambridge, MA 02138

[†] Current address: The Rockefeller University, 1230 York Avenue, New York, NY 10065

# These authors contributed equally.

* Corresponding authors. Emails: vchandra1@fas.harvard.edu, mansi@oeb.harvard.edu

Abstract

Acoel worms belong to an enigmatic and understudied animal lineage in the phylum

Xenacoelomorpha. Sparse taxonomic and histological work suggests that these worms exhibit a

diversity of reproductive anatomies and likely a corresponding diversity in reproductive behavior.

Here, we study the reproductive life history of the three-banded panther worm *Hofstenia*

miamia, an acoel that is emerging as a lab-tractable model system. Using confocal microscopy

and histology, we describe *H. miamia*'s reproductive organs, identifying structures previously

unknown in acoels. Following a cohort of worms from zygote to adulthood, we quantify the

developmental dynamics of their reproductive organs, and find that these organs emerge in a

stereotyped sequence as a function of increasing body size. Studying the dynamics of organ

growth and de-growth during regeneration and in starvation, we show that reproductive organs

follow similar growth rules in these contexts, suggesting that they are regulated by a size-

associated program in all growth contexts. Finally, we study egg-laying behavior, finding that *H.*

miamia lay their eggs through their mouths after loading them into their pharynges. Worms lay

eggs for multiple months after a single mating, suggesting long-term sperm storage despite

lacking a storage organ; we also find that worms can lay viable eggs without mating, indicating a

capacity for self-fertilization. Further, we show that worms assess their environment to make

decisions about when and where to lay their eggs, and sometimes lay eggs in communal

clutches. Together, our work establishes foundational knowledge to enable the experimental

study of reproductive anatomy, physiology, and behavior in acoels.

Introduction

Reproduction is critical for evolutionary success, and animals display great diversity in their

reproductive life histories. The structures and strategies employed for reproduction have been

relatively well-studied in some animal lineages, such as arthropods and vertebrates (Davies et

al., 2013). However, a full understanding of reproductive biology is lacking in many animal

phyla. For example, Phylum Xenacoelomorpha, which includes acoels, nemertodermatids, and

xenoturbellids, is an early branching bilaterian lineage of primarily marine worms whose life

histories are largely unknown (Cannon et al., 2016; Kapli and Telford, 2020; Philippe et al.,

2011) (Fig. 1A).

Roughly 400 species of acoel worms have been described (Achatz et al., 2013;

Jondelius et al., 2011; Tyler, S., Schilling, S., Hooge, M., & Bush, L. F., 2012; Wallberg, 2012).

Analyses of histological sections of specimens collected in the field suggest that, with few

exceptions (Crezée, 1975; Mamkaev, 1965; Raikova et al., 1995), acoels are simultaneous

hermaphrodites. Unlike many marine animals that reproduce sexually through spawning and

external fertilization, e.g., cnidarians, poriferans, echinoderms, etc. (Giribet and Edgecombe,

2020), acoels have been reported to reproduce sexually through internal fertilization (Achatz et

al., 2013). They exhibit a striking diversity in their reproductive anatomy (Table S1). This

anatomical variation likely accompanies variation in reproductive physiology and behavior.

While a few species have been cultured in the lab (Bailly et al., 2014; De Mulder et al., 2009;

Shannon and Achatz, 2007; Zauchner et al., 2015), little is known about how acoels develop

their reproductive organs, mate, or lay eggs.

To improve our understanding of acoel life histories, we studied reproduction in a lab

tractable species, the three-banded panther worm *Hofstenia miamia* (Fig. 1B,C)*. H. miamia* is a

new research organism with features that allow us to study it in controlled conditions: its life

cycle can be closed in the lab (worms develop from embryo to gravid adult in 2-3 months), and

there is a growing array of experimental tools to observe and manipulate the worms (Gehrke et

al., 2019; Hulett et al., 2023; Kimura et al., 2022, 2021; Ricci and Srivastava, 2021; Srivastava,

2022). Histology-based work has described the coarse anatomy of reproductive organs in

animals in the genus *Hofstenia* (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge

et al., 2007; Hookabe et al., 2024; Steinböck, 1966). However, the fine structure of these

organs, how they grow and develop, and how they function in the course of reproductive

behavior remains unknown. To address these questions about the reproductive life history of *H.*

miamia, we use a combination of light and confocal microscopy, fluorescence *in situ*

hybridization, immunofluorescence, and histology, as well as observations of reproductive

development and behavior in controlled conditions. We reveal new facets of acoel reproductive

biology, show that active processes of growth and destruction ensure coordinated organ

development and regeneration, and establish a foundation for the mechanistic study of

reproduction in acoels.

Results

An overview of *Hofstenia miamia*'s reproductive anatomy

The genus *Hofstenia* (Bock, 1923) lies within the family Hofsteniidae. The family currently

contains four genera (Ahyong et al., 2024; Hookabe et al., 2024) and is an early-branching

lineage within acoels (Abalde and Jondelius, 2024; Jondelius et al., 2011). Three species are

currently recognized within the genus *Hofstenia*: *H. atroviridis*, *H. miamia*, and *H. arabiensis*

(Ahyong et al., 2024; Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et al.,

2007). Like other acoels, *Hofstenia* are considered to be simultaneous hermaphrodites (Beltagi

and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et al., 2007; Steinböck, 1966).. *Hofstenia*

miamia is likely the most common and widespread species within the genus, with a geographic

distribution in the Caribbean and the North Atlantic, including the Bahamas and the Florida

coast (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et al., 2007; Steinböck,

1966). All three *Hofstenia* species are described as having an anterior male reproductive

system including a seminal vesicle, prostatic vesicle, a penis with sclerotized needles, 'diffuse'

testes, and a ventral male gonopore near the mouth (Beltagi and Mandura, 1991; Bock, 1923;

Corrêa, 1960; Hooge et al., 2007; Steinböck, 1966). The female reproductive system, as

described, includes two ovaries spanning the posterior two-thirds of the worm. However, these

descriptions were all based on worms roughly 5-8mm in length (Beltagi and Mandura, 1991;

Bock, 1923; Corrêa, 1960; Hooge et al., 2007; Steinböck, 1966). In our lab cultures, fully grown

adult *H. miamia* can be over 1.4cm in length. We noticed that these large adults were more

likely to reproduce than smaller worms, and we reasoned that their anatomy could also be

different. We therefore began by re-examining the coarse anatomy of *Hofstenia miamia*.

We anesthetized adult worms, mounted them ventral side up on a slide, and imaged

them through a dissecting microscope and found, as expected, that *H. miamia*'s male

copulatory structures are located in the anterior of the animal, just posterior to the mouth (Fig.

1C,D; Fig. S1A). These male structures are located within a translucent cylindrical region close

to the ventral surface of the worm and immediately ventral to the pharynx. We observed two

opaque, white regions within the male copulatory apparatus, located towards its posterior (Fig.

1C; Fig. S1A). The larger, posterior oval structure corresponds to the seminal vesicle, while the

smaller, teardrop-shaped anterior structure corresponds to the prostatic vesicle (Fig. 1C,D; Fig.

S1A) (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et al., 2007; Steinböck,

1966). Previous work reported that *H. miamia* has a penis (Fig. S1A): a sperm-delivery organ

containing rigid needles known as 'stylets' (Hooge et al., 2007; Steinböck, 1966). Although

stylets are difficult to see in this imaging preparation, we identified a small aperture immediately

posterior and ventral to the mouth (Fig. S1A), and we observed that the worm extended its

penis through it, showing that this aperture is the male gonopore and confirming the location of

the penis in the anterior of the worm.

As previously described (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge

et al., 2007; Steinböck, 1966), we found that the female reproductive system occupies the

posterior two-thirds of the worm (Fig. 1C,D; Fig. S1B). This system consists of numerous eggs

grouped into three distinct clusters: one lateral group on each side of the body extends along

the anterior-posterior axis (Fig. S1B) and a medially-located cluster present just posterior to the

pharynx (Fig. S1C). The lateral clusters contain oocytes of various sizes, likely corresponding to

different stages of maturation. These oocyte-laden regions in the parenchyma have been

identified as the ovaries in historical reports (Beltagi and Mandura, 1991; Bock, 1923; Corrêa,

1960; Hooge et al., 2007; Steinböck, 1966). Within the ovaries, there is no obvious organization

of oocytes (by size or shape) along any of the worm's body axes. All oocytes within the ovaries

have a germinal vesicle, a nucleus arrested in prophase I of meiosis that is a common feature of

oocytes of nearly all animals (Grossman et al., 2017; Herlands and Maul, 1994; Liu et al., 2006;

Munro et al., 2023). This nucleus breaks down during early oogenesis or after fertilization

(Falleni and Gremigni, 1990; Sagata, 1996), suggesting that *Hofstenia*'s lateral oocytes are

likely unfertilized.

Posterior to the pharynx, there is often a cluster of uniformly sized, spherical eggs that

do not have a germinal vesicle (Fig. 1C,D; Fig. S1C). The number of eggs at this position is

highly variable, ranging from zero to over a dozen eggs. In *H. atroviridis*, Bock (Bock, 1923)

speculated that these eggs are mature, fertilized, and ready to be laid: i.e., zygotes. To test

whether these eggs were indeed zygotes, we dissected out eggs from this midline cluster, as

well as from the lateral ovaries. We found that 46/56 eggs (removed from the midline clusters of

9 worms) developed and hatched into juvenile worms, in line with previously established

developmental timing (Fig. S1D) (Kimura et al., 2021). Meanwhile, all eggs dissected from the

lateral ovaries disintegrated within 48 hours. The eggs in this medial cluster are therefore

mature, likely fertilized, and ready for laying, while eggs in the ovaries are immature or

unfertilized. Overall, these observations confirm and extend previous descriptions of *H. miamia*,

and provide a high-level overview of its reproductive anatomy. Next, taking advantage of *H.*

miamia's tractability, we sought to obtain a finer understanding of its reproductive structures and

their development.

Dynamics of reproductive organ development

How reproductive structures develop as animals reach sexual maturity is not well understood in

the Xenacoelomorpha. As in the acoel *Aphanostoma pulchra* (previously *Isodiametra pulchra*)

(Chiodin et al., 2013), hatchling *H. miamia* have no visible reproductive structures, and previous

observations suggested that these develop over a month after hatching as the worms grow

toward adulthood (Kimura et al., 2021; Srivastava, 2022). To determine the timeline and

dynamics of reproductive development, we reared 42 zygotes in isolation and monitored their

growth over time (Fig. 1E,F), precisely controlling their rearing environment such that each

animal had access to controlled amounts of seawater and food (see Methods). To assess

whether nutrition affects reproductive development, we split the cohort of hatchling worms into

two groups that were fed differently. For each group, we gradually increased the amount of food

they were given as they grew larger, to ensure that worms always had *ad libitum* food access

while avoiding unnecessary overfeeding. These food increases occurred periodically, but we

delayed them by two weeks for one feeding group (Fig. S2A,B). We reasoned that this

treatment would delay growth in one group, allowing us to decouple biological age from organ

and body growth. Twice weekly, we captured images of the ventral surface of each worm and

quantified the size of each visible reproductive structure.

We found that worms maintained their aspect ratio from hatchling to adult, growing

proportionally over time (Fig. 1E-H; Fig. S2C). At every stage of development, the size of each

reproductive organ scaled with body size once that organ appeared (Fig. 1I-K). Worms with

delayed increases in feeding displayed corresponding delays in their growth, both in their body

size (Fig. S2D) and in the timing of appearance of their ovary and seminal vesicle (Fig. 1L). The

penis and seminal vesicle developed at a smaller size in these worms compared to worms

without delayed feeding increases (Fig. 1M), suggesting that resource-limited worms may invest

disproportionately in male development. Multiple regression found that organ size was

explained by both age and body size, but that body size was a better predictor of the size of

each reproductive organ (Table S2). This suggests that the differentiation and growth of

reproductive structures is coupled to the worm's body size, rather than to its age.

We found that reproductive development generally occurs in a stepwise fashion, where

organs appear in a consistent sequence at serially increasing body sizes. Newly hatched

animals possess all major somatic organs (such as a brain, muscle system, pharynx, gut, etc.

(Kimura et al., 2022, 2021)), but lack all reproductive structures (Fig. 1E,F). 4-6 weeks after egg

laying, when the worm is about 3 ± 0.6 mm (mean ± SD) in length, the anterior-ventral tissue

housing the future male copulatory apparatus becomes visibly less dense, and a groove

becomes apparent in the ventral surface. Soon after, the male gonopore becomes detectable

posterior to the mouth. Sperm begin to accumulate in the seminal vesicle around 35 days post

laying, once a worm is about 4.4 ± 0.9 mm (mean ± SD) long. Shortly after, when a worm is 5.6

± 0.9 mm (mean ± SD), the first oocytes become visible in nascent ovaries. These ovaries grow

with the worm over time, with existing oocytes maturing and new ones continuously added.

Around two months after being laid, typically after a worm is 8.7 ± 1.2 mm in length, eggs

appear near the base of the pharynx (Fig. 1E,F,L,M). These eggs are laid within hours or days,

and they develop and hatch into juvenile worms. Given that these worms were isolated since

birth, this suggests that *H. miamia* can reproduce through self-fertilization.

To visualize the extent to which reproductive structures develop in sequence, we ranked

each structure in each worm by its date of appearance (Fig. 1N). All worms, without exception,

first developed a penis. A few worms simultaneously developed a seminal vesicle; in most other

worms, this structure appeared second, with a few developing their seminal vesicle concurrent

with nascent ovaries. Ovaries always appeared before the first fertilized eggs were seen in the

central region. Next, we asked whether the left and right ovaries of each worm developed in

synchrony. We found that on the first day that an ovary was visible, 34/34 worms had both

ovaries visible. Partial regression analysis revealed that the sizes of the two ovaries within a

worm are strongly correlated, even when correcting for body size (Fig. S2E). Consistent with

findings from many other bilaterally-symmetric organs (Allard and Tabin, 2009; Boulan and

Léopold, 2021; Harris et al., 2021; Vallejo et al., 2015; Wolpert, 2010), this suggests that an

active process may synchronize ovarian growth within individual worms.

Figure 1: Reproductive organs develop sequentially following scaling rules. (**A**) Xenacoelomorpha

is an early-branching bilaterian lineage of aquatic worms. Phylogeny modified from (Srivastava, 2022).

Animal icons are from phylopic.org, and are in the public domain; the dashed line reflects uncertainty in

the consensus phylogeny (Cannon et al., 2016; Kapli and Telford, 2020; Philippe et al., 2011). (**B**) Dorsal

view of an adult *Hofstenia miamia*. (**C**) Ventral view of an adult *Hofstenia miamia*; most reproductive

structures are visible in this view. (**D**) Schematized view of the ventral surface of a worm with known

reproductive structures illustrated. (**E**) Timecourse of a representative worm through development, from

hatchling to reproductively mature adult. (**F**) Schematic of timecourse shown in (**E**) with key reproductive

developments illustrated. The first appearance of each organ is highlighted in red. (**G**) The length of

worms increases over time (R^2 = 0.91, p < 0.0001), and (**H**) worms grow proportionally: their length scales

with their width (R^2 = 0.85, p < 0.0001). Error band shows 95% confidence interval. (**I-K**) The length of

each reproductive organ scales with increases in body size (penis: R^2 = 0.70, p < 0.0001; seminal vesicle:

R^2 = 0.63, p < 0.0001; ovaries: R^2 = 0.84, p < 0.0001). Error band shows 95% confidence interval, with

zero values excluded from these regressions. (**L**) Worms with delayed feeding increases had significant

delays in the appearance of their seminal vesicle and ovaries, but not the penis (Welch's t test for date of

appearance for penis: p = 0.08, seminal vesicle: p = 0.04, ovary: p < 0.0001; n ≥ 15). (**M**) Worms with

delayed feeding increases had a smaller body length when a penis and seminal vesicle appeared, but not

when ovaries appeared (Welch's t test for length on date of appearance for penis: p = 0.005, seminal

vesicle: p = 0.03, ovary: p = 0.74; n ≥ 15). Asterisks indicate statistical significance. (**N**) Ranking the order

in which reproductive organs appear (y-axis) in developing worms reveals a stepwise pattern of

reproductive differentiation. The x-axis shows individual worms. dpl = days post laying. Scale bars: 1 mm.

De-growth and regeneration of reproductive organs

Next, we asked whether these patterns of growth persisted in different contexts. *H. miamia*, like

other studied acoels, is an excellent regenerator and can regenerate all tissues from a wide

range of initial tissue configurations (Srivastava, 2022; Srivastava et al., 2014; Steinböck, 1966).

We amputated and followed adult worms in three ways: by isolating tail tips (which contain no

reproductive structures) and studying how they regenerated the majority of their organs, by

cutting worms sagittally and studying how they regenerated their missing half, and by isolating

heads from tails and studying how these head fragments regenerated their tails and the

posterior region of their ovaries (Fig. 2A). Head and tail fragments both gradually increased in

size, with tail fragments growing much faster in size (Fig. S3A-G). Within two weeks, most of

these fragments had regained the characteristic shape of intact worms. Sagittally-cut fragments

first shrank in size before subsequently growing (Fig. S3G), and appeared to take longer to fully

regain normal worm-like appearance (Fig. S3C,D). All reproductive organs gradually

regenerated, with growth dynamics contingent on the nature of the injury (Fig. 2B-J; Fig. S3A-

F,H,I). Additionally, we asked how reproductive organs change in another context: degrowth.

Anecdotal observations suggested *H. miamia* tolerates long periods of starvation but gradually

shrinks in size when deprived of food. We used this starvation-induced de-growth to ask

whether reproductive organs scale with decreasing body size. We starved a cohort of adult

worms and quantified the de-growth of their reproductive organs over time (Fig. 2K,L). We found

that all worms survived over three months of continuous starvation, gradually shrinking over

time (Fig. 2M). Worms maintained their aspect ratios as they shrank (Fig. 2K,L,N), and their

reproductive organs shrank correspondingly (Fig. S4A-E). Partial regression analysis of

shrinking ovaries showed that they shrink synchronously (Fig. S4F), suggesting active

coordination of de-growth across the left-right axis.

The stepwise growth relationships we found between reproductive organs and body size

during post-embryonic development largely generalized to both regeneration and de-growth

(Fig. S4G-I). Starving worms lost most reproductive organs in the opposite order to which they

developed them, first losing fertilized eggs, then ovaries, and eventually losing their seminal

vesicle (Fig. 2O,P). However, after three months of starvation, most worms continued to retain

their penes despite being below the threshold size at which we would expect to see loss of this

organ (Fig. 2O), perhaps because rigid, sclerotized structures degrade differently from soft

tissue. In regenerating tail fragments, the male reproductive structures generally developed

before the female reproductive structures, following similar stepwise patterns to developing

juvenile worms (Fig. 2Q). Scaling coefficients for each organ were statistically distinguishable

across development, regeneration, and de-growth, but the effect sizes of the differences

between coefficients were generally small (Fig. S4G-I, Table S3). Analyzing this effect in more

detail, we found that unsurprisingly, amputated worms initially display aberrant organ scaling but

recover typical scaling as they regenerate (Fig. S4J-L). Qualitatively, across all of these growth

contexts, reproductive organs scaled with body size in similar ways (Fig. S4G-I, Table S3).

In sagittally-cut worms, each worm fragment retained one of its two ovaries immediately

after amputation. These ovaries first partially degenerated before growing back (Fig. 2G; Fig.

S3C,D), consistent with the reported loss of germline tissue during early regeneration in heads

regrowing tails (Hulett et al., 2023). Body size also decreased after sagittal amputation (Fig.

S3G), but ovarian degrowth was disproportionate (Fig. S4M). The rate of ovarian degrowth

(normalized to body size) was significantly greater than the rate of normalized ovarian degrowth

during starvation (Fig. S4M). Consistent with studies showing extensive cell death during early

regeneration in other systems (Ballarin et al., 2008; Pellettieri et al., 2010; Rychel and Swalla,

2008), this suggests that an active destructive process is responsible for degrowth during ovary

regeneration in acoels. The missing ovary began to grow back roughly one month after

amputation. We found that the ovaries grew asymmetrically in these worms, with the new ovary

growing faster (Fig. 2G). This suggests the existence of an active growth mechanism to ensure

that both ovaries reach a symmetric target size.

Together, our results suggest the existence of a size-associated program that regulates

the development of reproductive organs, as well as active tissue growth and destruction

mechanisms to achieve organ symmetry. It is likely that this program regulates reproductive

organ growth in a variety of different developmental contexts, including the transition from

juvenile to adult worm, regeneration after injury and tissue loss, and during starvation-induced

de-growth. With this high-level understanding of male and female reproductive system anatomy

and the scaling relationships that govern their formation in hand, we next sought to understand

these systems at higher resolution in order to decipher their functional morphology.

Figure 2: Reproductive organ development follows similar patterns in different growth contexts.

(**A**) Schematic of regeneration of the penis, seminal vesicle, and ovaries following three different

amputations. Shading indicates the tissue that regenerates. (**B-J**) Growth dynamics of reproductive

organs (within column) for each of three amputations (within row). Error bands show SEM. (**K**) Time

course of a starving worm undergoing de-growth and step-wise loss of reproductive organs. (**L**)

Schematic of reproductive organ degradation as seen in (**K**) over the course of starvation-induced de-

growth. (**M**) Worm length decreases over the course of degrowth (R^2 = 0.85, p < 0.0001). Error band

shows 95% confidence interval. (**N**) Worms shrink as they grow; their lengths and widths decrease

proportionally (R^2 = 0.73, p < 0.0001). Error band shows 95% confidence interval. (**O**) Across different

growth contexts, reproductive organs appear or disappear at roughly consistent body lengths. Ranking

the order in which reproductive structures are gained in regenerating worms (**Q**) and lost in worms

undergoing de-growth (**P**) shows that organs are gained and lost in roughly the same order in all growth

contexts. The x-axis shows individual worms in these plots. dps = days post onset of starvation. dpa =

days post amputation. Scale bars: 1 mm.

Fine structure of the male reproductive system

Next, to understand the organization of male reproductive structures at high resolution, we

visualized them using a combination of histology, immunofluorescence, staining with live dyes,

and fluorescence *in situ* hybridization (FISH) (Fig. 3; Fig. S5A). The most anterior part of the

male copulatory apparatus is the penis. This structure is difficult to visualize under reflected

white illumination (Fig. 1C; Fig. S1A), but can be seen when worms extend it, which we have

observed them do both spontaneously and during mating (data not shown). Staining fixed

worms with the dye SiR-actin labels the structure clearly, and can be visualized well in smaller

adult worms more amenable to confocal microscopy, enabling a high-resolution view of the

structure (Fig. 3A-C). Components of the penis can also be seen clearly in histological sections

of adult worms (Fig. 3D). These show that the penis contains a bundle of rigid needle-like

structures (referred to as stylets) (Fig. 3E-G, Fig. S5B,C). Our observations of the stylets match

the description of sclerotized needles described in other species in the genus and in other

studies of *Hofstenia miamia* (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et

al., 2007; Steinböck, 1966). The stylets are situated immediately posterior to the base of the

penis sheath – a long, conical structure (Fig. 3C; Fig. S5E,F; Video 1). The cavity inside this

sheath has previously been referred to as the 'male antrum' (Beltagi and Mandura, 1991; Bock,

1923; Corrêa, 1960; Hooge et al., 2007; Steinböck, 1966). The base of the sheath is cup-

shaped, and is lined with posterior-facing hair-like extensions (Fig. 3G). At its anterior end, it

connects to the male gonopore (Fig. 3C). Both the gonopore and the sheath are lined with cilia

(Video 1,2), consistent with reports in other hofsteniid species (Beltagi and Mandura, 1991;

Bock, 1923; Corrêa, 1960; Hooge et al., 2007; Steinböck, 1966). During mating, the penis

stylets are likely pushed anteriorly, through the sheath and out of the male gonopore. This could

evert the sheath, resulting in a penis-extension state in which the sheath emerges through the

gonopore around the stylets, with the ring of protrusions now near the anterior tip of the penis. It

is possible that these protrusions have a sensory function, and could inform fine penis

movements during mating. Immunofluorescence with an antibody against FMRFamide (see

Methods) revealed a set of cells that resemble neurons encircling the base of the sheath (Fig.

S5E). Further studies will be needed to understand how these components regulate copulatory

behavior.

Next, we focused on the teardrop-shaped prostatic vesicle, which is situated anterior to

the seminal vesicle (Fig. 1C,D) and immediately posterior to the penis stylets (Fig. 3H,I). Our

histological studies show that the prostatic vesicle (Fig. S6C-E) is surrounded by glands (Fig.

S6F-H). The secretions of this organ are thought to mix with the sperm as it passes through

(Faubel, 1983; Hyman, 1951). Unexpectedly, our immunofluorescence experiments also

revealed a layer of cells enclosing the prostatic vesicle (Fig. 3I). This layer may be an epithelial

tissue most likely associated with the prostatic glands. The interior of the prostatic vesicle

contains densely-packed sperm cells (Fig. 3D,I; Fig. S6C-G).

The seminal vesicle is the posterior-most organ of the male copulatory apparatus (Fig.

3J). Histological staining as well as immunofluorescence staining of the muscle marker

Tropomyosin, showed that the seminal vesicle and prostatic vesicle are both encircled by layers

of muscle (Fig. S5F; Fig. S6C-H), consistent with previous descriptions (Beltagi and Mandura,

1991; Bock, 1923; Corrêa, 1960; Steinböck, 1966). Histological and nuclear staining of the

seminal vesicle in adult worms (Fig. 3J-L; Fig. S5F; Fig. S6G,H) reveal that it consists primarily

of sperm cells. To confirm this, we dissected the seminal vesicle out of adult worms, and stained

it with a nuclear dye. We found that individual cells emerged from the structure over time, and

high-magnification imaging showed that they were indeed sperm cells, each with a ~23µm-long

nucleus and a ~22µm-long flagellum (Fig. 3M). During one mating event, we observed a failed

attempt at insemination by a worm that resulted in the release of a packet of sperm outside its

partner. We recovered this 0.6 mm-long packet of sperm (Fig. S5G) and found that it consisted

of cells with the same morphology as those in the seminal vesicle (Fig. S5H), confirming that

these are indeed sperm. As an aside, acoel sperm has historically been described as

"biflagellate", intended to mean that sperm cells each contain two axonemes, and this property

is thought to be a defining synapomorphic character of acoels (Achatz et al., 2013; Petrov et al.,

2004). We show that *H. miamia* sperm have a single flagellum, and suggest that the axonemal

properties of the flagellum may be a more accurate taxonomic character than the external

flagellar morphology.

Next, we sought to identify the testes (the organs of sperm production). Consistent with

previous observations (Beltagi and Mandura, 1991; Bock, 1923; Corrêa, 1960; Hooge et al.,

2007; Steinböck, 1966), nuclear staining and histological studies of adult worms revealed dense

clusters of sperm in the anterior parenchyma, wrapping around the worm's pharynx (Fig. 3N-U).

Tropomyosin labeling showed that the testes are embedded within layers of muscle (Fig. S5I,J).

Next, inspecting germline markers identified in previous single-cell RNA sequencing data (Hulett

et al., 2023), we identified one likely to label testes: the gene *pa1b3-2*. FISH for *pa1b3-2* reveals

that this gene does indeed specifically label testes, and that its expression coincides well with

the dense cellular clusters visible through nuclear staining (Fig. S7A,B).

Finally, we studied the development of the male reproductive system, using confocal

microscopy to visualize the maturation of its components at high resolution. Hatchling worms do

not possess any visible reproductive structures. Actin staining reveals that, within the copulatory

apparatus, the sheath and stylets emerge first, followed by a gradually-inflating prostatic vesicle

(Fig. 4A). The hair-like extensions of the sheath grow and become more conspicuous over time

(Fig. 4A). Next, we used two methods to visualize testes maturation. First, reasoning that

nuclear staining may not allow us to visualize testes at the earliest developmental stages, we

performed FISH to detect mRNA for *pa1b3-2* in juvenile worms of various sizes. In the smallest

worms (i.e. ~<2 mm), there was no expression (Fig. 4B-C). In larger, reproductively immature

worms, we detected expression of *pa1b3-2* in two lateral clusters that expanded slightly along

the anterior-posterior axis as development progressed (Fig. 4B-C). Since adult testes are a

single cylindrical structure that wraps around the pharynx, this early developmental pattern

suggests that the two lateral structures must grow along the dorso-ventral and medio-lateral

axes until they meet in the middle. Second, to test this hypothesis, we performed nuclear

staining on transverse sections of worms across a broader section of development (Fig. 4D; Fig.

S7C). In small worms, the testes are lateralized and do not meet on the dorsal surface. In larger

worms, these lateral regions have expanded across the dorsal surface of the worm to form a

single apparently-continuous region (Fig. 4E). Many acoel species are thought to have paired,

lateral testes (Table S1), a feature of taxonomic importance (Jondelius et al., 2011). While

juvenile *H. miamia* have paired testes, these organs morph into a single cylindrical structure in

the transition to adulthood. As lab-reared worms can be substantially larger than those found in

the wild, these results raise the possibility that other acoels may undergo similar morphological

changes during development.

Together, our data provide a high-resolution view of *H. miamia's* elaborate male

reproductive morphology. Consistent with our quantitative data, we find that this morphology

emerges in a stereotyped developmental sequence during the transition to adulthood in which

the penis sheath develops first, followed by nascent testes, the stylets, and the membrane-like

structure enveloping the prostatic vesicle. Sperm presumably from the testes then travels to the

prostatic vesicle. Over time, as all structures grow, the hair-like projections on the back of the

sheath become more prominent, the testes extend both along the anterior-posterior axis and

dorsally to wrap around the head, and sperm cells fill the seminal vesicle. The prostatic and

seminal vesicles appear to be surrounded by muscle, epithelia, and gland cells. Our histological

sections suggest that at least in adult worms, sperm cells may enter through the posterior of the

seminal vesicle and travel forwards into the prostatic vesicle. We do not yet understand how this

occurs in juvenile worms, how sperm cells navigate to these vesicles from their varied points of

origin, or how they are hypodermically injected during mating.

Figure 3: The male reproductive system of *Hofstenia* includes a penis with a collection of needles,

a sperm containing organ (seminal vesicle), and two testes that span the dorsal surface. (**A**) A

schematized view of the ventral surface of the worm with male reproductive structures highlighted in red.

(**B**) Schematic of male reproductive structures with the copulatory apparatus (excluding the seminal

vesicle) highlighted. (**C**) Labeling with an actin dye (white) labels the male gonopore, sheath, penis

stylets, and prostatic vesicle. (**D**) A histological section also reveals these organs. (**E**) Schematic of the

male reproductive system, with the penis stylets highlighted. (**F**) The stylets are a bundle of needles

labeled by actin. (**G**) The posterior of the penis sheath terminates in a ring of hair-like projections, also

labeled by actin. (**H**) Schematic of the male copulatory apparatus, with the prostatic vesicle highlighted. (**I**)

Actin staining with a nuclear label (Hoechst) shows that the prostatic vesicle is enveloped by a thin

epithelium-like layer, and contains densely packed sperm. (**J**) Schematic of the male copulatory

apparatus, with the seminal vesicle highlighted. (**K**) The morphology of the copulatory apparatus in

mature, adult worms is similar to that of early adults (previous panels). (**L**) The seminal vesicle of this

adult worm contains densely-packed sperm. (**M**) Dissecting out an adult seminal vesicle allows labeling of

individual sperm cells, showing their distinctive morphology. (**N**) Schematic of a transverse view of an

adult worm's anterior, showing the relative organization of the seminal vesicle and testes. (**O**) Transverse

sections show that testes appear as a continuous structure that spans the dorsal surface of the worm. (**P**)

The testes extend through the dorso-ventral axis of the worm and wrap around the head. The pharynx

(labeled and circled with a dotted line) contains residual food. (**Q**) Schematic of the male copulatory

apparatus, with the testes highlighted. (**R-T**) Nuclear staining on an adult worm, cut sagittally, reveals the

testes, which contains dense bundles of sperm organized around clusters of cells in the parenchyma. (**U**)

Histological sections confirm this organization of the testes. Scale bars: 20µm (C, U), 10µm (F-G, M),

50µm (D,I,S, T), 100µm (K,L,R), 200µm (O, P).

Figure 4: Stepwise emergence of individual components of the male reproductive system. (**A**)

Actin-dye labeling shows how the male reproductive system changes over the course of post-embryonic

development (shown here from left to right). The sheath and stylets emerge first, followed by the

appearance of the prostatic vesicle. (**B**) FISH for the male germline marker *pa1b3-2* results in two regions

of ventrolateral expression that extend along the dorsal-ventral axis to different depths. Images are

organized by pseudo-time: from least-developed (and smallest) on the left, to most-developed (and

largest) on the right. Panels in (**C**) show depth-coloration, showing that the testes extend through the

dorso-ventral axis. (**D**) Cross-sections of worms at different points in reproductive development stained

with nuclear dye show that testes grow towards the dorsal surface and eventually meet to form one

continuous structure. (**E**) Cartoon schematic of cross-sections shown in (**D**). Scale bars: 20μm (A), 100μm

(B, C), 200μm (D). Estimated worm lengths (wl) are noted under each panel.

Fine structure of the female reproductive system

As observed through imaging of adult worms using a stereo microscope, the female

reproductive system in *H. miamia* consists of two lateral ovaries running longitudinally along

both sides of the body, and one medial cluster of mature eggs located posterior to the pharynx

(Fig. 5A-C). To visualize the structure of the ovaries, we used FISH to label mRNA of a

previously-identified germline marker: *cgnl1-2* (Hulett et al., 2023). We found that this gene

specifically labeled the oocytes (Fig. 5D). Moreover, immunostaining with a custom antibody

against Piwi-1, a stem cell and germline marker, also labeled the ovaries (Fig. 5E). We were

also able to visualize the ovaries in histological sections (Fig. 5C). These methods, together with

our earlier images of ovaries at different worm ages, showed that oocytes were not organized

by maturity within the ovary (Fig. S8A). We did not detect a membrane or lining that envelopes

the ovaries. This is consistent with previous work that suggests that acoels lack true ovaries

sensu stricto: oocytes appear and develop within the parenchymal tissue without a specialized

membrane that forms a discrete organ (Eckelbarger and Hodgson, 2021; Rieger et al., 1991;

Schmidt-Rhaesa, 2007).

Oocytes in *Hofstenia* are surrounded by follicular cells, cells lining developing oocytes

that provide nutrition and secrete the eggshell (Bock, 1923; Rieger et al., 1991). We visualized

this cell layer with immunofluorescence for Piwi and histology and found that it was consistently

present around every oocyte in the ovaries irrespective of their developmental stage (Fig. 5F-G;

Fig. S8B-D). We observed that many follicle cells have very large nuclei, perhaps the result of

the fusion of neighboring cells (Fig. 5G; Fig. S8B-C). We also found clusters of sperm in the

layer of follicular cells surrounding maturing oocytes of a variety of developmental stages (Fig.

5F,G, insets), consistent with Bock's (Bock, 1923) proposal that the follicle may 'trap' sperm and

control the fertilization of maturing oocytes. We observed significant morphological variation in

the cellular components of the oocytes within the ovaries. Early-stage oocytes appeared small,

round or oval, although others showed irregular shapes. All oocytes contained a granular

substance primarily distributed peripherally and a germinal vesicle occupying a substantial

portion of the oocyte (Fig. 5H-J). The presence of a germinal vesicle in the oocytes in *H.*

miamia's ovaries suggests that these oocytes are unfertilized. It is likely that oocytes mature in

the ovaries, the germinal vesicle breaks down, and then oocytes are fertilized. These mature,

fertilized eggs then travel from the ovary to the central cavity prior to being laid.

We could not determine conclusively where or when fertilization happens, or where the

egg capsule (chorion) is produced. However, the spatial organization of the female reproductive

system suggests that an oocyte must mature, become fertilized, and then be transported to the

central cavity behind the pharynx. Across our internal examination, we did not identify any

oviducts, canals or additional structures that could facilitate this migration from the ovaries to the

central cluster. How eggs are transported thus remains unknown.

Figure 5: The female reproductive system includes embryos stored at the base of the pharynx and

oocytes in ovaries. (**A**) A schematized view of the ventral surface of the worm with female reproductive

structures highlighted. (**B**) Eggs near the pharynx of the worm (within the red circle) are fertilized and

mature while oocytes in ovaries (red arrow) are immature or unfertilized, with a visible germinal vesicle.

(**C**) A sagittal histological section shows that the ovaries contain oocytes of varied size and maturity

embedded in the parenchyma. (**D**) FISH shows that *cgnl1-2* labels immature oocytes in the ovaries. (**E**)

Oocytes in ovaries are also labeled by a Piwi-1 antibody. (**F**) A histological transverse section of an

immature oocyte encircled by follicular cells. Inset: sperm cells appear to be trapped in the follicle. (**G**)

Piwi-1 immunofluorescence confirms the organization of follicular cells, and nuclear staining sometimes

identifies sperm apparently trapped in its surface (inset). Histology also shows that immature oocytes may

have irregular shapes (**H**), contain a germinal vesicle (**H,I**), and possess an abundance of (likely yolk)

granules (**I,J**). Blue arrows label germinal vesicles in all relevant panels; yellow arrows label sperm; white

arrows label follicular cells. Scale bars: 100µm (B, C (inset), D-F), 500µm (C), 50µm (G, H-J)

Egg-laying behavior

Egg laying has only been directly observed a few times in acoels, mostly in species within the

Convolutidae (Costello and Costello, 1939; Gardiner, 1898). From these observations, acoels

are known to lay eggs through the female gonopore (if present) (Gardiner, 1898), a mode of egg

laying used by many flatworms (Tong and Ong, 2020), or through breaks in the body wall (Apelt,

1969; Costello and Costello, 1939). It has also been suggested that acoels could lay eggs

through the mouth; however, direct observation of such egg laying has been challenging (Apelt,

1969; Costello and Costello, 1939; Watzin, 1984).

Previous authors have speculated that *H. miamia* lay eggs through breaks in the body

wall, or perhaps partially through the mouth, but egg laying was not observed (Bock, 1923;

Steinböck, 1966). To observe how worms lay eggs, we isolated gravid adult worms and filmed

them from underneath for a 24 hour period. 3 worms laid a total of 25 eggs in these conditions.

We found that worms exclusively laid eggs through their mouths in events typically lasting less

than 2 minutes (Fig. 6A,B; Video 3; Fig. S9A). During this event, the worm performs a series of

muscle contractions to transfer a single egg from the ventral pocket of fertilized eggs into the

pharynx, and then applies further muscle contractions from posterior to anterior to move the egg

to the mouth. The worm then places each egg on the substrate with its mouth, likely secreting

mucus to attach the egg to the substrate (Fig. 6A,B; Video 3). Observations from these videos,

as well as of egg clutches in our culturing tanks, show that worms can either deposit a single

egg in one location or lay eggs in one or more clutches.

To understand the temporal dynamics of egg-laying in *H. miamia*, we then quantified the

time course of egg-laying in several contexts. We found that adult worms with previous access

to mates, once isolated, continued laying eggs for over a month (Fig. 6C). Juvenile worms

reared to adulthood in isolation also laid a single burst of eggs (Fig. 6D), multiple months after

first being isolated (although the timing of this burst may depend on rearing conditions and the

health of the worms). Consistent with our finding that worms reared entirely in isolation can lay

eggs (Fig. 1L,M), this burst of egg laying is also likely the result of a selfing event in which the

worms fertilize their own eggs, or possibly the result of a form of parthenogenesis allowing the

activation of unfertilized eggs. We also allowed virgin worms to mate once, in controlled

conditions, and subsequently isolated them. These mated and inseminated worms laid eggs for

over three months after a single mating, suggesting that received sperm may be stored in

follicular cells for several months (Fig. 6E).

Next, we asked how much control the worms had over their egg laying. To test whether

worms choose specific locations to lay eggs, we quantified the spatial positions of eggs laid in

their culturing tanks. We found that worms have a strong preference for laying eggs on the walls

of their containers, rather than on the floor (Fig. 6F). Eggs laid on walls are preferentially laid

close to the water line (Fig. 6G, Fig. S9B). This preference is not simply because the worms lay

eggs where they are: we observed that worms spend the majority of their time on the floors of

their containers, and seem to glide up the walls specifically to lay eggs (Fig. S9C). These spatial

preferences do not seem affected by food availability: when deprived of food, egg laying still

primarily occurs on the walls of culture boxes (Fig. S9D). These data suggest that worms make

active substrate choices for egg laying.

We also observed that worms lay 49% of their eggs in clutches. Individual worms can

produce these clutches (Video 3). Clutch sizes are often small (Fig. S9E), but the largest can

contain over 30 eggs deposited within a 3-4 day window in culture boxes containing 20-50

worms (Fig. 6H, Fig. S9E). The vast majority of individual worms lay fewer than 10 eggs in a 3-4

day period (Fig. S9F), suggesting that some clutches may be communal. Indeed, in other

culturing conditions with large groups of worms, clutch size often exceeds 140 eggs (Fig. 6I). To

test whether worms lay eggs in communal clutches, we allowed worms to lay eggs in containers

for 3 days. We then swapped worms between containers (or, for control containers, we

swapped worms and removed old eggs). We asked how worms interacted with egg clutches laid

in the first 3-4 days, and found that many new eggs were laid in pre-existing clutches (Fig. 6J,

Fig. S9G). Worms added new eggs to 42 percent of old egg clutches. This shows that worms

frequently lay eggs in communal egg clutches.

Finally, we asked whether worms withhold eggs in suboptimal environments. Our

observations (see Fig. 2K-L) suggested that worms are physiologically resilient to food stress.

We therefore conducted an experiment with a set of worms, half of which were deprived of food

for one feeding period (3-4 days). The other set was fed normally during this period. We found

that food-deprived worms laid significantly fewer eggs during this period (Fig. 6K). We

hypothesized that this reduction in egg-laying was because the worms were actively withholding

their eggs in the absence of food. Next, we repeated this experiment, first depriving some

worms of food and subsequently feeding them. After food-deprived worms were given food, they

laid an excess of eggs (Fig. 6L), showing that the worms withhold egg-laying in food-limited

environments. More generally, these data show that *H. miamia* assess their environments to

decide when and where to lay eggs.

Figure 6: *H. miamia* lays eggs through the mouth and exhibits environmental preferences in egg laying. (**A**) Sequence of images from a video of egg-laying through the mouth. Eggs in the pharynx and emerging through the mouth are shaded blue. (**B**) Schematic showing presumed process of embryo traveling from the cavity beneath the pharynx to the pharynx and then out through the mouth. (**C**) Histogram showing the timing of eggs laid by adult worms living in communal tanks and then isolated. (**D**) Histogram showing the timing of eggs laid by worms that undergo reproductive development in isolation and then self-fertilize. (**E**) Histogram showing the timing of eggs laid by worms that are allowed to mate once. (**F**) Scatterplot of the percentage of eggs found on the floor of communal tanks (n = 30). This is significantly different from the expected percentage of eggs based on tank surface area (t-test p < 0.0001). (**G**) Kernel density estimate of egg locations on a subset of tank surfaces with similar dimensions (n = 2144 eggs). (**H**) Density-based spatial clustering of egg coordinates shows that eggs are laid in clutches. Number of eggs in a clutch shown in white. (**I**) In some culturing conditions, worms lay clutches of up to 145 eggs. (**J**) New worms add eggs to pre-existing clutches laid by other worms. (**K**) Worms that are unfed for 4 days lay fewer eggs than fed worms (n = 9 tanks, t-test p < 0.0001). (**L**) Unfed worms that

are subsequently fed lay more eggs than worms that are continuously fed (n ≥ 8 tanks, t-test p < 0.0001).

Discussion

Our work describes the reproductive life history of an acoel across its life cycle (Fig. 7A). We

reveal many new facets of acoel biology, describe the structure and dynamics of growth,

regeneration, and egg-laying behavior, and establish a foundation for the experimental study of

reproductive biology in acoels. Below, we discuss the significance of our findings for

reproductive development, behavior, and the evolution of life history strategies.

Reproductive development and regeneration

Little is known about the development or regeneration of reproductive structures in acoels. Our

data show that in *H. miamia*, reproductive structures develop and regenerate in a stereotyped

sequence, with male organs appearing before female ones. During starvation-induced de-

growth, these organs degenerate in the opposite sequence. This stereotypy is unexpected and

may not be a universal feature of acoels: sparse data from three other acoel species,

Solenofilomorpha funilis (Crezée, 1975), *Otocelis luteola* (Kozloff, 2000a) and *Aphanostoma*

pulchra (Perea-Atienza et al., 2013), indicate that these species do not exhibit identical

sequences. The stereotypy we observe is consistent with the idea that a single, size-associated

program regulates reproductive organ development in *H. miamia*, and that it may be deployed in

a variety of growth contexts.

Our results also show that *H. miamia* is capable of regenerating all reproductive structures from

a variety of initial tissue configurations. The trajectories toward organ replacement vary based

on this initial state, and reveal important features of the regenerative process. For instance, we

found that worms missing their sagittal halves or their tails first shrink in size before re-growing.

This may be in part because the early stages of wound closure and regeneration require the

reorganization of existing tissues. In any case, this reduction in body size is associated with

disproportionate reductions in reproductive organ size, demonstrating active tissue destruction

mechanisms, possibly mediated by apoptotic processes similar to those reported in other

regenerative species (Ballarin et al., 2008; Pellettieri et al., 2010; Rychel and Swalla, 2008). In

addition, we found that the two ovaries within a worm grow in a significantly correlated manner,

and after sagittal amputation, the missing ovary grows disproportionately to regenerate

symmetry. In principle, such bilateral symmetry could be achieved passively, without any

feedback (Reddien, 2018). However, the strong within-worm correlation, even after controlling

for body size, suggests that there is either organism-specific coordination of growth or death

rates, or more likely a feedback process that ensures bilateral symmetry.

In many animal lineages such as arthropods, vertebrates, planarians, and nematodes, organs

typically grow symmetrically, and scale with body size (Boulan and Léopold, 2021; Ko et al.,

2024; Uppaluri and Brangwynne, 2015; Wolpert, 2010). The mechanisms underlying this

coordinated, scaled growth are not fully understood, and may vary across tissues and species.

For instance, some vertebrate tissues appear to grow using cell-intrinsic programs; others - and

many insect tissues - appear to rely on central coordinating mechanisms that read signals of

growth from the circulation or from unknown sources (Allard and Tabin, 2009; Boulan and

Léopold, 2021; Harris et al., 2021; Trible and Kronauer, 2017; Vallejo et al., 2015; Wolpert,

2010). How different organs ensure appropriate, coordinated scaling remains poorly

understood. We find that *H. miamia* must also solve this problem: their reproductive organs

scale with body size during development, regeneration, and de-growth, and an active

coordination mechanism ensures that the ovaries return to symmetry after perturbations. How

do tissues across the body 'read' the same indicators of size, and how is their growth

coordinated? Our work establishes a foundation for the experimental study of these and other

developmental phenomena.

Egg-laying physiology and behavior

Despite speculation, few acoels have been directly observed laying eggs (Costello and Costello,

1939; Gardiner, 1895; Watzin, 1984). We find that *H. miamia* lays fertilized eggs through its

mouth: a mode of egg-laying not found in other animals. This unusual behavior raises multiple

questions. First, where in the animal does fertilization occur? Our observations suggest that

follicular cells surrounding immature oocytes may act as a selective barrier to sperm, preventing

many sperm cells from reaching the oocyte, a function suggested by Bock (1923). Additionally,

given that *H. miamia* can lay eggs for months after a single mating despite lacking a seminal

bursa, it is likely that the follicular cells function as a sperm storage organ. Second, we find that

H. miamia appears capable of self-fertilization, as it can lay eggs without mating. While some

acoels are capable of asexual reproduction through fissioning and budding (Åkesson et al.,

2001; Ax and Schulz, 1959; du Bois-Reymond Marcus, 1955; Hanson, 1960; Ishikawa and

Yamasu, 1992; Zabotin and Evtugyn, 2021), self-fertilization has not been reported previously.

How does sperm travel to the ovaries? It is likely that sperm cells migrate posteriorly from the

testes towards the ovaries in isolated worms, and from arbitrary body regions towards the

ovaries after mating. In addition, sperm must travel from all regions of the testes to the seminal

and prostatic vesicles. Whether they use chemical cues to navigate towards oocytes and these

vesicles, and the mechanics of this process, remain unknown. Third, how do mature eggs travel

from the lateral ovaries to the central cavity, and how are the eggs in this cavity physically

loaded into the pharynx? Our anatomical observations have so far failed to reveal obvious

oviducts, or other tube-like structures that could facilitate movement of eggs into the pharynx.

The only known opening at the posterior of the pharynx is the pharyngeal sphincter that allows

food to pass into the gut. We therefore speculate that eggs travel through the gut to the central

cavity, which may simply be a pocket within the gut. Eggs may then be moved into the pharynx

through a form of reverse peristalsis (and observation of the worm prior to egg-laying shows

waves of muscle contraction proceeding posterior-to-anterior as each egg is loaded into the

pharynx).

We also find that *H. miamia* appears to make active choices about when and where to lay

fertilized eggs. *H. miamia* has clear spatial preferences for egg-laying in our culture chambers.

Worms often lay eggs in communal clutches, sometimes adding to another worm's clutch. They

also have environmental preferences, withholding egg-laying when deprived of food and then

quickly laying eggs once food is present again. Together, these results show that *H. miamia*

surveys its environment and then integrates this assessment into whether - and where - it is

suitable to lay an egg. More generally, the existence of communal egg clutches raises the

possibility that acoels may have a rich and unexplored social repertoire.

Evolution of reproductive strategies within acoels

Placed in a comparative context, our description of *H. miamia*'s reproductive life history reveals

that acoels and their sister lineage, the nemertodermatids, display a striking diversity of

reproductive morphologies, and likely a corresponding diversity in reproductive behavior (Fig.

7B). This diversity reveals correlated suites of reproductive traits that suggest a small number of

life history strategies. For instance, the presence of a bursa and female gonopore are

associated with a muscular penis, suggestive of cooperative, genital-handshake style mating.

The absence of a bursa and female gonopore are associated with a needle-laden penis,

suggestive of competitive, hypodermic-insemination style mating. Comparative work in the

distantly-related flatworms shows that these anatomical and behavioral traits indeed coevolve

similarly (Brand et al., 2022a, 2022b), in accordance with social evolutionary theory on sexual

conflict (Charnov, 1979; Michiels and Newman, 1998; Ramm et al., 2015). Within the genus

Macrostomum, a competitive hypodermic insemination syndrome has evolved at least 14 times

from ancestral cooperative, reciprocal insemination (Brand et al., 2022a, 2022b). This syndrome

involves the correlated evolution of a sharpened penis, a simplified female sperm-receiving

organ, sperm cells lacking bristles and other adaptations for post-copulatory sexual conflict, and

associated behavioral changes (Brand et al., 2022a, 2022b, 2022c; Schärer et al., 2014, 2011).

Whether similar anatomical associations truly predict mating strategies in acoels remains

unknown. In addition, the variability of some morphological features does not obviously fit this

pattern of cooperative vs competitive reproductive strategies. For example, the relative locations

of the testes and ovaries is highly variable, as is the number of bursal nozzles, suggesting that

there may be further evolutionary patterns awaiting explanation. Our work establishes an

approach to study reproductive anatomy and behavior in a model acoel. This approach can be

applied to many other acoel species. Given their rich diversity, acoels are a promising clade in

which to study the evolution of reproductive strategies, and in which to test the generality of

theories of sexual conflict.

Conclusion

These findings establish foundational knowledge of anatomical, physiological, and behavioral

elements of *H. miamia*'s reproductive life history. This enables future work on the molecular

genetics of reproductive organ development and regeneration, on the physiological processes

involved in egg maturation and oviposition, and on the neuroscience of reproductive behavior.

Figure 7: Reproductive life histories in Acoelomorpha. (A) The life cycle of *Hofstenia miamia*,

with major reproductive events displayed. **(B)** Family-level phylogeny of Acoelomorpha

(Nemertodermatida, Acoela) showing anatomical and reproductive life history traits (Table S1): position of

the mouth, whether gonads are mixed or separated by sex, penis type, paired or unpaired testes, paired

or unpaired ovaries, presence or absence of a female gonopore, presence or absence of a seminal

bursa, the number of associated bursal nozzles, egg-laying mode, mode of sexual reproduction,

alternative reproductive strategies, and regenerative capacity (see Table S4 for definitions of terms and

categories). Schematic diagram of the reproductive anatomy of representative species from each family

within Acoelomorpha with specific structures colored: male copulatory organ (purple), sperm in testes

and/or seminal vesicle (blue), oocytes (red), female or shared gonopore and/or bursa (green). (Table S1).

White boxes represent unknown phenotypic states, and in the case of asexual reproduction, its possible

absence. Phenotypic classifications are from (Achatz and Hooge, 2006; Apelt, 1969; Bailly et al., 2014;

Beltagi and Mandura, 1991; Bock, 1923; Boone et al., 2011; Bush, 1975; Costello and Costello, 1939,

1938; Crezee, 1978; Dörjes, 1968, 1966; Faubel, 1976, 1974; Faubel and Cameron, 2001; Gardiner,

1895; Grae and Kozloff, 1999; Hooge, 2003; Hooge et al., 2007; Hooge and Smith, 2004; Hyman, 1937;

Kostenko, 1989; Kozloff, 2000b; Meyer-Wachsmuth et al., 2014; Peebles, 1915; Perea-Atienza et al.,

2013; Raikova et al., 1995; Riser, 1987; Shannon and Achatz, 2007; Steinböck, 1966; Sterrer, 1998;

Watzin, 1984)**.**

Methods

Animal husbandry

We reared gravid *H. miamia* in communal plastic 2.25 L boxes with approximately 1.25 L of

artificial seawater (37 ppt, pH 7.8-8.2). These communal tanks were kept in incubators held at

21°C. Twice weekly, we collected embryos, changed their seawater, and fed them with *Artemia*

sp. (brine shrimp). The embryos were raised in petri dishes. Once they hatched, we transferred

juvenile worms to tanks maintained at room temperature and fed them with marine rotifers

(*Brachionus plicatilis*). When reproductive organs began to develop, we transferred worms to

larger tanks, each typically housing 30-50 worms. To pair animals for mating, we transferred

juvenile worms from tanks to 24-well plates and reared them in isolation following similar

feeding and cleaning routines as listed above. In order to study the regeneration of reproductive

systems, we isolated adult worms from communal tanks, anesthetized them in 15% tricaine

(ethyl 3-aminobenzoate methanesulfonic acid), and then amputated them with micro knives

(Fine Science Tools #10316-14). We maintained regenerating worm fragments in 6-well plates

and cleaned plates twice weekly. Five days after amputation, we fed these worms *Artemia* sp.

(brine shrimp) or rotifers. We also maintained starving animals in 6-well plates that we cleaned

twice weekly but that were not fed. All animals are derived from an inbreeding population of

worms collected in 2010 from Bermuda.

Histology

We fixed adult specimens of *H. miamia* in 4% paraformaldehyde in artificial seawater for 24

hours at room temperature. Following fixation, specimens were washed twice with 70% ethanol

to remove the fixative and then preserved in a third change of 70% ethanol for histological

preparation. Animals were dehydrated through a graded ethanol series (95% and 100%),

cleared in Histoclear, and embedded in Leica Surgipath (Paraplast) with a melting point of 56°C.

We made longitudinal, transverse, and sagittal sections at a thickness of 5 μm using a Leitz

1512 microtome. We stained these sections using a standard protocol with hematoxylin and

eosin, mounted them on glass slides with Permount, and imaged them using a Zeiss Axio

Scan.Z1, and an Olympus BX50 microscope.

Fluorescence *in situ* hybridization (FISH)

We synthesized riboprobes following protocols established in (Srivastava et al., 2014).

Following 1-2 weeks of starvation, we fixed whole worms in 4% paraformaldehyde in artificial

sea water for one hour at room temperature. We then washed the fixed animals with PBST

(PBS + 0.1% Triton-X-100) and transferred them to 24-well plates in small baskets with a mesh

bottom, with 4-6 animals in each basket. To remove pigment autofluorescence, we treated the

animals with a bleach solution (containing 4% hydrogen peroxide, SSC, and formamide) and left

them under a light for two hours. For all washes, 800 µL of solution was used. We first

permeabilized the animals using proteinase K solution (0.1% SDS, 1 µL/10 mL proK in PBSTx).

After 10 minutes, we post-fixed the worms in 4% formaldehyde in PBST, washed twice in PBST,

and then washed in a 1:1 PBST:PreHyb solution for 10 minutes. We then incubated the

samples in PreHyb solution (50% DI formamide, 25% 20X SSC, 0.05% 10% Tween-20, 1mg/mL

yeast tRNA, 20% water) for two hours in a 56°C hybridization oven and then transferred them to

a hybridization solution (50% DI formamide, 25% 20X SSC, 0.05% 10% Tween-20, 1 mg/mL

yeast tRNA, 10% 50% dextran sulfate, 10% water) containing riboprobe(s) overnight. We then

put the specimens through several 30-minute washes at 56°C, starting with two PreHyb

washes, two 1:1 PreHyb:2XSSCT washes, two 2XSSCT (2XSSC with 0.1% TritonX-100)

solution washes, two 0.2XSSCT (0.2X SSC with 0.1% TritonX-100) solution washes. We cooled

the specimens to room temperature and then washed them in two ten-minute PBST washes.

We then performed a 1-hour blocking at room temperature with blocking solution (5% horse

serum and 5% casein in PBST). We incubated specimens with anti-Digoxigenin-POD (1:1500

dilution; Roche, 11633716001) in a blocking buffer overnight. The following day, we washed

animals 6 times with PBST, incubated with a tyramide buffer (1.1688g/mL NaCl, 6.18mg/mL

boric acid, filled with water and adjusted pH to 8.5) for 10 minutes, then developed with

rhodamine-conjugated tyramide solution for 10 minutes. We washed animals again with PBST

and incubated for 1 hour in 1:500 PBST:Hoechst to label nuclei. We mounted the animals on

glass slides with VECTASHIELD® PLUS Antifade Mounting Medium (Vector Laboratories, H-

1900). We estimated worm length (Fig. 4A-D) by measuring copulatory apparatus length

(gonopore to prostatic vesicle) or cross-sectional worm width within the image and comparing

that measurement to measurements of developing worms.

Whole-mount immunofluorescence

We fixed worms in 4% paraformaldehyde (Electron Microscopy Sciences, 15714) in artificial sea

water for 1-2 hours at room temperature (juveniles were fixed for one hour, adults were fixed for

two) before washing with PBST (PBS + 0.1% Triton-X-100). We anesthetized adult worms in 0.5

mg/mL tricaine for 5 minutes prior to fixation to minimize epidermal rupture. For

immunofluorescence, we washed worms in PBST, blocked for one hour at room temperature in

10% goat serum in PBST, and incubated them in primary antibody for 48-72 hours at 4°C on a

shaker (juveniles for 48 hours, adults for 72 hours). The following day, we washed worms

thoroughly in PBST (8 x 20 min washes) on a nutator before blocking for one hour at room

temperature on a shaker. We incubated worms in secondary antibody overnight at 4°C; the

following day, we washed them thoroughly in PBST (8 x 20 in washes) on a shaker before

adding direct conjugate dyes and mounting them on glass slides with VECTASHIELD®

HardSet™ Antifade Mounting Medium (Vector Laboratories, H-1400-10). We used the following

antibodies: Tropomyosin (custom) (Hulett et al., 2020), Piwi-1 (custom), Goat anti-Rabbit IgG

(H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568 (ThermoFisher, A-11011),

Hoechst (ThermoFisher, H1399), SiR-actin (Cytoskeleton, CY-SC001). We acquired all images

with FISH and immunofluorescence on a Leica SP8 point-scanning confocal microscope. We

generated the Piwi-1 (Rabbit polyclonal) custom antibody using GenScript as previously

described for Tropomyosin (Hulett et al., 2020). Briefly, a peptide was synthesized from the *H.*

miamia Piwi-1 protein sequence, expressed in *E. coli,* and used to immunize rabbits. The Piwi-1

antibody used in this paper was #SC1195 (0.845 mg/mL), used at the concentration of 2 ug/mL.

Quantitative analysis of development, regeneration, and de-growth

We collected 42 zygotes laid from wild-type worms. Two embryos did not hatch, and two worms

died during the course of the experiment (likely due to handling error). Four worms were

removed from the experiment at different timepoints for additional imaging, resulting in 34

worms that we followed into adulthood (Fig. S2B). We transferred each embryo to a well of a

24-well cell culture plate with artificial seawater. Plates were stored in a temperature-controlled

incubator. Without removing the embryos from the plate, we imaged them twice weekly through

a dissection microscope with white illumination from LEDs mounted above the sample. We also

changed water on these days. We added the same volume of water to each well (2.35 mL in

each 12-well). Once at least one embryo had hatched, we added rotifers to each well. On each

feeding day, we calculated the concentration of rotifers by counting the number of rotifers in

diluted samples. Based on this concentration, we changed the volume of rotifers added to each

well to ensure a consistent number of rotifers was given to each animal. At different stages

during the experiment, we increased the number of rotifers (Fig. S2A). These increases were

gradual to ensure worms always had more food than they could eat, while minimizing excess

food which could affect water quality. We removed hatched worms from wells for imaging twice

weekly. We anesthetized worms in tricaine, and mounted them on microscope slides with a

20x20 mm sticker grid with 1mm resolution (ThomasScientific, #1207X53). Depending on the

worms' pigmentation, this grid is visible through the body of some worms, but this did not impact

the visibility of reproductive organs. When worms were larger, we moved them to larger 6-well

plates in 10 mL of artificial sea water and began to feed them brine shrimp in addition to rotifers

(Fig. S2A). We measured the number of brine shrimp added using the same concentration

procedure outlined above for rotifers. Once worms had matured, we increased the number of

brine shrimp.

We manually annotated images in FIJI. We measured lengths from the most posterior

part of the male reproductive system (e.g., base of the penis tube before sperm production

begins or base of the seminal vesicle) to the male gonopore, the length of the seminal vesicle,

the length of each ovary, the length of the worm's body, and the width of the worm's body. We

calculated penis length by subtracting the length of the seminal vesicle from the length of the

male gonad. All measurements were converted to millimeters using image-specific scale

information.

We repeated these imaging and measurement processing procedures with starving

worms and regenerating worm fragments. We anesthetized and imaged starved and fed control

worms roughly once a week for 11 weeks. 106 days after isolation, we imaged the starved

worms again, at which point one starved worm had died. We imaged regenerating worms

starting 2 days post-amputation and then again every 2-5 days until 25 days post-amputation at

which point we took images less frequently. We did not feed regenerating fragments for the first

four days after amputation, and then fed them normally. In these experiments, we used a two-

pointed line or multi-jointed spline to measure lengths. We performed all analyses in python,

with code assistance from GPT4.

Egg-laying behavior

We moved adult worms from a communal tank to individual wells in the center of a 24-well plate

or 12-well plate (Fisher Scientific #07-200-82). We mounted the plate on an elevated platform

within an enclosed behavioral rig. The plate was illuminated with uniform white light, from

addressable LEDs (Adafruit #SK6812RGBW, powered by 5V DC) arranged in a ring around the

plate behind a white acrylic diffuser (McMaster-Carr #8505K741). An infrared camera (Basler

Ace ac4024-29um USB 3.0 monochrome) was mounted underneath the plate to allow filming of

the worm's ventral surface. A USB fan was mounted on the floor of the behavioral chamber for

temperature regulation. We filmed worms at 1Hz for 8-12 hours. After filming, each worm was

returned to its original communal tank.

To determine the timing of egg laying in adults with continuous access to mates, we

collected adult worms from communal tanks and each was placed in an individual well of a 6-

well plate. Every 3-4 days, we counted the number of eggs laid by each worm, removed the

eggs, changed the water, and fed worms with brine shrimp or rotifers. To induce self-

fertilization, we moved juvenile worms with immature reproductive systems from communal

tanks to 24-well plates and fed them with rotifers. Once these worms had increased in size, we

moved them to 6-well plates and fed with artemia. We recorded egg counts every 3-4 days

when water and food were refreshed. To determine the time course of egg laying after mating

once, we isolated juvenile worms. Once these worms were reproductively mature, we paired

each worm with another isolated worm and allowed them to mate. After mating, we returned the

worms to isolation. We recorded egg counts as previously described. If worms stopped laying

eggs for more than two weeks, we allowed them to mate again before re-isolating them.

To find egg positions, we maintained communal tanks according to our husbandry

protocol. Every 3-4 days, before cleaning, we recorded images of each face of the tank (2 short

faces, 2 long faces, and the floor of the tank). We then scored the x- and y-coordinates of eggs

on each face using FIJI. Since few eggs were laid on the floors of the tanks and it was unclear

whether these eggs were attached or had been dislodged from the walls of the tanks, we did not

score their locations (Fig. 6F). We also scored the x- and y-coordinates of the boundaries of the

water in each image. From these image-specific boundaries, we calculated a common set of

bounds for each type of face (i.e., short, long). We scaled egg coordinate positions and aligned

them to common bounds for analysis of spatial egg laying preferences. To ask whether worms

add to existing clutches of eggs, we cared for communal tanks as previously described but did

not remove eggs from their faces. We took images of each face, and then we transferred each

cohort of worms in a tank to a different tank. After 3-4 days, we took images of tank faces again,

scored coordinates of each egg, and recorded whether each egg was old or newly laid.

For experiments measuring how worms altered their egg-laying behavior based on food

availability, we selected 17 and 18 communal worm tanks respectively. On the first day of the

experiment, half of these tanks were randomly selected to have food (brine shrimp) withheld.

After four days with or without food, we collected and counted eggs from all boxes. To test

whether egg laying recovers when unfed worms are fed, we randomly selected communal tanks

of adult worms to have food withheld. Four days after food was added or withheld, we removed

all eggs that were laid, and then fed all communal tanks. Three days later, eggs were collected

and counted.

Author Contributions

VC, SET, and MS designed the project; SET and VC performed development, physiology and

behavior experiments; VC and SET analyzed the data; APK performed and interpreted

immunofluorescence and imaging experiments; DMB performed and interpreted histology

experiments; VC, SET, and MS wrote the manuscript with input from all authors; and MS

supervised the project.

Acknowledgements

We thank Gonzalo Giribet, Matthew Hooge, James Hanken, Javier Ortega-Hernandez, Andrew

Berry, and all members of the Srivastava lab for helpful discussion. We also thank James

Hanken, Júlia Chaumel, and Christina Daly for histology help and infrastructure. VC and APK

are Fellows of the Jane Coffin Childs Fund for Medical Research. ST acknowledges funding

from the Harvard Museum of Comparative Zoology, Herchel Smith Undergraduate Science

Research Program, and the Program for Research in Science and Engineering. This project

was supported by grants from the National Institute of General Medical Sciences of the NIH to

MS: R35GM128817 and R35GM153252.

Supplement

909   **Table S1: Survey of reproductive traits within Acoelomorpha.** see attached .xlsx file

911   **Table S2: Body length is a better predictor of organ length than age in multiple**
912   **regression.** Multiple regression coefficients and associated p-values are reported for each
913   organ. Wald's test was used on coefficients to test whether they are significantly different from
914   each other.

organ	day coefficient	day p-value	worm length coefficient	worm length p-value	wald's test p-value
penis	0.0684	0.0002	0.1639	<0.0005	0.0082
seminal vesicle	0.0425	0.0003	0.0985	<0.0005	0.0128
ovary	0.3361	<0.0005	1.0029	<0.0005	<0.0005

915
916

917   **Table S3: ANCOVA of the relationship between organ size and body size.**

	F-statistic	p-value	η^2
penis	15.82	<0.0005	0.040
seminal vesicle	24.329	<0.0005	0.069
ovary	58.905	<0.0005	0.175

919   **Table S4: Glossary of anatomical and reproductive traits.**

mouth: approximate position of the mouth on the ventral surface

central	mouth located between the anterior end and midpoint of the ventral surface
anterior	mouth located between the anterior end and midpoint of the ventral surface
posterior	mouth located between the midpoint of the ventral surface and posterior end
none	no mouth
gonad(s): tissue or organ where gametes are specified	
mixed	sperm and eggs are produced within the same tissue or organ
separate	sperm and eggs are produced in distinct tissues or organs
penis: male copulatory organ	
needle-like	penis contains hardened, sclerotized elements
muscular	absence of hardened elements; soft or glandular penis
none	penis is absent
testes: tissue where sperm are produced	
paired	there are two distinct regions where sperm are produced
unpaired	there is a single region where sperm are produced
ovaries: tissue where oocytes are produced	
paired	there is are two distinct regions where oocytes are produced
unpaired	there is a single region where oocytes are produced
fem. gonopore: female genital opening	

present	a female genital opening is present
absent	a female genital opening is not present

bursa: an organ used to hold sperm received during mating	
present	a bursa is present
absent	a bursa is not present

# bursal nozzles: the number of channels accompanying the bursa; channels that facilitate sperm movement or modification	
0	the bursa is not accompanied by any sperm channels
1	the bursa has one channel used to move sperm
>1	the bursa has more than one channel used to move sperm

egg laying: the mode of oviposition that a species has been observed to use	
mouth	eggs are deposited via the mouth
fem. gonopore	eggs are deposited via the female genital opening
body wall rupture	eggs are deposited via breaks in the body wall

mating: the mode of sexual reproduction that a species has been observed to use	
hypodermic	a copulation mode where sperm is injected beneath the epidermis of a partner
hyperdermal	a copulation mode where sperm are deposited on the epidermis of a partner
true copulation	penis inserted in the female gonopore of a partner; insertion can be mutual or one-sided

alt. reproductive strategies: forms of reproduction other than two individuals mating	
fission	splitting of an individual into two or more individuals
budding	development of an outgrowth and then detachment of that outgrowth to form offspring
self-fertilization	zygote forms from fusion of sperm and egg from a single individual

Figure S1: Gross reproductive morphology is visible from the ventral surface of *H. miamia*. (A) The

male reproductive system includes the male gonopore, penis, prostatic vesicle, and seminal vesicle. (**B**)

H. miamia's ovaries are lateral, and organized along the anterior-posterior axis. There is no visible

organization of oocytes by size or stage within the ovaries. (**C**) A cluster of zygotes is visible medially, and

immediately posterior to the male copulatory apparatus. (**D**) Representative time course of embryonic

development of embryos dissected from the central cavity. Scale bars: 1 mm.

Figure S2: Growth and scaling. (**A**) Time course of feeding changes through juvenile development,

showing the delayed increases in feeding for one group, and the introduction of brine shrimp late in

development. (**B**) Survival curve showing the number of worms in each of the two feeding treatments over

time. (**C**) The aspect ratio (length:width) of worms is consistent over time (slope = 0.002, R^2 = 0.002, p =

0.31), showing that worms grow proportionally over development. Error band shows 95% confidence

interval. (**D**) Worm length increases over time, with worms growing more slowly when their increased

feeding was delayed. Error band shows SEM. (**E**) Residuals of ovary length (regressed against body

length) within each worm are highly correlated (Pearson correlation coefficient = 0.7, p < 0.0001). Error

band shows 95% confidence interval.

Figure S3: Across different growth contexts, body and organ scaling rules are conserved. (**A, C, E**)

Representative time course of regenerating tail tip, sagittal cut, and head, and (**B, D, F**) schematized

version with reproductive structures shown. (**G**) Regenerating heads and tail tips increase in length

following amputation while sagittally-cut fragments shrink and then begin to grow. Error bar shows SEM.

(**H**) Ovaries scale with body length in sagittally-cut fragments. (**I**) Ovaries regenerate once a tail fragment

reaches a certain length. Scale bars: 1 mm.

Figure S4: Across different growth contexts, body and organ scaling rules are conserved. (**A**)

Worm length decreases over time when worms are starved (starved: R^2 = 0.85, p < 0.0001; fed control:

R^2 = 0.004, p = 0.48). Error band shows 95% confidence interval. (**B-D**) Penis length (starved: R^2 = 0.54,

p < 0.0001; fed control: R^2 = 0.04, p = 0.03), seminal vesicle length (starved: R^2 = 0.60, p < 0.0001; fed

control: R^2 = 0.03, p = 0.06), and mean ovary length (starved: R^2 = 0.67, p < 0.0001; fed control: R^2 =

0.02, p = 0.15) decrease over time for worms experiencing de-growth but do not change in worms that

are continuously fed. Error band shows 95% confidence interval. (**E**) The number of zygotes in the central

cavity decreases over time in both starved and fed worms, likely because these worms are isolated

(starved: R^2 = 0.48, p < 0.0001; fed control: R^2 = 0.33, p < 0.0001). Error band shows 95% confidence

interval. (**F**) When controlling for the effect of body size, the residuals of ovary length (regressed against

body length) are strongly correlated between ovaries within worms over the course of de-growth (Pearson

correlation coefficient = 0.75, p < 0.0001). Error band shows 95% confidence interval. (**G-I**) The size of

reproductive structures scale similarly (see Table S3) across development, de-growth, and regeneration

(penis R^2 = 0.70, p < 0.0001; seminal vesicle: R^2 = 0.50, p < 0.0001; ovary: R^2 = 0.76, p < 0.0001). Error

band shows 95% confidence interval. (**J-L**) The residuals of organ length (regressed against body length)

generally converge towards zero over the course of regeneration despite sometimes displaying large

initial deviations. (**M**) Sagittal fragments degrade their existing ovary during initial regeneration at a faster

rate normalized to body size (slope = -0.015) than adult worms experiencing de-growth (slope = -0.003)

(Wald's test on slope of linear regression of ovary length and day: p=0.0006). Error band shows 95%

confidence interval.

Figure S5: Fine structure of *H. miamia*'s male reproductive system. (**A**) The gross morphology and

location of the reproductive system can be seen in sagittally-cut worms with nuclear staining. (**B-C**) Penis

stylets are arranged in a bundle, seen with differential interference contrast (DIC). (**D**) Schematic of the

male reproductive structures with the copulatory apparatus highlighted. (**E**) Antibody staining for

FMRFamide shows that the penis sheath is surrounded by presumptive neurons. (**F**)

Immunofluorescence for tropomyosin, and actin and nuclear labeling enables visualization of the

musculature surrounding the male structures. (**G**) A packet of sperm ejected during mating made up of

(**H**) sperm cells. (**I**) Schematic of the male reproductive structures with the testes highlighted. (**J**) Sperm in

the testes develop in clusters interspersed with the muscle of the body, visualized by

 immunofluorescence for tropomyosin, a muscle marker. Scale bars: 500µm (A), 100µm (B, G), 50µm (C,

981 E, J), 25µm (F), 10µm (H).

Figure S6: Histology of the male reproductive system reveals the organization of the male reproductive system along different body axes. (A) The male reproductive system in a sagittal section of a juvenile worm has a male gonopore but lacks a clear male copulatory apparatus, prostatic vesicle, and seminal vesicle. **(B)** A longitudinal section showing the male gonopore. **(C-E)** Sagittal sections through the male reproductive system along the medial-lateral axis showing a channel of sperm (yellow arrows) connecting the prostatic vesicle and seminal vesicle. **(F-H)** Transverse sections moving along the

anterior-posterior axis show the positions of the prostatic vesicle and seminal vesicle and the musculature

and glands surrounding the seminal vesicle. Scale bars: 100 um.

Fig. S7: Testes emerge as lateralized regions and grow to span the dorsal surface as

worms grow. (**A**) FISH for the male germline marker *pa1b3-2* results in two regions of

ventrolateral expression that extend toward the dorsal surfaces. (**B**) Hoechst and *pa1b3-2*

mRNA expression label sperm cells. (**C**) Nuclear staining of cross-sections at different points

along the anterior-posterior axis within the head shows that the testes emerge as two lateral

lobes that gradually form a continuous cylindrical structure. Cartoons depict the plane of

sectioning (red line) and reproductive structures (white). Scale bars: 250 µm (A), 50µm (B),

200µm (C).

1001

1002

1003

1004 **Figure S8: Oocytes in *Hofstenia*'s ovaries are surrounded by follicular cells regardless of position**

1005 **or maturity.** (**A**) Nuclear staining on a sagittal-cut worm confirms the presence of ovaries along the

1006 ventral side of the worm. Consistent with our other imaging, ovaries lack any sense of organization in

1007 regards to maturity along the anterior-posterior axis. (**B-D**) Oocytes are surrounded by a layer of follicular

1008 cells. Scale bars: 500µm (A), 100µm (D).

1009

1010

Figure S9: Worms have spatial preferences during egg-laying that are robust to suboptimal

conditions. (**A**) Quantifying the time from the start of visible muscle contractions to egg laying reveals

that the egg-laying process spans a timescale of seconds to minutes for each egg, with most eggs being

laid within 3 minutes. (**B**) Kernel density estimate of egg locations on the short walls of the worm tanks (n

= 1390 eggs). (**C**) Random observations of worms in their culture tanks (n = 12 tanks) finds only a

minority of worms on the upper half of tank walls, even in conditions where we expect them to be actively

laying eggs. (**D**) Worms preferentially avoid laying eggs on the floor of the tank regardless of whether they

are fed or unfed (two-sample t-test: p= 0.62, n ≥ 8). (**E**) Distribution of clutch sizes in communal tanks (n =

42 clutches) shows that clutches frequently contain more than 10 eggs laid within a 3-4 day window. (**F**)

Distribution of eggs laid by individual worms across different egg laying contexts shows that most worms

lay fewer than 10 eggs in a 3-4 day window (**G**) Worms often add eggs to clutches constructed by other

worms; the distribution shows the numbers of new eggs added to old clutches.

Video 1: The penis sheath is ciliated. Live-imaging of the ventral surface of a worm shows

beating cilia lining the sheath. This set of cilia is distinct from the cilia lining the epidermis of the

worm.

Video 2: The male gonopore is ciliated. Live imaging of the ventral surface of a worm shows

that the male gonopore has beating cilia.

Video 3: Eggs are laid through the mouth. Video of an adult worm laying eggs, filmed from

the ventral surface at 1Hz and sped up 2000x. The worm lowers its mouth and then contracts its

body. During these contractions, a cluster of mature eggs can be seen distending the body. An

individual egg is loaded into and travels through the pharynx eventually emerging through the

mouth.

1043

1044

1045

References

Abalde S, Jondelius U. 2024. A phylogenomic backbone for Acoelomorpha inferred from
transcriptomic data. *Syst Biol*. doi:10.1093/sysbio/syae057
Achatz JG, Chiodin M, Salvenmoser W, Tyler S, Martinez P. 2013. The Acoela: on their kind
and kinships, especially with nemertodermatids and xenoturbellids (Bilateria incertae
sedis). *Org Divers Evol* **13**:267–286.
Achatz JG, Hooge MD. 2006. Convolutidae (Acoela) from Tanzania. *Zootaxa* **1362**:1.
Ahyong S, Boyko CB, Bernot J, Brandão SN, Daly M, De Grave S, de Voogd NJ, Gofas S,
Hernandez F, Hughes L, Neubauer TA, Paulay G, van der Meij S, Boydens B, Decock W,
Dekeyzer S, Goharimanesh M, Vandepitte L, Vanhoorne B, Adlard R, Agatha S, Ahn KJ,
Alonso MV, Alvarez B, Amler MRW, Amorim V, Anderberg A, Andrés-Sánchez S, Ang Y,
Antić D, Antonietto LS, Arango C, Artois T, Atkinson S, Auffenberg K, Bailly N, Baldwin BG,
Bank R, Barber A, Barrett RL, Bartsch I, Bellan-Santini D, Bergh N, Bernard C, Berrios
Ortega FJ, Berta A, Bezerra TN, Bieler R, Blanco S, Blasco-Costa I, Blazewicz M, Błędzki
LA, Bock P, Bonifacino M, Böttger-Schnack R, Bouchet P, Boury-Esnault N, Bouzan R,
Boxshall G, Bray R, Brito Seixas AL, Bruce NL, Bruneau A, Budaeva N, Bueno-Villegas J,
Calvo Casas J, Cárdenas P, Carstens E, Cartwright P, Cedhagen T, Chan BK, Chan TY,
Chernyshev A, Choong H, Christenhusz M, Churchill M, Collins AG, Collins GE, Collins K,
Consorti L, Copilaș-Ciocianu D, Corbari L, Cordeiro R, Costa VM d. M, Costa Corgosinho
PH, Coste M, Costello MJ, Crandall KA, Cremonte F, Cribb T, Cuscó-Borràs J, Cutmore S,
Dahdouh-Guebas F, Daneliya M, Dauvin JC, Davie P, De Broyer C, de Lima Ferreira P, de
Mazancourt V, de Moura Oliveira L, Decker P, Defaye D, Dekker H, Di Capua I, Dippenaar
S, Dohrmann M, Dolan J, Domning D, Downey R, Dreyer N, Eisendle U, Eitel M, Eleaume
M, Enghoff H, Epler J, Esquete Garrote P, Evenhuis NL, Ewers-Saucedo C, Faber M,
Figueroa D, Fišer C, Fordyce E, Foster W, Fransen C, Freire S, Fujimoto S, Furuya H,
Galán-Díaz J, Galbany-Casals M, Gale A, Galea H, Gao T, García-Moro P, Garic R,
Garnett S, Gaviria-Melo S, Gerken S, Gibson D, Gibson R, Gil J, Gittenberger A, Glasby C,
Glenner H, Glover A, Gómez I, Gómez-Noguera SE, Gondim AI, Gonzalez BC, González-
Solís D, Goodwin C, Gostel M, Grabowski M, Gravili C, Grossi M, Guerra-García JM,
Guerrero JM, Guidetti R, Guiry, Gutierrez D, Hadfield KA, Hajdu E, Halanych K, Hallermann
J, Hayward BW, Hegna TA, Heiden G, Hendrycks E, Hennen D, Herbert D, Herrera
Bachiller A, Hodda M, Høeg J, Hoeksema B, Holovachov O, Hooge, Hooper JN, Horton T,
Houart R, Huys R, Hyžný M, Iniesta LFM, Iseto T, Iwataki M, Janssen R, Jaume D,
Jazdzewski K, Jersabek CD, Jiménez-Mejías P, Jóźwiak P, Kabat A, Kajihara H, Kakui K,
Kantor Y, Karanovic I, Karapunar B, Karthick B, Kathirithamby J, Katinas L, Kilian N, Kim
YH, King R, Kirk PM, Klautau M, Kociolek JP, Köhler F, Konowalik K, Kotov A, Kovács Z,
Kremenetskaia A, Kristensen RM, Kroh A, Kulikovskiy M, Kullander S, Kupriyanova E,
Lamaro A, Lambert G, Larridon I, Lazarus D, Le Coze F, Le Roux M, LeCroy S, Leduc D,
Lefkowitz EJ, Lemaitre R, Lichter-Marck IH, Lim SC, Lindsay D, Liu Y, Loeuille B, Lois R,
Lörz AN, Ludwig T, Lundholm N, Macpherson E, Maguilla E, Mah C, Mamos T, Manconi R,
Mapstone G, Marek PE, Markello K, Márquez-Corro JI, Marshall B, Marshall DJ, Martin P,
Martín-Bravo S, Martinez Arbizu P, Maslakova S, McFadden C, McInnes SJ, McKenzie R,
Means J, Mees J, Mejía-Madrid HH, Meland K, Merrin KL, Míguez M, Miller J, Mills C,

Moestrup Ø, Mokievsky V, Molodtsova T, Monniot F, Mooi R, Morales-Alonso A, Morandini
AC, Moreira da Rocha R, Morrow C, Mortelmans J, Müller A, Muñoz Gallego AR, Muñoz
Schüler P, Musco L, Nascimento JB, Nesom G, Neto Silva M d. S, Neubert E, Neuhaus B,
Ng P, Nguyen AD, Nielsen C, Nielsen S, Nishikawa T, Norenburg J, O'Hara T, Opresko D,
Osawa M, Osigus HJ, Ota Y, Páll-Gergely B, Panero JL, Patterson D, Pedram M, Pelser P,
Peña Santiago R, Perbiche-Neves G, Pereira J d. S, Pereira PHM, Pereira SGG, Perez-
Losada M, Petrescu I, Pfingstl T, Piasecki W, Pica D, Picton B, Pignatti J, Pilger JF,
Pinheiro U, Pisera AB, Poatskievick Pierezan B, Polhemus D, Poore GC, Potapova M,
Praxedes RA, Půža V, Read G, Reich M, Reimer JD, Reip H, Resende Bueno V, Reuscher
M, Reynolds JW, Richling I, Rimet F, Ríos P, Rius M, Rodríguez E, Rogers DC, Roque N,
Rosenberg G, Rützler K, Sa IR d. S, Sá HAB, Saavedra M, Sabater LM, Sabbe K, Sabroux
R, Saiz-Salinas J, Sala S, Samimi-Namin K, Sánchez-Villegas M, Sánchez-Villegas R,
Santagata S, Santos S, Santos SG, Santos Filho MAB d., Sanz Arnal M, Sar E, Saucède T,
Schärer L, Schierwater B, Schilling E, Schmidt-Lebuhn A, Schneider S, Schönberg C,
Schrével J, Schuchert P, Schweitzer C, Semple JC, Senna AR, Sennikov A, Serejo C,
Shaik S, Shamsi S, Sharma J, Shear WA, Shenkar N, Short M, Sicinski J, Sierwald P, Silva
DKF d., Silva ESS, Silva MLCN, Simmons E, Sinniger F, Sinou C, Sivell D, Smit H, Smit N,
Smol N, Sørensen MV, Souza-Filho JF, Spelda J, Sterrer W, Steyn HM, Stoev P, Stöhr S,
Suárez-Morales E, Susanna A, Suttle C, Swalla BJ, Taiti S, Tanaka M, Tandberg AH, Tang
D, Tasker M, Taylor J, Taylor J, Taylor K, Tchesunov A, Temereva E, ten Hove H, ter
Poorten JJ, Thirouin K, Thomas JD, Thuesen EV, Thurston M, Thuy B, Timi JT, Todaro A,
Todd J, Turon X, Tyler S, Uetz P, Urbatsch L, Uribe-Palomino J, Urtubey E, Utevsky S,
Vacelet J, Vader W, Väinölä R, Valdés-Florido A, Valls Domedel G, Van de Vijver B, van
Haaren T, van Soest RW, Vanreusel A, Vázquez-García B, Velandia J, Venekey V,
Verhoeff T, Villaverde T, Vinarski M, Vonk R, Vos C, Vouilloud AA, Walker-Smith G, Walter
TC, Watling L, Wayland M, Wesener T, Wetzel CE, Whipps C, White K, Wieneke U,
Williams DM, Williams G, Williams N, Wilson KL, Wilson R, Witkowski J, Wyatt N, Xavier J,
Xu K, Zanol J, Zeidler W, Zhao Z, Zullini A. 2024. World Register of Marine Species.
Åkesson B, Gschwentner R, Hendelberg J, Ladurner P, Müller J, Rieger R. 2001. Fission in
Convolutriloba longifissura: asexual reproduction in acoelous turbellarians revisited. *Acta*
Zool **82**:231–239.
Allard P, Tabin CJ. 2009. Achieving bilateral symmetry during vertebrate limb development.
Semin Cell Dev Biol **20**:479–484.
Apelt G. 1969. Fortpflanzungsbiologie, Entwicklungszyklen und vergleichende Frühentwicklung
acoeler Turbellarien. *Mar Biol* **4**:267–325.
Ax P, Schulz E. 1959. Ungeschlechtliche Fortpflanzung durch Paratomie bei acoelen
Turbellarien. *Biol Zent Bl* **78**:613–622.
Bailly X, Laguerre L, Correc G, Dupont S, Kurth T, Pfannkuchen A, Entzeroth R, Probert I,
Vinogradov S, Lechauve C, Garet-Delmas M-J, Reichert H, Hartenstein V. 2014. The
chimerical and multifaceted marine acoel Symsagittifera roscoffensis: from photosymbiosis
to brain regeneration. *Front Microbiol* **5**. doi:10.3389/fmicb.2014.00498
Ballarin L, Burighel P, Cima F. 2008. A tale of death and life: natural apoptosis in the colonial
ascidian Botryllus schlosseri (Urochordata, Ascidiacea). *Curr Pharm Des* **14**:138–147.
Beltagi S, Mandura A. 1991. Hofstenia arabiensis Nov. Sp. (Hofsteniidae): A New Species of
Acoelan Turbellaria from the Red Sea North of Jeddah. *Journal of King Abdulaziz*
University-science **3**:65–90.
Bock S. 1923. Eine neue marine Turbellariengattung aus Japan. Almqvist & Wiksell.
Boone M, Houthoofd W, Bert W, Artois T. 2011. First record of Nemertodermatida from Belgian
marine waters. *Belg J Zool* **141**:62–64.
Boulan L, Léopold P. 2021. What determines organ size during development and regeneration?
Development **148**:dev196063.

Brand JN, Harmon LJ, Schärer L. 2022a. Frequent origins of traumatic insemination involve
convergent shifts in sperm and genital morphology. *Evolution Letters* **6**:63–82.
Brand JN, Harmon LJ, Schärer L. 2022b. Mating behavior and reproductive morphology predict
macroevolution of sex allocation in hermaphroditic flatworms. *BMC Biol* **20**:35.
Brand JN, Viktorin G, Wiberg RAW, Beisel C, Schärer L. 2022c. Large-scale phylogenomics of
the genus Macrostomum (Platyhelminthes) reveals cryptic diversity and novel sexual traits.
Mol Phylogenet Evol **166**:107296.
Bush L. 1975. Biology of Neochilda fusca N. gen., n. sp. from the northeastern coast of the
United States (Platyhelminthes: Turbellaria). *Biol Bull* **148**:35–48.
Cannon JT, Vellutini BC, Smith J, Ronquist F, Jondelius U, Hejnol A. 2016. Xenacoelomorpha is
the sister group to Nephrozoa. *Nature* **530**:89–93.
Charnov EL. 1979. Simultaneous hermaphroditism and sexual selection. *Proc Natl Acad Sci U*
S A **76**:2480–2484.
Chiodin M, Børve A, Berezikov E, Ladurner P, Martinez P, Hejnol A. 2013. Mesodermal gene
expression in the acoel Isodiametra pulchra indicates a low number of mesodermal cell
types and the endomesodermal origin of the gonads. *PLoS One* **8**:e55499.
Corrêa DD. 1960. Two New Marine Turbellaria from Florida. *Bull Mar Sci* **10**:208–216.
Costello HM, Costello DP. 1939. EGG LAYING IN THE ACOELOUS TURBELLARIAN
POLYCHOERUS CARMELENSIS. *Biol Bull* **76**:80–89.
Costello HM, Costello DP. 1938. COPULATION IN THE ACŒLOUS TURBELLARIAN
POLYCHŒRUS CARMELENSIS. *Biol Bull* **75**:85–98.
Crezee M. 1978. Paratomella rubra Rieger and Ott, an amphiatlantic acoel turbellarian. *Cah Biol*
Mar **19**:1–9.
Crezée M. 1975. Monograph of the solenofilomorphidae (Turbellaria: Acoela). *Int Rev Gesamt*
Hydrobiol **60**:769–845.
Davies N, Krebs J, West S. 2013. An Introduction to Behavioural Ecology. Wiley-Blackwell.
De Mulder K, Kuales G, Pfister D, Willems M, Egger B, Salvenmoser W, Thaler M, Gorny A-K,
Hrouda M, Borgonie G, Ladurner P. 2009. Characterization of the stem cell system of the
acoel Isodiametra pulchra. *BMC Dev Biol* **9**:69.
Dörjes J. 1968. Die Acoela (Turbellaria) der Deutschen Nordseeküste und ein neues System
der Ordnung. *J Zoolog Syst Evol Res* **6**:56–452.
Dörjes J. 1966. Paratomella unichaeta, nov. gen. nov. spec., Vertreter einen neuen Familie der
Turbellaria Acoela mit asexueller Fortpflanzung durch Paratomie. *Veröffentlichungen des*
Instituts für Meeresforschung in Bremerhaven **II**:187–200.
du Bois-Reymond Marcus E. 1955. On Turbellaria and Polygordius from the Brazilian Coast. *Bol*
Fac Fil Ci Letr Univ Sao Paulo Zool **20**:19–53.
Eckelbarger K, Hodgson A. 2021. Invertebrate oogenesis – a review and synthesis: comparative
ovarian morphology, accessory cell function and the origins of yolk precursors. *Invertebr*
Reprod Dev **65**:71–140.
Falleni A, Gremigni V. 1990. Ultrastructural study of oogenesis in the acoel turbellarian
Convoluta. *Tissue Cell* **22**:301–310.
Faubel A. 1983. The Polycladida, Turbellaria. Proposal and establishment of a new system. Part
I. The Acotylea. *Mitt Hamb Zool Mus Inst* **80**:17–121.
Faubel A. 1976. Interstitielle Acoela (Turbellaria) aus dem Littoral der nordfriesischen Inseln Sylt
und Amrum. *Mitt Hamb Zool Mus Inst* **73**:17–56.
Faubel A. 1974. Die Acoela (Turbellaria) eines Sandstrandes der Nordseeinsel Sylt. *Mikrofauna*
Meeresbodens **32**:1–58.
Faubel A, Cameron B. 2001. Platyhelminthes from salt marshes of Coomera River,
Southeastern Queensland, Australia. *Mem Queensland Mus* **46**:511–519.
Gardiner E. 1898. The Growth of the Ovum, Formation of the Polar Bodies, and the Fertilization
in Polychaerus Caudatus.

Gardiner E. 1895. Early development of Polychoerus Caudatus, Mark. *J Morphol* **11**:155–176.

Gehrke AR, Neverett E, Luo Y-J, Brandt A, Ricci L, Hulett RE, Gompers A, Ruby JG, Rokhsar DS, Reddien PW, Srivastava M. 2019. Acoel genome reveals the regulatory landscape of whole-body regeneration. *Science* **363**:eaau6173.

Giribet G, Edgecombe GD. 2020. The Invertebrate Tree of Life. Princeton University Press.

Grae D, Kozloff E. 1999. Post-copulatory storage of sperm in the seminal bursa of the acoel flatworm Otocelis luteola. *Invertebr Reprod Dev* **35**:151–154.

Grossman H, Har-Paz E, Gindi N, Levi M, Miller I, Nevo N, Galiani D, Dekel N, Shalgi R. 2017. Regulation of GVBD in mouse oocytes by miR-125a-3p and Fyn kinase through modulation of actin filaments. *Sci Rep* **7**:2238.

Hanson ED. 1960. Asexual Reproduction in Acoelous Turbellaria. *Yale J Biol Med* **33**:107–111.

Harris MP, Daane JM, Lanni J. 2021. Through veiled mirrors: Fish fins giving insight into size regulation. *Wiley Interdiscip Rev Dev Biol* **10**:e381.

Herlands L, Maul GG. 1994. Characterization of a major nucleoplasmin-like germinal vesicle protein which is rapidly phosphorylated before germinal vesicle breakdown in Spisula solidissima. *Dev Biol* **161**:530–537.

Hooge M. 2003. Two new families, three new genera, and five new species of acoels (Acoela Platyhelminthes) from Queensland Australia. *Cah Biol Mar* **44**:275–298.

Hooge M, Smith JPS. 2004. New acoels (Acoela, Acoelomorpha) from North Carolina. *Zootaxa* **442**:1–24.

Hooge M, Wallberg A, Todt C, Maloy A, Jondelius U, Tyler S. 2007. A revision of the systematics of panther worms (Hofstenia spp., Acoela), with notes on color variation and genetic variation within the genus. *Hydrobiologia* **592**:439–454.

Hookabe N, Jimi N, Furushima Y, Fujiwara Y. 2024. Discovery of deep-sea acoels from a chemosynthesis-based ecosystem. *Biol Lett* **20**:20230573.

Hulett RE, Kimura JO, Bolaños DM, Luo Y-J, Rivera-López C, Ricci L, Srivastava M. 2023. Acoel single-cell atlas reveals expression dynamics and heterogeneity of adult pluripotent stem cells. *Nat Commun* **14**:2612.

Hulett RE, Potter D, Srivastava M. 2020. Neural architecture and regeneration in the acoel Hofstenia miamia. *Proceedings of the Royal Society B: Biological Sciences* **287**:20201198.

Hyman LH. 1951. The invertebrates: Platyhelminthes and Rhynchocoela. The acoelomate Bilateria. McGraw-Hill, New York.

Hyman LH. 1937. Reproductive system and copulation in amphiscolops langerhansi (turbellaria acoela). *Biol Bull* **72**:319–326.

Ishikawa K, Yamasu T. 1992. An acoel flat worm species related closely to the species Convolutriloba retrogemma Hendelberg and Akesson occurs in Okinawa Island, Ryukyu Archipelago. *Zoological Science* **9**:1281.

Jondelius U, Wallberg A, Hooge M, Raikova OI. 2011. How the Worm Got its Pharynx: Phylogeny, Classification and Bayesian Assessment of Character Evolution in Acoela. *Syst Biol* **60**:845–871.

Kapli P, Telford MJ. 2020. Topology-dependent asymmetry in systematic errors affects phylogenetic placement of Ctenophora and Xenacoelomorpha. *Sci Adv* **6**:eabc5162.

Kimura JO, Bolaños DM, Ricci L, Srivastava M. 2022. Embryonic origins of adult pluripotent stem cells. *Cell* **185**:4756–4769.e13.

Kimura JO, Ricci L, Srivastava M. 2021. Embryonic development in the acoel Hofstenia miamia. *Development* **148**:dev188656.

Ko JM, Reginato W, Wolff A, Lobo D. 2024. Mechanistic regulation of planarian shape during growth and degrowth. *Development* **151**:dev202353.

Kostenko AG. 1989. The position of Taurida fulvomaculata (Ax, 1959) in the system of Acoela. In: Morphology of turbellarians. *Akad Nauk USSR Trudy Zool* **195**:14–25.

Kozloff EN. 2000a. Differentiation, dedifferentiation, and redifferentiation of reproductive

structures of the acoel flatworm*Otocelis luteola*, and notes on longevity of this species.
Invertebr Reprod Dev **37**:95–106.
Kozloff EN. 2000b. A new genus and five new species of acoel flatworms from the Pacific coast
of North America, and resolution of some systematic problems in the families Convolutidae
and Otocelididae. *Cah Biol Mar* **41**:281–293.
Liu J-L, Buszczak M, Gall JG. 2006. Nuclear bodies in the Drosophila germinal vesicle.
Chromosome Res **14**:465–475.
Mamkaev YV. 1965. Etude morphologique d'Actinoposthia beklemischevi n. sp. (Turbellaria
Acoela). *Cah Biol Mar* **6**:23–50.
Meyer-Wachsmuth I, Curini Galletti M, Jondelius U. 2014. Hyper-cryptic marine meiofauna:
species complexes in Nemertodermatida. *PLoS One* **9**:e107688.
Michiels NK, Newman LJ. 1998. Sex and violence in hermaphrodites. *Nature* **391**:647–647.
Munro C, Cadis H, Pagnotta S, Houliston E, Huynh J-R. 2023. Conserved meiotic mechanisms
in the cnidarian *Clytia hemisphaerica* revealed by *Spo11* knockout. *Sci Adv* **9**:eadd2873.
Peebles F. 1915. A Description of three Acoela from the Gulf of Naples. *Mitteilungen a.d Zool
Station zu Neapel* **22**:292–312.
Pellettieri J, Fitzgerald P, Watanabe S, Mancuso J, Green DR, Sánchez Alvarado A. 2010. Cell
death and tissue remodeling in planarian regeneration. *Dev Biol* **338**:76–85.
Perea-Atienza E, Botta M, Salvenmoser W, Gschwentner R, Egger B, Kristof A, Martinez P,
Achatz JG. 2013. Posterior regeneration in Isodiametra pulchra (Acoela, Acoelomorpha).
Front Zool **10**:64.
Petrov A, Hooge M, Tyler S. 2004. Ultrastructure of sperms in Acoela (Acoelomorpha) and its
concordance with molecular systematics. *Invertebr Biol* **123**:183–197.
Philippe H, Brinkmann H, Copley RR, Moroz LL, Nakano H, Poustka AJ, Wallberg A, Peterson
KJ, Telford MJ. 2011. Acoelomorph flatworms are deuterostomes related to Xenoturbella.
Nature **470**:255–258.
Raikova O, Falleni A, Gremigni V. 1995. Oogenesis in Actinoposthia beklemischevi
(Platyhelminthes, Acoela): an ultrastructural and cytochemical study. *Tissue Cell* **27**:621–
633.
Ramm SA, Schlatter A, Poirier M, Schärer L. 2015. Hypodermic self-insemination as a
reproductive assurance strategy. *Proceedings of the Royal Society B: Biological Sciences*
282:20150660.
Reddien PW. 2018. The cellular and molecular basis for planarian regeneration. *Cell* **175**:327–
345.
Ricci L, Srivastava M. 2021. Transgenesis in the acoel worm Hofstenia miamia. *Dev Cell*
56:3160–3170.e4.
Rieger RM, Tyler S, Smith JPS, Rieger GE. 1991. Platyhelminthes and NemertineaMicroscopic
Anatomy of Invertebrates. New York: Wiley-Liss. pp. 7–140.
Riser NW. 1987. Nemertinoides elongatus gen. n., sp. n. (Turbellaria: Nemertodermatida) from
coarse sand beaches of the Western North Atlantic. *Proc Helminthol Soc Wash* **54**:60–67.
Rychel AL, Swalla BJ. 2008. Anterior regeneration in the hemichordate Ptychodera flava. *Dev
Dyn* **237**:3222–3232.
Sagata N. 1996. Meiotic metaphase arrest in animal oocytes: its mechanisms and biological
significance. *Trends Cell Biol* **6**:22–28.
Schärer L, Janicke T, Ramm SA. 2014. Sexual conflict in hermaphrodites. *Cold Spring Harb
Perspect Biol* **7**:a017673.
Schärer L, Littlewood DTJ, Waeschenbach A, Yoshida W, Vizoso DB. 2011. Mating behavior
and the evolution of sperm design. *Proceedings of the National Academy of Sciences*
108:1490–1495.
Schmidt-Rhaesa A. 2007. The Evolution of Organ Systems. OUP Oxford.
Shannon T, Achatz J. 2007. Convolutriloba macropyga sp. nov., an uncommonly fecund acoel

(Acoelomorpha) discovered in tropical aquaria. *Zootaxa* **1525**:1–17.

Srivastava M. 2022. Studying development, regeneration, stem cells, and more in the acoel
Hofstenia miamia. *Curr Top Dev Biol* **147**:153–172.

Srivastava M, Mazza-Curll KL, van Wolfswinkel JC, Reddien PW. 2014. Whole-Body Acoel
Regeneration Is Controlled by Wnt and Bmp-Admp Signaling. *Curr Biol* **24**:1107–1113.

Steinböck O. 1966. Grundsätzliches zur Evolution der Turbellarien. *Zeitschrift für Zoologische*
Systematik und Evolutionsforschung **4**:58–195.

Sterrer WE. 1998. New and known Nemertodermatida (Platyhelminthes- Acoelomorpha) - a
revision. *Belgian Journal of Zoology* **128**:55–92.

Tong SJW, Ong RSL. 2020. Mating behavior, spawning, parental care, and embryonic
development of some marine pseudocerotid flatworms (Platyhelminthes: Rhabditophora:
Polycladida) in Singapore. *Invertebr Biol* **139**:e12293.

Trible W, Kronauer DJC. 2017. Caste development and evolution in ants: it's all about size. *J*
Exp Biol **220**:53–62.

Tyler, S., Schilling, S., Hooge, M., & Bush, L. F. 2012. Turbellarian taxonomic database. Version
1.7. http://turbellaria.umaine.edu

Uppaluri S, Brangwynne CP. 2015. A size threshold governs Caenorhabditis elegans
developmental progression. *Proc R Soc B* **282**:20151283.

Vallejo DM, Juarez-Carreño S, Bolivar J, Morante J, Dominguez M. 2015. A brain circuit that
synchronizes growth and maturation revealed through Dilp8 binding to Lgr3. *Science*
350:aac6767.

Wallberg A. 2012. The Stylet - Diversity and Systematics of Acoela and Nemertodermatida.
http://acoela.myspecies.info

Watzin MC. 1984. Egg Laying in the Acoel Turbellarian Neochildia fusca. *Trans Am Microsc Soc*
103:322–325.

Wolpert L. 2010. Arms and the man: the problem of symmetric growth. *PLoS Biol* **8**:e1000477.

Zabotin YI, Evtugyn VG. 2021. Ultrastructure of spermatozoa and female copulatory organs in
preferably asexually-reproducing acoel Convolutriloba retrogemma (Acoelomorpha).
Zoomorphology **140**:19–26.

Zauchner T, Salvenmoser W, Egger B. 2015. A cultivable acoel species from the
Mediterranean, Aphanostoma pisae sp. nov. (Acoela, Acoelomorpha). *Zootaxa* **3941**:401–
413.

1325