Reproductive life history of an acoel worm

Vikram Chandra[#][*], Samantha E. Tseng[#], Allison P. Kann, D. Marcela Bolaños, Mansi

Srivastava[*]

Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology, Harvard

University, Cambridge, MA 02138

# These authors contributed equally

* Corresponding authors. Emails: vchandra1@fas.harvard.edu, mansi@oeb.harvard.edu

Abstract

Acoel worms belong to an enigmatic and understudied animal lineage (Phylum

Xenacoelomorpha). Sparse taxonomic and histological work suggests that these worms exhibit

a diversity of reproductive anatomies and likely also a corresponding diversity in reproductive

behavior. However, few acoels have been studied in detail. Here, we study the reproductive life

history of the three-banded panther worm *Hofstenia miamia*, an acoel that is emerging as a new

lab-tractable model system for regenerative biology. Using molecular labeling and histology, we

describe *H. miamia*'s reproductive organs, identifying structures previously unknown in acoels.

Following a cohort of worms from zygote to adulthood, we quantify the developmental dynamics

of their reproductive organs, and find that these organs are assembled step-wise, at increasing

body sizes. Studying the dynamics of organ growth during regeneration and de-growth in

starving adult worms, we show that reproductive organs follow similar growth rules in these

contexts, suggesting that their development and maintenance is regulated by a size-dependent

program. Finally, we study egg-laying behavior, finding that *H. miamia* lay their eggs through

their mouths after loading them into their pharynges. Experimentally varying the worms'

environment reveals that the worms make active decisions about when and where to lay their

eggs. In summary, our work establishes foundational knowledge to enable the experimental

study of reproductive anatomy, physiology, and behavior in acoels.

Introduction

Reproduction is critical for evolutionary success, and animals display great diversity in their

reproductive life histories. The structures and strategies employed for reproduction have been

relatively well-studied in some animal lineages, such as arthropods and vertebrates (1).

However, the complexities of reproductive biology in many animal phyla remain poorly studied.

For example, Phylum Xenacoelomorpha, which includes acoels, nemertodermatids, and

xenoturbellids, represents a deep-diverging lineage of animals with bilateral symmetry (2–4)

(Fig. 1A). Whereas reproductive structures have been reported in a handful of species in this

phylum of largely marine worms, many aspects of reproduction such as sexual maturation,

aging, mating, and egg laying are unexplored (Fig. 1B). Here, we apply a diverse toolkit for a

systematic study of reproductive biology in the acoel *Hofstenia miamia*, which has emerged as a

new laboratory research organism.

There are roughly 400 described species of acoel worms (5–7). Traditionally, acoel

taxonomy is done through the description and analysis of morphological characters observable

in histology slice preparations of specimens collected from the field (8). A few exceptions

notwithstanding, these data also constitute the majority of what we know about reproduction in

acoels. Acoels are thought to be simultaneous hermaphrodites (with few exceptions (9, 10));

once mature, each worm has both male and female reproductive structures (8). Unlike many

other aquatic animals, e.g., cnidarians, poriferans, echinoderms, that reproduce sexually

through spawning and external fertilization (11), acoels have been reported to reproduce

sexually through internal fertilization (6). Acoel worms exhibit a large diversity in their

reproductive anatomy (Fig. 1B, Table S1). This anatomical variation likely predicts variation in

reproductive physiology and behavior, but little is known about how most acoels develop their

reproductive organs, mate, or lay eggs. For instance, species lacking a female gonopore and a

seminal bursa (a sperm storage organ) likely mate through competitive hypodermic

insemination, but mating behavior has not been observed in the majority of these species. The

lack of a female gonopore in these species predicts that eggs must be laid either through

rupture of the body wall (12–14) or through the mouth, but few species have been observed

laying eggs at all, and despite speculation, no animal has ever been shown to lay eggs through

its mouth.

The majority of observations of reproductive behavior in acoels are based on a small

handful of species within one family (the Convolutidae) that has several derived traits (8).

Historically, it has been challenging to study acoel reproduction in controlled conditions because

few acoels have been cultured in the lab (12, 15–17), and most descriptions of acoel

reproduction occurred at a time when modern imaging methods did not exist or were

inaccessible. To improve our understanding of acoel life histories, we studied reproduction in an

early-diverging acoel, the three-banded panther worm *Hofstenia miamia* (Fig. 1C)*. H. miamia* is

an emerging model system with several features that allow us to study it in controlled

conditions: *H. miamia*'s life cycle can be closed in the lab (rearing worms from embryo to gravid

adult takes 2-3 months), and a growing array of experimental tools is available to observe and

manipulate the worm (18–23).

H. miamia is a member of the family Hofsteniidae, which contains three genera (5, 24)

and is placed as an early-diverging lineage among acoels (8). Five species have been

described within the genus *Hofstenia*. Histology-based work has described the coarse anatomy

of reproductive organs in *Hofstenia* (24–28). However, the fine structure of these organs, how

they grow and develop, and how they function in the course of reproductive behavior remains

unknown. Here, to understand the reproductive life history of *H. miamia*, we use a combination

of light and confocal microscopy, fluorescence in situ hybridization, immunofluorescence, and

histology, as well as observations of reproductive development and behavior in controlled

conditions. We describe the fine structure of *H. miamia*'s reproductive organs, its developmental

and regenerative trajectory, and how this worm lays eggs. These findings establish a foundation

for the mechanistic study of reproduction in an acoel worm.

Results

An overview of *Hofstenia miamia*'s reproductive anatomy

The genus *Hofstenia* was first described by Bock (25) and based on studies of the three

currently-recognized species, *H. atroviridis*, *H. miamia*, and *H. arabiensis*, *Hofstenia* are

considered to be simultaneous hermaphrodites (24–28). The three species are also described

as having an anterior male reproductive system including a seminal vesicle, prostatic vesicle, a

penis with sclerotized needles, diffuse ventrolateral testes, and a male gonopore near the mouth

(24–28). The female reproductive system, as described, includes two ovaries spanning the

posterior two-thirds of the worm. *Hofstenia miamia* is likely the most common and widespread

species within the genus, with a geographic distribution including the North Atlantic, the

Caribbean, the Bahamas, and the Florida coast (24–28). Although multiple authors have

described elements of *H. miamia*'s reproductive systems, these descriptions were all based on

worms roughly 5-8mm in length (24–28). In our lab cultures of *H. miamia*, fully grown adult

worms can be over 1.4cm in length, and we reasoned that their anatomy could be subtly

different. We therefore began by re-examining the coarse anatomy of *Hofstenia miamia*.

We anesthetized adult worms, mounted them ventral side up on a slide, and imaged

them through a dissecting microscope to reveal coarse reproductive organization. As expected,

we found that *H. miamia*'s male copulatory structures are located in the anterior of the animal,

just posterior to the mouth (Fig. 1D,E). These male structures are located within a translucent

tube-like region, located close to the ventral surface of the worm, immediately ventral to the

pharynx. We observed two white opaque regions within the male copulatory apparatus, located

towards its posterior (Fig. S1A). The larger, posterior oval structure corresponds to the seminal

vesicle, while the smaller, teardrop-shaped anterior structure corresponds to the prostatic

vesicle (Fig. 1D,E; Fig. S1A), traditionally referred to as the vesicula granulorum (24–28).

Previous work reported that *H. miamia* has a penis or stylet: a sperm-delivery organ containing

rigid needles. This structure is typically difficult to see in these imaging conditions. However, we

identified a small aperture immediately posterior and ventral to the mouth (Fig. S1A), and we

observed that the worm extended its penis through it, suggesting that this aperture is the male

gonopore, and confirming the location of the penis in the anterior of the worm.

As previously described (24–28), we found that the female reproductive system occupies

the posterior two-thirds of the worm. This system appears relatively simple, with numerous eggs

grouped into three distinct clusters (Fig. S1B). One lateral group on each side of the body, near

the margins, extends along the anterior-posterior axis. These lateral clusters contain oocytes of

various sizes, likely corresponding to different stages of maturation, each with a visible nucleus,

and have been identified as the ovaries in historical reports (24–28). There is no obvious

organization of oocytes in the ovaries along the anterior-posterior, dorso-ventral, or medio-

lateral axes.

Posterior to the pharynx, there is often a cluster of spherical eggs (Fig. 1D,E; Fig. S1C).

The number of eggs at this position is highly variable, ranging from zero to over a dozen eggs.

In *H. atroviridis*, Bock (25) speculated that these eggs are mature, fertilized, and ready to be

laid. This is consistent with our observations: these eggs appeared uniformly sized and lacked a

visible nucleus, in contrast to the oocytes in the ovaries, suggesting that eggs in this area may

already be fertilized. To test whether these eggs were indeed zygotes, we dissected out eggs

from this central cluster, as well as from the ovaries. We found that 46/56 eggs removed from

the central cluster developed and hatched into juvenile worms, in line with previously

established embryonic trajectories (Fig. S1D) (23). Meanwhile, all eggs dissected from the

ovaries disintegrated within 48 hours. The eggs in this central cluster are therefore mature, likely

fertilized, and ready for laying while eggs in the ovaries are immature or unfertilized. Overall,

these observations confirm and extend previous descriptions of *H. miamia*, and provide a high-

level overview of *H. miamia*'s reproductive anatomy. Next, we sought to apply a variety of

experimental approaches to obtain a detailed understanding of these structures and their

development.

Figure 1: *Hofstenia miamia* is a model acoel worm for studying Xenacoelomorph life history. (**A**)

Xenacoelomorpha is a deep-diverging bilaterian lineage of aquatic worms. Phylogeny modified from (18).

Animal icons are from phylopic.org, and are in the public domain. (**B**) Family-level phylogeny of

Acoelomorpha (Nemertodermatida, Acoela) showing anatomical and reproductive life history traits (Table

S1): position of the mouth, whether gonads are mixed or separated by sex, penis type, paired or unpaired

testes, paired or unpaired ovaries, ovary organization, presence or absence of a female gonopore,

presence or absence of a seminal bursa, the number of associated bursal nozzles, egg-laying mode,

mode of sexual reproduction, and mode of asexual reproduction (see Table S2 for definitions of terms

and categories). Schematic for each family shows reproductive anatomy of a representative species

(Table S1). White boxes represent unknown phenotypic states, and in the case of asexual reproduction,

its possible absence. Phenotypic classifications are from (10, 12–15, 24, 25, 27–51) (**C**) Dorsal view of

Hofstenia miamia. (**D**) Ventral view of an adult *Hofstenia miamia*; most reproductive structures are visible

in this view. (**E**) Schematized view of the ventral surface of a worm with known reproductive structures

illustrated. Scale bars: 1 mm (C, D).

Development of reproductive organs

How reproductive structures develop as animals reach sexual maturity is not well understood in

any xenacoelomorph species. In *H. miamia*, juvenile worms that hatch from eggs have no

reproductive structures, and previous observations suggested that these develop over a month

later as the worms grow toward adulthood (18, 23). Our initial observations (Fig. 1D) showed

that we could identify major reproductive structures in a non-invasive manner, simply by imaging

the ventral surfaces of anesthetized worms using a stereo microscope. Using this method in

combination with our ability to study all stages of the life cycle in *H. miamia*, we sought to

determine the timeline and dynamics of reproductive development. We reared 42 zygotes in

isolation and monitored their growth through development (Fig. 2A), precisely controlling their

rearing environment such that each animal had access to a standardized amount of seawater

and a defined amount of food (see Methods). To determine how nutrition affects reproductive

development, we split the cohort of zygotes into two groups that were fed differently. For each

group, we gradually increased the amount of food they were given as they grew larger, to

ensure that worms had enough food to minimize developmental time. These food increases

occurred periodically, but were delayed by two weeks for one feeding group (Fig. S2A). Twice

weekly, we captured images of the ventral surface of each worm and quantified the size of each

visible reproductive structure. Two embryos did not hatch, and two worms died during the

course of the experiment (likely due to handling error). Four worms were removed from the

experiment at different timepoints for additional imaging, resulting in 34 worms that we followed

into adulthood (Fig. S2B).

We found that worms maintained their aspect ratio from hatchling to adult, growing

proportionally over time (Fig. 2A-D; Fig. S2C). At every stage of development, the size of each

reproductive organ scaled with body size once that organ appeared (Fig. 2E-G). Worms with

delayed increases in feeding displayed corresponding delays in their growth, both in their body

size (Fig. S2D) and in the timing of appearance of their ovary and seminal vesicle (Fig. 2H,I).

Worms with delayed feeding developed a penis and a seminal vesicle at a slightly smaller body

size, but did not significantly differ in the body size at which their ovaries first appeared. Overall,

these data suggest that the growth and differentiation of reproductive structures is generally

coupled to the worm's body size.

For both treatments, we found that reproductive development generally occurs in a

stepwise fashion. Newly hatched animals possess all major somatic organs (such as a brain,

muscle system, pharynx, gut, etc. (21, 23)), but lack all reproductive structures (Fig. 2A,B). 4-6

weeks after being laid, when the worm is about 3.02 ± 0.6 mm (mean $\pm$ SD) in length, the

anterior-ventral tissue housing the future male copulatory apparatus becomes visibly less

dense, and a groove becomes apparent in the ventral surface. Soon after, the male gonopore

becomes detectable posterior to the mouth. Sperm begin to accumulate in the seminal vesicle

around 35 days post laying (dpl), once a worm is about 4.41 ± 0.9 mm (mean $\pm$ SD) long.

Shortly after, when a worm is 5.6 ± 0.9 mm (mean $\pm$ SD), the first oocytes become visible in

nascent ovaries. These ovaries grow with the worm over time, with existing oocytes maturing

and new ones continuously added. Around two months after being laid, typically after a worm is

8.73 ± 1.2 mm in length, eggs appear near the base of the pharynx (Fig. 2A,B,H,I). These eggs

are laid within hours or days, and they develop and hatch into juvenile worms. Given that these

worms were isolated since birth, this suggests that *H. miamia* can reproduce through both

mating and self-fertilization.

To visualize the extent to which reproductive structures develop in a sequence, we

ranked each structure in each worm by its date of appearance (Fig. 2J). All worms, without

exception, first developed a penis. A few worms simultaneously developed a seminal vesicle; in

most other worms, this structure appeared second, with a few developing their seminal vesicle

concurrent with nascent ovaries. Ovaries always appeared before the first fertilized eggs are

seen in the central region. Next, we asked whether the left and right ovaries of each worm

developed in synchrony. We found that on the first day that an ovary was visible, 34/34 worms

had both ovaries visible. Partial regression analysis revealed that the sizes of the two ovaries

within a worm are strongly correlated, even when correcting for body size (Fig. S2E). This

suggests that an active developmental process synchronizes the ovaries within individual

worms. Taken together, these observations suggest that the development of reproductive

organs is strongly coupled to body size.

Figure 2: Reproductive organs develop stepwise following scaling rules. (A) Time course of one

worm through developmental time from hatchling to reproductively mature adult. (B) Schematic of

timecourse shown in (A) with key reproductive developments illustrated. (C) The length of worms

increases over time (R^2 = 0.91, p < 0.001), and worms grow proportionally: their length scales with their

width (R^2 = 0.85, p < 0.0001) (D).The length of each reproductive organ scales with increases in body

size (penis: R^2 = 0.85, p < 0.0001; seminal vesicle: R^2 = 0.82, p < 0.0001; ovaries: R^2 = 0.82, p < 0.0001)

(E-G). Comparing the date of first appearance of reproductive organs between worms fed differently

reveals that these organs develop as a function of body size, but not age. (H) Worms with delayed

feeding increases had significant delays in the appearance of their seminal vesicle and ovaries, but not

the penis (Welch's t test for date of appearance for penis: p = 0.08, seminal vesicle: p = 0.04, ovary: p <

10^{-6}; n ≥ 15). (I) Worms with delayed feeding increases had a smaller body length when a penis and

seminal vesicle appeared, but not when ovaries appeared (Welch's t test for length on date of

appearance for penis: p = 0.005, seminal vesicle: p = 0.03, ovary: p = 0.74; n ≥ 15). Asterisks indicate

statistical significance. (**J**) Ranking the order in which reproductive organs appear (y-axis) in developing

worms reveals a stepwise pattern of reproductive differentiation. The x-axis shows individual worms. dpl =

days post laying. Scale bars: 1 mm.

De-growth and regeneration of reproductive organs

Next, we asked whether these patterns of growth persisted in different contexts. Anecdotal

observations suggested *H. miamia* tolerates long periods of starvation but gradually shrinks in

size when deprived of food. We starved a cohort of adult worms and quantified the de-growth of

their reproductive organs over time (Fig. 3A,B). We found that all worms survived over three

months of continuous starvation, gradually shrinking over time (Fig. 3C). Worms maintained

their aspect ratios as they shrank (Fig. 3A,B,D), and their reproductive organs shrank

correspondingly (Fig. S3A-E). Partial regression analysis of shrinking ovaries showed that they

shrink synchronously (Fig. S3F), suggesting active coordination of de-growth.

We also asked how reproductive organs change in another growth context. *H. miamia*,

like many acoels, is an excellent regenerator, and can regenerate all tissues from a wide range

of initial tissue configurations (18, 52). We amputated and followed adult worms in three ways:

by separating the tip of the tail (which contained no reproductive structures) and studying how

they regenerated the majority of their organs, by cutting worms sagittally and studying how they

regenerated their missing half, and by separating heads from tails and studying how these head

fragments regenerated their tails and the posterior region of their ovaries (Fig. 3E). Head and

tail fragments both gradually increased in size, with tail fragments growing much faster (Fig.

S3G-M). Within two weeks, most of these fragments had regained the characteristic shape of

intact homeostatic worms, and were foraging apparently normally. Sagittally-cut fragments first

shrank in size before subsequently growing (Fig. S3M), and appeared to take longer to fully

regain normal worm-like appearance (Fig. S3I,J). All reproductive organs gradually regenerate,

with growth dynamics contingent on the nature of the injury (Fig. 3F-N; Fig. S3G-L,N,O).

The scaling relationships we found between reproductive organs and body size during

post-embryonic development and juvenile growth generalized to both de-growth and

regeneration (Fig. 3O-Q). We then studied the smallest body size at which starving worms

retained their organs, as well as the body size at which reproductive organs first appeared in

regenerating fragments, comparing them to the sizes at which these organs first appear during

the normal developmental transition from juvenile to adult (Fig. 3R). In general, starving worms

lost reproductive organs in the opposite order to which they gained them, first losing fertilized

eggs, then ovaries, and eventually losing their seminal vesicle (Fig. 3S). However, after three

months of starvation, most worms continued to retain their penes despite being below the

threshold size at which we would expect to see loss of this organ (Fig. 3R). In regenerating tail

fragments, the male reproductive structures generally developed before the female reproductive

structures, following similar stepwise patterns to developing juvenile worms (Fig. 3T).

In regenerating head fragments and in sagittally-cut worms, each worm fragment

retained one of its two ovaries immediately after amputation. These ovaries first partially

degenerated before growing back (Fig. 3J), suggesting that there is active destruction of tissue

during the early stages of regeneration. This is consistent with the loss of germline marker

expression during early regeneration (19). The missing ovary began to grow back roughly one

month after amputation. We found that the ovaries grew asymmetrically in these worms, with

the new ovary growing faster (Fig. 3J). This suggests the existence of an active growth

mechanism to ensure that both ovaries reach a symmetric target size.

Together, our results suggest the existence of a size-dependent differentiation program

that regulates the development of reproductive organs, as well as active tissue growth and

destruction mechanisms to ensure organ symmetry. It is likely that these processes regulate

reproductive organ growth in a variety of different developmental contexts, including the

transition from juvenile to adult worm, regeneration after injury and tissue loss, and during

starvation-induced de-growth.

Figure 3: Reproductive organ development follows similar rules in different growth contexts.

(**A**) Time course of a starving worm undergoing de-growth and step-wise loss of reproductive organs. (**B**)

Schematic of reproductive organ degradation over the course of starvation-induced de-growth. (**C**) Worm

length decreases over the course of degrowth (R^2 = 0.85, p < 0.0001). (**D**) Worms shrink as they grow;

their lengths and widths decrease proportionally (R^2 = 0.73, p < 0.0001). dps = days post onset of

starvation. (**E**) Schematic of regeneration of the penis, seminal vesicle, and ovaries following three

different amputations. (**F-N**) Growth dynamics of reproductive organs (within column) for each of three

amputations (within row). (**O-Q**) The size of reproductive structures scale according to patterns conserved

across development, de-growth, and regeneration (penis: R^2 = 0.75, p < 0.0001; seminal vesicle: R^2 =

0.68, p < 0.0001; ovary: R^2 = 0.8, p < 0.0001). (**R**) Across different growth contexts, reproductive organs

appear or disappear at roughly consistent body lengths. Ranking the order in which reproductive

structures are lost in worms undergoing de-growth (**S**) and regenerating worms (**T**) shows that organs are

gained and lost in roughly the same order in all growth contexts. The x-axis shows individual worms in

these plots. dpa = days post amputation. Scale bars: 1 mm.

The male reproductive system

The results above show that imaging using a stereo microscope, even at low resolution,

provides a wealth of information about reproductive morphology and development. However,

the detailed morphology of these structures cannot be ascertained based on these images. We

therefore used a combination of histology, immunofluorescence, staining with live dyes, and

fluorescence *in situ* hybridization (FISH) to visualize male reproductive structures at high

resolution in worms at different stages of development (Fig. 4A; Fig. S4A,B; S5A).

The most anterior part of the male copulatory apparatus is the penis. This structure is

difficult to visualize under conventional illumination, but can be seen when worms extend it,

which we have observed them do both spontaneously and during mating (data not shown).

Staining fixed worms with the dye SiR-actin labels the structure clearly, and in early adult worms

lacking ovaries, we were able to use confocal imaging to visualize it at high resolution (Fig. 4B-

C). Components of the penis can also be seen clearly in histological sections of adult worms

(Fig. 4D). These show that the penis consists of a bundle of rigid needle-like structures (referred

to as stylet needles) (Fig. 4E-G). Our observations of the stylet match the description of the

cuticular rods described in other species in the genus and in other studies of *Hofstenia miamia*

(24–28). The stylet is situated immediately posterior to the base of the penis sheath - a long,

conical structure (Fig. 4C,D; Fig. S4E,H,J; Video 1). The base of the sheath is cup-shaped, and

is lined by posterior-facing hair-like extensions (Fig. 4G). At its anterior end, it connects to the

male gonopore (Fig. 4C, Fig. S5B), both of which are lined with cilia (Video 1,2), consistent with

reports of a ciliated sheath in other hofsteniid species (24–28). Immunofluorescence with an

antibody against Par3 (see Methods) revealed a set of cells that resemble neurons encircling

the base of the sheath (Fig. S4C,D). Further studies will be needed to establish the true identity

and function of these cells.

Next, we focused on the teardrop-shaped prostatic vesicle, which is situated anterior to

the seminal vesicle and is connected to the posterior end of the penis stylet (Fig. 4H-I). Our

histological studies shows that the prostatic vesicle is surrounded by glands that likely produce

granular secretions (Fig. S5,H). The secretions of this organ mix with the sperm as it passes

through, and it has been speculated that it is important for the health and vitality of the sperm,

although the nature of this secretion has not been identified (53, 54). Unexpectedly, our

immunofluorescence experiments revealed a thin layer of cells enclosing the prostatic vesicle

(Fig. 4I). This layer may be an epithelial tissue most likely associated with the extravesicular

prostatic glands. The interior of the prostatic vesicle contains densely-packed sperm cells (Fig.

4D,I; Fig. S4E; Fig. S5C-E; Fig. S5F-G).

The posterior of the male copulatory apparatus contains the seminal vesicle (Fig. 4J).

Histology, as well as nuclear staining of this organ in adult worms (Fig. 4K-L; Fig. S5C-E,G,H)

reveals that it consists entirely of sperm cells. To confirm this, we dissected the seminal vesicle

out of adult worms, and stained it with a nuclear dye – we found that individual cells emerged

from the structure over time, and high-magnification imaging showed that they were indeed

sperm cells, each with a long (~23um) nucleus, and a 22um-long flagellum (Fig. 4M). During

one mating event, we observed a failed attempt at insemination by a worm that resulted in the

release of a packet of sperm outside its partner. We recovered this 0.6 mm-long packet of

sperm (Fig. S4F-G) and found that it consisted of cells with the same morphology as those in

the seminal vesicle (Fig. S4H), confirming that these are indeed sperm. Histological staining as

well as immunofluorescence staining against the muscle marker tropomyosin, showed that the

seminal vesicle and prostatic vesicle are both encircled by layers of muscle (Fig. S4E; Fig. S5C-

H).

We then turned to the testes (Fig. 4N). Nuclear staining of adult worms (cut sagittally

prior to fixation) revealed dense bundles of sperm in the parenchyma, extending much of the

way around the circumference of the worm (Fig. O-Q). The location of the testes is consistent

with previous descriptions (24–28). Our histology of transverse sections of adult worms

confirmed these observations (Fig. 4R-T). Immunofluorescence staining against tropomyosin

showed that the testes are embedded within muscle layers in the parenchyma (Fig. S4L). To

understand the structure of the testes better, we used FISH for putative germline markers

revealed by our previous single-cell RNA-sequencing data (19). We found that one of these

marker genes, *pa1b3-2*, specifically labeled the testes (Fig. 4U), showing them to be paired,

lateralized structures in the worm's anterior.

Finally, we conducted a developmental time course study to visualize the maturation of

the male reproductive system. In juvenile worms, at early developmental stages, actin staining

reveals the prostatic vesicle is virtually absent and progressively emerges as the animal

matures (Fig. S4A). These data also show that the sheath grows progressively. We also

performed FISH to detect mRNA for *pa1b3-2*, using worms across multiple developmental

stages. In juvenile worms (smaller and lacking mature reproductive structures), there was no

expression (Fig. S4I-K). In older worms we began to see expression of *pa1b3-2*, with the

expression domain expanding along the anterior-posterior axis with age (Fig. S4I-K).

364   Together, our data provide a high-resolution view of *H. miamia's* elaborate male

365   reproductive morphology, and allow a more comprehensive understanding of its organization

366   and function.

Figure 4: The male reproductive system of *Hofstenia* includes a penis with a collection of needles,

a sperm containing organ (seminal vesicle), and two testes that span the dorsal surface. (**A**) A

schematized view of the ventral surface of the worm with male reproductive structures highlighted in red.

(**B**) Schematic of male reproductive structures with the copulatory apparatus highlighted. (**C**) Labeling

with an actin dye (white) labels the male gonopore, sheath, penis stylet, and prostatic vesicle: all

components of the male copulatory apparatus. (**D**) A sagittal histological section also reveals these

organs. (**E**) Schematic of the male copulatory apparatus, with the penis stylet highlighted. (**F**) The stylet

includes a bundle of needles labeled by actin. (**G**) The posterior of the penis sheath terminates in a ring of

hair-like projections, also labeled by actin. (**H**) Schematic of the male copulatory apparatus, with the

prostatic vesicle highlighted. (**I**) Actin staining with a nuclear label (Hoechst) shows that the prostatic

vesicle is enveloped by a thin epithelium-like layer, and contains densely packed sperm. (**J**) Schematic of

the male copulatory apparatus, with the seminal vesicle highlighted. (**K**) The morphology of the copulatory

apparatus in mature, adult worms is similar to that of early adults (previous panels). (**L**) The seminal

vesicle of this adult worm contains densely-packed sperm. (**M**) Dissecting out an adult seminal vesicle

allows labeling of individual sperm cells, showing their distinctive morphology. (**N**) Schematic of the male

copulatory apparatus, with the testes highlighted. (**O-Q**) Nuclear staining on an adult worm, cut sagittally,

reveals the testes, containing dense bundles of sperm organized around clusters of cells in the

parenchyma. (**R-S**) Transverse histological sections confirm this organization of the testes. (**T**) Schematic

of a transverse view of an adult worm's anterior, showing the relative organization of the seminal vesicle

and testes. FISH for the male germline marker *pa1b3-2* labels much of the testes (**U**), showing that they

are paired organs that extend through the dorso-ventral axis of the worm (shown through depth-coloring,

right panel) and at least partially wrap around the head (**V**). Scale bars: 20µm (C), 10µm (F-G, M), 50µm

(D, I,L,P), 100µm (K,O,U), 25µm (Q), 200µm (S).

The female reproductive system

As observed through imaging of adult worms using a stereo microscope, the female

reproductive system in *H. miamia* consists of two lateral ovaries running longitudinally down

either side of the body, and one central cluster of mature eggs located posterior to the pharynx

(Fig. 5A-C). To visualize the structure of the ovaries, we used FISH to label mRNA of another

previously-identified germline marker: *cgnl1-2* (19). We found that this gene specifically labeled

the ovaries (Fig. 5D). Moreover, immunofluorescence staining with a custom antibody against

Piwi, a stem cell and germline marker, also labeled the ovaries (Fig. 5E). We were also able to

visualize the ovaries in our histological sections (Fig. 5C). These methods, together with our

earlier images of ovaries at different stages of development and maturity, showed that oocytes

were not organized by maturity within the ovary (Fig. S6A). We did not detect a membrane or

lining that envelopes the ovaries. This is consistent with previous work that suggests that all

acoels lack true ovaries *sensu stricto*: the oocytes are thought to appear and develop within the

parenchymal tissue without a specialized membrane or tunica to form an organ. Thus, we

conclude that *H. miamia*'s ovaries are asacular, in contrast to true 'sacular' gonads, which are

enclosed in specialized membranes (55–57).

Oocytes in *Hofstenia* are 'follicular'; i.e., they are surrounded by specialized cells known

as follicular cells (25, 55). We visualized this cell layer with immunofluorescence for Piwi and

histology and found that it was consistently present in every oocyte irrespective of its

developmental stage (Fig. 5F-G; Fig. S6B-D). We observed that many follicle cells have very

large nuclei, perhaps the result of the fusion of neighboring cells (Fig. 5G; Fig. S6B-C). We also

found clusters of sperm in the layer of follicular cells surrounding maturing oocytes of a variety

of developmental stages (Fig. 5F,G, insets), in agreement with Bock's (25) suggestion that the

follicle may 'trap' sperm and regulate the fertilization of maturing oocytes. We observed

significant morphological variation in the cellular components of the oocytes within the ovaries.

Early-stage oocytes appeared small, round or oval, although others showed irregular shapes.

All oocytes exhibited a granular substance primarily distributed peripherally and a large nucleus

- a germinal vesicle - occupying a substantial portion of the oocyte (Fig. 5H-J). The germinal

vesicle is arrested in prophase I of meiosis, and is found in the oocytes of nearly all animals

(58–61). This nucleus breaks down during early oogenesis or after fertilization (62)(63). The

presence of a germinal vesicle in the oocytes in *H. miamia*'s ovaries suggests that these

oocytes are unfertilized. It is likely that oocytes mature in the ovaries, the germinal vesicle

breaks down, and then oocytes are fertilized. These mature, fertilized eggs then travel from the

ovary to the central cavity prior to being laid.

We could not determine conclusively where or when fertilization happens, or where the

egg capsule (chorion) is produced. However, the spatial organization of the female reproductive

system suggests that an oocyte must mature, become fertilized, and then be transported to the

central cavity behind the pharynx. Across our internal examination, we did not identify any

oviducts, canals or additional structures that could facilitate this migration from the ovaries to the

central cluster. How eggs are transported thus remains unknown.

Figure 5: The female reproductive system includes embryos stored at the base of the pharynx and

oocytes in ovaries. (**A**) A schematized view of the ventral surface of the worm with female reproductive

structures highlighted. (**B**) Eggs near the pharynx of the worm (within the red circle) are fertilized and

mature while oocytes in ovaries (red arrow) are immature or unfertilized, with a visible germinal vesicle.

(**C**) A sagittal histological section reveals the organization of the follicles within an ovary. (**D**) FISH shows

that *cgnl1-2* labels immature oocytes (white arrow: germinal vesicle) in the ovaries. (**E**) Oocytes in ovaries

are also labeled by a piwi antibody (white arrow: germinal vesicle). (**F**) Histology of an ovarian follicle

shows that follicle cells encircle the immature oocyte. Inset: sperm cells appear to be trapped in the

follicle. (**G**) piwi immunofluorescence confirms the organization of the follicle (white arrow: germinal

vesicle)), and nuclear staining identifies trapped sperm in its surface. Histology also shows that immature

oocytes may have irregular shapes (**H**), contain a germinal vesicle (yellow arrow) (**H,I**), and possess an

abundance of (likely yolk) granules (**I,J**). Scale bars: 100µm (B, C (inset), D-F), 500µm (C), 50µm (G, H-J)

Egg-laying behavior

Egg laying has only been directly observed a few times in acoels, mostly in species belonging to

more recently diverged clades (13, 64) (Fig. 1B). From these observations, acoels are known to

lay eggs through the female gonopore (if present) (64), a mode of egg laying used by many

flatworms (65), or through breaks in the body wall (13, 14). It has also been suggested that

acoels could lay eggs through the mouth; however, attempts at direct observation of egg laying

have been inconclusive (13, 41).

Previous authors have speculated that *H. miamia* lay eggs through breaks in the body

wall, or perhaps partially through the mouth, but egg laying was not observed (25, 27). To

observe how worms lay eggs, we isolated gravid adult worms and filmed them from underneath

for a 24 hour period. 3 worms laid 25 eggs in these conditions. We found that worms exclusively

laid eggs through their mouths (Fig. 6A,B; Video 3). Egg-laying occurs in a movement lasting

121 ± 114 (mean ± SD) seconds, during which the worm performs a series of muscle

contractions to transfer a single egg from the ventral pocket of fertilized eggs into the pharynx,

and then further applies muscle contractions from posterior to anterior to move the egg to the

mouth. The worm then places each egg on the substrate with its mouth, likely secreting mucus

to attach the egg to the substrate (Fig. 6A,B; Video 3). Observations from these videos, as well

as of egg clutches in our culturing tanks, show that worms can either deposit a single egg in one

location or lay eggs in one or more clutches.

To understand the temporal dynamics of egg-laying in *H. miamia*, we then quantified the

time course of egg-laying in several contexts. We found that gravid worms with access to

mates, once isolated, continued laying eggs for over a month (Fig. 6C). It is likely that all worms

had recently mated given that all worms laid eggs at the onset of isolation. Juvenile worms

reared to adulthood in isolation also laid a single burst of eggs (Fig. 6D), multiple months after

first being isolated (although the timing of this burst appears to depend on rearing conditions

and the health of the worms). This burst of egg laying is likely the result of a selfing event in

which the worms fertilize their own eggs, or possibly the result of a form of parthenogenesis

allowing the activation of unfertilized eggs. We also allowed virgin worms to mate once, in

controlled conditions, and subsequently isolated them. These mated and inseminated worms

laid eggs for over three months after a single mating, suggesting that sperm may be stored in

follicular cells for several months. (Fig. 6E).

Finally, we asked how much control the worms had over their egg laying. Anecdotal

observations suggested that worms are selective about the quality of their environments, and

about when and where they choose to lay eggs. By quantifying the spatial positions of eggs laid

in their culturing tanks, we found that worms have a strong preference for laying eggs on the

walls of their containers, rather than on the floor (Fig. 6F). Eggs laid on walls are preferentially

laid close to the water line (Fig. 6G, Fig. S7A). This preference is not simply because the worms

lay eggs where they are: we observed that worms spend the majority of their time on the floors

of their boxes, and seem to rise to the surface specifically to lay eggs. These spatial

preferences do not seem affected by food availability: when deprived of food, egg laying still

primarily occurs on the walls of culture boxes (Fig. S7B). Together, this shows that worms

actively prefer to lay eggs close to the water line, and more generally, that substrate choice is

an active decision-making behavior.

We then asked whether worms actively choose to withhold eggs in suboptimal

environments. Our observations (see Fig. 3A) suggested that worms are physiologically resilient

to food stress, but not to reductions in water quality (data not shown). We therefore conducted

an experiment with a set of worms, half of which were deprived of food for one feeding period

(3-4 days). The other set was fed normally during this period. We found that food-deprived

worms laid significantly fewer eggs during this period (Fig. 6H). We hypothesized that this

reduction in egg-laying was because the worms were actively withholding their eggs in the

absence of food. Next, we repeated this experiment, first depriving some worms of food and

subsequently feeding them. After food-deprived worms were given food, they laid an excess of

eggs (Fig. 6I), suggesting that the worms actively choose to withhold egg-laying in food-limited

environments. More generally, these data show that *H. miamia* assess their environments to

make active decisions about when and where to lay eggs.

**Figure 6: *H. miamia* lays eggs through the mouth and exhibits environmental preferences in egg

laying.** (**A**) Sequence of images from a video of egg-laying. Eggs in the pharynx and emerging through

the mouth are shaded blue. (**B**) Schematic showing presumed process of embryo traveling from the cavity

beneath the pharynx to the pharynx and then out through the mouth. (**C**) Histogram showing the timing of

eggs laid by adult worms living in communal enclosures and then isolated. (**D**) Histogram showing the

timing of eggs laid by worms that undergo reproductive development in isolation and then self-fertilize. (**E**)

Histogram showing the timing of eggs laid by worms that are allowed to mate once. (**F**) Scatterplot of the

percentage of eggs found on the floor of communal enclosures (n = 30). This is significantly different from

the expected percentage of eggs based on enclosure surface area (t-test p < 0.0001). (**G**) Kernel density

estimate of egg locations on a subset of enclosure surfaces with similar dimensions (n = 2144 eggs). (**H**)

Worms that are unfed for 4 days lay fewer eggs than fed worms (n = 9, t-test p < 0.0001). (**I**) Unfed worms

that are then fed lay more eggs than worms that are continuously fed (n ≥ 8, t test = < 0.0001).

Discussion

Our results provide a quantitative study of acoel reproduction for the complete life cycle from

embryo to adult egg-laying at unprecedented resolution, filling a gap in our knowledge of animal

reproduction in an understudied animal clade and laying a foundation for further study of

anatomy, physiology, and behavior in acoels. These data have general implications for our

understanding of reproductive anatomy in acoels, and for physiological and behavioral elements

of reproduction.

Implications for the reproductive anatomy of acoels

Our description of *H. miamia*'s anatomy affirms many previous descriptions, including Bock's

(25) observations of male and female reproductive systems of *Hofstenia atroviridis*. By using

molecular techniques and high-resolution imaging we were able to expand our understanding of

the functional organization of reproductive structures and to reveal aspects of *H. miamia*'s

physiology. For example, in addition to locating the seminal vesicle, prostatic vesicle, and

needle-like penis, we identified a ring of hair-like protrusions lining the posterior of the sheath.

We speculate that during mating, muscles push the penis stylet anteriorly, through the sheath

and out of the male gonopore. This movement could evert the sheath, resulting in a penis-

extension state in which the sheath emerges through the gonopore around the stylet, with the

ring of protrusions now near the anterior tip of the penis. It is possible that these protrusions

have a sensory function, and could inform fine penis movements during hypodermic

insemination. However, the behavior of the sheath has not yet been observed, and the function

of the ring remains unknown.

Taxonomic implications of sperm with a single flagellum

Acoels are reported as having biflagellate sperm (6, 66), but the exact definition of "biflagellate"

is unclear. This has been interpreted as the presence of two flagella (67), but it has also been

argued that only spermatids have two free flagella (6). Under this argument, the two flagella

fuse during spermatogenesis, yielding a sperm with a single flagellum that has two axonemes

(6, 68). This argument has been supported by ultrastructural studies of sperm, where two

axonemes are present in a single sperm cell (66, 69, 70) and imaging of whole sperm cells from

Paratomella rubra (69) and *Amphiscolops oni* (70), where there appears to be a single

flagellum. In our imaging of *H. miamia* sperm, we also observed sperm with a single flagellum.

The presence of a single flagellum in these three distantly related acoels supports the

interpretation that biflagellate refers to the presence of two axonemes rather than two free

flagella.

Selfing

Asexual reproduction in acoels has been demonstrated in just a handful of species that typically

use fission and budding to reproduce (66, 71–75). Although many other invertebrate

hermaphrodites, such as snails (76), tapeworms (77), and flatworms (78), are known to

reproduce through self-fertilization, it is thought that acoels typically reproduce through mating.

Our observation that *H. miamia* raised in isolation will lay eggs once reproductively mature

suggests that it is capable of self-fertilization (or possibly of asexual reproduction through

parthenogenesis). How self-fertilization could occur is unclear, but as sperm cells are observed

scattered through the head, it is possible they are able to migrate to the eggs in the posterior of

the worm. Nearly all acoels are hermaphroditic (8), and many share anatomical features with

Hofstenia (Fig. 1B). It is possible that reproduction via self-fertilization is a common life history

strategy across the acoels.

Development, regeneration, and allometry

Little is known about differentiation of reproductive structures in most acoels. One exception is

Otocelis luteola, an acoel that likely mates through a "genital handshake" (79) that results in

fertilization (45) Previous work found that the female gonopore and vagina develop first (42).

Sperm cells in the testes and the seminal vesicle then become visible. Roughly at the same

stage, oocytes also begin to develop. The ovaries later expand in size and produce increasingly

mature oocytes, and the vagina, female gonopore, and seminal bursa develop further. After

mating, the seminal bursa finally grows into its mature form. When *Otocelis luteola* is starved, it

experiences de-growth and de-differentiation of reproductive structures. Mature oocytes

disappear first, followed by sperm within the seminal vesicle. The testes are lost next, and the

female gonopore and other structures follow. In regenerating fragments of *Isodiametra pulchra*,

male structures develop and differentiate earlier than the female reproductive structures (43).

While sparse, these data and ours suggest that the regulation of reproductive organ growth may

vary across acoel species. Our data in *H. miamia* show that reproductive organs grow and scale

with body size, that they appear stepwise at increasing body sizes during the developmental

transition to adulthood, and that most of these organs are lost at the same sizes during

starvation-induced de-growth. Moreover, there may be size-dependent growth mechanisms that

ensure synchronized, symmetric organ growth. These findings enable future study of how

growth-relevant information is transmitted to all regions of the body to coordinate organ size

during growth and de-growth.

Egg-laying dynamics and implications for life history

Due to difficulties observing egg-laying, egg-laying modes are only known for a few species (13,

37, 41). Here, we filmed *H. miamia* laying eggs. It has been proposed that some acoels may lay

eggs through their mouths, a mode of egg-laying not found in other animals; here we provide

direct evidence that *H. miamia* do indeed lay eggs this way. We suspect that other acoels with a

similar suite of reproductive traits (i.e., lack of a female gonopore, a cluster of mature eggs near

the pharynx) could also lay eggs through their mouths. This unusual mode of egg laying raises

several questions. First, where in the animal does fertilization occur? Our observations suggest

that follicular cells surrounding immature oocytes act as a selective barrier to sperm, preventing

many sperm cells from reaching the oocyte, a function suggested by Bock (1923). The presence

of a germinal vesicle in all oocytes within the ovaries, but not in eggs in the central cavity, raises

the possibility that the follicle may regulate the timing of fertilization within the ovary, soon after

which the egg travels to the central cavity. Second, how do mature eggs travel from the lateral

ovaries to the central cavity, and how are the eggs in this cavity physically loaded into the

pharynx? Our anatomical observations have so far failed to reveal obvious oviducts, or other

tube-like structures that could facilitate movement of eggs into the pharynx. The only known

opening at the posterior of the pharynx is the pharyngeal sphincter that allows food to pass into

the gut. We therefore speculate that eggs travel through the gut to the central cavity, which may

simply be a pocket within the gut. Eggs may then be moved into the pharynx through a form of

reverse peristalsis (and observation of the worm prior to egg-laying shows waves of muscle

contraction proceeding posterior-to-anterior as each egg is loaded into the pharynx).

In addition to identifying the mode of egg laying in *H. miamia*, we also examined egg-

laying patterns over time. After a single mating, we find a peak in egg-laying around 2-3 weeks

later, but eggs are laid continuously for nearly 200 days. When isolated from a communal

enclosure where mating can happen continuously, eggs are laid for around 40 days. These

differences in the duration of egg laying may be in part due to corresponding differences in how

frequently worms mate (and in how much sperm is transferred during a single mating), and in

part due to cohort-specific fluctuations in water quality and other environmental factors.

Nonetheless, these data show that worms are able to lay eggs for months after mating,

suggesting that they are capable of some form of long-term sperm storage. *H. miamia* lacks an

anatomically-distinct organ to store sperm received during mating. We speculate that persistent

egg-laying following mating could be enabled by sperm trapped by the follicular cells

surrounding immature oocytes. As the oocytes mature, the sperm could fertilize the oocyte,

enabling fertilization weeks after the mating event.

Egg-laying behavior

Our results show that *H. miamia* likely makes an active choice of when and where to lay an egg

once that egg is mature and fertilized. *H. miamia* has clear spatial preferences when laying an

egg, preferring the area near the surface of the water. Although the worms seem to mostly be

found on the floor of their enclosures, they do not prefer to lay eggs on the floor. This suggests

that a worm likely navigates to a suitable position before laying an egg. This preference is

robust to food deficiency. *H. miamia* also has environmental preferences, suppressing egg-

laying when deprived of food and then quickly laying eggs once food is present again. Together,

these results show that *H. miamia* surveys its environment and then integrates this assessment

into whether - and where - it is suitable to lay an egg.

Conclusion

635 These findings establish a foundation of knowledge of anatomical, physiological, and behavioral

636 elements of *H. miamia*'s reproductive life history. This enables future work on the molecular

637 genetics of reproductive organ development and regeneration, on the physiological processes

638 involved in egg maturation and oviposition, and on the neuroscience of reproductive behavior.

Methods

Animal husbandry

Gravid *H. miamia* are reared in communal plastic 2.25 L boxes with about 1.25 L of artificial

seawater (37 ppt, pH 7.8-8.2). These communal enclosures are kept in incubators held at 21°C.

Twice weekly, embryos are collected, seawater is changed, and *Artemia* sp. (brine shrimp) are

added. Embryos are raised in petri dishes. Once hatched, juvenile worms are raised in tanks at

room temperature and are fed marine rotifers *Brachionus plicatilis*. When the reproductive

organs begin to develop, worms are transferred to larger tanks, each typically housing 30-50

worms. Animals paired for mating were transferred from juvenile tanks to 24-well plates and

reared in isolation following similar feeding and cleaning routines as listed above. In order to

assay regeneration of reproductive systems, adult worms were isolated from communal

enclosures, anesthetized in 15% tricaine (ethyl 3-aminobenzoate methanesulfonic acid), and

then amputated with a dissection blade. Regenerating fragments were maintained in 6-well

plates and cleaned twice weekly. Five days after amputation, fragments were fed *Artemia* sp. or

rotifers. Animals that were starved were also maintained in 6-well plates that were cleaned twice

weekly but with food omitted. All animals are derived from an inbred population of worms

collected in 2010 from Bermuda.

Histology

Adult specimens of *H. miamia* were fixed in 4% paraformaldehyde in artificial seawater for 24

hours at room temperature. Following fixation, specimens were washed twice with 70% ethanol

to remove the fixative and then preserved in a third change of 70% ethanol for histological

preparation. Animals were dehydrated through a graded ethanol series (95% and 100%),

cleared in Histoclear, and embedded in Leica Surgipath (Paraplast) with a melting point of 56°C.

Longitudinal, transverse, and sagittal sections were obtained at a thickness of 5 μm using a

Leitz 1512 microtome. The sections were stained using a standard protocol with hematoxylin

and eosin and mounted on glass slides with Permount. Imaging of the sections was performed

using a Zeiss Axio Scan.Z1, and an Olympus BX50 microscope.

Fluorescence in situ hybridization (FISH)

Riboprobes were synthesized following protocols established in Srivastava et al. (2014).

Following 1-2 weeks of starvation, whole worms were fixed in 4% paraformaldehyde in artificial

sea water for one hour at room temperature. Fixed animals were then transferred to 24-well

plates in small baskets with a mesh bottom, with 4-5 animals in each basket. To remove

pigment autofluorescence, animals were then transferred to a bleach solution (containing 4%

hydrogen peroxide, SSC, and formamide) and left under a light for two hours. For all washes,

800 μL of solution was used. The animals were then permeabilized using proteinase K solution

(0.1% SDS, 1 μL/10 mL proK in PBSTx). After 10 minutes, worms were fixed again in 4%

formaldehyde in PBST, washed twice in PBST, and then washed in a 1:1 PBST:PreHyb solution

for 10 minutes. Specimens were then incubated in the PreHyb solution (50% DI formamide,

25% 20X SSC, 0.05% 10% Tween-20, 1mg/mL yeast tRNA, 20% water) for two hours in a 56°C

hybridization oven and then transferred to a hybridization solution (50% DI formamide, 25% 20X

SSC, 0.05% 10% Tween-20, 1 mg/mL yeast tRNA, 10% 50% dextran sulfate, 10% water)

containing riboprobe(s) and left overnight. The specimens were then put through several 30-

minute washes at 56°C, starting with two PreHyb washes, two 1:1 PreHyb:2XSSCT washes,

two 2XSSCT (2XSSC with 0.1% TritonX-100) solution washes, two 0.2XSSCT (0.2X SSC with

0.1% TritonX-100) solution washes. Specimens were then left to cool at room temperature and

then washed in two ten-minute PBST washes. A 1-hour blocking at room temperature was then

performed with blocking solution (5% horse serum and 5% casein in PBST). Animals were then

incubated with anti-Digoxigenin-POD (1:1500 dilution; Roche, 11633716001) in a blocking

buffer overnight. The following day, animals were washed 6 times with PBST, incubated with a

tyramide buffer (1.1688g/mL NaCl, 6.18mg/mL boric acid, filled with water and adjusted pH to

8.5) for 10 minutes, then developed with rhodamine-conjugated tyramide solution for 10

minutes. Animals were washed again with PBST and incubated for 1 hour in 1:500

PBST:Hoechst to label nuclei. Animals were then mounted on glass slides with

VECTASHIELD® PLUS Antifade Mounting Medium (Vector Laboratories, H-1900).

Whole-mount immunofluorescence

Worms were fixed in 4% paraformaldehyde (Electron Microscopy Sciences, 15714) in artificial

sea water for 1-2 hours at room temperature (juveniles were fixed for one hour, adults were

fixed for two) before washing with PBST (PBS + 0.1% Triton-X-100). Adult worms were

anesthetized in 15% tricaine for 5 minutes prior to fixation to minimize epidermal rupture. For

immunofluorescence, worms were washed in PBST, blocked for one hour at room temperature

in 10% goat serum in PBST, and incubated in primary antibody for 48-72 hours at 4C on a

shaker (juveniles only require 48 hours, adults need the full 72 hours). The following day, worms

were washed thoroughly in PBST (8 x 20 min washes) on a nutator before blocking for one hour

at room temperature on a shaker. Worms were incubated in secondary antibody overnight at

4C; the following day, worms were washed thoroughly in PBST (8 x 20 in washes) on a shaker

before adding direct conjugate dyes and mounting on glass slides with VECTASHIELD®

HardSet™ Antifade Mounting Medium (Vector Laboratories, H-1400-10). Antibodies used:

Tropomyosin (custom, (80), Piwi-1 (custom), Par3 (St. John's Laboratory, STJ502116), Goat

anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568 (ThermoFisher,

A-11011), Hoechst (ThermoFisher, H1399), SiR-actin (Cytoskeleton, CY-SC001). All images

with FISH and immunofluorescence were acquired with a Leica SP8 point-scanning confocal

microscope.

Quantitative analysis of development, regeneration, and de-growth

We collected 42 zygotes laid from wild-type worms. Each embryo was transferred to a well of a

24-well cell culture plate with artificial seawater. Plates were stored in a temperature-controlled

incubator. Without removing the embryos from the plate, they were imaged twice weekly

through a dissection microscope with white illumination from LEDs mounted above the sample.

Water was also changed on these days. The same volume of water was added to each well

(2.35 mL in each 12-well). Once at least one embryo had hatched, rotifers were added to each

well. On each feeding day, the concentration of rotifers was calculated by counting the number

of rotifers in diluted samples. Based on this concentration, the volume of rotifers was altered to

ensure a consistent number of rotifers was given to each animal. At different stages during the

experiment, the number of rotifers (Fig. S2A) fed increased. Increases were gradual to ensure

worms always had more food than they could eat, while minimizing excess food which could

affect water quality. Hatched worms were removed from wells for imaging twice weekly. Worms

were anesthetized in tricaine (15% ethyl 3-aminobenzoate, methanesulfonic acid salt; Sigma-

Aldrich #E10521), and mounted on microscope slides with a 20x20 mm sticker grid with 1mm

resolution (ThomasScientific, #1207X53). When worms were larger, they were moved to larger

6-well plates in 10 mL of artificial sea water and began to be fed brine shrimp as well (Fig.

S2A). The number of brine shrimp added was measured using the same concentration

procedure outlined above for rotifers. Once worms had matured, the number of brine shrimp

increased.

Images were manually annotated in FIJI. A line was drawn to capture the length of the

sticker grid to define an image-specific scale. Additional lines were drawn to measure the length

from the most posterior part of the male reproductive system (e.g., base of the penis tube before

sperm production begins or base of the seminal vesicle) to the male gonopore, the length of the

seminal vesicle, the length of each ovary, the length of the worm's body, and the width of the

worm's body. Measurements were converted to mm using image-specific scale information.

Penis length was calculated by subtracting the length of the seminal vesicle from the length of

the male gonad.

These imaging and measurement processing procedures were repeated with starving

worms and regenerating worm fragments. Starved and fed control worms were anesthetized

and imaged roughly once a week for 11 weeks. 106 days after isolation, starved worms were

imaged again, at which point one starved worm had died. Regenerating worms were imaged

starting 2 days post-amputation and then every 2-5 days until 25 days post-amputation at which

point images were taken less frequently. Regenerating fragments were not fed for the first four

days after amputation, and were then fed normally. In these experiments, a two-pointed line or

multi-jointed spline was used to measure lengths. All analyses were then performed in python,

with code assistance from GPT4.

Egg-laying behavior

Adult worms from a communal tank were moved to individual wells in the center of a 24-well

plate or 12-well plate (Fisher Scientific #07-200-82). We mounted the plate on an elevated

platform within an enclosed behavioral rig. The plate was illuminated with uniform white light,

from addressable LEDs (Adafruit #SK6812RGBW, powered by 5V DC) arranged in a ring

around the plate behind a white acrylic diffuser (McMaster-Carr #8505K741). An infrared

camera (Basler Ace ac4024-29um USB 3.0 monochrome) was mounted underneath the plate to

allow filming of the worm's ventral surface. A USB fan (Amazon #B0C27NGKCV) was mounted

on the floor of the behavioral chamber for temperature regulation. We filmed worms at 1Hz for

8-12 hours. After filming, each worm was returned to its original communal tank.

To find egg positions, communal tanks were maintained according to our husbandry

protocol. Every 3-4 days, before cleaning, images were taken of each face of the box (2 short

faces, 2 long faces, and the floor of the box). The x- and y-coordinates of the eggs in small and

long face images were scored in FIJI. The positions of eggs laid on the floor were not scored

due to low density. The x- and y-coordinates of the boundaries of the water in each image were

also scored. From these image-specific boundaries, a common set of bounds was calculated for

each type of face (i.e., short, long). Egg coordinate positions were then scaled and aligned to

these common bounds for analysis.

For experiments measuring when worms laid eggs, communal enclosures of adult

worms were cleaned twice weekly and fed well. Three days later, eggs were collected from

each box and counted. On this day, half of these enclosures were randomly selected to have

food withheld. After four days with or without food, eggs were collected from all boxes and

counted. To test whether egg laying is recovered when unfed worms are fed, communal

enclosures of adult worms were randomly selected to have food or to have food withheld. Four

days later, eggs were collected and counted and then all communal enclosures were fed. Three

days later, eggs were collected and counted.

Author Contributions

VC, SET, and MS designed the project; SET and VC performed development, physiology and

behavior experiments; VC and SET analyzed the data; APK performed and interpreted

immunofluorescence and imaging experiments; DMB performed and interpreted histology

experiments in the context of prior anatomical literature; VC, SET, and MS wrote the manuscript

with input from all authors; and MS supervised the project.

Acknowledgements

We thank Gonzalo Giribet, Matthew Hooge, James Hanken, Javier Ortega-Hernandez, Andrew

Berry, and all members of the Srivastava lab for helpful discussion. We also thank James

Hanken, Júlia Chaumel, and Christina Daly for histology help and infrastructure. VC and APK

are Fellows of the Jane Coffin Childs Fund for Medical Research. ST acknowledges funding

from the Harvard Museum of Comparative Zoology, Herchel Smith Undergraduate Science

Research Program, and the Program for Research in Science and Engineering. This project

was supported by grants from the National Institute of General Medical Sciences of the NIH to

MS: R35GM128817 and R35GM153252.

References

1.   N. Davies, J. Krebs, S. West, *An Introduction to Behavioural Ecology* (Wiley-Blackwell,

2013).

2.   J. T. Cannon, *et al.*, Xenacoelomorpha is the sister group to Nephrozoa. *Nature* **530**, 89–93

(2016).

3.   P. Kapli, M. J. Telford, Topology-dependent asymmetry in systematic errors affects

phylogenetic placement of Ctenophora and Xenacoelomorpha. *Sci Adv* **6**, eabc5162

(2020).

4.   H. Philippe, *et al.*, Acoelomorph flatworms are deuterostomes related to Xenoturbella.

Nature **470**, 255–258 (2011).

5.   Tyler, S., Schilling, S., Hooge, M., & Bush, L. F., Turbellarian taxonomic database. Version

1.7. (2012). Available at: http://turbellaria.umaine.edu.

6.   J. G. Achatz, M. Chiodin, W. Salvenmoser, S. Tyler, P. Martinez, The Acoela: on their kind

and kinships, especially with nemertodermatids and xenoturbellids (Bilateria incertae

sedis). *Org. Divers. Evol.* **13**, 267–286 (2013).

7.   A. Wallberg, The Stylet - Diversity and Systematics of Acoela and Nemertodermatida.

(2012). Available at: http://acoela.myspecies.info.

8.   U. Jondelius, A. Wallberg, M. Hooge, O. I. Raikova, How the Worm Got its Pharynx:

Phylogeny, Classification and Bayesian Assessment of Character Evolution in Acoela. *Syst.*

Biol. **60**, 845–871 (2011).

9.   Y. V. Mamkaev, Etude morphologique d'*Actinoposthia beklemischevi* n. sp. (Turbellaria

Acoela). *Cah. Biol. Mar.* **6**, 23–50 (1965).

10. O. Raikova, A. Falleni, V. Gremigni, Oogenesis in *Actinoposthia beklemischevi*

(Platyhelminthes, Acoela): an ultrastructural and cytochemical study. *Tissue Cell* **27**, 621–

633 (1995).

11. G. Giribet, G. D. Edgecombe, *The Invertebrate Tree of Life* (Princeton University Press,

2020).

12. X. Bailly, *et al.*, The chimerical and multifaceted marine acoel *Symsagittifera roscoffensis*:

from photosymbiosis to brain regeneration. *Front. Microbiol.* **5**:498 (2014).

13. H. M. Costello, D. P. Costello, Egg laying in the acoelous turbellarian *Polychoerus*

carmelensis. *Biol. Bull.* **76**, 80–89 (1939).

14. G. Apelt, Fortpflanzungsbiologie, Entwicklungszyklen und vergleichende Frühentwicklung

acoeler Turbellarien. *Mar. Biol.* **4**, 267–325 (1969).

15. T. Shannon, J. Achatz, *Convolutriloba macropyga* sp. nov., an uncommonly fecund acoel

(Acoelomorpha) discovered in tropical aquaria. *Zootaxa* **1525**, 1–17 (2007).

16. T. Zauchner, W. Salvenmoser, B. Egger, A cultivable acoel species from the

Mediterranean, *Aphanostoma pisae* sp. nov. (Acoela, Acoelomorpha). *Zootaxa* **3941**, 401–

413 (2015).

17. K. De Mulder, *et al.*, Characterization of the stem cell system of the acoel *Isodiametra*

pulchra. *BMC Dev. Biol.* **9**, 69 (2009).

18. M. Srivastava, Studying development, regeneration, stem cells, and more in the acoel

Hofstenia miamia. *Curr. Top. Dev. Biol.* **147**, 153–172 (2022).

19. R. E. Hulett, *et al.*, Acoel single-cell atlas reveals expression dynamics and heterogeneity of

adult pluripotent stem cells. *Nat. Commun.* **14**, 2612 (2023).

20. L. Ricci, M. Srivastava, Transgenesis in the acoel worm *Hofstenia miamia*. *Dev. Cell* **56**,

3160–3170.e4 (2021).

21. J. O. Kimura, D. M. Bolaños, L. Ricci, M. Srivastava, Embryonic origins of adult pluripotent

stem cells. *Cell* **185**, 4756–4769.e13 (2022).

22. A. R. Gehrke, *et al.*, Acoel genome reveals the regulatory landscape of whole-body

regeneration. *Science* **363**: eaau6173 (2019).

23. J. O. Kimura, L. Ricci, M. Srivastava, Embryonic development in the acoel *Hofstenia

miamia*. *Development* **148**, dev188656 (2021).

24. M. Hooge, *et al.*, A revision of the systematics of panther worms (*Hofstenia* spp., Acoela),

with notes on color variation and genetic variation within the genus. *Hydrobiologia* **592**,

439–454 (2007).

25. S. Bock, *Eine neue marine Turbellariengattung aus Japan* (Almqvist & Wiksell, 1923).

26. D. D. Corrêa, Two New Marine Turbellaria from Florida. *Bull. Mar. Sci.* **10**, 208–216 (1960).

27. O. Steinböck, Grundsätzliches zur Evolution der Turbellarien. *Zeitschrift für Zoologische

Systematik und Evolutionsforschung* **4**, 58–195 (1966).

28. S. Beltagi, A. Mandura, *Hofstenia arabiensis* Nov. Sp. (Hofsteniidae): A New Species of

Acoelan Turbellaria from the Red Sea North of Jeddah. *Journal of King Abdulaziz

University-science* **3**, 65–90 (1991).

29. A. Faubel, B. Cameron, Platyhelminthes from salt marshes of Coomera River,

Southeastern Queensland, Australia. *Mem Queensland Mus* **46**, 511–519 (2001).

30.   J. Dörjes, Die Acoela (Turbellaria) der Deutschen Nordseeküste und ein neues System der

Ordnung. *J. Zoolog. Syst. Evol. Res.* **6**, 56–452 (1968).

31.   M. Boone, W. Houthoofd, W. Bert, T. Artois, First record of Nemertodermatida from Belgian

marine waters. *Belg. J. Zool.* **141**, 62–64 (2011).

32.   W. E. Sterrer, New and known Nemertodermatida (Platyhelminthes- Acoelomorpha) - a

revision. *Belgian Journal of Zoology* **128**, 55–92 (1998).

33.   J. G. Achatz, M. D. Hooge, Convolutidae (Acoela) from Tanzania. *Zootaxa* **1362**, 1 (2006).

34.   M. Hooge, Two new families, three new genera, and five new species of acoels (Acoela

Platyhelminthes) from Queensland Australia. *Cah. Biol. Mar.* **44**, 275–298 (2003).

35.   M. Hooge, J. P. S. Smith, New acoels (Acoela, Acoelomorpha) from North Carolina.

Zootaxa **442**, 1–24 (2004).

36.   H. M. Costello, D. P. Costello, Copulation in the acoelous turbellarian *Polychoerus*

carmelensis. *Biol. Bull.* **75**, 85–98 (1938).

37.   E. Gardiner, Early development of *Polychoerus caudatus*, Mark. *J. Morphol.* **11**, 155–176

(1895).

38.   L. H. Hyman, Reproductive system and copulation in *Amphiscolops langerhansi*

(Turbellaria Acoela). *Biol. Bull.* **72**, 319–326 (1937).

39.   F. Peebles, A Description of three Acoela from the Gulf of Naples. *Mitteilungen a.d. Zool.*

Station zu Neapel **22**, 292–312 (1915).

40.   L. Bush, Biology of *Neochilda fusca* N. gen., n. sp. from the northeastern coast of the

United States (Platyhelminthes: Turbellaria). *Biol. Bull.* **148**, 35–48 (1975).

41. M. C. Watzin, Egg laying in the acoel turbellarian *Neochildia fusca*. *Trans. Am. Microsc.*

Soc. **103**, 322–325 (1984).

42. E. N. Kozloff, A new genus and five new species of acoel flatworms from the Pacific coast

of North America, and resolution of some systematic problems in the families Convolutidae

and Otocelididae. *Cah. Biol. Mar.* **41**, 281–293 (2000).

43. E. Perea-Atienza, *et al.*, Posterior regeneration in *Isodiametra pulchra* (Acoela,

Acoelomorpha). *Front. Zool.* **10**, 64 (2013).

44. N. W. Riser, *Nemertinoides elongatus* gen. n., sp. n. (Turbellaria: Nemertodermatida) from

coarse sand beaches of the Western North Atlantic. *Proc. Helminthol. Soc. Wash.* **54**, 60–

67 (1987).

45. D. Grae, E. Kozloff, Post-copulatory storage of sperm in the seminal bursa of the acoel

flatworm *Otocelis luteola*. *Invertebr. Reprod. Dev.* **35**, 151–154 (1999).

46. I. Meyer-Wachsmuth, M. Curini Galletti, U. Jondelius, Hyper-cryptic marine meiofauna:

species complexes in Nemertodermatida. *PLoS One* **9**, e107688 (2014).

47. M. Crezee, *Paratomella rubra* Rieger and Ott, an amphiatlantic acoel turbellarian. *Cah. Biol.*

Mar. **19**, 1–9 (1978).

48. A. G. Kostenko, The position of Taurida fulvomaculata (Ax, 1959) in the system of Acoela.

In: Morphology of turbellarians. *Akad Nauk USSR Trudy Zool* **195**, 14–25 (1989).

49. J. Dörjes, *Paratomella unichaeta*, nov. gen. nov. spec., Vertreter einen neuen Familie der

Turbellaria Acoela mit asexueller Fortpflanzung durch Paratomie. *Veröffentlichungen des*

Instituts für Meeresforschung in Bremerhaven **II**, 187–200 (1966).

50. A. Faubel, Die Acoela (Turbellaria) eines Sandstrandes der Nordseeinsel Sylt. *Mikrofauna*

Meeresbodens **32**, 1–58 (1974).

51. A. Faubel, Interstitielle Acoela (Turbellaria) aus dem Littoral der nordfriesischen Inseln Sylt

und Amrum. *Mitt. Hamb. Zool. Mus. Inst.* **73**, 17–56 (1976).

52. M. Srivastava, K. L. Mazza-Curll, J. C. van Wolfswinkel, P. W. Reddien, Whole-Body Acoel

Regeneration Is Controlled by Wnt and Bmp-Admp Signaling. *Curr. Biol.* **24**, 1107–1113

(2014).

53. L. H. Hyman, *The invertebrates: Platyhelminthes and Rhynchocoela. The acoelomate*

Bilateria (McGraw-Hill, New York, 1951).

54. A. Faubel, The Polycladida, Turbellaria. Proposal and establishment of a new system. Part

I. The Acotylea. *Mitt. Hamb. Zool. Mus. Inst.* **80**, 17–121 (1983).

55. R. M. Rieger, S. Tyler, J. P. S. Smith, G. E. Rieger, "Platyhelminthes and Nemertinea" in

Microscopic Anatomy of Invertebrates, (Wiley-Liss, 1991), pp. 7–140.

56. A. Schmidt-Rhaesa, *The Evolution of Organ Systems* (OUP Oxford, 2007).

57. K. Eckelbarger, A. Hodgson, Invertebrate oogenesis – a review and synthesis: comparative

ovarian morphology, accessory cell function and the origins of yolk precursors. *Invertebr.*

Reprod. Dev. **65**, 71–140 (2021).

58. C. Munro, H. Cadis, S. Pagnotta, E. Houliston, J.-R. Huynh, Conserved meiotic

mechanisms in the cnidarian *Clytia hemisphaerica* revealed by *Spo11* knockout. *Sci Adv* **9**,

eadd2873 (2023).

59. J.-L. Liu, M. Buszczak, J. G. Gall, Nuclear bodies in the *Drosophila* germinal vesicle.

Chromosome Res. **14**, 465–475 (2006).

60. H. Grossman, *et al.*, Regulation of GVBD in mouse oocytes by miR-125a-3p and Fyn

kinase through modulation of actin filaments. *Sci. Rep.* **7**, 2238 (2017).

61. L. Herlands, G. G. Maul, Characterization of a major nucleoplasmin-like germinal vesicle

protein which is rapidly phosphorylated before germinal vesicle breakdown in *Spisula*

solidissima. *Dev. Biol.* **161**, 530–537 (1994).

62. N. Sagata, Meiotic metaphase arrest in animal oocytes: its mechanisms and biological

significance. *Trends Cell Biol.* **6**, 22–28 (1996).

63. A. Falleni, V. Gremigni, Ultrastructural study of oogenesis in the acoel turbellarian

Convoluta. *Tissue Cell* **22**, 301–310 (1990).

64. E. Gardiner, *The Growth of the Ovum, Formation of the Polar Bodies, and the Fertilization*

in Polychaerus Caudatus (1898).

65. S. J. W. Tong, R. S. L. Ong, Mating behavior, spawning, parental care, and embryonic

development of some marine pseudocerotid flatworms (Platyhelminthes: Rhabditophora:

Polycladida) in Singapore. *Invertebr. Biol.* **139**, e12293 (2020).

66. Y. I. Zabotin, V. G. Evtugyn, Ultrastructure of spermatozoa and female copulatory organs in

preferably asexually-reproducing acoel *Convolutriloba retrogemma* (Acoelomorpha).

Zoomorphology **140**, 19–26 (2021).

67. S. J. Bourlat, A. Hejnol, Acoels. *Curr. Biol.* **19**, R279–280 (2009).

68. A. Petrov, M. Hooge, S. Tyler, Ultrastructure of sperms in Acoela (Acoelomorpha) and its

concordance with molecular systematics. *Invertebr. Biol.* **123**, 183–197 (2004).

69. O. I. Raikova, A. Falleni, J.-L. Justine, Spermiogenesis in *Paratomella*

rubra(Platyhelminthes, Acoela): Ultrastructural, immunocytochemical, cytochemical studies

and phylogenetic implications. *Acta Zool.* **78**, 295–307 (1997).

70. M. Asai, H. Miyazawa, R. Yanase, K. Inaba, H. Nakano, A new species of Acoela

possessing a middorsal appendage with a possible sensory function. *Zoolog. Sci.* **39**, 147–

156 (2022).

71. B. Åkesson, *et al.*, Fission in *Convolutriloba longifissura*: asexual reproduction in acoelous

turbellarians revisited. *Acta Zool.* **82**, 231–239 (2001).

72. P. Ax, E. Schulz, Ungeschlechtliche Fortpflanzung durch Paratomie bei acoelen

Turbellarien. *Biol. Zent. Bl.* **78**, 613–622 (1959).

73. E. du Bois-Reymond Marcus, On Turbellaria and Polygordius from the Brazilian Coast. *Bol*

Fac Fil Ci Letr Univ Sao Paulo Zool **20**, 19–53 (1955).

74. E. D. Hanson, Asexual reproduction in Acoelous Turbellaria. *Yale J. Biol. Med.* **33**, 107–111

(1960).

75. K. Ishikawa, T. Yamasu, An acoel flat worm species related closely to the species

Convolutriloba retrogemma Hendelberg and Akesson occurs in Okinawa Island, Ryukyu

Archipelago. *Zoological Science* **9**, 1281 (1992).

76. P.-Y. Henry, L. Vimond, T. Lenormand, P. Jarne, Is delayed selfing adjusted to chemical

cues of density in the freshwater snail *Physa acuta*? *Oikos* **112**, 448–455 (2006).

77. M. Milinski, Fitness consequences of selfing and outcrossing in the cestode

Schistocephalus solidus. *Integr. Comp. Biol.* **46**, 373–380 (2006).

78. A. Giannakara, S. A. Ramm, Self-fertilization, sex allocation and spermatogenesis kinetics

in the hypodermically inseminating flatworm *Macrostomum pusillum*. *J. Exp. Biol.* **220**,

1568–1577 (2017).

79. N. K. Michiels, L. J. Newman, Sex and violence in hermaphrodites. *Nature* **391**, 647–647

(1998).

80. R. E. Hulett, D. Potter, M. Srivastava, Neural architecture and regeneration in the acoel

Hofstenia miamia. Proceedings of the Royal Society B: Biological Sciences **287**, 20201198

(2020).

980