**Title:** Impact of Equilibrative Nucleoside Transporters on *Toxoplasma gondii* Infection

and Differentiation.

Gabriel Messina[1], Amber Goerner[1] Charlotte Bennett[1], Euwen Brennan[1], Vern

B. Carruthers[2], Bruno Martorelli Di Genova[1*]

[1] – Department of Microbiology and Molecular Genetics, University of Vermont,

Burlington, VT 05405, USA

[2] - Department of Microbiology and Immunology, University of Michigan School of

Medicine, Ann Arbor, MI 48109, USA

[*] – corresponding author: bruno.mdigenova@med.uvm.edu

**Abstract**

*Toxoplasma gondii* cannot synthesize purines de novo and must import them; yet, the

functional interplay among its four equilibrative nucleoside transporters (ENTs)

homologs remains unclear. We systematically deconstructed this network by combining

CRISPR-Cas9 knockouts with an auxin-inducible degron. Across all phenotypic assays,

tachyzoite replication, nucleoside-analogue sensitivity, alkaline-stress–induced

differentiation, and murine cyst formation, the Δ*TgENT2* strain was indistinguishable

from the parental line, indicating that TgENT2 is dispensable under the conditions

tested. In contrast, the double mutant Δ*TgAT1*Δ*TgENT3* exhibited delayed bradyzoite

differentiation in vitro and produced smaller brain cysts in vivo. This double deletion

triggered a ~3-fold transcriptional up-regulation of *TgENT1*, whose product we partially

localized to the plant-like vacuolar compartment (PLVAC). Conditional depletion of

TgENT1 caused complete intracellular growth arrest, PLVAC swelling, and a purine-

starvation-like transcriptomic program enriched for nucleoside phosphatases and cyclic-

nucleotide phosphodiesterases. These findings reveal a compensatory salvage pathway

in which the parasite reroutes purine acquisition through a vacuolar route when plasma-

membrane import is compromised. Although this adaptation sustains tachyzoite

proliferation, it fails during the energetically demanding transition to bradyzoites,

creating a metabolic bottleneck that impairs chronic infection. Our work reveals an

adaptable yet ultimately limited purine-import network and identifies TgENT1, along with

the vacuolar salvage axis it mediates, as a promising therapeutic target for blocking

lifelong toxoplasmosis.

## Introduction

*Toxoplasma gondii* is an obligate intracellular parasite that causes toxoplasmosis, a disease affecting approximately one-third of the global population and posing risks to immunocompromised individuals and developing fetuses. The parasite's lifecycle includes acute (tachyzoite) and chronic (bradyzoite) stages, each characterized by distinct metabolic demands [1]. A critical adaptation of *T. gondii* is its complete reliance on host-derived purines due to the absence of a *de novo* purine synthesis pathway [2]. Understanding how *T. gondii* transports and manages purine resources is crucial for deciphering its survival strategies and identifying potential therapeutic targets. Equilibrative nucleoside transporters (ENTs) are key proteins involved in this purine acquisition from host cells [3]. Here we investigate the roles of four putative *T. gondii* ENTs: TgENT1 (TGME49_288540), TgENT2 (TGME49_500147), TgENT3 (TGME49_233130), and TgAT1 (TGME49_244440) across both acute and chronic stages, aiming to elucidate their distinct and overlapping functions in parasite survival and development.

*De novo* nucleic acid synthesis and salvage pathways are fundamental for cellular growth and survival, allowing cells to adapt to varying environmental conditions [4]. Higher eukaryotes possess sophisticated mechanisms to modulate nucleoside transport, salvage, and metabolism, integrating intracellular and extracellular cues to maintain nucleotide homeostasis [5–9].

Many early branching eukaryotes have evolved complex mechanisms to adapt to fluctuating nucleoside availability [10–12]. *T. gondii* exemplifies this adaptation through its complete dependence on host-derived purines, having lost the capacity for *de novo*

purine synthesis during evolution [2,11,13]. This reliance underscores the importance of

understanding the mechanisms of purine acquisition in *T. gondii*. In host cells such as

macrophages and neurons, purine levels are tightly regulated, presenting a challenging

environment for the parasite [9,14].

The parasite employs equilibrative nucleoside transporters (TgENTs) as its primary

means of scavenging purines from the host [15]. *T. gondii* utilizes two purine salvage

pathways: the phosphorylation of adenosine by the enzyme adenosine kinase and the

phosphoribosylation of purine nucleobases by hypoxanthine-guanine

phosphoribosyltransferase (HXGPRT) [11]. These pathways enable the parasite to

salvage both adenosine and other purine nucleobases efficiently. Additionally, *T. gondii*

possesses a functional *de novo* pyrimidine biosynthesis pathway, with uracil

phosphoribosyltransferase (UPRT) facilitating pyrimidine salvage by converting uracil to

uridine monophosphate [11].

The initial identification of equilibrative nucleoside transporter (ENT) activity in *T. gondii*

was demonstrated in extracellular tachyzoites, where adenosine, inosine, and

hypoxanthine were transported into the parasite [16]. Subsequent studies identified TgAT1

via insertional mutagenesis[17]. Radioactive nucleoside incorporation assays revealed the

presence of multiple nucleoside transporters in *T. gondii*, suggesting a complex system

with at least one high-affinity adenosine transporter yet to be characterized [18]. TgAT1

remains the only TgENT studied *in vitro* using heterologous expression systems [15].

In this study, we aim to elucidate the roles of TgAT1 and three other homologs,

TgENT1, TgENT2, and TgENT3, across different developmental stages of *T. gondii*. We

hypothesized that the network of purine transporters in *T. gondii* is characterized by

functional redundancy and compensatory regulation, ensuring parasite survival under

purine stress. To test this, we employed a systematic genetic approach to deconstruct

this network by the investigation of the parasite's adaptive response to each TgENT

disruption; and assess the importance of this transporter network for the critical

developmental transition from the acute tachyzoite to the chronic bradyzoite stage. This

strategy was designed to unmask the latent functions and regulatory connections within

the purine acquisition machinery.

**Results:**

**TgENT BLAST Analysis:** We performed protein-protein BLAST or position-specific

iterated BLAST on NCBI using Human ENT-1[3] and *Plasmodium falciparum* PfENT1[12]

nucleoside transporters as query sequences to search for *T. gondii* sequences with high

similarity. Phylogenetic analysis (**Fig. 1A**) places TgAT1 and TgENT1–3 in a single,

well-supported clade with apicomplexan and mammalian ENTs. Conserved-domain

searches and AlphaFold modelling suggest that TgAT1 and TgENT1-3 harbor the

hallmark 11- ENT fold with an open central pore (**Fig. 1B,C**, and **Table 1**). Pairwise

structural comparisons to the HsENT1 crystal structure (PDB 6OB7) show pore

dimensions within the expected range for functional ENT homologues (**Fig. 1C**),

supporting their candidacy as bona fide transporters. Finally, re-analysis of the

transcriptomic data[19,20] (**Fig. 1D**) reveals distinct but overlapping expression patterns;

TgAT1 and TgENT1 are abundant throughout enteric, tachyzoite, and bradyzoite

stages, TgENT3 is moderately expressed across the life cycle, and TgENT2 peaks

during sporulation, suggesting stage-specific expression. A multiple-sequence

alignment of TgENT1, TgENT2, TgENT3, TgAT1, and other characterized ENT

homologs from different organisms is provided in Supplementary **Figures S1**.

**Nucleoside analogs growth assay:** We hypothesized that the deletion of genes

encoding nucleoside transporter homologs would negatively affect parasite growth.

However, none of the single knockout strains displayed significant differences in either

short-term or long-term growth rates compared to the parental strain (**Fig. S2**). The

growth phenotype of TgENT1 knockdown parasites is described separately below (**Fig.**

**4**).

Among the four ENT homologs in *Toxoplasma gondii*, only TgAT1 has been

biochemically characterized[15]. To determine whether deletion of TgENT2, TgENT3, or

TgAT1 alters parasite sensitivity to the toxic nucleoside analogs 9-β-D-

arabinofuranosyladenine (Ara-A)[18] and 5-fluorouracil (5-FU)[21], we grew the respective

knockout strains in 96-well plates for three days in the presence of increasing

concentrations of each analog. Infected monolayers were then fixed and stained with a

rabbit anti-*Toxoplasma* (anti-Tg) antibody. Using immunofluorescence microscopy, we

measured the area of at least 50 parasitophorous vacuoles (PVs) per condition; the

mean PV size served as a quantitative read-out of growth inhibition (**Fig. 2**).

Consistent with TgAT1's role as a purine transporter[15], ΔTgAT1 parasites showed

increased resistance to the adenosine analog Ara-A (**Fig. 2A**). Unexpectedly, ΔTgAT1

also exhibited resistance to the pyrimidine analog 5-FU (**Fig. 2B**), suggesting broader

substrate specificity than previously recognized. In contrast, the ΔTgENT2 strain did not

differ from the parental line (**Fig. 2**), likely because TgENT2 is not expressed during the

tachyzoite stage (**Fig. 1D**). Similarly, the ΔTgENT3 single knockout showed no

difference in sensitivity relative to the parental line under drug treatment (**Fig. 2**).

However, the double knockout ΔTgAT1ΔTgENT3 formed significantly larger PVs than

ΔTgAT1 under 10 μM Ara-A, indicating that deletion of TgENT3 further enhances

resistance to this analog. Conversely, ΔTgAT1ΔTgENT3 vacuoles were significantly

smaller than those of ΔTgAT1 under 5 μM 5-FU, although they remained larger than

parental vacuoles. Collectively, these observations suggest that while TgENT3 deletion

may confer increased resistance to Ara-A in the ΔTgAT1ΔTgENT3 background, it also

leads to heightened sensitivity to 5-FU.

**Role of TgENTs homologs in chronic infection:** We successfully generated knockout

strains for TgAT1, TgENT2, and TgENT3 using CRISPR/Cas9-mediated gene

disruption in the ME49 background. Multiple attempts to generate a TgENT1 knockout

strain were unsuccessful, suggesting this gene may be essential for parasite viability.

This prompted us to instead use the mini auxin-inducible degron (mAID) system for

conditional knockdown analysis (**Fig. 4**) in the RH background.

Based on published transcriptomic data[19,20] (**Fig. 1D**), TgENT1, TgENT3, and TgAT1

are constitutively expressed throughout the parasite life cycle, whereas TgENT2

expression is restricted to bradyzoite and sexual developmental stages, explaining the

lack of phenotype in tachyzoite growth assays. Given this distinct expression pattern, as

well as the unexpected lack of growth defects in single knockout strains, we next

investigated potential changes in cyst formation using an *in vitro* and *in vivo* model.

We assessed the impact of TgENT deletion on parasite differentiation by culturing

human foreskin fibroblasts (HS27) infected monolayers cultured in RPMI medium

adjusted to pH 8.2 and atmospheric $CO_2$. Monolayers were fixed after two, three, or

seven days, and differentiation rates were quantified using immunofluorescence assays

(IFA) with the lectin Dolichos Biflorus Agglutinin (DBA), which stains *T. gondii's* cyst

wall[22]. Parasites were co-stained with anti-Tg antibody, which identifies both tachyzoites

and bradyzoites, to quantify all vacuoles in the monolayers and calculate differentiation

ratios among strains. ΔTgAT1 and ΔTgAT1ΔTgENT3 parasites exhibited significantly

slower differentiation. Both ΔTgAT1 and ΔTgAT1ΔTgENT3 have delayed differentiation,

exemplified by lower number of DBA vacuoles in the population (**Fig. 3A**). Moreover,

after 3 days in pH 8.2 medium, the DBA-positive vacuoles formed by ΔTgAT1 and

ΔTgAT1ΔTgENT3 parasites are smaller than parental and complement strains (**Fig.**

**S3**). By day 7, only 88% of ΔTgAT1 and 63% of ΔTgAT1ΔTgENT3 parasites were DBA-

positive, compared to approximately 99% in the parental and ΔTgENT3 strains (**Fig.**

**3A**). As we noticed at day 3, the ΔTgAT1 and ΔTgAT1ΔTgENT3 DBA positive vacuoles

are significantly smaller at day 7 as well compared to other strains (**Fig. 3B**). While

ΔTgENT3 did not show a reduction in differentiation rate, the accentuated phenotype in

the double deleted strain suggests an interplay between TgAT1 and TgENT3 in parasite

differentiation under alkaline stress.

To assess long-term differentiation *in vivo*, male C57BL/6 mice were intraperitoneally

infected with 250 parasites. Four weeks post-infection, mice were euthanized in

accordance with Institutional Animal Care and Use Committee (IACUC) guidelines.

Brains were extracted surgically, homogenized in PBS through sequential passage

using 18-, 20-, and 22-gauge needles, fixed, and stained with DBA. Cyst numbers and

sizes were quantified for each strain, as previously done[23], (**Fig. S4**). This experiment

was independently replicated twice, with four mice per strain in each replicate. No

significant differences in cyst counts were observed among the strains (**Fig. S4A**).

These findings suggest that, despite initially impaired differentiation observed in

ΔTgAT1 and ΔTgAT1ΔTgENT3 parasites, these mutants ultimately compensate for this

deficit during chronic infection in mice. The cyst sizes for ΔTgENT3 and

ΔTgAT1ΔTgENT3 were significantly lower compared to all other strains, suggesting that

TgENT3 may have important roles during bradyzoite replication, as cyst sizes are

proportional to parasite growth (**Fig. S4B**).

To investigate whether the comparable tachyzoite growth rate (**Fig. S2**) observed in

ΔTgAT1ΔTgENT3 was due to compensatory expression of another TgENT family

member, we performed quantitative PCR (qPCR) analyses. We compared the

expression levels of the remaining TgENT transporters, TgENT1 and TgENT2, across

tachyzoites derived from ΔTgAT1ΔTgENT3, single knockout strains, and the parental

strain. qPCR analysis revealed a ~3-fold upregulation of TgENT1 transcripts specifically

in ΔTgAT1ΔTgENT3 parasites compared to parental controls ($p < 0.05$), while TgENT2

expression remained unchanged (**Fig. 4A**). Notably, single knockout strains, ΔTgAT1

and ΔTgENT3, showed no compensatory changes in TgENT1 or TgENT2 (**Fig. 4A**).

These data indicate that the absence of both TgAT1 and TgENT3 may trigger

compensatory upregulation of TgENT1, maintaining tachyzoite growth rates similar to

single knockout strains.

**TgENT1 conditional deletion results in complete growth arrest:** TgENT1 has a markedly negative genome-wide CRISPR phenotype score of −3.68[24]. On this scale, increasingly negative values typically denote progressively stronger fitness costs, underscoring TgENT1's critical importance for parasite survival (**Table 1**). To investigate the role of TgENT1, we developed a parasite line, TgENT1-mAID-HA, in RH backgrond with regulated TgENT1 expression via 3-indoleacetic acid (IAA) treatment [25]. We assessed the impact of TgENT1 depletion on parasite growth by culturing the parasites with IAA for 48 hours. The treated parasites exhibited reduced replication when invading new HS27 monolayers (**Fig. 4B and C**), indicating that TgENT1 expression is crucial for normal growth and replication. Additionally, a long-term growth defect was observed over nine days of continuous culture, as evidenced by the formation of smaller and fewer plaques (**Fig. 4D**).

**TgENT1 partially colocalizes to the PLVAC:** Using the TgENT1-mAID-HA construct, we were able to visualize the association of TgENT1 within *T. gondii* parasites. When cultured with 500 µM indole-3-acetic acid (IAA), we observed the disappearance of the TgENT1 signal, confirming the system functionality (**Fig. 5**). Surprisingly, TgENT1 localization differed markedly from typical plasma membrane-associated ENTs. Instead, HA-TgENT1 displayed a distinctive punctate pattern within the parasite cytoplasm, suggesting it is localized within intracellular organelles (**Fig. 5**).

To pinpoint the intracellular compartment that hosts TgENT1, we performed immunofluorescence assays (IFA) on parasites expressing HA-tagged TgENT1 alongside well-validated organelle markers. Co-staining was carried out with antibodies

against cathepsin protease L (TgCPL) and the chloroquine-resistance transporter

(TgCRT), both definitive markers of the plant-like vacuole (PLVAC)[26], as well as the

propeptide of microneme protein 2 (TgProM2AP), which labels the endosome-like

compartment [27], and an anti-apicoplast (TgAtrx1) antibody[28]. Here, M(X) represents

Manders' colocalization coefficients which quantify the fraction of antibody X's

fluorescence that shares the same pixels as the HA signal (1.0 = complete overlap; 0 =

none). Across all markers, in control conditions, the TgENT1 HA-tagged protein shows

partial colocalization with PLVAC markers: TgCPL (Pearson's r =0.84; M(CPL) =0.90)

and TgCRT (Pearson's r =0.79; M(CRT) =0.54), weak overlap by the microneme protein

TgProM2AP (Pearson's r =0.61; M(M2AP) =0.09), and virtually no overlap with

TgAtrx1(Pearson's r =0.10; M(Atrx1) =0.05). After IAA-induced depletion, Manders'

coefficients collapse to near background (TgCPL 0.08; TgCRT 0.02; TgProM2AP 0.09;

TgAtrx1 0.00) and Pearson's r values likewise fall (CPL 0.21; CRT 0.03; ProM2AP 0.25;

Atrx1 0.05), indicating loss of specific HA signal and that the residual fluorescence no

longer maps to these organelles. Manders and Pearson's colocalization coefficients are

based on three different biological experiments. Together, these results identify the

PLVAC as the principal compartment for TgENT1.

**PLVAC measurement:** Given the partial colocalization of TgENT1 and TgCRT and

TgCPL (**Fig. 5**), we evaluated PLVAC morphology following TgENT1 depletion by

performing knockdown experiments in replicating parasites 36 hours post-infection.

Parasites were mechanically liberated and subsequently inoculated onto fresh HS27

monolayers for 30 minutes before fixation and staining. Importantly, TgENT1

knockdown parasites retained their ability to invade host cells efficiently.

Immunofluorescence assays were employed to measure PLVAC dimensions using an

anti-TgCPL antibody (**Fig. 6**). Area from 50 TgCPL puncta per experimental condition

were measured in triplicate, and comparative statistical analyses were performed using

t-test. Consistent with our hypothesis, TgENT1 depletion resulted in measurable

swelling of the PLVAC, further implicating TgENT1 as important for maintaining PLVAC

homeostasis.

**Transcriptomic analysis of TgENT1 knockdown:** Transcriptomic profiling following

conditional knockdown of TgENT1 revealed significant alterations in genes associated

with purine metabolism, reinforcing TgENT1's proposed function as a nucleoside

transporter (**Fig. 7**). Principal-component analysis (**Fig. 7A**) cleanly separates IAA-

treated and control tachyzoites along PC1 and shows a further time-dependent shift on

PC2, underscoring a rapid and progressive transcriptional response to TgENT1 loss. An

accompanying heat-map of (**Fig. 7B**) highlights the 50 most variable transcripts.

Volcano plots of differentially expressed genes (**Fig. 7C**) display a broader and more

pronounced response by 48 h, with the number and magnitude of regulated transcript

expanding and purine-stress markers (e.g., nucleoside phosphatases and

phosphodiesterases) among the most strongly up-regulated. For example,

TGGT1_225290 (GDA1/CD39 nucleoside phosphatase) demonstrated progressive

upregulation from day 1 ($log_2$ fold change = 2.98) to day 2 ($log_2$ fold change = 6.21).

Similarly, TGGT1_259960 (putative nucleoside-diphosphatase) exhibited strong

induction from day 1 ($log_2$ fold change = 2.81) to day 2 ($log_2$ fold change = 5.92),

suggesting a sustained compensatory effort to restore purine homeostasis when

nucleoside uptake is impaired. Importantly, other TgENT family members showed no

significant expression changes.

In total, 208 transcripts were already altered at 24 h and this number more than doubled

by 48 h, with 96 up- and 58 down-regulated genes shared between both time points

(**Fig. 7D**), defining a core TgENT1-dependent regulon. Among the induced genes were

several 3'5'-cyclic nucleotide phosphodiesterases (PDEs), notably PDE3

(TGGT1_233065; $\log_2$ fold change = 1.30 at day 1, significantly increased to 3.88 at day

2), and PDE17 (TGGT1_257945; $\log_2$ fold change = 2.92), suggesting heightened cyclic

nucleotide turnover and signaling adaptation [29].

KEGG pathway analyses (**Fig. 6E**) revealed significant differential expression in

pathways related to purine metabolism, nitrogen compound metabolism, and

oxidoreductase activities. The enrichment of the purine metabolism pathway further

intensified at day 2, according to KEGG pathway analysis (**Fig. 7E**) (fold enrichment =

2.39, p-value = 0.0008), featuring upregulated genes like TGGT1_230555[30]

(adenylate/guanylate cyclase; $\log_2$ fold change = 3.84) and further emphasizing

signaling changes upon TgENT1 knockdown. Additionally, TGGT1_290970[31] (8-amino-

7-oxononanoate synthase), involved in biotin biosynthesis, was substantially induced

from day 1 ($\log_2$ fold change = 2.57) to day 2 ($\log_2$ fold change = 5.22), suggesting

enhanced biotin-dependent metabolic processes.

Collectively, these transcriptomic findings suggest TgENT1's central role in purine

transport and homeostasis within *T. gondii*. The extensive and progressive

compensatory responses triggered by TgENT1 knockdown, including purine salvage,

metabolic reprogramming, and signaling adaptations, provide new insights into the

parasite's ability to adapt to disrupted nucleoside transport.

**Discussion**

In this study, we have expanded the model of purine acquisition in *Toxoplasma gondii*,

moving beyond a simple linear import pathway to reveal a dynamic, regulated network.

HyperLOPIT spatial-proteomics has previously assigned both TgAT1 and TgENT3 to

the parasite plasma membrane[36] **(Table 1)**. Although our experiments did not verify this

localization directly, the functional data reveal a broader principle of *metabolic*

*resilience*: *Toxoplasma gondii* can rapidly re-wire its purine-uptake network to

compensate for the loss of TgAT1 and TgENT3. Such plasticity is indispensable for an

organism that is purine-auxotrophic and therefore wholly dependent on the host for

nucleoside acquisition [32]. Importantly, we also demonstrate that this adaptive capacity is

limited during differentiation (**Fig. 3**). Understanding the mechanics, function, and

ultimate failure point of this network provides unprecedented insight into the metabolic

strategies governing the parasite's lifecycle.

Our data show that deletion of the TgAT1 and TgENT3 triggers specific transcriptional

upregulation of TgENT1 (**Fig. 4A**), a transporter that our localization studies showed to

partially localize in the Plant-Like Vacuolar Compartment (PLVAC) (**Fig. 5**). These

findings support a model where the parasite mobilizes an alternative salvage pathway.

This pathway might rely on acquiring purines from the degradation of host-derived

nucleic acids within the acidic, digestive environment of the PLVAC[26] or from the

recycling of its own nucleic acids, a function consistent with the PLVAC's role in

trafficking and catabolism [26]. The adaptive function of the PLVAC and TgENT1 creates

a crucial lifeline that sustains a baseline of purine homeostasis.

While this adaptive axis provides metabolic flexibility for tachyzoites, its capacity is

finite, and its functional limit is most profoundly revealed during the critical

developmental transition to the chronic stage. The failure of the ΔTgAT1ΔTgENT3 strain

to develop as bradyzoites *in vitro* (**Fig. 3A and C,** and **S3**) likely stems from two

interconnected deficiencies: a substrate and a signaling deficit. The substrate deficit

may deplete critical precursors for cyst formation, including the sugar-nucleotide donors

uridine diphosphate N-acetyl-D-glucosamine (UDP-GlcNAc) and uridine diphosphate N-

acetyl-D-galactosamine (UDP-GalNAc). These activated sugars supply N-

acetylglucosamine and N-acetylgalactosamine residues that decorate cyst-wall

glycoconjugates and are important for proper wall assembly [33]. More critically, however,

this purine starvation likely precipitates a failure in cellular signaling. Purine salvage is

the sole source of the adenosine required for ATP synthesis. ATP, in turn, is the

exclusive substrate for adenylyl cyclases that generate cyclic AMP (cAMP), a master

second messenger in *T. gondii*. The cAMP signaling pathway is known to be an

important regulator of stage conversion, with fluctuations in cAMP levels acting as a key

signal for maintaining the tachyzoite state or initiating differentiation into bradyzoites [34].

Therefore, we propose that the ΔTgAT1ΔTgENT3 strain, while receiving the external

cues to differentiate (e.g., alkaline stress), is metabolically incapable of mounting the

appropriate cAMP signaling response required to execute this complex developmental

program. Interestingly, although ΔTgAT1ΔTgENT3 mutants form smaller cysts than

parental strains in mice, the total cyst burden remains unchanged, suggesting that the

30-day *in vivo* period allows the parasites to overcome the differentiation delay

observed *in vitro* (**Fig. S4**).

Our study illuminates a clear transcriptional and phenotypic signature of nucleoside

starvation in the absence of TgENT1, and several open questions make the next

experimental steps especially compelling. First, the robust up-regulation of TgENT1 is

currently inferred to reflect enhanced host-nucleoside scavenging; confirming this

hypothesis with direct radiolabeled-substrate uptake assays will test our model and

establish quantitative benchmarks for future transporter engineering. Second, the

nucleoside-depletion phenotype reported for Δgra14 parasites, which are defective in

parasite endocytosis of material from the cytosol of infected cells, suggests that the

PLVAC has a central role in nucleoside salvage and invites a systematic dissection of

each vacuolar effector in that pathway [35]. Finally, untargeted metabolomics coupled to

parallel measurements of high-energy substrates such as ATP and GTP and their

signaling derivatives would provide a systems-level view of how disrupted uptake

reverberates through the parasite's metabolic and regulatory networks.

Ultimately, by deconstructing this purine acquisition network, we have revealed a

dynamic interplay between metabolic adaptation and developmental signaling. The

biological consequence of this network's failure is unambiguous. By uncovering a

scenario where the parasite's adaptive capacity breaks down at the nexus of

metabolism and developmental signaling, we have exposed a significant vulnerability.

This metabolic bottleneck, where purine insufficiency hypothetically precipitates a

signaling failure that blocks the establishment of chronic infection, represents a highly

promising and conceptually novel target for therapeutic intervention. Further

investigation offers the potential to develop strategies aimed specifically at disrupting

the lifelong persistence that is the clinical hallmark of toxoplasmosis.

**References:**

1. Dubey JP, Jones JL. Toxoplasma gondii infection in humans and animals in the United States. Int J Parasitol. England; 2008 Sept;38(11):1257–1278. PMID: 18508057

2. Krishnan A, Kloehn J, Lunghi M, Chiappino-Pepe A, Waldman BS, Nicolas D, Varesio E, Hehl A, Lourido S, Hatzimanikatis V, Soldati-Favre D. Functional and Computational Genomics Reveal Unprecedented Flexibility in Stage-Specific Toxoplasma Metabolism. Cell Host Microbe. United States; 2020 Feb 12;27(2):290-306.e11. PMID: 31991093

3. Wright NJ, Lee SY. Structures of human ENT1 in complex with adenosine reuptake inhibitors. Nature Structural & Molecular Biology. 2019 July 1;26(7):599–606.

4. Pareek V, Pedley AM, Benkovic SJ. Human de novo purine biosynthesis. Crit Rev Biochem Mol Biol. 2021 Feb;56(1):1–16. PMCID: PMC7869020

5. Verrier F, An S, Ferrie AM, Sun H, Kyoung M, Deng H, Fang Y, Benkovic SJ. GPCRs regulate the assembly of a multienzyme complex for purine biosynthesis. Nat Chem Biol. 2011 Oct 23;7(12):909–915. PMID: 22020552

6. Huang Z, Xie N, Illes P, Di Virgilio F, Ulrich H, Semyanov A, Verkhratsky A, Sperlagh B, Yu SG, Huang C, Tang Y. From purines to purinergic signalling: molecular functions and human diseases. Signal Transduct Target Ther. 2021 Apr 28;6(1):162. PMID: 33907179

7. Mason S. Signaling for purines. Nature Structural & Molecular Biology. 2011 Dec 1;18(12):1322–1322.

8. Jain S, Jacobson KA. Purinergic signaling in diabetes and metabolism. Biochem Pharmacol. 2021 May;187:114393. PMID: 33359363

9. Soler C, García-Manteiga J, Valdés R, Xaus J, Comalada M, Casado FJ, Pastor-Anglada M, Celada A, Felipe A. Macrophages require different nucleoside transport systems for proliferation and activation.

10. Carter NS, Yates PA, Gessford SK, Galagan SR, Landfear SM, Ullman B. Adaptive responses to purine starvation in Leishmania donovani. Mol Microbiol. 2010 Oct;78(1):92–107. PMCID: PMC2964060

11. Fox BA, Bzik DJ. Biochemistry and metabolism of Toxoplasma gondii: purine and pyrimidine acquisition in Toxoplasma gondii and other Apicomplexa. 2020 Jan 1;397–449.

12. Wang C, Yu L, Zhang J, Zhou Y, Sun B, Xiao Q, Zhang M, Liu H, Li J, Li J, Luo Y, Xu J, Lian Z, Lin J, Wang X, Zhang P, Guo L, Ren R, Deng D. Structural basis of the substrate recognition and inhibition mechanism of Plasmodium falciparum

nucleoside transporter PfENT1. Nat Commun. England; 2023 Mar 28;14(1):1727.
PMCID: PMC10050424

13.   Kloehn J, Hammoudi PM, Soldati-Favre D. Metabolite salvage and restriction
during infection — a tug of war between Toxoplasma gondii and its host. Current
Opinion in Biotechnology. 2021 Apr 1;68:104–114. PMID: 33202353

14.   Dunwiddie TV, Masino SA. The Role and Regulation of Adenosine in the Central
Nervous System. Annu Rev Neurosci. Annual Reviews; 2001 Mar 1;24(1):31–55.

15.   Campagnaro GD, Elati HAA, Balaska S, Martin Abril ME, Natto MJ, Hulpia F, Lee
K, Sheiner L, Van Calenbergh S, de Koning HP. A Toxoplasma gondii Oxopurine
Transporter Binds Nucleobases and Nucleosides Using Different Binding Modes.
International Journal of Molecular Sciences. 2022 Jan;23(2):710. PMID: 35054895

16.   Conrad Schwab J, Afifi Afifi M, Pizzorno G, Handschumacher RE, Joiner KA.
Toxoplasma gondii tachyzoites possess an unusual plasma membrane adenosine
transporter. Molecular and Biochemical Parasitology. 1995 Mar 1;70(1):59–69.
PMID: 7637715

17.   Chiang CW, Carter N, Sullivan WJ, Donald RGK, Roos DS, Naguib FNM, Kouni
MH el, Ullman B, Wilson CM. The Adenosine Transporter of Toxoplasma gondii:
IDENTIFICATION BY INSERTIONAL MUTAGENESIS, CLONING, AND
RECOMBINANT EXPRESSION *. Journal of Biological Chemistry. Elsevier; 1999
Dec 3;274(49):35255–35261. PMID: 10575012

18.   De Koning HP, Al-Salabi MI, Cohen AM, Coombs GH, Wastling JM. Identification
and characterisation of high affinity nucleoside and nucleobase transporters in
Toxoplasma gondii. International Journal for Parasitology. 2003 July 30;33(8):821–
831.

19.   Hehl AB, Basso WU, Lippuner C, Ramakrishnan C, Okoniewski M, Walker RA,
Grigg ME, Smith NC, Deplazes P. Asexual expansion of Toxoplasma gondii
merozoites is distinct from tachyzoites and entails expression of non-overlapping
gene families to attach, invade, and  replicate within feline enterocytes. BMC
Genomics. England; 2015 Feb 13;16(1):66. PMCID: PMC4340605

20.   Ramakrishnan C, Maier S, Walker RA, Rehrauer H, Joekel DE, Winiger RR, Basso
WU, Grigg ME, Hehl AB, Deplazes P, Smith NC. An experimental genetically
attenuated live vaccine to prevent transmission of Toxoplasma gondii by cats. Sci
Rep. England; 2019 Feb 6;9(1):1474. PMCID: PMC6365665

21.   Elati HAA, Goerner AL, Martorelli Di Genova B, Sheiner L, de Koning HP.
Pyrimidine salvage in Toxoplasma gondii as a target for new treatment. Front Cell
Infect Microbiol. 2023;13:1320160. PMCID: PMC10755004

22. Zhang YW, Halonen SK, Ma YF, Wittner M, Weiss LM. Initial characterization of
CST1, a Toxoplasma gondii cyst wall glycoprotein. Infect Immun. United States;
2001 Jan;69(1):501–507. PMCID: PMC97909

23. Watts E, Zhao Y, Dhara A, Eller B, Patwardhan A, Sinai AP. Novel Approaches
Reveal that Toxoplasma gondii Bradyzoites within Tissue Cysts Are Dynamic and
Replicating Entities In Vivo. mBio. United States; 2015 Sept 8;6(5):e01155-01115.
PMCID: PMC4600105

24. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, Thiru P, Saeij
JPJ, Carruthers VB, Niles JC, Lourido S. A Genome-wide CRISPR Screen in
Toxoplasma Identifies Essential Apicomplexan Genes. Cell. 2016 Sept
8;166(6):1423-1435.e12. PMID: 27594426

25. Brown KM, Long S, Sibley LD. Conditional Knockdown of Proteins Using Auxin-
inducible Degron (AID) Fusions in Toxoplasma gondii. Bio Protoc. 2018 Feb
20;8(4):e2728. PMCID: PMC5890294

26. Stasic AJ, Moreno SNJ, Carruthers VB, Dou Z. The Toxoplasma plant-like vacuolar
compartment (PLVAC). J Eukaryot Microbiol. United States; 2022
Nov;69(6):e12951. PMCID: PMC10576567

27. Gaji RY, Flammer HP, Carruthers VB. Forward targeting of Toxoplasma gondii
proproteins to the micronemes involves conserved aliphatic amino acids. Traffic.
England; 2011 July;12(7):840–853. PMCID: PMC3115430

28. DeRocher AE, Coppens I, Karnataki A, Gilbert LA, Rome ME, Feagin JE, Bradley
PJ, Parsons M. A thioredoxin family protein of the apicoplast periphery identifies
abundant candidate transport vesicles in Toxoplasma gondii. Eukaryot Cell. United
States; 2008 Sept;7(9):1518–1529. PMCID: PMC2547066

29. Moss WJ, Patterson CE, Jochmans AK, Brown KM. Functional Analysis of the
Expanded Phosphodiesterase Gene Family in Toxoplasma gondii Tachyzoites.
mSphere. United States; 2022 Feb 23;7(1):e0079321. PMCID: PMC8809380

30. Brown KM, Sibley LD. Essential cGMP Signaling in Toxoplasma Is Initiated by a
Hybrid P-Type ATPase-Guanylate Cyclase. Cell Host Microbe. United States; 2018
Dec 12;24(6):804-816.e6. PMCID: PMC6292738

31. Asady B, Dick CF, Ehrenman K, Sahu T, Romano JD, Coppens I. A single Na+-Pi
cotransporter in Toxoplasma plays key roles in phosphate import and control of
parasite osmoregulation. PLoS Pathog. United States; 2020 Dec;16(12):e1009067.
PMCID: PMC7817038

32. Schwartzman JD, Pfefferkorn ER. Toxoplasma gondii: Purine synthesis and
salvage in mutant host cells and parasites. Experimental Parasitology. 1982 Feb
1;53(1):77–86.

33. Caffaro CE, Koshy AA, Liu L, Zeiner GM, Hirschberg CB, Boothroyd JC. A
nucleotide sugar transporter involved in glycosylation of the Toxoplasma tissue
cyst wall is required for efficient persistence of bradyzoites. PLoS Pathog. United
States; 2013;9(5):e1003331. PMCID: PMC3642066

34. Sugi T, Ma YF, Tomita T, Murakoshi F, Eaton MS, Yakubu R, Han B, Tu V, Kato K,
Kawazu SI, Gupta N, Suvorova ES, White MW, Kim K, Weiss LM. Toxoplasma
gondii Cyclic AMP-Dependent Protein Kinase Subunit 3 Is Involved in the Switch
from Tachyzoite to Bradyzoite Development. mBio. American Society for
Microbiology; 2016 May 31;7(3):10.1128/mbio.00755-16.

35. Rimple PA, Olafsson EB, Markus BM, Wang F, Augusto L, Lourido S, Carruthers
VB. Metabolic adaptability and nutrient scavenging in Toxoplasma gondii: insights
from ingestion pathway-deficient mutants. mSphere. United States; 2025 Apr
29;10(4):e0101124. PMCID: PMC12039266

36. Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, Gupta V, Tromer
E, Mourier T, Stevens TJ, Breckels LM, Pain A, Lilley KS, Waller RF. A
Comprehensive Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT
Provides Spatial Context for Protein Functions. Cell Host & Microbe. 2020 Nov
11;28(5):752-766.e9.

37. Tobin Magle C, Pittman KJ, Moser LA, Boldon KM, Knoll LJ. A toxoplasma patatin-
like protein changes localization and alters the cytokine response during
toxoplasmic encephalitis. Infect Immun. United States; 2014 Feb;82(2):618–625.
PMCID: PMC3911373

**Figure Legends**

491 **Table 1 – BLAST results for *T. gondii* sequences with high similarity with Human**

492 **ENT1 and *Plasmodium falciparum* ENT1.**

| Gene ID | Name | Localization by hyperLOPIT[36] | PFam ID | PFam Description | Phenotype Score[24] |
|---|---|---|---|---|---|
| **TGME49_233130** | TgENT3 | PM integral, Golgi | PF01733 | Equilibrative nucleoside transporter | 0.04 |
| **TGME49_244440** | TgAT1 | PM integral, Golgi | PF01733 | Equilibrative nucleoside transporter | 1.02 |
| **TGME49_288540** | TgENT1 | None | PF01733 | Equilibrative nucleoside transporter | -3.68 |
| **TGME49_500147** | TgENT2 | None | PF01733 | Equilibrative nucleoside transporter | 0.47 |

**Figure 1 – Figure 1. Evolutionary, structural, and transcriptional overview of the**

***Toxoplasma gondii* equilibrative nucleoside transporter (TgENT) family.**

**(A)** Phylogenetic relationships: Full-length protein sequences of the four *T. gondii* ENTs

(TgAT1, TgENT1–3) were aligned with representative ENT homologs from

apicomplexan, fungal, plant, trypanosomatid, and mammalian lineages (CpAT,

*Cryptosporidium parvum*; PfENT1, *Plasmodium falciparum*; ScFUN26, *Saccharomyces*

*cerevisiae*; AtENT1, *Arabidopsis thaliana*; NNT, *Naegleria* sp.; LmNnt1-1, *Leishmania*

*major*; HsENT1/2, MmENT1, human and mouse ENT1/2; HsCNT1/2, human

concentrative nucleoside transporters—used as the out-group). MUSCLE-aligned

sequences were subjected to neighbor-joining analysis; node labels give bootstrap

support from 1,000 replicates (values < 30% omitted). Scale bar, 0.5 substitutions per

site.

**(B)** Predicted tertiary structures: AlphaFold models of the four TgENT paralogues are

shown in ribbon representation (top-view, upper row; side-view, lower row). Each

adopts the canonical 11–13 trans-membrane helix bundle typical of ENT family

members, with the central permeation pore clearly visible in the side view. Colors match

the gene names in panel A.

**(C)** Quantitative structural comparison to human ENT1 (PDB 6OB7).

Pore dimensions were measured along the long and short axes (Å) for each TgENT

model and for HsENT1, and Cα-root-mean-square deviations (RMSDs) were calculated

after SuperPose alignment to 6OB7. Heat-map shading reflects magnitude (scale bar,

right). Cartoon underneath illustrates the pore-width measurements on the HsENT1

template.

**(D)** Stage-specific expression of TgENT transcripts.

Heat-map of log2-transformed TPM values (+1) extracted from the ToxoDB

developmental transcriptome. EES 1–5, enteroepithelial stages in cat epithelium;

sporulating oocysts; tachyzoites; tissue cyst bradyzoites; unsporulated oocysts [19,20].

TgAT1 is highest in enteroepithilial and early tachyzoite stages, TgENT1 is broadly

expressed, TgENT2 peaks during sporulation, and TgENT3 shows moderate

expression throughout.

**Figure 2. Deletion of TgAT1 and TgENT3 alters *Toxoplasma* sensitivity to toxic**

**nucleoside analogs.** Parental ME49 parasites and the indicated mutants were allowed

to replicate for 72 h in HS27 monolayers in the presence of increasing concentrations of

**(A)** the adenosine analog 9-β-D-arabinofuranosyl-adenine (Ara-A) or **(B)** the pyrimidine

analog 5-fluorouracil (5-FU) Cultures were fixed, stained with anti-*Toxoplasma* (anti-Tg)

antibody, and the mean area of ≥50 parasitophorous vacuoles (PVs) per condition was

measured by automated image analysis. Growth inhibition (% ± SEM) was calculated

relative to drug-free controls and fitted to a four-parameter logistic curve (GraphPad

Prism). Data are from three biological replicates; each performed in technical triplicate.

Panel layout: For each drug, individual dose–response curves are shown for the

parental line, single knockouts (ΔTgENT3, ΔTgENT2, ΔTgAT1), the double knockout

(ΔTgAT1ΔTgENT3), and the corresponding epitope-tagged complementation lines

(ΔTgENT3::TgENT3-Ty1, ΔTgAT1::TgAT1-myc). The right-most graph in each row

overlays all strains for direct comparison (color key, inset).

**Figure 3 – Simultaneous deletion of TgENT3 and TgAT1 delays differentiation. (A)**

Percentage of differentiation of the parental line, single knockouts (ΔTgENT3,

ΔTgENT2, ΔTgAT1), the double knockout (ΔTgAT1ΔTgENT3), and the corresponding

epitope-tagged complementation lines TgENT3-Ty1 (ΔTgENT3::TgENT3-Ty1) and

TgAT1-myc (ΔTgAT1::TgAT1-myc) at day 2, 3, and 7 post invasion of HS27 monolayers

cultured in alkaline stress conditions (pH 8.1 and atmospheric $CO_2$). Parasites were

stained with immunofluorescent antibodies DAPI, anti-Tg, and DBA. Bars show the

percentage of DBA positive vacuoles (thresholds set based on parental line intensity)

over total vacuoles (anti-Tg). Slides were imaged with BioTEK Cytation 7 and quantified

using the Gen5 software (≥100 vacuoles were counted per replicate, n = 3 biological

replicates). Statistical significance was assessed using one-way ANOVA, (****P <

0.0001, ns = non-significant). **(B)** Average sizes of each cyst (DBA positive vacuole) on

day 7 post invasion.  **(C)** Representative images of vacuoles from each parasite line.

Scale bar, 10 µm

**Figure 4. TgENT1 is transcriptionally upregulated in ΔTgAT1ΔTgENT3 parasites**

**and is indispensable for intracellular replication and plaque formation.**

**(A)** qRT-PCR analysis of compensatory TgENT expression. Total RNA from tachyzoites

of the indicated mutant lines was analyzed by qRT-PCR with primers specific for

**TgENT1** and **TgENT2**. Bars show $2^{\Delta\Delta Ct}$ values (mean ± SEM, n = 3 biological

replicates) normalized to the parental line (dotted line = 1). Only the double knockout

(ΔTgAT1ΔTgENT3, open bars) exhibits a ~3-fold induction of TgENT1 (one-way

ANOVA with ***P < 0.001); TgENT2 is unchanged (ns).

**(B)** Acute intracellular growth after TgENT1 depletion. TgENT1-mAID-HA parasites

were pre-treated for 48 h with vehicle or 500 µM indole-3-acetic acid (+IAA) to trigger

auxin-inducible degradation, then allowed to invade fresh HS27 monolayers for a further

48 h under the same conditions. Stacked bars depict the distribution of vacuoles

containing 1, 2, 4, 8, 16, 32, 64 or >64 parasites (≥50 vacuoles counted per replicate; 3

replicates).

**(C)** Plaque-formation assay. Wells were inoculated with 50 tachyzoites of the indicated

strains and cultured for 9 days ± 500 µM IAA before fixation and crystal-violet staining.

Representative wells are shown.  **(D)** Quantification of plaque area. Individual plaque

areas from TgENT1-mAID-HA wells (vehicle vs +IAA) are plotted (median ± inter-

quartile range; n = 15–20 plaques). Student t-test, ****P < 0.0001. **(E)** Quantification of

plaque number. Mean ± SEM plaque counts per well from three independent

experiments. T-test, ****, P < 0.0001.

**Figure 5. TgENT1 resides in the PLVAC and is eliminated by auxin-induced**

**degradation.** Representative immunofluorescence images of TgENT1-mAID-HA

tachyzoites pre-treated for 48 h with vehicle or 500 µM indole-3-acetic acid (+IAA), then

grown in fresh HS27 monolayers for a further 48 h under the same conditions. Parasites

were fixed and stained with anti-HA (second column) plus one of the indicated

compartment markers (third column): *T. gondii* cathepsin protease L (TgCPL) or *T.*

*gondii* chloroquine-resistance transporter (TgCRT) for the PLVAC, TgAtrx1 for the

apicoplast, and TgProM2AP for the endosome-like/microneme compartment. Columns

show (left to right) phase-contrast, anti-HA, co-stain, DAPI, and the merged overlay

(anti-HA = green, co-stain = magenta, DAPI = blue). Under vehicle conditions TgENT1-

HA displays a punctate pattern that co-localizes with PLVAC markers TgCPL and

TgCRT, but only weakly with TgProM2AP and not with TgAtrx1, consistent with

predominant residence in PLVAC. Upon IAA treatment the HA signal is virtually

abolished, confirming efficient auxin-inducible degradation. Scale bar, 10 µm.

**Figure 6. Loss of TgENT1 causes swelling of the PLVAC. (A).** TgENT1-mAID-HA

parasites were grown for 48 h post-infection, mechanically released, and allowed to

reinvade new HS27monolayers for 30 minutes in the continued presence of vehicle or

500 µM IAA. Cells were fixed and co-stained with anti-HA (TgENT1, green) and anti-

TgCPL (PLVAC marker, magenta); nuclei were counter-stained with DAPI (blue). TgENT1 is readily detected in the central PLVAC punctum under vehicle conditions but is efficiently depleted after auxin treatment, coincident with an enlarged TgCPL-positive compartment. Scale bar, 10 μm.

**(B) Quantification of PLVAC size.** Area of 50 individual CPL-positive structures per condition (three independent experiments) was measured in FIJI. Box-and-whisker plots show median, inter-quartile range, and 5th–95th percentiles. Auxin-induced TgENT1 depletion increased PLVAC area nearly five-fold, student t-test, **** $P < 0.0001$, suggesting that TgENT1 activity is required to maintain normal vacuolar homeostasis.

**Figure 7. Transcriptomic response to acute TgENT1 depletion reveals a progressive, purine-centric stress program.** TgENT1-mAID-HA tachyzoites (three biological replicates) were pre-treated for 48 h with vehicle or 500 μM IAA, then allowed to invade fresh HS27 monolayers under the same conditions and harvested 24 h (Day 1) or 48 h (Day 2) after invasion. Poly-(A) RNA was sequenced (Illumina NovaSeq; 150 bp PE), reads were mapped to the GT1 genome, and differential expression was analyzed with DESeq2 ($|\log_2 FC| \geq 1$; FDR < 0.01).

**(A) Principal-component analysis (PCA).** PC1 (40%) cleanly separates treated from control samples, whereas PC2 (28%) differentiates time points, indicating a robust, time-dependent transcriptional shift upon TgENT1 knockdown.

**(B) Heat-map of the 50 most variable genes.** Z-scored expression values highlight a concerted up-regulation of purine- and nitrogen-metabolism genes (reds) and down-regulation of cytoskeletal / motility genes (blues) that intensifies from Day 1 to Day 2.

**(C) Volcano plots of differentially expressed genes.** Pink, up-regulated; blue, down-regulated; grey, non-significant. The magnitude and number of responsive genes expand markedly by Day 2. The complete list of genes and fold change is presented of **supplemental table 4**.

**(D) Overlap of DE genes between time points.** Venn diagrams show 96 genes up-regulated at both Day 1 and Day 2, and 58 genes consistently down-regulated, underscoring a core TgENT1-dependent regulon that is sustained and amplified over time.

**(E) Functional enrichment analysis.** Dot-plot of the top Gene Ontology / KEGG terms (size = gene count; color = $-\log_{10}$FDR). Up-regulated sets are dominated by **purine metabolism**, **nitrogen-compound metabolism**, and **cyano-/taurine-derivative pathways**, whereas down-regulated sets are enriched for **microtubule-based movement** and related processes, consistent with a shift from proliferation to metabolic stress adaptation.

**Supplemental material:**

**Supplemental Figure 1: Multiple sequence alignment of equilibrative nucleoside transporter (ENT/AT) homologs highlighting conservation across apicomplexan,**

**kinetoplastid, and human sequences, with human ENT1 transmembrane topology**

**reference.**

Shown are *Toxoplasma gondii* TgAT1 (TGME49_244440), TgENT1 (TGME49_288540),

TgENT2 (TGME49_500147), TgENT3 (TGME49_233130); *Plasmodium falciparum*

PfENT1; *Cryptosporidium parvum* CpNT1; *Leishmania major* LmNT1; *Saccharomyces*

*cerevisiae* FUN26; *Arabidopsis thaliana* ENT1; *Naegleria fowleri* NT1-like; and *Mus*

*musculus* ENT1, and human ENT1. Sequence labels list protein shorthand (species in

parentheses omitted for concision here) followed by aligned residue span (e.g., "/1–

696"). Sequences were aligned with MAFFT L-INS-i; no manual internal residue edits

were introduced (only removal of obvious trailing low-complexity overhangs where

noted). Amino acids are colored by physicochemical class (ClustalX scheme) to

facilitate rapid visual inspection of conserved motifs. Histogram bars beneath each

block depict position-specific conservation (gold = high conservation / strongly similar

physicochemical class; gray = variable) and gap frequency (black). Black horizontal

rectangles above the alignment mark the experimentally validated transmembrane

helices of human ENT1 (SLC29A1), provided as a structural topology reference against

which parasite, plant, protozoan, and kinetoplastid homologs can be compared.

**Supplemental Figure 2: Intracellular replication (parasites per vacuole) and multi-**

**cycle lytic growth (plaque numbers) of parental and ENT mutant strains.**

**(A)** Distribution of vacuole sizes (stacked categories: 2, 4, 8, 16, 32, 64, 128

parasites/vacuole) at 2 days (left) and 3 days (right) post-infection under standard

tachyzoite conditions for parental ME49Δku80, ΔTgAT1, complemented

ΔTgAT1::TgAT1-myc, ΔTgENT2, ΔTgENT3, complemented ΔTgENT3::TgENT3-ty, and

the double mutant ΔTgAT1ΔTgENT3. For each strain and time point at least 100

vacuoles were scored per biological replicate (n=3 independent infections). Bars show

mean percentage ± SEM of total vacuoles; categories sum to ~100%. No significant

differences in overall replication distribution between strains were detected by two-way

ANOVA (factors: strain, vacuole size category) strain comparisons at each time point).

**(B)** Plaque formation after 10 days of continuous lytic growth in HFF monolayers

(crystal violet staining). Bars indicate mean plaque number ± SEM (n=3 independent

plates per strain; each point = one plate). One-way ANOVA revealed no significant

difference among strains (p>0.05). Data indicate that deletion (or complementation) of

individual ENTs or combined loss of TgAT1 and TgENT3 does not measurably impair

tachyzoite intracellular replication over 72 h nor cumulative multi-cycle growth under

these assay conditions.

**Supplementary Figure S3. Reduced cyst size in ΔTgAT1 and ΔTgAT1ΔTgENT3**

**relative to parental and single ENT mutants. (A)** Quantification of cyst diameters

(median ± Standard Error of the Mean) for in vitro differentiated bradyzoites (alkaline

pH, Day 3) of the indicated strains: parental ME49Δku80, ΔTgAT1, complemented

ΔTgAT1::TgAT1-myc, ΔTgENT2, ΔTgENT3, complemented ΔTgENT3::TgENT3-ty, and

the double mutant ΔTgAT1ΔTgENT3. Each point represents the mean of one biological

replicate (n=3 independent differentiations); ≥150 DBA+ cysts were measured per strain

per replicate using an automated ImageJ macro**.** One-way ANOVA multiple

comparisons test detected significantly smaller cysts in ΔTgAT1 and ΔTgAT1ΔTgENT3

compared with parental and respective complemented strains (**** p<0.0001); "ns" =

not significant. **(B)** Representative high-magnification images of DBA-stained cysts

(magenta in merge; direct grayscale shown in second column) from each strain at Day

3. Columns: transmitted light (Phase), DBA (cyst wall), anti-Tg (pan-parasite antibody

marking internal bradyzoites), DAPI (host and parasite nuclei), and merged composite

(DBA green, anti-Tg magenta, DAPI blue). Complemented lines restore cyst sizes and

wall morphology towards parental phenotype. Scale bars: 10 μm. Images acquired

under identical exposure and processing settings across strains. Data support a

delay/reduction in cyst maturation rather than a complete differentiation block in

transporter mutants.

**Supplementary Figure S4. In vivo brain cyst burden and morphology after 30 days**

**of infection with parental and ENT mutant strains. (A)** Total brain cyst counts (cysts

per mouse) at 30 days post intraperitoneal infection (250 tachyzoites) with the indicated

strains: parental ME49Δku80, ΔTgAT1, complemented ΔTgAT1::TgAT1-myc,

ΔTgENT2, ΔTgENT3, complemented ΔTgENT3::TgENT3-ty, and the double mutant

ΔTgAT1ΔTgENT3. Each point = one mouse brain; bars show mean ± SEM. Two

independent experiments were performed (4 male C57BL/6J mice per strain per

experiment; symbols pooled after confirming no experiment–strain interaction by two-

way ANOVA). No significant difference in total cyst burden among strains (one-way

ANOVA, p>0.05). **(B)** Mean cyst diameter (μm) ± Standard Error of the Mean (SEM) for

DBA-positive cysts from the same brains (≥50 cysts measured per strain across mice;

automated ImageJ pipeline with uniform thresholding). ΔTgENT3 and ΔTgAT1ΔTgENT3

display significantly smaller cysts relative to parental and their respective complemented lines (one-way ANOVA with Tukey's multiple comparisons). *p<0.05, **p<0.01, ns = not significant.

**(C)** Representative fluorescence images of individual brain cysts stained with Dolichos biflorus agglutinin (DBA; cyst wall, green in merge), anti-Toxoplasma (pan-parasite antibody, magenta), and DAPI (nuclei, blue). Smaller, less fully expanded cysts are evident in ΔTgENT3 and most pronounced in the double mutant. Complementation ( TgENT3-ty1) restores cyst morphology toward parental appearance. Images acquired under identical acquisition settings; Scale bar: 10 µm.

**Supplementary Figure S5. PCR validation of TgENT knockouts, double knockout, and epitope/complementation lines.**

**(A) Diagnostic PCR across endogenous loci.** Genomic DNA from eight parasite lines (lane order, left to right: 1 ΔTgENT2; 2 ΔTgAT1ΔTgENT3; 3 parental ME49; 4 ΔTgENT3; 5 ΔTgAT1; 6 ΔTgENT3::TgENT3-Ty1; 7 ΔTgAT1::AT1-myc; 8 $H_2O$ control) was amplified with primer pairs specific for GRA17 (loading control) (P7/P8), TgENT1 (P9/P10), TgENT2 (P11/P12), TgENT3 (P13/P14) or TgAT1 (P15/P16), as indicated above each gel strip. Absence of the wild-type amplicon (or size-shift where a repair cassette was inserted) confirms successful disruption, whereas parent and complemented strains retain the expected band.

**(B) Verification of selectable-marker and complementation inserts.** Individual PCRs using primers flanking the repair junctions demonstrate correct integration of the chloramphenicol-acetyl-transferase (CAT) cassette into ΔTgAT1 (top)

(P17/P18) and ΔTgENT3 (fourth panel) (P19/P20), of the AT1-myc (P21/P22) and

ENT3-Ty (P23/P24) complementation cassettes into their respective loci (second and

third panels), and of a DHFR cassette into the ΔTgAT1ΔTgENT3 line (bottom)

(P25/P26). Lane order for each gel: 1 edited line; 2 parental ME49; 3 water control. The

PCR product was Sanger-sequenced as confirmation.

**(C) Integration of the mAID-HA tag at the TgENT1 locus.**

PCR with locus-specific primers yields the diagnostic HA-junction product exclusively in

the TgENT1-mAID-HA strain (lane 1) but not in RH-TIR1 (lane 2) or water control (lane

3), confirming single-locus tagging (P27/P28).

Molecular-weight ladders are shown in the first lane of each gel; expected band sizes

are indicated by yellow arrowheads. Together, these assays verify the correct genetic

architecture of all strains used in this study.

**Supplemental Table 1:** Oligo list detailing ssDNA sequences used for cloning,

genotyping, qPCR, and guide RNAs.

**Supplemental Table 2:** Full DNA sequences of our plasmids used for creation of

mutant strains.

**Supplemental Table 3:** List of antibodies used for immunofluorescence detection.

**Supplemental Table 4:** Differentiation expression analyzes of TgENT1 knockdown 24-

and 48-hours post-infection.

**Methods**

**Parasite Maintenance:** *Toxoplasma gondii* tachyzoites, ME49 and RH strains, were

propagated in human foreskin fibroblasts (HS27; ATCC CRL-1634) at 37 °C in DMEM

supplemented with 10 % FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin. Cells

and parasite lines were tested for mycoplasma monthly using a PCR test (Boca

Scientific).

**Plasmid Construction:** Final sequences are listed in **Supplementary Table 2**. PCRs

employed Q5 High-Fidelity (long fragments) or Taq DNA polymerase (routine

amplifications). Amplicons and vector backbones were assembled with Gibson

Assembly (NEB) unless noted.

- **ΔTgENT3 donor** – synthetic plasmid (GenScript).

- **ΔTgENT3::TgENT3-Ty1 donor** – 23-DD plasmid digested with BglI. Synthetic
insert (B1) ordered and inserted (Twist Bioscience).

- **ΔTgAT1 donor** – ΔTgENT3 donor digested with AvrII/NotI (insert) and SpeI/AflII
(backbone); 5′ and 3′ TgAT1 homology arms amplified with P1/P2 and P3/P4.

- **TgENT3-DHFR-KO donor** – DHFR cassette (Addgene #80329) amplified with
P5/P6 and inserted into ΔTgENT3 backbone (NotI/EcoRV).

- **mAID-HA-TgENT1 donor** – synthetic plasmid (GenScript).

- **ΔTgAT1::TgAT1-myc donor** – 24-DD plasmid digested with BssHI/ScaI.
Synthetic inserts (B2 and B3) ordered and inserted (Twist Bioscience).

- **ΔTgENT2** – ΔTgENT3 donor digested with AflII/SnaBI. Synthetic insert (B4)
ordered and inserted (Twist Bioscience).

• **Guide RNA plasmids** – pSS013 vector (Addgene) was BsaI-linearised;
annealed IDT oligos were inserted by Gibson Assembly, transformed into *E. coli*
DH5α, sequence-verified, and prepared with GeneJET Maxi kits.

**Genomic Editing and Transfection:** Knock-outs, knock-ins, and mAID tagging were
generated with dual-guide CRISPR–Cas9 as described (voltage 1500 V, 0.1 ms, two
pulses; BTX ECM™ 830, 4-mm cuvette). Parasites harvested from confluent T25 flasks
were washed once in cytomix (120 mM KCl, 0.15 mM CaCl$_2$, 10 mM K$_2$HPO$_4$/KH$_2$PO$_4$
pH 7.6, 25 mM HEPES, 2 mM EGTA, 5 mM MgCl$_2$). Electroporation mixtures contained
50 µg linearized donor DNA and 25 µg of each sgRNA plasmid in 400 µl cytomix.
Parasites were immediately added to fresh HS27 monolayers. Drug selection began 24
h post-transfection (40 µM chloramphenicol or 3 µM pyrimethamine). Drug-resistant
populations were single-cell-sorted (or cloned by limiting dilution), expanded, and
validated by diagnostic PCR, Western blot, and immunofluorescence (primer sets P7–
P28; see Supplementary Table 2).

**AlphaFold Modeling:** Predicted protein structures were retrieved from the AlphaFold
Protein Structure Database (AlphaFold DB) using their UniProt accession code. These
predicted structures were aligned with human ENT1 (PDBID: 6OB7) using PyMOL
version 3.1.4.1 (Schrodinger). Root Mean Square Deviation (RMSD) of each transporter
relative to human ENT1 was calculated using PyMOL. Atomic measurements of pore
width were also performed in PyMOL. As shown using HsENT1 in **Fig. 1C**, pore width
was measured across the widest and narrowest points (yellow and red dashed lines).
TgENT1, TgENT2, TgENT3, and TgAT1 pores were measured in the same manner.

**Immunofluorescence Assay:** Cells were fixed in 3.7% formaldehyde for 20 minutes, washed twice with 1x PBS, blocked and permeabilized with 3% BSA and 0.2% Triton X-100 for 2 hours at room temperature (or, alternatively, overnight at 4°C). Primary antibodies were diluted 1:500 (unless otherwise stated in Supplemental Table 3) in blocking/permeabilization solution and incubated for 1 hour at room temperature (or overnight at 4°C). Cells were washed twice with 1x PBS, secondary antibodies were applied at 1:500 dilution (DAPI at 1:200) in blocking/permeabilization solution for one hour at room temperature, washed three times for 5 minutes, and glass coverslips were mounted and sealed.

**Pearson and Manders' colocalization coefficients:** Pearson's *r* and Manders' coefficients assess spatial overlap between two fluorescence signals after background removal. Pearson's correlation coefficient (*r*) measures the linear proportionality of intensities on a pixel-by-pixel basis (−1 to 1), whereas Manders' M1 and M2 report the fraction of fluorophore A (or B) signal that coincides with the other channel independent of intensity proportionality. Colocalization was quantified in raw 16-bit TIFF images using Fiji (ImageJ) Coloc2. After identical acquisition (no saturation; sequential capture to avoid bleed-through), images were opened without gamma adjustment, a uniform rolling-ball background subtraction was applied to both channels, and no nonlinear filtering or manual thresholding was performed. Reported values in the manuscript are the averages of 50 vacuoles per condition pooled from three independent biological experiments.

**Bradyzoite *in vitro* differentiation:** The *in vitro* differentiation was performed as previously [34]. Confluent HS27 fibroblast monolayers on glass coverslips were infected with freshly egressed tachyzoites (MOI ≈1; 2 h invasion + gentle wash; total pre-induction incubation 4 h). Cultures were then shifted to pre-warmed alkaline induction medium (RPMI 1640 supplemented with 1% heat-inactivated FBS, 50 mM HEPES, pH 8.20 ± 0.05 at 37 °C) and incubated in ambient air (≈0.04% $CO_2$) in a humidified chamber (no external $CO_2$) to maintain pH ≥8.0. Induction medium was replaced every 24 h with freshly adjusted medium; pH was spot-checked at each change. After 2-, 3-, or 7-days post-infection, coverslips were fixed (4% paraformaldehyde/PBS, 20 min), permeabilized (0.2% Triton X-100), and dual-stained with rabbit anti-*Toxoplasma* antibody (Thermo Fisher; 1:500) to delineate vacuoles and Dolichos biflorus agglutinin (DBA; 2–5 µg/ml) to label the cyst wall. A vacuole was scored "differentiated" if ≥75% of its perimeter showed a continuous DBA rim. For each condition ≥100 vacuoles per coverslip were scored blindly (≥3 biological replicates).

**Tachyzoite Intracellular Growth and Plaque Assays:** Confluent HS27 (human foreskin fibroblast) monolayers on glass coverslips were infected with each *T. gondii* strain at an MOI yielding mostly isolated vacuoles (typically MOI 0.5–1.0) in standard pH 7.4 medium (DMEM + 10% FBS, 5% $CO_2$). After allowing invasion for 2 h at 37 °C, extracellular parasites were removed by two PBS washes and fresh medium was added. At the indicated post-infection time point in each experiment, cells were fixed in 4% paraformaldehyde (15 min), permeabilized (0.1% Triton X-100, 10 min), and

blocked (3% BSA/PBS, 30 min). Parasites were labeled with anti-*Toxoplasma* (followed

by Alexa Fluor–conjugated secondary antibody). For each biological replicate, 100

intact, non-overlapping parasitophorous vacuoles were randomly selected using a

Cytation 7 imager (Agilent) under consistent exposure settings, and the number of

tachyzoites per vacuole was recorded. Three biological replicates were performed per

experiment. Vacuoles at the edge of fields, obviously disrupted, or with uncertain

boundaries were excluded during counting.

For **plaque assays**, confluent HS27 monolayers in 6-well plates were infected with

1,000 of freshly egressed tachyzoites in 2 mL of complete medium. At endpoint,

monolayers were fixed in 100% ethanol (10 min) and stained with 0.1% crystal violet (in

20% methanol) for 15–20 min, rinsed, and air-dried. Plaques (zones of host cell

clearance) were imaged, counted manually and plaque areas were measured after

background subtraction and uniform threshold application. At least three biological

replicates (independent parasite preparations on different days) per strain were

analyzed. Statistical analysis used one-way ANOVA with appropriate multiple

comparison correction. **Quality controls.** Parasite viability and inoculum accuracy were

confirmed by counting extracellular tachyzoites with a hemocytometer.

**RNA extraction, mRNA sequencing and transcriptomics analyses:** Total RNA was

extracted from *T. gondii* infected cells using the GeneJET RNA Purification Kit (Thermo

Fisher Scientific) according to the manufacturer's protocol. RNA quality and

concentration were confirmed, and poly(A)-selected mRNA libraries were subsequently

prepared by Novogene. Libraries were sequenced on an Illumina NovaSeq X Plus

platform, generating 150-bp paired-end reads. All experiments were performed in

biological triplicate, and the resulting data are publicly available in the NCBI Sequence

Read Archive under BioProject accession PRJNA1247533. Reads were adapter- and

quality-trimmed with Cutadapt, mapped to the reference genome using STAR in splice-

aware mode, and uniquely aligned reads were summarized to gene-level counts with

featureCounts. Differential expression was then evaluated in R with DESeq2, which

applied median-of-ratios normalization, negative-binomial GLM fitting, and Benjamini–

Hochberg correction to identify significantly regulated genes.

**Immunofluorescence Assay for PLVAC Morphology:** PLVAC morphology was

assessed 36 h post-infection in TgENT1 AID-HA tachyzoites ± IAA. Parasites were

mechanically released (27-g needle) from host cells and immediately inoculated onto

fresh confluent HS27 monolayers for a 30-min invasion pulse (37 °C). Cultures were

fixed in 4% paraformaldehyde (20 min, RT) and stained with primary anti-TgCPL

antibody (PLVAC marker) followed by appropriate fluorophore-conjugated secondary

antibodies and DAPI. Imaging was performed under identical acquisition settings across

conditions. For each condition, the cross-sectional area of the first 50 TgCPL-positive

PLVAC puncta associated with intracellular parasites was quantified (Fiji) in each of

three biological replicates (n = 50 vacuoles/condition). Areas were compared by t-test.

**Animal experiments**

All animal work was approved by the University of Vermont IACUC (protocol

PROTO202100038) and performed in accordance with institutional guidelines. Mice

were housed under a 12 h light/12 h dark cycle with daily health monitoring that began

one week before infection and continued until study end; animals were euthanized if

predefined humane endpoints were met.

Two independent experiments were conducted. In each, four adult male CBA/J mice

(The Jackson Laboratory) per parasite strain received an intraperitoneal inoculation of

250 freshly-egressed tachyzoites suspended in 200 µl sterile PBS. At 30 days post-

infection (dpi), mice were euthanized following IACUC protocol, and whole cortices were

promptly harvested for downstream analyses.

Brain immunofluorescence was conducted as previously reported [37]. Briefly, the tissue

was homogenized using serial needles passing through 18-, 20-, and 22-gauge

needles, pelleted at 1000xg for 5 minutes, and fixed with 3.7% formaldehyde for 20

minutes. Pellets were washed twice with 1x PBS, blocked overnight in 1% BSA and

0.2% Triton X-100 in 1x PBS, primary antibodies diluted 1:500 were incubated

overnight, washed twice with 1x PBS, secondary antibodies (1:500 dilution) and DAPI

were added for 1 hour, washed twice with 1x PBS, and imaged using a microscope.

Cyst stained with DBA were wet mounted onto coverslips and scanned for DBA using a

Cytation 7 imager (Agilent). 50 randomly found cysts per sample were measured and

recorded using Agilent BioTek Gen5 software.

**qPCR**

RNA was extracted from freshly lysed tachyzoites using the GeneJet RNA Purification

Kit. Complementary DNA (cDNA) was subsequently synthesized employing the Maxima

H Minus First Strand cDNA Synthesis Kit. Quantitative PCR (qPCR) reactions were then

set up with primers Q1 – Q8 using the PowerSYBR Green PCR Master Mix.

**Statistical Analysis**

GraphPad Prism 9 was used for all statistics. Data are mean ± SEM of ≥3 independent

experiments. Group differences were analyzed by one-way ANOVA or unpaired two-

tailed Student's *t*-test (as indicated). *P* < 0.05 (*), 0.01 (**), and 0.0001 (****) were

considered significant.

**Acknowledgments**

This research was supported by a research grant from National Institutes of Health (P20

GM125498). We thank Dr. Harry de Koning for helpful discussions.

**A.**

**B.**

TgAT1    TgENT1    TgENT2    TgENT3

Top View

Side view

**C.**

Structural Comparison Across ENT Family Members

|        | Pore width (long)(Å) | Pore width (short)(Å) | RMSD (relative to 6OB7) (Å) |
|--------|----------------------|------------------------|------------------------------|
| hENT1  | 30.00                | 18.30                  | 0.00                         |
| TgENT1 | 30.30                | 18.00                  | 3.32                         |
| TgENT2 | 28.80                | 18.10                  | 4.31                         |
| TgENT3 | 30.10                | 18.50                  | 6.54                         |
| TgAT1  | 28.50                | 18.80                  | 4.41                         |

6OB7 (HsENT1)

Width - long

Width - short

**D.**

Expression of TgENT Family Members Across Developmental Stages

| Gene   | EES1  | EES2 | EES3 | EES4 | EES5 | Sporulating | Tachyzoites | Tissue Cysts | Unsporulated |
|--------|-------|------|------|------|------|-------------|-------------|--------------|--------------|
| TgAT1  | 10.15 | 8.45 | 8.92 | 8.64 | 7.51 | 3.86        | 5.93        | 3.58         | 0.00         |
| TgENT1 | 7.04  | 6.94 | 6.95 | 7.27 | 7.62 | 4.38        | 6.69        | 5.67         | 4.07         |
| TgENT2 | 1.53  | 0.00 | 0.00 | 0.00 | 0.00 | 7.34        | 0.00        | 2.44         | 0.04         |
| TgENT3 | 3.53  | 3.76 | 3.71 | 3.97 | 3.58 | 4.54        | 6.56        | 6.19         | 0.15         |

Developmental Stage

**A.**

| | Phase | anti-HA | anti-TgCPL | Merge |
|---|---|---|---|---|
| Vehicle | | | | |
| IAA | | | | |

**B.**