A reaction norm for flowering time plasticity reveals physiological footprints of maize adaptation

Justine Drouault[1], Carine Palaffre[2], Emilie J. Millet[1], Jonas Rodriguez[1], Pierre Martre[1], Kristian
Johnson[1], Boris Parent[1], Claude Welcker[1], Randall J. Wisser[1,3*]

[1]LEPSE, University of Montpellier, INRAE, Institut Agro Montpellier, Montpellier, FR

[2]Experimental Unit of Maize, INRAE, 40390 Saint Martin De Hinx, FR

[3]Department of Plant and Soil Sciences, University of Delaware, Newark, Delaware, USA

*Correspondence: Randall J. Wisser, randall.wisser@inrae.fr

ORCIDs: EJM: 0000-0002-2913-4892; JR: 0000-0001-8693-1022; PM: 0000-0002-7419-6558;
CW: 0000-0002-8275-1259; RJW: 0000-0003-1075-0115

Current address:

Emilie J. Millet: INRAE, UR1052 GAFL, 84143 Montfavet, France

Jonas Rodriguez: Syngenta Seeds, P.O Box 18300, Greensboro, North Carolina, USASyngenta Seeds,
P.O Box 18300, Greensboro, North Carolina, USA

keywords: adaptation, common garden trial, envirotyping, flowering time, maize, photoperiod,
reaction norm

Running title: Physiological footprints of flowering time adaptation in maize

Date of submission: July 9, 2024

Word count: 4,418

Number of tables and figures
Tables: 2
Figures: 4

Supplementary data
Figures: 2
Files: 6

Abstract

Understanding how plant phenotypes are shaped by their environments is crucial for addressing
questions about crop adaptation to new environments. This study investigated the interplay between
developmental responses to temperature fluctuations and photoperiod perception in maize that
contribute to genotype-by-environment variation in flowering time. We present a physiological
reaction norm for flowering time plasticity (PRN-FTP) for studying large collections of genotypes
tested in multi-environment trial (MET) networks. Using a new variable for computational
envirotyping of sensed photoperiod, it was found that, at high latitudes, different genotypes in the same
environment can experience hours-long differences in photoperiod. This emphasizes the importance of
considering genotype-specific differences in the experienced environment when investigating
plasticity. A statistical framework is introduced for modeling the PRN-FTP as a non-linear response
function, with parameters putatively linked to different regulatory modules for flowering time.
Applying the PRN-FTP to a sample of global breeding material for maize showed that tropical and
temperate maize occupy distinct territories of the trait space for PRN-FTP parameters, supporting that
the geographical spread and adaptation of maize was differentially mediated by exogenous and
endogenous pathways for flowering time regulation. Our results have implications for understanding
crop adaptation and for future crop improvement efforts.

Introduction

Plants exhibit vast phenotypic diversity, the principal axis for adaptation and evolution. The classic formalism $P = G + E + G \times E$ describes how phenotypes are determined by combinations of genetic (G) and environmental (E) effects. Different environmental conditions can produce distinct phenotypic outcomes for the same genotype, a phenomenon known as plasticity (Bradshaw, 1965). As 21st-century climate markedly shifts the environmental dimension to new frontiers, it is an important and opportune time to investigate mechanisms of environmental plasticity (E) and genetic variation in plastic responses ($G \times E$). Harnessing plastic responses to dominant shifts in climate conditions can facilitate the adaptation of crops into new territories or to alternative cropping systems (Rising and Devineni, 2020; Sloat *et al.*, 2020; Zabel *et al.*, 2021). New approaches for understanding and predicting the responses of traits driven by combinations of climate and weather conditions are also encouraging (Millet *et al.*, 2019; Cooper *et al.*, 2020; van Voorn *et al.*, 2023).

The functional relationship that defines a pattern of phenotypic change across an environmental gradient is known as a reaction norm, which provides information about the direction and magnitude of plasticity (Roof, 1997). To construct norms of reactions, phenotypic values must be recorded in a large number of settings presenting environmental variation, and a relevant variable or index is used to define the gradient of environmental conditions. There are different ways of formulating the response (trait variation) and predictor (environment variation) variables for modeling a reaction norm. Strictly speaking, reaction norms are defined as response functions to single environmental factors, but the term has also been used to describe responses to environments *per se*. A common approach in quantitative genetics uses the *Finlay–Wilkinson* model (Finlay and Wilkinson, 1963), in which the phenotypic values of individual genotypes are regressed on an index defined by the mean values of all genotypes per environment. This method provides a convenient generalization for using multi-environment trials (METs) with large numbers of genotypes to dissect and predict variation in plasticity (Jarquín *et al.*, 2014; Kusmec *et al.*, 2017; Guo *et al.*, 2023). Clearly, however, the environment is abstractly defined, leading to a lack of biological interpretability. In contrast, under controlled settings with fine-scale phenotyping, ecophysiologists have determined response functions of traits to environmental variables that allows for the definition of environmental sensitivities of different plant processes, such as the non-linear response of growth and development to temperature (Parent *et al.*, 2010; Parent and Tardieu, 2012). This provides biological perspective, but upscaling to study broad diversity in METs remains challenging. By extension, crop modelers have tied together these governing principles as mathematical models of ecophysiological processes, leading to systems for predicting the phenotypes of emergent characteristics as a function of genetics and environments (Muller and Martre, 2019). Concepts from these domains can be combined to define and model biologically-informed reaction norms in the design and analysis of MET networks.

In plants, the control of flowering time is regulated by a modular molecular network controlling physiological responses to multiple external signals, particularly temperature and photoperiod (Simpson and Dean, 2002; Colasanti and Coneva, 2009; Blümel *et al.*, 2015; Lee *et al.*, 2021). Responses to temperature occur throughout a plant's life cycle, while the response to photoperiod is growth-stage specific. In maize (*Zea mays* subsp. *mays*), a facultative short-day (SD) species, the stage at which plants become sensitized to photoperiod is not directly observable, but has been found to occur just prior to the timing of tassel initiation for plants grown in SD conditions (Kiniry *et al.*, 1983*a,b*). Above a critical photoperiod, tassel initiation is delayed beyond the sensitized stage for photoperiod sensitive genotypes. As physiological responses to temperature fluctuations interact with genotypic variations in the rate and duration and of plant development, different genotypes can reach the sensitized stage on different days, and therefore can perceive different photoperiods, even within

the same environment. For photoperiod sensitive genotypes, genotypic variation in critical thresholds
and degrees of photoperiod sensitivity create additional response variation, culminating in the
formation of $G \times E$ interactions in flowering time across environments. Extending previous work
(Kiniry *et al.*, 1983*a,b*; Choquette *et al.*, 2023), the current study is centered on the modeling and
analysis of a physiological reaction norm for flowering time plasticity (PRN-FTP) that captures the
combined effects of physiological sensing and responses to temperature and photoperiod underlying
variation in flowering time across environments. We formulated this as a bilinear, threshold-dependent
function for the relationship between thermal time from crop emergence to male flowering (anthesis;
T_t^{em-ant}) and a new environmental index for sensed photoperiod (DL^s). We also demonstrate how
statistical modeling of the PRN-FTP allows the decoupling of different components of flowering time
regulation: (i) flowering time *per se*; (ii) critical photoperiod; and (iii) photoperiod sensitivity.

Constructing PRN-FTPs from METs is the most practical approach for studying flowering time
plasticity in large collections of genotypes, but this is difficult due to the geographical structure of
photoperiod and the non-linear reaction norm. This requires testing across numerous field sites
spanning nearly an entire hemisphere to capture a sufficient range and density of DL^s needed to
estimate the non-linear function. A better understanding of the variation in DL^s within and across
environments could aid in the efficient design of METs for studying the PRN-FTP. A particular
difficulty is determining critical photoperiods, which in turn affect estimates of flowering time *per se*
and photoperiod sensitivity. Also, detecting the true PRN-FTP based on thermal time can be obscured
by the influence of other factors besides temperature on growth and development, such as drought (e.g.,
Harrison *et al.*, 2014; Herrero and Johnson 1981). Additionally, since genotypes can become sensitized
to photoperiod at different points in time within each environment, it must be considered whether a
common reference value or genotype-specific values for DL^s should be used for the PRN-FTP. As a
result, a full understanding about the diversity of flowering time plasticity in maize and other species
has yet to be realized. As another aspect of the current study, new methods are introduced for
envirotyping DL^s at a global scale, in addition to techniques for statistically analyzing the PRN-FTP.

Even if METs span photoperiod thresholds for a species, sparse coverage of DL^s can hinder the ability
to fit accurate functions for the PRN-FTP. For instance, an MET network may consist of environments
that can only be split into sets of long-day (LD) and SD scenarios by assuming a common threshold
for photoperiod sensitivity across all genotypes. In such cases, a simplified LD-SD method may be
required, where photoperiod sensitivity is estimated as the difference between mean T_t^{em-ant} in sets of
LD and SD environments (Coles *et al.*, 2010; Hung *et al.*, 2012). In this case, flowering time *per se*
can also be estimated as the mean T_t^{em-ant} among SD environments. Hereafter, this approach is called
the LD-SD model.

Finally, parameters of the PRN-FTP may be used to understand how components of flowering time
plasticity mediate adaptation to novel environments, and how this in turn changes genotype plasticity
for the future. Historical varieties of maize that spread 2000-5000 years ago from their tropical origins
into temperate zones required flowering time adaptations to cooler climates with longer photoperiods
(Tenaillon and Charcosset, 2011). Using the LD-SD method for genetic dissection, selection on genes
controlling photoperiod sensitivity have been identified as key factors for the tropical-to-temperate
transition of maize (Hung *et al.*, 2012; Yang *et al.*, 2013). Varieties adapted to temperate zones have
effectively lost their sense of photoperiod, such that flowering times are largely determined by
temperature interactions with variation in flowering time *per se*. Similar results from the adaptation of
tropical populations to temperate environments have been observed over short timescales (within 10
generations). When tropical populations have been selected for early flowering time in LD temperate
environments, photoperiod sensitivity was found to rapidly diminish while flowering time *per se* was

gradually reduced (Teixeira *et al.*, 2015; Wisser *et al.*, 2019; Choquette *et al.*, 2023). In contrast,
selection in SD environments reduced only flowering time *per se*, since photoperiod sensitivity was
not induced (Choquette *et al.*, 2023). Here, we show that relating the variation in flowering time *per*
se and photoperiod sensitivity for separate breeding pools can be used to understand how selection
history has shaped the plasticity of modern maize.

Materials and Methods

Experimental datasets

Multiple experiments and datasets from different projects were assembled to: (D1) select a temperature
response function for thermal time estimation; (D2) estimate the thermal time duration between crop
emergence and tassel initiation (T_t^{em-TI}) for diverse maize lines; (D3) envirotype sensed photoperiod
(DL^s) in maize fields across the Northern Hemisphere; (D4) demonstrate methods for modeling a
physiological reaction norm for flowering time plasticity (PRN-FTP); and (D5) compare the trait space
of parameters estimated from the PRN-FTP in a diversity panel constituting separate breeding pools of
maize. The current study focuses on days to anthesis (male flowering) as the final characteristic.

(D1) Three temperature response functions (Methods S1, Eq. S1-S3) for measuring thermal time were
compared using a dataset with 19 hybrids tested in at least 34 field environments, spanning latitudes
from 30.5371°N to 51.5016°N (File S1; data sourced from the Genomes-To-Fields Initiative). For each
environment, the sowing date, field coordinates, and best linear unbiased estimates (BLUEs) for
calendar days from planting to anthesis were obtained from McFarland *et al.*, 2020 and Rogers *et al.*,
2021. Using temperature data, the genotype-specific BLUEs for calendar days to anthesis were
converted to thermal time values.

(D2) The T_t^{em-TI} was estimated for a collection of 239 maize inbred lines (panel 1) using a
physiological formalism (Eq. 1), based on data for crop emergence, phyllochron, and final leaf number
recorded in two experiments under short-day (SD) conditions (File S2). The T_t^{em-TI} measured in SD
conditions is referred to as the basic vegetative phase to tassel initiation (BVP-ti; Kiniry *et al.*, 1983*a*).
Panel 1 primarily consisted of semi-tropical or tropical lines but included some temperate lines. Crop
emergence and phyllochron were measured in a greenhouse phenotyping platform at the LEPSE of
INRAE (PhenoArch, Montpellier, FR; Cabrera-Bosquet *et al.*, 2016) under well-watered conditions,
while final leaf number was recorded in a field experiment in Puerto Vallarta, MX, also in well-watered
conditions. For both experiments, daylengths throughout the growth cycle never exceeded 12.5 h,
providing SD conditions in which photoperiod was not expected to suppress tassel initiation.
Additional details for these experiments can be found in Supplemental Information (Methods S2).

(D3) Using the BVP-ti as a proxy for DL^s, computational envirotyping of DL^s was performed for fields
across the Northern Hemisphere (latitudinal range of 13.7578 °N to 54.2900 °N) where maize was
previously grown. For this, the geographical coordinates and planting dates of 680 field environments
were assembled from past publications and ongoing projects (File S3). Crop emergence dates available
from 15 field environments sourced from the Maize ATLAS project (Choquette *et al.*, 2023) were used
to compute a mean thermal time from sowing to emergence of 76 °Cd (estimated using Eq. S2), which
was used for predicting times to emergence in the remaining environments where only the planting
date was available.

(D4) To test modeling procedures for the PRN-FTP, flowering time data from 37 field trials were
combined into a multi-environment trial (MET) dataset (File S4). The dataset included observations

that maximized observations for a common set of genotypes also present in dataset D2, including seven
temperate inbred lines (B37, B73, M37W, Mo17, Oh43, LH123Ht, and 2369) and seven tropical inbred
lines (CML10, CML258, CML277, CML341, CML373, Tzi8, and Tzi9). Still, representation of these
lines across the 37 environments was not fully balanced. The temperate lines were present in a
minimum of 13 and a maximum of 22 environments. The tropical lines were present in a minimum of
18 and a maximum of 37 environments.

(D5) To examine the relationship between PRN-FTP parameters associated with flowering time *per se*
and photoperiod sensitivity, MET data for a collection of 111 temperate, 52 tropical, and 51 admixed
inbred lines constituting separate breeding pools (panel 2; Flint-Garcia *et al*., 2005) was assembled
from previous experiments (File S5). These lines were observed across a maximum of three SD
environments (maximum photoperiod <12.3 h) and 16 long-day (LD) environments (minimum
photoperiod of >14.5 h) spanning 18.00 °N to 42.76 °N latitude.

Environmental data

For field trials considered in this study, temperature data were acquired from ground and satellite data
sources for a common reference point of 2 m height. Ground data were obtained mostly from weather
stations at a given field location. In two cases, ground data from a nearby airport within 5 km of the
experimental field site was obtained from Weather Underground (https://www.wunderground.com/).
Satellite data (weather variable 'T2M'; spatial resolution of ½° latitude x ⅝° longitude) was acquired
from the R package *nasapower* (Sparks, 2018) using the latitude and longitude coordinates for each
field location.

The photoperiod based on civil twilight at a given location for a given day of the year was acquired
from the R package *TrenchR* (Buckley *et al*., 2023). This uses the CBM daylength model described by
Forsythe *et al.* (1995).

Comparison of temperature response functions

Three temperature response functions were compared for calculating thermal time (Eq. S1-S3).
Comparisons were made based on the normalized root mean square error (NRMSE, normalized by the
difference between maximum and minimum values) for the overall mean thermal time from sowing to
anthesis. The NRMSE was estimated for each of the 19 hybrids in dataset D1 in order to identify the
response function that minimized the mean NRMSE among all hybrids. The analyzed set of hybrids
were from crosses between temperate-adapted lines classified as photoperiod insensitive, such that
variation in anthesis dates (calendar days) across environments were expected to be primarily due to
differences in temperature. Therefore, the magnitude of residual variation per hybrid was expected to
reflect deviations from the model for temperature response. For the purpose of comparing different
models for thermal time estimation, we assumed the influence of any latent variables on growth and
development could be ignored.

Physiological reaction norm for flowering time

The PRN-FTP defined here was reformulated from a previous model by Kiniry *et al.* (1983*a*) that
described the relationship between photoperiod conditions (x-axis) and thermal time from crop
emergence to tassel initiation (T_t^{em-TI}; y-axis). Kiniry *et al.* (1983*a*) used growth chambers in which
constant levels of photoperiod were applied throughout the plant's growth cycle. However, in the
setting of MET networks considered here, photoperiods are not constant (DL^s depends on the timing
at which a plant become sensitized to photoperiod in a given environment) and estimating tassel

initiation from dissections of the shoot apical meristem is not practical at large scale (rather, in METs,
days to flowering are routinely recorded). Since there is no visible marker for when DL^s occurs, we
devised a proxy measure based on previous results showing that the initial stage of sensitization to
photoperiod closely corresponds to the BVP-ti (i.e., the T_t^{em-TI} measured under SD conditions; Kiniry
et al., 1983*b*). Thus, here we define the PRN-FTP as the relationship between the proxy measure for
DL^s (x-axis) and thermal time from crop emergence to anthesis (T_t^{em-ant}; y-axis)

DL^s was obtained by first calculating T_t^{em-TI} (Eq. 1) for genotypes grown in SD conditions (Methods
S2). Considering that tassel initiation occurs on the day when the last leaf primordium is formed
(Lejeune and Bernier, 1996) and that rates of leaf primordia formation and leaf tip appearance are
proportional, T_t^{em-TI} can be calculated as:

$$T_{t_i}^{em-TI} = \frac{P_{tip_i}}{\alpha_{phyll/plast}}\left(L_{F_i} - L_{pri}^{em}\right) \qquad \text{(Eq. 1)}$$

where, $T_{t_i}^{em-TI}$ (°Cd) is the cumulative thermal time from crop emergence to tassel initiation of the ith
genotype, P_{tip_i} (leaf °Cd^{-1}) is the phyllochron of the ith genotype (i.e., the thermal time between the
appearance of two successive leaf tips), L_{F_i} (leaf) is the final leaf number of the ith genotype,
$\alpha_{plast/phyll}$ (dimensionless) is a constant ratio of phyllochron to plastochron (i.e., the thermal time
between the formation of two successive leaf primordia), and L_{pri}^{em} (primordium) is a constant number
of primordia formed at crop emergence. Genotype-specific values for P_{tip_i} and L_{F_i} were obtained from
dataset D2 on panel 1 (Methods S2). For maize, $\alpha_{plast/phyll}$ may range from 1.58 to 1.85 (Kiniry, 1991;
Lejeune and Bernier, 1996; Padilla and Otegui, 2005; Andrieu *et al.*, 2006). In this study, we set it at
the average value reported in the literature (1.73). There are four to five leaf primordia in the embryo
of maize seeds (Bonnett, 1954; Padilla and Otegui, 2005) and between germination and crop emergence
about two other leaf primordia are formed (Padilla and Otegui, 2005). Here, L_{pri}^{em} was set equal to 5.5
primordia.

Finally, given a sowing date, the thermal time from sowing to crop emergence, and the T_t^{em-TI} from
Eq. 1, the temperature records for each environment were used to predict the calendar day that a given
genotype becomes sensitized to photoperiod. In turn, the photoperiod of the specific calendar day in
that environment was predicted using the R package *TrenchR* (Buckley *et al.*, 2023) to obtain the DL^s.

Envirotyping of sensed photoperiod

Dataset D3 was used to computationally envirotype DL^s for maize across cultivation areas in the
Northern Hemisphere. In addition, differences in DL^s were quantified for three maturity groups,
defined by the lower (5th), median (50th), and upper (95th) percentiles of the T_t^{em-TI} distribution
estimated for panel 1 (Table 1). The thermal time from sowing to crop emergence (fixed value based
on mean from observations in D3), plus the percentile-maturity group value of T_t^{em-TI} was used to
predict corresponding DL^s values per environment. In each environment, the rate of photoperiodic
change was also computed between single days flanking the day of DL^s for the 50th percentile-maturity
group.

In a typical application, the PRN-FTP would be constructed for each of the genotypes tested in the
MET. While T_t^{em-ant} are specific to the observations on each genotype, the DL^s could be defined for
each specific genotype or a common reference genotype. The choice depends on whether different
genotypes in the same environment experience large or small differences in photoperiod. Therefore, a

sensitivity analysis was performed to assess the impact of using genotype-specific values versus a
common reference value for DL^s.

Statistical analysis of the PRN-FTP

260   With a sufficient coverage of DL^s spanning SD and LD environments, we modeled the PRN-FTP using
261   a continuous two-phase (bilinear) regression model with threshold detection (Fong *et al.*, 2017*a*). Here,
262   using the R package *chngpt* (Fong *et al.*, 2017*b*), the "hinge model" was fit to the PRN-FTP for each
263   genotype in dataset D4.

$$T_t^{\text{em}-\text{ant}} = \alpha + \beta(DL^s - e)_+ \qquad \text{(Eq. 2)}$$

where $T_t^{\text{em}-\text{ant}}$ is the response variable; e is the threshold parameter; DL^s is the sensed daylength with
a threshold effect; $I(DL^s > e) = 1$ when $DL^s > e$ and 0 if $DL^s \leq e$; $(DL^s - e)_+$ denotes the hinge
function, which equals $DL^s - e$ for $DL^s > e$ and 0 otherwise; and α and β are the intercept and slope
parameters, respectively . The model was fitted using default settings, with no constraints set for
estimating the threshold value (breaking point of the bilinear model). Estimated effects include an
intercept, threshold value, and slope, which correspond to physiological components of flowering time:
the flowering time *per se*, the critical threshold for photoperiod sensitivity, and the degree of
photoperiod sensitivity, respectively.

Correction for latent variables affecting thermal times to anthesis. If only temperature influenced the
flowering time of a photoperiod insensitive genotype, their $T_t^{\text{em}-\text{ant}}$ would be nearly the same across
an MET. In our experience, this simplistic assumption often does not hold (as confirmed with dataset
D1), but the cause for deviations can be due to various possibilities, including the influence of other
environmental factors, differences in crop management practices, differences between air temperature
and the temperature sensed by the crop, imprecision and biases in weather data, or other latent
variables. Therefore, with dataset D4, we tested an alternative approach using photoperiod insensitive
control lines to adjust for across environment latent variable effects prior to constructing the PRN-FTP.

Seven photoperiod insensitive temperate genotypes were defined as control lines. The 37 environments
for dataset D4 were assembled from different experiments but unfortunately did not include all seven
control lines in all environments, so two sets of control groups were defined (set 1: B37, B73, M37W,
Mo17, and Oh43; set 2: LH123Ht and 2369). Sets 1 and 2 were associated with 22 and 13 of the
environments, respectively. For each set, the overall mean of $T_t^{\text{em}-\text{ant}}$ was calculated across
environments of the MET in which they were associated. The environment-specific residuals were then
used to adjust the thermal times for the remaining non-control genotypes in each of the corresponding
environments of the MET. We note that using two sets of control lines that were also imbalanced across
environments (see above) could introduce some bias and is not ideal, but our aim was to test this
approach.

Analysis of flowering time plasticity in a diversity panel

We sought to investigate the relationship between flowering time *per se* and photoperiod sensitivity in
panel 2 (dataset D5). The MET lacked sufficient coverage of DL^s to estimate critical threshold values
for the PRN-FTP using the hinge model. Therefore, the parameters were estimated by an LD-SD
method using simple linear regression of the BLUEs for $T_t^{\text{em}-\text{ant}}$ (response variable) on the means of
the DL^s values for the set of SD (range of 11.9 h to 12.3 h) and LD (range of 15.4 to 16.0 h)
environments (predictor variable). Not all genotypes were observed in all environments. Therefore,

DL^s means used for regression analysis varied slightly by genotype according to the sets of
environments for which data were available. When fitting the LD-SD linear regression model, the DL^s
means were offset to zero so that the intercept was estimated at the mean short-day index for the
corresponding genotype. The degree of photoperiod sensitivity was determined as the slope of the
regression.

Results

Choosing a temperature response function to construct the PRN-FTP

Thermal time is used in the estimation of DL^s for envirotyping and formulating the PRN-FTP function,
such that the choice of a temperature response function could result in different outcomes. A
comparison of different temperature response functions showed no difference in the NRMSEs for
thermal times to anthesis for 19 photoperiod-insensitive hybrids tested in more than 34 environments
in North America. Regardless of the temperature response function that was tested, for each hybrid,
the environment-specific thermal time to anthesis deviated, on average, by ~10% (range 7 to 12%)
from the across-environment mean (Fig. S1). Therefore, for the current study, we chose the non-linear
response function by Wang and Engel (1998) adapted for maize (Eq. S2) due to its common
implementation in crop growth models (Wang *et al.*, 2017), its higher physiological realism compared
with the linear response function (Eq. S1), and its greater flexibility compared to the Arhenus model
proposed by (Eq. S3; Parent *et al.*, 2010).

Predicting when maize becomes sensitized to photoperiod

There is no known visible marker to measure the timing when maize plants become sensitized to
photoperiod. Here, we defined this timing using $T_t^{\text{em}-\text{TI}}$ measured under SD conditions since this
closely corresponds to the photoperiod sensitized phase (Kiniry *et al.*, 1983*b*). An issue is that directly
observing tassel initiation requires temporal dissections of shoot apical meristems which is not possible
at a large scale. To circumvent this, we used an ecophysiological formalism to predict genotype-
specific values for $T_t^{\text{em}-\text{TI}}$ (Eq. 1) in panel 1 grown under SD conditions (dataset D2; File S2). A
summary of the variation in $T_t^{\text{em}-\text{TI}}$ is shown in Table 1. In a separate experiment, using a phenotyping
platform with a constant 11 h photoperiod, longitudinal dissections of shoot apical meristems were
performed on eight of the lines from the same panel (Methods S2). The Pearson correlation coefficient
between the shoot apical meristem transition and the predicted BVP-ti was 0.70, indicating the
formalism-based predictions for $T_t^{\text{em}-\text{TI}}$ were reasonably approximated.

Variation in the sensed photoperiod for maize

DL^s integrates information on sowing dates, emergence dates, developmental responses to temperature,
the temperature and photoperiodic conditions of an environment, and the genetic diversity for the
physiological traits underlying flowering time. This can be used to understand how different genotypes
sense photoperiod within and across environments, and how it varies according to spatiotemporal
patterns in climate and weather. To examine how photoperiods are perceived by maize, we
computationally envirotyped DL^s for 680 field environments across the Northern Hemisphere where
maize has been previously grown (Fig. 1).

We first summarized the spatial axis of variation in DL^s using the 50[th] percentile of $T_t^{\text{em}-\text{TI}}$ (319 °Cd)
in panel 1 as a common reference for envirotyping (Table 1; Fig. S2). The maximum difference in DL^s
among all of the environments was 6.4 hours (range: 11.9 to 18.3 h; Fig. 1a). This was primarily driven

by climatic variation in photoperiod across latitude and seasonality for maize cultivation. Temperature
variation and different sowing dates across years at the same planting location or fields at similar
longitudes also led to differences in DL^s, but to a lesser extent. Among 125 field sites where maize was
planted for at least two years, DL^s varied by as much as 1.7 hours, with a mean difference of 0.41 h.
Similarly, considering a 5-degree band of latitude covering -105° to 5° E longitude, traversing maize
fields from the USA and Europe, DL^s ranged from 14.9 to 16.7 h, a difference of 1.8 h.

Next, we questioned the extent to which multiple genotypes grown within the same field environment
perceive a different photoperiod. Beyond the equator, photoperiod is not constant across days of the
year, with increasing rates of daily change in association with latitude. Therefore, genotypes differing
in their T_t^{em-TI} will reach their photoperiod sensitized stage on different days and hence perceive a
different photoperiod (Fig. 1b-c), but to our knowledge the extent to which this varies has not been
investigated. Using the 5th (236 °Cd) and 95th (430 °Cd) percentiles of T_t^{em-TI} in panel 1 to represent
early and late percentile-maturity groups, the DL^s at these limits frequently differed by more than 1 h,
with the largest difference reaching 3.4 h (Fig. 1b). Differences in DL^s above 2 h were rare and occurred
only at high latitudes where late genotypes are not cultivated for maize production. Nevertheless, this
highlights how the genetic diversity of maize can intersect with environmental variation within the
same field, leading to differences in the photoperiod to which genotypes respond.

Considering the use of the 50th percentile as a common reference for all genotypes, the relative
difference in DL^s for early and late percentile-maturity groups varied by as little as no difference to as
much as 1.35 h and 2.04 h, respectively (File S3). The largest differences in DL^s among genotypes
within the same field occurred at high latitudes where rates of photoperiodic change were greatest. This
reduced to zero as a function of the rate of photoperiodic change (Fig. 1d).

Using METs for modeling the PRN-FTP

To demonstrate and test approaches for modeling the PRN-FTP, we combined flowering time data
from multiple experiments into an MET (dataset D4) for a common set of seven genotypes of tropical
origin that were expected to be photoperiod sensitive. This was used to estimate the flowering time *per*
se (intercept), the critical photoperiod (threshold), and the photoperiod sensitivity (slope) using a two-
phase regression model (see methods; Fig. 2).

Since different genotypes can experience different photoperiods across common environments, we
tested the impact of using values of DL^s estimated from a common reference (i.e., the 50th percentile
of T_t^{em-TI} in panel 1) versus separate, genotype-specific values when constructing the PRN-FTP.
Compared to the common reference, the T_t^{em-TI} values used for estimating DL^s ranged from the 37th
to 95th percentiles of the T_t^{em-TI} distribution in panel 1. Comparing DL^s values across the MET of
dataset D4, the photoperiods perceived by specific genotypes relative to the common reference differed
by no more than 0.47 h. In this case, there was no change in the estimated thresholds for sensitivity
and differences in the coefficient estimates were negligible.

Using MET data to construct the PRN-FTP makes the simplistic assumption that physiological
sensitivities to temperature and photoperiod are the only environmental factors influencing flowering
time. Although temperature and photoperiod are dominant factors, other variables can also affect maize
growth and development, but the specific causes are often unknown and likely to vary from one
environment to another. Therefore, we devised an approach to control for this. Using a set of
photoperiod insensitive genotypes present in the same MET, their overall mean T_t^{em-ant} was defined
as an expected value for the common effect of temperature across environments (i.e., for photoperiod

insensitive genotypes, $T_t^{\mathrm{em-ant}}$ is expected to be the same across environments, with deviations attributed to latent variables; see Materials and Methods). Next, environment-specific deviations from the overall mean of the insensitive lines were used to adjust the observed $T_t^{\mathrm{em-ant}}$ for the remaining lines in each of the corresponding environments. Fig. 2 portrays this procedure and how it affects estimates for the PRN-FTP parameters.

When tested on the set of seven tropical genotypes, using adjusted values led to changes in the estimated thresholds of sensitivity, which in turn resulted in differences in the estimated flowering times *per se* and slopes of sensitivity (Table 2). On average, latent variable adjustment enhanced the precision of the overall model fit of the PRN-FTP for each genotype by two-fold (Table 2). Even for this small sample of genotypes, Fig. 3 highlights the diversity of flowering time plasticity in maize consisting of unique parameterizations of the PRN-FTP.

Relating components of the PRN-FTP to understand maize adaptation

Since flowering time *per se* and photoperiod sensitivity underlie flowering time adaptation across environments (Choquette *et al.*, 2023), we questioned how these PRN-FTP components differ between primary breeding pools of maize. For this, we assembled an MET dataset (D5) from 19 field trials with a common set of inbred lines consisting of tropical, temperate, and admixed breeding material. Computational envirotyping showed that DL^s for this MET was restricted to two small intervals of photoperiod conditions, with three environments of 11.9 to 12.3 h photoperiod and 16 environments of 14.5 to16.5 h photoperiod. With a large gap in coverage of DL^s, the two-phase regression function for the PRN-FTP could not be reliably fit. In this case, the PRN-FTP was modeled using an LD-SD method based on simple linear regression after adjusting for latent variables. Examining the relationship between flowering time *per se* and photoperiod sensitivity parameters of the PRN-FTP showed that temperate and tropical material occupy distinct territories of the trait space (Fig. 4; File S6). Temperate lines tend to have a lower flowering time *per se* and nearly no photoperiod sensitivity, while tropical lines tend to have a higher flowering time *per se* coupled with moderate to high degrees of photoperiod sensitivity. Fitting expectations, genotypes with admixed ancestry are positioned between the two.

Discussion

Understanding trait plasticity is important for adapting crops to new environments. To study the variation in plastic responses, METs allow large collections of genotypes to be characterized across environmental gradients. However, due to phenological variation and stage-specific sensitivities, different genotypes within each environment do not experience identical environmental conditions (e.g., Millet *et al.*, 2019). This complicates the definition of environmental indices for modeling reaction norms from METs, challenging standard approaches that assume a common environment is sensed by all genotypes. Based on the ecophysiology of maize flowering time, the current study defines a new, genotype-specific metric for quantifying the photoperiod sensed by diverse genotypes within and across environments. Using DL^s to envirotype maize fields in the Northern Hemisphere, we showed that, at high latitudes, where cooler temperatures that slow development combine with relatively large daily rates of photoperiodic change, genotypes in the same environment can experience more than a 1 h difference in DL^s (Fig. 1). Thus, to provide more accurate information about the variation in trait plasticity, the construction of genotype-specific PRNs for flowering time and other traits should be considered. Assembling several experiments, we also found that MET datasets for maize have limited to no coverage of DL^s from 12.5 to 14.0 h for characterizing variation in critical thresholds for the photoperiod sensitivity of maize. With scant information on variation in critical

photoperiods, and given the dependence of accurate threshold estimation for regression analysis of the
PRN-FTP (Table 2), envirotyping with DL^s can be used to strategically sample environments for more
efficient designs of new METs.

The PRN-FTP provides a parsimonious model for flowering time plasticity, taking into account the
dominant effects of temperature and photoperiod. However, in METs, additional latent variables can
affect flowering time, requiring appropriate adjustments. To address this issue, we applied a between-
environment covariate adjustment using photoperiod-insensitive control lines that were expected to
have the same thermal time to anthesis across the full range of DL^s. This resulted in a more precise fit
of the bilinear response function for the PRN-FTP (Table 2; Fig. 2). Nevertheless, it is important to
acknowledge that the choice of a control group could introduce some bias, as it may uniquely vary in
other ecophysiological responses. Another challenge in modeling the PRN-FTP is that not all
genotypes exhibit a clear bilinear response. For example, photoperiod-insensitive genotypes only
display differences in flowering time *per se*, rendering the two-phase regression function invalid. In
such cases, linear regression using the LD-SD method is the most appropriate way of capturing the
relationship between T_t^{em-ant} and DL^s. Thus, while the PRN-FTP model offers a valuable framework
for understanding flowering time plasticity, careful consideration of latent variables, choice of control
group, and genotype-specific responses is necessary for accurate modeling. Incorporating additional
factors and exploring alternative models may further improve the utility of the PRN-FT model for
applications in plant science.

A large body of work has elucidated a generally conserved molecular network for flowering time
regulation in plants, involving a set of semi-independent pathways that converge on the genetic control
of shoot apical meristem identity and anthesis date (Srikanth and Schmid, 2011; Dong *et al.*, 2012). In
several crops including maize, there has been significant progress in understanding the multigenic
architectures for variation in days to flowering time and photoperiod sensitivity, reflective of the
complex network structure of flowering time regulation. We propose that constructing and
decomposing the PRN-FTP from METs on large collections of genotypes can be used to unravel the
variation linked to some of these specific pathways. Estimates for critical photoperiods and photoperiod
sensitivities from the PRN-FTP can be expected to capture interaction variation between the circadian
clock and the photoperiod response pathways (Itoh *et al.*, 2010), along with the possible partial effects
of the gibberellic acid pathway suggested by some observations (Johansson and Staiger, 2015; Izawa,
2021). Correcting for latent variables should remove some of the effects of other environment-
dependent responses that influence flowering time, leading to more accurate estimates of the variation
in the photoperiodic responses. We expect that flowering time *per se* primarily reflects endogenous
pathways including the autonomous, aging, and gibberellic acid pathways, whereby flowering time is
regulated independent of daylength (Minow *et al.*, 2018). The use of thermal time, however, negates
the analysis of variation arising from temperature effects, which is an emerging pathway in the
regulation of flowering time (Lee *et al.*, 2013; Jin *et al.*, 2023).

Predating the advent of modern varieties, populations of maize expanded from their tropical origins
into temperate zones (Tenaillon and Charcosset, 2011; Swarts *et al.*, 2017). This was facilitated by
flowering time adaptations, evidenced by selection on separate modules of the gene regulatory network
for flowering time (Gaudinier and Blackman, 2020). Distinct breeding pools for maize were then
established from locally preadapted populations, which have largely persisted as separate pools of
tropical and temperate maize. Thus, we reasoned that the PRN-FTP trait space of photoperiod
sensitivity and flowering time *per se* for modern varieties could be used to infer about historical
selection on flowering time pathways in maize. Figure 4 corroborates that distinct selection on these
pathway-associated components of flowering time have structured the plasticity of tropical and

temperate maize. However, the overall trend of reducing photoperiod sensitivity with flowering time *per se* is not necessarily indicative of genetic or physiological coupling of the regulatory pathways, such that genotypes with short flowering time *per se* and high photoperiod sensitivity could exist. Instead, this appears to be a byproduct of the preadaptation history and subsequent breeding of maize. This is supported by results from experimental evolution for early flowering time in tropical populations where photoperiod sensitivity is rapidly eliminated while flowering time *per se* is continually shortened by selection in LD environments of the temperate zone (Teixeira *et al.*, 2015; Choquette *et al.*, 2023). In contrast, selection in SD environments of the tropical zone has no effect on photoperiod sensitivity, where the longer growing seasons have limited the selection for a reduced flowering time *per se*. This highlights how the PRN-FTP can be used for studying crop diversity, and that knowledge about the historical context of adaptation and breeding is crucial for interpreting the observed patterns in the PRN-FTP-derived trait space.

Conclusions

In this study, we explored various aspects of a physiological reaction norm for modeling flowering time plasticity in maize. Envirotyping with DL^s is a valuable tool for examining the global distribution of DL^s in maize and for designing structured METs to investigate the diversity of components of the PRN-FTP. Notably, current MET data is sparse which makes it difficult to characterize the variation in critical thresholds for maize, warranting further experimentation.

The PRN-FTP model provides an ecophysiological framework for understanding (GxE) variation in flowering time. For a specific genotype, the pace of development towards reproductive transition is affected by temperature fluctuations within an environment, with the magnitude of this influence varying across environments based on the unique temperature patterns. When contrasting genotypes are present in the same environments, interactions between temperatures and genetic variation in rates of development and T_t^{em-TI} gives rise to variation in the timing of photoperiod sensitization. In turn, genetic variation in critical thresholds and photoperiod sensitivities lead to secondary interactions with daily rates of photoperiodic change within and across environments. In essence, as genotypes become sensitized to photoperiod at different times, genetic variation in their response to photoperiod further alters the timing of reproductive transition and flowering time, whether or not they are exposed to the same photoperiod within an environment. This highlights the intricate interplay between genetic and environment effects that shape the complex trait variation of flowering time.

Lastly, as the PRN-FTP model is based on common ecophysiological processes, we expect that it can be extended to other crops. Investigating the trait space of PRN-FTP components can provide valuable insights into flowering time adaptations in various species. Our findings underscore the significance of considering the historical context of adaptation and breeding in unraveling the genetic and physiological underpinnings of complex traits. This, in turn, holds promising implications for future crop improvement endeavors.

Acknowledgements: This study was made possible thanks to the production and availability of datasets from prior and ongoing projects: the EXPOSE and PhotoGrid projects at INRAE-LEPSE (ANR-16-IDEX-0006; France 2030 program; Post AgreenSkills Fund), the INVITE project (European Union's Horizon 2020 research and innovation program, grant no. 817970), the Maize ATLAS project (USDA-ARS, grant no. 2011-67003-30342), the Multiple Disease Resistance project (North Carolina State University, Cornell University, and University of Delaware collaboration), the Genomes-to-Fields Initiative (https://www.genomes2fields.org/home/), and the Panzea project (NSF-PGRP, grant

no. 1238014). We wish to thank a number of people who contributed to this study: (i) Mainassara
Abdou Zaman-Allah, Terence Molnar, and Prasanna Boddupalli from CIMMYT for providing much
of the germplasm used for panel 1; (ii) Alexis Bédiée, Llorenç Cabrera-Bosquet, Nicole Choquette,
Arnaud Desbiez-Piat, Thomas Laisné, Noé Lalouette-Marrier d'Unienville, Maëlle Lis, Anthony
Rosello, Javier Soto Mendoza, and Benoît Suard for supporting the PhenoArch platform experiments
at Montpellier, FR; (iii) Arnaud Brunet, Bernard Lagardere, and Jean-Rene Loustalot from the
experimental station at Saint-Martin-de-Hinx, FR for supporting germplasm management and field
testing; (iv) the ISRA team Ibrahima Sarr, Malick Ndiaye, Awa Mbengue, Abdoulaye Hoth, and Alain
Audebert (CIRAD) for field trial data from Nioro du Rip, SN; and (v) the Moreno Retis team José
Moreno and Luis Rodríguez for field trial data from Puerto Vallarta, MX.

Author Contributions: JD, RJW: conceptualization; CP, CW, RJW: resources; JD, CP, EM, RJW:
investigation; JD, CP, JR, RJW: data curation; JD, PM, BP, RJW: methodology; JD, JR, KJ, RJW:
formal analysis; JD, RJW: writing - original draft; EM, JR, PM, KJ: writing - review & editing; BP,
CW, RJW: supervision; EM, RJW: funding acquisition.

Conflict of Interest: The authors declare that the research was conducted in the absence of any
commercial or financial relationships that could be construed as a potential conflict of interest.

Funding Statement: This work was supported by the French National Research Agency (ANR-16-
IDEX-0006), the France 2030 program, project EXPOSE (PAF_18) supported by INRAE, and the
European Union's Horizon 2020 research and innovation program, grant no. 817970.

Data Availability: Data and metadata from this study are available in Supplementary Files at Dryad
(https://doi.org/10.5061/dryad.x95x69pth). Sources of external data are indicated in the methods
sections. Code for fitting the PRN-FTP is available at: https://github.com/maizeatlas/prnftp.

References

Andrieu B, Hillier J, Birch C. 2006. Onset of sheath extension and duration of lamina extension are major determinants of the response of maize lamina length to plant density. Annals of Botany **98**, 1005–1016.

Blümel M, Dally N, Jung C. 2015. Flowering time regulation in crops-what did we learn from Arabidopsis? Current Opinion in Biotechnology **32**, 121–129.

Bonnett OT. 1954. The inflorescences of maize. Science **120**, 77–87.

Bradshaw AD. 1965. Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics **23**, 207–215.

Buckley LB, Briones Ortiz BA, Caruso I, John A, Levy O, Meyer A V., Riddell EA, Sakairi Y, Simonis JL. 2023. TrenchR: An R package for modular and accessible microclimate and biophysical ecology. PLoS Climate **2**, e0000139.

Cabrera-Bosquet L, Fournier C, Brichet N, Welcker C, Suard B, Tardieu F. 2016. High-throughput estimation of incident light, light interception and radiation-use efficiency of thousands of plants in a phenotyping platform. The New Phytologist **212**, 269–281.

Choquette NE, Holland JB, Weldekidan T, Drouault J, de Leon N, Flint-Garcia S, Lauter N, Murray SC, Xu W, Wisser RJ. 2023. Environment-specific selection alters flowering-time plasticity and results in pervasive pleiotropic responses in maize. The New Phytologist **238**, 737–749.

Colasanti J, Coneva V. 2009. Mechanisms of floral induction in grasses: Something borrowed, something new. Plant Physiology **149**, 56–62.

Coles ND, McMullen MD, Balint-Kurti PJ, Pratt RC, Holland JB. 2010. Genetic control of photoperiod sensitivity in maize revealed by joint multiple population analysis. Genetics **184**, 799–812.

Cooper M, Tang T, Gho C, Hart T, Hammer G, Messina C. 2020. Integrating genetic gain and gap analysis to predict improvements in crop productivity. Crop Science **60**, 582–604.

Dong Z, Danilevskaya O, Abadie T, Messina C, Coles N, Cooper M. 2012. A gene regulatory network model for floral transition of the shoot apex in maize and its dynamic modeling. PLoS ONE **7**, e43450.

Finlay KW, Wilkinson GN. 1963. The analysis of adaptation in a plant-breeding programme. Australian Journal of Agricultural Research **14**, 742–754.

Flint-Garcia SA, Thuillet AC, Yu J, Pressoir G, Romero SM, Mitchell SE, Doebley J, Kresovich S, Goodman MM, Buckler ES. 2005. Maize association population: A high-resolution platform for quantitative trait locus dissection. Plant Journal **44**, 1054–1064.

Fong Y, Chongzhi D, Huang Y, Gilbert PB. 2017*a*. Model-robust inference for continuous threshold regression models. Biometrics, **73**, 452–462.

Fong Y, Huang Y, Gilbert PB, Permar SR. 2017*b*. chngpt: Threshold regression model estimation and inference. BMC Bioinformatics **18**, 1–7.

Forsythe WC, Rykiel EJ, Stahl RS, Wu H i., Schoolfield RM. 1995. A model comparison for daylength as a function of latitude and day of year. Ecological Modelling **80**, 87–95.

Gaudinier A, Blackman BK. (2020). Evolutionary processes from the perspective of flowering time diversity. The New Phytologist, **225**, 1883–1898.

Guo T, Wei J, Li X, Yu J. 2023. Environmental context of phenotypic plasticity in flowering time in
sorghum and rice. Journal of Experimental Botany **75**, 1004–1015.

Harrison MT, Tardieu F, Dong Z, Messina CD, Hammer GL. 2014. Characterizing drought stress
and trait influence on maize yield under current and future conditions. Global Change Biology **20**, 867–
878.

Herrero MP, Johnson RR. 1981. Drought stress and its effects on maize reproductive systems. Crop
Science **21**, 105–110.

Hung HY, Shannon LM, Tian F, *et al.* 2012. *ZmCCT* and the genetic basis of day-length adaptation
underlying the postdomestication spread of maize. Proceedings of the National Academy of Sciences
109, E1913-E1921.

Itoh H, Nonoue Y, Yano M, Izawa T. 2010. A pair of floral regulators sets critical day length for
Hd3a florigen expression in rice. Nature Genetics **42**, 635–638.

Izawa T. 2021. What is going on with the hormonal control of flowering in plants? Plant Journal **105**,
431–445.

Jarquín D, Crossa J, Lacaze X, *et al.* 2014. A reaction norm model for genomic selection using high-
dimensional genomic and environmental data. Theoretical and Applied Genetics **127**, 595–607.

Jin M, Liu H, Liu X, *et al.* 2023. Complex genetic architecture underlying the plasticity of maize
agronomic traits. Plant Communications **4**, 100473.

Johansson M, Staiger D. 2015. Time to flower: Interplay between photoperiod and the circadian
clock. Journal of Experimental Botany **66**, 719–730.

Kiniry JR. 1991. Maize Phasic Development. In Modeling Plant and Soil Systems (eds J. Hanks and
J.T. Ritchie). pp 55–70.

Kiniry JR, Ritchie JT, Musser RL. 1983*a*. Dynamic nature of the photoperiod response in maize.
Agronomy Journal **10**, 10–13.

Kiniry JR, Ritchie JT, Musser RL, Flint EP, Iwig WC. 1983*b*. The photoperiod sensitive interval
in maize. Agronomy Journal **75**, 687–690.

Kusmec A, Srinivasan S, Nettleton D, Schnable PS. 2017. Distinct genetic architectures for
phenotype means and plasticities in *Zea mays*. Nature Plants **3**, 715–723.

Lee SH, Choi CW, Park KM, *et al.* 2021. Diversification in functions and expressions of soybean
FLOWERING LOCUS T genes fine-Tune seasonal flowering. Frontiers in Plant Science **12**, 1–16.

Lee JH, Ryu H-S, Chung KS, Posé D, Kim S, Schmid M, Ahn JH. 2013. Regulation of temperature-
responsive flowering by MADS-box transcription factor repressors. Science **342**, 628–632.

Lejeune P, Bernier G. 1996. Effect of environment on the early steps of ear initiation in maize (*Zea
mays* L.). Plant, Cell and Environment **19**, 217–224.

McFarland BA, Alkhalifah N, Bohn M, *et al.* 2020. Maize genomes to fields (G2F): 2014-2017 field
seasons: Genotype, phenotype, climatic, soil, and inbred ear image datasets. BMC Research Notes **13**,
4–9.

Millet EJ, Kruijer W, Coupel-Ledru A, Alvarez Prado S, Cabrera-Bosquet L, Lacube S,
Charcosset A, Welcker C, van Eeuwijk F, Tardieu F. 2019. Genomic prediction of maize yield
across European environmental conditions. Nature Genetics **51**, 952–956.

Minow MAA, Ávila LM, Turner K, Ponzoni E, Mascheretti I, Dussault FM, Lukens L, Rossi V,
Colasanti J. 2018. Distinct gene networks modulate floral induction of autonomous maize and

photoperiod-dependent teosinte. Journal of Experimental Botany **69**, 2937–2952.

Muller B, Martre P. 2019. Plant and crop simulation models: Powerful tools to link physiology,
genetics, and phenomics. Journal of Experimental Botany **70**, 2339–2344.

Padilla JM, Otegui ME. 2005. Co-ordination between leaf initiation and leaf appearance in field-
grown maize (*Zea mays*): Genotypic differences in response of rates to temperature. Annals of Botany
96, 997–1007.

Parent B, Tardieu F. 2012. Temperature responses of developmental processes have not been affected
by breeding in different ecological areas for 17 crop species. The New Phytologist **194**, 760–774.

Parent B, Turc O, Gibon Y, Stitt M, Tardieu F. 2010. Modelling temperature-compensated
physiological rates, based on the co-ordination of responses to temperature of developmental processes.
Journal of Experimental Botany **61**, 2057–2069.

Rising J, Devineni N. 2020. Crop switching reduces agricultural losses from climate change in the
United States by half under RCP 8.5. Nature Communications **11**, 1–7.

Rogers AR, Dunne JC, Romay C, *et al.* 2021. The importance of dominance and genotype-by-
environment interactions on grain yield variation in a large-scale public cooperative maize experiment.
G3: Genes, Genomes, Genetics **11**.

Roof DA. 1997. Evolutionary Quantitative Genetics. Springer New York, NY

Simpson GG, Dean C. 2002. Arabidopsis, the Rosetta stone of flowering time ? Science **2002**, 285–
289.

Sloat LL, Davis SJ, Gerber JS, Moore FC, Ray DK, West PC, Mueller ND. 2020. Climate
adaptation by crop migration. Nature Communications **11**, 1–9.

Sparks A. 2018. nasapower: A NASA POWER global meteorology, surface solar energy and
climatology data client for R. Journal of Open Source Software **3**, 1035.

Srikanth A, Schmid M. 2011. Regulation of flowering time: All roads lead to Rome. Cellular and
Molecular Life Sciences **68**, 2013–2037.

Swarts K, Gutaker RM, Benz B, *et al.* 2017. Genomic estimation of complex traits reveals ancient
maize adaptation to temperate North America. Science **357**, 512–515.

Teixeira JEC, Weldekidan T, De Leon N, *et al.* 2015. Hallauer's Tusón: A decade of selection for
tropical-to-temperate phenological adaptation in maize. Heredity **114**, 229–240.

Tenaillon MI, Charcosset A. 2011. A European perspective on maize history. Comptes Rendus
Biologies **334**, 221–228.

van Voorn GAK, Boer MP, Truong SH, Friedenberg NA, Gugushvili S, McCormick R, Bustos
Korts D, Messina CD, van Eeuwijk FA. 2023. A conceptual framework for the dynamic modeling
of time-resolved phenotypes for sets of genotype-environment-management combinations: a model
library. Frontiers in Plant Science **14**, 1–14.

Wang E, Engel T. 1998. Simulation of phenological development of wheat crops. Agricultural
Systems **58**, 1–24.

Wang E, Martre P, Zhao Z, *et al.* 2017. The uncertainty of crop yield projections is reduced by
improved temperature response functions. Nature Plants **3**, 1–11.

Wisser RJ, Fang Z, Holland JB, *et al.* 2019. The genomic basis for short-term evolution of
environmental adaptation in maize. Genetics **213**, 1479–1494.

Yang Q, Li Z, Li W, *et al.* 2013. CACTA-like transposable element in *ZmCCT* attenuated photoperiod

sensitivity and accelerated the postdomestication spread of maize. Proceedings of the National
Academy of Sciences **110**, 16969–16974.

Zabel F, Müller C, Elliott J, *et al.* 2021. Large potential for crop production adaptation depends on
available future varieties. Global Change Biology **27**, 3870–3882.

List of Tables and Figures

Table 1. Distribution percentiles for the basic vegetative phase and leaf tip appearance at tassel initiation of panel 1.

Table 2. Impact of latent variable adjustment on PRN-FTP parameter estimation for different maize lines.

Figure 1. Envirotyping of sensed photoperiod across and within maize fields in the Northern Hemisphere.

Figure 2. Adjusting for latent variable effects across a multi-environment trial network when fitting the PRN-FTP.

Figure 3. The PRN-FTP for seven tropical lines of maize.

Figure 4. The trait space for flowering time *per se* and photoperiod sensitivity in a diversity panel of maize.

678 **Table 1. Percentiles of the distribution for the basic vegetative phase at tassel initiation (BVP-ti)**
679 **in panel 1.**

	Min	5%	25%	50%	75%	95%	Max
BVP-ti (ºCd)	183	236	284	320	356	427	499

681 **Table 2. Impact of latent variable adjustment on PRN-FTP parameter estimation for different**
682 **maize lines.**

	Genotype	Intercept (°Cd)	Threshold (h)	Slope (°Cd h⁻¹)	RMSE
Unadjusted values					
	CML10	848.3 ± 28.7	12.4 ± 0.7	46.4 ± 10.0	81.6
	CML258	934.1 ± 33.4	12.3 ± 0.7	42.5 ± 11.4	95.2
	CML277	855.8 ± 22.0	14.5 ± 0.6	96.4 ± 18.7	75.4
	CML341	802.1 ± 24.6	12.3 ± 0.6	49.8 ± 8.5	71.4
	CML373	813.9 ± 30.5	12.3 ± 1.1	36.1 ± 10.6	67.3
	Tzi8	855.8 ± 29.5	12.3 ± 1.1	30.2 ± 9.6	86.2
	Tzi9	808.0 ± 29.2	12.2 ± 1.1	26.5 ± 9.2	84.0
Adjusted values					
	CML10	893.3 ± 14.1	14.5 ± 0.6	84.4 ± 12.9	47.9
	CML258	953.1 ± 21.3	12.3 ± 0.9	37.8 ± 7.3	60.7
	CML277	873.7 ± 8.7	14.7 ± 0.6	98.6 ± 8.8	31.3
	CML341	845.0 ± 10.1	14.3 ± 0.6	83.2 ± 7.7	34.8
	CML373	850.7 ± 11.9	12.3 ± 0.8	27.2 ± 4.1	26.2
	Tzi8	888.4 ± 14.8	14.5 ± 1.1	49.0 ± 11.9	53.3
	Tzi9	840.3 ± 11.1	14.4 ± 0.9	42.3 ± 8.7	39.6

683 Root mean squared error (RMSE) is reported for the full bilinear model.

Figure 1. Envirotyping of sensed photoperiod across and within maize fields in the Northern Hemisphere.

(A) Sensed photoperiod (DL^S for the 50th percentile maturity reference) across maize field sites in the Northern Hemisphere. (B) Within environment differences in hours of sensed daylength (ΔDL^S) between the 95th and 5th percentile maturity groups defined by the BVP-ti distribution of panel 1. (C) The figure portrays how different maturity groups can reach the photoperiod sensitized stage on different days of the year within the same environment, leading to differences in DL^S (the 5th, 50th, and 95th percentile values of the BVP-ti in panel 1are indicated as early, reference, and late maturity groups) (D) The relationship between within environment differences in DL^S and rates of daily photoperiodic change. Here, the differences are shown in minutes for comparisons between the reference maturity group relative to the early and late maturity groups. As reflected in (B), the boxplots show that genotypes within the same environment experience increasingly greater differences in photoperiods when rates of photoperiodic change increase, which occurs in association with increasing latitude.

Figure 2. Adjusting for latent variable effects across a multi-environment trial network when
fitting the PRN-FTP.

(A) The plot shows the PRN-FTP for genotype CML341, using unadjusted BLUEs for thermal time
from crop emergence to anthesis (T_t^{em-ant}; black points). The grey lines show the estimated flowering
time *per se* (horizontal line), critical threshold value (vertical line), and photoperiod sensitivity (sloped
line). (B) Deviations from the overall average T_t^{em-ant} for the control groups of photoperiod
insensitive lines estimated across the same environments, with the environment identifiers ordered as
in panels (A) and (B) according to sensed daylength (DL^s). (C) The PRN-FTP for genotype CML341
after adjusting the BLUEs by deviations in (B), leading to different estimates for the flowering time
per se, the critical threshold value, and the photoperiod sensitivity.

Figure 3. The PRN-FTP for seven tropical lines of maize.

(A) For each genotype, the plots show the three PRN-FTP parameters estimated from the bilinear
function fitted using latent-variable adjusted BLUEs for thermal time from crop emergence to anthesis
(T_t^{em-ant}) in relation to sensed daylength (DL^s). Parameter estimates are shown for the flowering time
per se (blue line), the critical threshold value (red line), and the photoperiod sensitivity (yellow line).

Figure 4. The trait space for flowering time *per se* and photoperiod sensitivity in a diversity panel of maize.

Flowering time *per se*

Points correspond to genotypic values for flowering time *per se* (ºCd) and photoperiod sensitivity (ºCd h⁻¹), estimated using the LD-SD regression method on latent-variable adjusted BLUEs. Breeding pools are represented by samples of inbred lines previously assigned to temperate (*n*=137), admixed (*n*=62), and tropical (*n*=66) sources based on structure analysis of marker data (Flint-Garcia *et al.*, 2005).