Dynamics and Activation of Membrane-Bound B Cell Receptor Assembly

Hung N. Do[1], Mingfei Zhao[1], S. Munir Alam[2,3], and S. Gnanakaran[1,*]

[1]Theoretical Biology and Biophysics Group, Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA; [2]Duke Human Vaccine Institute, Duke School of Medicine; [3]Department of Medicine, Duke School of Medicine, Durham, North Carolina 27710, USA

[*]Correspondence: gnana@lanl.gov

Abstract

B-cell receptor complexes (BCR) are expressed on the surface of a B-cell and are the critical regulators of adaptive immune response. Even though the relevance of antibodies has been known for almost a hundred years, the antigen-dependent activation of antibody-producing B-cells has remained elusive. Several models have been proposed for BCR activation, including cross-linking, conformation-induced oligomerization, and dissociation activation models. Recently, the first cryo-EM structure of the human B-cell antigen receptor of the IgM isotype was published. Given the new asymmetric BCR complex, we have carried out extensive molecular dynamics simulations to probe the conformational changes upon antigen binding and the influence of the membrane. We identified two critical dynamical events that could be associated with antigen-dependent activation of BCR. First, antigen binding caused increased flexibility in regions distal to the antigen binding site. Second, this increased flexibility led to the rearrangement of helices in transmembrane helices, including the relative interaction of Igα/Igβ, which has been responsible for intracellular signaling. Further, these transmembrane rearrangements led to changes in localized lipid composition. Even though the simulations considered only a single BCR complex, our work indirectly supports the dissociation activation model.

Introduction

      B-cell receptors (BCRs) stand sentry on the front lines of the body's defenses against infection, functioning by binding foreign substances known as antigens and activating the immune system[1]. They are a complex of *IgM* containing dimer of antigen-binding the Fab domains, Fc domains, transmembrane helices (TM), and the Igα/Igβ heterodimer signaling component[1]. Despite BCRs' importance in adaptive immunity, mechanistic details of antigen-dependent activation remain elusive. However, the BCRs are known to be activated and clustered on the plasma membrane upon antigen binding[2,3], which also triggers signal transduction by Igα and Igβ[3,4]. A recent study also showed that BCRs' activation depends on the lipid membrane domains, and activated BCRs prefer more ordered domains in the plasma membrane vesicles[5].

      Several different models have been proposed for the BCR activation mechanism[2,4,6,7]. The first model, the cross-linking model, assumed the initiation of BCR signaling resulted from the aggregation of monomeric BCR as multivalent antigens were observed to dominantly activate B cells[7-10]. This model drew attention because it proposed no signal needed to be propagated from the extracellular domain to the cytoplasm of the BCR[7-10]. Instead, the aggregation of the Igα-Igβ intracellular domains caused the cross-phosphorylation of the ITAMs motifs (the immunoreceptor tyrosine-based activation motif) by associated tyrosine kinases[7-10]. Two alternative models, namely the conformation-induced oligomerization model and dissociation activation model, have been proposed since the classical cross-linking model was considered too simplistic to account for diverse types of antigens that bound to the BCRs[4,7,11-15]. Based on the conformation-induced oligomerization model, the binding of membrane antigens activated the ectodomains of the BCRs and exposed an oligomerization interface in the membrane proximal external region (MPER)[11,12]. With the exposure of this interface, the BCRs oligomerized, leading to perturbations of the local lipid environment, opening of the cytoplasmic domains, and initiation of signaling[11,12]. In the dissociation activation model, antigen binding dissociated the auto-inhibitory clusters of the resting BCRs on the B-cell surface, causing the opening of the Igα/Igβ heterodimer intracellular ends to expose their ITAM phosphorylation motifs[7,13-15].

Recent cryo-EM shows that BCR is an asymmetric complex where the membrane-bound immunoglobulin molecule (IgM) binds to a single Igα/Igβ heterodimer with a 1:1 stoichiometry[3]. This contrasts with the previous conventional (textbook) belief that it is a symmetric molecule where it forms an assembly with two Igα/Igβ heterodimers in a 1:2 complex[16]. Given this confirmation of the structural arrangement of the BCR complex, we have carried out extensive molecular dynamics (MD) simulations to evaluate the dynamics of this asymmetric complex. We consider both all-atom (AA) and coarse-grained (CG) MD simulations of the CH31, a human B-cell antigen receptor of the IgM isotype, to profile the different protein conformations of this assembly in the presence/absence of the antigen, HIV-1 envelope (Env) protein (including the gp120 envelope protein), in a B-cell membrane mimic, which included 63.2% phosphatidylcholine (POPC), 12.6% phosphatidylethanolamine (POPE), 17.4% palmitoyl sphingomyelin (PSM), 0.5% ceramide (CER3), 2.2% diacylglycerol (DAGL), and 4.2% cholesterol (CHOL), based on the experimental percentage of displayed lipid types of the plasma membrane determined from a previous study[17] (**Figure 1**). We uncover never-before-seen molecular allosteric alterations in dynamics when the antigen is bound to the BCR complex. We expect this work to contribute to understanding the antigen-dependent activation mechanism of BCRs.

Results

Description of BCR complex and antigen

The initial model for the BCR complex was based on the cryo-EM structure of the human B-cell antigen receptor of the *IgM* isotype with the VRC01 Fab domain (PDB: 7XQ8)[3]. The BCR structure is a "Y"-shaped complex of *IgM* containing dimer of the VRC01 Fab domains connected to the Fc domains, which were then connected to the TM helices, which associate to the Igα/Igβ heterodimer signaling component[3,7]. Since our antibody of interest, VRC-CH31, was of the same isotype as the VRC01, we kept the sequences and structures of the Fc domains, TM helices, and Igα/Igβ heterodimer as in the 7XQ8[3] PDB structure and replaced the VRC01 Fab with the VRC-CH31 Fab to model the CH31 BCR using the SWISS-MODELLER[18] homology modelling webserver. The antigen HIV-1 Env was taken from the

6NNJ[19] PDB structure and computationally docked into the model CH31 BCR using the HDOCK[20] webserver to prepare the model CH31 BCR bound by the HIV-1 Env (**Figures 1** and **S2**). Two different simulation systems of the model CH31 BCR were prepared in the presence/absence of the antigen (**Figure 1**). A close look at the HIV-1 Env bound Fab domains in the starting model revealed that the HIV-1 Env primarily interacted with the CDR3 regions of one of the Fab domains while it had minor interactions with the other Fab domain as well. (**Figure S1**). Five different MD simulation replicas of 500 ns each were performed on each CH31 BCR simulation system (**Table S1**).

Increase in the flexibility of BCR upon antigen binding

We calculated the average root-mean-square fluctuations (RMSFs) from all five MD simulation replicas for each BCR simulation system. We then determined the changes in the flexibilities of the BCR upon antigen binding by subtracting the RMSF of the BCR system without the HIV-1 Env from that of the corresponding system bound by the HIV-1 Env (**Figure 2**).

Overall, binding of the antigen to the BCR increased the flexibility of the BCR (**Figure 2a**). Antigen binding also increased the flexibility of the Fab domains, except for both variable parts (Fv) where the HIV-1 Env bound to the receptor (**Figure 2b**), the whole Fc domains, including their MPERs, and the extracellular domains (ECDs) of Igα and Igβ (**Figure 2a** and **2c**). However, the membrane helices turned slightly more rigid upon antigen binding (**Figure 2a** and **2c**). Seemingly, antigen binding to the BCR at the extracellular Fab domains propagated dynamic changes throughout the BCR, opposing the classical cross-linking model of BCR activation[7-10], and increased the flexibility of the MPER (including the C_{H3} and C_{H4} regions of the Fc domains and ECDs of Igα and Igβ), supporting the conformation-induced oligomerization model[11,12].

Allosteric changes in BCR transmembrane helices upon antigen binding

We employed the HELANAL[21] module within the MDAnalysis[22,23] python package to calculate the global tilt angles with respect to the vertical axis of the four membrane helices

(TM1, TM2, Igα, and Igβ) from the MD simulations of the BCR simulation systems to determine the effects of antigen binding on the orientations of the receptor within the membrane (**Figure 3**). The average and standard deviations of the global tilt angles of the membrane helices calculated from the MD simulations of the BCR systems in the membrane were included in **Figure 3**. The time courses of the global tilt angles were plotted in **Figure S2**, and the distributions of the global tilt angles of the four membrane-bound domains sampled from the MD simulations were shown in **Figure S3**.

Overall, antigen binding in the BCR seemingly reduced the global tilt angles of the membrane helices (**Figure 3**). Noticeably, the TM2, which belonged to the second IgM subunit that showed the primary interactions with the HIV-1 Env, was much less tilted than TM1. The Igα, which interacted more closely with TM2 than TM1, was in turn much less tilted than Igβ, which lay closer to TM1 than TM2 (**Figure 3**). In particular, the average global tilt angle of the TM1 reduced from 36.1° ± 12.2° to 27.3° ± 6.5°, of the TM2 reduced from 15.1° ± 6.0° to 9.0° ± 4.3°, of the Igα reduced from 14.3° ± 6.1° to 11.2° ± 4.9°, and of the Igβ reduced from 34.8° ± 8.3° to 25.5° ± 8.8° upon antigen binding (**Figure 3**). Therefore, antigen binding at the extracellular Fab domains reoriented the membrane helices, again opposing the classical cross-linking model[7-10] and supporting the conformation-induced oligomerization and dissociation activation models of BCR activation[7,11-15].

We then performed principal component analysis (PCA) on the MD simulations of every BCR simulation system to identify the representative low-energy conformational states of the BCR complexes (**Figures S4-S6**). The BCR was relatively rigid as similar conformations were observed at the Fab domains between different states (**Figures S4-S6**). Due to the rigidity of the two Fab domains, they always stayed close to each other, causing the model antigen to interact with both Fab domains (**Figures S5-S6**). Furthermore, as antigen bound to the BCR, the membrane helices turned more upright (**Figures S4-S5**), being consistent with our calculations of the global tilt angles in **Figure 3**.

Induction of signaling motifs upon antigen binding

To determine the effects of the antigen binding on the signaling motif within the BCR, we calculated the changes in residue contact frequencies (Δ(frequency)) between Igα and Igβ upon antigen binding in the BCR. The changes in residue contact frequencies between the transmembrane regions of Igα and Igβ upon antigen binding are shown in **Figure 4**. Overall, we observed that the loop that connected the ECD and the membrane helix of Igα wrapped around the Igβ, significantly increased the frequencies of residue contacts within the Igα/Igβ around this region (**Figure 4a**). In particular, notable residue pairs that displayed significantly increased contacts upon antigen binding to the BCR included Igα residue L163 – Igβ residue P176 (with a Δ(frequency) of 0.25), Igα residue E138 – Igβ residue R154 (a possible ionic bond, with a Δ(frequency) of 0.32), Igα residue K141 – Igβ residue K158 (a possible hydrogen bond, with a Δ(frequency) of 0.47), Igα residue N142 – Igβ residue K158 (a possible hydrogen bond, with a Δ(frequency) of 0.51), Igα residue E138 – Igβ residue K158 (a possible ionic bond, with a Δ(frequency) of 0.76), Igα residue F133 – Igβ residue D159 (with a Δ(frequency) of 0.89), Igα residue G137 – Igβ residue N155 (with a Δ(frequency) of 0.93), and Igα residue L134 – Igβ residue D159 (with a Δ(frequency) of 1.0) (**Figure 4a** and **4c**). As residues F133, L134, and G137 of the Igα were all located at the loop connecting the ECD and membrane-bound helix of the Igα, their significant increases in contact with the Igβ confirmed our observation that antigen binding to the BCR led to the wrapping of the Igα connecting loop around Igβ. Furthermore, we observed noticeable reduced contacts towards the C-terminal ends of Igα and Igβ, including Igα residue P159 – Igβ residue V175 (with a Δ(frequency) of -0.25) (**Figure 4a-4b**). This result supported the dissociation activation model[7,13-15], consistent with the fact that the changes in Igα/Igβ heterodimer signaling component may lead to the exposure of their intracellular ITAM phosphorylation motifs (**Figure 4a-4b**).

Lipid reorganization in response to antigen-induced conformational changes in BCR

In addition to the AA MD simulations, we also performed long timescale coarse grained (CG) molecular dynamics simulations for 5μs, started from the final conformations of each AA MD simulation replica obtained from each BCR simulation system to determine

lipid reorganization upon antigen binding in the BCR (**Figure S7**). We counted the number of lipid molecules within 4Å of the BCR within the last 100 ns of the AA and CG MD simulations of the CH31 BCR simulation systems using the MDAnalysis[22,23] python package (**Figure 5a**).

The initial AA conformation of the CH31 BCR was surrounded by 35 POPC, 8 POPE, 5 PSM, 2 CHOL, 1 DAGL, and 0 CER3 lipid molecules (**Figure S8**). On average, 31.0 ± 2.6 POPC, 8.5 ± 1.7 POPE, 4.1 ± 1.7 PSM, 0.5 ± 0.5 CHOL, 1.3 ± 0.4 DAGL, and 0.3 ± 0.2 CER3 lipid molecules came into contacts with the BCR within the last 100ns of its five AA MD simulations (**Figure 5b**). At the end of the five CG MD simulations, an average number of 35.4 ± 3.5 POPC, 11.3 ± 1.6 POPE, 6.7 ± 2.4 PSM, 0.2 ± 0.8 CHOL, 2.5 ± 0.8 DAGL, and 0.6 ± 0.6 CER3 lipid molecules contacted the receptor (**Figure 5c**). Meanwhile, at the start of the AA MD simulation of the antigen bound BCR, 35 POPC, 7 POPE, 8 PSM, 2 CHOL, 1 DAGL, and 0 CER3 lipid molecules were in contact with the BCR (**Figure S8**). In the last 100ns of the five AA MD simulations of the system, 29.5 ± 2.1 POPC, 6.6 ± 1.4 POPE, 8.6 ± 1.2 PSM, 2.0 ± 0.5 CHOL, 0.6 ± 0.4 DAGL, and 1.0 + 0.1 CER3 lipid molecules surrounded the receptor (**Figure 5b**). At the end of the five CG MD simulations, on average 26.0 ± 3.2 POPC, 4.5 ± 2.3 POPE, 12.6 ± 1.9 PSM, 4.2 ± 1.0 CHOL, 0.9 ± 0.6 DAGL, and 2.5 ± 0.7 CER3 were within 4Å distance of the receptor (**Figure 5c**).

We observed noticeable increases in the numbers of ordered lipid molecules such as PSM, CHOL, and CER3 in the proximity of the BCR upon antigen binding while seeing significant decreases in the numbers of disordered lipid molecules, including POPC, POPE, and DAGL surrounding the BCR (**Figure 5b-5c**). Therefore, conformational changes that occurred due to the binding of the antigen led to lipid reorganization within the proximity of the BCR as the average numbers of different lipid molecules surrounding the BCR were different. This raises the possibility that activated BCR may prefer a different lipid composition than those that are in basal state, being consistent with previous studies[7,11-15].

Membrane lipid properties during the MD simulations of the BCR complexes

We used the LiPyphilic[24] python toolkit to calculate the area per lipid in different membrane environments for both the AA and CG simulations. The average area per lipid molecule was

58.0 $Å^2$ in the BCR simulation system, while the area per lipid molecule was 57.6 $Å^2$ in the antigen-bound BCR simulation system. During the CG MD simulations, the average areas per lipid molecule for the BCR and HIV-1 Env-bound BCR were 57.5 $Å^2$ and 57.2 $Å^2$, respectively. Seemingly, the average areas per lipid were comparable between the AA and CG MD simulations. Furthermore, antigen binding to the BCR appeared to slightly increase the order of membrane lipids in the bilayers as we saw small decreases in the average areas per lipid upon antigen binding.

To confirm our predictions regarding the membrane orders, we further employed the LiPyphilic[24] python toolkit to calculate the average membrane thicknesses obtained from the AA and CG MD simulations of the BCR in the presence/absence of the model HIV-1 Env antigen (**Table S2**). In particular, the average membrane thickness calculated from the AA simulations increased from 36.5 ± 1.4 Å for the BCR to 47.2 ± 1.2 Å for the HIV-1 Env bound BCR (**Table S2**). The average membrane thickness calculated from the CG simulations was comparable to the AA simulations, with the respective values being 38.1 ± 6.1 Å and 48.8 ± 13.7 Å for the BCR and HIV-1 Env bound BCR (**Table S2**). Seemingly, upon antigen binding to the BCR, the average membrane thickness increased (**Table S2**), being consistent with our finding that antigen-bound BCR preferred ordered to disordered lipid bilayers in **Figure 5**.

We also employed the *2D streamplots*[25] module within the *MDAnalysis*[22,23] simulation analysis toolkit to monitor the lipid diffusions during our AA and CG MD simulations (**Figures S9-S12**). Overall, we observed higher lipid diffusion speeds during the CG MD simulations than the AA MD simulations. In particular, the maximum speeds observed for lipid diffusions during the AA MD simulations of the BCR and HIV-1 Env-bound BCR were between ~2.6 - ~2.7 (**Figures S9-S10**), while the maximum speeds observed for lipid diffusions during the CG MD simulations were ~4.0-4.2 (**Figures S11-S12**). We then used the LiPyphilic[24] python toolkit to calculate the average lipid diffusion coefficients within the membranes of the different AA and CG BCR simulation systems (**Table S3**). On average, the CG lipid diffusion coefficients were higher than the AA lipid diffusion coefficients at factors of 2-10 times. In particular, the highest diffusion coefficients calculated from the AA and CG MD simulations of the BCR were $2.2{\times}10^{-6} \pm 4.7{\times}10^{-8}$ nm^2/ns and $1.9{\times}10^{-5} \pm 4.6{\times}10^{-7}$ nm^2/ns, and of the HIV-1 Env bound

BCR were $1.7\times10^{-6} \pm 1.0\times10^{-7}$ nm^2/ns and $9.7\times10^{-6} \pm 7.4\times10^{-7}$ nm^2/ns (**Table S3**). Therefore, the CG MD simulations facilitated lipid motions within the membranes compared to the AA MD simulations due to the flattened free energy surfaces as a result of the employed MARTINI 3[26] force field during the CG simulations.

Discussion

Recently, the first cryo-EM structure of the human BCR of the IgM isotype was published[3]. Given the new structural information of this asymmetric complex, we have carried out extensive molecular dynamics simulations on different complexes of the model CH31 BCR. We profiled the important protein conformations and protein-lipid interactions upon antigen (HIV-1 envelope protein) binding in a complex membrane (**Figure 1**). We related the resulting dynamics captured from the simulations to known models of B-cell activation.

Findings from our work support either the conformation-induced oligomerization model[11,12] or the dissociation activation model[7,13-15] while opposing the classical cross-linking model[7-10] of BCR activation. The classical cross-linking model of BCR activation states that no signal propagation from the extracellular domain to the cytoplasm was necessary for BCR activation[7-10]. Instead, BCR activation is a simple event of the Igα/Igβ aggregation accomplished by binding multivalent antigens to multiple BCRs[7-10]. This work shows that antigen binding at the extracellular Fab domains led to changes in the dynamics throughout the BCR. This is required for the conformation-induced oligomerization and dissociation activation models. Further, the flexibility of the BCR monomer and the lipid reorganization around the BCR upon antigen binding favors dissociation activation model even though our simulations were performed on a single BCR complex.

We identify three critical BCR conformational changes upon antigen binding that show the pathway to its activation. First, antigen binding increased the flexibility of the MPER (including the C_{H3} and C_{H4} regions of the Fc domains and the ECDs of Igα/Igβ) (**Figure 2**), being consistent with the conformation-induced oligomerization model[11,12] as the model proposed that antigen binding opened an oligomerization interface at the MPER. Second, antigen binding increased residue contacts in the transmembrane helices near the MPER of

Igα and Igβ, while reduced residue contacts of the Igα and Igβ in the inner membrane leaflet, being consistent with the dissociation activation model[7,13-15] (**Figure 4**). Third, antigen binding in the BCR clearly reduces the global tilt angles of the transmembrane helices (including TM1, TM2, Igα, and Igβ) of the BCR (**Figures 3**). This tilting of the transmembrane helices of Igα/Igβ heterodimer signaling component can alter the exposure of the ITAM motifs in the cytoplasmic tails required for the downstream signaling.

Lipid reorganization around BCR upon antigen-induced conformational changes indirectly supports the dissociation activation model. We observe that the antigen-bound BCR preferred to be surrounded by more ordered lipid molecules such as PSM, CHOL, and CER3, while the free BCR is surrounded by more disordered lipid molecules such as POPC, POPE, and DAGL (**Figure 5**). These lipid rearrangements are caused by changes in the MPER and transmembrane regions of BCR because of antigen binding. This indicates that, potentially, the antigen-bound BCR prefers a different lipid environment than the free one. Specifically, the tilting towards the normal of the membrane may prefer a composition that is a more ordered domain with a higher membrane thickness. This could be potentially driven by the hydrophobic mismatch caused by the transmembrane rearrangements (**Figures 2-4**). These observations are supported by previous studies that suggested a relocation of BCR from a disordered domain to an ordered domain[7,11-15].

In conclusion, we have uncovered three critical dynamical events that could be associated with antigen-dependent activation of BCR. First, antigen binding caused increased flexibility in regions distal to the antigen binding site. Second, this increased flexibility led to the alterations of IgM transmembrane helices and MPER regions. Third, these alterations, led to changes in relative interaction between Igα and Igβ and the orientations of their transmembrane helices. These changes are expected to influence the exposure or phosphorylation of ITAM motifs in the cytoplasmic tails of Igα/Igβ. These conformational changes could potentially relocate BCR in a different region of the membrane as indicated by the differential preferences of lipids before and after the antigen binding. Even though the simulations considered only a single BCR complex, our work indirectly supports the dissociation activation model. Further work with multiple BCRs and different membrane

environments are warranted to conclusively make the direct connection to dissociation activation model.

Materials and Methods

Simulation system setups

We started from the cryo-EM structure of human B-cell antigen receptor (BCR) of the IgM isotype with the Fab domains of the VRC01 antibody (PDB: 7XQ8)[3]. Since our antibody of interest VRC-CH31 was of the same isotype as the VRC01 antibody, the sequences and structures of the Fc domains, transmembrane helices, Igα, and Igβ domains in the BCR with the Fab domains of VRC-CH31 should be identical to those from the 7XQ8 PDB[3] structure. The sequences of the VRC-CH31 Fab heavy and light chains were retrieved from the UniProt database[27] with UniRef IDs of UPI00038A5F4F and UPI00038A5F3B, respectively, and attached to the rest of the BCR through sequence alignment and SWISS-MODEL template-based homology modelling webserver[18]. SWISS-MODEL did not model the intracellular loops of the Igα/Igβ heterodimer[3] even when we provided the full sequences from the 7XQ8 FASTA file (**Figure S13a**). The important ITAM motif in the intracellular loop of Igβ was also not present in the 7XQ8 FASTA file[3]. We also tried using AlphaFold2-Multimer[28,29] and AlphaFold3[30] in modelling the Igα/Igβ heterodimer with intracellular loops as well as the full BCR structure (**Figure S13b-S13d**). We observed that AlphaFold[28-30] predicted the Igα/Igβ heterodimer intracellular loop to "U-turn" into the membrane (**Figure S13b-S13d**). Therefore, we decided to use the BCR model built by SWISS-MODELLER[18] to proceed with our simulations. The sequences and structure of the HIV-1 envelope protein (including the gp120 envelope protein) were taken from the 6NNJ PDB structure[19] and remodeled to fill in the missing regions using the SWISS-MODEL webserver[18]. The HIV-1 envelope protein was docked into the CH31 BCR using the HDOCK integrated protein-protein docking webserver[20]. To examine both the effects of antigen binding on the dynamics mechanisms of the BCR with VRC-CH31 Fab domains, we set up two different simulation systems, including the CH31 BCR and CH31 BCR bound by the HIV-1 envelope protein (HIV-1 Env) in the complex membrane (**Figure 1**). The complex membrane was modeled based on the experimental

percentage of displayed lipid types of the plasma membrane determined from a previous study[17] to include 63.2% phosphatidylcholine (POPC), 12.6% phosphatidylethanolamine (POPE), 17.4% palmitoyl sphingomyelin (PSM), 0.5% ceramide (CER3), 2.2% diacylglycerol (DAGL), and 4.2% cholesterol (CHOL). The CHARMM-GUI webserver[31-34] was used to set up the initial atomistic simulation systems. We did not model the glycosylation of the CH31 BCR and the HIV-1 envelope protein. The CH31 BCR systems were embedded in the membrane lipid bilayer before being solvated in 0.15 M NaCl solutions. The resulting system sizes ranged from ~1.5 million (for the BCR without HIV-1 Env) to ~2.5 million atoms (for the BCR bound by HIV-1 Env). The CHARMM36m force field parameter sets[35] were used for the proteins and lipids in the all-atom (AA) molecular dynamics (MD) simulations. The elastic networks of MARTINI 3 force field parameter sets[26] were used in the coarse-grained (CG) MD simulations.

Multiscale molecular dynamics simulation protocols

In the multiscale MD simulations, AA simulations were iterated with long timescale CG simulations (**Figure S7**). The protein-protein interactions were refined with AA simulations, while lipid mixing was facilitated during the CG simulations. Here, however, CG simulations were only performed to examine the lipid mixing properties during the AA simulations. In the AA MD simulations, periodic boundary conditions were applied to the simulation systems, and bonds containing hydrogen atoms were restrained with the LINCS[36] algorithm. The electrostatic interactions were calculated using the particle mesh Ewald (PME) summation[37] and the Verlet cutoff scheme[38] with a cutoff distance of 12 Å for long-range interactions. The temperature was kept constant at 310 K using the Nose-Hoover thermostat[39] with a friction coefficient of 1.0 ps^{-1}. The pressure was kept constant at 1.0 bar using the Parrinello-Rahman barostat[40] with semi-isotropic coupling. The pressure coupling constant was set to 5 ps, and the compressibility was set to 4.5×10^{-5} bar^{-1}. The simulation systems were energetically minimized to a maximum of 5,000 steps using the steepest-descent algorithm. Position restraints were applied on the backbone atoms with a force constant of 4,000 kJ.mol^{-1}.nm^{-2}, on the side chain atoms with a force constant of 2,000 kJ.mol^{-1}.nm^{-2}, and on the lipids and

dihedral angles with a force constant of 1,000 $kJ.mol^{-1}.nm^{-2}$. The systems were then equilibrated with the constant number, volume, and temperature (NVT) ensemble for a total of 375,000 steps, with a time step of 1 fs used. The force constants for position restraints were gradually reduced from 4,000 to 2,000 to 1,000 $kJ.mol^{-1}.nm^{-2}$ for backbone atoms, from 2,000 to 1,000 to 500 $kJ.mol^{-1}.nm^{-2}$ for side chain atoms, from 1,000 to 400 $kJ.mol^{-1}.nm^{-2}$ for lipids, and from 1,000 to 400 to 200 $kJ.mol^{-1}.nm^{-2}$ for dihedral angles, after every 125,000 steps. The systems were further equilibrated with the constant number, pressure, and temperature (NPT) ensemble for a total of 750,000 steps, with a time step of 2 fs used. The force constants for position restraints were gradually reduced from 500 to 200 to 50 $kJ.mol^{-1}.nm^{-2}$ for backbone atoms, from 200 to 50 to 0 $kJ.mol^{-1}.nm^{-2}$ for side chain atoms, from 200 to 40 to 0 $kJ.mol^{-1}.nm^{-2}$ for lipids, and from 200 to 100 to 0 $kJ.mol^{-1}.nm^{-2}$ for dihedral angles, after every 250,000 steps. Finally, the systems were equilibrated with a short 25ns conventional MD (cMD) simulation using a time step of 2 fs. Five 500ns cMD production simulations were then performed on each of the BCR simulation systems. The final frame from each AA MD simulation replica of every BCR simulation system was extracted and stripped off all ions and water molecules. The protein and membrane lipid portions were converted to their corresponding CG representation using the martinize2 and vermouth framework (https://github.com/marrink-lab/vermouth-martinize)[41] for the protein, and the *backward.py* (https://github.com/Tsjerk/MartiniTools/blob/master/backward.py) script[42] for the membrane lipids. The CG models of the proteins were built with the side chain corrections (-scfix option)[43] and the elastic network (-elastic option)[44]. The elastic bond force constant was set to 700 $kJ.mol^{-1}.nm^{-2}$ (-ef 700), the lower and upper bound of the elastic bond cutoff were set to 0.5 and 0.9 nm, respectively (-el 0.5 -eu 0.9)[44]. The elastic bond decay factor and power were both set to 0 (-ea 0 -ep 0) to make the bond strengths independent of bond length[44]. The *Insane* (https://github.com/Tsjerk/Insane) software package[45] was used to re-solvate the CG systems of the BCR assemblies and lipids in 0.15 M NaCl solutions, with the box dimensions kept identical to the AA simulations. In the CG MD simulations, periodic boundary conditions were applied to the simulation systems. A time step of 20 fs was used. The electrostatic interactions were calculated using the reaction field method[46] and the

Verlet cutoff scheme[38] with a cutoff distance of 11 Å for long-range interactions. The temperature was kept constant at 310 K using the velocity rescaling thermostat[47] with a friction coefficient of 1.0 ps^{-1}. The pressure was kept constant at 1.0 bar first using the stochastic cell rescaling (C-rescale)[48] during the equilibration stage and then Parrinello-Rahman barostat[40] during the production simulations with semi-isotropic coupling. The pressure coupling constant was set to 5 ps, and the compressibility was set to 3×10^{-4} bar^{-1}. The simulation systems were energetically minimized to a maximum of 500,000 steps given the large simulation system sizes. They were then equilibrated with the constant number, pressure, temperature (NPT) ensemble for a total of 300,000 steps. Position restraints were applied on the systems, with a force constant gradually reduced by half starting from 1,000 to 50 $kJ.mol^{-1}.nm^{-2}$ after every 50,000 steps, except the lipid heads where the force constants were gradually reduced from 200 to 10 $kJ.mol^{-1}.nm^{-2}$. The simulation systems were further equilibrated with a short 100ns cMD simulation. Finally, one 5μs cMD production simulation was performed on each CG simulation system obtained from each replica of the previous AA simulations. All simulations were carried out using the *gmx grompp and gmx mdrun* commands in GROMACS 2022[49] on Los Alamos National Laboratory (LANL) High Performance Computing (HPC) Clusters (**Table S1**).

Simulation analysis

First, we used both the CPPTRAJ[50] simulation analysis tool and the GROMACS 2022[49] simulation package to calculate the changes in root-mean-square fluctuations (ΔRMSF) to determine the changes in flexibility in different domains of the BCR complexes upon the binding of the HIV-1 Env to the BCR. Second, we used the HELANAL[21] module within the MDAnalysis[22,23] python package to calculate the global tilt angles with respect to the vertical axis of the four membrane helices (TM1, TM2, and the Igα/Igβ heterodimer) of the CH31 BCR simulation systems. Third, we performed the principal component analysis (PCA) to determine the important low-energy conformational states observed for each BCR simulation system. Fourth, we used the native contacts module within the MDAnalysis[22,23] python package to calculate the changes in residue contact frequencies between the Igα

and Igβ upon antigen binding. The native contacts module[22,23] was also used to calculate the number of lipid molecules within 4Å distance of the BCR within the last 100ns of the simulation systems. Fifth, we used the LiPyphilic[24] python toolkit to calculate the average area per lipid, average membrane thickness, and average lipid diffusion coefficients in each BCR simulation system. Finally, we employed the *2D streamplots*[25] module within the *MDAnalysis*[22,23] simulation analysis toolkit to monitor the lipid diffusion speed during the AA and CG MD simulations.

Acknowledgements

We would like to thank Cesar Lopez and Tyler Reddy for providing us with valuable insights and suggestions. This study was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health and by the Duke Center for HIV Structural Biology, grant number U54-AI170752-01. The authors would also like to thank the computational resources provided by the Theoretical Biology and Biophysics Group at LANL and LANL Institutional Computing. This work was performed at the Los Alamos National Laboratory, which is operated by Triad National Security, LLC, for the National Nuclear Security Administration of the U.S. Department of Energy (contract 89233218CNA000001). The views expressed in this article are those of the authors and do not reflect the official policy or position of the U.S. Government.

References

1. Robinson, R. (2006). Distinct B Cell Receptor Functions Are Determined by Phosphorylation. PLOS Biology *4*, e231. https://doi.org/10.1371/journal.pbio.0040231.
2. Harwood, N.E., and Batista, F.D. (2010). Early events in B cell activation. Annual Review of Immunology *28*, 185-210. https://doi.org/10.1146/annurev-immunol-030409-101216.
3. Su, Q., Chen, M., Shi, Y., Zhang, X., Huang, G., Huang, B., Liu, D., Liu, Z., and Shi, Y. (2022). Cryo-EM structure of the human IgM B cell receptor. Science *377*, 875-880.
4. Treanor, B. (2012). B-cell receptor: from resting state to activate Immunology *136*, 21-27. 10.1111/j.1365-2567.2012.03564.x. .
5. Shelby, S.A., Castello-Serrano, I., Wisser, K.C., Levental, I., and Veatch, S.L. (2023). Membrane phase separation drives responsive assembly of receptor signaling

domains. Nature Chemical Biology *19*, 750-758. https://www.nature.com/articles/s41589-023-01268-8.

6. Harwood, N.E., and Batista, F.D. (2008). New insights into the early molecular events underlying B cell activation. Immunity *28*, 609-619. 10.1016/j.immuni.2008.04.007

7. Tolar, P., and Pierce, S.K. (2022). Unveiling the B cell receptor structure. Science *377*, 819-820. https://www.science.org/doi/10.1126/science.add8065.

8. Thyagarajan, R., Arunkumar, N., and Song, W. (2003). Polyvalent Antigens Stabilize B Cell Antigen Receptor Surface Signaling Microdomains. The Journal of Immunology *170*, 6099-6106. https://doi.org/10.4049/jimmunol.170.12.6099.

9. Minguet, S., Dopfer, E.-P., and Schamel, W.W.A. (2010). Low-valency, but not monovalent, antigens trigger the B-cell antigen receptor (BCR). International Immunology *22*, 205-212. https://doi.org/10.1093/intimm/dxp129.

10. Kim, Y.-M., Pan, J.Y.-J., Korbel, G.A., Peperzak, V., Boes, M., and Ploegh, H.L. (2006). Monovalent ligation of the B cell receptor induces receptor activation but fails to promote antigen presentation. PNAS *103*, 3327-3332. https://doi.org/10.1073/pnas.0511315103.

11. Reth, M., Wienands, J., and Schamel, W.W. (2000). An unsolved problem of the clonal selection theory and the model of an oligomeric B-cell antigen receptor. Immunological Reviews *176*, 10-18. 10.1034/j.1600-065x.2000.00610.x.

12. Tolar, P., and Pierce, S.K. (2010). A conformation-induced oligomerization model for B cell receptor microclustering and signaling. Current Topics in Microbiology and Immunology *340*, 155-169. 10.1007/978-3-642-03858-7_8.

13. Yang, J., and Reth, M. (2010). Oligomeric organization of the B-cell antigen receptor on resting cells. Nature *467*, 465-469. 10.1038/nature09357.

14. Yang, J., and Reth, M. (2010). The dissociation activation model of B cell antigen receptor triggering. FEBS Letters *584*, 4872-4877. https://doi.org/10.1016/j.febslet.2010.09.045.

15. Yang, J., and Reth, M. (2016). Receptor Dissociation and B-Cell Activation. Current Topics in Microbiology and Immunology *393*, 27-43. 10.1007/82_2015_482.

16. Huse, K., Bai, B., Hilden, V.I., Bollum, L.K., Thea K. Våtsveen, T.K., Munthe, L.A., Smeland, E.B., Irish, J.M., Wälchli, S., and Myklebust, J.H. (2022). Mechanism of CD79A and CD79B Support for IgM+ B Cell Fitness through B Cell Receptor Surface Expression The Journal of Immunology *209*, 2042-2053. https://doi.org/10.4049/jimmunol.2200144.

17. Kwon, H.-Y., Das, R.K., Jung, G.T., Lee, H.-G., Lee, S.H., Berry, S.N., Tan, J.K.S., Park, S., Yang, J.-S., Park, S., et al. (2021). Lipid-Oriented Live-Cell Distinction of B and T Lymphocytes. Journal of the American Chemical Society *143*, 5836-5844. https://doi.org/10.1021/jacs.1c00944.

18. Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F., de Beer, T., Rempfer, C., Bordoli, L., et al. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic Acids Research *46*, W296-W303.

19. Liu, Q., Lai, Y.T., Zhang, P., Louder, M.K., Pegu, A., Rawi, R., Asokan, M., Chen, X., Shen, C.H., Chuang, G.Y., et al. (2019). Improvement of antibody functionality by

structure-guided paratope engraftment. Nature Communications *10*, 721. https://doi.org/10.1038/s41467-019-08658-4.

20. Yan, Y., Tao, H., He, J., and Huang, S.-Y. (2020). The HDOCK server for integrated protein–protein docking. Nature Protocols *15*, 1829-1852. https://doi.org/10.1038/s41596-020-0312-x.

21. Bansal, M., Kumar, S., and Velavan, R. (2000). HELANAL: a program to characterize helix geometry in proteins. Journal of Biomolecular Structure and Dynamics *17*, 811-819. 10.1080/07391102.2000.10506570.

22. Michaud-Agrawal, N., Denning, E.J., Woolf, T.B., and Beckstein, O. (2011). MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. Journal of Computational Chemistry *32*, 2319-2327. https://doi.org/10.1002/jcc.21787.

23. Gowers, R.J., Linke, M., Barnoud, J., Reddy, T.J.E., Melo, M.N., Seyler, S.L., Domanski, J., Dotson, D.L., Buchoux, S., Kenney, I.M., and Beckstein, O. (2016). MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. Proceedings of the 15th Python in Science Conference, 98-105. 10.25080/Majora-629e541a-00e.

24. Smith, P., and Lorenz, C.D. (2021). LiPyphilic: A Python Toolkit for the Analysis of Lipid Membrane Simulations. Journal of Chemical Theory and Computation *17*, 5907-5919. https://doi.org/10.1021/acs.jctc.1c00447.

25. Chavent, M., Reddy, T., Dahl, C.E., Goose, J., Jobard, B., and Sansom, M.S.P. (2014). Methodologies for the analysis of instantaneous lipid diffusion in MD simulations of large membrane systems. Faraday Discussions *169*, 455-475. https://pubs.rsc.org/en/content/articlelanding/2014/fd/c3fd00145h.

26. Souza, P.C.T., Alessandri, R., Barnoud, J., Thallmair, S., Faustino, I., Grunewald, F., Patmanidis, I., Abdizadeh, H., Bruininks, B.M.H., Wassenaar, T.A., et al. (2021). Martini 3: a general purpose force field for coarse-grained molecular dynamics. Nature Methods *18*, 382-388. https://doi.org/10.1038/s41592-021-01098-3.

27. Consortium, T.U. (2023). UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Research *51*, D523-D531. https://doi.org/10.1093/nar/gkac1052.

28. Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Zidek, A., Bates, R., Blackwell, S., Yim, J., et al. (2022). Protein complex prediction with AlphaFold-Multimer. bioRxiv. https://doi.org/10.1101/2021.10.04.463034.

29. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583-589. https://doi.org/10.1038/s41586-021-03819-2.

30. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature, In Press. https://doi.org/10.1038/s41586-024-07487-w.

31. Jo, S., Kim, T., Iyer, V., and Im, W. (2008). CHARMM-GUI: A Web-based Graphical User Interface for CHARMM. Journal of Computational Chemistry *29*, 1859-1865.

32. Wu, E., Cheng, X., Jo, S., Rui, H., Song, K., Davila-Contreras, E., Qi, Y., Lee, J., Monje-Galvan, V., Venable, R., et al. (2014). CHARMM-GUI Membrane Builder Toward

Realistic Biological Membrane Simulations. Journal of Computational Chemistry *35*, 1997-2004.

33. Jo, S., Lim, J., Klauda, J., and Im, W. (2009). CHARMM-GUI Membrane Builder for Mixed Bilayers and Its Application to Yeast Membranes. Biophysical Journal *97*, 50-58.

34. Lee, J., Cheng, X., Swails, J., Yeom, M., Eastman, P., Lemkul, J., Wei, S., Buckner, J., Jeong, J., Qi, Y., et al. (2016). CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations using the CHARMM36 Additive Force Field. Journal of Chemical Theory and Computation *12*, 405-413.

35. Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., de Groot, B.L., Grubmuller, H., and MacKerell Jr, A.D. (2017). CHARMM36m: an improved force field for folded and intrinscially disordered proteins. Nature Methods *14*, 71-73.

36. Hess, B., Bekker, H., Berendsen, H.J.C., and Fraaije, J.G.E.M. (1998). LINCS: A linear constraint solver for molecular simulations. Journal of Computational Chemistry *18*, 1463-1472. https://doi.org/10.1002/(SICI)1096-987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-H.

37. Essmann, U., Perera, L., Berkowitz, M.L., Darden, T., Lee, H., and Pedersen, L.G. (1995). A Smooth Particle Mesh Ewald Method. Journal of Chemical Physics *103*.

38. Grubmuller, H., Heller, H., Windemuth, A., and Schulten, K. (1991). Generalized Verlet Algorithm for Efficient Molecular Dynamics Simulations with Long-Range Interactions. Molecular Simulation *6*, 121-142.

39. Evans, D.J., and Holian, B.L. (1985). The Nose-Hoover thermostat. The Journal of Chemical Physics *83*, 4069-4074. https://doi.org/10.1063/1.449071.

40. Parrinello, M., and Rahman, A. (1981). Polymorphic Transitions in Single Crystals: A New Molecular Dynamics Method. Journal of Applied Physics *52*, 7182-7190. https://doi.org/10.1063/1.328693.

41. Kroon, P.C., Grunewald, F., Barnoud, J., Tilburg, M.v., Souza, P.C.T., Wassenaar, T.A., and Marrink, S.J. (2023). Martinize2 and Vermouth: Unified Framework for Topology Generation. eLife. https://doi.org/10.7554/eLife.90627.1.

42. Wassenaar, T.A., Pluhackova, K., Böckmann, R.A., Marrink, S.J., and Tieleman, D.P. (2014). Going Backward: A Flexible Geometric Approach to Reverse Transformation from Coarse Grained to Atomistic Models. Journal of Chemical Theory and Computation *10*, 676-690. https://doi.org/10.1021/ct400617g.

43. Herzog, F.A., Braun, L., Schoen, I., and Vogel, V. (2016). Improved Side Chain Dynamics in MARTINI Simulations of Protein–Lipid Interfaces. Journal of Chemical Theory and Computation *12*, 2446-2458. https://doi.org/10.1021/acs.jctc.6b00122.

44. Periole, X., Cavalli, M., Marrink, S.-J., and Ceruso, M.A. (2009). Combining an Elastic Network With a Coarse-Grained Molecular Force Field: Structure, Dynamics, and Intermolecular Recognition. Journal of Chemical Theory and Computation *5*. https://doi.org/10.1021/ct9002114.

45. Wassenaar, T.A., Ingolfsson, H.I., Bockmann, R.A., Tieleman, D.P., and Marrink, S.J. (2015). Computational Lipidomics with insane: A Versatile Tool for Generating Custom Membranes for Molecular Simulations. Journal of Chemical Theory and Computation *11*, 2144-2155. https://doi.org/10.1021/acs.jctc.5b00209.

46. Tironi, I.G., Sperb, R., Smith, P.E., and Gunsteren, W.F. (1995). A generalized reaction field method for molecular dynamics simulations. The Journal of Chemical Physics *102*, 5451-5459. https://doi.org/10.1063/1.469273.

47. Bussi, G., Donadio, D., and Parrinello, M. (2007). Canonical sampling through velocity rescaling. The Journal of Chemical Physics *126*, 014101. https://doi.org/10.1063/1.2408420.

48. Bernetti, M., and Bussi, G. (2020). Pressure control using stochastic cell rescaling. The Journal of Chemical Physics *153*, 114107. https://doi.org/10.1063/5.0020514.

49. Abraham, M.J., Murtola, T., Schulz, R., Pall, S., Smith, J.C., Hess, B., and Lindahl, E. (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX *1-2*, 19-25. https://doi.org/10.1016/j.softx.2015.06.001.

50. Roe, D.R., and Cheatham, I.T.E. (2013). PTRAJ and CPPTRAJ: software for processing and analysis of molecular dynamics trajectory data. Journal of Chemical Theory and Computation *9*, 3084-3095.

51. Song, W., Corey, R.A., Ansell, T.B., Cassidy, C.K., Horrell, M.R., Duncan, A.L., Stansfeld, P.J., and Sansom, M.S.P. (2022). PyLipID: A Python Package for Analysis of Protein–Lipid Interactions from Molecular Dynamics Simulations. Journal of Chemical Theory and Computation *18*, 1188-1201. https://doi.org/10.1021/acs.jctc.1c00708.

Figures

Figure 1. Simulation systems of the model B cell receptor (BCR) complexes. (a) The CH31 BCR in the complex membrane. **(b)** The CH31 BCR bound by a monomer of HIV-1 envelope protein (HIV-1 Env) in the complex membrane. The Fab heavy chains are colored red, Fab light chains are colored blue, Fc domains are colored cyan, transmembrane helices of BCR are colored green, Igα is colored magenta, Igβ is colored brown, and the HIV-1 Env is colored gray. The POPC molecules are colored light blue, POPE are colored light orange, PSM are colored light pink, diacylglycerols (DAGL) are colored pale green, cholesterols (CHOL) are colored wheat, and ceramides (CER3) are colored pale cyan.

Figure 2. Changes in the system flexibility as measured by the changes in root-mean-square fluctuations (ΔRMSF) upon antigen binding to the BCR (a). The important domains that showed significant dynamic changes upon antigen binding in the BCR are zoomed in **(b-c)**. A color scale of blue (-3.0) – white (0.0) – red (3.0) is used to show the magnitudes of ΔRMSFs. The HIV-1 Env is colored gray.

Figure 3. Effects of antigen binding to the BCR on the global tilt angles of the transmembrane helices (including TM1, TM2, and the Igα/Igβ heterodimer) normal to membrane (z-axis) as calculated from the MD simulations of the BCR complexes. The antigen was found to mostly interact with the second IgM subunit (containing TM2, marked by *).

System		TM1	TM2*	Igα	Igβ
BCMM	- Antigen	36.1° ± 12.2°	15.1° ± 6.0°	14.3° ± 6.1°	34.8° ± 8.3°
	+ Antigen	27.3° ± 6.5°	9.0° ± 4.3°	11.2° ± 4.9°	25.5° ± 8.8°

Figure 4. Changes in residue contacts between Igα and Igβ upon binding of the antigen to the membrane bound BCR. (a) Changes in residue contact frequencies between Igα and Igβ upon HIV-1 Env binding to the BCR. A contact definition of ≤ 8.0 Å distance between Cα atoms was used. A color scale of blue (-0.2) – white (0) – red (0.2) was used to show the magnitudes of changes in residue contact frequencies. **(b)** Representative reduced contacts upon antigen binding in the BCR. **(c)** Representative increased contacts upon antigen binding in the BCR. The structural residue contacts colored blue were those that showed decreases in contact frequencies upon antigen binding, whereas those colored red showed increased in contact frequencies upon antigen binding.

Figure 5. Reorganization of the local lipid environments upon antigen binding to the BCR in the complex membrane. (a) All lipid molecules within 4Å distance of the membrane-peripheral region and membrane-bound helices of the BCR are considered. **(b-c)** The numbers of surrounding lipid molecules during the last 100 ns of the AA **(b)** and CG MD simulations **(c)** in the complex membrane are shown. The bars for the BCR without and with antigen bound are colored blue and orange, respectively.