AgRP1 modulates breeding season-dependent feeding behavior in female medaka

Yurika Tagui[1], Shingo Takeda[1], Hiroyo Waida[1], Shoichi Kitahara[1], Tomoki Kimura[2], Shinji Kanda[3],

Yoshitaka Oka[1], Yu Hayashi[1], Chie Umatani[1,4]*

1 Graduate School of Science, the University of Tokyo, Tokyo, Japan.

2 Department of Applied Physics, the University of Tokyo, Tokyo, Japan.

3 Atmosphere and Ocean Research Institute, The University of Tokyo, Chiba, Japan

4 Graduate School of Agriculture, Tokyo University of Agriculture and Technology, Tokyo, Japan

*Corresponding author: chie@go.tuat.ac.jp

Abstract

Feeding and reproduction are known to be closely correlated with each other, and the seasonal breeders show breeding season-dependent feeding behavior. However, most model animals do not have definite breeding seasonality, and the mechanisms for such feeding behavior remain unclear. Here, we focused on female medaka (*Oryzias latipes*); they show breeding season-dependent feeding behavior, and their condition of breeding season can be experimentally controlled by day-length. We first demonstrated that, among previously reported feeding-related peptides (neuropeptides involved in feeding), agouti-related peptide 1 (*agrp1*) and neuropeptide y b (*npyb*) show higher brain expression under the breeding condition than under the non-breeding one. Combined with analysis of *agrp1* knockout medaka, we obtained results to suggest that long day-induced sexually mature condition, especially ovarian estrogenic signals, increase the expressions of *agrp1* in the brain, which results in increased food intake to promote reproduction. Our findings advance the understanding of neural mechanisms of feeding behavior for reproductive success.

Introduction

Feeding behavior is essential to animals for their survival and reproduction and is known to be modulated by various internal and external factors: nutritional status, sexual maturity, temperature, seasonality, etc. This behavior is known to be closely correlated with reproduction (Kauffman & Rissman, 2004), which is an essential biological activity important for the animal life. Previous studies reported that nutritional-state modulates reproductive behaviors and functions (Chen et al., 2006; Amirjani et al., 2019; Volk et al., 2017; Lynn et al., 2010). For example, musk shrews show defective sexual behavior under fasted conditions (Temple & Rissman, 2000). In addition, not a few studies demonstrated that fasting-induced low energy condition suppresses reproduction (Evans & Anderson, 2012; Kalra & Kalra, 1996; Kirkwood et al., 1987; Merry & Holehan, 1979; Hasebe et al., 2016). Thus, it has been well investigated how nutritional status resulting from feeding modulates reproduction. On the other hand, it has been reported that some animals change their feeding behavior during the breeding season. For instance, the black seabream migrates to the shallow water during the breeding season (Tsuyuki, 2018; Kawai et al., 2020) where they can get more food, and the white-tailed deer spend more time for feeding under reproductive status (Stone et al., 2017). Such a close relationship between reproduction and feeding is thought to be important for biological fitness. However, the regulatory mechanisms for breeding season-dependent feeding behavior are still unknown. One possible reason is that most of the model animals appear to have lost the well-defined breeding season. Although the mammalian models, mice and rats, and teleost model zebrafish, have reproductive cycles of about 4-5-days (Nilsson et al., 2015; Peute et al., 1978), they do not have definite breeding seasonality. Thus, the mechanisms for long-term changes in feeding behavior according to the breeding season have not yet been studied in detail.

Here, as a model animal for the seasonal breeder, we used a teleost fish, medaka (*Oryzias latipes*). Medaka is a useful model animal, whose reproductive status can be experimentally controlled by day length (Robinson & Rugh, 1943; Egami, 1954) while keeping an appropriate temperature constant. In the long-day (LD) condition (14 h light/10 h dark), female medaka becomes reproductive and regularly spawns every day, while it becomes non-reproductive in the short-day (SD) condition (10 h light/14 h dark). In other words, LD or SD condition can induce breeding or non-breeding season of female medaka, respectively. Thus, medaka enables us to analyze the mechanisms of breeding season-dependent feeding behavior without consideration for possible changes in metabolism and gene expressions due to the changes in ambient temperature.

Regulatory mechanisms of feeding behavior have mainly been analyzed in mammals. These studies reported that some neuropeptides, such as agouti-related peptide (AgRP) and neuropeptide Y (NPY),

are involved in the control of feeding and called "feeding-related peptides" as key molecules for the

regulation of feeding behavior (Hahn et al., 1998; Aponte et al., 2011; Krashes et al., 2011;

Andermann & Lowell, 2017). Teleosts have also been thought to possess a regulatory mechanism for

feeding similar to mammals. In fact, expression of homologous genes coding for feeding-related

peptides have been reported (Rønnestad et al., 2017; Conde-Sieira & Soengas, 2016). On the other

hand, although administration of some of them have been suggested to induce feeding behavior in

teleosts as well (Rønnestad et al., 2017), their functions in feeding behaviors still remain unclear.

In the present study, to understand mechanisms of breeding season-dependent feeding behavior, we

focused on female medaka, which clearly show seasonal changes in breeding conditions by day

length (Mitani et al., 2010; Kanda et al., 2008) under the fixed appropriate temperature. We first

quantified changes in feeding behavior according to the breeding states and found that female

medaka under the condition of breeding season (LD) eat more than those under the condition of

non-breeding (SD). Therefore, we searched for genes that show breeding state-dependent changes in

expression and found some candidates for feeding-related peptides in medaka. We then analyzed

expressions of the candidate genes by using RNA-seq, *in situ* hybridization, and RT-qPCR, and

analyzed phenotypes of gene knockout medaka. These results led us to conclude that AgRP1 plays a

key role in the breeding season-dependent changes in feeding behavior via ovarian estrogenic

signals.

Results

Feeding behavior of female medaka is upregulated in breeding season

To analyze food intake of male and female medaka in breeding/non-breeding seasons, we first established a method for measuring food intake in medaka. In brief, we placed medaka in a white cup, fed brine shrimp to medaka in all-you-can-eat style for 10 min, and counted the leftover brine shrimp in the cup with a "shrimp-counter" system (called Japanese "Wanko-soba" like method, Figure 1-Source Code 1 and Figure 1-Supplementary Figure 1). We used this system to analyze food intake of male and female medaka under the breeding condition equivalent to that in breeding season (kept under long day (LD) condition) or under the non-breeding condition equivalent to that in non-breeding season (short day (SD) condition) (Figure 1A; Kanda et al., 2008). We found that female medaka under the breeding (LD) condition eat more than those under the non-breeding (SD) condition (Figure 1B; $p = 0.02519$). In contrast to female, in males there was no significant difference in food intake between the breeding and non-breeding condition (Figure 1C; $p = 0.6540$). Since these results demonstrated that females, not males, show breeding season-dependent feeding behavior, we focused only on female medaka in the following analyses on neuronal mechanism for breeding season-dependent feeding behavior. Next, to examine which gene products modulate feeding behavior of female medaka in breeding season, we performed mRNA-sequencing (RNA-seq) using the whole brain of female medaka in breeding condition (LD) and non-breeding condition (SD) (Figure 1-Supplementary Figure 2A). Overall, 1025 genes showed significantly different expression between LD and SD female medaka. Figure 1-Supplementary Figure 2B shows a heat map of representative genes mainly related to neuroendocrine system, which were differently expressed between LD and SD females. Among the conventional candidate feeding-related neuropeptides, we identified two kinds of neuropeptides, *agrp1* and *npyb*, both of which showed higher expression in LD than in SD (Figure 1-Supplementary Figure 2C and 2D). Both AgRP and NPY are known to have orexigenic effects mainly in mice (Schwartz et al., 2000; Andermann & Lowell, 2017). Therefore, in the subsequent analyses, we focused on *agrp1* and *npyb* as candidate genes that modulate breeding season-dependent feeding behavior in female medaka.

AgRP1, NPYa and NPYb may be the "feeding-related peptides" in female medaka

Medaka has two *agrp* paralogues, *agrp1* and *agrp2* and two *npy* paralogues, *npya* and *npyb*, which arose from third round whole genome duplication early in the teleost lineage (Liu et al., 2019; Sundstrom et al., 2008). Therefore, we next examined the anatomical distribution of neurons expressing *agrp1*, *npyb*, and their paralogs in the female brain by *in situ* hybridization (ISH). We

found that *agrp1*- and *npyb*-expressing neurons are distributed in local brain regions (Figures 2A, 2B

and 2E), while *npya*- and *agrp2*- expressing neurons are more widely distributed (Figures 2A, 2C

and 2D) in the brain. The *agrp1* neurons were distributed in the nucleus ventralis tuberis (NVT) of

the hypothalamus (Figure 2B), while *agrp2* neurons were expressed in the telencephalon and in the

hypothalamus (Figure 2C). On the other hand, *npya* neurons were distributed more widely from

telencephalon to hypothalamus (Figure 2D). *npyb* neurons were distributed locally in the nucleus

ventralis telencephali pars dorsalis (Vd) of the telencephalon (Figure 2E).

In mice, *agrp* is known to be only expressed in the hypothalamus and mostly co-expressed with *npy*

(Hahn et al., 1998), and these AgRP/NPY neurons are known to regulate mammal feeding behavior

(Shutter et al., 1997; Broberger et al., 1998; Ollmann et al., 1997; Takahashi & Cone, 2005). In

medaka, on the other hand, *agrp1* signals were not observed in *npya* neurons (Figure 2F left),

although the both genes were expressed in the hypothalamus. In addition, *agrp2* signals were not

observed in hypothalamic *npya* neurons (Figure 2F right), either. These results suggest that AgRP

and NPY are not co-expressed in medaka. Since AgRP and NPY of medaka showed different

expressing patterns compared with other animals such as mice, we examined whether they act as

modulators of feeding. We divided female medaka in LD condition into two groups; one group was

kept under normally fed condition (Fed), and the other was kept under 2-week food restricted

condition (FR). We then analyzed whole-brain expressions of these four genes. RT-qPCR analysis

demonstrated that 2-week FR increased the expression of *agrp1* (Figure 2G; $p = 0.001243$) but

decreased that of *npyb* (Figure 2J; $p = 0.0003108$), suggesting that the two peptides are involved in

feeding in an opposite manner. On the other hand, *agrp2* did not significantly change their

expressions between Fed and FR conditions (Figure 2H; $p = 0.9551$). Although *npya* did not

significantly change their expressions between Fed and FR conditions (Figure 2I; $p = 0.2319$), it may

be possible that *npya* expression changed in a specific brain region, since *npya* neurons are widely

distributed in various brain regions as described above (Figure 2D). Since it is suggested that NPY

released from hypothalamic *npy*-expressing neurons controls food intake in mice (Kohno & Yada,

2012), we also examined the *npya*-expression in medaka hypothalamus by ISH. We counted

npya-expressing neurons in each hypothalamic region and compared them between Fed and FR

female medaka (Figure 2-Supplementary Figure 1). We found that the total number of

npya-expressing neurons in the hypothalamus was significantly larger in Fed compared with FR

(Figure 2-Supplementary Figure 1A; $p = 0.02857$). Here, significant increase in cell number was

observed in NRL ($p = 0.02857$) and NAT ($p = 0.02857$), but not in NVT ($p = 0.1143$) (Figure

2-Supplementary Figure 1B-D). On the other hand, the expression of *agrp2* did not show remarkable

difference in the hypothalamus under food restriction or not (Figure 2-Supplementary Figure 1E).

Thus, the results suggest that AgRP1, NPYa and NPYb may be the "feeding-related peptides" in

female medaka.

Both *agrp1* and *npyb* show higher expression levels in LD than in SD female medaka

To further examine the result of RNA-sequencing (Figure 1-Supplementary Figure 2), we compared

expression of *agrp1* and *npyb* between the female medaka under the breeding condition (LD) and

those under the non-breeding condition (SD), using ISH and whole-brain RT-qPCR (Figure 3). First,

we performed whole-brain RT-qPCR and found that the expression level of *agrp1* was higher in LD

than in SD female (Figure 3A, $p = 0.001865$). In ISH experiments, we observed larger number of

agrp1-expressing neurons in LD than in SD females (Figure 3B and 3C ($p = 0.02828$), Figure

3-Supplementary Figure 1). Since the expression level of *agrp1* was higher in LD than that in SD

(Figure 3A), higher expression of *agrp1* under the breeding condition may be due to the increase in

the number of neurons expressing *agrp1*. On the other hand, *npyb* expression in RT-qPCR was

significantly higher in LD than that in SD (Figure 3D; $p = 0.0001554$), although ISH analysis

demonstrated that *npyb*-expressing cell number was not significantly different between LD and SD

(Figure 3E; $p = 0.4206$). These results suggest that the expression level for each neuron increased in

LD compared with SD. Thus, higher expression of *npyb* under the breeding condition may be due to

the increase of expressions in each neuron expressing *npyb*.

In juvenile female medaka, expression levels of neither *agrp1* nor *npyb* shows significant

change according to the day-length

The results thus far indicates that expressions of *agrp1* and *npyb* are upregulated in female medaka

under the condition of breeding season. Since the breeding/non-breeding state is experimentally

controlled by day-length (LD/SD) in the present study, we examined which of the two factors,

day-length itself or substance(s) from LD-induced mature ovary, regulates the expression of *agrp1*

and *npyb*. Here, we used sexually immature juvenile medaka and compared their whole-brain

expressions of *agrp1* and *npyb* under LD/SD conditions using RT-qPCR (Figure 4). We found that

expression levels of neither *agrp1* nor *npyb* show significant difference between LD and SD (Figure

4A ($p = 0.4179$) and 4B ($p = 0.3357$)). Furthermore, food intake of juvenile female was not different

between LD and SD (Figure 4-Supplementary Figure 1; $p = 0.7197$). These results suggest that

neither of them is regulated directly by day length itself. Instead, the gene expression is suggested to

be regulated by LD-induced sexual maturity.

Estrogen, which is released from mature ovary, may affect the expression of *agrp1*

Among various factors associated with ovarian maturity, estrogens are known to be abundantly

released from mature ovary and play important roles in reproductive readiness, sexual behavior, and

so on (Jennings & de Lecea, 2020; Naftolin et al., 2007; Adachi et al., 2007; Clarkson & Herbison,

2009; Wintermantel et al., 2006; Micevych & Meisel, 2017; Melo & Ramsdell, 2001). Among the

ovarian estrogens, 17β-estradiol (E2) is the major factor important for reproduction (Kanda et al.,

2011; Kelly & Qiu, 2010), and the blood E2 concentration of LD-conditioned female medaka is

higher than those of SD (Ikegami et al., 2022). Thus, we hypothesized that E2 regulates the

expression of *agrp1* and *npyb* under the condition of breeding season. We analyzed the expression of

agrp1 and *npyb* in sham-operated (Sham), ovariectomized (OVX, fish with surgical ablation of the

ovary), and OVX medaka with E2-administration (OVX+E) (Figure 5A-5B). The OVX medaka

were allowed to survive at least for two weeks to clear the endogenous E2 (Kayo et al., 2020), and

spawning of all the Sham medaka were confirmed for three consecutive days. By using whole-brain

RT-qPCR, we found that OVX induces significantly lowered *agrp1* expression than Sham, and

OVX+E shows a tendency to recover *agrp1* expression compared with OVX (Figure 5A; Sham vs

OVX: $p = 0.04310$, OVX vs OVX+E: $p = 0.05790$, Sham vs OVX+E: $p = 0.2000$), which suggests

that the ovarian E2 regulates *agrp1* expression. On the other hand, the expression levels of *npyb* did

not show significant differences among the three groups (Figure 5B; Sham vs OVX: $p = 0.1386$,

OVX vs OVX+E: $p = 0.9991$, Sham vs OVX+E: $p = 0.08120$). In addition, food intake of OVX

female was not significantly different between LD and SD (Figure 5-Supplementary Figure 1; $p =$

0.7308), which suggests that ovarian signal may be important for breeding season-dependent feeding

behavior. Therefore, we focused more on the estrogenic regulation of *agrp1* expression.

Estrogens act mainly by interacting with estrogen receptors (Chen et al., 2022). Medaka has three

kinds of estrogen receptors; *esr1*, *esr2a* and *esr2b* (Tohyama et al., 2016; Kinoshita et al., 2009), and

all of them have been reported to be expressed in NVT (Zempo et al., 2013), in which *agrp1* was

also expressed (Figure 2B). Therefore, we examined co-expression of these *esr* genes and *agrp1*. As

shown in Figure 5C and 5D, *esr2a* signal was clearly co-expressed in some *agrp1*-expressing

neurons, which strongly suggests that E2 affects the expression of *agrp1* via *esr2a* in those neurons

of NVT.

$agrp1^{-/-}$ female medaka show a decrease in food intake and in the number of fertilized eggs

Our present experimental evidence thus far highlights the importance of *agrp1* as the factor

modulating the season-dependent feeding behavior in medaka. To analyze the function of AgRP1 in

medaka, we generated knockout medaka of *agrp1* ($agrp1^{-/-}$) by using CRISPR/Cas9. The designed

CRISPR guide RNA cleaved targeted sites of *agrp1* (exon3, Figure 6-Supplementary Figure 1A),

and we obtained $agrp1^{-/-}$ medaka, which has lots of amino acid changes in functional site for AgRP1

(Figure 6-Supplementary Figure 1B). In $agrp1^{-/-}$ brain, AgRP1 immunoreactive signals, which were

observed in WT, were not found (Figure 6-Supplementary Figure 1C). These suggested that $agrp1^{-/-}$

possess nonfunctional AgRP. As for phenotype of the knockout, the $agrp1^{-/-}$ female medaka

appeared skinny and the body weight was significantly lower than that of $agrp1^{+/+}$ (Figure 6A; body

weight: $p = 0.04113$; abdominal length: $p = 0.002165$). In addition, abdominal height of $agrp1^{-/-}$

was also smaller than that of $agrp1^{+/+}$, while the body length was not significantly different (Figure

6A; body length: $p = 0.3939$). We next analyzed food intake of $agrp1^{-/-}$ female medaka in LD

condition (breeding). As shown in Figure 6B ($p = 0.004329$) and Figure 6-Supplementary Figure 2

($agrp1^{+/+}$ vs $agrp1^{-/+}$: $p = 0.5470$, $agrp1^{+/+}$ vs $agrp1^{-/-}$: $p = 0.009353$, $agrp1^{-/+}$ vs $agrp1^{-/-}$: $p = $

0.01234), we found that LD $agrp1^{-/-}$ female medaka eat less than $agrp1^{+/+}$. Then, we kept $agrp1^{-/-}$

medaka in LD or SD condition and compared their food intake. In contrast to $agrp1^{+/+}$, $agrp1^{-/-}$ in

LD condition did not show a significant increase in food intake compared with SD (Figure 6C; $p = $

0.5953). We also examined whether loss of AgRP1 affects reproductive function. Whereas the

$agrp1^{-/-}$ females were fertile, the pairs of $agrp1^{-/-}$ female and $agrp1^{+/+}$ male resulted in fewer

spawned eggs than $agrp1^{+/+}$ pairs (Figure 6D; $p = 0.008658$). In addition, the ovarian size of $agrp1^{-/-}$

appeared to be smaller than $agrp1^{+/+}$ (Figure 6E left). In particular, since relative ovarian weight

normalized by body weight (Gonadosomatic index: GSI) of $agrp1^{-/-}$ female tended to be marginally

smaller than $agrp1^{+/+}$ (Figure 6E right; $p = 0.06494$), the smaller body size of $agrp1^{-/-}$ (Figure 6A)

may drastically affect ovarian morphology. Since the number of spawned eggs was decreased in

$agrp1^{-/-}$ female, we analyzed gene expressions of gonadotropins which should affect ovarian

maturation. Oocyte maturation and ovulation are known to be regulated by gonadotropins,

follicular-stimulating hormone (FSH) and luteinizing hormone (LH). As shown in Figure 6F,

$agrp1^{-/-}$ females showed lower levels of expression of gonadotropin genes (*fshb* and *lhb*; *lhb*: $p = $

0.008658; *fshb*: $p = 0.02597$), which suggests that loss of function of *agrp1* impaired breeding

season-dependent feeding behavior and led to attenuation of reproductive functions, especially the

decrease in number of spawned eggs in the breeding season.

Discussion

In the present study, we took advantage of female medaka which clearly shows breeding season-dependent feeding behavior and found that neuropeptides, *agrp1* and *npyb,* show higher expression under the breeding condition than under the non-breeding condition. We also obtained results to suggest that the expression of both *agrp1* and *npyb* changes depending on nutritional status of female medaka. In addition, ovariectomy and E2 administration changed expression of *agrp1* but not *npyb*, suggesting that increased release of ovarian E2 in breeding season upregulates the *agrp1* expression, which results in facilitation of female feeding behavior. Finally, loss-of-function mutation of AgRP1 decreased the amount of food intake and the number of spawned eggs. The present results suggest that breeding season-dependent feeding behavior can be modulated by the increased expression of *agrp1* upregulated by increased release of ovarian estrogen in the breeding season (Figure 7). To date, not a few previous reports have shown the influence of nutritional status on reproduction (Evans & Anderson, 2012; Kalra & Kalra, 1996; Kirkwood et al., 1987; Merry & Holehan, 1979; Hasebe et al., 2016). On the other hand, although seasonal breeders have been reported to show changes in feeding behavior during the breeding season (Tsuyuki, 2018; Kawai et al., 2020), its neuroendocrine mechanism have largely remained enigmatic. Our present results may provide a neuroendocrinological model for the mechanisms that play a key role in the control of breeding season-dependent feeding behavior in teleosts.

Feeding-related peptides AgRP and NPY in medaka

Here, we demonstrated that female medaka eat more under the condition of breeding season (Figure 1B). Various kinds of neuropeptides in the brain have been suggested to control feeding, and these are generally called "feeding-related peptides" (Funahashi et al., 2003). In the present study, we first used a seasonally breeding model teleost medaka and searched for the "feeding-related peptides" involved in seasonal feeding behavior. A whole-brain RNA-seq analysis using female medaka under the breeding condition (LD) and non-breeding condition (SD) revealed two kinds of feeding-related peptides, *agrp1* and *npyb*, which show different expression levels between LD and SD (Figure 1-Supprementary Figure 2). In mammals, AgRP and NPY are known to have orexigenic function and are co-expressed in hypothalamic neurons (Schwartz et al., 2000; Hahn et al., 1998). Previous studies in mammals (Schwartz et al., 2000; Andermann & Lowell, 2017; Muroi & Ishii, 2016) have suggested neural mechanisms of appetite including functions of AgRP and NPY. However, such mechanisms in non-mammalian vertebrates such as teleosts (Rønnestad et al., 2017; Blanco & Soengas, 2021) have not yet been clarified. Our present study using medaka has shown possible functions of AgRP and NPY in teleost feeding behavior, especially in a breeding season-dependent

manner.

Our present study using medaka showed that female medaka express *agrp1* in hypothalamus, and

food restriction increases the *agrp1* expression (Figure 2B and 2G). It has been reported that leptin

receptor-knockout medaka show higher food intake and higher expressions of *agrp1* and *npya* than

wild type, whereas the expression of *agrp2* and *npyb* remained to be analyzed (Chisada et al., 2014).

Zebrafish has also been used as a model animal in teleosts. In zebrafish, food restriction increased

agrp1 (Song et al., 2003; Opazo et al., 2018) expression, and transgenic overexpression of *agrp1*

caused gain of body weight (Song & Cone, 2007), as in mammals (Graham et al., 1997; Adam et al.,

2002; Hahn et al., 1998; Ilnytska & Argyropoulos, 2008). It has also been reported that *agrp1*

knockout zebrafish eat less than the wild type (Shainer et al., 2019), although loss of AgRP in mice

showed little effect on food intake (Qian et al., 2002). Our present results and these previous studies

strongly support that *agrp1* regulates feeding and may act as an orexigenic factor in teleosts. On the

other hand, *agrp2* neurons were distributed mainly in telencephalon, and its weak expressions were

also observed in the POA and the hypothalamus (Figure 2C), which is different from the results in

zebrafish (Shainer et al., 2017). In the present study, food restriction did not remarkably affect the

agrp2 expression in medaka (Figure 2 and Figure 2-Supplementary Figure 1E), and AgRP2 in

zebrafish is suggested to play an important role in stress response, not feeding (Shainer et al., 2019).

Thus, it is highly probable that *agrp2* is involved in functions other than feeding in female medaka.

Furthermore, we demonstrated that *npya* is expressed in multiple brain regions including

hypothalamus in medaka (Figure 2D), which is similar to mammals (Gray & Morley, 1986) and

zebrafish (Yokobori et al., 2012; Jeong et al., 2018). We also showed that *npya* is not co-expressed

with *agrp1* nor *agrp2* (Figure 2F) as in the zebrafish (Jeong et al., 2018), suggesting that The

relationship between NPY and AgRP of teleosts may be different from that of mammals, in which

most of the *agrp*-expressing neurons co-express *npy* (Hahn et al., 1998). Moreover, our present study

also showed that *npyb* expression is localized in telencephalon (Figure 2E), which is similar to the

previous report using tiger puffer (Kamijo et al., 2011). Previous studies of NPY in zebrafish showed

that zebrafish has only one type of NPY (NPYa) (Söderberg et al., 2000; Larsson et al., 2009) and

has lost NPYb during evolution. Like in mammals (Marks et al., 1992; Clark et al., 1984; Glenn

Stanley et al., 1986; Marks & Waite, 1997; Baldock et al., 2009), food restriction in zebrafish

increased the *npya* expression in the hypothalamus (Song et al., 2003; Opazo et al., 2018), and

intracerebroventricular administration of NPYa increased food intake (Yokobori et al., 2012).

Although these zebrafish studies suggest that NPYa may increase food intake, it is still debatable,

since body weight was not significantly different between *npya* knockout zebrafish and wild type

(Shiozaki et al., 2020). Interestingly, by analyzing both *npya* and *npyb* expression in medaka of

different nutritional conditions, we found that food restriction decreased the *npya*-expressing cell
number in the hypothalamus (Figure 2-Supplementary Figure 1) and *npyb* expression level (Figure
2J). These changes in *npya* and *npyb* expressions are not consistent with previous studies using other
conventional model animals described above (Yokobori et al., 2012; Marks et al., 1992; Clark et al.,
1984; Glenn Stanley et al., 1986; Marks & Waite, 1997; Baldock et al., 2009). Our present study
may suggest that the function of *npy* may be different among teleosts. In addition, *npyb* expression
was increased under the breeding condition (LD), while LD female showed increase in food intake
(Figure 3F). Thus, decreased expression of *npyb* by food restriction (Figure 2J) may suggest that the
change in *npyb* expression reflects nutritional condition in medaka. Thus, future study of *npya* and
npyb functions in the control of feeding will be necessary.
337
As described above, we found that *agrp1*, *npya*, and *npyb* change expression levels in response to
nutritional status. Among these three genes, we suggest that *agrp1* most probably affects relatively
long-term feeding in the breeding season, which agrees well with the recent studies in mice showing
the function of AgRP as a long-term orexigenic factor. In mice, it has been reported that
intracerebroventricular administration of AgRP increases food intake for one week (Hagan et al.,
2000), and stimulation of receptors expressed in AgRP neurons triggers AgRP release, leading to an
increase in food intake for three days (Nakajima et al., 2016). Our present study also suggests a
long-term (seasonal) orexigenic effect of AgRP in teleosts and may also provide an important insight
into the understanding of common regulatory mechanisms of feeding by AgRP among various
animal species.
348
349 **High concentration of E2 in breeding season facilitates *agrp1* expression**
350
Our results suggest that *agrp1* and *npyb* show higher expressions under the breeding condition (LD)
(Figure 3), but the experiments using juvenile female medaka (Figure 4) showed that expression
levels of these two genes do not change according to the day-length itself but to the LD-induced
sexual maturity. In addition, the present results indicate that the ovarian estrogen E2 upregulates
agrp1 expression mainly via the estrogen receptors *esr2a* that are co-expressed in some population
of *agrp1* neurons in the hypothalamic nucleus NVT (Figure 5). Since LD female medaka (breeding)
shows high blood concentration of E2 (Ikegami et al., 2022), this pathway may be important for
breeding season-dependent feeding behavior. Especially, in teleost, main egg protein for nutrition is
vitellogenin, whose expression is also facilitated by E2 (Tohyama et al., 2017). Taken together, it is
suggested that E2 may synchronously regulate amount of food intake and female-specific
reproductive signals (vitellogenin production and oocyte maturation), which plays a key role in
reproductive success in oviparous animals.

In mammals, previous studies have reported on inconsistent effects of ovary and E2 on feeding. Ablation of ovary caused suppression of food intake in mice (Yu et al., 2020), whereas it caused no change in rats (Roesch, 2006). On the other hand, administration of E2 decreased food intake in both mice (Yu et al., 2020) and rats (Roesch, 2006). It should be noted that these laboratory rodents only exhibit short estrous cyclicity and have lost breeding seasonality, and the blood E2 concentration drastically changes in a few days (Nilsson et al., 2015). Thus, it is possible that the control mechanisms of feeding may be different between animals with short estrous cyclicity and those with breeding seasonality.

Furthermore, our present study suggests different control mechanisms of feeding between the animals with breeding seasonally and those without. Here, we showed that E2 directly modulates *agrp1* expression via *esr2a* receptors co-expressed in the *agrp1* neurons (Figure 5C), while in mice, *AgRP/NPY* neurons are reported to be suppressed by E2 indirectly via *esr1* (*erα*)-expressing *Kiss1* neurons located in the hypothalamic arcuate nucleus (Qiu et al., 2018; Dubois et al., 2016; Yang et al., 2017). On the other hand, in medaka, expressions of all kinds of estrogen receptors are reported to be localized in NVT (Zempo et al., 2013), in which *agrp1* expression is also localized (Figure 2B). In addition, *esr2a* has been reported to be involved in the feedback regulation of follicle stimulating hormone in the pituitary and in the development of oviduct, and *esr2a* knockout females are completely infertile (Kayo et al., 2019). Our hypothesis that estrogen signaling via *esr2a* affects *agrp1* expression may highlight another important function of *esr2a* for reproduction, while a possibility still remains that *esr1-* and *esr2b*-expressing neurons also affect *agrp1* expression indirectly.

AgRP1 changes feeding behavior depending on LD-induced sexual maturity, which causes the increase in food intake in breeding season

Since our results thus far indicate the importance of *agrp1*, which shows upregulated expression directly stimulated by the ovarian E2 in breeding season, we examined phenotypes of *agrp1* knockout (*agrp1$^{-/-}$*) medaka. We found that *agrp1$^{-/-}$* medaka under the condition of breeding season eat less (Figure 6B) and spawn a smaller number of eggs (Figure 6D) than WT. Furthermore, the *agrp1$^{-/-}$* females did not show significant difference of food intake in LD and SD (Figure 6C). These results strengthen our hypothesis that *agrp1* is involved in the increased food intake in breeding season. Furthermore, *agrp1$^{-/-}$* female displayed light body weight (Figure 6A), accompanied by smaller ovary (Figure 6E) and low level of expression of gonadotropins, *fshb* and *lhb* (Figure 6F), which are considered to have caused smaller number of spawned eggs. All of these results support

our hypothesis that AgRP1 plays an important role in breeding season-dependent feeding behavior,

which culminates in normal reproduction.

In summary, by using a seasonal breeder medaka, we found evidence to suggest that long day-length

facilitates ovarian maturation and E2 release, which upregulates *agrp1* expression of hypothalamic

neurons to activate neural circuitry for feeding behavior and boost oocyte maturation. We propose

that this kind of positive feedback control may be important for animals that spawn many eggs every

day in the breeding season (Figure 7); medaka needs plenty of food for production of many eggs. In

other words, the metabolic costs of producing eggs on a daily basis in medaka would inevitably

require increased food intake. Indeed, previous study showed a need for high food intake for

reproduction (Hasebe et al., 2016). It is expected that future studies will elucidate whether or not the

present findings in medaka are applicable to other seasonal breeders as well.

Material and Methods

Animals

Female and male wild-type d-rR medaka (*Oryzias latipes*) and *agrp1* knockout (*agrp1⁻/⁻*) medaka were maintained in pairs or shoals at 27□. Fish were fed three times a day with brine shrimp and flake food (Otohime B-2; San-u Fish Farm, Osaka, Japan). Their reproductive status was controlled by day-length (long day (LD; 14-hour light/10-hour dark): reproductive, short day (SD; 10-hour light/14-hour dark): non-reproductive). The light-on time was 8:00 AM. We used juvenile medaka (~5 weeks after fertilization) and adult medaka (>3 months after fertilization). Female medaka, which spawned at least three consecutive days, were used as reproductive ones. For the analysis of the effect of food restriction, reproductive and non-reproductive female medaka were fasted for 14 days or 10 days. Note that all medaka survived after food restriction. For the analyses of food intake, we food-restricted medaka for 6 hours after 10-min feeding in the morning and sampled their whole brains for subsequent experiments. Food-restricted medaka were sampled at the same time as the other fed medaka. All experiments and fish maintenance were conducted in accordance with the Guidelines for Proper Conduct of Animal Experiments (Science Council of Japan) and the protocols approved by the Animal Care and Use Committee of Graduate School of Science, the University of Tokyo (Permission number, 17-1, 20-6) and the Animal Care and Use Committee of Graduate School of Agriculture, Tokyo University of Agriculture and Technology (Permission number, R05-15, R06-27).

Food intake assay

Each 6-hour food-restricted medaka was put into a white cup with 100 mL breeding water and was habituated for 5 min. Then, we fed medaka by application of 200 μL aliquots of food water containing brine shrimp in all-you-can-eat style and serve another aliquot once done with it, which is repeated N times (like the Japanese "Wanko soba"; so-called Japanese "Wanko soba" method). 10 min after the start, we stopped feeding medaka, placed a magnetic bar to stir the breeding water so that the shrimp concentration will be constant. Then we collected 10 mL aliquot from the experimental cup by using a micro pipette and transferred it to a conical tube. The conical tube was frozen overnight, and the leftover brine shrimp sunk in the bottom were counted by "shrimp-counter". We counted the number of brine shrimp in the 200 μL solution three times before and after the experiments, and the average number was used. The food intake was calculated as follows.

(Food Intake)

= (The average number of brine shrimp in the solution) * (number of aliquots, N)

- (number of leftover brine shrimp sunk in the bottom) * 10

Food intake was normalized by the average of LD or WT medaka.

"Shrimp-counter" system

The number of shrimps in the solution was counted using OpenCV3 library (Intel, Santa Clara, CA)

run under a python script (Figure 1-Source code 1). This script was run under Anaconda 4.4.0 for

windows running python 3.5.

RNA-sequencing

We collected two whole brains of LD or SD female in one tube (Note that pituitary was confirmed

not to be included), and extracted total RNA by using NucleoSpin® RNA Plus kit

(MACHEREY-NAGEL, Düren, Germany). cDNA was obtained by KAPA Stranded mRNA-Seq Kit

(Nippon Genetics Co., Ltd, Tokyo, Japan) and KAPA Library preparation kit (Nippon Genetics Co.,

Ltd). Then, it was applied to a next generation sequencer Hiseq 2500 (Illumina, San Diego, CA),

following the standard protocol of Illumina system. We selected the candidate genes judging from

transcripts per million (TPM) for expression value in the obtained data using CLC workbench. We

made volcano plot using R (R core team 2023) and Rstudio (2023) and colored dots, which indicate

P-value < 0.05 and |log FC| > 1. In addition, we made a heatmap of genes related to neuroendocrine

system using DESeq2 (Love et al., 2014).

Histological analysis of the distribution of *agrp*- and *npy*-neurons in the brain

To analyze the distribution of *agrp*- and *npy*-expressing neurons, we performed *in situ* hybridization

(ISH) for *agrp* and *npy* on frozen sections of reproductive medaka. In brief, female medaka was

anesthetized (FA100, Bussan Animal Health Co., Ltd., Osaka, Japan), and its brain was picked up

and fixed with 4% paraformaldehyde (PFA)/PBS. In analyses on *agrp2* expression, we performed

perfusion-fixation by using 4% PFA/PBS. After incubation with 30% sucrose/PBS, brains were

embedded in 5% low melting agar/ 20% sucrose/ PBS and sectioned at a thickness of 25 µm. The

sections were hybridized with *agrp1* (ENSORLG00000000398, 177 bases), *agrp2*

(ENSORLG00000029106, 303 bases), *npya* (ENSORLG00000004649, 288 bases) and *npyb*

(ENSORLG00000007880, 288 bases) specific digoxigenin (DIG)-labeled RNA probes and

performed nitro blue tetrazolium (NBT)/ 5-bromo-4-chloro-3-indolyl-phosphate color-reaction

(BCIP) after wash and incubation with anti-digoxigenin antibody (catalog. no. 11093274910; Roche;

RRID: AB_514497) as previously reported (Zempo et al., 2013). Photographs were taken with a

digital camera (DFC310FX; Leica Microsystems, Wetzlar, Germany) attached to an upright

microscope (DM5000B; Leica Microsystems).

Histological analysis of *agrp1*-, *npy*- and estrogen receptor (*esr*)-expressing neurons

To examine whether *agrp1*-expressing neurons co-express *npya* and *esr*, we prepared *agrp1*

fluorescein-labeled RNA probe and carried out double ISH as previously reported (Umatani et al.,

2022). *esr* DIG-labeled probes were kindly given by Dr. Kayo (Kyoto Univ.), and we used *npya*

DIG-labeled probe described in the previous paragraph. In brief, we made brain sections as

described above and applied both *agrp1* fluorescein-labeled and each DIG-labeled RNA probes.

Signals for *npya*, *esr1*, *esr2a* and *esr2b* were visualized by incubation with anti-digoxigenin

antibody (catalog. no. 11207733910; Roche; RRID: AB_514500) and TSA Plus Cy3 System

(TSA-Plus Cyanine 3 system, catalog. no. NEL744001KT, Akoya Biosciences, Marlborough, MA).

After inactivation of Cy3 system by 3% H_2O_2, we applied peroxidase-conjugated anti-fluorescein

antibody (catalog. no. 11426346910, Roche; RRID: AB_840257) on sections and performed TSA

Plus biotin system (catalog. no. NEL749A001KT, Akoya Biosciences). Then, signals for *agrp1* were

visualized by Alexa 488 conjugated streptavidin (catalog. no. S11223, Invitrogen). For

counter-staining of cell nuclei, DAPI in PBS was applied on section. On the other hand, to examine

whether *agrp2*-expressing neurons co-express *npya*, we used *npya* fluorescein-labeled RNA probe

and *agrp2* DIG- labeled one. Double ISH of them was performed according to the same method

described above. Fluorescent images were acquired with a confocal laser-scanning microscope

(AXR, Nikon, Tokyo, Japan) using excitation and emission wavelengths of 405 nm and 429-474 nm

for DAPI, 488 nm and 512-526 nm for Alexa 488, and 561 nm and 571-625 nm for Cy3, respectively.

These were photographed at Tokyo University of Agriculture and Technology for Smart Core facility

Promotion Organization.

Quantitative Real-time Polymerase Chain Reaction (RT-qPCR)

A Whole brain or a pituitary was collected from each medaka and total RNA was extracted by using

FastGene^{TM} RNA basic kit (Nippon Genetics Co., Ltd) according to the manufacturer's instructions.

For the juvenile medaka, we checked their sex as previously reported, and two samples of the same

sex were mixed and used as one sample. Total RNA samples were reverse transcribed by

FastGene^{TM} cDNA synthesis 5× ReadyMix OdT according to the manufacturer's instructions. For

the analyses of the brain, 1μL of cDNA diluted with 10-fold MQ was mixed with KAPA SYBR Fast

qPCR kit and amplified with Lightcycler96 [Roche; 95□ 150s (95□ 10 sec, 60□ 10 sec, 72□ 15

sec) × 45 cycles]. For the analysis of the pituitary, 1 μL of cDNA diluted with 5-fold MQ was mixed

with KAPA SYBR Fast qPCR kit and amplified with Lightcycler96 [Roche; 95□ 150s (95□ 10 sec,

60□ 10 sec, 72□ 10 sec) × 45 cycles]. The data was normalized by housekeeping gene, ribosomal

protein s13 (*rps13*). Primer sequences were as follows:

AgRP1 RT-PCR F1 CCAATTTCCAGTCACCGAAG

AgRP1 RT-PCR R1 CTGGGTCCAACACAGAATCA

AgRP2 RT-PCR F1 TTGTTGTGCTTCTTGCTGCT

AgRP2 RT-PCR R1 ACAGAGCTCCAAACGGTGTC

NPYa SE CTCATCACAAGACAGAGGTATGGG

NPYa AS GGGTTGTAACTTGACTGTGGAAGTG

NPYb SE CTGCCTGCTCCTCTGTTTTTTCTC

NPYb AS CACAGTGTCTGGGTTGTCTCTCTTTC

qPCR FSHb Fw new TGGAGATCTACAGGCGTCGGTAC

qPCR FSHb Fw new AGCTCTCCACAGGGATGCTG

qPCR LHb Fw new AGGGTATGTGACTGACGGATCCAC

qPCR LHb Rv new TGCCTTACCAAGGACCCCTTGATG

RPS13 SE GTGTTCCCACTTGGCTCAAGC

RPS13 AS CACCAATTTGAGAGGGAGTGAGAC

Sham operations, ovariectomy, and E2 administrations

Ovariectomy and E2 administration were performed according to a previous study (Kayo *et al.*,

2020). Briefly, reproductive female medaka were anesthetized with 0.02% MS-222 (Sigma-Aldrich,

St. Louis, MO) and their ovaries were excised via intraperitoneal operation. Sham operation group

was anesthetized, received an abdominal incision without removing the ovaries, and received skin

suture by using a silk thread. After checking that all Sham females spawn, we prepared three tanks;

two tanks contained 7-8 OVX medaka, and one tank contained Sham medaka in 2 L breeding water

in it. We dissolved β-Estradiol 1.4 mg in 1 mL EtOH (E2 stock) and dispensed 2 μL of E2 stock or

the same amount of 100% Ethanol for the control tank. Ethanol or E2 containing water were

changed every day. After the steroid treatment for 5 days, the medaka were anesthetized, and their

whole brains were collected for RT-qPCR analysis.

Generation of *agrp1* KO medaka lines

We generated *agrp1* KO medaka lines by using CRISPR/Cas9. Cas9 mRNA and tracer RNA were

purchased from Integrated DNA Technologies (IDT, Coralville, IA). The guide RNA sequence for

digestion by CRISPR/Cas9 complex was "CCTCACCAGCAGTCCTGCCTGG".

Mixture of Cas9 protein, tracer RNA, CRISPR RNA, GFP mRNA diluted with PBS and 0.02%

phenol red (final concentration: Cas9 protein; 500 ng/μL, tracer RNA; 100 ng/μL, CRISPR RNA;

50 ng/μL, GFP mRNA; 5 ng/μL) was injected into the cytoplasm of one- or two-cell-stage

embryos (F0). To obtain homozygous transgenic offspring, the carriers were crossed with each other.

Measurement of body size and ovary size

We took photographs of fish bodies from the lateral side by using a digital camera MC120HD

(Leica) and calculated the abdominal and body length by ImageJ. For the analysis of body length, we

measured the length from the mouth to the base of the tail. Gonadosomatic index (GSI) was
calculated as ovary weight/ body weight*100.

Immunohistochemistry using AgRP1 antibody

We made brain sections of WT and *agrp1* KO as described in "*Histological analysis of the*
distribution of agrp- and npy-neurons in the brain". After washing with PBST two times for 10 mins,
sections were incubated with AgRP1 antibody (1:1000; Rabbit polyclonal AgRP (83-132) amide
(Human), Cat# H-003-53; RRID: AB_2313908, Phoenix Pharmaceutical, Burlingame, CA) /10%
goat serum/PBS overnight. In the next day, slides were washed with PBST and incubated with
anti-rabbit biotinylated goat antibody (1:200; Cat. #BA-1000; Vector Laboratories, Burlingame, CA)
for 1 hour. Then we applied Alexa 488 conjugated streptavidin (1:500, Cat. #S11223, Invitrogen) and
DAPI (1:2000; Dojin, Kumamoto, Japan). Photographs were taken with a digital camera (Leica
Microsystems) attached to an upright microscope (Leica Microsystems).

Statistics

For the statistical analysis, we used Kyplot 5.0 software (Kyence, Tokyo, Japan) or R software (R
core team 2023) with R studio (version 2023.06.0+421). For the comparison of the TPM, we used
Student's t-test. For the comparison of *agrp1* and *npyb* expressions in OVX and E2 administrated
medaka, Steel-Dwass test was used for multiple comparison of the expression level. In the other
experiments, we used Mann-Whitney U test. In all statistical analysis, significance levels were
described as follows: *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$.

Data availability

All data supporting the findings of our present study and its supplementary information are available
in the article or from the corresponding author upon reasonable request.

Code availability

The code of "Shrimp-counter" system is available in the present study, Figure 1-Source code 1 (see
Methods).

Acknowledgements

We thank Drs. Mikoto Nakajo (Osaka Med and Pharm Univ), Soma Tomihara (Nagahama Inst of
Bio-Sci and Tech), and Kana Ikegami (Kitasato Univ) for helpful discussion. We also thank Dr.
Daichi Kayo (Kyoto Univ.) for his kind supply of *esr* DIG-labeled probes. In addition, we deeply
appreciate Dr. Hiroyuki Takeda (U Tokyo) for his continued kind support and encouragement during

our experiments. We also thank Dr. Yutaka Miura (Tokyo Univ of Agri and Tech) for his kind support of our experiments. As for the help of animal care, we thank Ms. Hisako Kohno, Miho Kyokuwa, Hiroko Tsukamoto, and Risa Nakaba. We thank Ms. Maiko Matsuda (Vesper studio) for her kind gift of medaka illustrations to us. The present study was supported by JSPS KAKENHI (JP21J20864 and JP22KJ0597 to YT, JP26221104 and JP21K06262 to YO, JP20H03071 to CU), and MEXT Initiative for Realizing Diversity in the Research Environment (Leadership training for women) (CU).

Author contributions

Conceptualization: Y.T. and C.U. ; Investigation: Y.T., S.T., H.W., S.Ki., S.Ka, and C.U. ; Methodology: Y.T., H.W., T.K., S.Ka., and C.U. ; Resources: Y.O., Y.H., and C.U.; Writing—original draft: Y.T. and C.U.; Writing—review and editing: Y.T., S.Ka., Y.O., Y.H., and C.U.; Project administration: C.U.; Supervision: S.Ka., Y.O., Y.H., and C.U.; and Funding acquisition: Y.T., S.Ka., Y.O., Y.H., and C.U.

Competing interests

The authors declare no competing interests.

References

Adachi, S., Yamada, S., Takatsu, Y., Matsui, H., Kinoshita, M., Takase, K., Sugiura, H., Ohtaki, T.,
Matsumoto, H., Uenoyama, Y., Tsukamura, H., Inoue, K., & Maeda, K.-I. (2007).
Involvement of Anteroventral Periventricular Metastin/Kisspeptin Neurons in Estrogen
Positive Feedback Action on Luteinizing Hormone Release in Female Rats. *Journal of*
Reproduction and Development, 53(2), 367-378.

Adam, C. L., Archer, Z. A., Findlay, P. A., Thomas, L., & Marie, M. (2002). Hypothalamic gene
expression in sheep for cocaine- and amphetamine-regulated transcript,
pro-opiomelanocortin, neuropeptide Y, agouti-related peptide and leptin receptor and
responses to negative energy balance. *Neuroendocrinology, 75*(4), 250-256.

Amirjani, S., Asemi, Z., Bazarganipour, F., Aramesh, S., Allan, H., Sayadi, M., Tabatabaei, M. S.,
Mohamadian, Z., Zabti, F., Iranpak, N., Heydarzadeh, A., Taghavi, S. A., Badehnoosh, B., &
Khashavi, Z. (2019). Dietary intake and lifestyle behaviour in different phenotypes of
polycystic ovarian syndrome: a case-control study. *J Hum Nutr Diet, 32*(4), 413-421.

Andermann, M. L., & Lowell, B. B. (2017). Toward a Wiring Diagram Understanding of Appetite
Control. *Neuron, 95*(4), 757-778.

Aponte, Y., Atasoy, D., & Sternson, S. M. (2011). AGRP neurons are sufficient to orchestrate feeding
behavior rapidly and without training. *Nature Neuroscience, 14*(3), 351-355.

Baldock, P. A., Lee, N. J., Driessler, F., Lin, S., Allison, S., Stehrer, B., Lin, E. J., Zhang, L., Enriquez, R.
F., Wong, I. P., McDonald, M. M., During, M., Pierroz, D. D., Slack, K., Shi, Y. C.,
Yulyaningsih, E., Aljanova, A., Little, D. G., Ferrari, S. L., Sainsbury, A., Eisman, J. A., &
Herzog, H. (2009). Neuropeptide Y knockout mice reveal a central role of NPY in the
coordination of bone mass to body weight. *PLoS One, 4*(12), e8415.

Blanco, A. M., & Soengas, J. L. (2021). Leptin signalling in teleost fish with emphasis in food intake
regulation. *Mol Cell Endocrinol, 526*, 111209.

Broberger, C., De Lecea, L., Sutcliffe, J. G., & Hökfelt, T. (1998). Hypocretin/orexin- and
melanin-concentrating hormone-expressing cells form distinct populations in the rodent
lateral hypothalamus: relationship to the neuropeptide Y and agouti gene-related protein
systems. *J Comp Neurol, 402*(4), 460-474.

Chen, P., Li, B., & Ou-Yang, L. (2022). Role of estrogen receptors in health and disease. *Front*
Endocrinol (Lausanne), 13, 839005.

Chen, S. E., McMurtry, J. P., & Walzem, R. L. (2006). Overfeeding-Induced Ovarian Dysfunction in
Broiler Breeder Hens Is Associated with Lipotoxicity12. *Poultry Science, 85*(1), 70-81.

Chisada, S., Kurokawa, T., Murashita, K., Rønnestad, I., Taniguchi, Y., Toyoda, A., Sakaki, Y., Takeda,
S., & Yoshiura, Y. (2014). Leptin receptor-deficient (knockout) medaka, Oryzias latipes,
show chronical up-regulated levels of orexigenic neuropeptides, elevated food intake and
stage specific effects on growth and fat allocation. *Gen Comp Endocrinol, 195*, 9-20.

Clark, J. T., Kalra, P. S., Crowley, W. R., & Kalra, S. P. (1984). Neuropeptide Y and human pancreatic
polypeptide stimulate feeding behavior in rats. *Endocrinology, 115*(1), 427-429.

Clarkson, J., & Herbison, A. E. (2009). Oestrogen, kisspeptin, GPR54 and the pre-ovulatory
luteinising hormone surge. *J Neuroendocrinol, 21*(4), 305-311.

Conde-Sieira, M., & Soengas, J. L. (2016). Nutrient Sensing Systems in Fish: Impact on Food Intake
Regulation and Energy Homeostasis. *Frontiers in Neuroscience, 10*, 603.

Dubois, S. L., Wolfe, A., Radovick, S., Boehm, U., & Levine, J. E. (2016). Estradiol Restrains
Prepubertal Gonadotropin Secretion in Female Mice via Activation of ERα in Kisspeptin
Neurons. *Endocrinology, 157*(4), 1546-1554.

Egami, N. (1954). Effect of artificial photoperiodicity on time of oviposition in the fish, Oryzias
latipes. *Annot Zool Japonenses, 27*(2), 57-62.

Evans, J. J., & Anderson, G. M. (2012). Balancing ovulation and anovulation: integration of the
reproductive and energy balance axes by neuropeptides. *Human Reproduction Update, 18*(3),
313-332.

Funahashi, H., Takenoya, F., Guan, J. L., Kageyama, H., Yada, T., & Shioda, S. (2003). Hypothalamic
neuronal networks and feeding-related peptides involved in the regulation of feeding. *Anat*
Sci Int, 78(3), 123-138.

Glenn Stanley, B., Kyrkouli, S. E., Lampert, S., & Leibowitz, S. F. (1986). Neuropeptide Y chronically
injected into the hypothalamus: A powerful neurochemical inducer of hyperphagia and
obesity. *Peptides, 7*(6), 1189-1192.
686

687 Graham, M., Shutter, J. R., Sarmiento, U., Sarosi, I., & Stark, K. L. (1997). Overexpression of Agrt
688   leads to obesity in transgenic mice. *Nat Genet, 17*(3), 273-274.
689

690 Gray, T. S., & Morley, J. E. (1986). Neuropeptide Y: Anatomical distribution and possible function in
691   mammalian nervous system. *Life Sciences, 38*(5), 389-401.
692

693 Hagan, M. M., Rushing, P. A., Pritchard, L. M., Schwartz, M. W., Strack, A. M., Van Der Ploeg, L. H.,
694   Woods, S. C., & Seeley, R. J. (2000). Long-term orexigenic effects of AgRP-(83---132) involve
695   mechanisms other than melanocortin receptor blockade. *Am J Physiol Regul Integr Comp*
696   *Physiol, 279*(1), R47-52.
697

698 Hahn, T. M., Breininger, J. F., Baskin, D. G., & Schwartz, M. W. (1998). Coexpression of Agrp and
699   NPY in fasting-activated hypothalamic neurons. *Nat Neurosci, 1*(4), 271-272.
700

701 Hasebe, M., Kanda, S., & Oka, Y. (2016). Female-Specific Glucose Sensitivity of GnRH1 Neurons
702   Leads to Sexually Dimorphic Inhibition of Reproduction in Medaka. *Endocrinology, 157*(11),
703   4318-4329.
704

705 Ikegami, K., Kajihara, S., Umatani, C., Nakajo, M., Kanda, S., & Oka, Y. (2022). Estrogen upregulates
706   the firing activity of hypothalamic gonadotropin-releasing hormone (GnRH1) neurons in the
707   evening in female medaka. *Journal of Neuroendocrinology, 34*(4), e13101.
708

709 Ilnytska, O., & Argyropoulos, G. (2008). The Role of the Agouti-Related Protein in Energy Balance
710   Regulation. *Cellular and Molecular Life Sciences, 65*(17), 2721.
711

712 Jennings, K. J., & de Lecea, L. (2020). Neural and Hormonal Control of Sexual Behavior.
713   *Endocrinology, 161*(10).
714

715 Jeong, I., Kim, E., Kim, S., Kim, H.-K., Lee, D.-W., Seong, J. Y., & Park, H.-C. (2018). mRNA
716   expression and metabolic regulation of npy and agrp1/2 in the zebrafish brain. *Neuroscience*
717   *Letters, 668*, 73-79.
718

719   Kalra, S. P., & Kalra, P. S. (1996). Nutritional Infertility: The Role of the Interconnected
720   Hypothalamic Neuropeptide Y–Galanin–Opioid Network. *Frontiers in Neuroendocrinology,*
721   *17*(4), 371-401.

723   Kamijo, M., Kojima, K., Maruyama, K., Konno, N., Motohashi, E., Ikegami, T., Uchiyama, M., Shioda,
724   S., Ando, H., & Matsuda, K. (2011). Neuropeptide Y in tiger puffer (Takifugu rubripes):
725   distribution, cloning, characterization, and mRNA expression responses to prandial
726   condition. *Zoolog Sci, 28*(12), 882-890.

728   Kanda, S., Akazome, Y., Matsunaga, T., Yamamoto, N., Yamada, S., Tsukamura, H., Maeda, K.-i., &
729   Oka, Y. (2008). Identification of KiSS-1 Product Kisspeptin and Steroid-Sensitive Sexually
730   Dimorphic Kisspeptin Neurons in Medaka (Oryzias latipes). *Endocrinology, 149*(5),
731   2467-2476.

733   Kanda, S., Okubo, K., & Oka, Y. (2011). Differential regulation of the luteinizing hormone genes in
734   teleosts and tetrapods due to their distinct genomic environments--insights into
735   gonadotropin beta subunit evolution. *Gen Comp Endocrinol, 173*(2), 253-258.

737   Kauffman, A. S., & Rissman, E. F. (2004). A critical role for the evolutionarily conserved
738   gonadotropin-releasing hormone II: Mediation of energy status and female sexual behavior.
739   *Endocrinology, 145*(8), 3639-3646.

741   Kawai, K., Fujita, H., Sanchez, G., Furusawa, S., & Umino, T. (2020). Estimating the spawning season
742   of black sea bream Acanthopagrus schlegelii in Hiroshima Bay, Japan, from temporal
743   variation in egg density. *Fisheries Science, 86*(4), 645-653.

745   Kayo, D., Oka, Y., & Kanda, S. (2020). Examination of methods for manipulating serum 17β-Estradiol
746   (E2) levels by analysis of blood E2 concentration in medaka (Oryzias latipes). *Gen Comp*
747   *Endocrinol, 285*, 113272.

749   Kayo, D., Zempo, B., Tomihara, S., Oka, Y., & Kanda, S. (2019). Gene knockout analysis reveals
750   essentiality of estrogen receptor β1 (Esr2a) for female reproduction in medaka. *Sci Rep, 9*(1),
751   8868.

753   Kelly, M. J., & Qiu, J. (2010). Estrogen signaling in hypothalamic circuits controlling reproduction.
754   *Brain Res, 1364*, 44-52.

Kinoshita, M., Murata, K., Naruse, K., & Tanaka, M. (2009). *Medaka: biology, management, and*
experimental protocols. John Wiley & Sons

Kirkwood, R., Cumming, D., & Aherne, F. (1987). Nutrition and puberty in the female. *Proceedings*
of the Nutrition Society, 46(2), 177-192.

Kohno, D., & Yada, T. (2012). Arcuate NPY neurons sense and integrate peripheral metabolic signals
to control feeding. *Neuropeptides, 46*(6), 315-319.

Krashes, M. J., Koda, S., Ye, C., Rogan, S. C., Adams, A. C., Cusher, D. S., Maratos-Flier, E., Roth, B.
L., & Lowell, B. B. (2011). Rapid, reversible activation of AgRP neurons drives feeding
behavior in mice. *Journal of Clinical Investigation, 121*(4), 1424-1428.

Larsson, T. A., Tay, B. H., Sundström, G., Fredriksson, R., Brenner, S., Larhammar, D., & Venkatesh,
B. (2009). Neuropeptide Y-family peptides and receptors in the elephant shark,
Callorhinchus milii confirm gene duplications before the gnathostome radiation. *Genomics,*
93(3), 254-260.

Liu, R., Kinoshita, M., Adolfi, M. C., & Schartl, M. (2019). Analysis of the Role of the Mc4r System in
Development, Growth, and Puberty of Medaka. *Frontiers in Endocrinology, 10*, 213.

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology, 15*(12), 550.

Lynn, S. E., Stamplis, T. B., Barrington, W. T., Weida, N., & Hudak, C. A. (2010). Food, stress, and
reproduction: short-term fasting alters endocrine physiology and reproductive behavior in
the zebra finch. *Horm Behav, 58*(2), 214-222.

Marks, J. L., Li, M., Schwartz, M., Porte, D., & Baskin, D. G. (1992). Effect of fasting on regional levels
of neuropeptide Y mRNA and insulin receptors in the rat hypothalamus: An
autoradiographic study. *Molecular and Cellular Neuroscience, 3*(3), 199-205.

Marks, J. L., & Waite, K. (1997). Intracerebroventricular neuropeptide Y acutely influences glucose
metabolism and insulin sensitivity in the rat. *J Neuroendocrinol, 9*(2), 99-103.

Melo, A. C., & Ramsdell, J. S. (2001). Sexual dimorphism of brain aromatase activity in medaka:
induction of a female phenotype by estradiol. *Environmental Health Perspectives, 109*(3),
257-264.
794
795   Merry, B., & Holehan, A. M. (1979). Onset of puberty and duration of fertility in rats fed a restricted
796   diet. *Reproduction, 57*(2), 253-259.
797
798   Micevych, P. E., & Meisel, R. L. (2017). Integrating Neural Circuits Controlling Female Sexual
799   Behavior. *Front Syst Neurosci, 11*, 42.
800
801   Mitani, Y., Kanda, S., Akazome, Y., Zempo, B., & Oka, Y. (2010). Hypothalamic Kiss1 but not Kiss2
802   neurons are involved in estrogen feedback in medaka (Oryzias latipes). *Endocrinology,*
803   *151*(4), 1751-1759.
804
805   Muroi, Y., & Ishii, T. (2016). A novel neuropeptide Y neuronal pathway linking energy state and
806   reproductive behavior. *Neuropeptides, 59*, 1-8.
807
808   Naftolin, F., Garcia-Segura, L. M., Horvath, T. L., Zsarnovszky, A., Demir, N., Fadiel, A., Leranth, C.,
809   Vondracek-Klepper, S., Lewis, C., Chang, A., & Parducz, A. (2007). Estrogen-induced
810   hypothalamic synaptic plasticity and pituitary sensitization in the control of the
811   estrogen-induced gonadotrophin surge. *Reprod Sci, 14*(2), 101-116.
812
813   Nakajima, K., Cui, Z., Li, C., Meister, J., Cui, Y., Fu, O., Smith, A. S., Jain, S., Lowell, B. B., Krashes, M.
814   J., & Wess, J. (2016). Gs-coupled GPCR signalling in AgRP neurons triggers sustained
815   increase in food intake. *Nat Commun, 7*, 10268.
816
817   Nilsson, M. E., Vandenput, L., Tivesten, Å., Norlén, A. K., Lagerquist, M. K., Windahl, S. H.,
818   Börjesson, A. E., Farman, H. H., Poutanen, M., Benrick, A., Maliqueo, M., Stener-Victorin, E.,
819   Ryberg, H., & Ohlsson, C. (2015). Measurement of a Comprehensive Sex Steroid Profile in
820   Rodent Serum by High-Sensitive Gas Chromatography-Tandem Mass Spectrometry.
821   *Endocrinology, 156*(7), 2492-2502.
822
823   Ollmann, M. M., Wilson, B. D., Yang, Y. K., Kerns, J. A., Chen, Y., Gantz, I., & Barsh, G. S. (1997).
824   Antagonism of central melanocortin receptors in vitro and in vivo by agouti-related protein.
825   *Science, 278*(5335), 135-138.
826

Opazo, R., Plaza-Parrochia, F., Cardoso Dos Santos, G. R., Carneiro, G. R. A., Sardela, V. F., Romero,

J., & Valladares, L. (2018). Fasting Upregulates npy, agrp, and ghsr Without Increasing

Ghrelin Levels in Zebrafish (Danio rerio) Larvae. *Front Physiol, 9*, 1901.

Peute, J., van der Gaag, M. A., & Lambert, J. G. D. (1978). Ultrastructure and lipid content of the liver

of the zebrafish, Brachydanio rerio, related to vitellogenin synthesis. *Cell and Tissue*

Research, 186(2), 297-308.

Qian, S., Chen, H., Weingarth, D., Trumbauer, M. E., Novi, D. E., Guan, X., Yu, H., Shen, Z., Feng, Y.,

Frazier, E., Chen, A., Camacho, R. E., Shearman, L. P., Gopal-Truter, S., MacNeil, D. J., Van

der Ploeg, L. H., & Marsh, D. J. (2002). Neither agouti-related protein nor neuropeptide Y is

critically required for the regulation of energy homeostasis in mice. *Mol Cell Biol, 22*(14),

5027-5035.

Qiu, J., Rivera, H. M., Bosch, M. A., Padilla, S. L., Stincic, T. L., Palmiter, R. D., Kelly, M. J., &

Rønnekleiv, O. K. (2018). Estrogenic-dependent glutamatergic neurotransmission from

kisspeptin neurons governs feeding circuits in females. *Elife, 7*.

Robinson, E. J., & Rugh, R. (1943). The reproductive processes of the fish, Oryzias latipes. *The*

Biological Bulletin, 84(1), 115-125.

Roesch, D. M. (2006). Effects of selective estrogen receptor agonists on food intake and body weight

gain in rats. *Physiol Behav, 87*(1), 39-44.

Rønnestad, I., Gomes, A. S., Murashita, K., Angotzi, R., Jonsson, E., & Volkoff, H. (2017).

Appetite-Controlling Endocrine Systems in Teleosts. *Frontiers in Endocrinology, 8*, 73.

Schwartz, M. W., Woods, S. C., Porte, D., Seeley, R. J., & Baskin, D. G. (2000). Central nervous

system control of food intake. *Nature, 404*(6778), 661-671.

Shainer, I., Buchshtab, A., Hawkins, T. A., Wilson, S. W., Cone, R. D., & Gothilf, Y. (2017). Novel

hypophysiotropic AgRP2 neurons and pineal cells revealed by BAC transgenesis in zebrafish.

Scientific Reports, 7, 44777.

Shainer, I., Michel, M., Marquart, G. D., Bhandiwad, A. A., Zmora, N., Ben-Moshe Livne, Z., Zohar,

Y., Hazak, A., Mazon, Y., Forster, D., Hollander-Cohen, L., Cone, R. D., Burgess, H. A., &

Gothilf, Y. (2019). Agouti-Related Protein 2 Is a New Player in the Teleost Stress Response
System. *Current Biology, 29*(12), 2009-2019 e2007.
865
866   Shiozaki, K., Kawabe, M., Karasuyama, K., Kurachi, T., Hayashi, A., Ataka, K., Iwai, H., Takeno, H.,
867   Hayasaka, O., Kotani, T., Komatsu, M., & Inui, A. (2020). Neuropeptide Y deficiency induces
868   anxiety-like behaviours in zebrafish (Danio rerio). *Sci Rep, 10*(1), 5913.
869
870   Shutter, J. R., Graham, M., Kinsey, A. C., Scully, S., Lüthy, R., & Stark, K. L. (1997). Hypothalamic
871   expression of ART, a novel gene related to agouti, is up-regulated in obese and diabetic
872   mutant mice. *Genes Dev, 11*(5), 593-602.
873
874   Söderberg, C., Wraith, A., Ringvall, M., Yan, Y. L., Postlethwait, J. H., Brodin, L., & Larhammar, D.
875   (2000). Zebrafish genes for neuropeptide Y and peptide YY reveal origin by chromosome
876   duplication from an ancestral gene linked to the homeobox cluster. *J Neurochem, 75*(3),
877   908-918.
878
879   Song, Y., & Cone, R. D. (2007). Creation of a genetic model of obesity in a teleost. *FASEB Journal,
880   21*(9), 2042-2049.
881
882   Song, Y., Golling, G., Thacker, T. L., & Cone, R. D. (2003). Agouti-related protein (AGRP) is
883   conserved and regulated by metabolic state in the zebrafish, Danio rerio. *Endocrine, 22*(3),
884   257-265.
885
886   Stone, D. B., Cherry, M. J., Martin, J. A., Cohen, B. S., & Miller, K. V. (2017). Breeding chronology
887   and social interactions affect ungulate foraging behavior at a concentrated food resource.
888   *PLoS One, 12*(6), e0178477.
889
890   Sundstrom, G., Larsson, T. A., Brenner, S., Venkatesh, B., & Larhammar, D. (2008). Evolution of the
891   neuropeptide Y family: new genes by chromosome duplications in early vertebrates and in
892   teleost fishes. *General and Comparative Endocrinology, 155*(3), 705-716.
893
894   Takahashi, K. A., & Cone, R. D. (2005). Fasting induces a large, leptin-dependent increase in the
895   intrinsic action potential frequency of orexigenic arcuate nucleus neuropeptide
896   Y/Agouti-related protein neurons. *Endocrinology, 146*(3), 1043-1047.
897
Temple, J. L., & Rissman, E. F. (2000). Brief refeeding restores reproductive readiness in

food-restricted female musk shrews (Suncus murinus). *Horm Behav, 38*(1), 21-28.

Tohyama, S., Miyagawa, S., Lange, A., Ogino, Y., Mizutani, T., Ihara, M., Tanaka, H., Tatarazako, N.,
Kobayashi, T., Tyler, C. R., & Iguchi, T. (2016). Evolution of estrogen receptors in ray-finned
fish and their comparative responses to estrogenic substances. *J Steroid Biochem Mol Biol,*
158, 189-197.

Tohyama, S., Ogino, Y., Lange, A., Myosho, T., Kobayashi, T., Hirano, Y., Yamada, G., Sato, T.,
Tatarazako, N., Tyler, C. R., Iguchi, T., & Miyagawa, S. (2017). Establishment of estrogen
receptor 1 (ESR1)-knockout medaka: ESR1 is dispensable for sexual development and
reproduction in medaka, Oryzias latipes. *Development Growth & Differentiation, 59*(6),
552-561.

Tsuyuki, A. T., U. (2018). Assessment of ichthyofauna at oyster rafts in Hiroshima Bay, Japan, using
underwater video cameras. *Aquaculture Science, 66*(4), 267-274.

Umatani, C., Yoshida, N., Yamamoto, E., Akazome, Y., Mori, Y., Kanda, S., Okubo, K., & Oka, Y.
(2022). Co-existing Neuropeptide FF and Gonadotropin-Releasing Hormone 3 Coordinately
Modulate Male Sexual Behavior. *Endocrinology, 163*(2).

Volk, K. M., Pogrebna, V. V., Roberts, J. A., Zachry, J. E., Blythe, S. N., & Toporikova, N. (2017).
High-Fat, High-Sugar Diet Disrupts the Preovulatory Hormone Surge and Induces Cystic
Ovaries in Cycling Female Rats. *J Endocr Soc, 1*(12), 1488-1505.

Wintermantel, T. M., Campbell, R. E., Porteous, R., Bock, D., Gröne, H. J., Todman, M. G., Korach, K.
S., Greiner, E., Pérez, C. A., Schütz, G., & Herbison, A. E. (2006). Definition of estrogen
receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone
neurons and fertility. *Neuron, 52*(2), 271-280.

Yang, J. A., Stires, H., Belden, W. J., & Roepke, T. A. (2017). The Arcuate Estrogen-Regulated
Transcriptome: Estrogen Response Element-Dependent and -Independent Signaling of ERα
in Female Mice. *Endocrinology, 158*(3), 612-626.

Yokobori, E., Azuma, M., Nishiguchi, R., Kang, K. S., Kamijo, M., Uchiyama, M., & Matsuda, K.
(2012). Neuropeptide Y stimulates food intake in the Zebrafish, Danio rerio. *Journal of*
Neuroendocrinology, 24(5), 766-773.

Yu, K., He, Y., Hyseni, I., Pei, Z., Yang, Y., Xu, P., Cai, X., Liu, H., Qu, N., Liu, H., He, Y., Yu, M.,
Liang, C., Yang, T., Wang, J., Gourdy, P., Arnal, J. F., Lenfant, F., Xu, Y., & Wang, C. (2020).
17β-estradiol promotes acute refeeding in hungry mice via membrane-initiated ERα
signaling. *Mol Metab, 42*, 101053.

Zempo, B., Kanda, S., Okubo, K., Akazome, Y., & Oka, Y. (2013). Anatomical distribution of sex
steroid hormone receptors in the brain of female medaka. *Journal of Comparative Neurology,*
521(8), 1760-1780.

944
945

 Figure Legends

Figure 1: Reproductive female medaka show larger amount of food intake.

(A) Light conditions for breeding and non-breeding status. (B) Food intake (10 mins) of female

medaka in LD (orange; n = 7) and SD (blue; n = 7) conditions, normalized to the amount of artemia

eaten by medaka in LD (breeding) condition ($p = 0.02519$, $U = 42.5$). (C) Food intake (10 mins) of

male medaka in LD (orange; n = 7) and SD (blue; n = 7) conditions, normalized to the amount of

artemia eaten by medaka in LD (breeding) condition ($p = 0.6540$, $U = 20.5$). Mann-Whitney U test,

* $p < 0.05$. n.s., not significant.

Figure 1-Source code 1: The code of "Shrimp-counter" system

Figure 1-Supplementary Figure 1 "Wanko-soba" method for calculation of the amount of food

intake of fish

(A) One medaka is put into a white cup with 100 mL breeding water and is habituated for 5 min. (B)

Medaka is fed by application of 200 μL aliquots of food water containing brine shrimp in

all-you-can-eat style and serve another aliquot once done with it, which is repeated N times (like the

Japanese "Wanko soba"). (C) Stop feeding 10 min after the start, take 10 mL out of the breeding

water and transfer it to a conical tube. The conical tube is frozen overnight and the leftover brine

shrimp sunk in the bottom are counted by "shrimp-counter". The food intake is calculated as follows

(also see Material and Methods).

(Food Intake)

= (The average number of brine shrimp in the solution) * (number of aliquots, N)

- (number of leftover brine shrimp sunk in the bottom) * 10

Food intake was normalized by the average of LD medaka.

Figure 1-Supplementary Figure 2: Whole-brain gene expression in LD (n = 3) and SD (n = 3)

female medaka.

(A) Volcano plot comparing whole-brain gene expression in LD (n = 3) and SD (n = 3) female

medaka. Genes with higher expression in LD than in SD (P-value < 0.05 and logFC < -1) are shown

in orange, and those with higher expression in SD than in LD (P-value < 0.05 and logFC > 1) are

shown in blue. (B) Heatmap of representative genes, which are mainly related to neuroendocrine

functions and show different expression patterns between LD and SD. (C and D) TPM value of

agrp1 and *npyb*.

Figure 2: *agrp1*- and *npyb*-**expressing neurons are distributed locally, while npya- and** *agrp2*-

expressing neurons are distributed more widely in the brain.

(A) Illustration of the lateral view of medaka brain and distributions of cell bodies expressing each

agrp or *npy* gene. The oblique lines labeled with a-e indicate the level of the frontal sections in a-e.

(a-e) Illustrations of frontal sections showing the distribution of neurons expressing each gene. The

localization of neurons is indicated in the right half of the illustrations. BO bulbus olfactorius, dDm

dorsal region of area dorsalis telencephali pars medialis, Dl area dorsalis telencephali pars lateralis,

lfb lateral forebrain bundle, NAT nucleus anterior tuberis, NRL nucleus recessus lateralis, NVT

nucleus ventralis tuberis, PIT pituitary, POA area preoptica, POp nucleus preopticus pars

paravocellularis, TO tectum opticum, Vd area ventralis telencephali pars dorsalis, Vl area ventralis

telencephali pars lateralis, Vs area ventralis telencephali pars supracommissuralis, Vv area ventralis

telencephali pars ventralis. *agrp1* (gray circle), *agrp2* (open triangle: high expression, dotted

triangle: low expression), *npya* (star), *npyb* (open square). (B) *agrp1*-expressing neurons are

localized in NVT. Scale bar: 100 µm. (C) *agrp2*-expressing neurons are observed in Vv, POA, and

NVT. Scale bar: 100 µm. (D) *npya*-expressing neurons are distributed in dDm, Dl, BO, Vs, Vl, lfb,

NRL, NAT, NVT, and TO. Scale bar: 100 µm. (E) *npyb*-expressing neurons are localized in Vd.

Scale bar: 100 µm. (F) Left: *agrp1* (green) and *npya* (magenta) are distributed in NVT, but the two

genes are not co-expressed. Right: *agrp2* (magenta) and *npya* (green) are distributed in NVT, but the

two genes are not co-expressed. Scale bars: 50 µm. (G-J) *agrp* and *npy* expressions in the whole

brain of female medaka under normally fed condition (Fed; white; n = 7) or 2-week food restricted

(FR; gray; n = 8). (G) *agrp1* ($p = 0.001243$, $U = 2$) (H) *agrp2* ($p = 0.9551$, $U = 29$) (I) *npya* ($p =$

0.2319, $U = 39$) (J) *npyb* ($p = 0.0003108$, $U = 56$) Mann-Whitney U test, ** $p < 0.01$, *** $p < 0.001$.

n.s., not significant.

1003

Figure 2-Supplementary Figure 1 The number of *npya*-expressing neurons in the

hypothalamus of fed and food-restricted (FR) medaka

(A) Total number of *npya*-expressing neurons in the hypothalamus of female medaka under normal

fed condition (Fed; white; n = 4) or 2-weeks food restriction (FR; gray; n = 4). $p = 0.02857$, $U = 16$.

(B) The number of *npya*-expressing neurons in the NVT of female medaka under normal fed

condition (Fed; white; n = 4) or 2-weeks food restriction (FR; gray; n = 4). $p = 0.1143$, $U = 14$. (C)

The number of *npya*-expressing neurons in the NRL of female medaka under normal fed condition

(Fed; white; n = 4) or 2-weeks food restriction (FR; gray; n = 4). $p = 0.02857$, $U = 16$. (D) The

number of *npya*-expressing neurons in the NAT of female medaka under normal fed condition (Fed;

white; n = 4) or 2-weeks food restriction (FR; gray; n = 4). $p = 0.02857$, $U = 16$. Mann-Whitney U

test, *$p < 0.05$. (E) *agrp2* expression of female medaka under normal fed condition (Fed) or 10 days

food restriction (FR). Arrowheads indicate cell bodies of *agrp2*-expressing neurons. Scale bar: 100

µm.

1017

Figure 3: *agrp1* and *npyb* show higher expression levels in LD than in SD female.

(A) *agrp1* expression in the whole brain of LD (orange; n = 8) and SD (blue; n = 8) female medaka.

($p = 0.001865$, $U = 60$) (B) ISH of *agrp1*-expressing neurons in LD and SD female medaka. Scale

bar: 100 μm. (C) The number of neurons expressing *agrp1* in LD (orange; n = 5) and SD (blue; n =

5). ($p = 0.02828$, $U = 23$) (D) *npyb* expression in the whole brain of LD (orange; n = 8) and SD

(blue; n = 8) female medaka. ($p = 0.0001554$, $U = 64$) (E) The number of neurons expressing *npyb* in

LD (orange; n = 5) and SD (blue; n = 5). ($p = 0.4206$, $U = 17$) The upper, middle, and lower bars

show the third quartile, median, and the first quartile, respectively. Mann-Whitney U test, * $p < 0.05$,

** $p < 0.01$, *** $p < 0.001$. n.s., not significant.

1027

Figure 3-Supplementary Figure1: Time course of the number of neurons showing ISH signals

for *agrp1*.

(A) Time course of the number of neurons showing ISH signals for *agrp1* at each 30-min time point

from the beginning of the chromogenic reaction in LD (orange; n = 5) and SD (blue; n = 5). The

positive neurons were counted from 0 min to 300 min after application of NBT/BCIP (see Material

and Methods), and the reaction saturated at 300 min onward. (B) Ratio of cell number at 90 min

divided by that at 300 min. The relative cell number was not significantly different between LD and

SD. ($p = 0.2948$, $U = 18$)

1036

Figure 4: In juvenile female medaka, expression levels of neither *agrp1* nor *npyb* shows

significant change according to the day-length.

(A) *agrp1* expression in the brain of juvenile female medaka. ($p = 0.4179$, $U = 21$) (B) *npyb*

expression in the brain of juvenile female medaka. ($p = 0.3357$, $U = 19$) LD: orange, n = 8; SD: blue,

n = 7. The upper, middle, and lower bars show the third quartile, median, and the first quartile,

respectively. Mann-Whitney U test. n.s., not significant.

1043

Figure 4-Supplementary Figure 1: Food intake of juvenile female does not show significant

change according to the day-length.

(A) Method for counting food intake of juvenile. A fish was placed in a white cup containing 50 mL

breeding water. After 5 mins habituation, 200 μL shrimp water was applied. Fish was picked up 10

mins later by a dropper, and its fin was cut for the genotype of sex. The food intake was calculated

by subtraction from the number of applied shrimps to that of the leftover shrimps. (B) Food intake of

juvenile female medaka in LD (n = 9) and SD (n = 10) conditions. Each food intake was normalized

to the average food intake of LD fish. ($p = 0.7197$, $U = 50$) Mann-Whitney U test. n.s., not

significant.

1053

Figure 5: Estrogen, which is secreted from mature ovary, may affect the expression of *agrp1*.

(A) Relative expression of *agrp1* in Sham (white; n = 6), OVX (Ovariectomized medaka, gray; n = 6), and OVX + E (OVX medaka kept in the tank containing E2, yellow; n = 7). Sham vs OVX: $p = 0.04310$, OVX vs OVX+E: $p = 0.05790$, Sham vs OVX+E: $p = 0.2000$. (B) Relative expression of *npyb* in Sham, OVX, and OVX + E. Sham vs OVX: $p = 0.1386$, OVX vs OVX+E: $p = 0.9991$, Sham vs OVX+E: $p = 0.08120$. The upper, middle, and lower bars show the third quartile, median, and the first quartile, respectively. Steel-Dwass test, * $p < 0.05$. n.s., not significant. (C) Photographs of brain slices after experiments of double *in situ* hybridization (*agrp1* (green) and *estrogen receptors* (magenta)). (D) Expanded photograph of *agrp1* and *esr2a* co-expressing neurons (white arrowhead). Scale bars: 50 µm.

Figure 5-Supplementary Figure 1: Food intake of OVX female does not show significant change according to the day-length.

Food intake of OVX female medaka in LD (n = 7) or SD (n = 6) conditions. Food intake was normalized to the average food intake of LD fish. ($p = 0.7308$, $U = 18$) Mann-Whitney U test. n.s., not significant.

Figure 6: *agrp1*$^{-/-}$ female medaka show decrease in food intake and the number of fertilized eggs.

 (A) Lateral views of representative *agrp1*$^{+/+}$ and *agrp1*$^{-/-}$ female medaka (Left), and body weight, body length, and abdominal height (right) of *agrp1*$^{+/+}$ (white; n = 6) and *agrp1*$^{-/-}$ (gray; n = 6). All the fish are 4-month-old adult medaka. Scale bar: 1 cm. body weight: $p = 0.04113$, $U = 31$; body length: $p = 0.3939$, $U = 24$; abdominal length: $p = 0.002165$, $U = 36$. (B) Food intake (10 min) of *agrp1*$^{+/+}$ (white; n = 6) and *agrp1*$^{-/-}$ (gray; n = 6) female medaka. Each amount of food intake is normalized by the average number of that of *agrp1*$^{+/+}$ medaka. ($p = 0.004329$, $U = 35$) (C) Food intake (10 min) of LD *agrp1*$^{-/-}$ (white; n = 9) and SD *agrp1*$^{-/-}$ (gray; n = 7) female medaka. Each amount of food intake is normalized by the average number of that of LD *agrp1*$^{-/-}$ medaka. ($p = 0.5953$, $U = 37$) (D) The number of eggs spawned by *agrp1*$^{+/+}$ (white; n = 6) and *agrp1*$^{-/-}$ (gray; n = 5) female medaka in a day. Each female was paired with a wildtype male. ($p = 0.008658$, $U = 28.5$). (E) Photograph of ovary in representative *agrp1*$^{+/+}$ and *agrp1*$^{-/-}$ female (left) and the gonado-somatic index (GSI, right). n = 6 of each group. ($p = 0.06494$, $U = 30$) Scale bar: 1 mm. (F) Expression of gonadotropin genes (*lhb* and *fshb*) in the pituitary of *agrp1*$^{+/+}$ (white; n = 6) and *agrp1*$^{-/-}$ (gray; n = 6) female medaka. *lhb*: ($p = 0.008658$, $U = 34$); *fshb*: ($p = 0.02597$, $U = 32$). The upper, middle, and lower bars show the third quartile, median, and the first quartile, respectively. Mann-Whitney U test, * $p < 0.05$, ** $p < 0.01$. n.s., not significant.

Figure 6-Supplementary Figure 1: Mutation site of *agrp1* knockout medaka

(A) The DNA sequence of WT (*agrp1* $^{+/+}$) and mutated (*agrp1* $^{-/-}$) exon3 of *agrp1* gene. The

nucleotide indicated with magenta represents an insertion. (B) The predicted amino acid sequence of

agrp1 $^{+/+}$ and *agrp1* $^{-/-}$. Mutated amino acids are indicated with magenta. (C) AgRP1

immunoreactive signals were found in WT female brain but not in *agrp1* $^{-/-}$ one. Upper row: WT,

lower row: *agrp1* $^{-/-}$. Green: AgRP1 immunoreactive signals, and Blue: nuclei (DAPI). Scale bar: 50

µm.

1097

Figure 6-Supplementary Figure 2: Food intake of WT, *agrp1* hetero and homo knockout

medaka

Food intake of WT (*agrp1* $^{+/+}$, n = 8), hetero (*agrp1* $^{+/-}$, n = 7) and homo knockout (*agrp1* $^{-/-}$, n = 5)

female medaka. *agrp1* $^{+/+}$ vs *agrp1* $^{-/+}$: $p = 0.5470$, *agrp1* $^{+/+}$ vs *agrp1* $^{-/-}$: $p = 0.009353$, *agrp1* $^{-/+}$ vs

agrp1 $^{-/-}$: $p = 0.01234$. The upper, middle, and lower bars show the third quartile, median, and the

first quartile, respectively. Steel-Dwass test, $* p < 0.05$, $** p < 0.01$. n.s., not significant.

1104

Figure 7: Illustration of mechanisms of breeding season-dependent feeding behavior in

medaka suggested by the present study

(A) Long day (LD) condition in breeding season induces ovarian maturation, which facilitates

release of estrogen (E2) from the mature ovary. (B) High concentration of serum E2 increases *agrp1*

expression in the brain via the estrogen receptors, especially, *esr2a*. (C) Higher expression of AgRP1

is suggested to activate neural circuitry for feeding, which leads to an increase in food intake and egg

spawning of female medaka in the breeding season.

1112

A

a b cd e f

- ● *agrp1*
- △ *agrp2*
- ★ *npya*
- □ *npyb*

a
dDm · Dl · BO

b
Vd · Vv

c
Vs · Ifb · VI · POA

d
POp

e
TO · NVT · PIT

f
NAT · NRL · NVT

B *agrp1*
NVT

C *agrp2*
Vv · POA · NVT

D *npya*
dDm · Dl · BO · Vs · Ifb · VI · NRL · NAT · NVT · TO

E *npyb*
Vd

F *agrp1*/*npya*/DAPI   *agrp2*/*npya*/DAPI

G *agrp1*

Relative expression (/*rps13*)

**

Fed   FR

H *agrp2*

Relative expression (/*rps13*)

n.s.

Fed   FR

I *npya*

Relative expression (/*rps13*)

n.s.

Fed   FR

J *npyb*

Relative expression (/*rps13*)

***

Fed   FR

A

agrp1⁺/⁺

agrp1⁻/⁻

Body Weight (mg) *
Body length (mm) n.s.
Abdominal hight (mm) **

+/+   -/-

B

Normalized food intake **

agrp1⁺/⁺   *agrp1⁻/⁻*

C *agrp1⁻/⁻*

Normalized food intake n.s.

LD   SD

D

Number of Eggs **

agrp1⁺/⁺   *agrp1⁻/⁻*

E

agrp1⁺/⁺

agrp1⁻/⁻

P = 0.06494

GSI (%)

+/+   -/-

F

lhb **

Relative exp. (/rps13)

+/+   -/-

fshb *

Relative Exp. (/rps13)

+/+   -/-

(C) Increase in food intake in breeding season (LD)

(B) E2 signal via Esr increases *agrp1* expression

Long Day (LD)
Light 14 h Dark 10 h

Many eggs

(A) Mature ovary in LD facilitates E2 production

ovary   release   E2

hypothalamus (NVT)

esr1

?

agrp1

esr2a

?

esr2b

AgRP1

brain