GlyT2-positive interneurons regulate timing and variability of information transfer in a cerebellar-behavioural loop

Ensor Rafael Palacios[1,2], Conor Houghton[2] and Paul Chadderton[1]

[1]School of Physiology Pharmacology and Neuroscience, University of Bristol, University Walk, Bristol, UK. BS8 1TD

[2]School of Engineering Mathematics and Technology, University of Bristol, Woodland Road, Bristol, UK. BS8 1UB

Correspondence should be addressed to:

Ensor Rafael Palacios (ensorrafael.palacios@bristol.ac.uk);
Paul Chadderton (p.chadderton@bristol.ac.uk)

No of pages: 47 (5 Figures; 1 Table)

Abstract word count: 160
Introduction word count: 608
Discussion word count: 1497

ACKNOWLEDGEMENTS

This work was supported by a Wellcome Trust Neural Dynamics PhD studentship to ERP, Leverhulme Research Fellowship (RF-2021-533) to CH and Wellcome Trust Investigator Award (209453/Z/17/Z) to PC. The schematic images in Figure 1 were generated and provided by Elisabeth Meyer whom we also thank for comments on the manuscript.

The authors declare no competing interests.

ABSTRACT

GlyT2-positive interneurons, Golgi and Lugaro cells, reside in the input layer of the cerebellar cortex in a key position to influence information processing. Here, we examine the contribution of GlyT2-positive interneurons to network dynamics in Crus 1 of mouse lateral cerebellar cortex during free whisking. We recorded neuronal population activity using NeuroPixels probes before and after chemogenetic downregulation of GlyT2-positive interneurons. Under resting conditions, cerebellar population activity reliably encoded whisker movements. Reductions in the activity of GlyT2-positive cells produced mild increases in neural activity which did not significantly impair these sensorimotor representations. However, reduced Golgi and Lugaro cell inhibition did increase the temporal alignment of local population network activity at the initiation of movement. These network alterations had variable impacts on behaviour, producing both increases and decreases in whisking velocity. Our results suggest that inhibition mediated by GlyT2-positive interneurons primarily governs the temporal patterning of population activity, which in turn is required to support downstream cerebellar dynamics and behavioural coordination.

SIGNIFICANCE STATEMENT

The cerebellum has a simple and conserved structure which has tantalised neurobiologists wishing to understand its function. Here we look at the role of granule cell layer inhibitory interneurons, Golgi and Lugaro cells, in the cerebellar cortex. We selectively turned down the activity of these cells in the awake cerebellum to characterise their influence on network activity and behaviour. We show that downregulation of Golgi and Lugaro cells has very little influence on sensorimotor representations in the cerebellum (i.e., *what* is represented), but instead modulates the timing of cortical population activity (i.e., *when* information is represented). Our results indicate that inhibitory interneurons in the granule cell layer are necessary to appropriately pace changes in cerebellar activity to match ongoing behaviour.

INTRODUCTION

The cerebellum is implicated in a diverse range of behaviours, extending from simple reflexes to complex functions such as language, and social interaction (Baumann & Mattingley, 2022; Kelly et al., 2020; Silveri, 2021). The circuit architecture underpinning this behavioural diversity is relatively simple and conserved across its extent, suggesting that different classes of input are subject to common processing mechanisms. The cerebellar cortex lacks recurrent excitation and almost all local feedback is mediated via GlyT2-positive inhibitory interneurons situated in the granule cell layer. Golgi cells receive direct input from MFs and excitatory feedback from PFs, enabling them to exert feedforward and feedback control over granule cells (Dieudonne, 1998; Eccles et al., 1966; Kanichay & Silver, 2008). Golgi cells are numerically sparse relative to granule cells and possess extended axonal arborisations, meaning each Golgi cell innervates many hundreds or thousands of granule cells. Inhibition via Golgi cells could thus potently impact the transformation of MF signals into PF activity. *In vivo* measurements confirm that inhibition gates the transformation of sensorimotor information in the granular layer (Chadderton et al., 2004; Duguid et al., 2012), sparsifying responses to sensory input, and facilitating pattern separation at the population level (Fleming et al., 2024). Lugaro cells make up approximately one third of granule cell layer inhibitory interneurons (Simat et al., 2007), and are known to make synaptic connections onto both Golgi cells and molecular layer interneurons (Eyre & Nusser, 2016). Although comparatively little is known about their function *in vivo* (Prestori et al., 2019), their sensitivity to serotonin levels in the cerebellar cortex (Dieudonne & Dumoulin, 2000) suggests Lugaro cells could regulate information processing in the granule cell layer based on behavioural context. Currently, we lack quantitative information about how changes in Golgi/Lugaro cell activity influences cerebellar cortical dynamics and affects sensorimotor representations in the molecular and Purkinje cell layers. Indeed, the net influence of granule cell layer inhibitory interneurons is not known but may be presumed to alter both the rate and timing of granule cell output. Resultant changes in PF excitation could have net excitatory or inhibitory effects at the level of Purkinje cells, and consequently for cerebellar nuclear neurons.

Here we investigate the influence of granule cell layer inhibitory interneurons on
cerebellar population activity in the context of whisker movement, to reveal how
inhibitory feedback influences sensorimotor transformations and ongoing behaviour.
Neuronal representations of whisking activity in the lateral cerebellum are widespread
(Bosman et al., 2010; Brown & Raman, 2018; Chen et al., 2016, 2017; Zhai et al.,
2024), and are partly based on a linear rate code, whereby the firing rates of cerebellar
neurons predict upcoming whisker position (Chen et al., 2016). These representations
are first seen in patterns of mossy fibre input and are passed forward at each stage of
cerebellar cortical processing (Chen et al., 2017). Golgi cell inhibition has the potential
to modulate how incoming whisking information is encoded and transformed within the
granule cell layer (Gurnani & Silver, 2021; Palacios et al., 2021). However, it is
unknown whether, and how, changing granule cell layer inhibition impacts neuronal
representations of whisker behaviour, and indeed whisking behaviour itself. We have
recorded whisker movements and population activity in the lateral cerebellum using
NeuroPixels probes in head-fixed mice expressing inhibitory DREADDs (designer
receptors exclusively activated by designer drugs) selectively in Golgi and Lugaro cells
(i.e. GlyT2-positive neurons) of lobule Crus 1. In these mice, it was possible to
downregulate GlyT2-positive cell activity by activating DREADDs with the agonist,
clozapine N-oxide (CNO), applied topically on the recording site. Our results reveal a
subtle but direct influence of GlyT2-positive cells on network dynamics and
consequently, the cerebellar-behavioural loop underlying voluntary whisking.

MATERIALS AND METHODS

Experiments were conducted on male and female C57BL/6 (wild-type) and heterozygous C57BL/6-Slc6a5<tm1.1(cre)Ksak> (GlyT2-Cre knockin) mice (Kakizaki et al., 2017) aged between 3-6 months. Animals were housed under a 12/12 h reversed light–dark cycle with food and water available *ad libitum*. Habituation and recordings were performed during the dark phase of the cycle. The C57BL/6-Slc6a5<tm1.1(cre)Ksak> mouse strain was provided by the RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan. All experiments were conducted under the United Kingdom Animals (Scientific Procedures) Act 1986 in accordance with Home Office guidelines.

Surgical procedures

Mice were anaesthetised for all surgical procedures. Mice were placed in an induction chamber, into which 5% v/v isoflurane (Harvard Apparatus Ltd) was administered via inhalation using oxygen (1.2–1.6 L/min). Once anaesthetized the concentration of isoflurane was lowered to 1–2% v/v to maintain anaesthesia for the duration of the surgery. The body temperature of the mouse was maintained at 37 ± 0.5 °C using a homeothermic heat mat (DC Temperature Control System, FHC). Corneal drying was prevented using an ocular lubricant (Lacrilube, Allergen). To induce analgesia, mice were injected subcutaneously with carprofen (Rimadyl, 5 mg/kg), buprenorphine (Vetergesic, 0.1 mg/kg) and lidocaine (2 mg/kg, locally). Surgery was carried out in a stereotaxic frame under aseptic conditions to prevent infections. After surgery, mice were placed in a warmed box (~37°C) and allowed to recover for as long as required. Mice were then returned to their home cages and monitored closely for at least 24 hrs.

GlyT2-Cre mice underwent two surgical operations: one to transduce cre-dependent expression of DREADDs in the cerebellar cortex via viral injection, and one to prepare the animal for head-fixed extracellular recording. During the first surgical session, a small craniotomy (~1 mm diameter) was performed over Crus 1 (6.36 mm posterior and 2.5 mm lateral of Bregma). AAV9-hSyn-DIO-hM4D(Gi)-mCherry (500-1000 nl)

was injected with a glass pipette at different depths (starting at 600µm retracting to
surface in steps of 100µm), waiting 2 minutes between pipette retractions. At the end
of the operation, skin over the head was sutured and animal was allowed to recover.

The second surgical session took place 8 weeks later to enable ample expression of
the DREADD receptor protein. First, neck muscles were gently moved to uncover the
cerebellum. A custom-made head implant was fixed to the exposed cranium using
tissue glue (Histoacryl, Braun Corporation) and dental cement (Associate Dental
Products Ltd). A small craniotomy (~1.5 mm diameter) was performed over the right
cerebellar hemisphere, the dura was removed, and a reference screw was inserted in
contact with the underlying brain. To secure both head place and reference screw,
dental cement (Associate Dental Products Ltd) was used to cover the skull, sparing a
recording well above Crus 1 of the left cerebellar hemisphere. A second craniotomy
(~2 mm diameter) was performed over left Crus 1, and the dura was removed. Finally,
a layer of agarose (1.5% in phosphate buffered saline; PBS) was used to cover the
brain, a layer of Kwik-Seal was used to protect the brain, and a layer of nail polish was
used to fix the Kwik-Seal. Wild-type mice underwent only the second surgical
operation in preparation for recordings. All mice had their whiskers trimmed on the left
(recording) side, preserving whiskers C1, C2 and C3 (posterior whiskers on the third
row from top).

Electrophysiological and video recordings

Mice were head-fixed in the recording apparatus in preparation for electrophysiological
recording. The Kwik-Seal was removed, the surface of the brain was cleaned from
agarose and kept moist with PBS. NeuroPixels probes were fixed to a
micromanipulator (IVM, Scientifica) via a custom-made 3D printed holder. Each probe
was coated with DiI stain (2.5 mg/ml) for *ex vivo* probe tracking before insertion at an
angle perpendicular to the cortical surface. The probe was advanced into the cortex
at a speed of 2-5 µm/s to a depth of 2500-3000 µm. Thereafter the probe was retracted
for ~100 µm and left to settle for 10-15 min, allowing the brain to relax. Under infrared
light illumination, whisker movements were filmed with a high-speed camera (Genie

HM640; Teledyne Dalsa Inc, USA) operating at 299 frames per second. Video

acquisitions were controlled by Streampix 6 software (Norpix, Canada). The open-

source software spikeGLX 3.0 (https://billkarsh.github.io/SpikeGLX/) was used to

record NeuroPixels data at 30 kHz. Video and electrophysiological recording were

synchronised by a TTL pulse originating from the video software. In chemogenetic

experiments, a baseline of 10 or 20 min was recorded prior to topical administration of

CNO (30 µM), delivered to the recording well. Previously Stachniak et al., 2014

showed that intracranial injection of 1 µM CNO is already sufficient to reliably inhibit

presynaptic neurotransmitter release within minutes. After CNO or vehicle delivery,

electrophysiological recordings continued for a further 40-50 mins, for a total recording

time of 1 hour. Animals underwent two recording sessions, the first performed at least

4 hours after recovery from the craniotomy, the second on the following day. After

recording, the probe was extracted, rinsed with deionised water and left in freshly

made 1% tergazyme solution for at least 24 hours. After the first recording, the

exposed brain surface was first rinsed with PBS, then covered with agarose, and finally

protected with a layer of Kwik-Seal that was fixed to the surrounding dental cement

with nail polish. Mice were returned to their cage. After the second recording session,

mice were humanely killed. See **Table 1** for full information on data included for

analysis.

Electrophysiological data processing

Output files (.imec) were preprocessed using the command-line tool CatGT

(https://billkarsh.github.io/ SpikeGLX/#catgt) to apply a high pass filter (cut-off 300 Hz)

and global demux filters, a common average referencing that takes in account the

probe channels subgrouping during data acquisition. Spike sorting was conducted

using the open-source software Kilosort 2 (https://github.com/MouseLand/ Kilosort) to

group spikes into units, each representing the activity of a distinct putative neuron.

Manual curation of sorted units was done with the open-source Python-library Phy2

(https://github.com/cortex-lab/phy), and consisted in merging and splitting units, as

well as categorising them into 'good', 'bad' and 'mua' (multiunit activity) units. In all

analysis, only good, i.e. well-isolated, units located in the cerebellar cortex were used.

Video processing and behavioural analysis

Video output (.avi) files were transcoded (.mp4, libx265 encoder), re-sampled at 299 Hz) and cropped using the ffmpeg software. We used the open-source toolbox DeepLabCut (https://github.com/DeepLabCut/DeepLabCut) to label four landmarks on whiskers C1, C2 and C3 (**Figure 1a**) across the entire recording; these landmarks were then used to compute whisker position. Specifically, the arctan function was used to measure the azimuthal angle between the first whisker segment (traced from the first, basal marker to the second marker) and the horizontal line passing through the basal marker. Increases in angle correspond to whisker protraction, whereas decreases correspond to whisker retraction. Next, the phase of the whisking cycles was extracted using the Hilbert transform, in order to compute the slowly changing whisking amplitude and setpoint across cycles (Hill et al., 2011). Finally, the whisking amplitude was low pass filtered and used to identify periods of whisking. A heaviside function with a threshold of 10°, was used to discriminate whisking from resting periods. All whisking analysis focused on the C3 (most anterior) whisker.

Histology

Brains were extracted and left in 4% PFA solution for 24-48 hours at 4°C. After washing with PBS, cerebellar coronal sections (50-100 μm) were sliced using a vibrating microtome (Leica VT1000S). Fluorescent images were acquired using a confocal microscope (Leica DM4000 B).

EXPERIMENTAL DESIGN AND STATISTICAL ANALYSES

Single unit analysis

To compute the tuning curve for each unit, we computed the total spike count and average whisker position (angle) in time bins of 33 ms. We then discretised the average whisking angles into 11 bins, ranging from the minimum to the maximum angle observed in each recording and paired the spike count in each time bin with the

corresponding whisker angle. These data were finally used to plot the firing rate,
together with their standard error, across the 11 angle bins.

Both the entropy of and KL-divergence between the pre- and post-drug tuning curves
for each unit was computed using the function entropy from the scipy.stats library. The
tuning curve for each unit was normalised to be the probability mass function of firing
rates over whisking position.

Clustering of tuning curves was performed with k-means algorithm
(sklearn.cluster.KMeans package), used on the spline coefficient fitted to the individual
unit tuning curves. The estimation of the tuning curves can be noisy due to, for
example, errors during the spike sorting process, which may lead to the contamination
of the spike history assigned to one unit by the activity of nearby neurons. This noise,
in turn, could affect clustering of tuning curves into meaningful groups. To reduce the
impact of this source of noise on our clustering analysis, we fitted a spline model to
each tuning curve: the model used seven cubic b-spline functions to smooth the firing
rate associated to each angle bin based on the firing rate of adjacent bins, while
maintaining the underlying structure of the tuning curves. The smoothed tuning curves
were then clustered using a k-means algorithm with $k = 8$. We finally computed the
average tuning curve from units within each cluster. Results were qualitatively the
same with fewer than $k = 8$ clusters.

Principal component analysis of population activity

We used principal component analysis to describe the population activity of each
recording (number of units per analysis can be found in **Table 1**). Projections onto the
first three eigenvectors (principal components; pcs) described the first three
orthogonal axes in neuronal space along which population activity varied the most. To
perform this analysis, we first segmented each recording into discrete whisking epochs
(from -2 to +3 seconds around the start of each whisking bout, defined using
behavioural analysis above), and computed the trial-averaged firing rate for each unit.
Firing rates were used to compute correlation matrices across units, from which

eigenvalues and eigenvectors were derived. Eigenvalues indicate the amount of
neuronal activity variability captured along the axis specified by the corresponding
eigenvectors. For each recording, the eigenvectors had length equal to the number of
units present in that recording. The values of an eigenvector are known as loads and
correspond to the weight associated to each unit when linearly combining the firing
rate from all units; hence, the loadings of an eigenvector correspond to how much
each unit contributes to the pc associated with that eigenvector. The eigenvectors can
be used to project either each unit's firing rates in pc space or single trial activity: to
compute the pc for single trial activity, we first smoothed single trial spike counts with
a Gaussian window of std = 20 (we also used std = 5 but results were equivalent; time
bin width = 3.3 ms).

Cross-correlation between whisking position and each pc was obtained using the
scipy.signal.correlate package. All cross-correlations were normalised by the absolute
maximum value of the first pc. For analysis involving CNO manipulation, cross-
correlation peaks for pre- and post-drop periods were computed, and the respective
distributions were compared using a two-sided Wilcoxon signed-rank test. To
compared cross-correlations across conditions, we first computed the within-recording
contrast between post- versus pre-drop cross correlation peaks; next, for each pc and
experimental condition, we fitted the contrasts with a Gaussian distribution, and finally
we took the difference in posterior samples of the mean between GlyT2-CNO and
control conditions.

Analysis to assess efficacy of DREADD manipulation

A multilevel model was used to account for the variability in total cerebellar cortical
spike counts detected over time across recordings; counts were the sum of the spikes
assigned to good (i.e. well-isolated) cortical units. Spike counts were computed for
successive 5 minute-time bins, with the 0-time bin including the time of drug
application. The counts in each bin were scaled with respect to the count in the -5-
minute bin.

In the model, spike counts were described using an inverse-Gamma distribution:

295                     $s \sim \text{Inverse-Gamma}(\alpha, \beta)$

296                     $\alpha = \exp(\theta_\alpha + \theta_{\alpha c} + \theta_{\alpha a} + \theta_{\alpha t})$ (1)

297                     $\beta = \exp(\theta_\beta + \theta_{\beta c} + \theta_{\beta a} + \theta_{\beta t})$

The inverse-Gamma distribution is used because it provides a good fit to the
population spike count data (**Figure 3e**); the two parameters α and β are usually
referred to as the shape and scale parameters and are related to the mean and
variance by $\mu = \beta/(\alpha - 1)$ and $\sigma^2 = \beta^2/[(\alpha-2)^2(\alpha-1)]$. Here α and β are each linked to a
linear model by a loglink function. Both linear models involve a coefficient θ for each
explanatory variable, namely, the experimental condition c, the time of the recording t
(dim = 9), and the animal ID a (dim = 17). The experimental conditions were, (i)
application of CNO to cerebellar surface of DREADD-transduced GlyT2-Cre mice
(GlyT2-CNO, n = 19 recordings), (ii) application of CNO to cerebellar surface of wild-
type mice (WT-CNO, n = 5 recordings), (ii) application of PBS the cerebellar surface
of wild-type or GlyT2 mice (WT-Veh, n = 9 recordings). Experimental conditions were
split by recording period into a pre-drug or baseline period (bins before 0-time bin) a
post-drug period (0-time bin and onward), for a total dimensionality of 3 x 2 (dim=6)
for the $\theta_{\alpha c}$ and $\theta_{\beta c}$ coefficient. The contrasts between pre- and post-drug coefficient θ_c
were used to assess the effect of the experimental manipulation.

The additional explanatory variables, t and a, were used to explain respectively spike
count variability within each pre-drug time bin due to natural changes in network
activity over recording time (n = 9, excluding -5-min time bin), and variability due to the
particular conditions of the network in each animal (n = 17), which might be affected,
for example, by the surgical procedure. Thus, the addition of these explanatory
variables allowed us to test within conditions for a difference in pre- and post-drop
spike count distribution, after accounting for variability explained by temporal
correlations and between-animal differences.

Each θ, except θ_t was drawn from a vector of independent normal distributions, of
length equal to the dimensionality of θ. The prior distribution for θ_t was a multivariate
normal distribution with covariance decaying with time distance, accounting for
temporal correlations within the data:

$$\Sigma_\alpha = I * \exp(-\delta_{tij}) \qquad \Sigma_\beta = I * \exp(-\delta_{tij})$$

$$\theta_\alpha \sim \mathrm{Normal}(\mu_\alpha, \sigma_\alpha) \qquad \theta_\beta \sim \mathrm{Normal}(\mu_\beta, \sigma_\beta)$$

$$\theta_{\alpha c} \sim \mathrm{Normal}(\mu_\alpha, \sigma_\alpha) \qquad \theta_{\beta c} \sim \mathrm{Normal}(\mu_\beta, \sigma_\beta) \qquad (2)$$

$$\theta_{\alpha a} \sim \mathrm{Normal}(\mu_\alpha, \sigma_\alpha) \qquad \theta_{\beta a} \sim \mathrm{Normal}(\mu_\beta, \sigma_\beta)$$

$$\theta_{\alpha t} \sim \mathrm{MVNormal}(\mu_\alpha, \Sigma_\alpha) \qquad \theta_{\beta t} \sim \mathrm{MVNormal}(\mu_\beta, \Sigma_\beta)$$

with δ_{tij} between any two periods i and j is defined as:

$$\delta_{tij} = exp\left(-\left|\frac{ti - tj}{5}\right| * 0.7\right)$$

$$(3)$$

Where $\left|t_i - t_j\right|/5$ is the absolute time difference between period i and j scaled to unity
(5 is the time width of each period in minutes).

Finally, the hyperpriors:

$$\mu_\alpha = 1 \qquad \mu_\beta = 1.2$$

$$(4)$$

$$\sigma_\alpha = 0.2 \qquad \sigma_\beta = 0.2$$

were chosen to sample prior predictive samples within a similar scale of the observed
population spike counts, including extreme values.

The posterior samples for the mean and variance spike count parameters were used
to compute the difference across conditions between post- and pre-drug contrasts in
mean and variance due to the experimental manipulation (**Figure 3f**). In practice, the
contrasts were computed using the same posterior samples, except for $\theta_{\alpha c}$ and $\theta_{\beta c}$,
which differed for pre- and post-drug mean and variance. The difference between
contrasts was used to assess changes in mean and variance across conditions
(**Figure 3g**).

Analysis of DREADD manipulation on neuronal and whisking behaviour

Comparisons were made from the pre- and post-drug time periods. The pre-drug
period lasted 20 minutes all but five recordings, where this period was 10 minutes. The
post-drug time period matched the pre-drug period and started 5 minutes after drug
application. Recordings that exhibited little or no whisking activity were excluded from
this analysis leaving N = 25 recordings in total.

For each population recording, we computed the absolute peak of the trial-averaged
neuronal activity (PETH) aligned to whisking onset. The subwindow used to compute
the absolute peaks spanned -0.7 to 1.3 sec around whisking onset, to focus the
subsequent analysis around the initial whisking protraction period. We used the
absolute peak of the PETH as a proxy of the timing of the neuronal response to
whisking-related input. For each recording we computed the standard deviation (std)
of the peaks, for pre- and post-drug data. For each experimental condition, the pre-
and post-drug stds were compared using a two-sided Wilcoxon signed-rank test.

For corresponding whisking data, we used a linear model to fit a line to the average
whisking position pre- and post-drug. Whisking data were delimited by a time window
ranging from -0.06 to 0.21 sec centred around whisking onset; the time window was
chosen to focus on the period of initial whisking protraction. We computed the
difference between post-drop and pre-drug slopes (i.e. velocity to compare the effects
of Golgi cell manipulation on the whisking protraction phase. To test whether the

variance of the distribution slope contrasts in the two experimental conditions were

different, we used Levene's test (scipy.stats.levene).

Code accessibility

Code used for the analysis is freely available

here: https://zenodo.org/doi/10.5281/zenodo.13327395

RESULTS

Population dynamics in lateral cerebellum during voluntary whisking

To understand the contribution of GlyT2-positive interneurons to cerebellar cortical

dynamics, we first set out to describe how cerebellar populations represent

sensorimotor information while Golgi and Lugaro cell inhibition was intact. We

recorded cerebellar population activity using NeuroPixels probes in lobule Crus 1 of

awake mice and performed spike sorting to isolate single units (see Materials and

Methods). We isolated an average of 42.3 (std = 25.3) cortical units per recording (N

= 12), for a total (n) of 508 putative units. To correlate spiking activity with whisker

movement, we simultaneously tracked spontaneous whisking via a high-speed

camera (**Figure 1a**). In all cases, bouts of whisking activity were associated with

pronounced alterations in cerebellar spiking activity (**Figure 1b, c**). Single cell

recordings previously revealed widespread tuning to whisker position in granule cells,

inhibitory interneurons and Purkinje cells (Bosman et al., 2010; Brown & Raman, 2018;

Chen et al., 2016, 2017; Zhai et al., 2024). We therefore measured tuning to whisker

position amongst single units of our population recordings. On a unit-by-unit basis, we

examined the relationship between firing rate and whisker position, to construct

individual tuning curves (see Materials and Methods). Consistent with previous work,

we observed robust tuning to whisker position, such that single units exhibited their

highest firing rates at preferred angles of whisker deflection.

Within locally recorded populations, we observed substantial diversity in the tuning to

whisker position. Indeed, units recorded in the same penetration displayed a broad

range of preferred angles and tuning profiles, including monotonic increases and

decreases to whisker protraction/retraction (**Figure 1d)**. We then recombined all single

unit spiking responses to calculate population tuning. Whereas individual units

exhibited diverse selectivity, the mean tuning function for the population displayed a

near-monotonic dependence upon whisker position, with rate changes in proportion to

the magnitude of protraction and retraction (**Figure 1e**).

To further assess tuning diversity at the population level, we clustered units based on

their tuning curves, testing whether we could discriminate discrete functional

subclasses of unit. We fitted a spline model with 7 cubic b-spline to each tuning curve

(n = 508, N = 12) to capture tuning curve shape and reduce variability due to noise

(see Materials and Methods). We then inspected the output of k-means clustering with

different numbers of expected clusters. With increasing cluster number, there was no

evidence of a change in the sum-of-squared-distances decaying pattern, or elbow,

which can be used as an indicator of the optimal number of clusters to be considered.

Visual inspection of the mean tuning curves for each cluster revealed a progressive

shift of preferred position from the most retracted to the most protracted whisker

locations (**Figure 1f**). Projection of tuning curves in 3D space using t-SNE, a non-

linear dimensionality reduction technique particularly suited to uncovering clustering

structure further confirmed that different groups of clusters are not well isolated (data

not shown). Thus, our analysis does not support discrete classes of tuning curve but

rather confirms that cerebellar neurons form a functional continuum in the space of

possible tuning curves. At the local population level, Crus 1 cerebellar cortical neurons

heterogeneously encode whisking position, but represent increasing protractive and

retractive movements via accumulating increases in firing rates.

Having established that concerted population activity is required for accurate

sensorimotor representations, we performed principal component analysis upon the

grouped spike trains from each recording. We initially preserved projections of

population activity onto the first three eigenvectors, which describe the three

orthogonal axes in neuronal space along which activity varies the most (**Figure 1g**).

Analysis was restricted around whisking bouts (-2 to +3 seconds about whisking

onset). Projections of population activity revealed clear structure in neuronal space

during transitions from quiescence to whisking activity and back again. This structure

in turn suggested that population activity contains accurate information about whisking

dynamics. We tested this proposal by linearly combining the first three principal

components, attempting to reconstruct movement trajectories (see Materials and

Methods). Using this approach, it was possible to decode whisking setpoint during

single trials (**Figure 1h**). In summary, our results show that the population activity in

the lateral cerebellar cortex accurately reflects upcoming slow whisking dynamics,

even when approximated within a low-dimensional space.

To quantify the relationship between different principal components and whisking, we

computed the normalised cross-correlation between whisker angle and each of the

first 3 principal components (pc) in each recording. Interestingly, pc1-3 showed distinct

temporal relationships with respect to whisker movement (**Figure 2a,b**). Information

related to whisker position contained in pc1 tended to anticipate behaviour, whereas

information in pc2 and pc3 lagged behaviour. Across all recordings (N = 12), the peak

signal of pc1 always preceded the whisking signal, indicating that, across recordings,

variation in the whisking-aligned average neuronal activity expressed by the pc1 tends

to anticipate behaviour (one sample t test, t = 4.72, p = 0.0006, mean = 36 ± 8 ms).

On the other hand, the peak signal for pc2 almost always occurred after whisker

movement, indicating that whisker-related information contained in pc2 lags whisking

activity (t = -4.43, p = 0.001, mean = -195 ± 44 ms). Finally, peak signals for pc3 were

broadly distributed around 0 ms (t = -1.38, p = 0.19, mean = -89 ± 65 ms). Notably,

pc1 peaks across recordings are narrowly distributed in time; this suggests that pc1

specifically contains precise information about whisking behaviour, i.e. upcoming

movement, occurring ~40 ms in the future (**Figure 2c**).

Whisking-related information contained in pc1-3 could originate from a small

proportion of units in the population, whose activity may be particularly responsive to

whisking; alternatively, information could be more distributed, whereby all units

contribute to some extent to the population encoding of behaviour. To measure how

behavioural information was distributed across the population, we measured the pc

loading of each unit (n = 508), indicating the contribution of a unit to the corresponding

pc (see Materials and Methods). The loading distribution for each pc was centred

around 0 and was roughly symmetric, meaning that units could contribute both

positively or negatively to each pc (**Figure 2d**). The number of outliers in each

distribution was used to quantify the extent to which each pc is dominated by the
activity of few units. We calculated the excess kurtosis of each distribution, which is a
measure of the frequency of outliers observed, using as a point of reference the
frequency expected from normally distributed data. The distribution for pc1 had an
excess kurtosis of -2.85, indicating infrequent outliers; this in turn suggests that the
contribution of units across recordings to pc1 is broadly distributed within the
population. In contrast, the excess kurtosis for pc3 was 4.54, indicating a high
frequency of outliers, and thus a subset of units which tend alone to contribute mostly
to pc3. The excess kurtosis for pc2 was -0.47, meaning that outliers are neither
frequent nor infrequent. Together, these results suggest that for pc1, and thus accurate
representation of whisking behaviour, information is broadly distributed across the
entire population. Finally, we noted that the first three eigenvalues accounted for only
a moderate amount of the total variance in the population activity (~30%) across all
recordings (**Figure 2e,f**). Therefore pc1-3 provide only a coarse description of
population activity, and whisker movement only partially describes neuronal dynamics
in Crus 1 despite restricting our analysis to the initial period of whisking activity. Indeed,
the large proportion of neuronal variability not captured in the first 3 principal
components supports the proposal that cerebellar cortex may represent additional
behaviourally relevant variables.

Chemogenetic downregulation of GlyT2-positive interneurons elevates
population activity

Having established the relationship between population activity and slow whisking
dynamics under baseline conditions, we next sought to understand how Golgi and
Lugaro cell inhibition contributes to these representations in the cerebellar cortex. We
adopted a chemogenetic strategy to selectively downregulate GlyT2-positive neuronal
activity in an otherwise intact network while simultaneously recording population
activity and whisker movement. Manipulation of GlyT2-positive cells was achieved via
activation of the inhibitory DREADD, hM4Di, which causes reductions in excitability
and neurotransmitter release (Stachniak et al., 2014) in cells expressing these

receptors. Localised expression of DREADDs (**Figure 3b**) was achieved by injecting
the AAV-DIO-hM4D(Gi)-mCherry virus into Crus 1 of GlyT2-Cre mice (cre-
recombinase restricted to GlyT2-positive cells in the cerebellar cortex; see Materials
and Methods; Kakizaki et al., 2017). DREADDs were activated by local, topical
delivery of the agonist, clozapine-N-oxide (CNO), on top of the Neuropixels
penetration site, which itself had been targeted as the site of virus injection (**Figure**
3a); this method was used to minimise potential off-site effects of the agonist and
afforded within-minute temporal precision to our manipulation. To confirm the fidelity
of our approach, we first tested the effects of topical delivery of muscimol (1.7 μg/μl),
a GABA$_A$-receptor agonist, during NeuroPixels recordings from wildtype mice (N = 3).
In all cases, within minutes of drug application, the total spike count recorded across
the cerebellar cortex started to decrease and reached its minimum within 10-30
minutes (data not shown). These results supported the viability of applying CNO
topically onto cerebellar cortex to locally activate DREADD receptors.

We next examined whether CNO-dependent reduction in GlyT2-positive cell activity
had a measurable impact on cerebellar cortical population activity. To do this, we
compared changes in cortical spike count before and after CNO administration across
three experimental groups. GlyT2-Cre mice, pre-injected with AAV9-hSyn-DIO-
hM4D(Gi)-mCherry virus into Crus 1, or uninjected C57BL6 mice received topical
delivery of CNO (GlyT2-CNO; *N* = 19 recordings) or saline vehicle (Veh; *N* = 9
recordings) to the cerebellar cortex. A third group of uninjected wildtype (C57BL6) mice
received topical delivery of CNO to the cerebellum (WT-CNO; *N* = 5 recordings). Initial
inspection of the data revealed an increase in normalised spike count in the GlyT2-
CNO groups, but not Veh and WT-CNO groups, shortly after application of CNO
(**Figure 3c,d**). Additionally, when comparing the mean of the normalised post-drop
cortical spike counts (5-minute bin width), we found a significant difference across
conditions (ANOVA, *F* = 10.91, *p* = 2.82e-05). However, significant variability in spike
counts were observed both within and between experimental groups. Therefore, to
confirm that alterations in spike count observed across recordings were due to
chemogenetic manipulation of GlyT2-positive cells, we used a multilevel model

approach; this approach was warranted by the temporal correlations between
repeated measures and clustering by animals. The model described the distribution of
spike counts for each time bin with an inverse-Gamma distribution, parameterised by
a shape parameter α and a scale parameter β. The model used as explanatory
variables the experimental conditions, associated to pre- and post-drug administration
periods, to find the α and β parameters that best explained the pre- and post-drug
administration data. For each parameter of the gamma distribution ($α$ and $β$) and each
group (WT-CNO, Veh, GlyT2-CNO), the model had a coefficient $θ$ associated to the
pre-drug recording period and one for the post-drug administration period. The
contrast between the two coefficients was used to measure the change in spike counts
due to the experimental manipulation. To confirm the fidelity of the model, we
compared the distribution of all experimentally observed spike counts together with
the model fit (**Figure 3e**): the overlap between these distributions indicated that the
model performed well in capturing the overall structure of the data.

Having confirmed the fidelity of the model, we then compared the contrasts between
model coefficients associated with the pre- and post-drug administration periods for
each experimental group. These contrasts reflect the effect of the experimental
manipulation on spike count distribution. For each experimental group there are two
contrasts, one associated with the shape parameter $α$ and one with the scale
parameter $β$ of the distribution describing spike counts. Comparing contrasts and
highest density intervals (hdi, 94%) for the parameters across groups revealed a
specific effect in GlyT2-CNO mice on both $α$ and $β$ (**Figure 3f**). To gain a better intuition
of the implications of these results, we used the posterior samples of $α$ and $β$ to derived
samples for the mean and variance parameters of the inverse-Gamma distribution; the
mean and variance offer an alternative parametrisation that characterise more
intuitively the centre of mass and spread of spike counts across recordings for each
condition, before and after drug application (see Materials and Methods). Thus, to
compare results across condition, we computed the pre- minus post-drug mean and
variance contrasts for each experimental condition, and the difference in contrasts
between conditions, for each parameter. This analysis revealed that both mean and

variance increased after drug application in the GlyT2-CNO group compared to the
two control groups. (**Figure 3g**; mean hdi 0.05- 0.46, 0.05 - 0.49 and variance hdi -
0.01 – 0.31, -0.02 – 0.31 for GlyT2-CNO versus WT-CNO and Veh respectively). It
further revealed that in the WT-CNO group, CNO application alone has no detectable
effect on network spiking activity (mean hdi -0.17– 0.21 and variance hdi -0.02 – 0.03
for WT-CNO versus Veh). Together our analyses confirm that chemogenetic inhibition
of Golgi cells causes a global increase in spiking activity within the cerebellar cortex.
566

567 **Limited influence of GlyT2-positive cell downregulation on sensorimotor**
568 **representation**
569

Having established that chemogenetic downregulation of GlyT2-positive cells
measurably altered cerebellar activity, we investigated how the perturbation affected
sensorimotor representations of whisking. Golgi cell inhibition acts via the granule cell
layer, but changes in the input-output transformation of granule cells will have knock
on effects upon the activity of molecular interneurons and Purkinje cells. Golgi cells
control spike output of granule cells through various mechanisms, including changes
in gain, i.e., slope of the input-output function (Brickley et al., 1996; Chadderton et al.,
2004; Hamann et al., 2002; Mitchell & Silver, 2003). By reducing Golgi cell inhibition,
granule cells may become more sensitive to excitatory inputs, increasing their gain.
We looked for changes in the sensitivity of cerebellar activity to whisker movement by
comparing population tuning curves before and after CNO administration in GlyT2
mice. We observed no striking differences in the overall profile of these curves (**Figure**
4a). Post-CNO, population firing rates were marginally higher for some whisker
positions, but tuning curves were essentially unchanged. This result suggests that
chemogenetic downregulation of GlyT2-positive cells has no substantial effect on gain
or overall representation of whisker position at the population level. In principle,
altering Golgi inhibition could cause reorganisation of the granule cell population code
by altering the requirements for granule cell integration of excitatory inputs from
different mossy fibres (Marr, 1969). We therefore explored whether perturbed GlyT2-
positive cell activity causes substantial reorganisation of whisking representation at

the single cell level. To do this, we measured the entropy of individual tuning curves before and after CNO/vehicle administration. Entropy is a measure of how informative units are about, in this case, whisking position, with higher-entropy neurons being less informative. Following CNO administration, we observed no changes in cumulative entropy distribution in GlyT2-CNO recordings (**Figure 4b**). Interestingly, under control conditions (i.e. Veh and WT-CNO recordings), units tended to have higher entropy values after vehicle/CNO administration, indicating that tuning curves become less informative about whisking position over time. We further compared the distributions of entropy differences pre- and post-drug delivery (**Figure 4c**). Under control conditions, tuning curve entropies typically remained unchanged or increase over time (control density skewness = 2.80). CNO administration in GlyT2 mice causes additional reorganisation - some units decrease their entropy - leading to an even distribution of entropy changes after drug application (GlyT2-CNO density skewness = 0.21). The cumulative entropy distribution in GlyT2-CNO recordings therefore remains similar across time suggesting that the firing rate of some units becomes more sensitive to changes in whisking position following GlyT2-positive cell manipulation. These results also imply that recording conditions in the control condition are associated with small but progressive losses of representational fidelity. To confirm this proposal, we again measured the correlation between each of the first three population principal components (pc1-3) and whisker position for GlyT2-CNO and control recordings (**Figure 4d,e**). We found no evidence for large differences in correlation pre- and post-drug administration. However, there was a general trend for correlations to decrease after drug delivery, except for pc1 and pc2 (i.e. fast and slower representations of whisking) in the GlyT2-CNO condition, where correlations increased slightly. Our results suggest that the fidelity of sensorimotor representation is subject to a mild downward drift under control conditions in the present experimental setting, and that GlyT2-positive cell downregulation may slightly improve representation, offsetting this drift. But overall, we did not observe notable changes in the quality of the encoding of whisking behaviour by population activity. Together our results indicate that individual sensorimotor tuning of cerebellar neurons is only weakly influenced by chemogenetic perturbation of GlyT2-positive interneurons.

GlyT2-positive cell downregulation alters cerebellar dynamics and weakens coupling to whisking behaviour

We further explored the impact of GlyT2-positive cell perturbation by comparing trial
averaged peri-event time histograms (PETH) and whisking activity from pre- and post-
drug administration periods (see Materials and Methods). Because cerebellar activity
can directly influence whisker movement (Proville et al., 2014), reduced GlyT2-positive
cell inhibition could influence both cerebellar dynamics and produce measurable
changes in whisking behaviour. We first considered cerebellar activity: reductions in
Golgi cell inhibition could alter the temporal dynamics of population activity in several
ways, acting via tonic and/or phasic inhibition (Brickley et al., 1996; Chadderton et al.,
2004; Duguid et al., 2015), and/or affecting Golgi cell synchrony (van Welie et al.,
2016). We therefore compared the patterns of population activity at movement onset
before and after chemogenetic perturbation. Typically population activity peaks close
to the onset of individual whisking bouts (**Figure 1b,c**). Therefore, for each recording,
we computed the standard deviation of the distribution of peak unit firing times aligned
to whisking onset, pre- and post-drug delivery (**Figure 5a**; see Materials and Methods).
In the GlyT2-CNO mice, but not controls, pre- and post- drug standard deviations were
significantly different (**Figure 5b**), with significantly lower variation following CNO
administration (two-sided Wilcoxon signed-rank test, T = 12, p = 0.017). Thus, GlyT2-
positive cell downregulation increased the temporal alignment of peak neuronal
activity around whisking onset. These results suggest that decreased GlyT2-positive
cell inhibition has an impact on the pacing of neuronal responses around the onset of
whisking activity, and in particular narrows the time window within which units fire most
strongly.

Finally, we measured the effects of GlyT2-positive cell manipulation on whisking
velocity at the onset of movement bouts (see Materials and Methods). For each
recording, we used a linear model to approximate the slope of the average whisking
bouts during its protraction phase (**Figure 5c**); the coefficient for each linear fit was

used as a read out for velocity during the movement initiation. Interestingly, unlike

control recordings, GlyT2-CNO mice showed changes in their whisking behaviour

following drug administration. However, the sign of these changes was not consistent

across animals, occurring in both increasing and decreasing directions (**Figure 5d**).

Individual onsets of voluntary whisking thus became faster or slower after drug

administration in GlyT2-CNO mice. The overall distribution of changes in whisker

onset velocity ('slope contrast') was therefore significantly broader in the GlyT2-CNO

mice compared to control (**Figure 5e**). Thus, GlyT2-positive cell downregulation

produces measurable but divergent effects on local cerebellar dynamics and whole

animal whisking behaviour. Local cerebellar dynamics were consistently compressed,

leading to a more tightly locked population activity, whereas whisking velocity was

either increased or reduced. These results suggest a breakdown in the link between

cerebellar dynamics and behaviour following GlyT2-positive cell downregulation. We

confirmed this by plotting changes in population activity and whisking velocity at

movement onset for individual recordings (**Figure 5f**). In control recordings,

differences in network activity and whisker movement before and after CNO/vehicle

delivery are small in magnitude but importantly, are well correlated (r^2 = 0.52). In

GlyT2-CNO recordings, changes in cerebellar dynamics and movement become

decoupled from one another, leading to break down in this correlation (r^2 = 0.15).

Together our results indicate GlyT2-positive cell inhibition regulates the temporal

patterning of population activity and that disruption of this patterning at the local level

can have diverse consequences for behaviour.

DISCUSSION

Granule cell layer inhibitory interneurons are proposed to play important roles in

cerebellar transformation (Albus, 1971; D'Angelo et al., 2013; Marr, 1969; Palacios et

al., 2021), but we lack detailed information about their influence on circuit activity. Here

we have tested the influence of Golgi and Lugaro cells on cerebellar population activity.

Our GlyT2-positive cell perturbations caused mild changes in neural population activity

without profoundly disrupting sensorimotor representations in the cerebellar cortex.

Such an alteration is likely within normal physiological operating limits, and the
changes we observe can thus be presumed to be ethologically relevant.

Cerebellar cortical neurons heterogeneously encode whisking behaviour

Lateral cerebellar neurons accurately represent whisking behaviour. Individual units
exhibit different tuning profiles to whisking position (**Figure 1d**), consistent with
previous work (Bosman et al., 2010; Brown & Raman, 2018; Chen et al., 2016, 2017),
enabling multidimensional representations of movement kinematics (Gurnani & Silver,
2021; Markanday et al., 2023). Our units are unlikely to be granule cells (Beau et al.,
2024), but rather inhibitory interneurons and Purkinje cells. However, as observed in
small virtual granule cell populations (Chen et al., 2017), ensemble activity displayed
a near-monotonic dependence upon whisker set point, with increasing activity in the
direction of both protraction and retraction (**Figure 1e**) in line with previous findings
(Brown & Raman, 2018; Chen et al., 2016, 2017; Lanore et al., 2021; Proville et al.,
2014).

The cerebellar cortex uses a distributed code to represent whisking dynamics

We used principal component analysis to describe population activity with three
independent variables. The first three pcs capture only a moderate amount of the total
variance in the population, suggesting that cerebellar activity reflects additional
behaviourally relevant variables (Chabrol et al., 2015; Ishikawa et al., 2015; Sobel et
al., 1998; Wagner & Luo, 2020). Nevertheless, neuronal dynamics had structure that
matched whisking dynamics (**Figure 1g**). Notably, pc1 dynamics consistently led (i.e.
anticipated) whisking behaviour by ~40ms whereas pc2 dynamics lagged behaviour
by a few hundred milliseconds (**Figure 2**). The former is consistent with a forward
model of planned/predicted movement, while the latter resembles a feedback signal.
These results support a role for the cerebellum in controlling future behaviour,
incorporating recent sensory feedback and further behaviourally relevant variables, in

line with theoretical treatments of the cerebellum (Palacios et al., 2024). Whisker-
related information, especially in pc1, was well distributed across units. Thus, most
units contribute to some extent to its encoding.

Decreasing GlyT2-positive cell inhibition does not saturate population activity

Next, we examined how altering GlyT2-positive cell activity affected neuronal
representations in the cerebellar cortex. This was achieved by activating inhibitory
receptors (hM4Di) on Golgi and Lugaro cells, through application of the agonist CNO
directly at the surface of the recording site. This approach afforded within-minute
temporal precision to the manipulation. hM4Di receptors inhibit neuronal activity via
moderate hyperpolarisation and reduced presynaptic release of neurotransmitters
(Roth, 2016). Topical application of CNO onto the cerebellar cortex produced an
increased spread of the population spike count distribution, with higher spike counts
becoming more likely across recordings (**Figure 3c,d**). However, manipulation of
GlyT2-positive cell inhibition did not lead to excitatory saturation of network dynamics,
i.e., a drastic and indiscriminate increase in neural activity. The absence of strong
saturation of network activity may be because granule cell excitability is partly
controlled via non-vesicular sources of GABA (Lee et al., 2010; Rossi et al., 2003; Wall
& Usowicz, 1997). In addition, Lugaro cells target inhibitory neurons, here potentially
producing compensatory upregulation of molecular layer interneuron activity (Eyre &
Nusser, 2016). The tuning of individual neurons was also largely unchanged (**Figure
4a**). Theoretical and *in vitro* evidence indicates that the gain of granule cell input/output
transformation may be under the control of local inhibitory processes. As such,
reduced inhibition is predicted to increase cortical sensitivity to incoming synaptic
input. Following our perturbation, we saw little evidence of increased sensitivity to
features of whisker movement (**Figure 4a**), although we did observe a small fraction
of neurons that exhibited reduced entropy following GlyT2-positive cell inhibition
(**Figure 4c**). These neurons may represent a subpopulation that receive subthreshold
whisker-related synaptic input. Here, small decreases in synaptic inhibition may result
in the generation of tuned action potential output (Chadderton et al., 2004; Chen et al.,
2017). However, our results suggest that dynamic GlyT2-positive cell inhibition may

not be necessary for maintaining excitation at the level required by the network to
perform its computations (Albus, 1971; Marr, 1969). Manipulation of GlyT2-positive
cell inhibition instead produces more fine-grained effects on cerebellar cortical
dynamics.

746 Decreased GlyT2-positive cell inhibition increases the temporal alignment of
747 neuronal activation during movement onset

Decreased GlyT2-positive cell inhibition was associated with increased temporal
alignment or synchronisation within neuronal populations around whisking onset. A
familiar downstream consequence of feedforward inhibition in many brain regions, and
at the molecular layer interneuron-Purkinje cell synapse, is increased temporal
precision of neuronal dynamics, as inhibition normally trails excitation and sharpens
the window of input integration (Gabernet et al., 2005; Mittmann et al., 2005; Wehr &
Zador, 2003). Here, notably, Golgi cells play the opposite role, temporally dispersing
or desynchronising neural activity (Vervaeke et al., 2010). Although Golgi cells provide
feedforward inhibition to the granule cell layer, evoked inhibition has been previously
shown to precede or be concomitant with mossy fibre excitation, under which
situations it may reduce the temporal precision of early granule cell responses (Duguid
et al., 2015), and decorrelate multisensory representations (Fleming et al., 2024). Our
observations are consistent with such a mode of operation, as decreasing GlyT2-
positive cell inhibition leads to an increased temporal alignment of the overall network
activity during movement initiation (Vervaeke et al., 2010). Changes in Golgi cell
activity can modulate the amplitude of tonic inhibitory conductances (Crowley et al.,
2009). Reduced Golgi cell activity may therefore directly reduce the tonic inhibitory
conductance, making granule cells more excitable and more likely to fire rapidly to a
burst of incoming mossy fibre synaptic inputs. Such control by Golgi cells of the timing
of network activity can support the specific cerebellar role in integrating information
and learning contexts (Fleming et al., 2024). Indeed, distinct temporal patterning of
granule cell responses to mossy fibre input is proposed to be important for how the
cerebellar network integrates and distinguishes parallel sensory and motor pathways
(Chabrol et al., 2015).

Decreasing GlyT2-positive cell inhibition decreases the stability of whisking behaviour

Finally, we found that reduced GlyT2-positive cell inhibition led to increased variability in whisking velocity at movement onset. The specific change in movement velocity varied amongst recordings: in some cases, the velocity of whisking increased while in other cases it decreased. We suggest that location of the site of perturbation, varying from mouse to mouse, is likely to have influenced the valence of the change in whisking behaviour (Apps et al., 2018). Importantly, the link between cerebellar cortical dynamics and whisker position is decoupled after GlyT2-positive cell perturbation (**Figure 5f**). This result highlights the importance of an intact cerebellum for precise whisking control, which is subject to the coordination with other motor and sensory systems (Deschenes et al., 2012; Kurnikova et al., 2017; McElvain et al., 2018; Romano et al., 2020). Increased variability in motor initiation may be linked with the possible contribution of Golgi cell inhibition to representations of multiple streams of information and behaviours – bound together at the level of the cerebellar cortex – improving accuracy as specific behaviours are performed.

Conclusions

Granule cell layer inhibition itself is under the control of many mechanisms, including feedback loops encompassing the cortex (Nietz et al., 2017; Witter et al., 2016), the cerebellar nuclei (Ankri et al., 2015), as well as neuromodulatory systems (Dieudonne & Dumoulin, 2000; Fleming et al., 2024; Fore et al., 2020). Thus, granule cell layer inhibitory interneurons sit in a key position to flexibly control cerebellar computations, based on the dynamic behavioural context, which may be required to focus on different aspects of behaviour at different times (Palacios et al., 2021). To fully understand the pervasive involvement of the cerebellum in behavioural control, it is critical to elucidate how different mechanisms contribute to fine tune Golgi and Lugaro cell activity in a context-dependent manner.

Our work shows how altering GlyT2-positive cell inhibition changes the response of
the network to its input. By controlling the timing of network activity, GlyT2-positive
interneurons can support cerebellar roles in integrating information and learning
contexts (Fleming et al., 2024). Distinct temporal patterning of granule cell responses
to mossy fibre input may enable the cerebellar cortex to integrate and distinguish
parallel sensory and motor pathways, which in turn may be relevant for pattern
separation (Chabrol et al., 2015). Cerebellar computations have also been described
as state estimation processes (Doya, 1999; Miall et al., 1993; Palacios et al., 2024;
Paulin, 1993) in which granule cell layer inhibition acts as a mechanism to fine tune
the integration of information from different mossy fiber pathways (Palacios et al.,
2021). In this scheme, temporal changes in cerebellar dynamics may result from tuned
integration of whisking-related information with other, temporally heterogeneous
streams of information.

AUTHOR CONTRIBUTIONS

ERP, CH and PC designed the experiments. All experiments were performed by ERP.
Data was analysed and interpreted by ERP, CH and PC. The authors declare no
competing interests.

REFERENCES

Albus, J. S. (1971). A theory of cerebellar function. *Math Biosci, 10*(1-2), 25-61.

Ankri, L., Husson, Z., Pietrajtis, K., Proville, R., Lena, C., Yarom, Y., Dieudonne, S., & Uusisaari, M. Y. (2015). A novel inhibitory nucleo-cortical circuit controls cerebellar Golgi cell activity. *eLife, 4*. https://doi.org/10.7554/eLife.06262

Apps, R., Hawkes, R., Aoki, S., Bengtsson, F., Brown, A. M., Chen, G., Ebner, T. J., Isope, P., Jorntell, H., Lackey, E. P., Lawrenson, C., Lumb, B., Schonewille, M., Sillitoe, R. V., Spaeth, L., Sugihara, I., Valera, A., Voogd, J., Wylie, D. R., & Ruigrok, T. J. H. (2018). Cerebellar Modules and Their Role as Operational Cerebellar Processing Units: A Consensus paper [corrected]. *Cerebellum, 17*(5), 654-682. https://doi.org/10.1007/s12311-018-0952-3

Baumann, O., & Mattingley, J. B. (2022). Cerebellum and Emotion Processing. *Adv Exp Med Biol, 1378*, 25-39. https://doi.org/10.1007/978-3-030-99550-8_3

Beau, M., Herzfeld, D. J., Naveros, F., Hemelt, M. E., D'Agostino, F., Oostland, M., Sanchez-Lopez, A., Chung, Y. Y., Michael, M., Kyranakis, S., Stabb, H. N., Martinez Lopera, M. G., Lajko, A., Zedler, M., Ohmae, S., Hall, N. J., Clark, B. A., Cohen, D., Lisberger, S. G., . . . Medina, J. F. (2024). A deep-learning strategy to identify cell types across species from high-density extracellular recordings. *bioRxiv*. https://doi.org/10.1101/2024.01.30.577845

Bosman, L. W., Koekkoek, S. K., Shapiro, J., Rijken, B. F., Zandstra, F., van der Ende, B., Owens, C. B., Potters, J. W., de Gruijl, J. R., Ruigrok, T. J., & De Zeeuw, C. I. (2010). Encoding of whisker input by cerebellar Purkinje cells. *J Physiol, 588*(19), 3757-3783. https://doi.org/10.1113/jphysiol.2010.195180

Brickley, S. G., Cull-Candy, S. G., & Farrant, M. (1996). Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA receptors. *J Physiol 497*(3), 753-759. https://doi.org/10.1113/jphysiol.1996.sp021806

Brown, S. T., & Raman, I. M. (2018). Sensorimotor Integration and Amplification of Reflexive Whisking by Well-Timed Spiking in the Cerebellar Corticonuclear Circuit. *Neuron, 99*(3), 564-575 e562. https://doi.org/10.1016/j.neuron.2018.06.028

Chabrol, F. P., Arenz, A., Wiechert, M. T., Margrie, T. W., & DiGregorio, D. A. (2015). Synaptic diversity enables temporal coding of coincident multisensory inputs in single neurons. *Nat Neurosci, 18*(5), 718-727. https://doi.org/10.1038/nn.3974

865 Chadderton, P., Margrie, T. W., & Hausser, M. (2004). Integration of quanta in
866   cerebellar granule cells during sensory processing. *Nature*, *428*(6985), 856-
867   860. https://doi.org/10.1038/nature02442
868
869 Chen, S., Augustine, G. J., & Chadderton, P. (2016). The cerebellum linearly encodes
870   whisker position during voluntary movement. *eLife*, *5*, e10509.
871   https://doi.org/10.7554/eLife.10509
872
873 Chen, S., Augustine, G. J., & Chadderton, P. (2017). Serial processing of kinematic
874   signals by cerebellar circuitry during voluntary whisking. *Nat Commun*, *8*(1),
875   232. https://doi.org/10.1038/s41467-017-00312-1
876
877 Crowley, J. J., Fioravante, D., & Regehr, W. G. (2009). Dynamics of fast and slow
878   inhibition from cerebellar golgi cells allow flexible control of synaptic integration.
879   *Neuron*, *63*(6), 843-853. https://doi.org/10.1016/j.neuron.2009.09.004
880
881 D'Angelo, E., Solinas, S., Mapelli, J., Gandolfi, D., Mapelli, L., & Prestori, F. (2013).
882   The cerebellar Golgi cell and spatiotemporal organization of granular layer
883   activity. *Front Neural Circuits*, *7*, 93. https://doi.org/10.3389/fncir.2013.00093
884
885 Deschenes, M., Moore, J., & Kleinfeld, D. (2012). Sniffing and whisking in rodents.
886   *Curr Opin Neurobiol*, *22*(2), 243-250.
887   https://doi.org/10.1016/j.conb.2011.11.013
888
889 Dieudonne, S. (1998). Submillisecond kinetics and low efficacy of parallel fibre-Golgi
890   cell synaptic currents in the rat cerebellum. *J Physiol*, *510*(3), 845-866.
891   https://doi.org/10.1111/j.1469-7793.1998.845bj.x
892
893 Dieudonne, S., & Dumoulin, A. (2000). Serotonin-driven long-range inhibitory
894   connections in the cerebellar cortex. *J Neurosci*, *20*(5), 1837-1848.
895   https://doi.org/10.1523/JNEUROSCI.20-05-01837.2000
896
897 Duguid, I., Branco, T., Chadderton, P., Arlt, C., Powell, K., & Hausser, M. (2015).
898   Control of cerebellar granule cell output by sensory-evoked Golgi cell
899   inhibition. *Proc Natl Acad Sci U S A*, *112*(42), 13099-13104.
900   https://doi.org/10.1073/pnas.1510249112
901
902 Duguid, I., Branco, T., London, M., Chadderton, P., & Hausser, M. (2012). Tonic
903   inhibition enhances fidelity of sensory information transmission in the
904   cerebellar cortex. *J Neurosci*, *32*(32), 11132-11143.
905   https://doi.org/10.1523/JNEUROSCI.0460-12.2012
906
907 Eccles, J. C., Llinas, R., & Sasaki, K. (1966). The mossy fibre-granule cell relay of the
908   cerebellum and its inhibitory control by Golgi cells. *Exp Brain Res*, *1*(1), 82-101.
909   https://doi.org/10.1007/BF00235211
910

911 Fleming, E., & Hull, C. (2019). Serotonin regulates dynamics of cerebellar granule cell
912   activity by modulating tonic inhibition. *J Neurophysiol*, *121*(1), 105-114.
913   https://doi.org/10.1152/jn.00492.2018
914
915 Fleming, E. A., Field, G. D., Tadross, M. R., & Hull, C. (2024). Local synaptic inhibition
916   mediates cerebellar granule cell pattern separation and enables learned
917   sensorimotor associations. *Nat Neurosci*, *27*(4), 689-701.
918   https://doi.org/10.1038/s41593-023-01565-4
919
920 Fore, T. R., Taylor, B. N., Brunel, N., & Hull, C. (2020). Acetylcholine Modulates
921   Cerebellar Granule Cell Spiking by Regulating the Balance of Synaptic
922   Excitation and Inhibition. *J Neurosci*, *40*(14), 2882-2894.
923   https://doi.org/10.1523/JNEUROSCI.2148-19.2020
924
925 Gabernet, L., Jadhav, S. P., Feldman, D. E., Carandini, M., & Scanziani, M. (2005).
926   Somatosensory integration controlled by dynamic thalamocortical feed-
927   forward inhibition. *Neuron*, *48*(2), 315-327.
928   https://doi.org/10.1016/j.neuron.2005.09.022
929
930 Hamann, M., Rossi, D. J., & Attwell, D. (2002). Tonic and spillover inhibition of granule
931   cells control information flow through cerebellar cortex. *Neuron*, *33*(4), 625-633.
932   https://doi.org/10.1016/s0896-6273(02)00593-7
933
934 Hill, D. N., Curtis, J. C., Moore, J. D., & Kleinfeld, D. (2011). Primary motor cortex
935   reports efferent control of vibrissa motion on multiple timescales. *Neuron*, *72*(2),
936   344-356. https://doi.org/10.1016/j.neuron.2011.09.020
937
938 Ishikawa, T., Shimuta, M., & Hausser, M. (2015). Multimodal sensory integration in
939   single cerebellar granule cells in vivo. *eLife*, *4*.
940   https://doi.org/10.7554/eLife.12916
941
942 Kakizaki, T., Sakagami, H., Sakimura, K., & Yanagawa, Y. (2017). A glycine transporter
943   2-Cre knock-in mouse line for glycinergic neuron-specific gene manipulation.
944   *IBRO Rep*, *3*, 9-16. https://doi.org/10.1016/j.ibror.2017.07.002
945
Kelly, E., Meng, F., Fujita, H., Morgado, F., Kazemi, Y., Rice, L. C., Ren, C.,
Escamilla, C. O., Gibson, J. M., Sajadi, S., Pendry, R. J., Tan, T., Ellegood, J.,
Basson, M. A., Blakely, R. D., Dindot, S. V., Golzio, C., Hahn, M. K., Katsanis,
N., . . . Tsai, P. T. (2020). Regulation of autism-relevant behaviors by
cerebellar-prefrontal cortical circuits. *Nat Neurosci*, *23*(9), 1102-1110.
https://doi.org/10.1038/s41593-020-0665-z
952
953 Kurnikova, A., Moore, J. D., Liao, S. M., Deschenes, M., & Kleinfeld, D. (2017).
954   Coordination of Orofacial Motor Actions into Exploratory Behavior by Rat. *Curr
955   Biol*, *27*(5), 688-696. https://doi.org/10.1016/j.cub.2017.01.013
956

957 Lanore, F., Cayco-Gajic, N. A., Gurnani, H., Coyle, D., & Silver, R. A. (2021). Cerebellar
958   granule cell axons support high-dimensional representations. *Nat Neurosci*,
959   *24*(8), 1142-1150. https://doi.org/10.1038/s41593-021-00873-x
960

961 Lee, S., Yoon, B. E., Berglund, K., Oh, S. J., Park, H., Shin, H. S., Augustine, G. J., &
962   Lee, C. J. (2010). Channel-mediated tonic GABA release from glia. *Science*,
963   *330*(6005), 790-796. https://doi.org/10.1126/science.1184334
964

965 Marr, D. (1969). A theory of cerebellar cortex. *J Physiol*, *202*(2), 437-470.
966   https://doi.org/10.1113/jphysiol.1969.sp008820
967

968 McElvain, L. E., Friedman, B., Karten, H. J., Svoboda, K., Wang, F., Deschenes, M.,
969   & Kleinfeld, D. (2018). Circuits in the rodent brainstem that control whisking in
970   concert with other orofacial motor actions. *Neuroscience*, *368*, 152-170.
971   https://doi.org/10.1016/j.neuroscience.2017.08.034
972

973 Mitchell, S. J., & Silver, R. A. (2003). Shunting inhibition modulates neuronal gain
974   during synaptic excitation. *Neuron*, *38*(3), 433-445.
975   https://doi.org/10.1016/s0896-6273(03)00200-9
976

977 Mittmann, W., Koch, U., & Hausser, M. (2005). Feed-forward inhibition shapes the
978   spike output of cerebellar Purkinje cells. *J Physiol*, *563*(2), 369-378.
979   https://doi.org/10.1113/jphysiol.2004.075028
980

981 Nietz, A. K., Vaden, J. H., Coddington, L. T., Overstreet-Wadiche, L., & Wadiche, J. I.
982   (2017). Non-synaptic signaling from cerebellar climbing fibers modulates Golgi
983   cell activity. *eLife*, *6*. https://doi.org/10.7554/eLife.29215
984

985 Palacios, E. R., Houghton, C., & Chadderton, P. (2021). Accounting for uncertainty:
986   inhibition for neural inference in the cerebellum. *Proc Biol Sci*, *288*(1947),
987   20210276. https://doi.org/10.1098/rspb.2021.0276
988

Chabrol, F. P., Arenz, A., Wiechert, M. T., Margrie, T. W., & DiGregorio, D. A. (2015).
Synaptic diversity enables temporal coding of coincident multisensory inputs in
single neurons. *Nat Neurosci*, *18*(5), 718-727. https://doi.org/10.1038/nn.3974
Dieudonne, S. (1998). Submillisecond kinetics and low efficacy of parallel fibre-Golgi
cell synaptic currents in the rat cerebellum. *J Physiol*, *510 (Pt 3)*(Pt 3), 845-
866. https://doi.org/10.1111/j.1469-7793.1998.845bj.x
Doya, K. (1999). What are the computations of the cerebellum, the basal ganglia and
the cerebral cortex? *Neural Netw*, *12*(7-8), 961-974.
https://doi.org/10.1016/s0893-6080(99)00046-5
Eccles, J. C., Llinas, R., & Sasaki, K. (1966). The mossy fibre-granule cell relay of the
cerebellum and its inhibitory control by Golgi cells. *Exp Brain Res*, *1*(1), 82-101.
https://doi.org/10.1007/BF00235211
Eyre, M. D., & Nusser, Z. (2016). Only a Minority of the Inhibitory Inputs to Cerebellar
Golgi Cells Originates from Local GABAergic Cells. *eNeuro*, *3*(2).
https://doi.org/10.1523/ENEURO.0055-16.2016

Fleming, E. A., Field, G. D., Tadross, M. R., & Hull, C. (2024). Local synaptic inhibition mediates cerebellar granule cell pattern separation and enables learned sensorimotor associations. *Nat Neurosci*, *27*(4), 689-701. https://doi.org/10.1038/s41593-023-01565-4

Gurnani, H., & Silver, R. A. (2021). Multidimensional population activity in an electrically coupled inhibitory circuit in the cerebellar cortex. *Neuron*, *109*(10), 1739-1753 e1738. https://doi.org/10.1016/j.neuron.2021.03.027

Kanichay, R. T., & Silver, R. A. (2008). Synaptic and cellular properties of the feedforward inhibitory circuit within the input layer of the cerebellar cortex. *J Neurosci*, *28*(36), 8955-8967. https://doi.org/10.1523/JNEUROSCI.5469-07.2008

Markanday, A., Hong, S., Inoue, J., De Schutter, E., & Thier, P. (2023). Multidimensional cerebellar computations for flexible kinematic control of movements. *Nat Commun*, *14*(1), 2548. https://doi.org/10.1038/s41467-023-37981-0

Miall, R. C., Weir, D. J., Wolpert, D. M., & Stein, J. F. (1993). Is the cerebellum a smith predictor? *J Mot Behav*, *25*(3), 203-216. https://doi.org/10.1080/00222895.1993.9942050

Palacios, E. R., Chadderton, P., Friston, K., & Houghton, C. (2024). Cerebellar state estimation enables resilient coupling across behavioural domains. *Sci Rep*, *14*(1), 6641. https://doi.org/10.1038/s41598-024-56811-x

Palacios, E. R., Houghton, C., & Chadderton, P. (2021). Accounting for uncertainty: inhibition for neural inference in the cerebellum. *Proc Biol Sci*, *288*(1947), 20210276. https://doi.org/10.1098/rspb.2021.0276

Paulin, M. G. (1993). The role of the cerebellum in motor control and perception. *Brain Behav Evol*, *41*(1), 39-50. https://doi.org/10.1159/000113822

Prestori, F., Mapelli, L., & D'Angelo, E. (2019). Diverse Neuron Properties and Complex Network Dynamics in the Cerebellar Cortical Inhibitory Circuit. *Front Mol Neurosci*, *12*, 267. https://doi.org/10.3389/fnmol.2019.00267

Simat, M., Parpan, F., & Fritschy, J. M. (2007). Heterogeneity of glycinergic and gabaergic interneurons in the granule cell layer of mouse cerebellum. *J Comp Neurol*, *500*(1), 71-83. https://doi.org/10.1002/cne.21142

Vervaeke, K., Lorincz, A., Gleeson, P., Farinella, M., Nusser, Z., & Silver, R. A. (2010). Rapid desynchronization of an electrically coupled interneuron network with sparse excitatory synaptic input. *Neuron*, *67*(3), 435-451. https://doi.org/10.1016/j.neuron.2010.06.028

Proville, R. D., Spolidoro, M., Guyon, N., Dugue, G. P., Selimi, F., Isope, P., Popa, D., & Lena, C. (2014). Cerebellum involvement in cortical sensorimotor circuits for the control of voluntary movements. *Nat Neurosci*, *17*(9), 1233-1239. https://doi.org/10.1038/nn.3773

Romano, V., Reddington, A. L., Cazzanelli, S., Mazza, R., Ma, Y., Strydis, C., Negrello, M., Bosman, L. W. J., & De Zeeuw, C. I. (2020). Functional Convergence of Autonomic and Sensorimotor Processing in the Lateral Cerebellum. *Cell Rep*, *32*(1), 107867. https://doi.org/10.1016/j.celrep.2020.107867

Rossi, D. J., Hamann, M., & Attwell, D. (2003). Multiple modes of GABAergic inhibition of rat cerebellar granule cells. *J Physiol*, *548*(1), 97-110. https://doi.org/10.1113/jphysiol.2002.036459

Roth, B. L. (2016). DREADDs for Neuroscientists. *Neuron*, *89*(4), 683-694. https://doi.org/10.1016/j.neuron.2016.01.040

Silveri, M. C. (2021). Contribution of the Cerebellum and the Basal Ganglia to Language Production: Speech, Word Fluency, and Sentence Construction-Evidence from Pathology. *Cerebellum*, *20*(2), 282-294. https://doi.org/10.1007/s12311-020-01207-6

Sobel, N., Prabhakaran, V., Hartley, C. A., Desmond, J. E., Zhao, Z., Glover, G. H., Gabrieli, J. D., & Sullivan, E. V. (1998). Odorant-induced and sniff-induced activation in the cerebellum of the human. *J Neurosci*, *18*(21), 8990-9001. https://doi.org/10.1523/JNEUROSCI.18-21-08990.1998

Stachniak, T. J., Ghosh, A., & Sternson, S. M. (2014). Chemogenetic synaptic silencing of neural circuits localizes a hypothalamus-->midbrain pathway for feeding behavior. *Neuron*, *82*(4), 797-808. https://doi.org/10.1016/j.neuron.2014.04.008

van Welie, I., Roth, A., Ho, S. S., Komai, S., & Hausser, M. (2016). Conditional Spike Transmission Mediated by Electrical Coupling Ensures Millisecond Precision-Correlated Activity among Interneurons In Vivo. *Neuron*, *90*(4), 810-823. https://doi.org/10.1016/j.neuron.2016.04.013

Wagner, M. J., & Luo, L. (2020). Neocortex-Cerebellum Circuits for Cognitive Processing. *Trends Neurosci*, *43*(1), 42-54. https://doi.org/10.1016/j.tins.2019.11.002

Wall, M. J., & Usowicz, M. M. (1997). Development of action potential-dependent and independent spontaneous GABAA receptor-mediated currents in granule cells of postnatal rat cerebellum. *Eur J Neurosci*, *9*(3), 533-548. https://doi.org/10.1111/j.1460-9568.1997.tb01630.x

Wehr, M., & Zador, A. M. (2003). Balanced inhibition underlies tuning and sharpens spike timing in auditory cortex. *Nature*, *426*(6965), 442-446. https://doi.org/10.1038/nature02116

Witter, L., Rudolph, S., Pressler, R. T., Lahlaf, S. I., & Regehr, W. G. (2016). Purkinje Cell Collaterals Enable Output Signals from the Cerebellar Cortex to Feed Back to Purkinje Cells and Interneurons. *Neuron*, *91*(2), 312-319. https://doi.org/10.1016/j.neuron.2016.05.037

1096   Zhai, P., Romano, V., Soggia, G., Bauer, S., van Wingerden, N., Jacobs, T., van der
1097   Horst, A., White, J. J., Mazza, R., & De Zeeuw, C. I. (2024). Whisker kinematics
1098   in the cerebellum. *J Physiol, 602*(1), 153-181. https://doi.org/10.1113/JP284064

1099

1100

1101

FIGURE LEGENDS

Figure 1 Single unit and population activity in Crus 1 during voluntary whisking

a. Monitoring whisker movement and neuronal population activity in lateral cerebellar
cortex using high-speed videography and NeuroPixels recording in head-fixed mice.
Left: Whiskers were labelled and reconstructed using DeepLabCut to recover
movement trajectories. Image credit: Elisabeth Meyer. Right: Brightfield image of
coronal cerebellar section superimposed fluorescent image tract left by NeuroPixels
probe coated with DiI localised to lobule Crus 1.

b. 30 second segment of simultaneous whisking (top) and population activity (bottom;
n = 88 units) from a single recording session.

c. Trial-averaged whisker position and neuronal activity from the same recording,
aligned to onset of whisking bout.

d. Tuning curves of four individual cerebellar units from the same recording. Each unit
displays a different relationship between firing rate and whisker position. Blue lines
indicate fraction of time spent at different whisker positions; empty grey bars show
tuning curves for shuffled data. Left inset: colour-coded centre of mass of each unit on
NeuroPixels probe.

e. Average tuning curve for all recorded units, from all recordings (black line; n = 508,
N = 12). Population firing rate increases monotonically from whisker resting point
(white triangle) in both protractive and retractive directions. Blue line indicates fraction
of time spent at different whisker positions; shading represents standard deviation;
white triangle indicates whisker resting point; grey line shows tuning for shuffled data.

f. Average tuning curves for all units clustered in eight groups. Units were clustered
using k-mean algorithm based on individual unit tuning curves. Heterogeneity in tuning
curves enables a continuous encoding of whisking position.

g. Single trial of whisker movement (top) with corresponding projections in 3D principal
component space for average (bottom left) and single trial (bottom right) population
activity. Population activity in this space has structure that reflects whisking dynamics,
both on average and at a single trial level.

h. Recovered movement trajectory using cerebellar population activity. Reconstruction
of the whisking setpoint using a linear combination of the first three principal
components computed from neuronal populations. Whisker setpoint (brown line)
during two trials, together with predictions (red) and the highest density interval (hdi,
red shading).

Figure 2 Population representations of whisker movement at different timescales

a. Trial-averaged whisking position (top) and projected population activity (bottom) for one recording.

b. Cross-correlations between pc1-3 and whisking activity for the same recording; vertical lines indicate times of peak correlation.

c. Cumulative variance explained by pc1-3 across all recordings (N = 12). Only a modest amount of variability in neuronal activity is accounted for by first three principal components.

d. Distribution of unit loadings (n = 508) for pc1-3. Kurtosis measurement, to quantify the number of outliers in each distribution, reveals fewer outliers for pc1 than what would be expected if the data were normally distributed (excess kurtosis -2.76). This indicates that information contained in pc1, which best reflects whisking activity, tends to be distributed across neurons. The loading distributions for pc2 and pc3 have respectively a similar or higher number of outliers compared to normally distributed data (excess kurtosis -0.43 and 4.97, respectively), indicating that information in pc2 and pc3 is increasingly concentrated in fewer units.

e. Time of peak correlation for pc1-3 across all recordings. The distribution for pc1 is tightly concentrated around ~40 ms, meaning that information captured by pc1 tends to anticipate whisking activity with high temporal precision (one sample t test, t = 4.71, p = 0.0006). The information contained in pc2 and pc3 shows the opposite trend, with more variability (pc2: mean -194 ms, t = -4.44, p = 0.0009; pc3: mean -92 ms, t = -1.44, p = 0.17), suggesting that these components might reflect different aspects of behaviour.

f. Mean decrease in unexplained variance (unexp. var.) across recordings with increasing numbers of principal components (#pcs); standard deviation shaded in grey.

Figure 3 Chemogenetic inhibition of GlyT2-positive cells increases neuronal activity in cerebellar cortex

a. Left: targeted expression of hM4Di receptors was achieved via injection of AAV-DIO-hM4Di-mCherry into the lateral cerebellar cortex of GlyT2-Cre mice, selectively expressing Cre-recombinase in GlyT2-positive cells in this brain region. Right: NeuroPixels probes were targeted to site of viral injection. During electrophysiological recordings, the exogenous drug, clozapine-N-oxide (CNO) was topically applied to activate hM4Di receptors.

b. Left: GlyT2-Cre mouse brain selectively expressing hM4Di in GlyT2-positive cells of the lateral cerebellar cortex. Right: tract left by NeuroPixels probe coated with DiI, showing co-localisation of the site of recording and hM4Di expression.

c. Changes in population spike count before and after CNO/vehicle delivery (at 0 min) normalised by count at -5 min, for three experiment conditions. Pink, GlyT2-Cre mice and CNO (N = 19); Blue, C57BL6 mice and CNO (N = 5); Green, GlyT2-Cre/C57BL6 mice and saline vehicle only (N = 9).

d. Box plots showing pooled data for each experimental condition.

e. Multilevel modelling approach to assess effect of drug delivery on cerebellar population activity. Population spike count distributions for each condition (top) were modelled using an inverse-Gamma distribution, described by shape parameter α and scale parameter β. Each parameter was modelled as a linear combination of different coefficients, including two θ_{cond}, one for α and one for β, which captured the specific effect of the experimental manipulation on total spike counts. Comparing the empirical distribution with the distribution of posterior predictive checks (samples from the model after fitting) shows that the model captures the overall structure of the data (bottom).

f. 94% highest density intervals (hdi) of the contrasts between post- and pre-drop posterior samples for θ_{cond} for α and β. The contrasts highlight a specific effect of GlyT2-positive cell manipulation (GlyT2 + CNO) on both α and β parameters.

g. Contrast difference (mean and variance) between post- and pre-drop of the inverse-Gamma distribution fitted to total spike counts. Bars indicate the 94% hdi of the posterior differences. Left, contrast difference between GlyT2 + CNO condition and each control condition. Right, contrast difference between the two control conditions. GlyT2-positive cell manipulation produces positive contrast differences versus both control conditions.

Figure 4 Weak influence of GlyT2-positive cell perturbation on cerebellar sensorimotor representations

a. Average tuning curves for GlyT2-CNO recordings before (grey) and after (red) topical administration of the drug.

b. Entropy cumulative density functions for all units recorded in the GlyT2-CNO (top) and control (bottom) conditions. Under control conditions, but not GlyT2-CNO, unit entropy tends to increase after drug/vehicle delivery (post-drop), suggesting that tuning curves become less informative about whisking position over time.

c. Probability mass function of unit entropy differences before and after CNO/vehicle administration ('post-drop' minus 'pre-drop' entropy). Under control conditions, tuning curve entropy tends to remain stable or increase over time (density skewness = 2.79). In the GlyT2-CNO recordings, a fraction units exhibit decreased entropy after drug application (density skewness = 0.21), suggesting that some units become more sensitive to changes in whisking position following GlyT2-positive cell inhibition.

d. Comparison of peak pre- and post-drop correlation values between principal components 1-3 (pc1-3) and whisker position for control recordings.

e. Comparison of peak pre- and post-drop correlation values between principal components 1-3 (pc1-3) and whisker position for GlyT2-CNO recordings.

Figure 5 GlyT2-positive cell inhibition reduces temporal variability in neuronal populations and weakens behavioural coupling at movement onset

a. Peri-event time histograms aligned to onset of whisker movement for pre- (top) and post-drop (bottom) periods for a GlyT2-CNO recording. Grey and red dots indicate the absolute peak of neuronal activity centred around whisking onset for the pre- and post-drop period, respectively. Units ranked according to their peak firing rate. Temporal jitter peak activity decreases after CNO application.

b. Temporal dispersion of neuronal population activity before and after drug/vehicle administration (pre- and post-drop) for control (n = 12) and GlyT2-CNO (n = 13) recordings. Each data point represents the standard deviation of the distribution of absolute peak times of neuronal activity for each recorded population. Reduction of GlyT2-positive cell inhibition decreases the temporal dispersion of neuronal activity during whisking initiation (two-sided Wilcoxon signed-rank test, T = 12, p = 0.017).

c. Average whisking onsets for two representative GlyT2-CNO recordings before (grey) and after (red) drug administration. Dashed lines represent linear fits of initial whisker protraction. CNO delivery was associated with both increased (top) and decreased (bottom) slope of protraction.

d. Change in slope of whisker protraction at movement onset before and after drug/vehicle administration (pre- and post-drop) for control and GlyT2-CNO recordings.

e. Box plot showing differences in slope of whisker protraction at movement onset between pre- and post-drug/vehicle administration for control and GlyT2-CNO recordings. The standard deviation of the slope distribution in the GlyT2-CNO condition is higher than controls (Levene's test W = 8.39, p = 0.008) suggesting that decreasing local GlyT2-positive cell inhibition produces variable changes in the dynamics of movement onset across recordings.

1240 **f.** Relationship between change in cerebellar population dynamics and whisker
1241 movement following drug/vehicle administration for all recordings. Under control
1242 conditions, measured changes in neural activity and movement are small in magnitude
1243 and well correlated (r^2 = 0.52). In GlyT2-CNO recordings, relationships between
1244 cerebellar activity and movement are decoupled (r^2 = 0.15).

Figure 1 - Palacios et al. 2024

Figure 2 - Palacios et al. 2024

Figure 3 - Palacios et al. 2024

Figure 4 - Palacios et al. 2024

Figure 5 - Palacios et al. 2024