Golgi cells regulate timing and variability of information transfer in a cerebellar-behavioural loop

Ensor Rafael Palacios[1,2], Conor Houghton[2] and Paul Chadderton[1]

[1]School of Physiology Pharmacology and Neuroscience, University of Bristol, University Walk, Bristol, UK. BS8 1TD

[2]School of Engineering Mathematics and Technology, University of Bristol, Woodland Road, Bristol, UK. BS8 1UB

Correspondence should be addressed to:

Ensor Rafael Palacios (ensorrafael.palacios@bristol.ac.uk);
Paul Chadderton (p.chadderton@bristol.ac.uk)

No of pages: 47 (5 Figures; 1 Table)

Abstract word count: 168
Introduction word count: 573
Discussion word count: 1500

ACKNOWLEDGEMENTS

This work was supported by a Wellcome Trust Neural Dynamics PhD studentship to ERP, Leverhulme Research Fellowship (RF-2021-533) to CH and Wellcome Trust Investigator Award (209453/Z/17/Z) to PC. The schematic images in Figure 1 were generated and provided by Elisabeth Meyer whom we also thank for comments on the manuscript.

The authors declare no competing interests.

ABSTRACT

Golgi cells are inhibitory interneurons residing in the input layer of the cerebellar cortex. These neurons sit in a key position to govern the transformation of incoming information from extracerebellar regions and influence downstream cerebellar processing. Here, we examine the contribution of Golgi cells to network dynamics in Crus 1 of mouse lateral cerebellar cortex during free whisking. We recorded neuronal population activity using NeuroPixels probes before and after chemogenetic downregulation of Golgi cell activity. Under resting conditions, cerebellar population activity reliably encoded whisker movements. Reductions in Golgi cell activity produced mild increases in neural activity which did not significantly impair these sensorimotor representations. However, reduced Golgi cell inhibition did increase the temporal alignment of local population network activity at the initiation of movement. These network alterations had variable impacts on behaviour, producing both increases and decreases in whisking velocity. Our results suggest that Golgi cell inhibition primarily governs the temporal patterning of population activity, which in turn is required to support downstream cerebellar dynamics and behavioural coordination.

SIGNIFICANCE STATEMENT

The cerebellum is known to have a simple and highly conserved structure which has tantalised neurobiologists wishing to understand its function. Here we look at the role of one specific class of inhibitory interneuron, Golgi cells, in the input layer of the cerebellar cortex. We selectively turned down Golgi cell activity in the awake cerebellum to characterise the influence on network activity and behaviour. Here we show that downregulation of Golgi cells has very little influence on sensorimotor representations in the cerebellum (i.e., *what* is represented), but instead modulates the timing of cortical population activity (i.e., *when* information is represented). Our results indicate that Golgi cells are necessary to appropriately pace changes in cerebellar activity to match ongoing behaviour.

INTRODUCTION

The cerebellum is implicated in a diverse range of behaviours, extending from simple reflexes to complex functions such as language, and social interaction (Baumann & Mattingley, 2022; Kelly et al., 2020; Silveri, 2021). The circuit architecture underpinning this behavioural diversity is relatively simple and conserved across its extent, suggesting that different classes of input are subject to common processing mechanisms. The cerebellar cortex lacks recurrent excitation and almost all local feedback is mediated via inhibitory Golgi cells situated in the granule cell layer. Excitatory signals arriving in the granule cell layer via mossy fibres (MFs) are combined and propagated to the molecular layer through parallel fibres (PFs). Excitatory PF signals and recruited feedforward inhibition modulate Purkinje cell activity, the cortical output signal. Golgi cells receive direct input from MFs and excitatory feedback from PFs, enabling them to exert feedforward and feedback control over granule cells (Dieudonne, 1998; Eccles et al., 1966; Kanichay & Silver, 2008). Golgi cells are numerically sparse relative to granule cells and possess extended axonal arborisations, meaning each Golgi cell innervates many hundreds or thousands of granule cells. Inhibition via Golgi cells could thus potently impact the transformation of MF signals into PF activity. *In vivo* measurements confirm that inhibition gates the transformation of sensorimotor information in the granular layer (Chadderton et al., 2004; Duguid et al., 2012), sparsifying responses to sensory input, and facilitating pattern separation at the population level (Fleming et al., 2024). However, we lack quantitative information about how changes in Golgi cell activity influences cerebellar cortical dynamics and affects sensorimotor representations in the molecular and Purkinje cell layers. Indeed, the net influence of Golgi cell activity in the cerebellar cortex is not known but may be presumed to alter both the rate and timing of granule cell output. Resultant changes in PF excitation could have net excitatory or inhibitory effects at the level of Purkinje cells, because of molecular layer inhibition (Jelitai et al., 2016), and consequently for cerebellar nuclear neurons.

Here we investigate the influence of Golgi cell activity on cerebellar population activity in the context of whisker movement, to reveal how inhibitory feedback influences sensorimotor transformations and ongoing behaviour. Neuronal representations of

whisking activity in the lateral cerebellum are widespread (Bosman et al., 2010; Brown

& Raman, 2018; Chen et al., 2016, 2017; Zhai et al., 2024), and are partly based on a

linear rate code, whereby the firing rates of cerebellar neurons predict upcoming

whisker position (Chen et al., 2016). These representations are first seen in patterns

of mossy fibre input and are passed forward at each stage of cerebellar cortical

processing (Chen et al., 2017). Golgi cell inhibition has the potential to modulate how

incoming whisking information is encoded and transformed within the granule cell layer

(Gurnani & Silver, 2021; Palacios et al., 2021). However, it is unknown whether, and

how, changing levels of Golgi cell inhibition impacts neuronal representations of

whisker behaviour, and indeed whisking behaviour itself. We have recorded whisker

movements and population activity in the lateral cerebellum using NeuroPixels probes

in head-fixed mice expressing inhibitory DREADDs (designer receptors exclusively

activated by designer drugs) selectively in Golgi cells of lobule Crus 1. In these mice,

it was possible to downregulate Golgi cell activity by activating DREADDs with the

agonist, clozapine N-oxide (CNO), applied topically on the recording site. Our results

reveal a subtle but direct influence of Golgi cells on network dynamics and

consequently, the cerebellar-behavioural loop underlying voluntary whisking.

MATERIALS AND METHODS

Experiments were conducted on male and female C57BL/6 (wild-type) and heterozygous C57BL/6-Slc6a5<tm1.1(cre)Ksak> (GlyT2-Cre knockin) mice (Kakizaki et al., 2017) aged between 3-6 months. Animals were housed under a 12/12 h reversed light–dark cycle with food and water available *ad libitum*. Habituation and recordings were performed during the dark phase of the cycle. The C57BL/6-Slc6a5<tm1.1(cre)Ksak> mouse strain was provided by the RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan. All experiments were conducted under the United Kingdom Animals (Scientific Procedures) Act 1986 in accordance with Home Office guidelines.

Surgical procedures

Mice were anaesthetised for all surgical procedures. Mice were placed in an induction chamber, into which 5% v/v isoflurane (Harvard Apparatus Ltd) was administered via inhalation using oxygen (1.2–1.6 L/min). Once anaesthetized the concentration of isoflurane was lowered to 1–2% v/v to maintain anaesthesia for the duration of the surgery. The body temperature of the mouse was maintained at 37 ± 0.5 °C using a homeothermic heat mat (DC Temperature Control System, FHC). Corneal drying was prevented using an ocular lubricant (Lacrilube, Allergen). To induce analgesia, mice were injected subcutaneously with carprofen (Rimadyl, 5 mg/kg), buprenorphine (Vetergesic, 0.1 mg/kg) and lidocaine (2 mg/kg, locally). Surgery was carried out in a stereotaxic frame under aseptic conditions to prevent infections. After surgery, mice were placed in a warmed box (~37°C) and allowed to recover for as long as required. Mice were then returned to their home cages and monitored closely for at least 24 hrs.

GlyT2-Cre mice underwent two surgical operations: one to transduce cre-dependent expression of DREADDs in cerebellar Golgi cells via viral injection, and one to prepare the animal for head-fixed extracellular recording. During the first surgical session, a small craniotomy (~1 mm diameter) was performed over Crus 1 (6.36 mm posterior and 2.5 mm lateral of Bregma). AAV9-hSyn-DIO-hM4D(Gi)-mCherry (500-1000 nl) was injected with a glass pipette at different depths (starting at 600µm retracting to

surface in steps of 100µm), waiting 2 minutes between pipette retractions. At the end

of the operation, skin over the head was sutured and animal was allowed to recover.

The second surgical session took place 8 weeks later to enable ample expression of

the DREADD receptor protein. First, neck muscles were gently moved to uncover the

cerebellum. A custom-made head implant was fixed to the exposed cranium using

tissue glue (Histoacryl, Braun Corporation) and dental cement (Associate Dental

Products Ltd). A small craniotomy (~1.5 mm diameter) was performed over the right

cerebellar hemisphere, the dura was removed, and a reference screw was inserted in

contact with the underlying brain. To secure both head place and reference screw,

dental cement (Associate Dental Products Ltd) was used to cover the skull, sparing a

recording well above Crus 1 of the left cerebellar hemisphere. A second craniotomy

(~2 mm diameter) was performed over left Crus 1, and the dura was removed. Finally,

a layer of agarose (1.5% in phosphate buffered saline; PBS) was used to cover the

brain, a layer of Kwik-Seal was used to protect the brain, and a layer of nail polish was

used to fix the Kwik-Seal. Wild-type mice underwent only the second surgical

operation in preparation for recordings. All mice had their whiskers trimmed on the left

(recording) side, preserving whiskers C1, C2 and C3 (posterior whiskers on the third

row from top).

Electrophysiological and video recordings

Mice were head-fixed in the recording apparatus in preparation for electrophysiological

recording. The Kwik-Seal was removed, the surface of the brain was cleaned from

agarose and kept moist with PBS. NeuroPixels probes were fixed to a

micromanipulator (IVM, Scientifica) via a custom-made 3D printed holder. Each probe

was coated with DiI stain (2.5 mg/ml) for *ex vivo* probe tracking before insertion at an

angle perpendicular to the cortical surface. The probe was advanced into the cortex

at a speed of 2-5 µm/s to a depth of 2500-3000 µm. Thereafter the probe was retracted

for ~100 µm and left to settle for 10-15 min, allowing the brain to relax. Under infrared

light illumination, whisker movements were filmed with a high-speed camera (Genie

HM640; Teledyne Dalsa Inc, USA) operating at 299 frames per second. Video

acquisitions were controlled by Streampix 6 software (Norpix, Canada). The open-

source software spikeGLX 3.0 (https://billkarsh.github.io/SpikeGLX/) was used to

record NeuroPixels data at 30 kHz. Video and electrophysiological recording were
synchronised by a TTL pulse originating from the video software. In chemogenetic
experiments, a baseline of 10 or 20 min was recorded prior to topical administration of
CNO (30 µM), delivered to the recording well. Previously Stachniak et al., 2014
showed that intracranial injection of 1 µM CNO is already sufficient to reliably inhibit
presynaptic neurotransmitter release within minutes. After CNO or vehicle delivery,
electrophysiological recordings continued for a further 40-50 mins, for a total recording
time of 1 hour. Animals underwent two recording sessions, the first performed at least
4 hours after recovery from the craniotomy, the second on the following day. After
recording, the probe was extracted, rinsed with deionised water and left in freshly
made 1% tergazyme solution for at least 24 hours. After the first recording, the
exposed brain surface was first rinsed with PBS, then covered with agarose, and finally
protected with a layer of Kwik-Seal that was fixed to the surrounding dental cement
with nail polish. Mice were returned to their cage. After the second recording session,
mice were humanely killed. See **Supplementary Table 1** for full information on data
included for analysis.

Electrophysiological data processing

Output files (.imec) were preprocessed using the command-line tool CatGT
(https://billkarsh.github.io/ SpikeGLX/#catgt) to apply a high pass filter (cut-off 300 Hz)
and global demux filters, a common average referencing that takes in account the
probe channels subgrouping during data acquisition. Spike sorting was conducted
using the open-source software Kilosort 2 (https://github.com/MouseLand/ Kilosort) to
group spikes into units, each representing the activity of a distinct putative neuron.
Manual curation of sorted units was done with the open-source Python-library Phy2
(https://github.com/cortex-lab/phy), and consisted in merging and splitting units, as
well as categorising them into 'good', 'bad' and 'mua' (multiunit activity) units. In all
analysis, only good, i.e. well-isolated, units located in the cerebellar cortex were used.

Video processing and behavioural analysis

Video output (.avi) files were transcoded (.mp4, libx265 encoder), re-sampled at 299
Hz) and cropped using the ffmpeg software. We used the open-source toolbox

DeepLabCut (https://github.com/DeepLabCut/DeepLabCut) to label four landmarks on
whiskers C1, C2 and C3 (**Figure 1a**) across the entire recording; these landmarks
were then used to compute whisker position. Specifically, the arctan function was used
to measure the azimuthal angle between the first whisker segment (traced from the
first, basal marker to the second marker) and the horizontal line passing through the
basal marker. Increases in angle correspond to whisker protraction, whereas
decreases correspond to whisker retraction. Next, the phase of the whisking cycles
was extracted using the Hilbert transform, in order to compute the slowly changing
whisking amplitude and setpoint across cycles (Hill et al., 2011). Finally, the whisking
amplitude was low pass filtered and used to identify periods of whisking. A heaviside
function with a threshold of 10°, was used to discriminate whisking from resting
periods. All whisking analysis focused on the C3 (most anterior) whisker.

Histology

Brains were extracted and left in 4% PFA solution for 24-48 hours at 4°C. After
washing with PBS, cerebellar coronal sections (50-100 μm) were sliced using a
vibrating microtome (Leica VT1000S). Fluorescent images were acquired using a
confocal microscope (Leica DM4000 B).

EXPERIMENTAL DESIGN AND STATISTICAL ANALYSES

Single unit analysis

To compute the tuning curve for each unit, we computed the total spike count and
average whisker position (angle) in time bins of 33 ms. We then discretised the
average whisking angles into 11 bins, ranging from the minimum to the maximum
angle observed in each recording and paired the spike count in each time bin with the
corresponding whisker angle. These data were finally used to plot the firing rate,
together with their standard error, across the 11 angle bins.

Both the entropy of and KL-divergence between the pre- and post-drug tuning curves
for each unit was computed using the function entropy from the scipy.stats library. The
tuning curve for each unit was normalised to be the probability mass function of firing
rates over whisking position.

Clustering of tuning curves was performed with *k*-means algorithm
(sklearn.cluster.KMeans package), used on the spline coefficient fitted to the individual
unit tuning curves. The estimation of the tuning curves can be noisy due to, for
example, errors during the spike sorting process, which may lead to the contamination
of the spike history assigned to one unit by the activity of nearby neurons. This noise,
in turn, could affect clustering of tuning curves into meaningful groups. To reduce the
impact of this source of noise on our clustering analysis, we fitted a spline model to
each tuning curve: the model used seven cubic b-spline functions to smooth the firing
rate associated to each angle bin based on the firing rate of adjacent bins, while
maintaining the underlying structure of the tuning curves. The smoothed tuning curves
were then clustered using a *k*-means algorithm with $k = 8$. We finally computed the
average tuning curve from units within each cluster. Results were qualitatively the
same with fewer than $k = 8$ clusters.

Principal component analysis of population activity

We used principal component analysis to describe the population activity of each
recording. Projections onto the first three eigenvectors (principal components; pcs)
described the first three orthogonal axes in neuronal space along which population
activity varied the most. To perform this analysis, we first segmented each recording
into discrete whisking epochs (from -2 to +3 seconds around the start of each whisking
bout, defined using behavioural analysis above), and computed the trial-averaged
firing rate for each unit. Firing rates were used to compute correlation matrices across
units, from which eigenvalues and eigenvectors were derived. Eigenvalues indicate
the amount of neuronal activity variability captured along the axis specified by the
corresponding eigenvectors. For each recording, the eigenvectors had length equal to
the number of units present in that recording. The values of an eigenvector are known
as loads and correspond to the weight associated to each unit when linearly combining
the firing rate from all units; hence, the loadings of an eigenvector correspond to how
much each unit contributes to the pc associated with that eigenvector. The
eigenvectors can be used to project either each unit's firing rates in pc space or single
trial activity: to compute the pc for single trial activity, we first smoothed single trial

spike counts with a Gaussian window of std = 20 (we also used std = 5 but results
were equivalent; time bin width = 3.3 ms).

Cross-correlation between whisking position and each pc was obtained using the
scipy.signal.correlate package. All cross-correlations were normalised by the absolute
maximum value of the first pc. For analysis involving CNO manipulation, cross-
correlation peaks for pre- and post-drop periods were computed, and the respective
distributions were compared using a two-sided Wilcoxon signed-rank test. To
compared cross-correlations across conditions, we first computed the within-recording
contrast between post- versus pre-drop cross correlation peaks; next, for each pc and
experimental condition, we fitted the contrasts with a Gaussian distribution, and finally
we took the difference in posterior samples of the mean between GlyT2-CNO and
control conditions.

Analysis to assess efficacy of DREADD manipulation

A multilevel model was used to account for the variability in total cerebellar cortical
spike counts detected over time across recordings; counts were the sum of the spikes
assigned to good (i.e. well-isolated) cortical units. Spike counts were computed for
successive 5 minute-time bins, with the 0-time bin including the time of drug
application. The counts in each bin were scaled with respect to the count in the -5-
minute bin.

In the model, spike counts were described using an inverse-Gamma distribution:

$$s \sim \text{Inverse-Gamma}(\alpha, \beta)$$

$$\alpha = \exp(\theta_\alpha + \theta_{\alpha c} + \theta_{\alpha a} + \theta_{\alpha t}) \qquad (1)$$

$$\beta = \exp(\theta_\beta + \theta_{\beta c} + \theta_{\beta a} + \theta_{\beta t})$$

The inverse-Gamma distribution is used because it provides a good fit to the
population spike count data (**Figure 3e**); the two parameters α and β are usually
referred to as the shape and scale parameters and are related to the mean and
variance by $\mu = \beta/(\alpha - 1)$ and $\sigma^2 = \beta^2/[(\alpha-2)^2(\alpha-1)]$. Here α and β are each linked to a
linear model by a loglink function. Both linear models involve a coefficient θ for each

explanatory variable, namely, the experimental condition c, the time of the recording t

(dim = 9), and the animal ID a (dim = 17). The experimental conditions were, (i)

application of CNO to cerebellar surface of DREADD-transduced GlyT2-Cre mice

(GlyT2-CNO, n = 19 recordings), (ii) application of CNO to cerebellar surface of wild-

type mice (WT-CNO, n = 5 recordings), (ii) application of PBS the cerebellar surface

of wild-type or GlyT2 mice (WT-Veh, n = 9 recordings). Experimental conditions were

split by recording period into a pre-drug or baseline period (bins before 0-time bin) a

post-drug period (0-time bin and onward), for a total dimensionality of 3 x 2 (dim=6)

for the $\theta_{\alpha c}$ and $\theta_{\beta c}$ coefficient. The contrasts between pre- and post-drug coefficient θ_c

were used to assess the effect of the experimental manipulation.

The additional explanatory variables, t and a, were used to explain respectively spike

count variability within each pre-drug time bin due to natural changes in network

activity over recording time (n = 9, excluding -5-min time bin), and variability due to the

particular conditions of the network in each animal (n = 17), which might be affected,

for example, by the surgical procedure. Thus, the addition of these explanatory

variables allowed us to test within conditions for a difference in pre- and post-drop

spike count distribution, after accounting for variability explained by temporal

correlations and between-animal differences.

Each θ, except θ_t was drawn from a vector of independent normal distributions, of

length equal to the dimensionality of θ. The prior distribution for θ_t was a multivariate

normal distribution with covariance decaying with time distance, accounting for

temporal correlations within the data:

$$\Sigma_\alpha = I * \exp(-\delta_{tij}) \qquad \Sigma_\beta = I * \exp(-\delta_{tij})$$

$$\theta_\alpha \sim Normal(\mu_\alpha, \sigma_\alpha) \qquad \theta_\beta \sim Normal(\mu_\beta, \sigma_\beta)$$

$$\theta_{\alpha c} \sim Normal(\mu_\alpha, \sigma_\alpha) \qquad \theta_{\beta c} \sim Normal(\mu_\beta, \sigma_\beta) \qquad (2)$$

$$\theta_{\alpha a} \sim Normal(\mu_\alpha, \sigma_\alpha) \qquad \theta_{\beta a} \sim Normal(\mu_\beta, \sigma_\beta)$$

$$\theta_{\alpha t} \sim MVNormal(\mu_\alpha, \Sigma_\alpha) \qquad \theta_{\beta t} \sim MVNormal(\mu_\beta, \Sigma_\beta)$$

with δ_{tij} between any two periods i and j is defined as:

$$\delta_{tij} = exp\left(-\left|\frac{ti-tj}{5}\right| * 0.7\right)$$

330   (3)

331   Where $\left|t_i - t_j\right|/5$ is the absolute time difference between period i and j scaled to unity

332   (5 is the time width of each period in minutes).

333   Finally, the hyperpriors:

334                          $\mu_\alpha = 1$ $\mu_\beta = 1.2$

335   (4)

336                          $\sigma_\alpha = 0.2$ $\sigma_\beta = 0.2$

were chosen to sample prior predictive samples within a similar scale of the observed

population spike counts, including extreme values.

The posterior samples for the mean and variance spike count parameters were used

to compute the difference across conditions between post- and pre-drug contrasts in

mean and variance due to the experimental manipulation (**Figure 3f**). In practice, the

contrasts were computed using the same posterior samples, except for $\theta_{\alpha c}$ and $\theta_{\beta c}$,

which differed for pre- and post-drug mean and variance. The difference between

contrasts was used to assess changes in mean and variance across conditions

(**Figure 3g**).

Analysis of DREADD manipulation on neuronal and whisking behaviour

Comparisons were made from the pre- and post-drug time periods. The pre-drug

period lasted 20 minutes all but five recordings, where this period was 10 minutes. The

post-drug time period matched the pre-drug period and started 5 minutes after drug

application. Recordings that exhibited little or no whisking activity were excluded from
this analysis leaving N = 25 recordings in total.

For each population recording, we computed the absolute peak of the trial-averaged
neuronal activity (PETH) aligned to whisking onset. The subwindow used to compute
the absolute peaks spanned -0.7 to 1.3 sec around whisking onset, to focus the
subsequent analysis around the initial whisking protraction period. We used the
absolute peak of the PETH as a proxy of the timing of the neuronal response to
whisking-related input. For each recording we computed the standard deviation (std)
of the peaks, for pre- and post-drug data. For each experimental condition, the pre-
and post-drug stds were compared using a two-sided Wilcoxon signed-rank test.

For corresponding whisking data, we used a linear model to fit a line to the average
whisking position pre- and post-drug. Whisking data were delimited by a time window
ranging from -0.06 to 0.21 sec centred around whisking onset; the time window was
chosen to focus on the period of initial whisking protraction. We computed the
difference between post-drop and pre-drug slopes (i.e. velocity to compare the effects
of Golgi cell manipulation on the whisking protraction phase. To test whether the
variance of the distribution slope contrasts in the two experimental conditions were
different, we used Levene's test (scipy.stats.levene).

Code accessibility

Code used for the analysis is freely available
here: https://zenodo.org/doi/10.5281/zenodo.13327395

RESULTS

Population dynamics in lateral cerebellum during voluntary whisking

To understand the contribution of Golgi cells to cerebellar cortical dynamics, we first set out to describe how cerebellar populations represent sensorimotor information when Golgi cell inhibition remains intact. We therefore recorded cerebellar population activity using NeuroPixels probes in lobule Crus 1 of awake mice and performed spike sorting to isolate single units (see Materials and Methods). We isolated an average of 42.3 (std = 25.3) cortical units per recording (N = 12), for a total (n) of 508 putative units. To correlate spiking activity with whisker movement, we simultaneously tracked spontaneous whisking via a high-speed camera (**Figure 1a**). In all cases, bouts of whisking activity were associated with pronounced alterations in cerebellar spiking activity (**Figure 1b, c**). Single cell recordings previously revealed widespread tuning to whisker position in granule cells, inhibitory interneurons and Purkinje cells (Bosman et al., 2010; Brown & Raman, 2018; Chen et al., 2016, 2017; Zhai et al., 2024). We therefore measured tuning to whisker position amongst single units of our population recordings. On a unit-by-unit basis, we examined the relationship between firing rate and whisker position, to construct individual tuning curves (see Materials and Methods). Consistent with previous work, we observed robust tuning to whisker position, such that single units exhibited their highest firing rates at preferred angles of whisker deflection.

Within locally recorded populations, we observed substantial diversity in the tuning to whisker position. Indeed, units recorded in the same penetration displayed a broad range of preferred angles and tuning profiles, including monotonic increases and decreases to whisker protraction/retraction (**Figure 1d**). We then recombined all single unit spiking responses to calculate population tuning. Whereas individual units exhibited diverse selectivity, the mean tuning function for the population displayed a near-monotonic dependence upon whisker position, with rate changes in proportion to the magnitude of protraction and retraction (**Figure 1e**).

To further assess tuning diversity at the population level, we clustered units based on
their tuning curves, testing whether we could discriminate discrete functional
subclasses of unit. We fitted a spline model with 7 cubic b-spline to each tuning curve
(n = 508, N = 12) to capture tuning curve shape and reduce variability due to noise
(see Materials and Methods). We then inspected the output of k-means clustering with
different numbers of expected clusters. With increasing cluster number, there was no
evidence of a change in the sum-of-squared-distances decaying pattern, or elbow,
which can be used as an indicator of the optimal number of clusters to be considered.
Visual inspection of the mean tuning curves for each cluster revealed a progressive
shift of preferred position from the most retracted to the most protracted whisker
locations (**Figure 1f**). Projection of tuning curves in 3D space using t-SNE, a non-
linear dimensionality reduction technique particularly suited to uncovering clustering
structure further confirmed that different groups of clusters are not well isolated (data
not shown). Thus, our analysis does not support discrete classes of tuning curve but
rather confirms that cerebellar neurons form a functional continuum in the space of
possible tuning curves. At the local population level, Crus 1 cerebellar cortical neurons
heterogeneously encode whisking position, but represent increasing protractive and
retractive movements via accumulating increases in firing rates.

Having established that concerted population activity is required for accurate
sensorimotor representations, we performed principal component analysis upon the
grouped spike trains from each recording. We initially preserved projections of
population activity onto the first three eigenvectors, which describe the three
orthogonal axes in neuronal space along which activity varies the most (**Figure 1g**).
Analysis was restricted around whisking bouts (-2 to +3 seconds about whisking
onset). Projections of population activity revealed clear structure in neuronal space
during transitions from quiescence to whisking activity and back again. This structure
in turn suggested that population activity contains accurate information about whisking
dynamics. We tested this proposal by linearly combining the first three principal
components, attempting to reconstruct movement trajectories (see Materials and
Methods). Using this approach, it was possible to decode whisking setpoint during
single trials (**Figure 1h**). In summary, our results show that the population activity in

the lateral cerebellar cortex accurately reflects upcoming slow whisking dynamics,

even when approximated within a low-dimensional space.

To quantify the relationship between different principal components and whisking, we

computed the normalised cross-correlation between whisker angle and each of the

first 3 principal components (pc) in each recording. Interestingly, pc1-3 showed distinct

temporal relationships with respect to whisker movement (**Figure 2a,b**). Information

related to whisker position contained in pc1 tended to anticipate behaviour, whereas

information in pc2 and pc3 lagged behaviour. Across all recordings (N = 12), the peak

signal of pc1 always preceded the whisking signal, indicating that, across recordings,

variation in the whisking-aligned average neuronal activity expressed by the pc1 tends

to anticipate behaviour (one sample t test, t = 4.72, p = 0.0006, mean = 36 ± 8 ms).

On the other hand, the peak signal for pc2 almost always occurred after whisker

movement, indicating that whisker-related information contained in pc2 lags whisking

activity (t = -4.43, p = 0.001, mean = -195 ± 44 ms). Finally, peak signals for pc3 were

broadly distributed around 0 ms (t = -1.38, p = 0.19, mean = -89 ± 65 ms). Notably,

pc1 peaks across recordings are narrowly distributed in time; this suggests that pc1

specifically contains precise information about whisking behaviour, i.e. upcoming

movement, occurring ~40 ms in the future (**Figure 2c**).

Whisking-related information contained in pc1-3 could originate from a small

proportion of units in the population, whose activity may be particularly responsive to

whisking; alternatively, information could be more distributed, whereby all units

contribute to some extent to the population encoding of behaviour. To measure how

behavioural information was distributed across the population, we measured the pc

loading of each unit (n = 508), indicating the contribution of a unit to the corresponding

pc (see Materials and Methods). The loading distribution for each pc was centred

around 0 and was roughly symmetric, meaning that units could contribute both

positively or negatively to each pc (**Figure 2d**). The number of outliers in each

distribution was used to quantify the extent to which each pc is dominated by the

activity of few units. We calculated the excess kurtosis of each distribution, which is a

measure of the frequency of outliers observed, using as a point of reference the

frequency expected from normally distributed data. The distribution for pc1 had an

excess kurtosis of -2.85, indicating infrequent outliers; this in turn suggests that the

contribution of units across recordings to pc1 is broadly distributed within the

population. In contrast, the excess kurtosis for pc3 was 4.54, indicating a high

frequency of outliers, and thus a subset of units which tend alone to contribute mostly

to pc3. The excess kurtosis for pc2 was -0.47, meaning that outliers are neither

frequent nor infrequent. Together, these results suggest that for pc1, and thus accurate

representation of whisking behaviour, information is broadly distributed across the

entire population. Finally, we noted that the first three eigenvalues accounted for only

a moderate amount of the total variance in the population activity (~30%) across all

recordings (**Figure 2e,f**). Therefore pc1-3 provide only a coarse description of

population activity, and whisker movement only partially describes neuronal dynamics

in Crus 1 despite restricting our analysis to the initial period of whisking activity. Indeed,

the large proportion of neuronal variability not captured in the first 3 principal

components supports the proposal that cerebellar cortex may represent additional

behaviourally relevant variables.

Chemogenetic downregulation of Golgi cells elevates population activity

Having established the relationship between population activity and slow whisking

dynamics under baseline conditions, we next sought to understand how Golgi cell

inhibition contributes to these representations in the cerebellar cortex. We adopted a

chemogenetic strategy to selectively downregulate Golgi cell activity in an otherwise

intact network while simultaneously recording population activity and whisker

movement. Manipulation of Golgi cells was achieved via activation of the inhibitory

DREADD, hM4Di, which causes reductions in excitability and neurotransmitter release

(Stachniak et al., 2014) in cells expressing these receptors. Localised expression of

DREADDs (**Figure 3b**) was achieved by injecting the AAV-DIO-hM4D(Gi)-mCherry

virus into Crus 1 of GlyT2-Cre mice (cre-recombinase restricted to Golgi cells in the

cerebellar cortex; see Materials and Methods; Kakizaki et al., 2017). DREADDs were

activated by local, topical delivery of the agonist, clozapine-N-oxide (CNO), on top of

the Neuropixels penetration site, which itself had been targeted as the site of virus

injection (**Figure 3a**); this method was used to minimise potential off-site effects of the

agonist and afforded within-minute temporal precision to our manipulation. To confirm

the fidelity of our approach, we first tested the effects of topical delivery of muscimol (1.7 μg/μl), a GABA$_A$-receptor agonist, during NeuroPixels recordings from wildtype mice (N = 3). In all cases, within minutes of drug application, the total spike count recorded across the cerebellar cortex started to decrease and reached its minimum within 10-30 minutes (data not shown). These results supported the viability of applying CNO topically onto cerebellar cortex to locally activate DREADD receptors.

We next examined whether CNO-dependent reduction in Golgi cell activity had a measurable impact on cerebellar cortical population activity. To do this, we compared changes in cortical spike count before and after CNO administration across three experimental groups. GlyT2-Cre mice, pre-injected with AAV9-hSyn-DIO-hM4D(Gi)-mCherry virus into Crus 1, or uninjected C57BL6 mice received topical delivery of CNO (GlyT2-CNO; *N* = 19 recordings) or saline vehicle (Veh; *N* = 9 recordings) to the cerebellar cortex. A third group of uninjected wildtype (C57BL6) mice received topical delivery of CNO to the cerebellum (WT-CNO; *N* = 5 recordings). Initial inspection of the data revealed an increase in normalised spike count in the GlyT2-CNO groups, but not Veh and WT-CNO groups, shortly after application of CNO (**Figure 3c,d**). Additionally, when comparing the mean of the normalised post-drop cortical spike counts (5-minute bin width), we found a significant difference across conditions (ANOVA, *F* = 10.91, *p* = 2.82e-05). However, significant variability in spike counts were observed both within and between experimental groups. Therefore, to confirm that alterations in spike count observed across recordings were due to chemogenetic manipulation of Golgi cells, we used a multilevel model approach; this approach was warranted by the temporal correlations between repeated measures and clustering by animals. The model described the distribution of spike counts for each time bin with an inverse-Gamma distribution, parameterised by a shape parameter α and a scale parameter β. The model used as explanatory variables the experimental conditions, associated to pre- and post-drug administration periods, to find the α and β parameters that best explained the pre- and post-drug administration data. For each parameter of the gamma distribution (*α* and *β)* and each group (WT-CNO, Veh, GlyT2-CNO), the model had a coefficient *θ* associated to the pre-drug recording period and one for the post-drug administration period. The contrast between the two coefficients was used to measure the change in spike counts due to the experimental manipulation. To

confirm the fidelity the model, we compared the distribution of all experimentally observed spike counts together with the model fit (**Figure 3e**): the overlap between these distributions indicated that the model performed well in capturing the overall structure of the data.

Having confirmed the fidelity of the model, we then compared the contrasts between model coefficients associated with the pre- and post-drug administration periods for each experimental group. These contrasts reflect the effect of the experimental manipulation on spike count distribution. For each experimental group there are two contrasts, one associated with the shape parameter α and one with the scale parameter β of the distribution describing spike counts. Comparing contrasts and highest density intervals (hdi, 94%) for the parameters across groups revealed a specific effect in GlyT2-CNO mice on both α and β (**Figure 3f**). To gain a better intuition of the implications of these results, we used the posterior samples of α and β to derived samples for the mean and variance parameters of the inverse-Gamma distribution; the mean and variance offer an alternative parametrisation that characterise more intuitively the centre of mass and spread of spike counts across recordings for each condition, before and after drug application (see Materials and Methods). Thus, to compare results across condition, we computed the pre- minus post-drug mean and variance contrasts for each experimental condition, and the difference in contrasts between conditions, for each parameter. This analysis revealed that both mean and variance increased after drug application in the GlyT2-CNO group compared to the two control groups. (**Figure 3g**; mean hdi 0.05- 0.46, 0.05 - 0.49 and variance hdi - 0.01 – 0.31, -0.02 – 0.31 for GlyT2-CNO versus WT-CNO and Veh respectively). It further revealed that in the WT-CNO group, CNO application alone has no detectable effect on network spiking activity (mean hdi -0.17– 0.21 and variance hdi -0.02 – 0.03 for WT-CNO versus Veh). Together our analyses confirm that chemogenetic inhibition of Golgi cells causes a global increase in spiking activity within the cerebellar cortex.

Limited influence of Golgi cell downregulation on sensorimotor representation

Having established that chemogenetic downregulation of Golgi cells measurably alters cerebellar activity, we investigated how the perturbation affected sensorimotor

representations of whisking. Golgi cell inhibition acts via the granule cell layer, but
changes in the input-output transformation of granule cells will have knock on effects
upon the activity of molecular interneurons and Purkinje cells. Golgi cells control spike
output of granule cells through various mechanisms, including changes in gain, i.e.,
slope of the input-output function (Brickley et al., 1996; Chadderton et al., 2004;
Hamann et al., 2002; Mitchell & Silver, 2003). By reducing Golgi cell inhibition, granule
cells may become more sensitive to excitatory inputs, increasing their gain. We looked
for changes in the sensitivity of cerebellar activity to whisker movement by comparing
population tuning curves before and after CNO administration in GlyT2 mice. We
observed no striking differences in the overall profile of these curves (**Figure 4a**). Post-
CNO, population firing rates were marginally higher for some whisker positions, but
tuning curves were essentially unchanged. This result suggests that chemogenetic
Golgi cell downregulation has no substantial effect on gain or overall representation of
whisker position at the population level. In principle, altering Golgi inhibition could
cause reorganisation of the granule cell population code by altering the requirements
for granule cell integration of excitatory inputs from different mossy fibres (Marr, 1969).
We therefore explored whether perturbed Golgi cell activity causes substantial
reorganisation of whisking representation at the single cell level. To do this, we
measured the entropy of individual tuning curves before and after CNO/vehicle
administration. Entropy is a measure of how informative units are about, in this case,
whisking position, with higher-entropy neurons being less informative. Following CNO
administration, we observed no changes in cumulative entropy distribution in GlyT2-
CNO recordings (**Figure 4b**). Interestingly, under control conditions (i.e. Veh and WT-
CNO recordings), units tended to have higher entropy values after vehicle/CNO
administration, indicating that tuning curves become less informative about whisking
position over time. We further compared the distributions of entropy differences pre-
and post-drug delivery (**Figure 4c**). Under control conditions, tuning curve entropies
typically remained unchanged or increase over time (control density skewness = 2.80).
CNO administration in GlyT2 mice causes additional reorganisation - some units
decrease their entropy - leading to an even distribution of entropy changes after drug
application (GlyT2-CNO density skewness = 0.21). The cumulative entropy distribution
in GlyT2-CNO recordings therefore remains similar across time suggesting that the
firing rate of some units becomes more sensitive to changes in whisking position

following Golgi cell manipulation. These results also imply that recording conditions in the control condition are associated with small but progressive losses of representational fidelity. To confirm this proposal, we again measured the correlation between each of the first three population principal components (pc1-3) and whisker position for GlyT2-CNO and control recordings (**Figure 4d,e**). We found no evidence for large differences in correlation pre- and post-drug administration. However, there was a general trend for correlations to decrease after drug delivery, except for pc1 and pc2 (i.e. fast and slower representations of whisking) in the GlyT2-CNO condition, where correlations increased slightly. Our results suggest that the fidelity of sensorimotor representation is subject to a mild downward drift under control conditions in the present experimental setting, and that Golgi cell downregulation may slightly improve representation, offsetting this drift. But overall, we did not observe notable changes in the quality of the encoding of whisking behaviour by population activity. Together our results indicate that individual sensorimotor tuning of cerebellar neurons is only weakly influenced by chemogenetic Golgi cell perturbation.

Golgi cell downregulation alters cerebellar dynamics and weakens coupling to whisking behaviour

We further explored the impact of Golgi cell perturbation by comparing trial averaged peri-event time histograms (PETH) and whisking activity from pre- and post-drug administration periods (see Materials and Methods). Because cerebellar activity can directly influence whisker movement (Proville et al., 2014), reduced Golgi cell inhibition could influence both cerebellar dynamics and produce measurable changes in whisking behaviour. We first considered cerebellar activity: reductions in Golgi cell inhibition could alter the temporal dynamics of population activity in several ways, acting via tonic and/or phasic inhibition (Brickley et al., 1996; Chadderton et al., 2004; Duguid et al., 2015), and/or affecting Golgi cell synchrony (van Welie et al., 2016). We therefore compared the patterns of population activity at movement onset before and after chemogenetic perturbation. Typically population activity peaks close to the onset of individual whisking bouts (**Figure 1b,c**). Therefore, for each recording, we computed the standard deviation of the distribution of peak unit firing times aligned to whisking onset, pre- and post-drug delivery (**Figure 5a**; see Materials and Methods).

In the GlyT2-CNO mice, but not controls, pre- and post- drug standard deviations were significantly different (**Figure 5b**), with significantly lower variation following CNO administration (two-sided Wilcoxon signed-rank test, $T = 12$, $p = 0.017$). Thus, Golgi cell downregulation increased the temporal alignment of peak neuronal activity around whisking onset. These results suggest that decreased Golgi cell inhibition has an impact on the pacing of neuronal responses around the onset of whisking activity, and in particular narrows the time window within which units fire most strongly.

Finally, we measured the effects of Golgi cell manipulation on whisking velocity at the onset of movement bouts (see Materials and Methods). For each recording, we used a linear model to approximate the slope of the average whisking bouts during its protraction phase (**Figure 5c**); the coefficient for each linear fit was used as a read out for velocity during the movement initiation. Interestingly, unlike control recordings, GlyT2-CNO mice showed changes in their whisking behaviour following drug administration. However, the sign of these changes was not consistent across animals, occurring in both increasing and decreasing directions (**Figure 5d**). Individual onsets of voluntary whisking thus became faster or slower after drug administration in GlyT2-CNO mice. The overall distribution of changes in whisker onset velocity ('slope contrast') was therefore significantly broader in the GlyT2-CNO mice compared to control (**Figure 5e**). Thus, Golgi cell downregulation produces measurable but divergent effects on local cerebellar dynamics and whole animal whisking behaviour. Local cerebellar dynamics were consistently compressed, leading to a more tightly locked population activity, whereas whisking velocity was either increased or reduced. These results suggest a breakdown in the link between cerebellar dynamics and behaviour following Golgi cell inhibition. We confirmed this by plotting changes in population activity and whisking velocity at movement onset for individual recordings (**Figure 5f**). In control recordings, differences in network activity and whisker movement before and after drug are small in magnitude but importantly, are well correlated ($r^2 = 0.52$). In GlyT2-CNO recordings, changes in cerebellar dynamics and movement become decoupled from one another, leading to break down in this correlation ($r^2 = 0.15$). Together our results indicate Golgi cell inhibition regulates the temporal patterning of population activity and that disruption of this patterning at the local level can have diverse consequences for behaviour.

DISCUSSION

Golgi cells are proposed to play important roles in cerebellar transformation (Albus, 1971; D'Angelo et al., 2013; Marr, 1969; Palacios et al., 2021), but we lack detailed information about their influence on circuit activity. Here we have tested directly the contribution of Golgi cells to cerebellar population activity by analysing whisking behaviour and its neuronal representations in intact networks and those where Golgi cell inhibition was reduced. To do this, we used hM4i receptor activation, targeted specifically to cerebellar Golgi cells, locally and selectively downregulating these neurons during voluntary behaviour. We used multi-level modelling methods to demonstrate that Golgi cell perturbations cause subtle changes in neural population activity. Our perturbation did not profoundly disrupt or alter sensorimotor representations in the cerebellar cortex and may be considered relatively mild. Such an alteration of network dynamics is likely within normal physiological operating limits, and the changes we observe can thus be presumed to be ethologically relevant.

Cerebellar cortical neurons heterogeneously encode whisking behaviour

Population recordings from intact networks show that lateral cerebellar neurons accurately represent whisking behaviour. Individual units exhibit different tuning profiles to whisking position, including monotonic increases, decreases, and localised increases/decreases in firing rate (**Figure 1d**), consistent with previous work (Bosman et al., 2010; Brown & Raman, 2018; Chen et al., 2016, 2017), enabling multidimensional representations of movement kinematics (Gurnani & Silver, 2021; Markanday et al., 2023). Due their small size and high packing density, our units are unlikely to be granule cells (the postsynaptic target of Golgi cells) (Beau et al., 2024), but rather inhibitory interneurons and Purkinje cells. However, as observed in small virtual granule cell populations (Chen et al., 2017), neuronal ensemble activity displayed a near-monotonic dependence upon whisker set point change, with increasing activity in the direction of both protraction and retraction (**Figure 1e**). These results are in line with previous findings (Brown & Raman, 2018; Chen et al., 2016, 2017; Lanore et al., 2021; Proville et al., 2014).

The cerebellar cortex uses a distributed code to represent whisking dynamics

We used principal component analysis to describe population activity with three independent variables. The first three pcs capture only a moderate amount of the total variance in the population activity, suggesting that cerebellar activity integrates and reflects multiple other behaviourally relevant variables (Chabrol et al., 2015; Ishikawa et al., 2015; Sobel et al., 1998; Wagner & Luo, 2020). Nevertheless, neuronal dynamics projected into 3D component space still had structure that well matched whisking dynamics (**Figure 1g**). Notably, the dynamics captured by pc1 consistently led (i.e. anticipated) whisking behaviour by ~40ms whereas pc2 typically lagged behaviour, more variably, by a few hundred milliseconds (**Figure 2**). The former is consistent with a forward model of planned or predicted movement, while the latter resembles a feedback signal. These results suggest a role for the cerebellum in controlling future behaviour, incorporating recent sensory feedback in addition to other behaviourally relevant variables, in line with theoretical treatments of the cerebellum (Palacios et al., 2024). Whisker-related information, especially in the first principal component, was well distributed across units. Thus, information about whisking is broadly distributed in the recorded neuronal populations, with most units contributing to some extent to its encoding.

Decreasing Golgi cell inhibition does not saturate population activity

Next, we examined how altering Golgi cell activity affected neuronal representations in the cerebellar cortex. This was achieved by activating inhibitory receptors (hM4Di) on Golgi cells, through application of the agonist CNO directly at the surface of the recording site. This approach afforded within-minute temporal precision to the manipulation. hM4Di receptors inhibit neuronal activity via moderate hyperpolarisation and reduced presynaptic release of neurotransmitters (Roth, 2016). Topical application of CNO onto the cerebellar cortex produced an increased spread of the population spike count distribution, with higher spike counts becoming more likely across recordings (**Figure 3c,d**). However, manipulation of Golgi cell inhibition did not lead to excitatory saturation of network dynamics, i.e., a drastic and indiscriminate increase in neural activity, and instead weakly affected the magnitude of population

activity. The absence of strong saturation of network activity may be because granule
cell excitability is under strong control of tonic inhibition, which arises in part from non-
vesicular sources of GABA (Lee et al., 2010; Rossi et al., 2003; Wall & Usowicz, 1997),
and/or compensatory upregulation of molecular layer interneuron activity. The tuning
of individual neurons to whisker position was also largely unchanged (**Figure 4a**).
Theoretical and *in vitro* evidence has indicated that the gain of granule cell input/output
transformation may be under the control of local inhibitory processes. As such,
reduced inhibition is predicted to increase cortical sensitivity to incoming synaptic
input. Following our perturbation, we saw little evidence of increased sensitivity to
features of whisker movement (**Figure 4a**), although we did observe a small fraction
of neurons that exhibited reduced entropy following Golgi cell inhibition (**Figure 4c**).
These may represent a subset of neurons that receive subthreshold whisker-related
synaptic input. Here, small decreases in synaptic inhibition may result in the
generation of tuned action potential output (Chadderton et al., 2004; Chen et al.,
2017). However, our results as a whole suggest that dynamic Golgi cell inhibition may
not be necessary for maintaining network excitation at the level required for the
network to perform its computations (Albus, 1971; Marr, 1969); instead, manipulation
of Golgi cell inhibition produces subtle effects on cerebellar cortical neuronal
dynamics.

Decreased Golgi cell inhibition increases the temporal alignment of neuronal
activation during movement onset

Decreased Golgi cell inhibition was associated with increased temporal alignment or
synchronisation within neuronal populations around whisking onset. A familiar
downstream consequence of feedforward inhibition in many brain regions, and at the
molecular layer interneuron-Purkinje cell synapse, is increased temporal precision of
neuronal dynamics, as inhibition normally trails excitation and sharpens the window of
input integration (Gabernet et al., 2005; Mittmann et al., 2005; Wehr & Zador, 2003).
Here, notably, Golgi cells play the opposite role, temporally dispersing or
desynchronising neural activity (Vervaeke et al., 2010). Although Golgi cells provide
feedforward inhibition to the granule cell layer, evoked inhibition has been previously
shown to precede or be concomitant with mossy fibre excitation, under which

situations it may reduce the temporal precision of early granule cell responses (Duguid

et al., 2015), and decorrelate multisensory representations (Fleming et al., 2024). Our

observations are consistent with such a mode of operation, as decreasing Golgi cell

inhibition leads to an increased temporal alignment of the overall network activity

during movement initiation (Vervaeke et al., 2010). Changes in Golgi cell activity can

modulate the amplitude of tonic inhibitory conductances (Crowley et al., 2009).

Reduced Golgi cell activity may therefore directly reduce the tonic inhibitory

conductance, making granule cells more excitable and more likely to fire rapidly to a

burst of incoming mossy fibre synaptic inputs. Such control by Golgi cells of the timing

of network activity can support the specific cerebellar role in integrating information

and learning contexts (Fleming et al., 2024). Indeed, distinct temporal patterning of

granule cell responses to mossy fibre input is proposed to be important for how the

cerebellar network integrates and distinguishes parallel sensory and motor pathways

(Chabrol et al., 2015).

Decreasing Golgi cell inhibition decreases the stability of whisking behaviour

Finally, we found that reduced Golgi cell inhibition led to increased variability in

whisking velocity at movement onset. The specific change in movement velocity varied

amongst recordings: in some cases, the velocity of whisking increased while other

cases it decreased. We suggest that location of the site of perturbation, varying from

mouse to mouse, is likely to have influenced the valence of the change in whisking

behaviour (Apps et al., 2018). Importantly, the link between cerebellar cortical

dynamics and whisker position is decoupled after Golgi cell perturbation (**Figure 5f**).

This result highlights the importance of an intact cerebellum for precise whisking

control, which is subject to the coordination with other motor and sensory systems

(Deschenes et al., 2012; Kurnikova et al., 2017; McElvain et al., 2018; Romano et al.,

2020). Increased variability in motor initiation may be linked with the possible

contribution of Golgi cell inhibition to representations of multiple streams of information

and behaviours – bound together at the level of the cerebellar cortex – improving

accuracy as specific behaviours are performed.

Conclusions

Our work shows how altering Golgi cell inhibition changes the response of the network to its input. Golgi cell inhibition itself is under the control of many mechanisms, including feedback loops encompassing the cortex (Dieudonne & Dumoulin, 2000; Nietz et al., 2017; Witter et al., 2016), the cerebellar nuclei (Ankri et al., 2015), as well as neuromodulatory systems (Fleming & Hull, 2019; Fore et al., 2020). Thus, Golgi cells sit in a key position to flexibly control cerebellar computations, based on the dynamic behavioural context, which may be required to focus on different aspects of behaviour at different times (Palacios et al., 2021). To fully understand the pervasive involvement of the cerebellum in behavioural control, it is critical to elucidate how different mechanisms contribute to fine tune the Golgi cell network in a context-dependent manner.

AUTHOR CONTRIBUTIONS

ERP, CH and PC designed the experiments. All experiments were performed by ERP. Data was analysed and interpreted by ERP, CH and PC. The authors declare no competing interests.

REFERENCES

Albus, J. S. (1971). A theory of cerebellar function. *Math Biosci*, *10*(1-2), 25-61.

Ankri, L., Husson, Z., Pietrajtis, K., Proville, R., Lena, C., Yarom, Y., Dieudonne, S., & Uusisaari, M. Y. (2015). A novel inhibitory nucleo-cortical circuit controls cerebellar Golgi cell activity. *eLife*, *4*. https://doi.org/10.7554/eLife.06262

Apps, R., Hawkes, R., Aoki, S., Bengtsson, F., Brown, A. M., Chen, G., Ebner, T. J., Isope, P., Jorntell, H., Lackey, E. P., Lawrenson, C., Lumb, B., Schonewille, M., Sillitoe, R. V., Spaeth, L., Sugihara, I., Valera, A., Voogd, J., Wylie, D. R., & Ruigrok, T. J. H. (2018). Cerebellar Modules and Their Role as Operational Cerebellar Processing Units: A Consensus paper [corrected]. *Cerebellum*, *17*(5), 654-682. https://doi.org/10.1007/s12311-018-0952-3

Baumann, O., & Mattingley, J. B. (2022). Cerebellum and Emotion Processing. *Adv Exp Med Biol*, *1378*, 25-39. https://doi.org/10.1007/978-3-030-99550-8_3

Beau, M., Herzfeld, D. J., Naveros, F., Hemelt, M. E., D'Agostino, F., Oostland, M., Sanchez-Lopez, A., Chung, Y. Y., Michael, M., Kyranakis, S., Stabb, H. N., Martinez Lopera, M. G., Lajko, A., Zedler, M., Ohmae, S., Hall, N. J., Clark, B. A., Cohen, D., Lisberger, S. G., . . . Medina, J. F. (2024). A deep-learning strategy to identify cell types across species from high-density extracellular recordings. *bioRxiv*. https://doi.org/10.1101/2024.01.30.577845

Bosman, L. W., Koekkoek, S. K., Shapiro, J., Rijken, B. F., Zandstra, F., van der Ende, B., Owens, C. B., Potters, J. W., de Gruijl, J. R., Ruigrok, T. J., & De Zeeuw, C. I. (2010). Encoding of whisker input by cerebellar Purkinje cells. *J Physiol*, *588*(19), 3757-3783. https://doi.org/10.1113/jphysiol.2010.195180

Brickley, S. G., Cull-Candy, S. G., & Farrant, M. (1996). Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA receptors. *J Physiol* *497*(3), 753-759. https://doi.org/10.1113/jphysiol.1996.sp021806

Brown, S. T., & Raman, I. M. (2018). Sensorimotor Integration and Amplification of Reflexive Whisking by Well-Timed Spiking in the Cerebellar Corticonuclear Circuit. *Neuron*, *99*(3), 564-575 e562. https://doi.org/10.1016/j.neuron.2018.06.028

Chabrol, F. P., Arenz, A., Wiechert, M. T., Margrie, T. W., & DiGregorio, D. A. (2015). Synaptic diversity enables temporal coding of coincident multisensory inputs in single neurons. *Nat Neurosci*, *18*(5), 718-727. https://doi.org/10.1038/nn.3974

Chadderton, P., Margrie, T. W., & Hausser, M. (2004). Integration of quanta in cerebellar granule cells during sensory processing. *Nature*, *428*(6985), 856-860. https://doi.org/10.1038/nature02442

Chen, S., Augustine, G. J., & Chadderton, P. (2016). The cerebellum linearly encodes whisker position during voluntary movement. *eLife*, *5*, e10509. https://doi.org/10.7554/eLife.10509

Chen, S., Augustine, G. J., & Chadderton, P. (2017). Serial processing of kinematic signals by cerebellar circuitry during voluntary whisking. *Nat Commun*, *8*(1), 232. https://doi.org/10.1038/s41467-017-00312-1

Crowley, J. J., Fioravante, D., & Regehr, W. G. (2009). Dynamics of fast and slow inhibition from cerebellar golgi cells allow flexible control of synaptic integration. *Neuron*, *63*(6), 843-853. https://doi.org/10.1016/j.neuron.2009.09.004

D'Angelo, E., Solinas, S., Mapelli, J., Gandolfi, D., Mapelli, L., & Prestori, F. (2013). The cerebellar Golgi cell and spatiotemporal organization of granular layer activity. *Front Neural Circuits*, *7*, 93. https://doi.org/10.3389/fncir.2013.00093

Deschenes, M., Moore, J., & Kleinfeld, D. (2012). Sniffing and whisking in rodents. *Curr Opin Neurobiol*, *22*(2), 243-250. https://doi.org/10.1016/j.conb.2011.11.013

Dieudonne, S. (1998). Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell synaptic currents in the rat cerebellum. *J Physiol*, *510*(3), 845-866. https://doi.org/10.1111/j.1469-7793.1998.845bj.x

Dieudonne, S., & Dumoulin, A. (2000). Serotonin-driven long-range inhibitory connections in the cerebellar cortex. *J Neurosci*, *20*(5), 1837-1848. https://doi.org/10.1523/JNEUROSCI.20-05-01837.2000

Duguid, I., Branco, T., Chadderton, P., Arlt, C., Powell, K., & Hausser, M. (2015). Control of cerebellar granule cell output by sensory-evoked Golgi cell inhibition. *Proc Natl Acad Sci U S A*, *112*(42), 13099-13104. https://doi.org/10.1073/pnas.1510249112

Duguid, I., Branco, T., London, M., Chadderton, P., & Hausser, M. (2012). Tonic inhibition enhances fidelity of sensory information transmission in the cerebellar cortex. *J Neurosci*, *32*(32), 11132-11143. https://doi.org/10.1523/JNEUROSCI.0460-12.2012

Eccles, J. C., Llinas, R., & Sasaki, K. (1966). The mossy fibre-granule cell relay of the cerebellum and its inhibitory control by Golgi cells. *Exp Brain Res*, *1*(1), 82-101. https://doi.org/10.1007/BF00235211

Fleming, E., & Hull, C. (2019). Serotonin regulates dynamics of cerebellar granule cell activity by modulating tonic inhibition. *J Neurophysiol*, *121*(1), 105-114. https://doi.org/10.1152/jn.00492.2018

Fleming, E. A., Field, G. D., Tadross, M. R., & Hull, C. (2024). Local synaptic inhibition mediates cerebellar granule cell pattern separation and enables learned

sensorimotor associations. *Nat Neurosci*, *27*(4), 689-701. https://doi.org/10.1038/s41593-023-01565-4

Fore, T. R., Taylor, B. N., Brunel, N., & Hull, C. (2020). Acetylcholine Modulates Cerebellar Granule Cell Spiking by Regulating the Balance of Synaptic Excitation and Inhibition. *J Neurosci*, *40*(14), 2882-2894. https://doi.org/10.1523/JNEUROSCI.2148-19.2020

Gabernet, L., Jadhav, S. P., Feldman, D. E., Carandini, M., & Scanziani, M. (2005). Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition. *Neuron*, *48*(2), 315-327. https://doi.org/10.1016/j.neuron.2005.09.022

Hamann, M., Rossi, D. J., & Attwell, D. (2002). Tonic and spillover inhibition of granule cells control information flow through cerebellar cortex. *Neuron*, *33*(4), 625-633. https://doi.org/10.1016/s0896-6273(02)00593-7

Hill, D. N., Curtis, J. C., Moore, J. D., & Kleinfeld, D. (2011). Primary motor cortex reports efferent control of vibrissa motion on multiple timescales. *Neuron*, *72*(2), 344-356. https://doi.org/10.1016/j.neuron.2011.09.020

Ishikawa, T., Shimuta, M., & Hausser, M. (2015). Multimodal sensory integration in single cerebellar granule cells in vivo. *eLife*, *4*. https://doi.org/10.7554/eLife.12916

Jelitai, M., Puggioni, P., Ishikawa, T., Rinaldi, A., & Duguid, I. (2016). Dendritic excitation-inhibition balance shapes cerebellar output during motor behaviour. *Nat Commun*, *7*, 13722. https://doi.org/10.1038/ncomms13722

Kakizaki, T., Sakagami, H., Sakimura, K., & Yanagawa, Y. (2017). A glycine transporter 2-Cre knock-in mouse line for glycinergic neuron-specific gene manipulation. *IBRO Rep*, *3*, 9-16. https://doi.org/10.1016/j.ibror.2017.07.002

Kelly, E., Meng, F., Fujita, H., Morgado, F., Kazemi, Y., Rice, L. C., Ren, C., Escamilla, C. O., Gibson, J. M., Sajadi, S., Pendry, R. J., Tan, T., Ellegood, J., Basson, M. A., Blakely, R. D., Dindot, S. V., Golzio, C., Hahn, M. K., Katsanis, N., . . . Tsai, P. T. (2020). Regulation of autism-relevant behaviors by cerebellar-prefrontal cortical circuits. *Nat Neurosci*, *23*(9), 1102-1110. https://doi.org/10.1038/s41593-020-0665-z

Kurnikova, A., Moore, J. D., Liao, S. M., Deschenes, M., & Kleinfeld, D. (2017). Coordination of Orofacial Motor Actions into Exploratory Behavior by Rat. *Curr Biol*, *27*(5), 688-696. https://doi.org/10.1016/j.cub.2017.01.013

Lanore, F., Cayco-Gajic, N. A., Gurnani, H., Coyle, D., & Silver, R. A. (2021). Cerebellar granule cell axons support high-dimensional representations. *Nat Neurosci*, *24*(8), 1142-1150. https://doi.org/10.1038/s41593-021-00873-x

Lee, S., Yoon, B. E., Berglund, K., Oh, S. J., Park, H., Shin, H. S., Augustine, G. J., & Lee, C. J. (2010). Channel-mediated tonic GABA release from glia. *Science*, *330*(6005), 790-796. https://doi.org/10.1126/science.1184334

Marr, D. (1969). A theory of cerebellar cortex. *J Physiol*, *202*(2), 437-470. https://doi.org/10.1113/jphysiol.1969.sp008820

McElvain, L. E., Friedman, B., Karten, H. J., Svoboda, K., Wang, F., Deschenes, M., & Kleinfeld, D. (2018). Circuits in the rodent brainstem that control whisking in concert with other orofacial motor actions. *Neuroscience*, *368*, 152-170. https://doi.org/10.1016/j.neuroscience.2017.08.034

Mitchell, S. J., & Silver, R. A. (2003). Shunting inhibition modulates neuronal gain during synaptic excitation. *Neuron*, *38*(3), 433-445. https://doi.org/10.1016/s0896-6273(03)00200-9

Mittmann, W., Koch, U., & Hausser, M. (2005). Feed-forward inhibition shapes the spike output of cerebellar Purkinje cells. *J Physiol*, *563*(2), 369-378. https://doi.org/10.1113/jphysiol.2004.075028

Nietz, A. K., Vaden, J. H., Coddington, L. T., Overstreet-Wadiche, L., & Wadiche, J. I. (2017). Non-synaptic signaling from cerebellar climbing fibers modulates Golgi cell activity. *eLife*, *6*. https://doi.org/10.7554/eLife.29215

Palacios, E. R., Houghton, C., & Chadderton, P. (2021). Accounting for uncertainty: inhibition for neural inference in the cerebellum. *Proc Biol Sci*, *288*(1947), 20210276. https://doi.org/10.1098/rspb.2021.0276

Dieudonne, S. (1998). Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell synaptic currents in the rat cerebellum. *J Physiol*, *510 (Pt 3)*(Pt 3), 845-866. https://doi.org/10.1111/j.1469-7793.1998.845bj.x

Eccles, J. C., Llinas, R., & Sasaki, K. (1966). The mossy fibre-granule cell relay of the cerebellum and its inhibitory control by Golgi cells. *Exp Brain Res*, *1*(1), 82-101. https://doi.org/10.1007/BF00235211

Gurnani, H., & Silver, R. A. (2021). Multidimensional population activity in an electrically coupled inhibitory circuit in the cerebellar cortex. *Neuron*, *109*(10), 1739-1753 e1738. https://doi.org/10.1016/j.neuron.2021.03.027

Kanichay, R. T., & Silver, R. A. (2008). Synaptic and cellular properties of the feedforward inhibitory circuit within the input layer of the cerebellar cortex. *J Neurosci*, *28*(36), 8955-8967. https://doi.org/10.1523/JNEUROSCI.5469-07.2008

Markanday, A., Hong, S., Inoue, J., De Schutter, E., & Thier, P. (2023). Multidimensional cerebellar computations for flexible kinematic control of movements. *Nat Commun*, *14*(1), 2548. https://doi.org/10.1038/s41467-023-37981-0

Palacios, E. R., Chadderton, P., Friston, K., & Houghton, C. (2024). Cerebellar state estimation enables resilient coupling across behavioural domains. *Sci Rep*, *14*(1), 6641. https://doi.org/10.1038/s41598-024-56811-x

Palacios, E. R., Houghton, C., & Chadderton, P. (2021). Accounting for uncertainty: inhibition for neural inference in the cerebellum. *Proc Biol Sci*, *288*(1947), 20210276. https://doi.org/10.1098/rspb.2021.0276

Vervaeke, K., Lorincz, A., Gleeson, P., Farinella, M., Nusser, Z., & Silver, R. A. (2010). Rapid desynchronization of an electrically coupled interneuron network with sparse excitatory synaptic input. *Neuron*, *67*(3), 435-451. https://doi.org/10.1016/j.neuron.2010.06.028

Proville, R. D., Spolidoro, M., Guyon, N., Dugue, G. P., Selimi, F., Isope, P., Popa, D., & Lena, C. (2014). Cerebellum involvement in cortical sensorimotor circuits for the control of voluntary movements. *Nat Neurosci*, *17*(9), 1233-1239. https://doi.org/10.1038/nn.3773

Romano, V., Reddington, A. L., Cazzanelli, S., Mazza, R., Ma, Y., Strydis, C., Negrello, M., Bosman, L. W. J., & De Zeeuw, C. I. (2020). Functional Convergence of Autonomic and Sensorimotor Processing in the Lateral Cerebellum. *Cell Rep*, *32*(1), 107867. https://doi.org/10.1016/j.celrep.2020.107867

Rossi, D. J., Hamann, M., & Attwell, D. (2003). Multiple modes of GABAergic inhibition of rat cerebellar granule cells. *J Physiol*, *548*(1), 97-110. https://doi.org/10.1113/jphysiol.2002.036459

Roth, B. L. (2016). DREADDs for Neuroscientists. *Neuron*, *89*(4), 683-694. https://doi.org/10.1016/j.neuron.2016.01.040

Silveri, M. C. (2021). Contribution of the Cerebellum and the Basal Ganglia to Language Production: Speech, Word Fluency, and Sentence Construction-Evidence from Pathology. *Cerebellum*, *20*(2), 282-294. https://doi.org/10.1007/s12311-020-01207-6

Sobel, N., Prabhakaran, V., Hartley, C. A., Desmond, J. E., Zhao, Z., Glover, G. H., Gabrieli, J. D., & Sullivan, E. V. (1998). Odorant-induced and sniff-induced activation in the cerebellum of the human. *J Neurosci*, *18*(21), 8990-9001. https://doi.org/10.1523/JNEUROSCI.18-21-08990.1998

Stachniak, T. J., Ghosh, A., & Sternson, S. M. (2014). Chemogenetic synaptic silencing of neural circuits localizes a hypothalamus-->midbrain pathway for feeding behavior. *Neuron*, *82*(4), 797-808. https://doi.org/10.1016/j.neuron.2014.04.008

van Welie, I., Roth, A., Ho, S. S., Komai, S., & Hausser, M. (2016). Conditional Spike Transmission Mediated by Electrical Coupling Ensures Millisecond Precision-Correlated Activity among Interneurons In Vivo. *Neuron*, *90*(4), 810-823. https://doi.org/10.1016/j.neuron.2016.04.013

Wagner, M. J., & Luo, L. (2020). Neocortex-Cerebellum Circuits for Cognitive Processing. *Trends Neurosci*, *43*(1), 42-54. https://doi.org/10.1016/j.tins.2019.11.002

1059   Wall, M. J., & Usowicz, M. M. (1997). Development of action potential-dependent and
1060   independent spontaneous GABAA receptor-mediated currents in granule cells
1061   of postnatal rat cerebellum. *Eur J Neurosci*, *9*(3), 533-548.
1062   https://doi.org/10.1111/j.1460-9568.1997.tb01630.x
1063
1064   Wehr, M., & Zador, A. M. (2003). Balanced inhibition underlies tuning and sharpens
1065   spike timing in auditory cortex. *Nature*, *426*(6965), 442-446.
1066   https://doi.org/10.1038/nature02116
1067
1068   Witter, L., Rudolph, S., Pressler, R. T., Lahlaf, S. I., & Regehr, W. G. (2016). Purkinje
1069   Cell Collaterals Enable Output Signals from the Cerebellar Cortex to Feed Back
1070   to Purkinje Cells and Interneurons. *Neuron*, *91*(2), 312-319.
1071   https://doi.org/10.1016/j.neuron.2016.05.037
1072
1073   Zhai, P., Romano, V., Soggia, G., Bauer, S., van Wingerden, N., Jacobs, T., van der
1074   Horst, A., White, J. J., Mazza, R., & De Zeeuw, C. I. (2024). Whisker kinematics
1075   in the cerebellum. *J Physiol*, *602*(1), 153-181. https://doi.org/10.1113/JP284064
1076
1077
1078

 FIGURE LEGENDS

 Figure 1 Single unit and population activity in Crus 1 during voluntary whisking

a. Monitoring whisker movement and neuronal population activity in lateral cerebellar
cortex using high-speed videography and NeuroPixels recording in head-fixed mice.
Left: Whiskers were labelled and reconstructed using DeepLabCut to recover
movement trajectories. Image credit: Elisabeth Meyer. Right: Brightfield image of
coronal cerebellar section superimposed fluorescent image tract left by NeuroPixels
probe coated with DiI localised to lobule Crus 1.

b. 30 second segment of simultaneous whisking (top) and population activity (bottom;
n = 88 units) from a single recording session.

c. Trial-averaged whisker position and neuronal activity from the same recording,
aligned to onset of whisking bout.

d. Tuning curves of four individual cerebellar units from the same recording. Each unit
displays a different relationship between firing rate and whisker position. Blue lines
indicate fraction of time spent at different whisker positions; empty grey bars show
tuning curves for shuffled data. Left inset: colour-coded centre of mass of each unit on
NeuroPixels probe.

e. Average tuning curve for all recorded units, from all recordings (black line; n = 508,
N = 12). Population firing rate increases monotonically from whisker resting point
(white triangle) in both protractive and retractive directions. Blue line indicates fraction
of time spent at different whisker positions; shading represents standard deviation;
white triangle indicates whisker resting point; grey line shows tuning for shuffled data.

f. Average tuning curves for all units clustered in eight groups. Units were clustered
using k-mean algorithm based on individual unit tuning curves. Heterogeneity in tuning
curves enables a continuous encoding of whisking position.

g. Single trial of whisker movement (top) with corresponding projections in 3D principal
component space for average (bottom left) and single trial (bottom right) population

activity. Population activity in this space has structure that reflects whisking dynamics, both on average and at a single trial level.

h. Recovered movement trajectory using cerebellar population activity. Reconstruction of the whisking setpoint using a linear combination of the first three principal components computed from neuronal populations. Whisker setpoint (brown line) during two trials, together with predictions (red) and the highest density interval (hdi, red shading).

Figure 2 Population representations of whisker movement at different timescales

a. Trial-averaged whisking position (top) and projected population activity (bottom) for one recording.

b. Cross-correlations between pc1-3 and whisking activity for the same recording; vertical lines indicate times of peak correlation.

c. Cumulative variance explained by pc1-3 across all recordings (N = 12). Only a modest amount of variability in neuronal activity is accounted for by first three principal components.

d. Distribution of unit loadings (n = 508) for pc1-3. Kurtosis measurement, to quantify the number of outliers in each distribution, reveals fewer outliers for pc1 than what would be expected if the data were normally distributed (excess kurtosis -2.76). This indicates that information contained in pc1, which best reflects whisking activity, tends to be distributed across neurons. The loading distributions for pc2 and pc3 have respectively a similar or higher number of outliers compared to normally distributed data (excess kurtosis -0.43 and 4.97, respectively), indicating that information in pc2 and pc3 is increasingly concentrated in fewer units.

e. Time of peak correlation for pc1-3 across all recordings. The distribution for pc1 is tightly concentrated around ~40 ms, meaning that information captured by pc1 tends to anticipate whisking activity with high temporal precision (one sample t test, t = 4.71, p = 0.0006). The information contained in pc2 and pc3 shows the opposite trend, with more variability (pc2: mean -194 ms, t = -4.44, p = 0.0009; pc3: mean -92 ms, t = -1.44, p = 0.17), suggesting that these components might reflect different aspects of behaviour.

f. Mean decrease in unexplained variance (unexp. var.) across recordings with increasing numbers of principal components (#pcs); standard deviation shaded in grey.

Figure 3 Chemogenetic inhibition of Golgi cells increases neuronal activity in cerebellar cortex

a. Left: targeted expression of hM4Di receptors was achieved via injection of AAV-DIO-hM4Di-mCherry into the lateral cerebellar cortex of GlyT2-Cre mice, selectively expressing Cre-recombinase in Golgi cells in this brain region. Right: NeuroPixels probes were targeted to site of viral injection. During electrophysiological recordings, the exogenous drug, clozapine-N-oxide (CNO) was topically applied to activate hM4Di receptors.

b. Left: GlyT2-Cre mouse brain selectively expressing hM4Di in Golgi cells of the lateral cerebellar cortex. Right: tract left by NeuroPixels probe coated with DiI, showing co-localisation of the site of recording and hM4Di expression.

c. Changes in population spike count before and after CNO/vehicle delivery (at 0 min) normalised by count at -5 min, for three experiment conditions. Pink, GlyT2-Cre mice and CNO (N = 19); Blue, C57BL6 mice and CNO (N = 5); Green, GlyT2-Cre/C57BL6 mice and saline vehicle only (N = 9).

d. Box plots showing pooled data for each experimental condition.

e. Multilevel modelling approach to assess effect of drug delivery on cerebellar population activity. Population spike count distributions for each condition (top) were modelled using an inverse-Gamma distribution, described by shape parameter α and scale parameter β. Each parameter was modelled as a linear combination of different coefficients, including two θ_{cond}, one for α and one for β, which captured the specific effect of the experimental manipulation on total spike counts. Comparing the empirical distribution with the distribution of posterior predictive checks (samples from the model after fitting) shows that the model captures the overall structure of the data (bottom).

f. 94% highest density intervals (hdi) of the contrasts between post- and pre-drop posterior samples for θ_{cond} for α and β. The contrasts highlight a specific effect of Golgi cell manipulation (GlyT2 + CNO) on both α and β parameters.

g. Contrast difference (mean and variance) between post- and pre-drop of the inverse-

Gamma distribution fitted to total spike counts. Bars indicate the 94% hdi of the

posterior differences. Left, contrast difference between GlyT2 + CNO condition and

each control condition. Right, contrast difference between the two control conditions.

Golgi cell manipulation produces positive contrast differences versus both control

conditions.

Figure 4 Weak influence of Golgi cell perturbation on cerebellar sensorimotor representations

a. Average tuning curves for GlyT2-CNO recordings before (grey) and after (red) topical administration of the drug.

b. Entropy cumulative density functions for all units recorded in the GlyT2-CNO (top) and control (bottom) conditions. Under control conditions, but not GlyT2-CNO, unit entropy tends to increase after drug/vehicle delivery (post-drop), suggesting that tuning curves become less informative about whisking position over time.

c. Probability mass function of unit entropy differences before and after CNO/vehicle administration ('post-drop' minus 'pre-drop' entropy). Under control conditions, tuning curve entropy tends to remain stable or increase over time (density skewness = 2.79). In the GlyT2-CNO recordings, a fraction units exhibit decreased entropy after drug application (density skewness = 0.21), suggesting that some units become more sensitive to changes in whisking position following Golgi cell inhibition.

d. Comparison of peak pre- and post-drop correlation values between principal components 1-3 (pc1-3) and whisker position for control recordings.

e. Comparison of peak pre- and post-drop correlation values between principal components 1-3 (pc1-3) and whisker position for GlyT2-CNO recordings.

Figure 5 Golgi cell inhibition reduces temporal variability in neuronal populations and weakens behavioural coupling at movement onset

a. Peri-event time histograms aligned to onset of whisker movement for pre- (top) and post-drop (bottom) periods for a GlyT2-CNO recording. Grey and red dots indicate the absolute peak of neuronal activity centred around whisking onset for the pre- and post-drop period, respectively. Units ranked according to their peak firing rate. Temporal jitter peak activity decreases after CNO application.

b. Temporal dispersion of neuronal population activity before and after drug/vehicle administration (pre- and post-drop) for control (n = 12) and GlyT2-CNO (n = 13) recordings. Each data point represents the standard deviation of the distribution of absolute peak times of neuronal activity for each recorded population. Reduction of Golgi cell inhibition decreases the temporal dispersion of neuronal activity during whisking initiation (two-sided Wilcoxon signed-rank test, T = 12, p = 0.017).

c. Average whisking onsets for two representative GlyT2-CNO recordings before (grey) and after (red) drug administration. Dashed lines represent linear fits of initial whisker protraction. CNO delivery was associated with both increased (top) and decreased (bottom) slope of protraction.

d. Change in slope of whisker protraction at movement onset before and after drug/vehicle administration (pre- and post-drop) for control and GlyT2-CNO recordings.

e. Box plot showing differences in slope of whisker protraction at movement onset between pre- and post-drug/vehicle administration for control and GlyT2-CNO recordings. The standard deviation of the slope distribution in the GlyT2-CNO condition is higher than controls (Levene's test W = 8.39, p = 0.008) suggesting that decreasing local Golgi cell inhibition produces variable changes in the dynamics of movement onset across recordings.

f. Relationship between change in cerebellar population dynamics and whisker movement following drug/vehicle administration for all recordings. Under control conditions, measured changes in neural activity and movement are small in magnitude

1220   and well correlated (r^2 = 0.52). In GlyT2-CNO recordings, relationships between

1221   cerebellar activity and movement are decoupled (r^2 = 0.15).

Figure 1 - Palacios et al. 2024

Figure 2 - Palacios et al. 2024

Figure 3 - Palacios et al. 2024

Figure 4 - Palacios et al. 2024

Figure 5 - Palacios et al. 2024

Recording	Animal	Strain	Drug	Whisking	# cortical units	Time CNO/PBS drop (min)
1	1	GlyT2	CNO	poor	18	10
2	1	GlyT2	CNO	good	5 (4)	10
3	2	GlyT2	CNO	good	8 (7)	10
4	2	GlyT2	CNO	good	12 (11)	10
5	3	GlyT2	CNO	good	10	10
6	3	GlyT2	CNO	good	10 (6)	10
7	4	GlyT2	CNO	good	8	20
8	5	GlyT2	CNO	absent	27	20
9	6	GlyT2	CNO	good	61	20
10	7	GlyT2	CNO	good	76	20
11	7	GlyT2	CNO	good	24	20
12	8	GlyT2	CNO	absent	5	20
13	8	GlyT2	CNO	poor	30	20
14	9	GlyT2	CNO	good	58	20
15	9	GlyT2	CNO	good	23	20
16	10	GlyT2	CNO	good	46	20
17	10	GlyT2	CNO	good	49	20
18	11	GlyT2	CNO	poor	9	20
19	11	GlyT2	CNO	poor	29	20
20	12	WT	CNO	good	96 (88)	10
21	13	WT	CNO	good	28	20
22	13	WT	CNO	good	31	20
23	14	WT	CNO	good	14	20
24	14	WT	CNO	good	55 (54)	20
25	4	GlyT2	PBS	good	8	20
26	5	GlyT2	PBS	absent	23	20
27	6	GlyT2	PBS	good	31	20
28	15	WT	PBS	good	88	20
29	15	WT	PBS	poor	38	20
30	16	WT	PBS	good	61	20
31	16	WT	PBS	good	28	20
32	17	WT	PBS	good	52	20
33	17	WT	PBS	good	25	20

Supplementary Table 1: Neuronal and behavioural data. Each animal underwent two recordings (except animal 12). Each recording was performed on either GlyT2-Cre or C56BL6 (wild-type; WT) mice; GlyT2 mice expressed Cre-recombinase selectively in Golgi cells in the cerebellar cortex, and therefore only in these mice CNO could selectively decrease Golgi cell inhibition. The three experimental conditions are GlyT2+CNO, in which CNO was used on GlyT2-Cre mice, WT+CNO in which CNO was used on WT mice, and vehicle, in which phosphate-buffered saline (PBS) was used on either injected GlyT2-Cre mice (3 recordings) or WT mice (6 recordings); the WT+CNO and vehicle conditions were pooled into one control condition, after assessing for the specific effect of our manipulation on total cerebellar cortical spike counts using the statistical model described in equations 1-4 (see *Materials & Methods*). For this statistical analysis of spike counts, we used all units from all recordings (n = 1086 units, N = 33 recordings). In all other analyses, instead, we excluded recordings with absent/poor whisking activity (N = 25 after exclusion), as the analyses required concomitant behavioural and neuronal information; for the same reason, we additionally excluded a small number of units (n = 16, remaining number of units in parenthesis) whose activity was absent or too sparse during whisking periods in order to compute the peri-event time histogram. Poor whisking behaviour was defined as a flat trial-averaged whisker position trace for either or both pre- and post- drop periods. Control mice only were used for analysis included in Figures 1 and 2 (n = 508, N = 12 recordings after exclusion).