1 **Increasing intraspecific plant chemical diversity at plot and**
2 **plant level affects herbivorous, predatory, and pollinating**
3 **arthropod communities**
4

5 Lina Ojeda-Prieto[1], Eliecer L. Moreno[1], Robin Heinen[1*], Wolfgang
6 W. Weisser[1*]
7
8 [1] Terrestrial Ecology Research Group, Department for Life Science
9 Systems, TUM School of Life Sciences, Technical University of
10 Munich, 85354 Freising, Germany.
11 *These authors supervised the project equally and share the last
12 authorship
13
Author for correspondence
Lina Ojeda-Prieto
Terrestrial Ecology Research Group, Department for Life Science
Systems
Technical University of Munich
Hans Carl-von-Carlowitz-Platz 1. 85354, Freising, Germany
lina.ojeda@tum.de
21
22 **ORCIDs**
23 Lina Ojeda-Prieto: 0009-0003-6069-901X
24 Eliecer L. Moreno: 0009-0008-3798-3326
25 Robin Heinen: 0000-0001-9852-1020
26 Wolfgang W. Weisser: 0000-0002-2757-8959

27
Additional Information:
Main text word count: 4937 words.
Number of References: 82 references.
Number of figures and tables: 3 main figures (in color), 3
supplementary figures (in color), and 15 supplementary
tables (10 black and white, 5 in color).
34

Summary

36 1. Plant chemistry mediates interactions between plants and their environment. While intraspecific chemodiversity at the plant level is well-studied, the effects of groups of plants differing in chemistry on interactions need more attention. We conducted a field experiment to test how intraspecific chemodiversity affects plant-arthropod interactions.

42 2. We manipulated plots of *Tanacetum vulgare* L., differing in chemotype richness and composition, and monitored four arthropod groups (herbivores, flower visitors, predators, and ants) over three seasons. We hypothesized that higher plot-level chemotype richness enhances occurrence across all studied arthropod groups and that the effect on abundance would be negative for herbivore and ant abundance and positive for flower visitors and predators.

50 3. Our findings, as revealed by mixed models, show that with increasing plot-level chemotype richness, herbivore abundance decreased, flower visitor abundance increased, and predatory arthropods and ants remained largely unaffected.

54 4. These results underscore the role of plant chemical diversity in shaping insect communities and contributing to ecosystem dynamics.

57 **Key words:** aphids, flower visitors, herbivores, insects, predators, plant-insect interactions, *Tanacetum vulgare* (tansy), terpenoids

Introduction

Biodiversity is recognized as a driver of ecosystem functioning and stability (Mora et al., 2014; Espinosa-García, 2022). Biodiversity studies have focused on species richness, examining how increasing plant community diversity affects ecosystem-level processes (Schulze & Mooney, 1992; Cardinale et al., 2006; Eisenhauer et al., 2011; Weisser et al., 2017). Plant communities with greater species richness are linked to a more diverse community of organisms (Ebeling et al., 2018), alterations in food web structure (Giling et al., 2019), and increased ecosystem process rates (Hertzog et al., 2017; Meyer et al., 2017). While species-level diversity has been well-studied, recent research explores biodiversity within species (Crutsinger et al., 2006; Koricheva & Hayes, 2018; Raffard et al., 2019; Noto & Hughes, 2020; Zerebecki & Hughes, 2023; Govaert et al., 2024; Konefal et al., 2024).

One component of within-species diversity is chemodiversity (Müller et al., 2020; Wetzel & Whitehead, 2020). Chemodiversity refers to the variety of chemical compounds produced by organisms (Whitehead et al., 2021; Walker et al., 2022). These compounds, including terpenoids, alkaloids, and phenolics, play diverse roles in plant defense (Hare, 2011; Wetzel et al., 2023), communication, and reproduction (Hartmann, 2007; Whitehead et al., 2022), potentially altering ecosystem dynamics like herbivory rates and pollinator preferences (Eisenring et al., 2021; Müller & Junker, 2022). Chemodiversity affects ecological interactions, impacting richness, diversity, and community structure (Dyer & Jeffrey, 2021; Fernandez-Conradi et al., 2022; Petrén et al., 2023). Studies have shown cascading chemodiversity effects within food webs (Kos et al., 2011; Bálint et al., 2016; Senft et al., 2019).

Intraspecific chemodiversity can be investigated at different scales,
from individual plants to larger groups like plots or populations
(Müller et al., 2020; Ojeda-Prieto et al., 2024). Research at the plant
level has focused on chemodiversity effects on arthropod-plant
interactions (Kos et al., 2011; Senft et al., 2017; Benedek et al.,
2019). For instance, field observations show certain compounds
influence herbivore preference and performance (Kleine & Müller,
2011; Clancy et al., 2018), as well as natural enemies' behavior and
reproductive success (Benedek et al., 2015; Zytynska et al., 2019).
Manipulative studies reveal that plants with different chemical
compositions differ in attractiveness to aphids (Neuhaus-Harr et al.,
2024) and herbivore performance (Jakobs & Müller, 2018),
reinforcing the role of chemodiversity in plant-arthropod
interactions.

Larger-scale studies in chemodiversity are scarce and
predominantly correlative. Studies have examined how local
chemotype richness influences arthropod communities (Hauri et al.,
2021; Espinosa-García, 2022), but questions remain on the
responses of different arthropod groups. Manipulative experiments,
directly testing the consequences of a gradient of chemotype
richness at the group level, show that increasing chemodiversity
correlates with higher herbivore diversity but lower plant damage
(Bustos-Segura et al., 2017). By systematically manipulating plot-
level chemotype richness (CR), we aim to understand how
chemodiversity at plant and plot levels shapes arthropod abundance
and occurrence. Due to their distinct ecological roles,
chemodiversity's effects are expected to vary among arthropod
groups. This complexity has not been fully addressed, highlighting
the need for comprehensive experimental approaches on arthropod
groups and their interactions with plants.

Different plant-interacting groups, such as herbivores, flower
visitors, predators, and ants, may share different co-evolutionary
histories with – and offer different selective pressures to - the plant
that have likely shaped its chemical composition by selecting
chemical profiles that serve optimal plant fitness (Dyer & Jeffrey,
2021; Schneider et al., 2021; Whitehead et al., 2021, 2022).
Arthropod **abundance** is influenced by plant attractiveness and
resource suitability, but it is also shaped by a complex interplay of
factors, including predator-prey dynamics, trophic cascades, and
habitat structure. However, conceptually, these relationships are
best understood from a plant-centered perspective. For herbivores,
plants may benefit from having a diverse bouquet of chemicals to
deter herbivores and reduce their abundance and impact on the plant
(Glassmire et al., 2019; Kariñho-Betancourt, 2020; Zu et al., 2020;
Rabelo et al., 2023). For predatory arthropods, plants may benefit
from their presence as predators may control herbivore populations
(Barbosa et al., 2009; Turlings & Erb, 2018). Therefore, it is likely
that more diverse blends of compounds may have evolved in part to
attract predatory arthropods (De Boer et al., 2004, 2008; Clavijo
McCormick et al., 2012). Similarly, many flower visitors play a
crucial role in pollination (Jacobsen & Raguso, 2018; Dantas et al.,
2020). Plants may evolve specific compounds to attract more flower
visitors, and more diverse profiles attract more visitors. Ants may
have positive, negative, or no effects on plants (Yamawo et al.,
2021; Ramos et al., 2022). Given that many ants engage in
mutualisms with herbivorous aphids, it may be beneficial for plants
to evolve repellant compounds. On the other hand, if ants engage in
protective mutualisms with plants, it may be beneficial for plants to
evolve attractant compounds. Therefore, hypotheses regarding
relationships between plant chemodiversity and ant abundance will
likely depend on the role of the ants in the study system. Concerning

the **occurrence** of arthropods, higher chemodiversity likely favors arthropod occurrence by increasing the likelihood that a specific compound can be used as a host recognition cue (Lampert et al., 2010; Dyer, 2018; Jacobsen & Raguso, 2018). As a chemical profile is, by definition, a result of cumulative selection processes that have taken place over evolutionary history, compounds may serve unique or dual purposes depending on the interactions. Understanding how plant chemodiversity impacts arthropod communities will provide insights into the evolutionary forces shaping it.

Despite extensive research on plant chemodiversity's effects on arthropod communities, gaps remain. Studies often test only one arthropod group, focus on short-term effects, or compare high vs. low chemodiversity without examining a gradient. Our study addresses these gaps by manipulating the chemodiversity gradient and observing the responses of different arthropod groups over several seasons.

We used *Tanacetum vulgare* L. (tansy), an herbaceous plant with high intraspecific variation in terpenoids, to investigate how variation in the number of tansy chemotypes in a plot affects associated arthropod communities. In a field experiment with 84 plots (six plants each), we manipulated plot-level chemotype richness (CR; i.e., 1, 2, 3, or 6 chemotypes present in a plot) and studied its effects on herbivores, flower visitors, predators, and ants (Fig. 1). We examined the effects of chemodiversity on arthropod occurrence and abundance at the plot level, recognizing that responses may differ across groups due to their distinct ecological roles and interactions with plants. Additionally, we investigated the effects of each of the six chemotypes on arthropod communities, anticipating differential responses based on compound attractiveness or repellence.

We specifically tested the following hypotheses:

H1: Plots with higher chemotype richness will have
increased probabilities of arthropod occurrence across
groups.

H2: Plots with higher chemotype richness will have
decreased herbivore and ant abundances but increased
predator and flower visitor abundances.

H3: The presence of certain chemotypes in a plot will have
specific effects on herbivores, predators, flower visitors, and
ants, potentially influencing their abundance and occurrence
patterns.

Materials and Methods

Experimental design

Tansy chemotypes were characterized based on their leaf terpenoid
compositions. We selected six chemotypes ('Athu-Bthu', 'Bthu-
high', 'Bthu-low', 'Chrys-acet', 'Mixed-high', and 'Mixed-low') with
three biological replicates each (Fig. S1-1). These chemotypes were
propagated via stem cuttings and utilized in a field experiment at the
Jena Experiment site in Germany in 2021. The experiment consisted
of 84 1m^2 plots, each containing six tansy plants. The plots were
designed to vary in chemotype richness levels ranging from 1 to 6
(Fig. S1-2), with 12, 30, 30, and 12 replicate plots, respectively, for
each level. Each plot's chemotype composition was pre-assigned,
ensuring biological replication of each chemotype combination
across the experiment. Plots were distributed equally among six
randomized blocks, with specific numbers of plots allocated per

chemotype richness level within each block (more detail in: Ojeda-
Prieto et al., 2024)

Insect sampling

From 2021 to 2023, we assessed herbivores, flower visitors,
predators, and ants. Specific sampling dates are shown in Table S1-
1. Each year, herbivore communities were monitored by counting
individual aphids on the plants from their first appearance until
nearly none were found. Although the focus was on aphids, other
herbivores were scarce. Therefore, the generalization to herbivores
here reflects predominantly aphids. Counts were conducted 13 times
from June to August 2021, eight times from May to August 2022,
and three times from June to August 2023. Each count, executed
within a day between 7h and 19h, involved a meticulous scan of
each plant, which started from the tip of each stem and proceeded to
the end, examining both sides of each leaf. Identification of aphid
species followed the key provided in Supplementary Information 1
(S1). When multiple aphid species were present on a plant, we
recorded the total number of each. In cases of high aphid densities
(>100), we estimated the total by counting aphids on 1 cm of stem
or 1 cm^2 of leaf three times in different stems and leaves for each
plant, then multiplying the average by the estimated lengths of stems
(in cm) or the estimated areas of leaves (in cm^2) colonized by the
aphids. We used plot-level occupancy and cumulative plot-level
number of aphids to analyze occurrence and abundance.

Flower visitors were assessed by counting the flower heads of
individual plants during the flower-visiting recording. Flower
visitors were counted three times in July 2021 and once in August
2021. Surveys lasted three minutes per plot and were conducted in
random block order from 9h to 15h under optimal weather

conditions (no gusts of wind or precipitation). Flower visitors were

identified at the order level. We used presence absence data to

analyze the effects of CR and chemotype presence on occurrence.

However, as CR and individual chemotypes influence reproductive

traits in our system (Ojeda-Prieto et al., 2024), flower visitor

abundance per flower head (visitation rate) assessed effects on

abundance.

Predatory arthropods and ants on individual plants were recorded

during herbivore counting for 2021 and 2022 and the day after in

2023. Predators were classified into taxa/feeding guilds (Araneae,

Coleoptera, Dermaptera, Diptera, Hemiptera, and Hymenoptera),

specifying aphid parasitoids and parasitized aphids (counted as

mummies found). Ant presence on plants, but not abundance, was

noted. Predator and ant occurrences and parasitized aphid

abundance were used for analyses for the three seasons. In 2023,

predator and ant abundances were also analyzed.

Statistical analyses

Statistical analyses were conducted using R version 4.1.2 and

RStudio 2023.09.1 (R Core Team, 2021). All statistics were

performed at the plot level. The effects of chemotype presence and

CR on the occurrence of herbivores, flower visitors, predators, and

ants were assessed through binomial Generalized Linear Mixed-

effect Models (GLMM) using the '*lme4*' package (Bates et al.,

2015). Linear Mixed-Effect Models (LMM) and Zero-Inflated

Negative Binomial Models (ZINB), using the "*glmmTMB*" package

(Brooks et al., 2017), modeled the effect on abundances. P-values

were estimated by type II Wald-Chi-Squared tests using the Anova()

function ('*car*' package; Fox & Weisberg, 2019). Data visualization

used academic-licensed Biorender® and '*ggplot2*' (Wickham,

2009), *'grid'* (R Core Team, 2021)*, 'gridExtra'* (Auguie et al.,
2017), *'ggpubr'* (Kassambara, 2020), and *'DHARMa'* (Hartig,
2022) packages.

Effects of chemotype richness and chemotype presence on arthropod
taxa occurrence and abundance

Binomial GLMMs assessed the effect of CR and chemotype
presence on occurrence. LMMs and ZINB tested the effects on
abundance. Data were analyzed at the arthropod group and lower
taxonomic levels (when indicated). Data were analyzed
cumulatively for each year and for each independent sampling time.
Detailed model conditions and transformations for each group are
given in Supplementary Information 1 (S1).

Results

During our sampling campaigns, we registered the following
arthropod numbers: For 2021, 166.630 herbivores in total for 13
sampling dates (58.550 *Aphis fabae* (Scopoli, 1763)*,* 22.691
Brachycaudus cardui (Linnaeus, 1758), 382 *Macrosiphoniella*
tanacetaria (Kaltenbach, 1843), 62.055 *Metopeurum fuscoviride*
(Stroyan, 1950), and 22.952 *Uroleucon tanaceti* (Linnaeus, 1758)),
974 flower visitors in four sampling dates (316 Coleoptera, 236
Diptera, 134 Hemiptera, 237 Hymenoptera, and 51 Lepidoptera),
and 4.233 parasitized aphids in 10 sampling dates. For nine
sampling dates in 2022, 11'171.066 herbivores (1.610 *A. fabae,*
3.321 *B. cardui*, 35.902 *Ma. tanacetaria*, 37.196 *Me. fuscoviride*,
and >1.1 Mio *U. tanaceti*), and 341 parasitized aphids. For three
sampling dates in 2023, 234.386 herbivores (249 *A. fabae,* 217 *B.*
cardui, 0 *Ma. tanacetaria*, 9.535 *Me. fuscoviride*, and 224.385 *U.*
tanaceti), 9.940 predators (1.573 Araneae, 4.770 Coleoptera, 508

Dermaptera, 1.032 Diptera, 531 Hemiptera, 5.309 Hymenoptera)
and 3.783 ants (*Lasius niger* (Linnaeus, 1758)).

Seasonal patterns in the herbivore community

Phloem feeders heavily dominated the free-living insect herbivore
community in the tansy model system, whereas chewing herbivores
were largely absent. We observed two generalist aphid species in
our field study (*A. fabae* and *B. cardui*) and three tansy-specialist
aphid species (*Ma. tanacetaria, Me. fuscoviride,* and *U. tanaceti)*.
Our observations revealed the species-specific seasonal pattern of
aphid populations. Populations grew exponentially during the
warmer months of June and July and collapsed in August each year
(Fig. S1-2). However, the timing and height of peak abundance
varied among the species. The two generalist species (*A. fabae* and
B. cardui) exhibited a similar pattern; they were the first to colonize
the tansy plants and reached peak abundance in early July 2021 (Fig.
S1-2a). However, they were much less abundant in 2022, with a
peak occurring at the end of May. Conversely, the specialist species
were less abundant in 2021, reaching their peak in mid-June for *Ma.*
tanacetaria and in mid-July for *Me. fuscoviride* and *U. tanaceti.* In
2022, these species were more abundant, displacing the generalist
species, and reached peak abundance in mid to late June (Fig. S1-
2b). Generalist aphid species were even less common in the field in
2023, while *U. tanaceti* dominated the plots with increased
abundance in July (Fig. S1-2c).

Effects of chemotype richness and chemotype on herbivore
occurrence and abundance

Our study first investigated the effect of CR on plot-level I)
herbivore occurrence and II) herbivore abundance across samplings

in each experimental year. All plots were occupied by at least one

herbivore each year, but the herbivore species-specific occurrence

in plots varied. For instance, *Me. fuscoviride* was present in all plots

in 2021 (Table S1-2), for *U. tanaceti* this was the case in 2022 (Table

S1-3) and 2023 (Table S1-4). The only significant effect of CR on

aphid occurrence was observed for *Me. fuscoviride* in 2023, which

decreased when CR increased (X^2_1=9.42, P=0.002).

We then examined how specific tansy chemotypes influenced

herbivore occurrence at the plot level using individual models for

each chemotype (S1). Certain chemotypes were associated with

higher or lower likelihoods of colonization, i.e., occurrence, and

higher or lower herbivore abundance, depending on the year and

herbivore species (Table S1-11). *A. fabae* occurrence in 2022 was

higher when 'Bthu-low' (X^2_1=4.50, P=0.034) or 'Mixed-high' was

in the plot (X^2_1=6.68, P=0.010), but lower when 'Mixed-low' was

in the plot (X^2_1=4.89, P=0.027). In 2023, it was lower when 'Bthu-

low' (X^2_1=4.50, P=0.034) or 'Mixed-low' (X^2_1=4.89, P=0.027) was

present and higher when 'Mixed-high' was present (X^2_1=6.68,

P=0.010). Certain chemotypes within the plot affected *B. cardui*

occurrence in 2022; 'Bthu-high' increased *B. cardui* occurrence

(X^2_1=4.16, P=0.041), while 'Bthu-low' decreased it (X^2_1=6.42,

P=0.011). *Ma. tanacetaria* occurrence was not affected by any

chemotype. *Me. fuscoviride* occurrence in 2023 decreased when

'Athu-Bthu' (X^2_1=5.23, P=0.022) or 'Mixed-high' (X^2_1=9.42,

P=0.002) was present in the plot. *U. tanaceti* occurrence in 2021

decreased when 'Mixed-high' was present (X^2_1=7.47, P=0.002).

Concerning abundance, there was no effect of CR on cumulative

aphid abundance across all species (X^2_1=0.29, P=0.588) nor within

each of the five species in the first year (*A. fabae:* X^2_1=0.03,

P=0.870; *B. cardui:* X^2_1=1.51, P=0.219; *Ma. tanacetaria:* X^2_1=1.13,

P=0.288; *Me. fuscoviride:* $X^2_1=0.52$, P=0.471; *U. tanaceti:* $X^2_1=0.38$,

P=0.536; Fig. 3a, Table S1-2). However, increasing CR negatively

affected herbivore abundance in the following years (2022:

$X^2_1=7.63$, P=0.006; 2023: $X^2_1=4.39$, P=0.036; Fig. 2a, Tables S1-3,

S1-4), mainly driven by the negative effect of CR on *U. tanaceti*

populations in 2022 ($X^2_1=5.93$, p=0.015) and on *U. tanaceti*

($X^2_1=4.44$, P=0.035) and *B. cardui* abundances in 2023 ($X^2_1=5.56$,

P=0.018; Fig. 3a). Notably, *Ma. tanacetaria* aphids were not

observed in the field in 2023 at all. Plot-level chemotypic diversity

had little effect on whether a plot was occupied by aphids during the

year, mainly because all plots were occupied. However, there were

negative effects on the abundance of several herbivore species.

If there was an effect of a specific chemotype present in a plot on

herbivore abundance in the year 2021, it was always positive.

However, all significant effects of particular chemotypes observed

in 2022 or 2023 were negative (Table S1-11). Herbivore abundance

was lower in 2022 in either 'Chrys-acet' or 'Mixed-high' presence

and lower in 2023 when 'Athu-Bthu' or 'Mixed-low' was in the plot

(Tables S1-3 and S1-4). When investigating species-specific effects,

we found that *A. fabae* abundance in 2021 increased when 'Mixed-

high' was present ($X^2_1=6.77$, P=0.009), *B. cardui* when either

'Bthu-high' ($X^2_1=3.97$, P=0.046) or 'Chrys-acet' were present

($X^2_1=6.16$, P=0.013), and *U. tanaceti* when 'Bthu-high' was in the

plot ($X^2_1=5.98$, P=0.014). Lower *A. fabae* abundance was observed

in 2022 when 'Bthu-low' was in the plot ($X^2_1=3.91$, P=0.048) and

in 2023 when 'Athu-Bthu' ($X^2_1=7.39$, P=0.006) or 'Chrys-acet'

($X^2_1=19.14$, P<0.001) were part of the community. *Me. fuscoviride*

abundance in 2022 was negatively associated with 'Bthu-low'

($X^2_1=5.85$, P=0.016), and *U. tanaceti* abundance in 2022 was

negatively associated with either 'Chrys-acet' ($X^2_1=12.77$, P<0.001)

or 'Mixed-high' (X^2_1=5.58, P=0.018). In 2023, *B. cardui* was lower

when 'Athu-Bthu', 'Bthu-high', or 'Mixed-high' was present (Athu-

Bthu: X^2_1=4.42, P=0.035; Bthu-high: X^2_1=40.71, P<0.001; Mixed-

high: X^2_1=79.26, P<0.001), and *U. tanaceti* abundance was

negatively associated to the presence of 'Athu-Bthu' (X^2_1=5.08,

P=0.024) or 'Mixed-low' (X^2_1=3.92, P=0.048). The effects of

specific chemotypes on herbivore abundance varied across different

years, with positive impacts observed in 2021 and predominantly

negative impacts in 2022 and 2023.

Effects of chemotype richness and chemotype identity on flower

visitors

We first analyzed the effect of CR on flower visitor occurrence.

Higher CR increased the occurrence of Lepidoptera, i.e., adult

butterflies (X^2_1=3.87, P=0.049), but not other taxa. Hymenoptera

and Lepidoptera flower visitor occurrences increased when 'Bthu-

low' and 'Athu-Bthu' were in the plot, respectively (Tables S1-5,

S1-11).

We evaluated the effect of CR on the visitation rate of flower

visitors. We found a positive effect on all orders (X^2_1=10.71,

P=0.001, Fig. 2b), and specifically for Coleoptera (X^2_1=16.78,

P<0.001), Hymenoptera (X^2_1=11.56, P<0.001), and Lepidoptera

(X^2_1=8.80, P=0.003, Fig. 3b). Concerning the presence of particular

chemotypes within plots, 'Chrys-acet' (X^2_1=5.98, P=0.014) and

'Mixed-high' (X^2_1=6.95, P=0.008) positively affected the visitation

rate across all the orders. While Diptera and Hemiptera abundances

were not influenced by the presence/absence of any chemotypes,

Coleoptera was influenced by the presence of 'Bthu-low' and

'Chrys-acet' (Table S1-5); Hymenoptera by 'Bthu-low' (X^2_1=7.58,

P=0.006), and Lepidoptera by 'Athu-Bthu' (X^2_1=7.53, P=0.006,
Table S1-12).

Analysis of the effect on the total number of individuals (not the
visitation rate) is in Table S1-6. We found positive effects on total
number of individuals of all orders (X^2_1=10.71, P=0.001),
Coleoptera (X^2_1=16.78, P<0.001), Hymenoptera (X^2_1=11.56,
P<0.001), and Lepidoptera (X^2_1=8.80, P=0.003), as found by the
analysis of CR on visitation rate. However, when using the total
number as the response variable, particular chemotypes had
different effects on different groups. The abundance of all orders
was affected by the presence of any of the chemotypes except for
'Chrys-acet', 'Athu-Bthu' affected the number of flower visitors
belonging to Coleoptera and Lepidoptera, 'Bthu-low' to Coleoptera
and Hemiptera, and 'Bthu-high' and 'Mixed-low' to Coleoptera
(Table S1-6).

Effects of chemotype richness and chemotype on predators and ants

Predatory arthropods occurred in all plots in all years (Tables S1-7
to S1-9), and ants (*L. niger*) occurred in all plots in 2023 (Table S1-
10). The only effect of CR on predator taxa/feeding guild occurrence
was observed in predatory Hemiptera in 2022 (X^2_1=6.54, P=0.010),
which increased with CR. For chemotype identity effects, the
occurrence of parasitized aphids in 2021 decreased when 'Mixed-
low' was in the plot and in 2022 when 'Chrys-acet' was present
(2021: X^2_1=4.71, P=0.030; 2022: X^2_1=5.55, P=0.018, Tables S1-7,
S1-8).

Increasing CR did not increase predator abundance (overall -Fig. 2c-
nor by taxa/feeding guild -Fig. 3c-). However, there was a positive
effect of 'Chrys-acet' presence on the number of parasitized aphids

in 2021 (X^2_1=6.30, P=0.012, Table S1-7) and ant abundance in 2023

(X^2_1=6.58, P=0.010, Table S1-8), but a negative effect on total

predator abundance in 2023 (X^2_1=4.16, P=0.041) and on abundance

of predatory beetles in 2023 (Coleoptera: X^2_1=3.85, P=0.050, Table

S1-9). A positive effect on Dermaptera abundance in 2023 was

found when 'Athu-Bthu' was present (X^2_1=8.51, P=0.004).

Temporal effects

Analyses at independent sampling times, available in

Supplementary Information 1 (S1), revealed that while relationships

between chemotype richness and the occurrence of various

ecological groups were generally weak, the temporal patterns in

abundance were clear. Effect on abundance showed a shift in slope

from positive to negative for herbivores over time (particularly in

specialist aphids); while it remained mostly positive relationships

for flower visitors, ants, and predators (Supplementary Information

2).

Discussion

Intraspecific chemodiversity is a key driver of interaction variation

within plant species. Our understanding mainly comes from

observational and correlative studies, with controlled studies in

natural or semi-natural settings limited due to the challenges of

manipulating plant chemistry. In this study, we manipulated plant

chemical diversity at the plot level by selecting plant chemotypes

with varying chemical blends and manipulating the number of

chemotypes within plots (i.e., plot-level chemotype richness -CR-).

Using tansy (*Tanacetum vulgare* L.), we explored the effect of

chemodiversity on natural arthropod communities. Our study

provides evidence that intraspecific chemodiversity affects

processes at the plot level. We showed that CR had many negative

effects on herbivores but positive effects on flower visitors, with

minimal effects on predators or ants. Additionally, there were strong

effects of chemotype identity, i.e., the composition of the plots, on

plant-arthropod interactions at the plot level, an effect also often

found in biodiversity experiments (e.g., Scherber et al., 2010). Our

study thus highlights the role of individual chemotypes in shaping

interactions, contributing to our understanding of the ecological

significance of chemical diversity in plants.

Effects of chemotype richness on the occurrence of arthropod

groups

Contrary to our expectations, we did not find strong evidence that

CR affects plot-level arthropod occurrences **(H1)**. While some

significant relationships between CR and specific taxa occurrence

were observed, most were weak. One reason may be that many

arthropod taxa occurred in all plots or were absent during certain

samplings, limiting the variation needed to test hypotheses. Another

reason is that specialized herbivore populations build up over the

season before they decline, and in years when aphids are abundant,

total population sizes are very high, and almost all plants are

infested. We observed significant positive relationships for

polyphagous aphid species such as *A. fabae* and *B. cardui* in the

individual samplings during the first sampling year. These species

only feed on tansy for several weeks. Significant negative

relationships were observed for monophagous species, particularly

in *Me. fuscoviride* in 2023, suggesting that polyphagous and

monophagous species may respond differently to chemical cues in

their host-seeking behavior (Gripenberg et al., 2010; Benedek et al.,

2015; Ziaja & Müller, 2023). It also emphasizes the strong shift

from polyphagous to monophagous herbivores in the system over

the season. It could be that during the establishment phase,
polyphagous herbivores were more abundant in the surrounding
grassland and that monophagous species required time to colonize
from the nearest tansy hosts. All specialized tansy aphids overwinter
on or next to the host plant; hence, once the field plots are colonized,
local overwintering and early colonization in the season by the
specialized aphids are possible.

For flower-visiting taxa, we consistently observed positive
relationships between CR and occurrence, indicating the role of
specialized chemistry in these mutualists (Silva et al., 2018; Farré-
Armengol et al., 2020; Sasidharan et al., 2023). In contrast, the
relationship between CR and other groups' occurrence was weak.
These findings suggest that plant chemistry differentially affects
arthropod groups based on their interaction with the plant. While
plant chemistry significantly influences directly interacting
organisms like flower visitors, it does not show a similar significant
impact on indirectly interacting organisms such as predators and
ants (Ali et al., 2023).

Effects of chemotype richness on the abundance of arthropod taxa

We found strong support for CR affecting the abundance of
herbivores and flower-visiting arthropods but not predatory
arthropods and ants, partially in line with **(H2)**. Higher CR plots
supported fewer herbivores in 2022 and 2023, although this
relationship varied between aphid species. In general, the effects of
CR on herbivore abundance were driven by its effects on the
dominant monophagous aphid species, *U. tanaceti*, for both years
and *B. cardui* for 2023. An interesting observation is that the slope
between CR and herbivore abundance, despite the relationships not
always being significant, started consistently positive in 2021 but

turned increasingly negative over the first season and across the

years. It might be that the shift towards monophagous aphids over

time emphasized the negative effect of CR on herbivore abundance.

Since *U. tanaceti* had not yet successfully colonized the experiment

in the first year, and polyphagous species were more common, this

may explain the lack of CR effects in the first year. Monophagous

herbivores may be more strongly affected by chemical defenses in

high CR (Ziaja & Müller, 2023) in concordance with the

neighborhood-mediated associational resistance hypothesis

(Randlkofer et al., 2010).

Higher CR plots had more flower visitors, with significant positive

effects observed for Coleoptera, Hymenoptera, and Lepidoptera but

not for Diptera and Hemiptera, suggesting that CR may

differentially affect flower visitors (Sasidharan et al., 2023).

Chemotypes exhibit variations in their phenology, influenced by the

chemical diversity of their growth environment (Ojeda-Prieto et al.,

2024). These variations can extend the flowering period within a

plot containing a greater number of different chemotypes.

Consequently, CR will affect the occurrence and abundance of

flower visitors.

Our study revealed that the chemodiversity of the plot did not affect

predator or ant abundance. The precise reasons underlying these

findings remain unclear. However, it is plausible that additional

factors, such as prey availability, which may be influenced by

herbivore-host chemical composition (Benedek et al., 2019), or

alternative foraging strategies (Wäschke et al., 2013), contribute to

shaping these patterns. Further research is warranted to elucidate the

interplay between these factors and their collective impact on

predator and ant abundance in response to chemodiversity.

	Effects of chemotype presence on arthropod communities

We found chemotype-specific impacts on plot-level occurrence and
abundance across arthropod taxa, in line with our hypothesis **(H3)**.
In herbivores, patterns were rather incongruent and differed strongly
with the year of sampling. Herbivore occurrence data showed few
chemotype-specific effects in 2021, the first season after
establishment. Different chemotypes positively and negatively
affected the species observed in the following two years. This
suggests that most aphids seem to respond to plant-chemical cues as
expected (Mehrparvar et al., 2014; Clancy et al., 2018). Herbivore
abundance data show a clear pattern, where in 2021, chemotype
effects were typically positive, which may suggest that colonizers
were using specific terpenoid cues initially. Interestingly, in the
subsequent years, chemotype-specific effects, when significant,
were always negative. A possible scenario suggests that chemotypes
may serve as a mosaic that partly regulates aphid densities, which
aligns with the idea that specialized chemistry evolved with a
defensive function (Fox, 1981; Wolf et al., 2011). This idea is
further reinforced by the fact that the overall effect of chemotypes
on herbivore abundance tends to be neutral to negative.

For flower-visiting arthropods, chemotype-specific patterns were
most common in abundance data. Most flower visitors are highly
mobile and do not linger in the same plot for long (Burdon et al.,
2020; Knauer et al., 2021), suggesting that specific chemotypes lead
to higher abundance in certain plots. Significant chemotypes
positively affected flower visitors, with low-concentration B-
thujone chemotypes having the most robust and consistent effects
on different taxa, aligning partially with previous results
(Sasidharan et al., 2023). Mixed chemotypes had few effects, and
high-concentration B-thujone chemotype did not affect flower

visitors. Although B-thujone can signal nectar or pollen rewards
(Metcalf & Kogan, 1987), its effects depend on the overall blend of
volatile compounds and the preferences of local flower-visiting
species. We found stronger effects of chemotype identity on the total
number of individuals than on the visitation rate (Tables S1-5 and
S1-6). CR and individual chemotypes influence reproductive traits
in our system (Ojeda-Prieto et al., 2024). These effects might be
confounded by the influence of CR and chemotype on reproductive
traits rather than reflecting true effects on flower visitors.

Fewer and weaker chemotype-specific effects were found for
predators and ants, suggesting that these groups could be influenced
by different mechanisms, including modified behaviors, such as
predation activity or aphid-tending in ants (Senft et al., 2017;
Mehrparvar et al., 2018; Hauri et al., 2022).

Future perspectives for chemodiversity research

Exploring several key areas of chemodiversity will deepen our
understanding. Scaling up experiments to entire landscapes will
elucidate how chemodiversity influences ecosystems and
biodiversity across diverse habitats. Assessing its effect on plant-
insect interactions should include ecosystem functions like
predation rates and herbivore performance. Studying how
chemodiversity influences herbivore behavior, including responses
from generalist and specialist species, will offer insights with
potential applications in pest management. Further exploring the
interplay between plant chemistry and arthropod communities will
advance our understanding of ecosystem functioning and guide
conservation and management practices.

Conclusion

Previous studies have focused on intraspecific variability in ecosystem functioning. Our research demonstrates that plot-level chemotypic diversity impacts arthropod colonization patterns. Our results emphasize the role of intraspecific chemodiversity in shaping arthropod communities on plants. Our work shows that plots with higher intraspecific chemodiversity accumulate fewer herbivores and attract more flower visitors. The presence of specific chemotypes within a plot exerted highly variable impacts on abundance but also on occurrence patterns of various arthropod groups, suggesting that specific chemotypes, and potentially specific components they produce, may be a driver of the occurrence of arthropod taxa.

Acknowledgments

We thank the personnel at TUM PTC Dürnast, especially Sabine Zuber and Petra Scheuerer, for their work during the plant propagation and preparation phase. We thank Caroline Müller and Elisabeth Eilers for their contributions to chemotyping the plants, and to Anne Ebeling and Nico Eisenhauer for granting us access to core facilities within the Jena Experiment, facilitated under the guidance of Gerlinde Kratzsch. We acknowledge the assistance of Nafiseh Mahdavi in establishing the field experiment, and we are thankful to the Jena Experiment gardening staff for their guidance and support in maintaining the experiment. Special thanks are extended to Beate Piecha, Annika Neuhaus-Harr, and other friends and colleagues who assisted during the harvests. This study received funding from the German Research Foundation (DFG) under project 415496540 (WE3081/40-1) as part of the Research Unit (RU) FOR 3000, and we are grateful to all members of the RU for their valuable discussions.

Competing interests

All authors declare no conflict of interest to disclose.

Author contributions

WWW conceived the original idea and designed the experiment. Research conducted by: RH prepared the field experiment and propagation of plants. LOP, RH, and WWW planted the field experiment. LOP and ELM collected all data and maintained the field experiment. LOP analyzed ecological data with input from RH and WWW. LOP wrote the manuscript with input from ELM, RH, and WWW. All authors contributed to the final version and approved the final submission.

Data availability

Data and R code scripts supporting the results are archived in Dryad.

References

Fox, L. R. (1981). Defense and dynamics in plant-herbivore systems. *American Zoologist*, *21*(4), 853–864. https://doi.org/10.1093/icb/21.4.853

Metcalf, R. L., & Kogan, M. (1987). Plant volatiles as insect attractants. *Critical Reviews in Plant Sciences*, *5.3*, 251–301.

Schulze, E. D., & Mooney, H. A. (1992). *Biodiversity and ecosystem function*. Ecol Stud, Heidelberg: Springer.

De Boer, J. G., Posthumus, M. A., & Dicke, M. (2004).

Identification of volatiles that are used in discrimination

between plants infested with prey or nonprey herbivores by

a predatory mite. *Journal of Chemical Ecology*, *30*(11),

2215–2230.

https://doi.org/10.1023/B:JOEC.0000048784.79031.5e

Cardinale, B. J., Srivastava, D. S., Emmett Duffy, J., Wright, J. P.,

Downing, A. L., Sankaran, M., & Jouseau, C. (2006). Effects

of biodiversity on the functioning of trophic groups and

ecosystems.   *Nature*,   *443*(7114),   989–992.

https://doi.org/10.1038/nature05202

Crutsinger, G. M., Collins, M. D., Fordyce, J. A., Gompert, Z., Nice,

C. C., & Sanders, N. J. (2006). Plant genotypic diversity

predicts community structure and governs an ecosystem

process.   *Science*,   *313*(5789),   966–968.

https://doi.org/10.1126/science.1128326

Hartmann, T. (2007). From waste products to ecochemicals: Fifty

years   research   of   plant   secondary   metabolism.

Phytochemistry,   *68*(22–24),   2831–2846.

https://doi.org/10.1016/j.phytochem.2007.09.017

De Boer, J. G., Hordijk, C. A., Posthumus, M. A., & Dicke, M.

(2008). Prey and non-prey arthropods sharing a host plant:

Effects on induced volatile emission and predator attraction.

Journal of Chemical Ecology, *34*(3), 281–290.

https://doi.org/10.1007/s10886-007-9405-z

Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A., &

Szendrei, Z. (2009). Associational resistance and

associational susceptibility: Having right or wrong

neighbors. *Annual Review of Ecology, Evolution, and*

Systematics, *40*(1), 1–20.

https://doi.org/10.1146/annurev.ecolsys.110308.120242

Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis*.

Springer New York. https://doi.org/10.1007/978-0-387-

98141-3

Gripenberg, S., Mayhew, P. J., Parnell, M., & Roslin, T. (2010). A

meta-analysis of preference–performance relationships in

phytophagous insects. *Ecology Letters*, *13*(3), 383–393.

https://doi.org/10.1111/j.1461-0248.2009.01433.x

Lampert, E. C., Dyer, L. A., & Bowers, M. D. (2010). Caterpillar

chemical defense and parasitoid success: *Cotesia*

congregata parasitism of *Ceratomia catalpae*. *Journal of*

Chemical Ecology, *36*(9), 992–998.

https://doi.org/10.1007/s10886-010-9840-0

Scherber, C., Eisenhauer, N., Weisser, W. W., Schmid, B., Voigt,

W., Fischer, M., Schulze, E.-D., Roscher, C., Weigelt, A.,

Allan, E., Beßler, H., Bonkowski, M., Buchmann, N.,

Buscot, F., Clement, L. W., Ebeling, A., Engels, C., Halle,

S., Kertscher, I., … Tscharntke, T. (2010). Bottom-up effects

of plant diversity on multitrophic interactions in a

biodiversity experiment. *Nature*, *468*(7323), 553–556.

https://doi.org/10.1038/nature09492

Eisenhauer, N., Milcu, A., Sabais, A. C. W., Bessler, H., Brenner,

J., Engels, C., Klarner, B., Maraun, M., Partsch, S., Roscher,

C., Schonert, F., Temperton, V. M., Thomisch, K., Weigelt,

A., Weisser, W. W., & Scheu, S. (2011). Plant diversity

surpasses plant functional groups and plant productivity as

driver of soil biota in the long term. *PLoS ONE*, *6*(1),

e16055. https://doi.org/10.1371/journal.pone.0016055

Hare, J. D. (2011). Ecological role of volatiles produced by plants

in response to damage by herbivorous insects. *Annual

Review   of   Entomology*,   *56*(1),   161–180.

https://doi.org/10.1146/annurev-ento-120709-144753

Kleine, S., & Müller, C. (2011). Intraspecific plant chemical

diversity and its relation to herbivory. *Oecologia*, *166*(1),

175–186. https://doi.org/10.1007/s00442-010-1827-6

Kos, M., Broekgaarden, C., Kabouw, P., Oude Lenferink, K.,

Poelman, E. H., Vet, L. E. M., Dicke, M., & Van Loon, J. J.

A. (2011). Relative importance of plant-mediated bottom-up

and top-down forces on herbivore abundance on *Brassica*

oleracea: Bottom-up and top-down effects on herbivores.

Functional Ecology, *25*(5), 1113–1124.

https://doi.org/10.1111/j.1365-2435.2011.01871.x

Wolf, V. C., Berger, U., Gassmann, A., & Müller, C. (2011). High

chemical diversity of a plant species is accompanied by

increased chemical defence in invasive populations.

Biological Invasions, *13*(9), 2091–2102.

https://doi.org/10.1007/s10530-011-0028-5

Clavijo McCormick, A., Unsicker, S. B., & Gershenzon, J. (2012).

The specificity of herbivore-induced plant volatiles in

attracting herbivore enemies. *Trends in Plant Science*, *17*(5),

303–310. https://doi.org/10.1016/j.tplants.2012.03.012

Wäschke, N., Meiners, T., & Rostás, M. (2013). Foraging strategies

of parasitoids in complex chemical environments. In E.

Wajnberg & S. Colazza (Eds.), *Chemical Ecology of Insect*

Parasitoids (1st ed., pp. 37–63). Wiley.

https://doi.org/10.1002/9781118409589.ch3

Mehrparvar, M., Mansouri, S. M., & Weisser, W. W. (2014).

Mechanisms of species-sorting: Effect of habitat occupancy

on aphids' host plant selection: Host selection by competing

herbivores. *Ecological Entomology*, *39*(3), 281–289.

https://doi.org/10.1111/een.12096

Mora, C., Danovaro, R., & Loreau, M. (2014). Alternative

hypotheses to explain why biodiversity-ecosystem

functioning relationships are concave-up in some natural

ecosystems but concave-down in manipulative experiments.

Scientific Reports, *4*(1), 5427.

https://doi.org/10.1038/srep05427

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting

linear mixed-effects models using lme4. *Journal of*

Statistical Software, *67*(1), 1–48.

Benedek, K., Bálint, J., Salamon, R. V., Kovács, E., Ábrahám, B.,

Fazakas, C., Loxdale, H. D., & Balog, A. (2015). Chemotype

of tansy (*Tanacetum vulgare* L.) determines aphid genotype

and its associated predator system. *Biological Journal of the*

Linnean Society, *114*(3), 709–719.

Bálint, J., Zytynska, S. E., Salamon, R. V., Mehrparvar, M.,

Weisser, W. W., Schmitz, O. J., Benedek, K., & Balog, A.

(2016). Intraspecific differences in plant chemotype

determine the structure of arthropod food webs. *Oecologia*,

180(3), 797–807. https://doi.org/10.1007/s00442-015-3508-

y

Auguie, B., Antonov, A., & Auguie, M. B. (2017). *gridExtra:*

Miscellaneous functions for "grid" graphics.

Brooks, M. E., Kristensen, K., Van Benthem, J., Magnusson, A.,

Berg, C. W., Nielsen, A., Skaug, H. J., Maechler, M., &

Bolker, B. M. (2017). glmmTMB balances speed and

flexibility among packages for zero-inflated generalized

linear mixed modeling. *The R Journal*, *9*(2), 378–400.

Bustos-Segura, C., Poelman, E. H., Reichelt, M., Gershenzon, J., &

Gols, R. (2017). Intraspecific chemical diversity among

neighbouring plants correlates positively with plant size and

herbivore load but negatively with herbivore damage.

Ecology Letters, *20*(1), 87–97.

https://doi.org/10.1111/ele.12713

Hertzog, L. R., Ebeling, A., Weisser, W. W., & Meyer, S. T. (2017).

Plant diversity increases predation by ground-dwelling

invertebrate predators. *Ecosphere*, *8*(11), e01990.

https://doi.org/10.1002/ecs2.1990

Meyer, S. T., Scheithe, L., Hertzog, L., Ebeling, A., Wagg, C.,

Roscher, C., & Weisser, W. W. (2017). Consistent increase

in herbivory along two experimental plant diversity

gradients over multiple years. *Ecosphere*, *8*(7), e01876.

https://doi.org/10.1002/ecs2.1876

Senft, M., Weisser, W. W., & Zytynska, S. E. (2017). Habitat

variation, mutualism and predation shape the spatio-

temporal dynamics of tansy aphids. *Ecological Entomology*,

42(4), 389–401. https://doi.org/10.1111/een.12396

Weisser, W. W., Roscher, C., Meyer, S. T., Ebeling, A., Luo, G.,

Allan, E., Beßler, H., Barnard, R. L., Buchmann, N., Buscot,

F., Engels, C., Fischer, C., Fischer, M., Gessler, A.,

Gleixner, G., Halle, S., Hildebrandt, A., Hillebrand, H., De

Kroon, H., … Eisenhauer, N. (2017). Biodiversity effects on

ecosystem functioning in a 15-year grassland experiment:

Patterns, mechanisms, and open questions. *Basic and*

Applied Ecology, *23*, 1–73.

https://doi.org/10.1016/j.baae.2017.06.002

Clancy, M. V., Zytynska, S. E., Moritz, F., Witting, M., Schmitt-

Kopplin, P., Weisser, W. W., & Schnitzler, J. (2018).

Metabotype variation in a field population of tansy plants

influences aphid host selection. *Plant, Cell & Environment*, *41*(12), 2791–2805. https://doi.org/10.1111/pce.13407

Dyer, L. A. (2018). Multidimensional diversity associated with plants: A view from a plant–insect interaction ecologist. *American Journal of Botany*, *105*(9), 1439–1442. https://doi.org/10.1002/ajb2.1147

Ebeling, A., Hines, J., Hertzog, L. R., Lange, M., Meyer, S. T., Simons, N. K., & Weisser, W. W. (2018). Plant diversity effects on arthropods and arthropod-dependent ecosystem functions in a biodiversity experiment. *Basic and Applied Ecology*, *26*, 50–63. https://doi.org/10.1016/j.baae.2017.09.014

Jacobsen, D. J., & Raguso, R. A. (2018). Lingering effects of herbivory and plant defenses on pollinators. *Current Biology*, *28*(19), R1164–R1169. https://doi.org/10.1016/j.cub.2018.08.010

Jakobs, R., & Müller, C. (2018). Effects of intraspecific and intra-individual differences in plant quality on preference and performance of monophagous aphid species. *Oecologia*, *186*(1), 173–184. https://doi.org/10.1007/s00442-017-3998-x

Koricheva, J., & Hayes, D. (2018). The relative importance of plant

intraspecific diversity in structuring arthropod communities:

A meta-analysis. *Functional Ecology*, *32*(7), 1704–1717.

https://doi.org/10.1111/1365-2435.13062

Mehrparvar, M., Zytynska, S. E., Balog, A., & Weisser, W. W.

(2018). Coexistence through mutualist-dependent reversal

of competitive hierarchies. *Ecology and Evolution*, *8*(2),

1247–1259. https://doi.org/10.1002/ece3.3689

Silva, R. F., Rabeschini, G. B. P., Peinado, G. L. R., Cosmo, L. G.,

Rezende, L. H. G., Murayama, R. K., & Pareja, M. (2018).

The ecology of plant chemistry and multi-species

interactions in diversified agroecosystems. *Frontiers in*

Plant   Science,   *9*,   1713.

https://doi.org/10.3389/fpls.2018.01713

Turlings, T. C. J., & Erb, M. (2018). Tritrophic interactions

mediated by herbivore-induced plant volatiles: Mechanisms,

ecological relevance, and application potential. *Annual*

Review   of   Entomology,   *63*(1),   433–452.

https://doi.org/10.1146/annurev-ento-020117-043507

Benedek, K., Bálint, J., Máthé, I., Mara, G., Felföldi, T., Szabó, A.,

Fazakas, C., Albert, C., Buchkowski, R. W., Schmitz, O. J.,

& Balog, A. (2019). Linking intraspecific variation in plant

chemical defence with arthropod and soil bacterial

community structure and N allocation. *Plant and Soil*,

444(1–2), 383–397. https://doi.org/10.1007/s11104-019-

04284-7

Fox, J., & Weisberg, S. (2019). *An {R} companion to applied*

regression (Third Edition). Thousand Oaks CA: Sage.

Giling, D. P., Ebeling, A., Eisenhauer, N., Meyer, S. T., Roscher,

C., Rzanny, M., Voigt, W., Weisser, W. W., & Hines, J.

(2019). Plant diversity alters the representation of motifs in

food webs. *Nature Communications*, *10*(1), 1226.

https://doi.org/10.1038/s41467-019-08856-0

Glassmire, A. E., Philbin, C., Richards, L. A., Jeffrey, C. S., Snook,

J. S., & Dyer, L. A. (2019). Proximity to canopy mediates

changes in the defensive chemistry and herbivore loads of an

understory tropical shrub, *Piper kelleyi*. *Ecology Letters*,

22(2), 332–341. https://doi.org/10.1111/ele.13194

Raffard, A., Santoul, F., Cucherousset, J., & Blanchet, S. (2019).

The community and ecosystem consequences of

intraspecific diversity: A meta-analysis. *Biological Reviews*,

94(2), 648–661. https://doi.org/10.1111/brv.12472

Senft, M., Clancy, M. V., Weisser, W. W., Schnitzler, J., &

Zytynska, S. E. (2019). Additive effects of plant chemotype,

mutualistic ants and predators on aphid performance and

survival. *Functional Ecology*, *33*(1), 139–151.

https://doi.org/10.1111/1365-2435.13227

Zytynska, S. E., Guenay, Y., Sturm, S., Clancy, M. V., Senft, M.,

Schnitzler, J., Dilip Pophaly, S., Wurmser, C., & Weisser,

W. W. (2019). Effect of plant chemical variation and

mutualistic ants on the local population genetic structure of

an aphid herbivore. *Journal of Animal Ecology*, *88*(7), 1089–

1099. https://doi.org/10.1111/1365-2656.12995

Dantas, E., do Valle Capelli, N., Pastori, T., Alencar, A. C., Cole, T.

C. H., & Demarco, D. (2020). Diversity of floral glands and

their secretions in pollinator attraction. In *Co-evolution of*

secondary metabolites (Springer, pp. 403–425).

Farré-Armengol, G., Fernández-Martínez, M., Filella, I., Junker, R.

R., & Peñuelas, J. (2020). Deciphering the biotic and

climatic factors that influence floral scents: A systematic

review of floral volatile emissions. *Frontiers in Plant*

Science, *11*, 1154. https://doi.org/10.3389/fpls.2020.01154

892 Kariñho-Betancourt, E. (2020). Coevolution: Plant-herbivore

893   interactions and secondary metabolites of plants. In *Co-*

894   *evolution of secondary metabolites* (pp. 47–76). Springer.

895 Kassambara, A. (2020). *ggpubr: "ggplot2" based publication ready*

896   *plots*.

897 Müller, C., Bräutigam, A., Eilers, E., Junker, R., Schnitzler, J.-P.,

898   Steppuhn, A., Unsicker, S., Van Dam, N., Weisser, W., &

899   Wittmann, M. (2020). Ecology and evolution of intraspecific

900   chemodiversity of plants. *Research Ideas and Outcomes*, *6*,

901   e49810. https://doi.org/10.3897/rio.6.e49810

902 Noto, A. E., & Hughes, A. R. (2020). Genotypic diversity weakens

903   competition within, but not between, plant species. *Journal*

904   *of Ecology*, *108*(6), 2212–2220.

905   https://doi.org/10.1111/1365-2745.13465

906 Wetzel, W. C., & Whitehead, S. R. (2020). The many dimensions of

907   phytochemical diversity: Linking theory to practice. *Ecology*

908   *Letters*, *23*(1), 16–32. https://doi.org/10.1111/ele.13422

909 Zu, P., Boege, K., del-Val, E., Schuman, M. C., Stevenson, P. C.,

910   Zaldivar-Riverón, A., & Saavedra, S. (2020). Information

911   arms race explains plant-herbivore chemical communication

in ecological communities. *Science*, *368*(6497), 1377–1381.

https://doi.org/10.1126/science.aba2965

Dyer, L. A., & Jeffrey, C. S. (2021). Chemically mediated multi-

trophic interactions. In K. Del-Claro & H. M. Torezan-

Silingardi (Eds.), *Plant-Animal Interactions* (pp. 17–38).

Springer International Publishing.

https://doi.org/10.1007/978-3-030-66877-8_2

Eisenring, M., Unsicker, S. B., & Lindroth, R. L. (2021). Spatial,

genetic and biotic factors shape within-crown leaf trait

variation and herbivore performance in a foundation tree

species. *Functional Ecology*, *35*(1), 54–66.

https://doi.org/10.1111/1365-2435.13699

Hauri, K. C., Glassmire, A. E., & Wetzel, W. C. (2021). Chemical

diversity rather than cultivar diversity predicts natural

enemy control of herbivore pests. *Ecological Applications*,

31(3), e02289. https://doi.org/10.1002/eap.2289

R Core Team. (2021). *R: A language and environment for statistical*

computing. R Foundation for Statistical Computing.

Schneider, G. F., Salazar, D., Hildreth, S. B., Helm, R. F., &

Whitehead, S. R. (2021). Comparative Metabolomics of

Fruits and Leaves in a Hyperdiverse Lineage Suggests Fruits

933   Are a Key Incubator of Phytochemical Diversification.

934   *Frontiers in Plant Science*, *12*, 693739.

935   https://doi.org/10.3389/fpls.2021.693739

936 Whitehead, S. R., Bass, E., Corrigan, A., Kessler, A., & Poveda, K.

937   (2021). Interaction diversity explains the maintenance of

938   phytochemical diversity. *Ecology Letters*, *24*(6), 1205–

939   1214. https://doi.org/10.1111/ele.13736

940 Yamawo, A., Suzuki, N., & Tagawa, J. (2021). Species diversity and

941   biological trait function: Effectiveness of ant–plant

942   mutualism decreases as ant species diversity increases.

943   *Functional Ecology*, *35*(9), 2012–2025.

944   https://doi.org/10.1111/1365-2435.13859

945 Espinosa-García, F. J. (2022). The role of phytochemical diversity

946   in the management of agroecosystems. *Botanical Sciences*,

947   *100*(Special), 245–262.

948   https://doi.org/10.17129/botsci.3075

949 Fernandez-Conradi, P., Defossez, E., Delavallade, A., Descombes,

950   P., Pitteloud, C., Glauser, G., Pellissier, L., & Rasmann, S.

951   (2022). The effect of community-wide phytochemical

952   diversity on herbivory reverses from low to high elevation.

953   *Journal of Ecology*, *110*(1), 46–56.

954   https://doi.org/10.1111/1365-2745.13649

Hartig, F. (2022). *DHARMa: Residual diagnostics for hierarchical*
(multi-level / mixed) regression model. https://cran.r-
hub.io/web/packages/DHARMa/DHARMa.pdf

Hauri, K. C., Glassmire, A. E., Randall, B., Zehr, L. N., & Wetzel,
W. C. (2022). Plant chemical diversity and its frequency
have distinct but complementary effects on insect foraging.
Journal of Applied Ecology, 59(5), 1362–1371.
https://doi.org/10.1111/1365-2664.14151

Müller, C., & Junker, R. R. (2022). Chemical phenotype as
important and dynamic niche dimension of plants. *New*
Phytologist, 234(4), 1168–1174.
https://doi.org/10.1111/nph.18075

Ramos, L., Fagundes, M., Boanares, D., Fernandes, G. W., & Solar,
R. (2022). Experimental manipulation of biotic and abiotic
parameters changes the outcome of insect-plant interactions.
Basic and Applied Ecology, 65, 97–108.
https://doi.org/10.1016/j.baae.2022.04.002

Walker, T. W. N., Alexander, J. M., Allard, P., Baines, O., Baldy,
V., Bardgett, R. D., Capdevila, P., Coley, P. D., David, B.,
Defossez, E., Endara, M., Ernst, M., Fernandez, C.,
Forrister, D., Gargallo-Garriga, A., Jassey, V. E. J., Marr, S.,
Neumann, S., Pellissier, L., … Salguero-Gómez, R. (2022).

Functional Traits 2.0: The power of the metabolome for ecology. *Journal of Ecology*, *110*(1), 4–20. https://doi.org/10.1111/1365-2745.13826

Whitehead, S. R., Schneider, G. F., Dybzinski, R., Nelson, A. S., Gelambi, M., Jos, E., & Beckman, N. G. (2022). Fruits, frugivores, and the evolution of phytochemical diversity. *Oikos*, *2022*(2), oik.08332. https://doi.org/10.1111/oik.08332

Ali, M. Y., Naseem, T., Holopainen, J. K., Liu, T., Zhang, J., & Zhang, F. (2023). Tritrophic interactions among arthropod natural enemies, herbivores and plants considering volatile blends at different scale levels. *Cells*, *12*(2), 251. https://doi.org/10.3390/cells12020251

Dussarrat, T., Schweiger, R., Ziaja, D., Nguyen, T. T. N., Krause, L., Jakobs, R., Eilers, E. J., & Müller, C. (2023). Influences of chemotype and parental genotype on metabolic fingerprints of tansy plants uncovered by predictive metabolomics. *Scientific Reports*, *13*(1), 11645. https://doi.org/10.1038/s41598-023-38790-7

Petrén, H., Anaia, R. A., Aragam, K. S., Bräutigam, A., Eckert, S., Heinen, R., Jakobs, R., Ojeda-Prieto, L., Popp, M., Sasidharan, R., Schnitzler, J.-P., Steppuhn, A., Thon, F.,

Tschikin, S., Unsicker, S. B., Van Dam, N. M., Weisser, W.

W., Wittmann, M. J., Yepes, S., … Junker, R. R. (2023).

Understanding the phytochemical diversity of plants:

Quantification, variation and ecological function.

https://doi.org/10.1101/2023.03.23.533415

Rabelo, R. S., Dyer, L. A., Yamaguchi, L. F., Diniz, I., Simbaña,

W., Kussano, A. J. M., Kato, M. J., & Massad, T. J. (2023).

Plasticity in plant defense and the role of phytochemical

dissimilarity in limiting specialist herbivory. *Frontiers in*

Ecology and Evolution, *11*, 1175590.

https://doi.org/10.3389/fevo.2023.1175590

Sasidharan, R., Grond, S. G., Champion, S., Eilers, E. J., & Müller,

C. (2023). *Flower visitor groups show differential responses*

to individual and plot-level chemodiversity with

consequences for plant fitness.

https://doi.org/10.1101/2023.11.21.568056

Wetzel, W. C., Inouye, B. D., Hahn, P. G., Whitehead, S. R., &

Underwood, N. (2023). Variability in plant–herbivore

interactions. *Annual Review of Ecology, Evolution, and*

Systematics, *54*(1), 451–474.

https://doi.org/10.1146/annurev-ecolsys-102221-045015

Zerebecki, R. A., & Hughes, A. R. (2023). Ecological consequences

of genotypic diversity within a foundation plant, *Spartina*

alterniflora , are pervasive but not universal across multiple

stress gradients. *Journal of Ecology*, *111*(8), 1794–1810.

https://doi.org/10.1111/1365-2745.14147

Ziaja, D., & Müller, C. (2023). Intraspecific chemodiversity

provides plant individual- and neighbourhood-mediated

associational resistance towards aphids. *Frontiers in Plant*

Science,   *14*,   1145918.

https://doi.org/10.3389/fpls.2023.1145918

Govaert, L., Hendry, A. P., Fattahi, F., & Möst, M. (2024).

Quantifying interspecific and intraspecific diversity effects

on ecosystem functioning. *Ecology*, *105*(1), e4199.

https://doi.org/10.1002/ecy.4199

Konefal, A., Kirkland, A., Gilpin, R., Wyssmann, K., Anthony, N.

M., Cebrian, J., & Cox, T. E. (2024). The relationship

between genetic diversity, function, and stability in marine

foundation   species.   *BioScience*,   *74*(3),   187–206.

https://doi.org/10.1093/biosci/biad123

Neuhaus-Harr, A., Ojeda-Prieto, L., Eilers, E., Müller, C., Weisser,

W. W., & Heinen, R. (2024). Chemodiversity affects

preference for *Tanacetum vulgare* chemotypes in two aphid

species. *Oikos*, *2024*(3), e10437.

https://doi.org/10.1111/oik.10437

Ojeda-Prieto, L., Medina-van Berkum, P., Unsicker, S. B., Heinen,

R., & Weisser, W. W. (2024). Intraspecific chemical

variation of *Tanacetum vulgare* affects plant growth and

reproductive traits in field plant communities. *Plant Biology*,

plb.13646. https://doi.org/10.1111/plb.13646

1049
1050

Main Figures

Figure 1. Influence of *Tanacetum vulgare* intraspecific
chemodiversity on arthropod communities. The left panel
illustrates the gradient of plot-level chemotype richness, with each
color symbolizing a specific chemotype of *Tanacetum vulgare* in a
six-plant community. From levels 1 to 6, the chemodiversity
increases, indicating an increase in different chemotypes. The right
panel presents the evaluated arthropod communities: herbivores,
flower visitors, predators, and ants, which are expected to be
influenced by intraspecific chemodiversity. Created with
Biorender®. Agreement number: FE271K6HDD.

1062

Figure 2. Effect of plot-level chemotype richness on the
abundance of different arthropod communities over three years
(2021–2023). (a) Effect on herbivore abundance, represented by
five aphid species each year. **(b)** Effect on flower visitor abundance
(i.e., visitation rate), calculated as the cumulative count of flower-
visiting insects divided by the sum of flower heads counted during
each assessment date. **(c)** Effect on parasitized aphid abundance
(aphid mummies). **(d)** Effect on predator abundance. **(e)** Effect on
ant abundance. Significance is indicated as follows: n.s.=not
significant, * $P<0.05$, ** $P<0.01$, and *** $P<0.001$. Degrees of
freedom, Wald's Chi-square statistics, and p-values are reported in
Tables S1 – S8.

1075

Figure 3. Effect of plot-level chemotype richness on specific group abundances within each guild of arthropod communities over three years (2021–2023). (a) Effect on herbivore abundance of *Aphis fabae*, *Brachycaudus cardui*, *Macrosiphoniella tanacetaria*, *Metopeurum fuscoviride*, and *Uroleucon tanaceti* each year. **(b)** Effect on flower visitor abundance of Coleoptera, Diptera, Hemiptera, Hymenoptera, and Lepidoptera. **(c)** Effect on predator abundance of Araneae, Coleoptera, Dermaptera, Diptera, Hemiptera, Aphid parasitoids, and Other Hymenoptera for 2023. Significance is indicated as follows: n.s.=not significant, * P<0.05, ** P<0.01, and *** P<0.001. Degrees of freedom, Wald's Chi-square statistics, and p-values are reported in Tables S1 – S7.

Plot-level chemotype richness

Herbivores

Flower visitors

Predators

Ants