1

Polysaccharide quantification using microbial enzyme cocktails

Sammy Pontrelli[1], and Uwe Sauer[1]*

[1]Institute of Molecular Systems Biology, ETH Zürich, Zürich, Switzerland

*sauer@ethz.ch

Key words: Polysaccharide quantification, chitin, laminarin, marine enzymes, enzyme assay, enzyme cocktail, polysaccharide

Abstract:

Polysaccharide quantification is crucial for studying ecological and nutritional processes in microbes, plants, and animals, as well as their role in carbon cycling and sequestration in aquatic environments. Traditional methods hydrolyze polysaccharides into monomers for quantification but are limited to chemically hydrolysable polysaccharides or require characterized recombinant enzymes. Here we use undefined mixtures of enzymes that are secreted by marine microbes or seawater communities to hydrolyze polysaccharides, focusing on colloidal chitin and laminarin as model marine polysaccharides. Colloidal chitin was digested using an enzyme cocktail from a chitin-degrading *Psychromonas sp.* bacterial isolate. By quantifying mono- and oligomers with a 3,5-dinitrosalicylic acid reducing sugar assay and/or liquid chromatography-mass spectrometry, chitin could be quantified down to 62 mg/L and 15 mg/L, respectively, enabling us to monitor microbial chitin degradation. Laminarin was digested using enzymes from an undefined marine microbial community with a lower detection limit of 30 mg/L using a reducing sugar assay. Our findings demonstrate that individual microbes or undefined communities can be readily harnessed as enzyme sources, offering a flexible method for polysaccharide quantification that bypasses limitations in chemical hydrolysis and availability of characterized recombinant enzymes.

Introduction

Biological polysaccharides carry out vital roles, serving as energy storage molecules(Suzuki and Suzuki, 2013), structuring cell walls(Lerouxel et al., 2006) and biofilms(Raszka et al., 2006), and fueling carbon to microbial communities(Arnosti et al., 2021; Cerqueira et al., 2020; Enke et al., 2019). Their abundance in organic matter places them at the center of biogeochemical carbon cycling and carbon sequestration in terrestrial(Horwath, 2007) and aquatic environments(Arnosti et al., 2021). Quantifying polysaccharides is essential to study these biological processes, yet their diverse chemical properties make it difficult to develop universally applicable quantification methods.

The most common approaches to quantify polysaccharides involve hydrolysis into monosaccharides, followed by quantification using colorimetric or chromatographic techniques(Arnosti et al., 2021; Decamp et al., 2021; Schiavone et al., 2014). Typically, hydrolysis is performed with chemical or enzymatic methods. Acid hydrolysis presents an obstacle as different polysaccharides are hydrolyzed with varying efficiencies or not at all(Arnosti et al., 2021), and the harsh conditions can destroy acid-sensitive components(Cao et al., 2017; Lin and Guo, 2015). An important example for resistance to acid hydrolysis is chitin, a linear homopolymer of β-(1,4)-linked N-acetylglucosamine (GlcNAc) monomers(Peter, 1995). As the main constituent of mollusk and arthropod exoskeletons in marine environments(Keyhani and Roseman, 1999), it is one of the most abundant polysaccharides in the ocean. While chitin can be deacetylated into chitosan to enable acid hydrolysis to glucosamine(Yan and Evenocheck, 2012), this method is time-consuming and cannot be done in microtiter plates.

Enzymatic degradation enables the quantification of polysaccharides that cannot be hydrolyzed chemically(Becker et al., 2017; Schiavone et al., 2014; Steinke et al., 2022; Sun et al., 2024). However, a drawback is that identifying enzymes with desired activity may be

challenging, and enzyme expression and purification are time-consuming and may require

optimization. As an alternative, commercially produced enzymes may be used. However as in

the case of chitinases, they may be prohibitively expensive, restricting their accessibility.

A largely unexplored alternative is to use microbial enzyme cocktails. This involves growing

a polysaccharide-degrading bacterium or community on the target polysaccharide to induce

the expression of enzymes with desired hydrolytic activity. This mix of undefined secreted

enzymes, or cocktail, can be concentrated from the supernatant and used to digest a

polysaccharide followed by mono- or oligomer quantification. Enzyme cocktails have

previously been used to identify the monomeric composition of polysaccharide

substrates(Buck-Wiese et al., 2023; Sichert et al., 2020), and to identify the catalytic

capabilities(Pontrelli et al., 2022; Sichert et al., 2020; Unfried et al., 2018) or relative

activities(Pontrelli et al., 2022; Sichert et al., 2020) of secreted enzymes.

Here, we utilized enzyme cocktails for polysaccharide quantification, overcoming pitfalls of

chemical hydrolysis and limited recombinant enzyme availability. We collected an enzyme

cocktail from the chitin-degrading isolate, *Psychromonas sp. Psy*6C06, as a proof of concept

to quantify colloidal chitin. Additionally, we used seawater microbial communities to induce a

laminarin-degrading enzyme cocktail, which we further purified and used to quantify

laminarin. This method simplifies polysaccharide quantification and can be quickly adapted

for other polysaccharides and biological systems.

Methods

Chemicals

All chemicals were obtained from Sigma Aldritch unless otherwise specified.

Preparation of colloidal chitin

10 grams of powdered chitin from shrimp shells (Sigma) was dissolved in 100 mL of

concentrated phosphoric acid and placed at 4°C for 48 hrs. 500 ml of millQ water was added

to precipitate the chitin, which was then vacuum filtered and dialyzed for 72 hours in water

with a cellulose dialysis tubing membrane (Sigma). The colloidal chitin was then

homogenized using a Bosch SilentMixx Pro blender. The colloidal chitin was autoclaved

before use.

LC-MS quantification

Measurements were performed using Liquid Chromatography (Agilent Infinity II UHPLC

stack) coupled with an Agilent 6520 Time of Fight Quadrupole Time of Flight Mass

Spectrometer in negative mode, 4GHz, high resolution mode.

For quantification of chitin oligomers, an Agilent EC-CN Poroshell column (50x2.1 mm, 2.7

µM) was used isocratically to reduce interference of salts on metabolite ionization(Pontrelli

and Sauer, 2021). The buffer contained 90% water, 10% (v/v) acetonitrile

(CHROMOSOLVE) and 0.1% (v/v) formic acid. The flow rate was 0.35 mL/min. Prior to

measurement, the sample was diluted 20-fold in millQ water. 3 µL was injected every 2

minutes. Data analysis and quantification was performed using Agilent Quantitative

Masshunter software.

For time course laminarin pentose degradation, an Agilent EC-CN Poroshell column (50x2.1

mm, 2.7 µM) was used isocratically as described above. Here, the assay was initiated in the

Agilent Infinity autosampler by adding 50 µL of 10 mM laminarin pentose to 50 µL of

enzyme cocktail, and 3 µL of the enzyme reaction was injected every 3 hours. Data analysis

was performed through manual peak integration using Agilent Qualitative Masshunter

software.

For quantification of laminarin oligomers of different chain lengths, an Agilent HILIC-Z Poroshell column (100x2.1 mm, 1.8 µM) was used. Buffer A contains 10% (v/v) acetonitrile and 0.3% (v/v) ammonium hydroxide. Buffer B contains 90% acetonitrile (v/v) and 0.3% ammonium hydroxide (v/v). The flow was 0.9 mL/min and the column was kept at 35°C. The gradient is as follows. 0% A for 1 minute, followed by a gradient to 40% A until 5 minutes. This is held for 1 minute before dropping to 0% A and re-equilibration for 3 minutes. 3 µL of sample was injected. Data analysis and quantification was performed using manual peak integration with Agilent Qualitative Masshunter software.

Enzyme cocktail purification

*Psy*6C06 was streaked from -80°C glycerol stocks onto 1.5% agar plates of MB 2216 (Fisher) medium. A single colony was used for an overnight preculture in MB 2216, and the following day the preculture was inoculated 1% into 400 mL of simplified MBL medium with 2 g/l colloidal chitin and shaken at 200 rpm until early stationary phase. Simplified MBL medium is created from several stocks: 4-fold concentrated seawater salts (NaCl, 80 g/L; $MgCl_2*6H_2O$, 12 g/L; $CaCl_2*2H_2O$, 0.6 g/L; KCl, 2 g/L), 1000-fold diluted sodium sulfate (1 M), 500-fold diluted phosphate dibasic (0.5mM), and 20-fold diluted HEPES buffer (1 M pH 8.2).

After growth, the medium was centrifuged at 2800 relative centrifugal force for 20 minutes to remove any remaining chitin and cells, then sterile filtered using a 0.2 µM membrane. This supernatant was then concentrated 10 fold using an Amicon stirred cell (Millipore) with a 3 kDa cutoff filter. A protease inhibitor was added to the final concentrate (Roche cOmplete EDTA free protease inhibitor cocktail). 500 µL aliquots were added to 1.5 mL microcentrifuge tubes, which were then snap frozen by placing them in liquid nitrogen before storing at -80°C until use. The concentration of protein in the chitinase enzyme cocktail is 0.07 mg/mL as quantified using Bradford reagent.

For generation of laminarinases from seawater microbes, the same growth and purification

protocol was followed, except the medium contained 1 g/l Laminarin from *Laminaria digitata*

(Sigma) as the carbon source and 10 mM ammonium chloride as the nitrogen source.

Seawater was collected from the North Sea on the shore of Cuxhaven (53°53'04.3"N

8°37'57.1"E), passed through a 5 µM filter, mixed 1:1 with 80% (v/v) glycerol, and stored at -

80 °C until inoculation. The concentration of protein in the laminarinase enzyme cocktail is

0.012 mg/mL as quantified using Bradford reagent.

Chitin quantification assay

Frozen enzyme aliquots were thawed on ice. 100 µL of enzyme was added to 100 µL of

sample and placed at room temperature until sampling. If the sample is suspected to contain

more than 500 mg/L chitin, it was diluted to fall within this quantifiable range. To sample, the

assay was centrifuged at 2800 rcf for 1 minute and the supernatant was collected and stored at

-20°C until enzyme products are measured using DNS reagent or LC-MS.

Laminarin quantification assay

Frozen enzyme aliquots were thawed on ice. For laminarin oligomer formation, 50 µL of

enzyme was added to 50 µL of laminarin pentamer (Megazyme, dissolved in 40 mM

ammonium bicarbonate buffer pH 7.8) and placed at room temperature for 24 hours. The

reaction mixture was directly quantified using LC-MS as described above. For laminarin

quantification, 50 µL laminarin from *Laminaria digitata* was added at various concentrations

to 50 µL of enzyme cocktail. Samples were taken at 24 hours and directly quantified using

DNS reagent.

DNS reagent preparation

100 mL of water was heated to 70-75°C, then poured over 2 g of NaOH pellets with constant

stirring. 2.18 g of 3,5-dinitrosalicylic acid was then dissolved in this mixture, and 30 g of

sodium potassium tartrate (Rochelle salts) were added. The solution was cooled to room

temperature and stored in the dark.

DNS reducing sugar assay

25 µL of sample was added to 75 µL of DNS reagent and the mixture was heated to 95°C for

15 minutes in a thermocycler. 80 µL was added to a 384 well plate and a colorimetric readout

was measured at OD_{540}.

Assessment

       Figure 1

Assay design

A polysaccharide-degrading microbe or complex microbial community is cultured on the

polysaccharide until early stationary phase to select for degraders and induce the expression

of hydrolytic enzymes. The cells are then filtered to obtain a cell-free supernatant containing

secreted enzymes. Enzymes are concentrated tenfold using a 3kDa molecular weight cutoff

filter, aliquoted, snap-frozen in liquid nitrogen, and stored at -80°C. This enzyme cocktail is

then thawed and mixed with a polysaccharide sample to initiate hydrolysis. After hydrolysis,

the supernatant is collected and quantification of monomers and oligomers is performed using

a reducing sugar assay or liquid chromatography mass spectrometry (LC-MS). Akin other

enzyme-based quantification methods(Becker et al., 2017; Deng et al., 2022), this assay relies

on identical enzyme activity across all samples, producing consistent sets of oligomers and

monomers. This provides a quantitative relationship between the amount of polysaccharide

digested and the number of reducing sugar ends and degradation products. A calibration curve

Figure 2

is thus calculated based on digestion of standards with known polysaccharide concentrations.

These standards are always digested alongside unknown samples to account for variations in

enzyme activity of different cocktail batches, their degradation during storage, or any mono-

or oligomers that may be residually present in the cocktails following their concentration.

Quantification of colloidal chitin using enzyme cocktails from a bacterial isolate

We tested the assay design to quantify colloidal chitin with an enzyme cocktail obtained from

the chitin-degrading *Psychromonas sp.* bacterium *Psy*6C06 that encodes 13

chitinases(Pontrelli et al., 2022). To verify chitinase activity, the cocktail was incubated for 72

h at equal volumes with a sample containing 2 g/L of colloidal chitin. Produced mono- and

oligomers were determined by LC-MS. Monomeric GlcNAc and oligomers with various

chain lengths were detected. The major products are monomeric GlcNAc and dimeric

chitobiose, mirroring previous characterization of *Psy*6C06 secreted chitinases(Pontrelli et al.,

2022) **(Fig. 2A)**. We tested the time required for the cocktail to completely hydrolyze samples

containing up to 2 g/L of colloidal chitin by measuring the formation of reducing sugar ends

over time using the 3,5-dinitrosalicylic acid (DNS) reagent **(Fig. 2B)**. Within 72 hours, only

samples with 500 mg/L colloidal chitin or less were completely hydrolyzed, as judged from a

stable signal. All subsequent experiments therefore used 72 hours for hydrolysis and samples

known, or suspected, to contain more than 500 mg/L colloidal chitin were diluted prior to

digestion.

Figure 3

Enzyme cocktails may contain multiple hydrolytic enzymes that degrade polysaccharides

other than the target polysaccharide. Thus, if an alternative polysaccharide is non-specifically

degraded into mono- and oligomers, colorimetric assays like DNS, which react with many

different reducing sugar ends, may overestimate concentrations of the target polysaccharide.

Since LC-MS or HPLC can distinguish mono- and oligomers after a sample is digested, these

degradation products can be used as specific readouts for the degradation of a target

polysaccharide. As previously demonstrated(Becker et al., 2017; Deng et al., 2022), this is

especially useful if the enzyme digest produces known, specific oligomers from a target

polysaccharide, which can help distinguish a specific polysaccharide's oligomers even in

samples with multiple polysaccharides present. Here, enzyme cocktails from *Psy*6C06

produce GlcNAc and chitobiose as major degradation products. Both can be used as readouts

for chitin quantification. However, in cases where it is unknown if other GlcNAc-containing

polysaccharides are present in a sample that may be non-specifically hydrolyzed, quantifying

chitin based on chitobiose provides a viable option. This is because chitobiose is verified to be

specifically produced by the enzyme digest, and direct GlcNAc-GlcNAc linkages are

generally rare in natural polysaccharides. We compared the use of chitobiose measured with

LC-MS as a readout against the other two readouts: GlcNAc measured with LC-MS and total

reducing sugar ends using DNS reagent. We determined the limit of detection (LOD), defined

as the lowest concentration with a signal above the background mean plus five standard

deviations. Using LC-MS, we obtained LODs of 15 mg/L colloidal chitin with chitobiose as a

readout or 500 mg/L with GlcNAc **(Fig. 3ABC)**. Using DNS, we determined an LOD of 62.5

mg/L. These results demonstrate that using chitobiose as a readout not only provides a more

specific quantification for chitin, but also allows for the most sensitive readouts.

Measuring colloidal chitin dynamics in bacterial cultures

Quantifying chitin degradation dynamics is important for the investigation of bacterial

cultures and marine communities involved in biogeochemical carbon cycling, allowing to link

changes in population dynamics with chitin degradation. Using the above enzyme cocktail,

we monitored the time-course consumption of colloidal chitin by the chitin-degrading *Vibrio*

*splendidus, Vib*1A01(Amarnath et al., 2023; Iffland-Stettner et al., 2023). Aliquots were

periodically sampled during growth, centrifuged, and the pellet was briefly boiled to denature

any *Vib*1A01-derived chitinases. The denatured pellet was then incubated for 72 h with a 1:1

ratio of the enzyme cocktail. To quantify degradation products, we first ensured that other

components in *Vib*1A01 biomass are not nonspecifically hydrolyzed by the enzyme digest. To

this end, we used the enzyme cocktail to digest biomass from a *Vib*1A01 culture grown on

Figure 4

20mM GlcNAc instead of chitin and observed no detectable background signal using DNS

reagent (data not shown). We therefore determined reducing sugars in the chitin culture using

DNS reagent. Results show a gradual reduction in colloidal chitin corresponding to increased

*Vib*1A01 growth **(Fig. 4)**. This highlights the method's utility in quantifying chitin degradation

dynamics in bacterial cultures, which has broader implications for researchers studying chitin

dynamics in other biological systems.

Laminarin quantification using enzyme cocktails from environmental communities

For polysaccharides where degrading microbes are unavailable or cannot be isolated from the

environment, collecting enzyme cocktails from complex environmental microbial

communities is an alternative. By growing environmental communities on the polysaccharide

of interest as a sole carbon source, degraders will be selected, and their secreted enzymes can

be concentrated into a cocktail. We tested this approach for the quantification of laminarin, an

algal storage molecule with a β-(1-3)-glucose backbone and single β-(1-6)-glucose side

branches. Seawater collected from the North Sea on the shore of Cuxhaven (53°53'04.3"N

8°37'57.1"E) was used to inoculate a medium with laminarin as the sole carbon source to

induce and concentrate laminarinases in a cocktail. To verify the activity of the enzyme

cocktail, we digested 5 mM of a β-(1-3)-glucose pentamer, representing the laminarin

backbone, for four hours and measured the formation of smaller oligomers using LC-MS **(Fig.**

5A). The formation of a mix of monomers and smaller oligomers indicates the presence of

multiple microbial laminarinases with different activities. We determined the time required to

completely digest 2 mM laminarin pentamer (1.6 g/L) using LC-MS and observed a stable

signal before 21 hours **(Fig. 5B)**, identifying 24 hours as a suitable digestion time for

quantities of laminarin of 1 g/L or less. To demonstrate that DNS reagent can also be used to

quantify laminarin, and hence provide a widely accessible measurement method beyond LC-

MS, we used it to detect laminarin concentrations between 1 g/L and 0.03 g/L after a 24 hr

digestion period and observed an LOD of 30 mg/L **(Fig. 5C)**. These results highlight the

utility of environmental microbial communities as a source of enzyme cocktails to quantify

polysaccharides.

Figure 5

Discussion

In this study, we demonstrate how microbial enzyme cocktails can be used to quantify

polysaccharides, exemplified with colloidal chitin and laminarin. Unlike other enzyme-based

polysaccharide quantification techniques, our approach eliminates the need to identify and

recombinantly express specific hydrolytic enzymes, by collecting them from cultures of

bacteria that are actively degrading a polysaccharide. This is particularly advantageous for

polymers whose degradation requires multiple enzymes such as fucoidan(Sichert et al., 2020).

Concentrating all secreted enzymes into single cocktails has the additional benefit that

potential non-catalytic proteins that enhance activities of hydrolytic enzymes are

maintained(Vaaje-Kolstad et al., 2005). Furthermore, we demonstrate that enzyme cocktails

can be derived from complex microbial communities, bypassing the need to obtain specific

polysaccharide degraders.

A limitation of this approach is that cocktails may contain other hydrolytic enzymes that break

down alternative polysaccharides in complex samples, inflating quantified values. One way to

overcome this is by performing controls to rule out background interference. For example, we

showed that the *Psy*6C06 enzyme cocktail does not digest polysaccharides in *Vib*1A01

bacterial biomass, meaning there is no interference when measuring bacterial degradation of

colloidal chitin. LC-MS can monitor specific mono- and oligomers, aiding in quantifying the

target polysaccharide, especially if the enzyme cocktail produces polysaccharide-specific

oligomers like chitobiose. However, LC-MS may still fall short for complex samples, as

different polysaccharides may have similar monosaccharide compositions, skewing readouts

due to nonspecific hydrolysis of alternative polysaccharides. This may occur in the case of

laminarin which is solely composed of glucose and would be difficult to quantify when

alternative glucans and glucanases are also present. Nevertheless, the presented approach is

suitable when non-specific hydrolysis can be ruled out or when measuring polysaccharide-

specific oligomers. It is also suitable when the polysaccharide composition is known, such as

when specific substrates are provided to microbial isolates as sole carbon sources or after

extraction procedures isolate the target polysaccharide from complex environmental samples.

This work showcases the utility of microbial enzyme cocktails as an easily accessible means

for quantifying polysaccharides, and provides a specific protocol for colloidal chitin that is

currently missing for this abundant marine polysaccharide(Pontrelli et al., 2022; Salas-Ovilla

et al., 2019; Shen et al., 2010; Szabo et al., 2022). By requiring only a microbial isolate or

community that secretes hydrolytic enzymes, our approach is highly transferable to other

polysaccharides and biological systems. Our findings highlight the potential to harness natural

microbial communities as enzyme sources, offering a flexible method for measuring a wide

range of polysaccharides.

Acknowledgements

We greatly acknowledge financial support from the Simons Foundation through the Principles

of Microbial Ecosystems (PriME) collaboration (grant 542395).

Bibliography

Amarnath, K., Narla, A.V., Pontrelli, S., Dong, J., Reddan, J., Taylor, B.R., Caglar, T., Schwartzman, J., Sauer, U., Cordero, O.X., Hwa, T., 2023. Stress-induced metabolic exchanges between complementary bacterial types underly a dynamic mechanism of inter-species stress resistance. Nature Communications 14, 3165. https://doi.org/10.1038/s41467-023-38913-8

Arnosti, C., Wietz, M., Brinkhoff, T., Hehemann, J.-H., Probandt, D., Zeugner, L., Amann, R., 2021. The Biogeochemistry of Marine Polysaccharides: Sources, Inventories, and Bacterial Drivers of the Carbohydrate Cycle. Annu. Rev. Mar. Sci. 13, 81–108. https://doi.org/10.1146/annurev-marine-032020-012810

Becker, S., Scheffel, A., Polz, M.F., Hehemann, J.-H., 2017. Accurate Quantification of Laminarin in Marine Organic Matter with Enzymes from Marine Microbes. Appl Environ Microbiol 83, e03389-16. https://doi.org/10.1128/AEM.03389-16

Buck-Wiese, H., Andskog, M.A., Nguyen, N.P., Bligh, M., Asmala, E., Vidal-Melgosa, S., Liebeke, M., Gustafsson, C., Hehemann, J.-H., 2023. Fucoid brown algae inject fucoidan carbon into the ocean. Proc. Natl. Acad. Sci. U.S.A. 120, e2210561119. https://doi.org/10.1073/pnas.2210561119

Cao, X., Mulholland, M.R., Helms, J.R., Bernhardt, P.W., Duan, P., Mao, J., Schmidt-Rohr, K., 2017. A Major Step in Opening the Black Box of High-Molecular-Weight Dissolved Organic Nitrogen by Isotopic Labeling of *Synechococcus* and Multibond Two-Dimensional NMR. Anal. Chem. 89, 11990–11998. https://doi.org/10.1021/acs.analchem.7b02335

Cerqueira, F.M., Photenhauer, A.L., Pollet, R.M., Brown, H.A., Koropatkin, N.M., 2020. Starch Digestion by Gut Bacteria: Crowdsourcing for Carbs. Trends in Microbiology 28, 95–108. https://doi.org/10.1016/j.tim.2019.09.004

Decamp, A., Michelo, O., Rabbat, C., Laroche, C., Grizeau, D., Pruvost, J., Gonçalves, O., 2021. A New, Quick, and Simple Protocol to Evaluate Microalgae Polysaccharide Composition. Marine Drugs 19, 101. https://doi.org/10.3390/md19020101

Deng, Y., Chen, L.-X., Zhu, B.-J., Zhao, J., Li, S.-P., 2022. A quantitative method for polysaccharides based on endo-enzymatic released specific oligosaccharides: A case of Lentinus edodes. International Journal of Biological Macromolecules 205, 15–22. https://doi.org/10.1016/j.ijbiomac.2022.02.048

Enke, T.N., Datta, M.S., Schwartzman, J., Cermak, N., Schmitz, D., Barrere, J., Pascual-García, A., Cordero, O.X., 2019. Modular assembly of polysaccharide-degrading marine microbial communities. Current Biology 29, 1528-1535.e6. https://doi.org/10.1016/j.cub.2019.03.047

Horwath, W., 2007. Carbon cycling and formation of soil organic matter, in: Soil Microbiology, Ecology and Biochemistry. Elsevier, pp. 303–339. https://doi.org/10.1016/B978-0-08-047514-1.50016-0

Iffland-Stettner, A., Okano, H., Gralka, M., Guessous, G., Amarnath, K., Cordero, O.X., Hwa, T., Bonhoeffer, S., 2023. A Genome-Scale Metabolic Model of Marine Heterotroph Vibrio splendidus Strain 1A01. mSystems 8, e00377-22. https://doi.org/10.1128/msystems.00377-22

Keyhani, N.O., Roseman, S., 1999. Physiological aspects of chitin catabolism in marine bacteria11This paper is publication 521 from the McCollum-Pratt Institute. Biochimica et Biophysica Acta (BBA) - General Subjects 1473, 108–122. https://doi.org/10.1016/S0304-4165(99)00172-5

Lerouxel, O., Cavalier, D.M., Liepman, A.H., Keegstra, K., 2006. Biosynthesis of plant cell wall polysaccharides — a complex process. Current Opinion in Plant Biology 9, 621–630. https://doi.org/10.1016/j.pbi.2006.09.009

Lin, P., Guo, L., 2015. Spatial and vertical variability of dissolved carbohydrate species in the
northern Gulf of Mexico following the Deepwater Horizon oil spill, 2010–2011.
Marine Chemistry 174, 13–25. https://doi.org/10.1016/j.marchem.2015.04.001
Peter, M.G., 1995. Applications and Environmental Aspects of Chitin and Chitosan. Journal of
Macromolecular Science, Part A 32, 629–640.
https://doi.org/10.1080/10601329508010276
Pontrelli, S., Sauer, U., 2021. Salt-tolerant metabolomics for exometabolomic measurements
of marine bacterial isolates. Anal. Chem. 93, 7164–7171.
https://doi.org/10.1021/acs.analchem.0c04795
Pontrelli, S., Szabo, R., Pollak, S., Schwartzman, J., Ledezma-Tejeida, D., Cordero, O.X.,
Sauer, U., 2022. Metabolic cross-feeding structures the assembly of polysaccharide
degrading communities. Sci. Adv. 8, eabk3076. https://doi.org/10.1126/sciadv.abk3076
Raszka, A., Chorvatova, M., Wanner, J., 2006. The role and significance of extracellular
polymers in activated sludge. Part I: Literature review. Acta hydrochim. hydrobiol. 34,
411–424. https://doi.org/10.1002/aheh.200500640
Salas-Ovilla, R., Gálvez-López, D., Vázquez-Ovando, A., Salvador-Figueroa, M., Rosas-
Quijano, R., 2019. Isolation and identification of marine strains of *Stenotrophomona
maltophilia* with high chitinolytic activity. PeerJ 7, e6102.
https://doi.org/10.7717/peerj.6102
Schiavone, M., Vax, A., Formosa, C., Martin-Yken, H., Dague, E., François, J.M., 2014. A
combined chemical and enzymatic method to determine quantitatively the
polysaccharide components in the cell wall of yeasts. FEMS Yeast Res 14, 933–947.
https://doi.org/10.1111/1567-1364.12182
Shen, C.-R., Chen, Y.-S., Yang, C.-J., Chen, J.-K., Liu, C.-L., 2010. Colloid Chitin Azure Is a
Dispersible, Low-Cost Substrate for Chitinase Measurements in a Sensitive, Fast,
Reproducible Assay. SLAS Discovery 15, 213–217.
https://doi.org/10.1177/1087057109355057
Sichert, A., Corzett, C.H., Schechter, M.S., Unfried, F., Markert, S., Becher, D., Fernandez-
Guerra, A., Liebeke, M., Schweder, T., Polz, M.F., Hehemann, J.-H., 2020.
Verrucomicrobia use hundreds of enzymes to digest the algal polysaccharide fucoidan.
Nat Microbiol 5, 1026–1039. https://doi.org/10.1038/s41564-020-0720-2
Steinke, N., Vidal☐Melgosa, S., Schultz☐Johansen, M., Hehemann, J., 2022. Biocatalytic
quantification of α☐glucan in marine particulate organic matter. MicrobiologyOpen
11, e1289. https://doi.org/10.1002/mbo3.1289
Sun, Y., Li, L., Zhang, Y., Xue, C., Chang, Y., 2024. An enzyme-pHBH method for specific
quantification of porphyran. International Journal of Biological Macromolecules 257,
128530. https://doi.org/10.1016/j.ijbiomac.2023.128530
Suzuki, E., Suzuki, R., 2013. Variation of Storage Polysaccharides in Phototrophic
Microorganisms. J. Appl. Glycosci. 60, 21–27. https://doi.org/10.5458/jag.jag.JAG-
2012_016
Szabo, R.E., Pontrelli, S., Grilli, J., Schwartzman, J.A., Pollak, S., Sauer, U., Cordero, O.X.,
2022. Historical contingencies and phage induction diversify bacterioplankton
communities at the microscale. Proc. Natl. Acad. Sci. U.S.A. 119, e2117748119.
https://doi.org/10.1073/pnas.2117748119
Unfried, F., Becker, S., Robb, C.S., Hehemann, J.-H., Markert, S., Heiden, S.E., Hinzke, T.,
Becher, D., Reintjes, G., Krüger, K., Avcı, B., Kappelmann, L., Hahnke, R.L., Fischer,
T., Harder, J., Teeling, H., Fuchs, B., Barbeyron, T., Amann, R.I., Schweder, T., 2018.
Adaptive mechanisms that provide competitive advantages to marine bacteroidetes
during microalgal blooms. The ISME Journal 12, 2894–2906.
https://doi.org/10.1038/s41396-018-0243-5

Vaaje-Kolstad, G., Horn, S.J., van Aalten, D.M.F., Synstad, B., Eijsink, V.G.H., 2005. The
Non-catalytic Chitin-binding Protein CBP21 from Serratia marcescens Is Essential for
Chitin Degradation. Journal of Biological Chemistry 280, 28492–28497.
https://doi.org/10.1074/jbc.M504468200
Yan, X., Evenocheck, H.M., 2012. Chitosan analysis using acid hydrolysis and HPLC/UV.
Carbohydrate Polymers 87, 1774–1778. https://doi.org/10.1016/j.carbpol.2011.09.091
407

Figure legends

Figure 1: Assay design and workflow A) Enzyme cocktail preparation involves growing a

microbe, or microbial community, on the polysaccharide as a sole carbon source, followed by

filtering cells and concentrating enzymes using a 3 kDa molecular weight cutoff membrane.

The concentrate is aliquoted and stored until use. B) Polysaccharide quantification entails

mixing the enzyme cocktail with a sample to initiate digestion, leading to the formation of

monomers or oligomers. Post-digestion, the products can be quantified using LC-MS, HPLC,

chemical, or other methods.

Figure 2: Activity of chitin enzyme cocktail. A) GlcNAc oligomers (Degree of

polymerization, DP 1-6) formed after 72 hr digestion of 2 g/L colloidal chitin by the enzyme

cocktail, measured using LC-MS. B) Samples with varying concentrations of colloidal chitin

digested with equal volumes of enzyme cocktail. Liberation of mono- and oligomers is

measured every 24 hours for 3 days using a DNS reducing sugar assay (with an OD_{540} signal).

Experiments were performed in triplicate. Error bars represent standard deviation from the

mean.

Figure 3: Assay sensitivity. Decreasing concentrations of colloidal chitin are applied to

enzyme cocktail and digested for 72 hours. A) Detection of total reducing sugar ends with the

DNS reagent. B) Detection of GlcNAc by LC-MS. C) Detection of dimeric chitobiose by LC-

MS. A linear regression of signal intensity per chitin concentration is shown.

Figure 4: Chitin concentrations during bacterial growth. OD_{600} of chitin degrader

*Vib*1A01 when growing on chitin as a sole carbon source (Red). Colloidal chitin concentration

at each of the measured timepoints (Blue). Error bars represent standard deviation of the mean

of three biological replicates.

Figure 5: Laminarin quantification with seawater community enzymes: A) Mass

spectrometry measurement of glucose oligomers (Degree of polymerization, DP1-4) after 4

hour digestion of a laminarin pentamer by a laminarinase enzyme cocktail expressed by a

seawater microbial community. The horizontal bar is fold change of 1 compared to a no-

enzyme control. B) Time course digestion of 2 mM laminarin pentamer using the enzyme

cocktail, measured using LC-MS. C) Digestion of different laminarin concentrations by an

enzyme cocktail, quantified with DNS reagent and a readout of OD_{540}. A polynomial

regression is shown and the LOD is 30 mg/L. Error bars represent standard deviation of the

mean of three biological replicates.