Inclusive, Exclusive and Hierarchical Atlas of NFATc1$^+$/PDGFR-α$^+$ Cells in Dental and Periodontal Mesenchyme

Xue Yang[1, 2, *], Chuyi Han[1, *], Changhao Yu[1, 2], Bin Zhou[3], Ling Ye[1, 2, #], Feifei Li[1, 4, #], Fanyuan Yu[1, 2, #]

1 State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases, West China Hospital of Stomatology, Sichuan University, China

2 Department of Endodontics, West China Hospital of Stomatology, Sichuan University, China

3 Center for Excellence in Molecular Cell Science, University of Chinese Academy of Sciences, Chinese Academy of Sciences, China

4 Department of Pediatric Dentistry, West China Hospital of Stomatology, Sichuan University, China

* These authors contribute equally.

Corresponding Author. E-mail: yeling@scu.edu.cn; feidepingfanshijie@163.com; fanyuan_yu@outlook.com

Abstract

Platelet-derived growth factor receptor alpha (PDGFR-α) activity is crucial in the process of dental and periodontal mesenchyme regeneration facilitated by autologous platelet concentrates (APCs), such as platelet-rich fibrin (PRF), platelet-rich plasma (PRP) and concentrated growth factors (CGF), as well as by recombinant PDGF drugs. However, it is largely unclear about the physiological patterns and cellular fate determinations of PDGFR-α$^+$ cells in the homeostasis maintaining of adult dental and periodontal mesenchyme. We previously identified NFATc1 expressing PDGFR-α$^+$ cells as a subtype of skeletal stem cells (SSCs) in limb bone, but their roles in dental and periodontal remain unexplored. To this end, in the present study we investigated the spatiotemporal atlas of NFATc1$^+$ and PDGFR-α$^+$ cells residing in dental and periodontal mesenchyme, their capacity for progeny cell generation, and their

inclusive, exclusive and hierarchical relations in homeostasis. We utilized CRISPR/Cas9-mediated gene editing to generate two dual recombination systems, which were *Cre/loxP-Dre/Rox* combined intersectional and exclusive reporters respectively, to concurrently demonstrate the inclusive, exclusive, and hierarchical distributions of NFATc1$^+$ and PDGFR-α$^+$ cells and their lineage commitment. By employing the state-of-the-art transgenic lineage tracing techniques in cooperating with tissue clearing-based advanced imaging and three-dimensional slices reconstruction, we systematically mapped the distribution atlas of NFATc1$^+$ and PDGFR-α$^+$ cells in dental and periodontal mesenchyme and tracked their *in vivo* fate trajectories. Our findings extend current understanding of NFATc1$^+$ and PDGFR-α$^+$ cells in dental and periodontal mesenchyme homeostasis, and furthermore enhance our comprehension of their sustained therapeutic impact for future clinical investigations.

Key words

Lineage tracking; genetic labeling; mesenchymal stromal cell; dental pulp; periodontal tissues

Introduction

Platelet-derived growth factor (PDGF), first identified in platelets, intricately involved in the process of tissue regeneration facilitated by autologous platelet concentrates (APCs)[1, 2]. The effects of PDGFs are mediated by two receptor tyrosine kinases, namely PDGFRα and PDGFRβ. The PDGFRα homodimer predominantly binds to the majority of PDGF isoforms (PDGF-AA, PDGF-AB, PDGF-BB, and PDGF-CC)[3]. In recent decades, APCs have gained widespread utilization in the field of tissue regenerative medicine, with notable generations including platelet-rich plasma (PRP), platelet-rich fibrin (PRF), and concentrated growth factors (CGF). Evidence-based medicine studies have demonstrated that these three APC techniques have been utilized in the field of periodontology, contributing to periodontal

regenerative therapy, albeit with varying efficacy[4]. Additionally, certain investigations

suggest that PDGF-BB also exerted a role in promoting regeneration of both

periodontal tissues and dental pulp[5, 6]. Teeth, being highly mineralized organs, enclose

a mineralized chamber housing the dental pulp mesenchyme, while the surrounding

tissue comprises periodontal soft tissue—periodontal ligament (PDL)[7]. Simply, teeth

can be regarded as hard tissues enveloped by mesenchyme. While the dental pulp and

periodontium fulfill distinct physiological roles, they also maintain

interconnectedness to collectively sustain the physiological function of the tooth.

Consequently, when pathological changes occur independently or concurrently in the

dental pulp and periodontal tissues, they may mutually influence each other[8].

Therefore, it is imperative to elucidate the expression and biological mechanism of

PDGFR-α^+ cells in PDL and dental pulp tissues in order to comprehensively

comprehend the physiological and therapeutic effects mentioned above.

PDGFR-α serves as a pivotal signaling molecule involved in mediating tooth

development, morphogenesis, and physiological responses[9-11]. Its potential role in

periodontal formation has also been suggested[12]. Within skeletal tissues, PDGFR-α

has been identified as a heterogeneous population, with not all PDGFR-α^+ cells

exhibiting mesenchymal stem cell characteristics[13-16]. Certain studies suggest that

subpopulations of PDGFR-α^+ cells are considered "true" stem cells. For instance,

co-labeling of Nestin and PDGFR-α has been observed in bone marrow stem cells

(BMSCs) derived from the neural crest[17], and the co-expression of PDGFR-α and

Sca-1 in the non-hematopoietic compartment has identified a distinct cell population

within the adult murine bone marrow[18]. Our prior investigations have also revealed a

population of NFATc1 and PDGFR-α double-positive skeletal stem cells (SSCs)

within long bones[19]. In long bones, NFATc1$^+$ cells are considered SSCs[19].

Furthermore, previous RNA-sequencing analyses have reported the expression of

NFATc1 in jawbones and periodontal tissues[20]. However, this evidence was limited to

RNA-level sequencing, lacking data support *in vivo*.

The conventional single-enzyme cutting system, represented by the *Cre-loxP*

system primarily focuses on individual cells[21]. The introduction of multiple DNA

recombinases has revolutionized lineage tracing by enabling a layered and

intersectional analysis that was previously unattainable[22, 23]. Notably, the utilization of

multiple recombinase-mediated recombination includes systems such as Intersectional

reports (IR) and exclusive reporters (ER), which facilitate the visualization of two sets

of orthogonal recombination or hierarchical relationships[23]. In this study, considering

significant tissue variations in dental pulp and PDL, traditional slicing methods may

inadequately represent their expression patterns[24]. To ensure accurate depiction of

expression patterns in spatial dimensions, we employed two strategies: tissue

transparency (TC) and three-dimensional (3D) slices reconstruction, guaranteeing

comprehensive spatial imaging technology for investigating tissue structural

architecture.

In summary, we utilized the advanced *Cre/loxP-Dre/Rox* combined intersectional

and exclusive reporters—IR and ER systems along with advanced imaging methods

to comprehensively examine the intricate relationships, including inclusive, exclusive

and hierarchical between two distinct cell populations within pulp and periodontal

mesenchyme for the first time. Furthermore, we demonstrated their capacity for

progeny cell generation and hierarchical positioning to depict their *in vivo* fate

determination trajectories. Additionally, we assessed the advantages and limitations of

TC and 3D slices reconstruction techniques. These findings not only furnish essential

tools and references for future investigations into MSC lineage tracing within dental

and periodontal tissues but also enhance our comprehension of biological research

and therapeutic applications in regenerative medicine.

RESULT

1. Establishment of multiple genetic recombination systems and advanced

imaging technology

To understand the cell fate, behaviors, and their spatiotemporal correlation of

PDGFR-α^+ and NFATc1$^+$ cells in dental and periodontal mesenchyme, systematic

genetic fate atlas were established by lineage tracing based on genetic DNA recombination technology. In conventional single recombinase-mediated genetic readouts, the *Cre-loxP* system is widely used for mammalian gene editing with Cre recombinase catalyzing recombination between *loxP* sites. This system, often activated by a ubiquitous promoter like Rosa26 and H11, removes a transcription stop cassette post-recombination to allow for reporter gene expression (Figure 1A). However, the drawback of the conventional approach is that it is impossible to observe the genetic lineage and spatial relationship of two specific cell type at the same time, also, targeting two gene promoters in one cell population could be more precise than relying on a single promoter as commonly employed in the conventional reporter system. Given that, diverse dual recombinase–mediated genetic labeling systems have been developed to enhance the specificity and the number of cell types being labeled simultaneously. *Cre-loxP, Flp-frt, Dre-rox,* and *Nigri-nox* have been respectively used for designing dual system. Using reporter as an entry site can categorize these systems into three different types for multiple recombinase–mediated fate mapping studies, including intersectional reporters, exclusive reporters, and nested reporters (Figure 1A-b, c). In this work, the most advanced transgenic technology, dual recombinase–mediated genetic labeling systems, including intersectional reporters and exclusive reporters, were constructed to simultaneously show the inclusive, exclusive and hierarchical distribution of PDGFR-α^+/NFATc1$^+$ cells in dental and periodontal mesenchyme. Specifically, multichromatic crossover intersectional reporters (H11-LGRT) systems, which structured as *loxP-Stop-loxP-ZsGreen-rox-Stop-rox-tdTomato*, enabled the tracing of three cell types: *Cre+&Dre-, Cre-&Dre+, and Cre+&Dre+*, corresponding to the colors ZsGreen, tdTomato, and yellow, respectively. This technology guaranteed us to precisely understand the respective and inclusive distribution of Pdgfr-α^+, NFATc1$^+$ cells and their progeny cells. In addition, IR1 system is an exclusive reporter expressed as *CAG-loxP-rox-Stop-loxP-ZsGreen-Stop-rox-tdTomato* (Figure 1A-c). The first *Cre-loxP* recombination results in ZsGreen expression and the removal of a

rox site, preventing subsequent *Dre-rox* recombination in the same cell. This means

that once a recombination event occurs, the expression of the corresponding reporter

gene eliminates a recognition site for the other recombination system, which could

provide generous information of the hierarchical distribution of PDGFR-α^+ and

NFATc1$^+$ cells in dental and periodontal mesenchyme.

Due to the peculiarity of dental/periodontal tissues, including pulp and

periodontal of maxillary and mandibular first molars (small volume, complex

structure and uneven spatial distribution of SSCs in mesenchyme), single-slice images

based on traditional frozen section and confocal microscope imaging were insufficient

to accurately presume the practical situation of the organization. Therefore, the

advanced tissue deep clearance using SUMIC procedure and 3D reconstruction was

exploited to observe the overall spatial distribution of PDGFR-α^+ and NFATc1$^+$ cells

in dental and periodontal mesenchyme of transgenic mice with multiple DNA

recombinases-based genetic lineage tracing system (Figure 1C-a). Simultaneously, as

an irreplaceable imaging technology, traditional sectioning combined confocal

imaging and procedure was also estimated (Figure 1C-b). These two

3D-reconstruction and imaging technologies complement each other to jointly address

the spatial positioning and hierarchical relationships of PDGFR-α^+, NFATc1$^+$, and

PDGFR-α^+ NFATc1$^+$ cells from multiple perspectives.

2. Exploration of NFATc1$^+$/PDGFR-α^+ cells and their progeny cells in H11-LGRT

system

First, we established *Cre-loxP* and *Dre-rox* reporter lineage-tracking model of

PDGFR-α and NFATc1 in adult mice to observe the spatial location of PDGFR-α^+ and

NFATc1$^+$ cells in pulp and periodontal of maxilla & mandible M1. The pulse timing

point we set was tamoxifen treated twice at day 1 and day 3, and then the mice were

sacrifice to obtain maxilla and mandible on day 5 (Figure 2A, B).

2.1 Dental and periodontal region of maxilla M1 and mandible M1

As one of the most advanced imaging technologies currently available, tissue

clearing/imaging allows for direct observation of the spatial location and relationships

of fluorescently labeled cells within the intact tissue. Therefore, according to the

existing SUMIC tissue deep clearing (TC) methods[25, 26], we modified and improved a

rapid and efficient procedure, which enable rapid single-cell resolution and

quantitative panoptic 3D light-sheet imaging. This procedure not only rapidly clear

maxillary hard tissue, but also directly penetrate the dentin to visualize the dental pulp.

The total time of this procedure was only 2 days including tissue fixation and

decalcification, this ultrafast clearing technique could minimize the impact on

transgenic mice. As shown in Figure S1A-B, we recorded bright-field images of the

maxilla before and after clearing, and our procedure achieved high transparency of the

whole tissue. On this basis, whole-tissue imaging can be achieved, with the

observation of different cell type distribution in spatial 3D structure. As such, dental

pulp and periodontal ligament (PDL) region of M1 were representatively delineated

for 3D reconstruction and analysis. Figure 2C and Video S1 showed the contoured

maxilla M1, including pulp and periodontal ligament (PDL) with virtual dentin shell

(white). The buccal view, coronal view, and radicular view of pulp and PDL showed

that PDGFR-α^+ cells were equidistributional in the whole area of M1 pulp (including

crown pulp and root pulp) and PDL area, almost constitute the whole dental and

periodontal tissue, while NFATc1$^+$ cells were almost exclusively distributed in crown

pulp and scattered existence in PDL. Across the board, the distribution of NFATc1$^+$

cells were significantly less than that of PDGFR-α^+ cells in both pulp and PDL area,

indicated that the expansive range of PDGFR-α^+ cells which may contain NFATc1$^+$

population. As mentioned above, studies have shown that with not all PDGFR-α^+ cells

within long bones exhibiting mesenchymal stem cell characteristics, and only certain

specific subpopulations of PDGFR-α^+ cells can be considered as "real" stem cells. In

addition, our previous study has evidenced NFATc1$^+$PDGFR-α^+ cells as skeletal stem

cells (SSCs) within long bones, played an indispensable role in bone homeostasis and

regeneration. As such, these phenomena we observed from the results of TC

preliminarily indicate a similar situation in dental and periodontal tissue, and the

biological role of NFATc1$^+$PDGFR-α$^+$ cells in homeostasis and regeneration situation

is our major focus in the future research.

In addition, from the section of XZ axis after 3D reconstruction, NFATc1$^+$ cells

were mainly distributed in the alveolar bone area, including the surface edge of

trabecular bone (Figure S1C), which was consistent with RNA-sequencing results in

the previous study[20]. For the cross-sectional analysis of the 3D reconstructed data, the

maximum cross-section of XZ axis which containing the whole pulp and PDL and the

XY axis of the root pulp were selected (Figure 2E). However, the image is not ideal,

which due to: 1) the editing efficiency DNA recombinase–mediated lineage-tracing

system have limitations; 2) the less amount presence of NFATc1$^+$ cells in the tissue

guaranteed the week signals; 3) the as-describe TC method may cause poor

penetration of the td-tomato fluorescence signals.

Therefore, in order to obtain high resolution image to understand more detailed

information, we still need to rely on the traditional serial section technology (Figure

3A). A total of 121 slices were collected in this maxilla of *Pdgfr-α^{CreER}; Nfatc1^{DreER};*

LGRT mice sample, from the jaw root PDL just appeared, to the buccal root PDL

completely disappeared (10 μm/slice) (Figure S2). As shown in Figure 3B and Figure

S3, although time consuming, the images obtained by confocal imaging can clearly

obtain the distribution of PDGFR-α$^+$ and NFATc1$^+$ cells from the single cell level.

However, it is difficult to determine the presence of PDGFR-α$^+$&NFATc1$^+$ cells

spatiotemporally. So, we also used Imaris to 3D-reconstruct these 121 images. Figure

3C and Video S2 showed the contoured M1 of maxilla, including pulp and PDL with

virtual dentin shell (white). Figure 3D was the buccal view, coronal view, and

radicular view of pulp and PDL, respectively. Different from the whole imaging

obtained by TC and light-sheet, the td-tomato signal reconstructed by traditional serial

section-based confocal imaging method was very conspicuous, even in root pulp,

which were hardly obtained by TC imaging (Figure S4). Same as TC-based

reconstructed results, PDGFR-α$^+$ cells almost constituted the whole structure of pulp

and PDL, with NFATc1$^+$ cells as subpopulation. However, due to the stratification of

slices, the sample integrity was poor, leading to discontinuities in the z-axis. (Figure

S5), which may render some biases in the results. To enhance the comprehensive and

accurate display of the reconstruction results and to mitigate the potential errors that

may arise from relying on single reconstruction method, we employed an alternative

3D reconstruction method—DICOM-3D[27]. This method is based on sequential 2D

DICOM images and utilizes 3D reconstruction and visualization technology to

generate a stereoscopic 3D image with intuitive effects, which was a comparatively

straightforward and highly efficient approach. We transformed the serial IF images

into DICOM format and subsequently reconstruct it, and the same conclusion can be

drawn, namely, PDGFR-α^+ cells almost constituted the whole structure of pulp and

PDL, with NFATc1$^+$ cells as subpopulation (Figure S6).

The previous sequencing analyses have reported the expression of NFATc1 in

mandible and periodontal tissues[20]. However, this evidence is limited to RNA-level

sequencing, lacking in vivo data support. Therefore, with assistance of multiple

genetic recombination systems, we next examined the expression pattern of NFATc1

in dental and PDL region of mandible M1, simultaneously observed the distribution

atlas of PDGFR-α^+ cells. As shown in Figure 4A, the pulse timing points were same

as above. Figure 4C and Video S3 showed the contoured mandible M1, including pulp

and PDL with virtual dentin shell (white) in the buccal view, coronal view, and

radicular view of pulp and PDL, respectively. According to the 3D-reconstruction

results of 88 consecutive slices, it can be concluded that PDGFR-α^+ cells almost

constituted the whole structure of dental pulp and periodontal ligament, while

NFATc1$^+$ cells most distributed in the pulp angle in the dental pulp tissue, and also

scattered distributed in PDL region (Figure 4C&E and Figure S7). As shown in Figure

4C and Figure S8, the images obtained by confocal imaging can clearly obtain the

distribution of PDGFR-α^+ and NFATc1$^+$ cells from the single cell level.

To further understand the presence of PDGFR-α^+ cells & NFATc1$^+$ cells and their

progeny cells after long-term tracking, we set a tracing timing points (Figure 5A). The

results showed that nearly the whole pulp and PDL were composed of PDGFR-α^+

cells, and NFATc1$^+$ cells were also scattered present in the root pulp and PDL region

of maxilla M1 (Figure 5B-D, Video S4). Statistical data quantitatively showed that

after 11 days of tracing, the number of PDGFR-α$^+$ cells, NFATc1$^+$ cells,

PDGFR-α$^+$NFATc1$^+$ cells, and their progeny cells increased in both dental pulp and

PDL (Figure S9). Specifically, the number of NFATc1$^+$ cells in PDL area after tracing

11 days was 2.19-fold that of pulse NFATc1$^+$ cells. Further, we also conducted

cross-sectional analysis of the 3D reconstructed data, and selected the maximum

cross-section of XZ axis which containing the whole pulp and PDL and the XY axis

of the root pulp, respectively (Figure 5E). The results were similar to Figure 2E. From

the cross section, only the Zs-Green fluorescence signal of PDGFR-α$^+$ cells can be

observed and analyze while the td-tomato signal was illegible in target area, which

was unable to distinguish from background interference signal. In the process of

sample preparation, the soft tissue on the surface needs to be removed as much as

possible before fixation, otherwise a similar situation will occur. However, for

periosteum and other membranous structures that needed to be studied, the method of

TC seems not suitable.

The tracing samples were also serially sliced and 3D reconstructed (Figure 6 and

Figure 7). A total of 117 slices were collected in maxilla of tracing sample (Figure

S10) and a total of 120 slices were collected in mandible (Figure S15) from the jaw

root PDL just appeared, to the buccal root PDL completely disappeared (10 μm/slice).

As shown in Figure 6B, Figure S11 and Figure 7B, Figure S16, the images obtained

by confocal imaging can clearly obtain the distribution of PDGFR-α$^+$ and NFATc1$^+$

cells from the single cell level. Consistent with the quantification of TC-based

imaging results (Figure S9), the number of PDGFR-α$^+$ cells and NFATc1$^+$ cells were

significantly higher than that in pulse group, which indicated that there were a large

number of PDGFR-α$^+$ & NFATc1$^+$ cells and their progeny cells in crown pulp, root

pulp and PDL after 11 days of tracing. In order to determine the existence of

PDGFR-α$^+$NFATc1$^+$ cells spatiotemporally, we also used software to 3D- reconstruct

these 117 images. Figure 6C-D and Video S5 showed the contoured M1 of maxilla,

including pulp and PDL, at buccal view, coronal view, and radicular view. Figure 7C-D and Video S6 showed the contoured mandible M1. Consistent with TC imaging and following quantitative results (Figure S9), the number of PDGFR-α^+NFATc1$^+$ cells increased significantly in both dental pulp and periodontal ligament, similar to the phenomenon previously observed in long bone, which further encouraged us to follow up with this cell population, explore whether it is a specific mesenchymal stem cell population in periodontal tissue, and explore the indispensable role in periodontal homeostasis and regeneration. Worth notably, different from the whole imaging obtained by TC and light-sheet, the tdTomato signal reconstructed by traditional serial section-based confocal imaging method was very conspicuous, no matter in coronal pulp, root pulp, or PDL (Figure S12). We hypothesize that the current light-sheet systems for intact tissue-imaging have inherent limitations in capturing tdTomato signals, which become more evident in tissues with inherently low fluorescence strengths (in this work, due to the limitations of editing efficiency in DNA recombinase mediated lineage-tracing system, which guaranteed weaker tdTomato signal compared to ZsGreen). In contrast, traditional confocal imaging techniques do not encounter such issues.

However, due to the stratification of slices, the sample integrity was poor, leading to discontinuities in the z-axis. (Figure S13), which may render some biases in the results. We also utilized 3D reconstruction of mandible M1 by DICOM-3D (Figure S14). After 11 days tracing, the number of PDGFR-α^+ & NFATc1$^+$ cells and PDGFR-α^+NFATc1$^+$ cells increased significantly (Figure 7), which further supported our above conjecture that NFATc1$^+$ cells were a subset of PDGFR-α^+ cells, and PDGFR-α^+NFATc1$^+$ cells may contribute to mesenchymal stem cell population in dental and PDL regions.

2.2 Cranium and cranial sutures region

As a part of craniomaxillofacial hard tissue, we also intended to explore whether the presence of NFATc1$^+$ and PDGFR-α^+ cells in cranial bone tissue/suture is different from dental and periodontal tissue (our previous study has identified the presence of

NFATc1$^+$ cells in the cranium by single-cell sequencing[28]), therefore, we also obtained

the cranium of *Pdgfr-α^{CreER}; Nfatc1^{DreER}; LGRT* mice (pulse) for advanced TC

imaging to familiarize NFATc1$^+$/PDGFR-α$^+$ cells in cranium and cranial sutures

region. As shown in Video S7-9, PDGFR-α$^+$ cells were widely distributed, including

cranial sutures and bone interstitium, while NFATc1$^+$ cells are mainly distributed in

cranial sutures, partially coincided with the spatial location of PDGFR-α$^+$ cells. The

PDGFR-α$^+$NFATc1$^+$ cells located in the cranial suture were likely to participate in the

physiological function of the cranial suture as a mesenchymal stem cell population.

The improved TC technology and following 3D imaging can deepen the

understanding of specific cell populations in craniomaxillofacial tissues to a certain

extent, and also lay a foundation for our future researches. However, it seemed not

suitable for the study of periosteum, due to the strong auto-fluorescence under the

light film system of soft tissues, which will affect the observation of target cells.

334   3. The hierarchical distribution of NFATc1$^+$/PDGFRα$^+$ cells in IR1 lineage

335   tracing system

From the above *Pdgfr-α^{CreER}; Nfatc1^{DreER}; LGRT* system, the wider range of

PDGFR-α$^+$ cells than NFATc1$^+$ cells were observed, the wider range of PDGFR-α$^+$

cells than NFATc1$^+$ cells were observed, we speculate that there may be a hierarchical

relationship between the two. To further explore the relationship between PDGFR-α$^+$

and NFATc1$^+$ cells in pulp and PDL and their hierarchical presence, respectively, we

established the IR1 lineage tracing model in adult mice to assess the existence of these

cells   in   the   dental   pulp   and   PDL.   The   structure   of   IR1   is

CAG-loxP-rox-Stop-loxP-ZsGreen-Stop-rox-tdTomato,   and   the   first   *Cre-loxP*

recombination would result in ZsGreen expression that removes a *rox* site, preventing

the subsequent Dre-rox recombination in the same cell. The genotypes used were

PDGFR-α^{CreER}; IR1, Nfatc1^{DreER}; IR1 and *PDGFR-α^{CreER}; Nfatc1^{DreER}; IR1* (Figure

8A-B) in which NFATc1$^+$ cells expressed tdTomato and PDGFR-α$^+$ cells were marked

by ZsGreen. Tracing timing was set based on preliminary experimental outcomes. To

comprehensively and accurately observe the distribution of NFATc1$^+$ and PDGFR-α$^+$

cells in the pulp and PDL, we employed 3D reconstruction technology based on serial

sections (Figure 8C-E, Video S10-12 and Figure S17-19, 21). As shown in Figure

8C-E and Figure S20, the images obtained by confocal imaging can clearly obtain the

distribution of ZsGreen$^+$ and tdTomato$^+$ cells from the single cell level. In *Nfatc1^{DreER};*

IR1 group, most tdTomato$^+$ cells distributed in PDL region, while only sparse and

punctate distribution in dental pulp, mostly in coronal pulp. These results were

consistent with *Pdgfr-α^{CreER}; Nfatc1^{DreER}; LGRT* mice. Moreover, since the

recombinase recognition sites are interleaved (*loxP–rox–loxP–rox*), recombination by

one system will naturally remove a recognition site of the other system, rendering its

reporter gene inactive for further recombination. The results showed no tdTomato$^+$

cells or ZsGreen$^+$ cells were detected in the *Pdgfr-α^{CreER}; IR1* or *Nfatc1^{DreER}; IR1*

group respectively demonstrating the feasibility and accuracy of the *IR1* system. In

the *PDGFR-α^{CreER}; Nfatc1^{DreER}; IR1* group, the distribution of Nfact1$^+$ cells in PDL

were significantly fewer in number than in the *Nfatc1^{DreER}; IR1* group, while no

significant difference in pulp area. In addition, PDGFR-α$^+$ cells were noted in both the

dental pulp and PDL, with a wider distribution than NFATc1$^+$ cells, predominantly in

the odontoblastic layer of the pulp and also present in the pulp core. In the PDL, they

were distributed from the gingival direction down to the root apex. These phenomena

illustrated that PDGFR-α$^+$ cells were the precursor cell of NFATc1$^+$ cells in PDL

region. By contrast, there was almost no difference in the number of NFATc1$^+$ cells

between *Nfatc1^{DreER}; IR1* group and *PDGFR-α^{CreER}; Nfatc1^{DreER}; IR1* group in dental

pulp tissue, which was insufficient to certificate the hierarchical relationship of

PDGFR-α$^+$ and NFATc1$^+$ cells in pulp area.

4. Identification of PDGFR-α$^+$ and NFATc1$^+$ population in pulp and PDL

4.1 Ablation of PDGFR-a$^+$ cells disrupt the morphology of dental pulp and

periodontal mesenchymal tissues

The results from the aforementioned lineage tracing experiments showed that

PDGFR-α$^+$ cells constitute a significant component of both dental pulp and

periodontal tissues. Additionally, the hierarchical relationship experiments revealed

that a portion of NFATc1$^+$ cells in the periodontal ligament derives from PDGFR-α$^+$

progenitor cells. Therefore, investigating the role of PDGFRα$^+$ cells in dental pulp and

periodontal tissues has become more urgent. To explore the role of PDGFR-α$^+$ cells in

dental pulp and PDL tissues, we performed the cell ablation assay using *Pdgfr-α^{CreER};*

DTA mice with tamoxifen administered for 12 weeks (Figure 9A). The *Pdgfr-αCreER;*

DTA mice strain harbors a *loxP*-flanked termination cassette within the ubiquitous

ROSA26 locus, controlling the expression of DTA. Tamoxifen administration was

employed for in vivo toxin-mediated cell ablation (Figure 9B). Through H&E staining

and Masson trichrome staining, we observed that in the control group, the odontoblast

cell layer adjacent to the dentin consisted of densely arranged columnar cells both in

the coronal or radicular pulp. However, this layer of odontoblasts was disrupted and

nearly disappeared after ablation. Additionally, ablation of PDGFR-α$^+$ cells in adult

DTA mice also showed significant shrinkage in the central area of dental pulp tissue,

specifically within the region of the pulp core, along with a substantial reduction in

cell numbers (Figure 9C-D). H&E staining confirmed a decrease in periodontal

composition following PDGFR-α$^+$ cell clearance (Figure 9E). Furthermore, Masson's

staining results revealed that the diminished portion primarily comprised periodontal

fibrous tissue (Figure 9F). Therefore, it can be inferred that PDGFR-α$^+$ cells

predominantly constitute the interstitial components of both the periodontium and

dental pulp. PDGFR-α$^+$ cells within the dental pulp play a crucial role in the

formation of the odontoblast cell layer and contribute significantly to most of the core

tissue, while PDGFR-α is primarily involved in shaping periodontal components,

particularly fibrous tissue.

4.2 PDGFR-α$^+$ & NFATc1$^+$ cells including MSCs and hematopoietic population in

pulp and PDL

To identify the population of PDGFR-α$^+$ and NFATc1$^+$ co-expressing cells in the pulp

and periodontal ligament (PDL), we generated *Pdgfr-a^{CreER}; Nfatc1^{DreER};*

R26-LSL-RSR-tdT-DTR (LRTD) mice. In these mice, Dre-rox and Cre-loxP

recombination events enable the simultaneous expression of tdTomato and DTR

specifically in pulp and PDL cells. In this experiment, we utilized the LRTD mice

solely to label double-positive cells, so DTX was not administered to ablate the cells.

To achieve high labeling efficiency, we treated the mice with tamoxifen twice (Figure

S22A). Strong tdTomato signals were detected in both the PDL (Figure S22B) and

pulp (Figure S22C). With respect to the MSC-specific marker AlphaV, we observed

AlphaV$^+$tdTomato$^+$ cells in both regions. Additionally, CD45$^+$ (hematopoietic marker)

tdTomato$^+$ cells were also present in these areas (Figure S22B, C). These findings

suggest that the population of PDGFR-α^+ and NFATc1$^+$ co-expressing cells is

heterogeneous.

Discussion

PDGFR-α and NFATc1, commonly acknowledged as markers for stem cells, are

recognized for their contributions to stem cell characteristics. Our previous study also

identified PDGFR-α and NFATc1 cells as SSCs in long bones. The conventional

approach to studying the expression patterns of two cell populations involved

techniques such as immunofluorescence or immunohistochemical staining, the

generation of single-enzyme-cut editing gene mice, or a combination thereof.

However, these traditional methods were often susceptible to factors like antibody

potency, leading to incomplete and inaccurate depiction of the relationship between

the two population of cells. In this study, we have developed two innovative genetic

instruments, H11-LGRT and IR1, to facilitate distinct in vivo genetic lineage tracing

assays. These tools were specifically designed for precision mapping of cellular

lineages and have been utilized, for the first time, within the oral and maxillofacial

regions. The arrangement of different recombination sites in relation to reporter genes

is crucial for their use in in vivo tracking of multiple cell lineages. The ablation assay

demonstrated that PDGFR-α nearly labeled almost mesenchyme cells, which was

similar to results of H11-LGRT, noting that H11-LGRT is an available and reliable

tool. Our results demonstrate that both IR1 and H11-LGRT are capable of tracing the

cell fate mapping of various cells within the oral and craniofacial regions.

Dental pulp and the periodontal supporting tissues surrounding the tooth can be

regarded as a relatively integrated biological functional unit. In the research of

regeneration and repair mechanisms of dental pulp and PDL, the roles of periodontal

and dental pulp stem cells are indispensable[8]. These cells play a vital role in the oral

and maxillofacial regions, not only participating in routine tissue maintenance but also

acting as key players in the self-repair process after injury[29, 30]. Dental pulp stem cells

have the potential to regenerate dental pulp tissue, while periodontal stem cells are

crucial for the regeneration of periodontal tissues, including alveolar bone, PDL and

gingiva[31, 32]. The interactions between these cell types and their prospects in tissue

engineering open new strategies and therapeutic approaches for modern dental

treatments.

In our study, H11-LGRT systems have illuminated distinct expression patterns of

PDGFR-α^+, NFATc1$^+$ and PDGFR-α^+&NFATc1$^+$ cells in PDL and dental pulp,

indicating their diverse characteristics and roles. In PDL tissue, pulse data

demonstrated widespread and abundant expression of PDGFR-α single-positive cells

as well as NFATc1 single-positive cells, with no significant alteration in expression

pattern or quantity after lineage tracing. Consequently, we conclude that in

periodontal ligament and dental pulp tissues, PDGFR-α single-positive and NFATc1

single-positive cells primarily label intrinsic periodontal mesenchyme in PDL.

Conversely, PDGFR-α^+&NFATc1$^+$ cells exhibited a more confined localization in

PDL. The tracing data clearly illustrated that PDGFR-α^+&NFATc1$^+$ cells successfully

gave rise to numerous progenies, which become predominant constituents within the

periodontal ligament. In pulp tissue, the distribution of PDGFR-α single-positive cells

was similar as that in PDL, primarily labeled odontoblast cell layer and there was not

a significant increase in ZsGreen signal after tracing assay (Figure. 5-7). However,

distinct outcomes were observed for NFATc1$^+$ single-positive cells, as tracing

experiments revealed an increased generation of progeny cells from those

homogeneous NFATc1 within the intrinsic dental pulp. Consistently, the results for

PDGFR-α^+&NFATc1$^+$ cells mirrored those of NFATc1$^+$ single-positive cells. This

observation may imply a niche-specific role for these cells especially

PDGFR-α^+&NFATc1$^+$ cells in maintaining critical functions such as regeneration.

This could indicate a more generalized role in tissue maintenance and repair,

potentially contributing to the regeneration of both soft and hard tissues.

In order to investigate the hierarchical relationship between PDGFR-α^+ and

NFATc1$^+$ in the dental pulp and PDL, we developed IR1 system (Figure. 8). Our

results revealed distinct hierarchical relationships in PDL and dental pulp. In

periodontal tissues, compared to the control group (*Pdgfr-α^{CreER}; IR1* and *Nfatc1^{DreER};*

IR1), there was significantly less tdTomato$^+$ cells distribution in the *Pdgfr-α^{CreER};*

Nfatc1^{DreER}; IR1 group while the signal of ZsGreen did not decrease significantly.

Based on the function of the IR1 system, we could conclude that in the periodontal

ligament, PDGFR-α^+ cells serve as precursor cells for some NFATc1$^+$ cells. However,

in dental pulp tissue, there was almost no difference in tdTomato$^+$ expression between

the experimental and control groups, indicating that PDGFR-α^+ cells cannot be

considered as precursor for NFATc1$^+$ cells in the dental pulp, or there may not be a

clear hierarchical relationship between the two. The sparse presence of NFATc1$^+$ cells

in both dental pulp and PDL, as observed in the *Nfatc1^{DreER}; IR1* group, raised

questions about the plasticity and differentiation potential of these cells. Are they

primarily quiescent, being activated only upon injury, or do they continuously

contribute to tissue homeostasis? Further studies, including functional assays and

injury models, would be instrumental in clarifying the dynamic nature of these cells.

Cell ablation and immunofluorescence staining experiments further characterized

the types and functions of PDGFR-α^+/PDGFR-α^+&NFATc1$^+$ populations. After

ablating PDGFR-α^+ cells, we observed damage to the odontoblast layer and shrinkage

of the pulp core in dental pulp tissue, indicating that PDGFR-α^+ cells contribute to the

composition of dental pulp tissue, particularly the odontoblast layer (Figure. 9C, D).

In the periodontal ligament, we noted a reduction and destruction of collagen fibers,

suggesting a role for PDGFR-α^+ cells in periodontal tissue structure (Figure. 9E, F).

Previous results confirmed the presence of double-positive cells in both dental pulp

and periodontal tissues and provided insights into their hierarchical relationships in

the periodontal ligament (Figure. 8). To further investigate the double-positive cell

population, we developed an inducible dual-editing enzyme reporter system to label

these cells with tdTomato signals. Using AlphaV as a marker for mesenchymal stem

cells (MSCs) and CD45 for hematopoietic cells, we found that double-positive cells

included components of both MSCs and hematopoietic cells (Figure S22B, C),

indicating a heterogeneous population. Further experiments are necessary to

determine whether the predominant role in this co-positive MSC population is played

by PDGFR-α^+ or NFATc1$^+$ and to clarify the specific functions of these cells in the

future.

For the traditional sectioning & confocal imaging procedure, it can obtain a

highly distinct single-layer image. However, it is troubled to determine whether the

different cell populations overlap in spatial location. At the same time, the

experimental results were deeply affected by the differences between different section,

which may cause error. It is also impossible to determine the amount and spatial

position of a certain cell population in the whole tissue. In this study, we utilized two

different techniques to map the distribution of aimed cells. We successfully achieved a

3D cellular-level visualization of PDL and pulp from *Pdgfr-α^{CreER}; Nfatc1^{DreER}; LGRT*

mice by TC-based advanced imaging and 3D reconstructions based on continuous

sections provide a more accurate representation of spatial distribution.TC-based

advanced imaging procedure can clearly visualize its 3D structure, reconstruct the

whole across latitudes, and understand the spatial position and expression of each

structure, which could avoid the bias of traditional single-layer slicing may cause, and

provides a more intuitive and objective description of the existing situation. However,

our results demonstrated TC still has some limitations. During the experiment, we

found that the signal of tdTomato in the pulp and PDL area was poor and could not be

distinguished from autofluorescence through the imaging of light-sheet system, which

may due to the limitations of editing efficiency in DNA recombinase–mediated

lineage-tracing system, which guaranteed weaker tdTomato signal compared to

Zsgreen. In addition, the total amount of NFATc1$^+$ SSCs in these regions was limited,

with the highly distribution in the alveolar bone, jaw bones, and trabecular bone

surface edge, which guaranteed less visible signal in the target region. Another

possibility is that the as-describe TC method may cause poor penetration of the

tdTomato fluorescence signals. The above results prompted that we should minimize

the sample processing time to avoid fluorescence depletion to the greatest extent when

using dual DNA recombinases-based genetic lineage tracing system for whole tissue

imaging after TC treatment.

The 3D sections reconstruction results, however, effectively addressed the issue

of weak tdTomato signal and provide a clearer visualization of the distribution of

ZsGreen and tdTomato signals. For example, the tdTomato signal in the root pump,

which was almost completely unobservable by TC-based imaging, can be clearly seen

using confocal imaging and 3D reconstruction (Figure 3C-D, Figure 6C-D, and Figure

S4, Figure S12). However, compared to TC, the quality of 3D reconstruction of

sections still relies on the angle and quality of the sections, with the section angle

having a significant impact on the reconstruction outcome. In addition, because the

slice itself has a certain thickness (10 μM in this study), which leads to the appearance

of discontinuous in the final reconstructed image, and the aesthetics and accuracy

could be affected to a certain extent. Also, unavoidable tissue damage during the

sectioning process may result in the loss of some information. Therefore, a variety of

different information could be obtained through two different imaging technologies,

which prompt us to use the advanced experimental procedure according to the actual

purpose.

The use of advanced genetic tools has also shed light on the limitations and

challenges associated with current lineage tracing methodologies[33]. For instance, the

potential for recombinase leakiness or off-target effects necessitates the inclusion of

stringent controls and corroborative experimental approaches[34]. As such, the

refinement of these tools and the development of additional specific markers are

required to enhance the accuracy of cell fate mapping.

In conclusion, our study developed multiple DNA recombinases based on genetic

lineage tracing and improved their accuracy and usability cooperating with advanced

imaging techniques in order to map the atlas of NFATc1$^+$/ PDGFR-α^+ inclusive,

exclusive and hierarchical distribution in dental and periodontal mesenchyme. Our

study underscored the importance of NFATc1$^+$ and PDGFRα^+ cells in the homeostasis

and regeneration of oral and maxillofacial tissues. The nuanced understanding of their

distribution and potential roles paves the way for novel regenerative strategies that

could harness the intrinsic capabilities of these cells. Future research should focus on

the functional characterization of these cells, their responsiveness to injury, and the

interplay of signaling pathways that regulate their activity, which will be crucial for

translating these findings into clinical applications.

Materials and methods

Methods

Ethics Statement and Animals

All animal procedures were reviewed and approved by Ethical Committees of West

China School of Stomatology, Sichuan University (WCHSIRB-D-2017-041). Relative

surgical models were conducted according to approved guidelines set by State Key

Laboratory of Oral Diseases, West China Hospital of Stomatology.

Nfatc1^{DreER} and *IR1* strain were kindly provided by Prof. Bin Zhou (Chinese Academy

of Sciences), which were also reported in our previous study (Yu et al. 2022).

Pdgfr-α^{CreER} (Stock 018280) was purchased from the JAX Lab,

LSL-ZsGreen-RSR-tdTomato (LGRT) was purchased from Shanghai Model

Organisms Center (Strain# NM-KI-200319), and DTA mice were purchased from

JAX Lab (Stock# 009669).

In vivo mouse studies

All the transgenic mice discussed in this work are adults around 12-14 weeks, which

were intraperitoneally injected with tamoxifen (TAM) according to our previous

report[19].

RT-qPCR

For RT-qPCR total RNA was extracted using the TRIzolTM (Invitrogen) according to

the manufacturer's protocol. Complementary DNA was synthesized by using the

HiScript III RT SuperMix for qPCR (Vazyme) in accordance to user manuals. Then

quantitative real-time PCR was performed in triplicate by using AceQ Universal

SYBR qPCR Master Mix (Vazyme) for PCR reactions on an iCycler Real-Time

Detection System (BioRad, USA).

Maxillae and cranium collection, fixation, and decalcification for light sheet

imaging

Fresh dissected tissues were washed with PBS (Solarbio), and then rapidly transferred

to 4% PFA (Sigma-Aldrich) for 4 hours. Before fixation, the muscle and fat attached

to the tissue surface should be removed as much as possible, otherwise there will be

strong interference signal during imaging. The tissues were then washed three times

with PBS at 4 °C on a rocker platform for 15 min each time. For decalcification,

tissues were incubated with 0.5 M EDTA solution (pH 7.4) at 4 °C for 24 h. After

decalcification, the samples were washed three times with PBS on the rocker (15 mi

each time). The tissues were then submersed in 50%, 80% and 100% ethanol gradient

for 30 min each to dehydrated. 100% ethanol was changed twice after every 20 min.

Then the samples were immersed in 5% (V/V) H_2O_2 for 2 h (Sigma-Aldrich, H1009)

for bleaching [35].

Clearing of maxillae and cranium

After bleaching, the samples were rinsed three times with Ethyl cinnamate (ECi)

(Sigma-Aldrich, 112372) for 5 min each at room temperature.

Histological Preparation and Staining

Specifically, the tissues were fixed in 4% PFA at 4 ☐ for 24 hours. After fixation, the maxilla was immersed in 30% sucrose solution at 4 ° C overnight. Then the samples were embedded in OCT tissue freezing medium (Leica) and placed on liquid nitrogen to conduct non-decalcified frozen sectioning at 7 to 10 μm thickness. The slices were treated with DAPI at room temperature for 1h. Finally, the slices were washed with PBS for 3 times, 10 minutes each time, and sealed for detection. For H&E staining and Masson trichrome staining, decalcified samples were dehydrated in graded ethanol and embedded in paraffin. Mandibles were sectioned into 6 μm slices using the microtome (Leica RM2255). H&E staining was performed according to the manufacturer's instruction (Biosharp). Masson trichrome staining was performed according to the manufacturer's instruction (Solarbio).

Imaging and 3D reconstruction of maxillae and mandible sections

Digital pathological system (Olympus FV3000) was used to scan all the stained sections and reconstructed by following the previously described protocol (Wu et al. 2020). The light sheet fluorescence microscopy-Zeiss Lightsheet Z.7 was used to acquire clearing samples microscopic image stacks.

Image analysis and quantifications

Slices/Z-stacks of images acquired on the light sheet and confocal microscope were processed and reconstructed in three dimensions with Imaris software (version 10.0.1). Imaris, Adobe Photoshop 25.0, and ImageJ were used for image processing and analysis in line with the journal's guidance for image processing. Quantification of cell numbers was done on a z stack of images in Imaris using the automatic spot detection feature.

3D surface reconstruction

3D surface rendering in images was applied using the surface module in Imaris.

Statistical analysis

All data were presented as the mean ± standard error of the mean (SEM). A two-tailed Student's t-test was used for comparison between groups. *P < 0.05* was considered statistically significant. N.S. stands for not significant.

Impact statement

This work utilized multi-enzyme cutting lineage tracing system along with advanced imaging methods to systematically map the distribution atlas of NFATc1$^+$ and PDGFR-α$^+$ cells in dental and periodontal mesenchyme.

Acknowledgment

This work was supported by National Key Research and Development Program of China 2023YFC3605600 (L.Y. and F.Y.); National Natural Science Foundation of China 82201045 (F.Y.) and 82100982 (F.L.); and Young Elite Scientist Sponsorship Program by CAST (2022QNRC001 to F.Y.).

Conflict of Interests

All authors declare no conflicts of interest.

Author Contributions

F.Y. conceived the study; X.Y., C.H., C.Y., and F.L. performed research; X.Y, C.H., and F.L. analyzed data; X.Y. and C.H. wrote the manuscript; and L.Y., F.L. and F.Y. reviewed and edited the manuscript. X.Y contributed equally with C.H.

References

1. Andrae J., Gallini R., Betsholtz C. (2008) **Role of platelet-derived growth factors in physiology and medicine** *Genes Dev* 22:1276-1312
 https://doi.org/10.1101/gad.1653708

2. Dereka X.E., Markopoulou C.E., Vrotsos I.A. (2006) **Role of growth factors on periodontal repair** *Growth Factors* 24:260-267
 https://doi.org/10.1080/08977190601060990

3. Heldin C.H., Lennartsson J. (2013) **Structural and functional properties of platelet-derived growth factor and stem cell factor receptors** *Cold Spring Harb Perspect Biol* 5:a009100

https://doi.org/10.1101/cshperspect.a009100

4. Mijiritsky E., Assaf H.D., Peleg O., Shacham M., Cerroni L., Mangani L. (2021) **Use of PRP, PRF**
and CGF in Periodontal Regeneration and Facial Rejuvenation-A Narrative Review
Biology (Basel) 10:317

https://doi.org/10.3390/biology10040317

5. Komatsu K., Ideno H., Shibata T., Nakashima K., Nifuji A. (2022) **Platelet-derived growth**
factor-BB regenerates functional periodontal ligament in the tooth replantation *Sci Rep*
12:3223

https://doi.org/10.1038/s41598-022-06865-6

6. Zhang M., Jiang F., Zhang X., Wang S., Jin Y., Zhang W., Jiang X. (2017) **The Effects of**
Platelet-Derived Growth Factor-BB on Human Dental Pulp Stem Cells Mediated
Dentin-Pulp Complex Regeneration *Stem Cells Transl Med* 6:2126-2134

https://doi.org/10.1002/sctm.17-0033

7. Woelfel J.B., Scheid R.C. (1997) **Dental anatomy** (M). Williams & wilkins

8. Ouchi T., Nakagawa T. (2020) **Mesenchymal stem cell-based tissue regeneration therapies for**
periodontitis *Regen Ther* 14:72-78

https://doi.org/10.1016/j.reth.2019.12.011

9. Chai Y., Bringas P., Jr., Mogharei A., Shuler C.F., Slavkin H.C. (1998) **PDGF-A and PDGFR-alpha**
regulate tooth formation via autocrine mechanism during mandibular morphogenesis in
vitro *Dev Dyn* 213:500-511

https://doi.org/10.1002/(SICI)1097-0177(199812)213:4<500::AID-AJA14>3.0.CO;2-A

10. Guo T., Pei F., Zhang M., Yamada T., Feng J., Jing J., Ho T.V., Chai Y. (2024) **Vascular**
architecture regulates mesenchymal stromal cell heterogeneity via P53-PDGF signaling
in the mouse incisor *Cell Stem Cell*   https://doi.org/10.1016/j.stem.2024.04.011

11. Xu X., Bringas P., Jr., Soriano P., Chai Y. (2005) **PDGFR-alpha signaling is critical for tooth**
cusp and palate morphogenesis *Dev Dyn* 232:75-84

https://doi.org/10.1002/dvdy.20197

12. Cui Y., Ji W., Gao Y., Xiao Y., Liu H., Chen Z. (2021) **Single-cell characterization of monolayer**
cultured human dental pulp stem cells with enhanced differentiation capacity *Int J Oral*
Sci 13:44

https://doi.org/10.1038/s41368-021-00140-6

13. Morikawa S., Mabuchi Y., Kubota Y., Nagai Y., Niibe K., Hiratsu E., Suzuki S., Miyauchi-Hara C.,
Nagoshi N., Sunabori T., Shimmura S., Miyawaki A., Nakagawa T., Suda T., Okano H.,
Matsuzaki Y. (2009) **Prospective identification, isolation, and systemic transplantation of**
multipotent mesenchymal stem cells in murine bone marrow *J Exp Med* 206:2483-2496

https://doi.org/10.1084/jem.20091046

14. Uezumi A., Fukada S., Yamamoto N., Ikemoto-Uezumi M., Nakatani M., Morita M., Yamaguchi A.,
Yamada H., Nishino I., Hamada Y., Tsuchida K. (2014) **Identification and characterization**
of PDGFRalpha+ mesenchymal progenitors in human skeletal muscle *Cell Death Dis*
5:e1186

https://doi.org/10.1038/cddis.2014.161

15. Kfoury Y., Scadden D.T. (2015) **Mesenchymal cell contributions to the stem cell niche** *Cell Stem*
Cell 16:239-253

https://doi.org/10.1016/j.stem.2015.02.019

16. Ning K., Yang B., Chen M., Man G., Liu S., Wang D.E., Xu H. (2022) **Functional Heterogeneity**
of Bone Marrow Mesenchymal Stem Cell Subpopulations in Physiology and Pathology
Int J Mol Sci 23:11928

https://doi.org/10.3390/ijms231911928

17. Changmeng Z., Hongfei W., Cheung M.C., Chan Y.S., Shea G.K. (2023) **Revealing the**
developmental origin and lineage predilection of neural progenitors within human bone
marrow via single-cell analysis: implications for regenerative medicine *Genome Med*
15:66

https://doi.org/10.1186/s13073-023-01224-0

18. Koide Y., Morikawa S., Mabuchi Y., Muguruma Y., Hiratsu E., Hasegawa K., Kobayashi M., Ando
K., Kinjo K., Okano H., Matsuzaki Y. (2007) **Two distinct stem cell lineages in murine bone**
marrow *Stem Cells* 25:1213-1221

https://doi.org/10.1634/stemcells.2006-0325

19. Yu F., Li F., Yu P., Zhou B., Ye L. (2022) **Identification and characterization of NFATc1(+)**
skeletal stem cells in bone regeneration *Cell Rep* 41:111599

https://doi.org/10.1016/j.celrep.2022.111599

20. Nassif A., Lignon G., Asselin A., Zadikian C.C., Petit S., Sun H.W., Klein C., Ferre F.C., Morasso
M.I., Berdal A., Fournier B.P.J., Isaac J. (2022) **Transcriptional Regulation of Jaw**
Osteoblasts: Development to Pathology *J Dent Res* 101:859-869

https://doi.org/10.1177/00220345221074356

21. Buckingham M.E., Meilhac S.M. (2011) **Tracing cells for tracking cell lineage and clonal**
behavior *Dev Cell* 21:394-409

https://doi.org/10.1016/j.devcel.2011.07.019

22. Anastassiadis K., Fu J., Patsch C., Hu S., Weidlich S., Duerschke K., Buchholz F., Edenhofer F.,
Stewart A.F. (2009) **Dre recombinase, like Cre, is a highly efficient site-specific**
recombinase in E. coli, mammalian cells and mice *Dis Model Mech* 2:508-515

https://doi.org/10.1242/dmm.003087

23. Liu K., Jin H., Zhou B. (2020) **Genetic lineage tracing with multiple DNA recombinases: A**
user's guide for conducting more precise cell fate mapping studies *J Biol Chem*
295:6413-6424

https://doi.org/10.1074/jbc.REV120.011631

24. Yao L., Li F., Yu C., Wang H., Wang Y., Ye L., Yu F. (2023) **Chronological and Replicative Aging**
of CD51(+)/PDGFR-alpha(+) Pulp Stromal Cells *J Dent Res* 102:929-937

https://doi.org/10.1177/00220345231158038

25. Antila S., Chilov D., Nurmi H., Li Z., Nasi A., Gotkiewicz M., Sitnikova V., Jantti H., Acosta N.,
Koivisto H., Ray J., Keuters M.H., Sultan I., Scoyni F., Trevisan D., Wojciechowski S.,
Kaakinen M., Dvorakova L., Singh A., Jukkola J., Korvenlaita N., Eklund L., Koistinaho J.,
Karaman S., Malm T., Tanila H., Alitalo K. (2024) **Sustained meningeal lymphatic vessel**
atrophy or expansion does not alter Alzheimer's disease-related amyloid pathology *Nat*
Cardiovasc Res 3:474-491

https://doi.org/10.1038/s44161-024-00445-9

26. Biswas L., Chen J., De Angelis J., Singh A., Owen-Woods C., Ding Z., Pujol J.M., Kumar N., Zeng
F., Ramasamy S.K., Kusumbe A.P. (2023) **Lymphatic vessels in bone support regeneration**

after injury *Cell* 186:382-397 e324

https://doi.org/10.1016/j.cell.2022.12.031

27. Popescu D., Marinescu R., Laptoiu D., Deac G.C., Cotet C.E. (2021) **DICOM 3D viewers, virtual**

reality or 3D printing - a pilot usability study for assessing the preference of orthopedic

surgeons *Proc Inst Mech Eng H* 235:1014-1024

https://doi.org/10.1177/09544119211020148

28. Yu F.Y., Li F.F., Yu P., Zhou B., Ye L. (2022) **Identification and characterization of**

NFATc1+skeletal stem cells in bone regeneration *Cell Reports* 41: https://doi.org/ARTN

111599

10.1016/j.celrep.2022.111599

29. Tomokiyo A., Wada N., Maeda H. (2019) **Periodontal Ligament Stem Cells: Regenerative**

Potency in Periodontium *Stem Cells Dev* 28:974-985

https://doi.org/10.1089/scd.2019.0031

30. Liang Y., Luan X., Liu X. (2020) **Recent advances in periodontal regeneration: A biomaterial**

perspective *Bioact Mater* 5:297-308

https://doi.org/10.1016/j.bioactmat.2020.02.012

31. Chalisserry E.P., Nam S.Y., Park S.H., Anil S. (2017) **Therapeutic potential of dental stem cells** *J*

Tissue Eng 8:2041731417702531

https://doi.org/10.1177/2041731417702531

32. Leyendecker Junior A., Gomes Pinheiro C.C., Lazzaretti Fernandes T., Franco Bueno D. (2018)

The use of human dental pulp stem cells for in vivo bone tissue engineering: A systematic

review *J Tissue Eng* 9:2041731417752766

https://doi.org/10.1177/2041731417752766

33. Liu J., Willet S.G., Bankaitis E.D., Xu Y., Wright C.V., Gu G. (2013) **Non-parallel recombination**

limits Cre-LoxP-based reporters as precise indicators of conditional genetic

manipulation *Genesis* 51:436-442

https://doi.org/10.1002/dvg.22384

34. Zhao H., Zhou B. (2019) **Dual genetic approaches for deciphering cell fate plasticity in vivo:**

more than double *Curr Opin Cell Biol* 61:101-109

https://doi.org/10.1016/j.ceb.2019.07.004

35. Biswas L., Chen J., De Angelis J., Singh A., Owen-Woods C., Ding Z., Pujol J.M., Kumar N., Zeng

F., Ramasamy S.K., Kusumbe A.P. (2023) **Lymphatic vessels in bone support regeneration**

after injury *Cell* 186:382-397.e324

https://doi.org/10.1016/j.cell.2022.12.031

  Figures and figure legends

  Figure 1

  Figure 1. (A) Genetic recombination systems. The initial column lists various reporter

  alleles, while the subsequent second and third columns present the corresponding

  reporter constructs and the recombinases utilized. The final column provides the

  outcomes associated with the respective reporter alleles. Traditional reporter

  mechanisms comprise both single-color and multicolor systems (not shown in the

  figure), created through a singular recombination event. Advanced dual-reporter

configurations encompass intersectional reporters, exclusive reporters, and nested

reporters. (B) The craniofacial hard tissue, including maxilla and mandible of

transgenic mice with multiple DNA recombinases-based genetic lineage tracing

system, were obtained for advanced imaging technologies. (C-a) The tissue deep

clearing procedure based on SUMIC, which needed to go through 4% PFA fixation,

0.5M EDTA decalcification, different concentration of methanol dehydration, 5%

H_2O_2 bleaching, ECI tissue clearing, and finally used the light-sheet system to directly

image the whole tissue. (C-b) The traditional frozen section technology needs fixation,

decalcification, dehydration and other steps, which takes about two weeks.

Figure 2

Figure 2. The whole-mount and high-speed imaging of (A) mouse molar with a tiling

light-sheet microscope. (B) Schematic illustration of lineaging tracing in *Pdgfr-α^{CreER};*

Nfatc1^{DreER}; LGRT mice. The mice were administrated with tamoxifen at D1 and D3,

and sacrificed at D5. (C) The contoured M1 of maxilla, including pulp and PDL with

virtual dentin shell (white) in buccal view (scale bar = 300 μm).   (D) Image stack

was displayed in buccal view, coronal view, and radicular view of pulp and PDL,

respectively (scale bar = 300 μm). (E) An optical slice was acquired on the X-Z (scale

bar = 300 μm) and X-Y direction (scale bar = 400 μm) to display the pulp and PDL.

Figure 3

Figure 3. Using traditional serial section for imaging of maxilla of *Pdgfr-α^{CreER}*;

Nfatc1^{DreER}; LGRT mice (pulse). (A) Operation process of frozen section. Using this

procedure, a total of 121 slices were collected in this sample. (B) Representative

images acquired by confocal microscopy (scale bar = 200 μm). Box 1: coronal pulp,

Box 2: root pulp, Box 3: PDL, scale bar = 100 μm. (C) Maxilla M1 after 3D

reconstruction by imaris, including pulp and PDL with virtual dentin shell (white) in

buccal view (scale bar = 300 μm). (D) Image stack was displayed in buccal view,

coronal view, and radicular view of pulp and PDL, respectively (scale bar = 300 μm).

Figure 4

Figure 4. Using traditional serial section for imaging of mandible of $Pdgfra^{CreER}$;

$Nfatc1^{DreER}$; $LGRT$ mice (pulse). (A) Schematic illustration of lineaging tracing in

$Pdgfr-\alpha^{CreER}$; $Nfatc1^{DreER}$; $LGRT$ mice. (B) Operation process of frozen section. Using

this procedure, a total of 88 slices were collected in this sample. (C) Mandible M1

after 3D reconstruction by Imaris, including pulp and PDL with virtual dentin shell

(white) in buccal view, coronal view and radicular view (scale bar = 200 μm). (E) 3D

reconstruction of mandible M1 by DICOM-3D; in PDL, ZsGreen+ cells in green,

tdTomato+ cells in rose red; in pulp, ZsGreen+ cells in purple, tdTomato+ cells in

blue. The image stack was also displayed in buccal view, coronal view, and radicular

view of pulp and PDL, respectively. (scale bar = 200 μm).

Figure 5

Figure 5. The observation and analysis of lineage tracing of PDGFR-α^+ & NFATc1$^+$

cells in M1 by whole-mount and high-speed imaging. (A) The flowchart of tracing.

The mice were administrated with tamoxifen at D1 and D3, and sacrificed at D11. (B)

The 3D images of contoured M1 of maxilla, including pulp and PDL with virtual

dentin shell (white) in buccal view (scale bar = 300 μm). (D) Image stack was

displayed in buccal view, coronal view, and radicular view of pulp and PDL, respectively. (E) An optical slice was acquired on the X-Z (scale bar = 400 μm) and X-Y direction to display the pulp and PDL (scale bar = 300 μm).

Figure 6

Figure 6. Using traditional serial section for imaging of maxilla of *Pdgfr-α^{CreER}; Nfatc1^{DreER}; LGRT* mice (tracing 11 days). (A) Using this procedure, a total of 117 slices were collected in this sample. (B) Representative images acquired by confocal microscopy (scale bar = 200 μm). Box 1: coronal pulp; Box 2: root pulp; Box 3: PDL (scale bar = 50 μm). (C) Maxilla M1 after 3D reconstruction by imaris, including pulp and PDL with virtual dentin shell (white) in buccal view (scale bar = 300 μm). (D) Image stack was displayed in buccal view, coronal view, and radicular view of pulp and PDL, respectively (scale bar = 300 μm).

Figure 7

Figure 7. H11-LGRT tracing distinct cell populations simultaneously in the pulp and PDL. (A) Schematic illustration of pulse and lineaging tracing in $Pdgfr\text{-}\alpha^{CreER}$; $Nfatc1^{DreER}$; $LGRT$. (B) 3D reconstruction of mandible M1 using Imaris, including pulp and PDL with virtual dentin shell (white) (scale bar = 200 μm). The image stack was displayed in buccal view, coronal view, and radicular view of pulp and PDL, respectively. (C) 3D reconstruction of mandible M1 using DICOM-3D, in which the distribution of NFATc1$^+$ cells in pulp and PDL tissues can be more precisely perceive (scale bar = 200 μm).

Figure 8

Figure 8. IR1 tracing distinct cell populations simultaneously in the pulp and PDL.

(A) Schematic illustration of lineaging tracing in *Pdgfr-α^CreER^; IR1, Nfatc1^DreER^; IR1*

and *Pdgfr-α^CreER^; Nfatc1^DreER^; IR1* mice. (B) Schematic diagram of the IR1 working

principle. (C) 3D reconstruction of maxilla M1 using Imaris, including pulp and PDL

with virtual dentin shell (white) (scale bar = 200 μm). The image stack was displayed

in buccal view, coronal view, and radicular view of PDL (D) and pulp (E),

respectively.

Figure 9

Figure 9. Ablation of PDGFR-α⁺ Cells Disrupts the morphology of dental pulp and

periodontal ligament tissues. A) Schematics of tamoxifen induction. B) Schematic

diagram of the DTA working principle. C, E) Representative H&E images of pulp C)

and PDL E) of mandible M1 in *Pdgfr-a^{CreER}; DTA* and control mice. D, F) Masson

trichrome staining of pulp D) and PDL F) of mandible M1 in *Pdgfr-a^{CreER}; DTA* and

control mice. Arrows C, D) indicate the odontoblast cell layer. Dotted lines E, F)

outline ROI of the PDL. D: dentin; AB: alveolar bone. Scale bar: 50 μm.