A functional interleukin-4 homolog is encoded in the genome of infectious laryngotracheitis virus: unveiling a novel virulence factor

Jeremy D. Volkening[a], Stephen J. Spatz[b],∗, Maricarmen García[c], Ana C. Zamora[c], Kenneth S. Rosenthal[d], April Skipper[c], Julia Blakey[b], Teresa A. Ross[b]

[a] BASE₂BIO, Oshkosh, WI, USA
[b] US National Poultry Research Center, ARS-USDA, Athens, GA, USA
[c] University of Georgia, College of Vetarinary Medicine, Department of Population Health, Poultry Diagnostic Research Center, Athens, GA, USA
[d] Augusta University/University of Georgia Medical Partnership, Department of Biomedical Sciences, Athens, GA, USA

Abstract

Herpesviruses have evolved numerous immune evasion tactics, persisting within their hosts through self-perpetuating strategies. One such tactic involves acquiring functional copies of host genes encoding cytokines such as IL-6 (HHV-8), IL-8 (GaHV-2), IL-10 (HHV-4, HHV-5), and IL-17 (SaHV-2). These viral mimics, or virokines, can bind to cellular receptors, modulating the natural cytokine signaling and manipulating the immune response to favor the virus. In the course of full-length cDNA sequencing of infectious laryngotracheitis virus transcripts, a previously unknown highly-spliced gene was discovered in the viral genome predicted to encode a 147 amino acid protein with similarity to vertebrate interleukin-4. The three-intron gene structure was precisely conserved with chicken and other vertebrate IL-4 homologs, and the amino acid sequence displayed structural conservation with vertebrate homologs at the primary, secondary, and tertiary levels based on computational modeling. The viral IL-4 gene was subsequently identified in all sequenced ILTV genomes. Phylogenetic analyses, along with the conserved gene structure, suggested direct capture from a *Galliformes* host. Functionally, an LPS-stimulation assay showed that the expressed viral IL-4 homolog stimulated nitric oxide production in a macrophage cell line at comparable levels to recombinant chicken IL-4. A recombinant virus lacking vIL-4 exhibited slightly higher titers in cell culture compared to the parental strain. *In vivo* bird studies demonstrated reduced pathogenicity of the vIL-4 knockout compared to wildtype. These results represent the first report of a previously unknown virokine encoded in the ILTV genome expressing a functional IL-4 homolog and virulence factor.

Keywords: infectious laryngotracheitis virus, virokine, interleukin-4, alphaherpesvirus, deletion mutant, virulence, host-pathogen interaction

∗Corresponding author
Email address: `stephen.spatz@usda.gov` (Stephen J. Spatz)

Introduction

Infectious laryngotracheitis virus (ILTV) causes acute upper respiratory disease that impacts chickens reared on high-density poultry farms, with severity of the disease dependent on the viral strain, infective dose, and age of the bird [1,2]. ILTV replicates to high titers in the laryngeal and tracheal mucosae, leading to severe cytolytic damage to mucosal epithelia and hemorrhaging. Mild disease causes coughing, nasal discharge, and conjunctivitis; severe disease is further characterized by marked dyspnea, gasping, open-mouth breathing, and expectoration of blood mucoid phlegm [1]. Mucoid plugs, likely the result of mucus hypersecretion by goblet cells in the trachea, often obstruct airways, predisposing birds to death due to asphyxiation [3]. If birds survive acute infection, a lifelong latent infection of the trigeminal ganglia will remain.

As with other α-herpesviruses, it is well-documented that cell-mediated immunity is the effective arm of the immune response to ILTV infection. It has been shown that chicks bursectomized within 24 hours post-hatch and vaccinated four weeks later were still resistant to ILTV challenge [2]. However, thymectomized chickens were more susceptible to disease and showed only slightly decreased protection in challenge experiments [4,5]. These results are consistent with previous findings from passive transfer experiments which showed that transferring antibodies from hyperimmune chickens did not provide protection against infection [4,6]. Sabir et al. recently found a weak association between protective immunity against ILTV and antibody titers to individual ILTV antigens such as glycoproteins C, D, E, G, I, and J [7]. These and other studies indicate a pivotal role for cell-mediated immunity in combating ILTV infection [8].

The process of host gene acquisition by viruses is well-known and widely studied — it is likely that most viral genes beyond the core replication machinery are a product of host gene capture at some point in viral evolution [9]. In larger DNA viruses with room for a relatively broad gene repertoire, the activities of these gene products extend beyond core replication and structural functions to specialized roles in immune invasion, pathogenicity, and other functions conveying an evolutionary advantage. Among these specialized host-acquired genes are those which manipulate the host immune response, including functional homologs of cytokines and their receptors which have agonistic or antagonistic effects relative to their host counterparts [10]. In herpesviruses alone, at least four viruses have been reported to harbor host-acquired cytokine homologs in their genomes: IL-6 in HHV-6; IL-17 in SaHV-2; and IL-10 homologs in both HHV-4 and HHV-5 [11–14].

The dsDNA type D genome of ILTV is approximately 151 Kbp in length and encodes approximately 80 putative protein-coding genes. With the exception of gene clusters near the termini of the unique long subgenomic region encoding eight ORFs (A, B, C, D, E, F, UL-0 and UL-1) that are unique to members of the *Iltovirus* genus, the majority of the remaining annotated CDS are homologous to genes of the α-herpesvirus prototype HSV-1. An earlier study classified the genes of ILTV into four categories based on their transcription kinetics: immediate early (α), early (β), leaky late (γ1) and true late (γ2) [15]. Immediate early genes are produced in the absence of *de novo* viral protein synthesis. Early viral genes typically code for

enzymes necessary for DNA replication, while late genes encode for the structural polypeptides of the virion. Transcription in ILTV is largely "leaky" (not strictly temporally controlled), with only ICP4 expression qualifying as truly immediate early. This is similar to equine herpesvirus 1 (EHV-1) which also encodes a single α gene (ORF62, an ICP4 homolog) [16]. In contrast, HSV-1 expresses four immediate-early genes (ICP4, ICP0, ICP22, and ICP27) [17].

Studies investigating the transcriptome of ILTV-infected cells have employed both *in vitro* and *in vivo* models and have relied on short-read sequencing (SRS) for assembling the viral and cellular transcripts [18–21]. Although SRS has a high yield and base accuracy, its short read length is ill-suited to identifying alternate splice forms and overlapping/co-terminal transcripts such as are common in herpesviruses [22]. Conversely, long-read sequencing (LRS) technologies such as Oxford Nanopore (ONT) and Pacific Biosciences (PacBio) are capable of resolving longer, in many cases full-length, transcript isoforms with varying rates of accuracy. LRS can be used to redefine the transcriptomes of virus-infected cells given sufficient viral coverage, sometimes in combination with deep SRS sequencing [23–25]. In the course of an effort to define the viral transcriptome of primary chicken kidney cells infected with either wild type (1874C5) ILTV, a US clade VI strain, or the vaccine CEO strain Laryngo-Vac (Zoetis) using PacBio IsoSeq full-length cDNA sequencing, we discovered a novel spliced transcript family mapping to a previously unannotated genomic region at the UL/IRS junction and predicted to encode a homolog to avian interleukin 4. We hypothesize that ILTV acquired an ancient avian interleukin 4 gene through a non-retrotranscription (direct) capture mechanism during host coevolution which potentially aided immune evasion. We demonstrate that vIL-4 exhibits structural and functional properties similar to those of chicken IL-4. To investigate the role of vIL-4 during ILTV infection, a deletion mutant virus lacking vIL-4 (ΔvIL-4) was generated and characterized in terms of its growth kinetics in cell culture and pathogenicity in birds.

Results

ILTV encodes a novel spliced transcript at the U_S/IR$_S$ junction

In the course of full-length cDNA sequencing of the ILTV transcriptome, a novel transcript family mapping to a previously unannotated region of the ILTV genome was revealed. The primary isoform consists of four exons transcribed from the end of the IRS, on the opposite strand to the surrounding UL(-1) and ICP4 genes (Figure 1). This isoform was detected in the IsoSeq transcripts from both virulent (1874C5) and attenuated (CEO Laryngo-Vac) strains. Less abundant junction-spanning splice forms were also observed in 1874C5, although at <1% of the abundance of the primary transcript. Post-discovery, we were also able to detect the transcript in existing short-read data from the NCBI Sequence Read Archive (an example is shown in Figure 1).

The novel transcript (named *vIL-4* as discussed below) was found to be highly abundant both *in vitro* and *in vivo*. In the *in vitro* semi-quantitative IsoSeq data, the *vIL-4* transcript was present at 3247 and 2606 viral transcripts per million (vTPM) in 1874C5 and CEO/LaryngoVac, respectively. By rank, these equated to the 18th and 19th most abundant transcript in each strain. Short-read RNA-Seq revealed the *vIL-4* transcript to be even more abundant *in vivo*,

Figure 1: Alignment of experimental IsoSeq transcripts to ILTV UL/IRS junction. IsoSeq models are colored by strand (blue: +, red: -) and shaded by supporting read count (darker = higher count). The primary 4-exon vIL-4 transcript (center, forward strand) is roughly 100× more abundant than the alternative isoform models in 1874C5. In CEO (LaryngoVac), where there was lower viral abundance overall, only the dominant isoform was detected. Also shown (bottom track) is an alignment of previously published short-read RNA-Seq data from strain LJS09 demonstrating expression of the vIL-4 transcript in an additional strain. Only a single replicate from this experiment is shown for the sake of brevity, but all four infected replicates showed similar coverage.

representing the fifth-most abundant viral transcript in birds inoculated with either 1874C5 or CEO/LaryngoVac (Figure 2).

The novel transcript encodes an IL-4 homolog

The longest open reading frame of the novel transcript codes for a 147 amino acid protein. A BLASTp search of the *nr* database suggested closest similarity to avian interleukin-4 homologs; the gene has thus been named *vIL-4* in keeping with nomenclature from other herpesvirus virokines. In order to carefully examine this relationship, IL-4 homologs from 29 sequenced avian and related crocodilian and turtle genomes were carefully curated. Species were chosen to broadly represent the avian tree of life, with the requirement that the full IL-4 gene structure was locatable in the genome. Some gene structures already existed in the genome annotations; in other cases, gene structures were determined by a combination of protein-to-nucleotide alignment and *de novo* gene prediction as described in the methods.

Secondary and tertiary structure prediction showed vIL-4 to be a classic four-helical cytokine with high confidence (Figures 3 and 4). It contains the expected N-terminal signal peptide and four universally conserved cysteine residues forming stabilizing disulfide bridges across the four-helical bundle. Predicted tertiary structures of vIL-4 and chicken IL-4 bound to the Type I receptor complex aligned closely (Figure 4C). In fact, spatial analysis of the predicted

Figure 2: Highly expressed in vivo viral transcripts. Shown are the top 8 ILTV transcripts by median TPM as a distribution of (A) viral TPM and (B) ordinal rank of TPM. RNA-Seq expression data is from ocular innoculation of birds with 1874C5 followed by sampling at 3, 5, and 7 d.p.i. In this experiment, vIL-4 is the fifth most abundant transcript expressed, following two origin-associated transcripts, glycoprotein G, and VP22.

models suggests a potentially stronger binding interaction between vIL-4 and its putative receptor IL-4Rα and γC (CD132) subunits than that of cIL-4 (21 and 17 predicted hydrogen bonds, respectively).

The genomic structures of the 30 curated IL-4 genes, including *vIL-4*, all contained three introns with identical locations relative to the amino acid alignment, and all intron boundaries occurred without exception at phase 0 in the coding sequence. The introns were located between core secondary structures (the first α-helix, first β-strand, α-helices two and three, and α-helix four) (Figure 3). All three of these characteristics (three coding sequence introns, aligned to codon phase 0, and occurring between adjacent secondary structures) have previously been shown to be conserved within the four-helical cytokine family as a whole [26]. Intron lengths varied widely, generally increasing with increasing genetic distance from vIL-4. Conservation of the vertebrate genomic structure in vIL-4 is in contrast to other herpesvirus-encoded cytokines, which are either unspliced or contain a single intron.

Viral IL-4 evolved within Galliformes

Viral IL-4 had a global amino acid identity to chicken IL-4 of 34.7% (max: turkey, 39.5%) and a global similarity to chicken IL-4 of 52.4% (max: Japanese_quail, 56.8%). Multiple alignment and phylogenetic analysis of the curated set of 30 sequences placed vIL-4 evolutionarily within this representative group (Figure 5). Trees produced from amino acid and nucleotide coding sequence alignments were highly congruent.

Viral IL-4 did not partition most closely with chicken, but rather emerged from a more distant common clade representing the Galliformes subsequent to divergence of brush-turkey (Fig. 5). The phylogenetic distance within this group was small overall. Based on the analysis of Chen et al. [27], this clade diverged roughly 75 Mya in the Late Cretaceous Epoch. There are several possible interpretations of this result. ILTV may have captured the host IL-4 from a distant ancestor during this period of divergence. This would place the origins of ILTV vIL-4 much further in the past than current estimates of divergence between sequenced present-day strains, which have been estimated at only 1–1.5 Mya [28], but on a similar time scale to the acquisition of a host PD-1 receptor ligand by TTGHV1 roughly 50 Mya [29]. Another possibility is that vIL-4 was captured more recently from an as-of-yet unsequenced avian host. While ILTV is most commonly associated with chickens, it has also been reported in pheasant, partridges and peafowl [1,30]. It is also possible that the different rates of evolution between viral and vertebrate homologs confounds the ability to correctly interpret the phylogenetic placement. The IL-4 trees suggest that vIL-4 is evolving at a rate at least twice as fast as the next most divergent sequence in the clade (guan), and roughly eight times as rapidly as chicken. The difficulty of interpretation in this scenario was discussed by Gibbs in the context of vertebrate and poxvirus thymidine kinases [31].

Viral IL-4 is highly conserved in ILTV

An intact vIL-4 gene was found in all 89 complete ILTV genomes published at the time of writing, after removing known recombinants, duplicates, and the original published mosaic sequence. After further filtering two sequences containing unknown (X) amino acids, 72 vIL-4

Figure 3: Primary, secondary, and genomic structure of the viral interleukin-4 homolog. (A) Amino acid alignment of vIL-4 with homologs from selected species. (B) Secondary structure probability of the vIL-4 amino acid sequence as predicted by JPRED2. (C) Signal peptide domain probability as predicted by SignalP 6.0; indicated are the N-terminal domain ['N'], hydrophobic domain ['H'], and C-terminal domain ['C']. (D) Location of genomic introns in a representative selection of homologs, translated into amino acid space. Black vertical bars indicate the intron position, with the intron length indicated to the right of each bar. Intron locations within the amino acid alignment were identical for all homologs examined, including those not shown.

Figure 4: Comparative tertiary structures. (A) Empirical crystal structure of human IL-4 (PDB:2B8U); cysteines/disulfide bonds shaded yellow. (B) Alphafold model of vIL-4 shaded by pLDDT from low (red) to high (blue); cysteines/disulfide bonds shaded yellow. ILTV vIL-4 is predicted to share the same four-alpha-helix, anti-parallel beta sheet arrangement as the empirically modeled hIL-4. Like all avian IL-4 homologs, it lacks the cysteines at both protein termini that form a third disulfide bond in human IL-4 (panel A lower-left). (C) Superimposed predicted structures of vIL-4 (blue) and chicken IL-4 (green) in complex with chicken IL-4Rα (transparent purple) and γc (transparent red) receptor subunits (Type I receptor complex), as predicted by Alphafold and aligned by the RCSB Pairwise Structure Alignment tool (jFATCAT/rigid). Disulfide bridges are shown in yellow.

Figure 5: ML trees of vIL-4 and interleukin-4 homologs from a selection of representative bird and reptilian genomes. (A) From amino acid alignment, built with IQTREE using the 'Q.bird+I+G4' model as selected by ModelFinder, ultrafast bootstrapping (1000 iterations). (B) From coding nucleotide alignment, built with IQTREE using the 'GTR+I+R' model as selected by ModelFinder, ultrafast bootstrapping (1000 iterations). Branches with UFBoot support (in red) < 90 were collapsed.

amino acid sequences were identical to 1874C5 and Laryngo-Vac. The other 15 sequences were in four groups that could be distinguished by only six amino acid substitutions (Figure 6).

Functional equivalence of vIL-4

We investigated the ability of both cIL-4 and vIL-4 to stimulate the production of nitric oxide (NO) in avian macrophage cells. Macrophages play a critical role in the immune system and are known to produce NO in response to certain stimuli. For this experiment, an avian macrophage cell line HD11 was utilized. Cells were stimulated with varying concentrations (ranging from 62.5 to 1000 ng/ml) of the tagged vIL-4 and cIL-4 proteins for a duration of 4 hours. This stimulation period allowed for interaction between the tagged proteins and their cognate receptors on the macrophage cell surface, potentially triggering downstream signaling pathways. Subsequently, *E. coli* lipopolysaccharide (LPS), a potent bacterial endotoxin known to stimulate NO production in macrophages, was added to the cultures to activate the macrophages. Following a 20-hour incubation period, the concentration of nitrite (NO_2^-) in the cell culture supernatant was measured using the Griess assay. Both chicken interleukin-4 (cIL-4) and viral interleukin-4 (vIL-4) exhibited similar stimulatory effects on nitric oxide (NO) synthesis in

Figure 6: ML tree of amino acid sequences of vIL-4 in sequenced ILTV genomes. An intact gene structure and in-frame coding sequence was found in all 89 complete ILTV genomes examined (after removing known recombinants, duplicates, and the original published mosaic genome). Most of the amino acid sequences were identical, and are shown as a single 'Serva-like' cluster. Only a handful of variants characterized the remaining sequences. Each branchpoint on the tree represents one or several amino acid variants, labeled in blue. The tree was built with IQTREE using the 'Q.bird' model as selected by ModelFinder, ultrafast bootstrapping (10000 iterations). Support values shown in gray are UFBoot estimations.

Figure 7: Stimulation by IL-4 of nitric oxide production in macrophages. The Griess assay was used to measure NO_2^- as an indicator of NO production at varying concentrations of either chicken IL-4 (cIL-4) or viral IL-4 (vIL-4) and in the presence of 2.5 μg/ml LPS. Inset: standard curve of Griess reagent used to calculate NO_2^- concentrations ($R^2 = 0.92$).

macrophages when co-stimulated with the microbial agonist lipopolysaccharide (LPS) (Figure 7). No statistically significant difference was observed in the amount of nitrite (NO2-), a stable breakdown product of NO, produced by macrophages treated at each concentration of either cIL-4 or vIL-4 in the presence of 2.5 μg/mL LPS. This suggests a functional similarity between these two interleukin-4 variants regarding their ability to induce NO production in macrophages upon LPS stimulation [32–35].

In vitro *replication kinetics of parent 1874C5 and ΔvIL4 strains*

The replication kinetics of parent 1874C5 and ΔvIL-4 ILTV strains were determined by measurement of $TCID_{50}$/ml (Figure 8A) and viral genome load (Figure 8B) from combined infected supernatant and cells collected at 24, 48, 72 and 96 hours post virus absorption. No measurable cytopathic effect was detected at 24 hours post-absorption for either 1874C5 or ΔvIL-4 strains, while by 48 hours post-absorption, the parent 1874C5 strain had an average titer of $\log_{10}$ 0.24 $TCID_{50}$/ml as compared to an average titer of $\log_{10}$ 2.99 $TCID_{50}$/ml for the ΔvIL-4 strain. At 72 and 96 hours the ΔvIL-4 strain titers were $\log_{10}$ 1.1 to 1.7 $TCID_{50}$/ml higher than the parent 1874C5 strain that reached an average titer of $\log_{10}$ 4.40 $TCID_{50}$/ml by 96 hours post-absorption. Despite the difference in viral $TCID_{50}$/ml, viral genome load for both viruses showed an almost equivalent rate of increase by 72 and 96 hours post-absorption (Figure 8B).

Figure 8: Growth kinetics of 1874C5 parental and ΔvIL-4 ILTV strains in the LMH cell line. LMH cells (1.4×10^6 cells per well) were inoculated in triplicate with 1874C5 parental strain and ΔvIL-4 strain at MOI of 0.002. After absorption for 1 hour at 39°C and 5% CO_2, cells were washed and incubated for 0, 24, 48, 72 and 96 hours. At each time point supernatant and cells were collected (n=3) and (A) viral titer and (B) viral genome copy number were determined for each strain at each time point. Individual replicates and corresponding nonlinear fit curves are shown.

Pathogenicity of parent 1874C5 and ΔvIL4 strains after intratracheal inoculation

The percent survivability of chickens inoculated with the ΔvIL-4 strain (n=17) was higher (47.1%) than the survivability of chickens (n=15) inoculated with the parent 1874C5 strain (33.3%). Mortalities induced by the parent 1874C5 strain peaked at day four, while mortalities for chickens inoculated with the ΔvIL-4 strain peaked at five d.p.i. (Figure 9A). Average clinical sign scores (CSS) induced by the parent 1874C5 and ΔvIL-4 strains at days 2 to 7 p.i. were not statistically different (Figure 9B). However, at days 4 and 5 p.i. clinical sign scores induced by the parent 1874C5 strain (4.60 and 5.00) were numerically higher than clinical signs scores induced by the ΔvIL-4 strain (3.55 & 3.89). The peak of clinical signs for both the parent 1874C5 and ΔvIL-4 groups of chickens were identified at day five p.i. (Figure 9B) which coincided with the ΔvIL-4 strain peak of mortality. However, the peak of mortality for the parent 1874C5 inoculated group of chickens was recorded earlier (day 4 p.i.) than for the ΔvIL-4 strain peak of mortality (day 5 p.i.) (Figure 9A). The average trachea viral genome load for the parent 1874C5 and ΔvIL-4 strains was measured at 3 and 5 d.p.i — no statistical differences were detected. (Figures 9C & D). At three days pi the parent 1874C5 and the ΔvIL-4 viral genome load in trachea reached an average of 5.339 and 5.773, respectively (Figure 9C). At 5 days p.i., although not statistically different (p>0.05), the average viral genome load for parent 1874C5 strain was higher (4.6) than that of the ΔvIL-4 strain (3.3) (Figure 9D). When comparing decrease of viral genome load of each strain in the trachea between days 3 and 5 p.i., the parent 1874C5 genome load showed no significant decrease (p>0.05) (Figure 9E) while the genome load of ΔvIL-4

strain showed a significant decrease (p=0.001) from 5.8 at day 3 to 3.3 at day 5 p.i. (Figure 9F).

Based on the Guy et al 1990 trachea lesion score system [36], all upper tracheas from the parental-inoculated group of chickens received the highest score of 5 while all trachea sections from the ΔvIL-4-inoculated group of chickens received a score of 4. Most of the trachea sections from parental-inoculated chickens displayed severe damage to the mucosa characterized by detachment of complete sections of the epithelium and abundant fibrino-haemorrhagic cellular infiltrates in trachea lumen, with either a slim submucosa layer remaining (Figure 10A) or sections of enlarged submucosa with no mucosal epithelium remaining (Figure 10B); in either instance, syncytia and inclusion bodies were rarely found (Figures 10A & B). Although the ΔvIL-4-inoculated upper trachea sections displayed damage and detachment of the mucosal epithelium, frequently trachea segments maintained an enlarged submucosa with some distal, thin remaining epithelium in which scattered syncytia with intranuclear inclusion bodies could be detected (Figure 10C; see inserted photomicrograph). Another histopathological presentation characteristic of ΔvIL-4 inoculated tracheas was the presence of slightly haemorrhagic cellular pseudomembranes associated to the enlarged submucosa (Figure 10D).

Discussion

Vertebrate immune systems have evolved intricate mechanisms to skillfully recognize and control viral infections. However, viruses have developed countermeasures to subvert host immunity and ensure their continued survival. A key driver in this host-pathogen co-evolution is horizontal gene transfer, where large DNA viruses acquire host genes. Over time, these captured genes are modified through mutation and selection to perform three main functions: (i) mimicry of host molecules to evade detection, (ii) antagonism of immune pathways for direct suppression, or (iii) development of novel immunoregulatory function to manipulate the host response. Notably, the acquisition and subsequent adaptation of antiviral defense components confer selective advantages on viruses. This allows viruses to directly target and disrupt both the innate and adaptive arms of the immune response, ultimately enhancing their survival and dissemination. Interleukins play a fundamental role in regulating the magnitude of these responses. Accordingly, several herpesviruses have been reported to encode functional homologous "mimics" of host cytokines (IL-6, IL-10, and IL-17). These virokines mimic their cellular counterparts and are more common in γ- than β-herpesviruses [37]. There are no reports of non-chemotactic cytokines encoded by α-herpesviruses until our recent discovery of an interleukin 4 mimic in the genome of ILTV. One chemokine, vIL8, is encoded by the related avian α-herpesvirus Gallid alphaherpesvirus 2 (GaHV-2), the causative agent of Marek's disease [38]. Generally, virokines can interact with immune cell receptors, transmit a signal through them via the JAK/STAT pathway and play a role in the pathogenesis of herpesvirus-associated diseases by modulating immune responses, promoting cell survival, and stimulating cell proliferation.

One of the most well-studied virokines is viral interleukin-6 (vIL-6), encoded by Kaposi's sarcoma-associated herpesvirus (KSHV). Despite sharing only 37% amino acid identity with human IL-6, vIL-6 exhibits potent pro-inflammatory activity. It acts in an autocrine manner, promoting both cell growth and survival by inducing expression of host IL-6 [39] and suppressing

Figure 9: Pathogenicity of 1874C5 (parent) and vIL-4 gene-deleted strain (ΔvIL-4) in specific pathogen free (SPF) chickens after intratracheal inoculation at 24 days of age. (A) Percent survival for chickens inoculated with the ΔvIL-4 strain (n=17) 47.1% and for chickens inoculated with the parent strain (n=15) 33.3%. (B) Average clinical sign scores per group of chickens from 2 to 7 days post-intratracheal inoculation. Although not significant, average clinical sign scores induced by the 1874C5 strain were higher than average scores induced by the knockout. (C) Viral genome load in trachea three days post-intratracheal inoculation. (D) Viral genome load in trachea five days post-intratracheal inoculation. (E) Comparison of parent 1874C5 genome load at 3- and 5-days post-inoculation. (F) Comparison of ΔvIL-4 genome load at 3- and 5-days post-inoculation. Individual replicate loads indicated by points; columns heights indicate the average trachea viral genome load per group. No significant (ns) differences (p>0.05) were detected between viral genome load of the 1874C5 parent and the ΔvIL-4 mutant at either 3- or 5-days post-inoculation. Significant differences (p<0.05) were detected for the average viral genome load of ΔvIL-4 in the trachea at 3- vs 5-days post-inoculation.

Figure 10: Histopathological lesions in the trachea induced by parental 1874C5 and ΔvIL-4 at four days post-intratracheal inoculation. Photomicrographs are H&E-stained upper trachea sections from chickens infected with parental 1874C5 (A & B) or ΔvIL-4 (C & D) at 20x magnification with image scale bar of 100 μm. (C) inset: Inserted photomicrograph at 40x magnification with image scale bar of 50 μm underscores the presence of mucosal epithelium and goblet cells with few syncytia and intranuclear inclusion bodies still attached distal of the submucosa.

production of IFNα, a key molecule for activating macrophages and enhancing their anti-viral activity.

A striking convergence in viral evolution is evident in the independent acquisition of interleukin 10 homologs by more than a dozen viruses belonging to three distinct viral families: eight in Herpesviridae, one in Alloherpesviridae and four in Poxviridae [40]. The widespread distribution of interleukin-10 (IL-10) mimics among diverse viral families highlights a profound selective pressure during viral-host co-evolution to acquire genes or cDNA copies encoding proteins capable of immunosuppression. The best-studied IL-10 homologs are those of Epstein-Barr virus (EBV) and cytomegalovirus (CMV) [41]. The gene encoding vIL-10 of EBV has over 80% amino acid identity with the human counterpart. Like vIL-6 of KSHV, EBV's vIL-10 inhibits IFN-γ synthesis in T cells [42] and potentially inhibits natural killer (NK) cells and cytotoxic T cells, as evidenced by reduced activities of these cells in mice infected with a recombinant vaccinia virus expressing vIL-10 of EBV [43]. This suppression hinders the immune system's ability to clear the virus and potentially promotes viral persistence within the host.

Cytomegaloviruses of primates have also adapted to manipulate the host immune response through an IL-10 homolog [44–46]. The protein cmvIL-10 in HCMV arose from a captured human IL-10 gene some >42 million years ago, evolving a molecule with only 27% amino acid sequence identity to its human counterpart. Despite this low homology, cmvIL-10 retained the ability to bind the human IL-10 receptor with high affinity [47] and exerts a multifaceted immunosuppressive effect depending on the target cell and the stage of infection.

Saimiriine herpesvirus 2 (HVS) further exemplifies viral mimicry with its unique IL-17 mimic, vIL-17 [12,48]. This virokine exhibits a high degree of structural similarity (74% amino acid identity) to its host counterpart and can bind the same cellular receptor [12]. While the precise role of vIL-17 in HVS infection remains to be fully elucidated, the established etiological role of human IL-17 in human idiopathic pulmonary fibrosis raises the intriguing possibility that vIL7 might serve a similar proinflammatory function, potentially creating an immunosuppressive environment beneficial for HVS persistence.

These observations collectively reinforce the broader concept that virally encoded cytokine mimics represent a strategic advantage for herpesviruses, depending on the mimicked function, to establish persistent infections within their hosts. Anti-inflammatory mimics like vIL-10 directly suppress the host immune response, promoting viral persistence. Even pro-inflammatory mimics can benefit the virus. Viral IL-6, while triggering inflammation, suppresses antiviral interferon production and promotes cell survival, potentially aiding the virus. The role of pro-inflammatory mimics like vIL-17 is less clear. While the human counterpart, IL-17, can contribute to inflammation, its specific function in viral infections remains under investigation. Saimiriine herpesvirus 2's vIL-17 might create a pro-inflammatory environment, but its ultimate effect on the host immune response and viral persistence needs further research. Overall, the benefit hinges on how the mimic manipulates the immune response.

Here, we report the discovery of a functional interleukin-4 (IL-4) homolog in an α-herpesvirus, adding to the list of known virokines and uncovering a potential new immunosuppressive tactic employed specifically by an avian herpesvirus. In mammals, interleukin-4 (IL-4) acts as a central regulator in type-2 immunity. It influences a variety of cells, both haematopoietic and

non-haematopoietic. Its effects include activating specific Th2 cells, differentiating B cells into plasma cells, promoting immunoglobulin E (IgE) class switching, generating alternatively activated macrophages, stimulating smooth muscle contraction and mucus production, and innate cell recruitment to the site of inflammation. Avian species, in contrast to mammals, lack the immunoglobulin E (IgE) class and express only IgY, IgM, and IgA antibodies. Unfortunately, research on the functional role of chicken IL-4 is less comprehensive compared to its mammalian counterpart.

Our study demonstrates that the four-exon genomic structure of vIL-4 is remarkably conserved with chicken and other vertebrate IL-4 homologs and the amino acid sequence displayed structural conservation with vertebrate homologs based on computer modeling. Phylogenetic analysis suggests its direct capture from a Galliformes host approximately 75 million years ago. *In vivo* RNA-Seq showed vIL-4 to be among the top five most abundantly transcribed viral genes during the sampling period from 3 to 7 days post-infection, which encompassed the period of virulent lytic replication. Functionally, vIL-4 stimulates nitric oxide production in macrophages to levels comparable to chicken IL-4. Notably, compared to the parent 1874C5 strain, an ILTV mutant lacking vIL-4 (ΔvIL-4) exhibited slightly higher viral titers but similar genome load in LMH cells. This result may suggest that *in vitro*, the absence of vIL-4 may favor a faster production of infectious viral particles, reflected by the increased early titers for the ΔvIL-4 strain. In contrast, the rate of viral genome replication *in vitro* was similar for both strains, which coincides with the *in vivo* results in which parent 1874C5 and ΔvIL-4 viral genome load in the trachea at 3- and 5-days p.i. showed no difference. However, when comparing each viral strain trachea genome load at 3 and 5 days, the genome load of ΔvIL-4 was significantly lower at day 5 pi. Also, the ΔvIL-4 strain induced lower mortalities and lower clinical signs of the disease. These subtle differences in pathogenicity were corroborated by the histopathological findings in the upper trachea of chickens inoculated with the ΔvIL-4 strain. Although the trachea lesions induced by the ΔvIL-4 strain were characterized by increased infiltration and epithelium damage, in most of the trachea rings examined the submucosa and lamina propria remained in place with some distal epithelium still attached. In the case of 1874C5-inoculated chickens, most of the trachea mucosa and submucosa were detached into the lumen. These findings collectively suggest that vIL-4 acts as a novel virokine, potentially modulating the host immune response and contributing to ILTV virulence.

The discovery of vIL-4 with sequence similarity to proinflammatory cIL-4 also raises questions about the connection between this viral mimic and other cytokines, particularly cIL-13. In mammals, both IL-4 and IL-13 can activate common pathways through usage of lineage-determining factors STAT6 and GATA3 via binding of shared receptor subunits. This raises the possibility that vIL-4 might emulate some functions of chicken IL-13 in addition to IL-4. Notably, mammalian IL-13 shares similar effects on lymphocytes with IL-4, both being classified as proinflammatory under normal circumstances. However, in chronic inflammatory diseases like asthma, IL-13 can exhibit seemingly paradoxical properties. Through an anti-inflammatory response, it can also induce the production of matrix metalloproteinases and goblet cell metaplasia, leading to mucus hypersecretion. This seemingly contradictory response might be a protective mechanism against excessive inflammation that could otherwise lead to asphyxiation, the

ultimate cause of mortality in ILTV infection.

In conclusion, this study presents the discovery of a previously unknown vIL-4 encoded by ILTV. It is likely that ILTV has developed a multifaceted approach to manipulate the host's immune response through the action of its encoded cytokines glycoprotein G (chemokine) and vIL-4 (virokine). This virokine may play a role in ILTV virulence, potentially through interactions with both IL-4 and IL-13 receptors. Paradoxically, vIL-4 might even promote specific aspects of the humoral immune response that don't effectively combat the virus through a shift in the Th1/Th2 paradigm. Further investigation into the complex interplay between vIL-4, cIL-4, cIL-13 and their receptors should pave the way to a better understanding of immune evasion strategies employed by ILTV, undoubtedly aiding efforts to develop more effective strategies to combat infectious laryngotracheitis.

Data availability

Sequences for the 1874C5 and Laryngo-Vac vIL-4 mRNA transcripts have been deposited in GenBank under accession numbers PP974316 and PP974317. All curated nucleotide and amino acid sequences referenced, along with GFF3 annotations for homologs in public vertebrate and ILTV genomes, are available at the manuscript GitLab repository (`https://gitlab.com/BASE2BIO/projects/vIL4`). The public git repository also includes all code used to perform statistical analyses, generate figures, and assemble the manuscript in RMarkdown.

Materials and Methods

Viruses and Cells

The ILTV virulent strain 1874C5 was originally isolated from the upper respiratory tract of broilers during outbreaks of the disease and subsequently classified as a member of the genotype VI [49]. The CEO vaccine strain Laryngo-Vac was a generous gift from Zoetis. The vIL-4 null mutant was derived from the genotype VI virulent 1874C5 strain. Leghorn male hepatoma (LMH) cells were propagated in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (ThermoFisher), 100 U/ml penicillin, 100 μg/ml streptomycin and 2mM L-glutamine. For virus infections, maintenance DMEM supplemented with 2% FBS was used for the generation of vIL-4 null mutant and replication kinetics studies. ILTV strains CEO, 1874C5 and ΔvIL-4 virus were also propagated in primary chicken kidney (CK) cells from 3- to 4-week-old chickens for the purpose of generating virus stocks and isolation of nucleic acids as previously described [50,51]. Titers were expressed as tissue culture infective dose (TCID$_{50}$) and calculated by the Reed and Muench method [52].

Avian macrophage cells (HD11) were cultured in DMEM supplemented with 10% fetal bovine serum, 100 U/ml penicillin, 100 μg/ml streptomycin, and 2.0 mM L-glutamine. For LPS/IL-4 stimulation assays, aliquots of 100 microliters of cell suspension, containing $2{\times}10^6$ cells/ml, were seeded into wells of a 96-well tissue culture plate. The cells were propagated until they reached approximately 85% confluence before further experimentation.

RNA isolation

Chicken kidney (CK) cells were infected with 1874C5 or LaryngoVac at a MOI of 0.001. RNA was isolated from infected CK cells using Trizol according to the manufacturer's recommendations. RNA was quantified using the Qubit RNA BR (Broad-Range) Assay Kit on a Qubit Fluorometer (ThermoFisher). The quality of RNA was assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies) with the RNA 6000 Nano LabChip kit according to the manufacturer's instructions. Briefly, the RNA samples were diluted in water and loaded onto the microfluidic chip along with RNA ladder standards. The resulting electropherograms were analyzed with Agilent 2100 Expert software to determine the RNA Integrity Number (RIN).

PacBio sequencing

RNA samples with RIN ≥ 7 were used to prepare the IsoSeq barcoded libraries following the PacBio protocol (Procedure & Checklist – Iso-Seq™ Express Template Preparation for Sequel® and Sequel II Systems, PN 101-763-800 Version 02) using the SMRTbell Express Template Prep kit 2.0 (PacBio). The individual libraries were barcoded using the Iso-Seq Express Oligo kit (Pacbio PN 101-737-500). The barcoded libraries were multiplexed in equimolar amounts into a final SMRTbell template mix, which was purified using 1× beads. The final library was sequenced on the Sequel II system for 20 hours using the Sequel II Binding Kit 2.1. The final loading concentration was 80 pM.

IsoSeq preprocessing and transcript building

Raw PacBio raw subreads were processed to circular consensus sequencing (CCS) reads using ccs v6.4.0 (`--min-rq 0.9`) (this and all other PacBio software referenced was installed from the PacBio Bioconda packages (`https://github.com/PacificBiosciences/pbbioconda`) [53]. Demultiplexing and adapter removal was performed using lima v2.7.1 (`--isoseq-peekguess`). Poly-A tail removal and concatemer filtering were performed using isoseq refine from the isoseq3 package v3.8.2 (`--require-polya`). Full-length, non-concatemeric (FLNC) reads were then clustered into non-redundant transcripts using isoseq cluster (`--use-qvs`). Transcript cluster sequences were mapped against the combined bGalGal1.mat.broiler.GRCg7b+virus genome reference using pbmm2 align v1.10.0 (`--preset ISOSEQ-sort`). Overlapping, exon-sharing transcripts were further collapsed using isoseq3 collapse (`--do-not-collapse-extra-5exons- max-5p-diff 200`). Only transcripts with a minimum support of four FLNC reads were kept. Further filtering of potential artifacts was performed using SQANTI3 v5.1.1 [54] to remove transcripts with a genomic adenine content downstream of the mapped TTS of 60% or greater (suggesting artifactual priming on the genome rather than a transcript poly-A tail) and spliced transcripts with non-canonical donor/acceptor sites. IGV [55] was used to visualize transcripts on the viral genome to first detect the presence of the vIL-4 transcript cluster.

Curation of IL-4 homologs

Initial BLASTP and TBLASTN [56] searches of the chicken IL-4 sequence (NP_001385388) against the *nr* and *nt* databases, respectively, were used to generate a candidate list of published homologs. In some cases, high-confidence annotations were found for a species; in others,

only partial TBLASTN matches to genomic regions were seen. In the latter case, the chromosome/scaffold/contig sequence was analysed using a combination of miniprot [57] protein-to-nucleotide alignment with GFF3 output and AUGUSTUS [58] *de novo* gene prediction (*Gallus gallus* model) to generate a preliminary gene structure. After multiple alignment of these preliminary sequences, several homologs from existing database entries contained questionable insertions/deletions in conserved regions. These homologs were analyzed either visually in IGV or using miniprot alignment to see if valid adjustments could be make to the splice junctions. In all such cases, valid adjustments using canonical donor/acceptor sites were found that brought the protein sequences into better structural agreement with the group as a whole. The final gene models in GFF3 format are available in the manuscript repository as described in *Data Availability*. Notes on the curation of each homolog sequence can be found in Table S1. Homolog sequences that were either absent from existing genome annotations or were modified from the existing annotation were submitted to GenBank as third-party annotations (TPA:inferential) with accession numbers XXX–XXX.

In vivo *RNA-Seq data analysis*

Raw FASTQ files were trimmed using Trim Galore (`https://github.com/FelixKrueger/TrimGalore`)(`--2colour 8 --stringency 4 --length 40 --trim-n --max_n 5`). Kallisto v0.50.1 was used for transcript quantification [59]. A reference index was generated using `kallisto index` on the combined bGalGal1.mat.broiler.GRCg7b + virus transcript sequences. For this purpose, a local, as-of-yet unpublished updated annotation of the 1874C5 genome was used, based on an (also unpublished) corrected genome sequence. These files in GFF3 and FASTA format are available upon request. Quantification was performed with the following non-default settings: `--rf-stranded`. TPM values were merged for all replicates and re-normalized using only viral transcripts.

Re-analysis of LJS09 sequencing

RNA-Seq reads originally published by Wang et al. [21] of treatment of chick embryo liver cells with ILTV strain LJS09 or mock were downloaded from the NCBI Sequence Read Archive (accessions SRR5515169–SRR5515172 and SRR5515181–SRR5515184). These reads were mapped directly against the combined bGalGal1.mat.broiler.GRCg7b + virus genome reference using HISAT2 v2.2.1 [60] (defaults) and converted to bedgraph coverage format for visualization using bedtools v2.31.1 (`genomecov -split -ibam <IN> -bg`) [61].

Structural modeling

The JPred4 web server was used for predicting secondary structure of the vIL-4 amino acid sequence [62]. The SignalP 6.0 web server was used to predict signal peptide regions [63]. Alphafold 2.0 [64] (`--max_template_date=2020-05-14`) was used for tertiary structure prediction of both vIL-4 and chicken IL-4, individually and in conjunction with the sequences of the Type I receptor complex (IL-4Rα and γc). The input files and output models used in prediction are available in the manuscript repository.

Phylogenetics and tree reconstruction

MAFFT v7.525 [65] (`--auto`) was used to build all CDS and amino acid multiple alignments used in the paper. Tree inference was performed using IQ-TREE v2.3.4 [66]. Models for each multiple alignment were initially chosen using the ModelFinder feature of IQ-TREE [67] ('Q.bird+I+G4' for IL-4 amino acid alignment; 'GTR+I+R' for IL-4 coding sequence alignment; 'Q.bird' for ILTV vIL-4 alignment. Bootstrap support was computed using UFBoot2 with 10000 iterations [68].

Generation of GFP/IL-4 fusion proteins

To evaluate the functional similarity between viral interleukin 4 (vIL-4) and chicken interleukin 4 (cIL-4), both proteins were N-terminally tagged with a green fluorescent protein (muGFP) [69] (Figure S2A). The amino terminus was chosen based on *in silico* spatial modeling of the IL-4 receptor complex (see Figure 4) in order to minimize potential interference with receptor binding. A flexible linker peptide consisting of 12 amino acids (GSAGSAAGS-GEF) [70] was inserted between the muGFP and the respective IL-4 coding sequences. This linker serves a crucial role in maintaining proper protein folding and minimizing steric hindrance, ensuring the tagged protein retains functionality. The vIL-4 signal peptide sequence was incorporated at the N-terminus of the muGFP moiety in both constructs. This sequence directs the tagged proteins towards the secretory pathway, enabling their efficient export from the cell. To monitor transfection efficiency, an internal ribosome entry site (IRES) was placed upstream of a separate reference fluorescent protein (red fluorescent protein; RFP). This allowed for independent translation of the RFP, providing a reliable indicator of successful vector delivery into the host cell, even in situations where the tagged IL-4 proteins might exhibit lower expression levels. To accomplish this, the vector pNonSynVP2-DsRed2 [71] containing a CMV immediate early promoter upstream of an IRES proceeding a DsRFP CDS was used. The coding sequence (828 bp) for monomeric GFP (muGFP) [69], flanked at its 5' terminus by 75 bp encoding the vIL4 signal sequence and flanked at the 3' terminus with 36 bp encoding a flexible linker, was synthesized commercially. Two constructs were generated using Gibson cloning in a 2 insert/1 vector scheme. The amplicons for the cIL-4 constructs were generated using primer Set 1: VTR muGFP FOR C-IL4/muGFP C-IL4REV cIL4 (muGFP amplicon); primer Set 2: C-IL4 MuGFP For/cIL4 Rev (cIL-4 amplicon); and primer Set 3: vtr cil4 for/vtrMuGFP REV C-IL4 (vector amplicon). The amplicon for the vIL4 constructs were generated using primer Set 4: VirIL4 For/muGFP vIL4 REV (muGFP amplicon); primers Set 5: vIL4 MuGFP For/VirIL4 Rev (vIL-4 amplicon); and primer Set 6: vtrVir IL4 for/vtr Vir IL rev (vector amplicon). For the sequences of the primers see Figure S1. Following in-fusion cloning the resulting cIL-4 and vIL-4 tagged constructs (Figure S2A) were transformed into NEB alpha electrocompetent E. coli and selected on LB agar plates containing kanamycin (50 µg/mL). Positive recombinants were identified through colony PCR and restriction enzyme digestion of mini-prep DNA.

Transient expression and quantification of secreted GFP-tagged proteins

LMH cells were seeded into 6-well plates at a density of 8.0×10^5 and transfected with 2.5 µg of GFP-tagged cIL-4 or GFP-tagged vIL-4 using TransIT (Mirus) according to the manu-

facturer's instructions. Forty-eight hours post transfection the supernatants were harvested and centrifuged at 300×g for 10 minutes to remove cellular debris. Purified GFP standards (ranging from 10 ng/μL to 350 ng/μL) and diluted samples were pipetted (100 μL) into separate wells of a black, flat-bottomed 96-well plate. Fluorescence intensity was measured using a SpectraMax fluorometer at excitation and emission wavelengths of 488 nm and 507 nm, respectively. A standard curve was generated by plotting the average fluorescence intensity of each standard against its corresponding GFP concentration. The concentration of the secreted GFP-tagged protein in the original media was then determined by interpolating the sample fluorescence intensity on the standard curve.

LPS stimulation of HD11 cells

HD 11 cells (2.0×10^5) were grown in 96 wells overnight. When they reached ~85-90% confluency the culture media were replaced with fresh DMEM media to remove any immunostimulatory factors present in the conditioned media. To assess the direct effects of the GFP-tagged cytokines vIL-4 and cIL-4 on NO production, cells were preincubated with varying concentration (1000, 500, 250, 125 or 62.5 ng/ml) of either GFP-tagged vIL-4 or GFP-tagged cIL-4 for 4 hours. Following pre-incubation, LPS (Escherichia coli Serotype 0111:B4) 2.5 μg/ml was added to each well, and the cells were further stimulated for an additional 20 hours. Nitrite accumulation in the cell culture supernatant was measured as an indirect indicator of NO production using the Griess assay. Briefly, after the stimulation period, culture supernatants were transferred to a new plate and mixed with sulfanilamide and naphthylenediamine solutions. The plate was incubated for 30 minutes at room temperature, and nitrite concentration was determined by measuring the absorbance at 595 nm. A standard curve generated using sodium nitrite solutions was used to convert absorbance values into nitrite concentrations in the culture supernatants.

Generation of vIL-4 null mutants

The sequence of the four vIL-4 exons were screened *in silico* using Geneious Prime software to identify optimal target sequences for Cas9 nuclease based on the scoring algorithms developed by Doench et al. [72]. Two sgRNAs ATCCTTAGGGAAGAACCCAGGT and ACCGTGCCCTATAACAGTGGG with scores >0.7 were chosen and cloned in a human-codon optimized cas9 expression vector (VectorBuilder) under the control of U6 promoters. LMH cells were seeded in 6-well plates at a density of 1×10^5 per well and incubated overnight at 39°C in 5% CO_2 atmosphere. Cells at 90% confluency were transfected using TransIT (Mirus) with 7.5 μg of plasmid containing the sgRNAs and cas9 gene according to the manufacturer's protocol. The transfected cells were incubated at 39°C in 5% CO_2 for 12 hours. The cells were then infected with strain 1874C5 using either an MOI of 0.001 or 0.01. After absorption for one hour, the media was removed and DMEM containing 2% FBS and phosphonacetic acid (100 μg/ml) was added and the cells were incubated for an additional 12 hours. The media was again replaced with DMEM containing 2% FBS with 2% methylcellulose. Transfected/infected cells were incubated for 3 days, and 20 plaques were picked using a Evos microscope. Picked plaques, stored in a volume of 100-200 μl and frozen at -80°C, were subjected to three freeze/thaw cycles to release the virions.

Plaque purification and PCR confirmation

To determine whether single plaques were isolated to homogeneity, a plaque PCR assay was employed. A single, isolated ILTV plaque from infected LMH cells overlayed with DMEM containing 2% methylcellulose was picked and suspended in complete DMEM medium. Virions were released after three freeze/thaw cycles at -80°C and 37°C, respectively. Following 5 rounds of plaque purification using limited dilutions (1:2) and expanding onto cells seeded in 24 well plates, the infected cells were trypsinized. To isolate the viral DNA, the cells were pelleted, washed with 1x PBS and lysed in 100 µl of 5mM Tris-HCl pH 8.0 at 95°C for 10 minutes. Proteinase K was added to a final concentration 0.3 µg/µl (1.5 µl of 20 mg/ml solution). After incubation at 55°C for 30 minutes, the reaction was stopped via heating at 95°C for 10 minutes and the cell debris removed by centrifugation. Five microliters were used in amplification reactions using LongAmp® Taq DNA Polymerase (NEB) and primers: F1-4 AAGGGCCG-CAATTAGGTGGT and R1-4 AGCCCCGT TGACTTTAGCTT in a standard reaction with the following cycling conditions after an initial denaturation at 94°C for 2 min: 94°C for 30 sec, 58°C for 30 sec and 65°C for 1 min (40 cycles). The final extension was at 65°C for 1 min. Two clones (17 and A) that generated single amplicons of the expected size (654 bp) were selected for further confirmation. Sanger sequencing of the amplified region encompassing the targeted sequence verified the presence of the desired vIL-4 deletion.

Replication kinetics

The replication of the ΔvIL4 (CK passage 2) as compared to its parental 1874C5 strain (CK passage 5) was evaluated in the LMH cell line. Briefly, LMH cells were seeded at 1.4×10^6 cells in 2.5 ml of growth media (DMEM supplemented with 10% v/v fetal bovine serum) into 0.5% gelatin-treated six-well-plates (Cell BIND, Corning) and incubated at 37°C for 24 h. The cell monolayers were inoculated at an MOI of 0.002 with the ΔvIL4 or the parental 1874C5 strain. Viruses were adsorbed for 60 min at 39°C. After absorption the inoculum was removed and replaced with 2.5 ml of growth media supplemented with 2% fetal bovine serum. At 0 (immediately after media replacement), 24, 48, 72 and 96 hours post-incubation after replacing growth media, supernatant and cells from three independent wells (n=3) were separately collected and divided into two 1-ml aliquots for storage at -80°C. One of the aliquots was used to determine viral titer in LMH cells. Viral titers were expressed as the tissue culture infectious dose 50 perl ml (TCID$_{50}$/ml) using the Reed and Meunch method [52], The second aliquot collected per well was utilized to extract total DNA and titrate viral genome copy numbers per sample.

Experimental design to assess in vivo *expression of vIL-4*

A total of 60 specific pathogen free (SPF) eggs were obtained from a commercial source (Charles River Laboratories Inc., Wilmington, MA, USA) and incubated at 37.5°C and 55% relative humidity (RH) in a small-scale hatcher (Natureform Inc., Jacksonville, FL, USA) at the Poultry Diagnostic Research Center – University of Georgia. At 19 days of embryonation, eggs were candled and infertile eggs and nonviable embryos were removed. At hatch, 54 chickens were distributed in three groups of 18 and each group was housed in separate isolation units with filtered negative pressure air flow (6 chickens per unit). The chickens received water and a

standard diet *ad libitum* during the duration of the experiment. At 18 days of age, two groups of chickens (n=18) were inoculated with either virulent strain 1874C5 or the CEO vaccine Laryngo-Vac® (Zoetis, Animal Health). Viruses were administered at a dose of $10^{3.8}$ TCID$_{50}$ in a 33 μL volume via eye drop. The third group of chickens (n=18) was mock inoculated in a similar fashion with cell culture media and served as the negative control (Mock). At 3-, 5-, and 7-days post-inoculation (p.i.), six chickens per group were humanely euthanized using CO_2 inhalation. The upper and lower conjunctiva from both eyes were collected in 2 mL microcentrifuge tubes with 1 mL of RNA Later (Invitrogen, Waltham, MA, USA) and stored at −80°C for RNA extraction, cDNA synthesis, and library preparation for sequencing.

Experimental design to assess parental 1874C5 and ΔvIL-4 pathogenicity

To evaluate whether the deletion of the vIL-4 gene changed the pathogenicity of the virulent 1874C5 strain, 24 days old specific pathogen free (SPF) chickens were inoculated with the parental 1874C5 or ΔvIL-4 strains via the intratracheal route. Briefly, seventy SPF eggs were obtained from a commercial source (ASV Bio Inc., Wilmington, MA, USA) and incubated at 37.5°C and 55% relative humidity (RH) in a small-scale hatcher (Natureform Inc., Jacksonville, FL, USA) at the Poultry Diagnostic Research Center – University of Georgia. At 19 days of embryonation, eggs were candled and infertile eggs and nonviable embryos were removed. At hatch, a total of 66 chickens were housed in isolation units with filtered negative pressure air flow. Six to seven chickens were housed per unit. Chickens received water and a standard diet *ad libitum* during the duration of the experiment. At 24 days of age, two groups of chickens (n=20 to 22) were inoculated with the parental 1874C5 or ΔvIL-4 strains. Chickens were inoculated with a 100 μl inoculum via the intratracheal route at a $\log_{10}$ 3.65 TCID$_{50}$ dose of either viral strain. A third group of chickens (n=19) was mock-inoculated intratracheally with cell culture media. This group served as the negative control (Mock). From 2 to 7 days post inoculation (p.i.), clinical signs scores and percent survivability were documented. At 3- and 5-days p.i. tracheal swabs were collected from each group of chickens (Mock, parental 1875C5, and ΔvIL-4, n= 14 to 17) to evaluate virus genome load by real-time PCR. At four days p.i. five chickens that were not previously used to collect tracheal swabs were humanly sacrificed by CO_2 inhalation from each group (Mock, parental 1875C5, and ΔvIL-4). The trachea was removed and an upper trachea segment (0.5 cm), caudal to the larynx, was collected from each chicken and fixed in 10% buffered formalin v/v, embedded in paraffin, sectioned (5 μm), and stained with hematoxylin and eosin (H&E) for examination of microscopic lesions and trachea epithelium changes induced after four days of ILTV inoculation. During the length of the experiment, chickens were given a standard feed diet, and water was provided *ad libitum*. All animal experiments described in this study were performed under the Animal Use Protocol A2018 06-009-Y2-A0 approved by the Animal Care and Use Committee (IACUC) in accordance with regulations of the Office of the Vice President for Research at the University of Georgia.

Clinical signs

Clinical signs were evaluated for the categories of conjunctivitis, dyspnea, and lethargy. Each category was scored on a scale of 0 to 3, as previously described [5]. Briefly, the scoring of

individual categories (of conjunctivitis, dyspnea, and lethargy) was considered 0 for normal; 0.5 or 1 for mild signs; 1.5 or 2 for moderate signs; and 2.5 or 3 for severe signs. The total clinical score per chicken was calculated based on the sum of the three categories scores. The sum of scores for individual clinical sign categories (total clinical signs) per chicken, and the average clinical sign total score per group per time point was calculated.

Trachea microscopic lesions.

Upper trachea sections collected at four days p.i. from Mock, parental 1874C5 and ΔvIL-4- inoculated chickens were microscopically examined. Each trachea section received a score to reflect the level of microscopic lesions induced by ILTV infection. Lesions were scored according to the criteria previously described by Guy et al [36]. Briefly, a score scale from 0 to 5 was utilized. Score 0: normal epithelium. Score 1: normal epithelium with mild to moderate lymphocytic infiltration and absence of syncytia with intranuclear inclusion bodies. Score 2: normal epithelium with mild to moderate lymphocytic infiltration and few foci of syncytia with intranuclear inclusion bodies. The latter are the two characteristic lesions that define ILTV lytic replication. Score 3: affected epithelium shows moderate to marked hyperemia, lymphocytic infiltration, and numerous syncytia with intranuclear inclusion bodies. Score 4: mucosa shows areas without epithelium and syncytia with intranuclear inclusion bodies sometimes present. Score 5: Mucosa has no residual epithelium and occasional presence of syncytia and intranuclear inclusion bodies.

Also, the amount of mucosa epithelium and submucosa remaining in the trachea rings collected from Parent 1874C5 IT and ΔvIL-4 IT groups of chickens was assessed microscopically. Briefly, each trachea ring (n=5 per group) was divided digitally in six segments (1 to 6 clockwise) and in each segment (n=30 per group) was scored. A score of 0 was conferred to trachea segments where only a thin layer of submucosa distal to the trachea cartilage remained. A score of 1 was conferred to trachea segments with expanded submucosa, due to increased infiltration, but without any mucosal epithelium remaining. A score of 2 was conferred to trachea segments where submucosa with a distal thin layer of mucosal epithelium remained in place after four days post inoculation. The number of trachea segments categorized as score of 0, 1, or 2 for chickens inoculated with the parental 1874C5 or the ΔvIL-4 strain are presented.

DNA extraction

Tracheal swabs were placed in 2 ml tubes with 1×PBS. All samples were stored at −80°C until processed. Total DNA from feather pulps and tracheas were extracted using the MagaZorb® DNA mini-prep kit (Promega, Madison, WI), following the manufacturer's recommendations with a few modifications. Briefly, using a 96-well plate, 7 μl of proteinase K (PK) solution was loaded into each well, followed by 70 μl of sample and 50 μl of lysis buffer per well. The plate was then incubated at 56°C for 10 min, 10 μl of magnetic beads and 125 μl of binding buffer were added per well, followed by rotating the plate for 10 min at room temperature. The supernatant and magnetic beads were separated using a magnetic stand and the supernatant was discarded. Finally, using 250 μl of washing buffer, the beads were washed twice, and DNA was

eluted from the beads with 100 μl of elution buffer after rotating the plate for 10 min at room temperature.

RNA extraction, cDNA synthesis, and RNA-Seq for in vivo *expression*

Conjunctivas were thawed and transferred into lysing bead matrix tubes (MP Biomedicals, Santa Ana, CA) containing 900 μL of Qiazol (Qiagen, Germantown, MD, USA), incubated for 30 minutes and homogenized using the FastPrep-24 5G (MP Biomedical, Santa Ana, CA, USA) at 6.0 m/s for 20 seconds. Total RNA from conjunctiva tissues was extracted using the RNeasy Plus universal kit (Qiagen, Germantown, MD, USA) according to the manufacturer's protocol. RNA concentrations and purity were determined with the Nanodrop 2000c spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA). cDNA was synthesized from 1 μg of total RNA using the RT2 First Strand Kit (Qiagen, Germantown, MD, USA), following the manufacturer's guidelines. Stranded sequencing libraries were prepared from ribo-depleted RNA preparations using KAPA Stranded mRNA kit and sequenced on an Illumina NextSeq 2000 instrument using a high output flow cell at the Georgia Genomics and Bioinformatic Core at the University of Georgia.

Duplex real-time PCR

ILTV genomes were quantified from LMH infected cultures and from tracheal swabs collected at days 3 and 5 p.i. using a duplex real-time PCR. The ILTV genomes were amplified with primers that target the UL44 ILTV gene, and chicken genome was detected with primers that target the chicken α2 collagen gene as previously described [73]. The relative amount of viral DNA per sample was calculated as the $\log_{10}$ 2-$\Delta\Delta C^t$ as previously described [74].

Statistical analysis of replication and in vivo *results*

The Shapiro–Wilk test was used to test the normality of the data. ILTV genome load data were analyzed using one-way ANOVA followed by Tukey's multiple comparisons test and with the non-parametric Mann-Whitney test. Clinical signs and trachea histopathological microscopic scores were evaluated using the Kruskal–Wallis test followed by the Dunn's multiple comparisons test and with the non-parametric Mann-Whitney test. All statistical analyses were performed using the GraphPad Prism software 10 (GraphPad Software, Inc., La Jolla, CA). All reported p-values were based on two-sided comparisons. Statistical differences were defined as significant when they reached p<0.05.

Manuscript preparation

The manuscript was prepared using RMarkdown with heavy use of the ggplot2 library [75] for generating figures. Figure 1 was generated using the Gviz R package v1.46.1 [76]. Pymol [77] was used for visualization of 3D modeling. The ggtree [78] R package was used for tree visualization. All supplementary code used for data format manipulation and processing is available from the manuscript git repository (see *Data Availability*).

References

[1]   Garcia M, Spatz S, Guy JS. Infectious laryngotracheitis. In: Swayne DE, Glisson JR, McDougald LR, Nolan LK, Suarez DL, Nair V, editors. Diseases of poultry, vol. 13th edition, Ames, IA: Wiley-Blackwell; 2013.

[2]   Fahey KJ, Bagust TJ, York JJ. Laryngotracheitis herpesvirus infection in the chicken: The role of humoral antibody in immunity to a graded challenge infection. Avian Pathology 1983;12:505–14.

[3]   Bagust TJ, Jones RC, Guy JS. Avian infectious laryngotracheitis. Revue Scientifique Et Technique 2000;19:483–92.

[4]   Honda T, Okamura H, Taneno A, Yamada S, Takahashi E. The role of cell-mediated immunity in chickens inoculated with the cell-associated vaccine of attenuated infectious laryngotracheitis virus. J Vet Med Sci 1994;56:1051–5.

[5]   Maekawa D, Whang P, Riblet SM, Hurley DJ, Guy JS, García M. Assessing the infiltration of immune cells in the upper trachea mucosa after infectious laryngotracheitis virus (ILTV) vaccination and challenge. Avian Pathology 2021;50:540–56. https://doi.org/10.1080/03079457.2021.1989379.

[6]   Fahey KJ, York JJ. The role of mucosal antibody in immunity to infectious laryngotracheitis virus in chickens. J Gen Virol 1990;71 (Pt 10):2401–5.

[7]   Sabir AJ, Adams TE, O'Rourke D, Devlin JM, Noormohammadi AH. Investigation onto the correlation between systemic antibodies to surface glycoproteins of infectious laryngotracheitis virus (ILTV) and protective immunity. Vet Microbiol 2019;228:252–8. https://doi.org/10.1016/j.vetmic.2018.12.010.

[8]   Wolfrum N. Infectious laryngotracheitis: An update on current approaches for prevention of an old disease. J Anim Sci 2020;98:S27–s35. https://doi.org/10.1093/jas/skaa133.

[9]   Koonin EV, Dolja VV, Krupovic M. The logic of virus evolution. Cell Host & Microbe 2022;30:917–29. https://doi.org/10.1016/j.chom.2022.06.008.

[10]   Alcami A. Viral mimicry of cytokines, chemokines and their receptors. Nature Reviews Immunology 2003;3:36–50. https://doi.org/10.1038/nri980.

[11]   Neipel F, Albrecht JC, Ensser A, Huang YQ, Li JJ, Friedman-Kien AE, et al. Human herpesvirus 8 encodes a homolog of interleukin-6. Journal of Virology 1997;71:839–42. https://doi.org/10.1128/jvi.71.1.839-842.1997.

[12]   Yao Z, Fanslow WC, Seldin MF, Rousseau A-M, Painter SL, Comeau MR, et al. Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. Immunity 1995;3:811–21. https://doi.org/10.1016/1074-7613(95)90070-5.

[13] Moore KW, Vieira P, Fiorentino DF, Trounstine ML, Khan TA, Mosmann TR. Homology of cytokine synthesis inhibitory factor (IL-10) to the Epstein-Barr virus gene BCRFI. Science (New York, NY) 1990;248:1230–4. `https://doi.org/10.1126/science.2161559`.

[14] Kotenko SV, Saccani S, Izotova LS, Mirochnitchenko OV, Pestka S. Human cytomegalovirus harbors its own unique IL-10 homolog (cmvIL-10). Proceedings of the National Academy of Sciences 2000;97:1695–700. `https://doi.org/10.1073/pnas.97.4.1695`.

[15] Mahmoudian A, Markham PF, Noormohammadi AH, Browning GF. Kinetics of transcription of infectious laryngotracheitis virus genes. Comp Immunol Microbiol Infect Dis 2012;35:103–15. `https://doi.org/10.1016/j.cimid.2011.11.001`.

[16] Smith RH, Caughman GB, O'Callaghan DJ. Characterization of the regulatory functions of the equine herpesvirus 1 immediate-early gene product. J Virol 1992;66:936–45. `https://doi.org/10.1128/JVI.66.2.936-945.1992`.

[17] Wagner EK, Ramirez JJ, Stingley SW, Aguilar SA, Buehler L, Devi-Rao GB, et al. Practical approaches to long oligonucleotide-based DNA microarray: Lessons from herpesviruses. Prog Nucleic Acid Res Mol Biol 2002;71:445–91.

[18] Lee J, Bottje WG, Kong BW. Genome-wide host responses against infectious laryngotracheitis virus vaccine infection in chicken embryo lung cells. BMC Genomics 2012;13:143. `https://doi.org/10.1186/1471-2164-13-143`.

[19] Lee JY, Song JJ, Wooming A, Li X, Zhou H, Bottje WG, et al. Transcriptional profiling of host gene expression in chicken embryo lung cells infected with laryngotracheitis virus. BMC Genomics 2010;11:445.

[20] Luo J, Carrillo JA, Menendez KR, Tablante NL, Song J. Transcriptome analysis reveals an activation of major histocompatibility complex 1 and 2 pathways in chicken trachea immunized with infectious laryngotracheitis virus vaccine. Poult Sci 2014;93:848–55. `https://doi.org/10.3382/ps.2013-03624`.

[21] Wang Z, Sun B, Gao Q, Ma Y, Liang Y, Chen Z, et al. Host Src controls gallid alpha herpesvirus 1 intercellular spread in a cellular fatty acid metabolism-dependent manner. Virology 2019;537:1–13. `https://doi.org/10.1016/j.virol.2019.08.011`.

[22] Boldogkői Z, Moldován N, Balázs Z, Snyder M, Tombácz D. Long-read sequencing – a powerful tool in viral transcriptome research. Trends Microbiol 2019;27:578–92.

[23] Rhoads A, Au KF. PacBio Sequencing and Its Applications. Genomics Proteomics Bioinformatics 2015;13:278–89. `https://doi.org/10.1016/j.gpb.2015.08.002`.

[24] Tombacz D, Csabai Z, Olah P, Balazs Z, Liko I, Zsigmond L, et al. Full-Length Isoform Sequencing Reveals Novel Transcripts and Substantial Transcriptional Overlaps in a Herpesvirus. PLoS One 2016;11:e0162868. `https://doi.org/10.1371/journal.pone.0162868`.

[25] Tombacz D, Torma G, Gulyas G, Fulop A, Dormo A, Prazsak I, et al. Hybrid sequencing discloses unique aspects of the transcriptomic architecture in equid alphaherpesvirus 1. Heliyon 2023;9:e17716. `https://doi.org/10.1016/j.heliyon.2023.e17716`.

[26] Betts MJ, Guigó R, Agarwal P, Russell RB. Exon structure conservation despite low sequence similarity: A relic of dramatic events in evolution? The EMBO Journal 2001;20:5354–60. `https://doi.org/10.1093/emboj/20.19.5354`.

[27] Chen D, Hosner PA, Dittmann DL, O'Neill JP, Birks SM, Braun EL, et al. Divergence time estimation of Galliformes based on the best gene shopping scheme of ultraconserved elements. BMC Ecology and Evolution 2021;21:209. `https://doi.org/10.1186/s12862-021-01935-1`.

[28] Lee S-W, Devlin JM, Markham JF, Noormohammadi AH, Browning GF, Ficorilli NP, et al. Phylogenetic and molecular epidemiological studies reveal evidence of multiple past recombination events between infectious laryngotracheitis viruses. PLoS ONE 2013;8:e55121. `https://doi.org/10.1371/journal.pone.0055121`.

[29] Martínez-Vicente P, Poblador F, Leitner J, Farré D, Steinberger P, Engel P, et al. Discovery of the first PD-1 ligand encoded by a pathogen. Frontiers in Immunology 2022;13. `https://doi.org/10.3389/fimmu.2022.1007334`.

[30] Ou SC, Giambrone JJ. Infectious laryngotracheitis virus in chickens. World J Virol 2012;1:142–9.

[31] Gibbs A. Molecular evolution of viruses; 'trees', 'clocks' and 'modules'. Journal of Cell Science 1987;1987:319–37. `https://doi.org/10.1242/jcs.1987.Supplement_7.22`.

[32] Huang H, Lavoie-Lamoureux A, Moran K, Lavoie JP. IL-4 stimulates the expression of CXCL-8, E-selectin, VEGF, and inducible nitric oxide synthase mRNA by equine pulmonary artery endothelial cells. Am J Physiol Lung Cell Mol Physiol 2007;292:L1147–54. `https://doi.org/10.1152/ajplung.00294.2006`.

[33] Paoliello-Paschoalato AB, Oliveira SH, Cunha FQ. Interleukin 4 induces the expression of inducible nitric oxide synthase in eosinophils. Cytokine 2005;30:116–24. `https://doi.org/10.1016/j.cyto.2005.01.001`.

[34] Xin J, Ohmori K, Nishida J, Zhu Y, Huang H. The initial response of CD4+ IL-4-producing cells. Int Immunol 2007;19:305–10. `https://doi.org/10.1093/intimm/dxl147`.

[35] Xu L, Huang Y, Yang J, Van Der Meide PH, Levi M, Wahren B, et al. Dendritic cell-derived nitric oxide is involved in IL-4-induced suppression of experimental allergic encephalomyelitis (EAE) in Lewis rats. Clin Exp Immunol 1999;118:115–21. `https://doi.org/10.1046/j.1365-2249.1999.01029.x`.

[36]   Guy JS, Barnes HJ, Morgan LM. Virulence of infectious laryngotracheitis viruses: Comparison of modified-live vaccine viruses and North Carolina field isolates. Avian Diseases 1990-01/1990-03;34:106–13.

[37]   Gorshkova EA, Shilov ES. Possible Mechanisms of Acquisition of Herpesvirus Virokines.   Biochemistry (Mosc) 2016;81:1350–7.   `https://doi.org/10.1134/s0006297916110122`.

[38]   Parcells MS, Lin S-F, Dienglewicz RL, Majerciak V, Robinson DR, Chen H-C, et al. Marek's disease virus (MDV) encodes an interleukin-8 homolog (vIL-8): Characterization of the vIL-8 protein and a vIL-8 deletion mutant MDV. Journal of Virology 2001;75:5159–73. `https://doi.org/10.1128/jvi.75.11.5159-5173.2001`.

[39]   Sin SH, Dittmer DP. Cytokine homologs of human gammaherpesviruses. J Interferon Cytokine Res 2012;32:53–9. `https://doi.org/10.1089/jir.2011.0083`.

[40]   Slobedman B, Barry PA, Spencer JV, Avdic S, Abendroth A. Virus-encoded homologs of cellular interleukin-10 and their control of host immune function. J Virol 2009;83:9618–29. `https://doi.org/10.1128/jvi.01098-09`.

[41]   Vieira P, de Waal-Malefyt R, Dang MN, Johnson KE, Kastelein R, Fiorentino DF, et al. Isolation and expression of human cytokine synthesis inhibitory factor cDNA clones: Homology to Epstein-Barr virus open reading frame BCRFI. Proc Natl Acad Sci U S A 1991;88:1172–6. `https://doi.org/10.1073/pnas.88.4.1172`.

[42]   Hsu DH, de Waal Malefyt R, Fiorentino DF, Dang MN, Vieira P, de Vries J, et al. Expression of interleukin-10 activity by Epstein-Barr virus protein BCRF1. Science 1990;250:830–2. `https://doi.org/10.1126/science.2173142`.

[43]   Kurilla MG, Swaminathan S, Welsh RM, Kieff E, Brutkiewicz RR. Effects of virally expressed interleukin-10 on vaccinia virus infection in mice. J Virol 1993;67:7623–8. `https://doi.org/10.1128/jvi.67.12.7623-7628.1993`.

[44]   Poole E, Neves TC, Oliveira MT, Sinclair J, da Silva MCC. Human Cytomegalovirus Interleukin 10 Homologs: Facing the Immune System. Front Cell Infect Microbiol 2020;10:245. `https://doi.org/10.3389/fcimb.2020.00245`.

[45]   Chang WL, Barry PA, Szubin R, Wang D, Baumgarth N. Human cytomegalovirus suppresses type I interferon secretion by plasmacytoid dendritic cells through its interleukin 10 homolog. Virology 2009;390:330–7. `https://doi.org/10.1016/j.virol.2009.05.013`.

[46]   Cheung AK, Gottlieb DJ, Plachter B, Pepperl-Klindworth S, Avdic S, Cunningham AL, et al. The role of the human cytomegalovirus UL111A gene in down-regulating CD4+ T-cell recognition of latently infected cells: Implications for virus elimination during latency. Blood 2009;114:4128–37. `https://doi.org/10.1182/blood-2008-12-197111`.

[47]   Lin YL, Chang PC, Wang Y, Li M. Identification of novel viral interleukin-10 isoforms of human cytomegalovirus AD169. Virus Res 2008;131:213–23. `https://doi.org/10.1016/j.virusres.2007.09.011`.

[48] Knappe A, Hiller C, Niphuis H, Fossiez F, Thurau M, Wittmann S, et al. The interleukin-17 gene of herpesvirus saimiri. J Virol 1998;72:5797–801. `https://doi.org/10.1128/jvi.72.7.5797-5801.1998`.

[49] Oldoni I, Rodriguez-Avila A, Riblet SM, Zavala G, Garcia M. Pathogenicity and growth characteristics of selected infectious laryngotracheitis virus strains from the United States. Avian Pathol 2009;38:47–53.

[50] Rodriguez-Avila A, Oldoni I, Riblet S, Garcia M. Replication and transmission of live attenuated infectious laryngotracheitis virus (ILTV) vaccines. Avian Diseases 2007;51:905–11.

[51] Rodriguez-Avila A, Oldoni I, Riblet S, Garcia M. Evaluation of the protection elicited by direct and indirect exposure to live attenuated infectious laryngotracheitis virus vaccines against a recent challenge strain from the United States. Avian Pathol 2008;37:287–92. `https://doi.org/10.1080/03079450802043742`.

[52] Reed LJ, Muench H. A simple method of estimating fifty percent endpoints. Am J Hygiene 1938;27,:493–7.

[53] Grüning B, Dale R, Sjödin A, Chapman BA, Rowe J, Tomkins-Tinch CH, et al. Bioconda: Sustainable and comprehensive software distribution for the life sciences. Nature Methods 2018;15:475–6. `https://doi.org/10.1038/s41592-018-0046-7`.

[54] Tardaguila M, Fuente L de la, Marti C, Pereira C, Pardo-Palacios FJ, Risco H del, et al. SQANTI: Extensive characterization of long-read transcript sequences for quality control in full-length transcriptome identification and quantification. Genome Research 2018;28:396–411. `https://doi.org/10.1101/gr.222976.117`.

[55] Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): High-performance genomics data visualization and exploration. Briefings in Bioinformatics 2013;14:178–92. `https://doi.org/10.1093/bib/bbs017`.

[56] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. Journal of Molecular Biology 1990;215:403–10. `https://doi.org/10.1006/jmbi.1990.9999`.

[57] Li H. Protein-to-genome alignment with miniprot. Bioinformatics (Oxford, England) 2023;39:btad014. `https://doi.org/10.1093/bioinformatics/btad014`.

[58] Stanke M, Schöffmann O, Morgenstern B, Waack S. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. BMC Bioinformatics 2006;7:62. `https://doi.org/10.1186/1471-2105-7-62`.

[59] Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nature Biotechnology 2016;34:525–7. `https://doi.org/10.1038/nbt.3519`.

[60] Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology 2019;37:907–15. https://doi.org/10.1038/s41587-019-0201-4.

[61] Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic features. Bioinformatics 2010;26:841–2. https://doi.org/10.1093/bioinformatics/btq033.

[62] Drozdetskiy A, Cole C, Procter J, Barton GJ. JPred4: A protein secondary structure prediction server. Nucleic Acids Research 2015;43:W389–94. https://doi.org/10.1093/nar/gkv332.

[63] Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et al. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nature Biotechnology 2022;40:1023–5. https://doi.org/10.1038/s41587-021-01156-3.

[64] Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein structure prediction with AlphaFold. Nature 2021;596:583–9. https://doi.org/10.1038/s41586-021-03819-2.

[65] Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Molecular Biology and Evolution 2013;30:772–80. https://doi.org/10.1093/molbev/mst010.

[66] Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution 2020;37:1530–4. https://doi.org/10.1093/molbev/msaa015.

[67] Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 2017;14:587–9. https://doi.org/10.1038/nmeth.4285.

[68] Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution 2018;35:518–22. https://doi.org/10.1093/molbev/msx281.

[69] Scott DJ, Gunn NJ, Yong KJ, Wimmer VC, Veldhuis NA, Challis LM, et al. A Novel Ultra-Stable, Monomeric Green Fluorescent Protein For Direct Volumetric Imaging of Whole Organs Using CLARITY. Sci Rep 2018;8:667. https://doi.org/10.1038/s41598-017-18045-y.

[70] Waldo GS, Standish BM, Berendzen J, Terwilliger TC. Rapid protein-folding assay using green fluorescent protein. Nat Biotechnol 1999;17:691–5. https://doi.org/10.1038/10904.

[71] Spatz SJ, Volkening JD, Mullis R, Li F, Mercado J, Zsak L. Expression of chicken parvovirus VP2 in chicken embryo fibroblasts requires codon optimization for production of naked DNA and vectored meleagrid herpesvirus type 1 vaccines. Virus Genes 2013;47:259–67. https://doi.org/10.1007/s11262-013-0944-9.

[72] Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, et al. Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol 2016;34:184–91. https://doi.org/10.1038/nbt.3437.

[73] Vagnozzi A, Riblet SM, Zavala G, Garcia M. Optimization of a duplex real-time PCR method for relative quantitation of infectious laryngotracheitis virus. Avian Diseases 2012;56:406–10.

[74] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 2001;25:402–8. https://doi.org/10.1006/meth.2001.1262.

[75] Wickham H. Ggplot2: Elegant Graphics for Data Analysis. Cham: Springer International Publishing; 2016. https://doi.org/10.1007/978-3-319-24277-4.

[76] Hahne F, Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor. In: Mathé E, Davis S, editors. Statistical Genomics: Methods and Protocols, New York, NY: Springer; 2016, p. 335–51. https://doi.org/10.1007/978-1-4939-3578-9_16.

[77] Schrödinger, LLC. The PyMOL molecular graphics system, version 1.8 2015.

[78] Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Current Protocols in Bioinformatics 2020;69:e96. https://doi.org/10.1002/cpbi.96.

Supplementary materials

cIL4 Inserts		
VTRmuGFPFORC-IL4	TCAGATCCGCTAGCGCCGCCATGCCAGGCAAGAGCAAGCC	Set 1
muGFP C-IL4 REV	CACAGACAGCTGCAGACACAGTGTGAACTCTCCGCTTCCAGCTGC	
C-IL4 MuGFP For	GCAGCTGGAAGCGGAGAGTTCACACTGTGTCTGCAGCTGTCTGTG	Set2
cIL4 Rev	TACCGTCGACTGCAGAATTCGAAGCTTGAGCTCGAGTCACTTGTTCTTGGCCAGC	
cIL4 vector		
vtr cIL4 for	CACCAGCTGGCCAAGAACAAGTGACTCGAGCTCAAGCTTCGAATTCTGCAGTC	Set 3
vtr MuGFP REV C-IL4	GGCTTGCTCTTGCCTGGCATGGCGGCGCTAGCGGATCTG	
vIL4 Inserts		
VirIL4 For	CAGATCCGCTAGCAGACGCCACCATGCCAGGCAAGAGCAAGCCAC	Set 4
muGFP vIL4 REV	TGGAGCGGCTCTAATGCTGCGGAACTCTCCGCTTCCAGCTGC	
vIL4 MuGFP For	GCAGCTGGAAGCGGAGAGTTCCGCAGCATTAGAGCCGCTCCA	Set 5
Vir IL4 Rev	AGCTTGAGCTCGAGAGTTACATCAGTACTGCCTGCTCTGTCTGACGT	
vIL4 vector		
vtr Vir IL4 for	GATACGTCAGACAGAGCAGGCAGTACTGATGTAACTCTCGAGCTCAAGCT	Set 6
vtr Vir IL4 rev	GTGGTGGCTTGCTCTTGCCTGGCATGGTGGCGTCTGCTAGCGGAT	

Figure S1: List of primers used to generate PCR amplicons assembled to create N-terminal tagged versions of cIL-4 and vIL-4.

Figure S2: (A) Diagram of the plasmid constructs expressing GFP-tagged cIL-4 and vIL-4 proteins. Fluorescent microscopy of LMH cells transfected with (B) GFP-cIL-4 and (C) GFP-vIL4. Magnification is 100×.

Table S1: Vertebrate IL-4 homologs used in the analysis. Accessions are from GenBank and are either the direct protein entries used or the genomic sequences on which the predicted genes are located (as indicated in the Notes).

Common name	Scientific name	Accession	Notes
vIL-4	Gallid alpha-herpesvirus 1	N/A	from GaHV-1 1874C5 IsoSeq and internal 1874C5 corrected genome
chicken	Gallus gallus	NM_001398459.1	used as-is
Chinese bamboo-partridge	Bambusicola thoracicus	POI22934.1	miniprot/AUGUSTUS prediction, matches indicated protein accession
turkey	Meleagris gallopavo	ENSMGAT00000023360.1	GenBank NM_001303181.1 is actually IL13
ring-necked pheasant	Phasianus colchicus	NW_022205427.1	miniprot/AUGUSTUS prediction
greater sage-grouse	Centrocercus urophasianus	JAHKSY010000026.1	miniprot/AUGUSTUS prediction
rock ptarmigan	Lagopus muta	CM041489.1	miniprot/AUGUSTUS prediction
green peafowl	Pavo muticus	JACDJE010000203.1	miniprot/AUGUSTUS prediction
red-legged partridge	Alectoris rufa	CAMZON010000379.1	miniprot/AUGUSTUS prediction
Japanese quail	Coturnix japonica	XM_015875546.1	used as-is
guineafowl	Numida meleagris	XM_021410541.1	used as-is
guan	Penelope pileata	WBMW01000786.1	miniprot/AUGUSTUS prediction
bobwhite	Colinus virginianus	AWGT02000115.1	miniprot/AUGUSTUS, structure differs from OXB76676
brush-turkey	Alectura lathami	VXAV01004783.1	miniprot/AUGUSTUS prediction
tufted duck	Aythya fuligula	XM_032196612.1	used as-is
mallard	Anas platyrhynchos	NC_051785.1	miniprot/AUGUSTUS, several nucleotide positions differ from MF346730.1
black swan	Cygnus atratus	XM_035560814.1	used as-is

Table S1: Vertebrate IL-4 homologs used in the analysis. Accessions are from GenBank *(continued)*

Common name	Scientific name	Accession	Notes
egret	Egretta garzetta	XM_009647167.1	used as-is
Adelie penguin	Pygoscelis adeliae	XM_009322783.1	used as-is
goshawk	Accipiter gentilis	XM_049829109.1	used as-is
golden eagle	Aquila chrysaetos	XM_029998407.1	used as-is
zebra finch	Taeniopygia guttata	NC_044225.2	miniprot/AUGUSTUS, structure differs from XM_041718987.1
song sparrow	Melospiza melodia maxima	RZID01002771.1	used as-is
ostrich	Struthio camelus	JJRT01005166.1	miniprot/AUGUSTUS, structure differs from XM_009679367.1
emu	Dromaius no-vaehollandiae	NW_020453704.1	miniprot/AUGUSTUS, structure differs from XP_025974110.1
tinamou	Nothoprocta perdicaria	NW_020454172.1	miniprot/AUGUSTUS, structure differs from XM_026045620.1
saltwater crocodile	Crocodylus porosus	NW_017728914.1	miniprot/AUGUSTUS prediction
alligator	Alligator sinensis	XM_006023567.1	used as-is
slider turtle	Trachemys scripta elegans	NC_048305.1	miniprot/AUGUSTUS, structure differs from XM_03477890.1
green sea turtle	Chelonia mydas	NC_057854.1	miniprot/AUGUSTUS prediction