Genetic and morphological characterization of *Metacollinia emscheri* n. sp., a

novel sanguicolous apostome ciliate of the freshwater amphipods *Gammarus*

pulex* and *G. fossarum

Sebastian Prati[1,2]*, Sonja Rückert[2,3], Daniel S. Grabner[1,2], Bernd Sures[1,2,4], Jamie Bojko[5,6]

[1]Aquatic Ecology, University of Duisburg-Essen, Universitaetsstr. 5, 45141 Essen, Germany.

[2]Centre for Water and Environmental Research, University of Duisburg-Essen, Universitaetsstr.

5, 45141 Essen, Germany. [3]Eukaryotic Microbiology, University of Duisburg-Essen,

Universitaetsstr. 5, 45141 Essen, Germany. [4]Research Center One Health Ruhr of the University

Alliance Ruhr, University of Duisburg-Essen, Universitätsstraße 5, 45141 Essen, Germany.

[5]National Horizons Centre, Teesside University, Darlington, DL1 1HG, United Kingdom. [6]School

of Health and Life Sciences, Teesside University, Middlesbrough, TS1 3BX, United Kingdom

*Corresponding author: sebastian.prati@uni-due.de

Keywords: Collinidae, endoparasitic ciliate, amphipod pathogen, crustacea, mitogenome

Abstract

We describe a novel sanguicolous parasitic ciliate, *Metacollinia emscheri* n. sp., found in the

freshwater amphipods *Gammarus pulex* and *G. fossarum*. This ciliate infected up to 20% of the

amphipods collected in a German stream catchment, the Boye, a tributary of the river Emscher.

The ciliate showed morphological characteristics fitting the genus *Metacollinia*. Different life

stages of variable size occurred simultaneously in the hemocoel throughout the hosts' body. The

tomont had 40-47 slightly spiraled kineties, a non-ciliated cortical band, a large macronucleus, and

contractile vacuoles arranged in rows or scattered throughout the cytoplasm. The

protomites/tomites with nine somatic kineties presented evidence of the buccal kineties x, y, and

z reminiscent of those of the order Foettingeriida. Phylogenetic analyses of the 18S rRNA and COI

regions confirm the ciliate placement in the Collinidae and a close relatedness to the type species

of the genus *Metacollinia*, *Metacollinia lucensis*. Hence, we formally describe this parasite as

Metacollinia emscheri n. sp. The systemic infections observed in histological preparations and the pathogenicity of *Metacollinia emscheri* n. sp. suggest that this parasite might influence host population dynamics. Given the ecological importance of amphipods as keystone species in freshwater ecosystems, an outbreak of this parasite might indirectly impact ecosystem functioning.

1. Introduction

Macroinvertebrates, including amphipods, are essential components of stream ecosystem food webs. Amphipods are highly diversified and ubiquitous keystone organisms that often represent the dominant element of invertebrate communities (Giari et al., 2020; Macneil et al., 1997). Their shredding activity can account for up to 75% of overall leaf-litter breakdown in some ecosystems, and they are an important food source for a wide range of predators, linking several trophic levels within riverine food webs (Giari et al., 2020). Their ecological niche makes them ideal hosts for a hyper-diverse range of pathogens ranging from viruses to trophically transmitted metazoans (Bojko and Ovcharenko, 2019). Yet most studies focused on metazoan parasites and, to a lesser extent, on microsporidians due to their abundance, diversity, and ecological relevance (Prati et al., 2023), as well as gregarine apicomplexans (McKinley et al. 2024). Thus, a large portion of less conspicuous but highly diverse pathogens is expected to remain unnoticed.

Protists, such as ciliates, are commonly found as commensal epibionts on the carapace, gills, and extremities of amphipods, where they benefit from the host movement to collect particular matter. However, few apostome ciliates have specialized symbiotic and parasitic relationships with crustaceans, including amphipods (Lynn, 2008). Among them, sanguicolous parasites are able to penetrate the host cuticle and feed on the cells and fluid of the hemocoel (Lynn et al., 2014; Lynn and Strüder-Kypke, 2019). Representatives of the Collinidae and the Pseudocollinidae families reproduce continuously in the host hemolymph, leading to systemic infections, which might ultimately result in host death (Chantangsi et al., 2013; Lynn et al., 2014; Lynn and Strüder-Kypke, 2019; Prokopowicz et al., 2010). For instance, Pseudocollinidae representatives have been linked to mass mortality events in krill in marine environments (Gómez-Gutiérrez et al., 2012; Gómez-Gutiérrez and Kawaguchi, 2017) and the Collinidae *Lynnia grapsolytica* strongly enhanced mortality rate in infected crabs under laboratory conditions (Metz and Hechinger, 2021).

The majority of Collinidae are parasites of freshwater amphipods and isopods (De Puytorac and Lom, 1962). One of the first observations in freshwater amphipods dates back to 1852 when an apostome ciliate, later described as *Paracollinia branchiarum* was found in the hemolymph of *Gammarus pulex* (De Puytorac and Lom, 1962). Successive investigations revealed the presence of Collinidae in freshwater amphipods from Lake Baikal and subterranean environments in Europe and the USA (Capriulo and Small, 1985). More recently, an unidentified ciliate has been observed in the hemolymph of *Gammarus roeselii* (Bojko et al., 2017). These findings indicate that these parasites might be ubiquitous but poorly investigated.

Given the ecological importance of amphipods and the possible high pathogenicity of ciliates belonging to the Collinidae family, deepening our knowledge of these sanguicolous parasites is paramount. If highly pathogenic, these parasites might strongly influence host population dynamics and, thus, indirectly, ecosystem functioning. We, therefore, document the finding of a novel Collinidae species in the hemocoel of the amphipods *Gammarus pulex* and *G. fossarum,* and formally describe it as *Metacollinia emscheri* n. sp. Besides documenting the morphological characteristics of this sanguicolous ciliate and exploring its pathology via histopathological examinations, we conducting molecular phylogenetic analyses.

2. Materials and methods

2.1 Sampling

Amphipods were collected between December 2022 and March 2024 from eight sampling locations across the Boye catchment in North Rhine-Westphalia, Germany (Table 1). The first collection of amphipods (December 2022) initially aimed at describing novel microsporidian parasites comprised 72 individuals (n = 52 from BOYohKI and n = 20 from VORohBOY, Table 1). Thus, only microsporidian-infected amphipods (n = 20; for molecular identification method, see Prati et al. 2023) were processed for histological analyses. After finding an individual infected with an unknown sanguicolous ciliate, we decided to carry out a non-targeted sampling in April 2023.

All amphipods were kept alive and dissected within 6h post collection. The head and anterior part of the thorax were fixed in 96% ethanol for molecular analyses. The proximate segments of the thorax (2 mm) were placed in 2.5% glutaraldehyde 0.1 M sodium cacodylate buffer (pH 7.4) for electron microscopy. The rest of the body was fixed in Davidson´s freshwater

fixative for 24 hours, then transferred to 70% ethanol for histological examination. Additionally,

given the high morphological variability observed in the ciliates, we collected 42 amphipods in

March 2024. We dissected them and separately stored the freshly isolated ciliates by life stages in

96% ethanol for molecular analyses to confirm their taxonomic placement.

2.2 Histological analyses

Amphipod tissues were infiltrated with paraffin wax in an automated tissue processor (ethanol-

xylene substitute wax), embedded into wax blocks, and sectioned to 4 µm thickness through the

animal center. Each section was mounted on a glass slide and stained with hematoxylin and eosin,

following a standard de-wax and rehydration protocol. The slides were screened for pathogens

using an Olympus CX 33 trinocular light microscope and photographed with a Sony Alpha 6600

mounted via extension tubes on the camera port.

2.3 Scanning Electron Microscopy

Haemocoel from glutaraldehyde-fixed or fresh amphipods was transferred into self-made filter

baskets with a 10 µm polycarbonate membrane filter (Millipore Corp.) attached to a cut-off

BEEM® conical tip capsule. These capsules were placed in 0.9% NaCl in a multi-well plate. A

piece of filter paper was mounted on the inside of the multi-well plate lid and saturated with 4%

(w/v) OsO_4. The ciliates were fixed with OsO_4 vapor for 30 min in the dark. Ten drops of 4% (w/v)

OsO_4 were added directly to the filter baskets, and the samples were fixed again for 30 min. The

filters with the ciliates were washed with distilled water, dehydrated with a graded series of ethyl

alcohol, and critical point dried with CO_2 (Leica EM CPD 300 critical point drier). Filters were

mounted on stubs, sputter coated with 5 nm platinum (Leica ACE 600 sputter coater), and viewed

using the Zeiss Crossbeam 540 scanning electron microscope.

2.4 Quantitative protargol staining

We employed a rapid Quantitative protargol staining (QPS) on ciliates isolated from

glutaraldehyde-fixed amphipods to better observe the internal structures of ciliates. The protocol

(see details in Supplementary material 1) is a modified version of those proposed by Skibbe (1994),

Kurilov (2017), and Ji and Wang (2018). Briefly, the steps employed involved in-house production

of liquid protargol (Kurilov, 2017), the concentration of fixed ciliates on cellulose filters (Skibbe,

1994), coverage of ciliates with a thin layer of agarose, hardening of agarose with formaldehyde,

fast impregnation in warm protargol solution (Skibbe, 1994), dehydration and mounting of the

filter on permanent slides. The process included steps to correct overbleaching and overstaining

(Ji and Wang, 2018).

2.5 Molecular analyses

DNA was isolated from muscular tissue following a modified salt precipitation protocol described

by Grabner et al. (2015). DNA extracted from the only amphipod infected with an unknown ciliate

in December 2022 was quantified using Qubit (Invitrogen) and sequenced using MinIon (Oxford

Nanopore) and Illumina to assemble the ciliate mitogenome. Based on the obtained sequences, we

then conducted a molecular screening of sanguicolous ciliates in amphipods collected in April

2023 using DNA barcoding (Sanger sequencing). DNA barcoding was also used to confirm that

all ciliate life stages isolated in March 2024 belonged to the same species. Furthermore, we

molecularly identified infected hosts using DNA barcodes.

2.6 DNA barcoding

For the molecular screening of sanguicolous ciliates in amphipods collected in April 2023 and for

the   ciliates   isolated   in   March   2024,   we   used   the   primer   pair   SB2F

(TGTAAAACGACGGCCAGTGTTCCCCTTGAACGAGGAATTC)   and   ITS2.2R

(CAGGAAACAGCTATGACCCTGGTTAGTTTCTTTTCCTCCGC) in conjunction with the

pair cil18SspannerF (TGTAAAACGACGGCCAGT-AATCAGAGTAATGATTAATAGG) and

cil18SspannerR (CAGGAAACAGCTATGAC-TGGTTCACCGGACCACTCGA) targeting the

18S rRNA and LSU rRNA region (Adlard and Lester, 1995; Metz and Hechinger, 2021) as these

primers bound well to the mitogenome sequences obtained from the ciliate collected in December

2022 and did not amplify amphipods DNA. All infected amphipods were identified using the

universal eukaryotic primers LCO1490 and HCO2198, which amplify the COI region.

149   Each PCR reaction volume used for the SB2F-ITS2.2R and cil18SspannerF-

cil18SspannerR primer pairs consisted of 20 μL composed of 10 μL of 2 × AccuStart II PCR

ToughMix (Quantabio), 1 μL of each primer (0.5 μM), 0.35 μL of 50 × GelTrack Loading Dye

(Quantabio), 6.65 μL MilliQ water and 1 μL of DNA template. For the LCO1490 and HCO2198

primer pair, the PCR reaction consisted of 20 μL assay with 10 μL of OneTaq Quick-Load 2x

Master mix (New England Biolabs), 1 µL of each primer (0.5 µM), 7 µL of nuclease-free water

and 1 µL of DNA template. The PCR settings for the SB2F- ITS2.2R and cil18SspannerF-

cil18SspannerR primers were as follows: initial denaturation at 94°C for 5 min, 35 cycles of

denaturation at 94°C for 45 sec, annealing at 55°C for 75 sec, and extension at 72°C for 90 sec,

with a final extension at 72°C for 7 min. PCR setting for LCO1490 and HCO2198 primers were

as follows: initial denaturation at 94 °C for 60 sec, five cycles of denaturation at 94 °C for 60 sec,

annealing at 45 °C for 90 sec, and extension at 68 °C for 90 sec, followed by 40 cycles at 94 °C

for 60 sec, 50 °C for 90 sec, 68 °C for 60 sec, with a final extension at 72 °C for 5 min. PCR

products were sent to Microsynth Seqlab (Germany) for sequencing using the forward primers

SB2F, cil18SspannerF, and LCO1490.

2.7 Genetic distance and phylogenetic analyses

Raw 18S rRNA sequences (infected amphipods collected in April 2023 and ciliate isolated in

March 2024) and a COI sequence (extracted from the mitogenome assembly) were quality-

checked and edited using Geneious Prime v2024.0.5 (Biomatters). The obtained sequences were

separately aligned with apostome ciliate sequences available on GenBank

(www.ncbi.nlm.nih.gov) using the MAFFT v7.490 algorithm with standard settings (Katoh et al.,

2019). All available 18S rRNA sequences with a minimum length of 1500 bp were used for genetic

distance analyses, whereas 18S rRNA and COI sequences with a minimum length of 750 bp were

used for phylogenetic analyses. Genetic distance was calculated in Mega11 v11.0.13 (Tamura et

al., 2021) from 1450 unambiguously aligned nucleotide positions using the Kimura two-parameter

(K2P) model with 1000 bootstrap replicates. Maximum likelihood phylogenetic trees with

bootstrap support values (1000 replicates) were produced in IQ-Tree 2.2.2.6 (Minh et al., 2020).

The HKY+F+I+R2 (18s) and TPM2u+F+I+G4 (COI) substitution models were selected based on

Bayesian information criterion scores. The ciliates *Cinetochilum ovale* (Genbank accession

number FJ870103) and *Paramecium bursaria* (JX082062) were used respectively as outgroups for

18S rRNA and COI sequences. Ciliates and their hosts were identified by comparing obtained

sequences against GenBank records using Blastn.

3. Results

A total of 256 amphipods collected between December 2022 and April 2023 from eight sampling sites were screened for sanguicolous ciliates molecularly and histologically (Table 1). All known amphipod species occurring in the Boye system (Prati et al., 2022) were represented in the sample. More specifically, *G. fossarum* clade 2 (n = 36, 100% similarity, 100% coverage, e-value = 0 to ON093817), *G. pulex* clade C (n = 96, 99.9-100% similarity, 100% coverage, e-value = 0 to ON093813), and clade E (n = 124, 99.9-100% similarity, 100% coverage, e-value = 0 to ON093815). The most common amphipods in the investigated sites were *G. pulex* clade E and C.

3.1 Prevalence

Half of the sampling sites had ciliate-infected amphipod populations, with overall prevalences ranging from 6.67 to 21.43% (Table 1). Among the 12 infected amphipods collected between December 2022 and April 2023, three were *G. fossarum* clade 2, four *G. pulex* clade C, and 5 *G. pulex* clade E. Two ciliate-infected individuals presented simultaneous infections with Microsporidium sp. 515. One of them was also infected with a trematode. The 9 infected amphipods collected in March 2024 and used to isolate fresh ciliates were neither molecularly identified nor screened for co-infections. Overall, the average size of infected amphipods (4th coaxal plate length 3.07 mm ± 0.44 SD, n = 21) was similar to that of uninfected ones (2.93 mm ± 0.49 SD, n = 277). No individual below 2.57 mm was infected.

3.2 Histology and pathology

Histological analyses indicate that the ciliate is a parasite of the hemolymph causing systemic infections. Freely dividing tomonts and protomites-tomites were visible in the host hemocoel, surrounding testis, muscles, the hepatopancreas, gill lamellae, and the outer host cuticle (Figure 1). We found no evidence that the ciliate infiltrated organs such as the hepatopancreas or the testis. However, compared to uninfected individuals, the muscular mass progressively reduced with increasingly high ciliate densities in infected individuals.

3.3 Morphology

Similar to the histological sections, we observed tomonts and protomites-tomites in the glutaraldehyde-fixed gammarids. The tomonts were subelliptical and had a dense ciliature. Their length ranged between 34.6 and 90.8 µm (66.8 µm ± 15.3 SD), and the width between 15.4 and

59.1 µm (32.5 µm ± 8 SD). Tomonts had a large central macronucleus (width 48.5 µm ± 13.2 SD

and length 15.9 µm ± 4.8 SD) and contractile vacuoles either arranged in rows parallel or slightly

diagonal to kineties or rather scattered throughout the cytoplasm (Figures 2 and 3). The largest

tomonts displayed 40 to 47 irregularly spaced kineties converging toward the extremities but

ending shortly before reaching the poles (Figures 2 and 3). However, the vast majority of tomonts

were along the path of regressive division; thus, it is possible that non-regressing cells might attain

a larger size and kineties number than what here reported.

Following palintomy, a process during which the parental cell freely undergoes a sequence

of repeated divisions without intervening growth in the host hemocoel, the size and the number of

kineties of the daughter cells are progressively reduced. Regressing tomonts showed a non-ciliated

band likely resulting from the reabsorption of kineties (Figures 2 and 3). The protomites-tomites

shape varied from pyriform to subelliptical, and their size ranged between 9.9 and 27.8 µm (20.7

± 4.6) in length and 9.3 to 23 µm (14 ± 2.9) in width. They had a large macronucleus (length 11.1

± 3.1 µm and width 6.73 ± 1.8 µm), nine somatic kineties, and in some cases, presented tree buccal

kineties indicated as x, y, and z (Figures 2 and 3).

3.4 Molecular phylogeny and genetics

Molecular identification of the collected sanguicolous ciliates using aligned 18S rRNA sequences

indicated that these were closely related to *Metacollinia luciensis* (MH200618, 95.90% similarity,

100% coverage, e-value = 0), and all belonged to the same haplotype (Figure 4). Similar results

were obtained for the COI region (MH182621, 75.50% similarity, 99% coverage, e-value = 1e-

140); however, the availability of sequences for this region was very limited (Figure 5). Molecular

identification of all ciliate life stages isolated from amphipods collected in March 2024 confirmed

that these belonged to the same species and haplotype. The genetic divergence (K2P) between the

sequences obtained in the current study (0 differing nucleotides) and those of the closest related

species *M. luciensis* (69 differing nucleotides) and *Lynnia grapsolytica* (74 differing nucleotides),

both belonging to the Collinidae family, were respectively 4% and 4.7%, suggesting a closer

genetic relationship with *M. luciensis*. Apostome ciliates belonging to other families diverged by

6.4-6.7% (92-100 differing nucleotides), supporting the placement of the ciliate in the Collinidae.

4. Discussion

This study describes a novel sanguicolous apostome ciliate that infects the amphipods *G. pulex*

and *G. fossarum* from four populations collected in a German stream catchment, the Boye (North

Rhine-Westphalia). Our phylogenetic analyses place *M. emscheri* n. sp. within the Collinidae

family. Due to morphological characteristics, phylogenetic closeness with the type species *M.*

lucensis, and the absence of DNA sequences of less molecularly divergent species, we

provisionally place this ciliate in the genus *Metacollinia.*

4.1 Prevalence of *Metacollinia emscheri* n. sp.

The presence of *M. emscheri* n. sp. in *G. pulex* clade E, *G. pulex* clade C, and *G. fossarum* clade

C2 indicate lower host specificity than other collinids, as these have mostly been reported from

single hosts. The prevalence of *M. emscheri* n. sp. across the four infected amphipod populations

indicates medium infection levels and, in the case of the population collected from the lower stretch

of the Haarbach (HAAun) temporal stability. This sanguicolous ciliate appears to be localized in

the Haarbach and the lower part of the Boye, as none of the amphipods collected upstream of

BOYohKI, which is located just above the junction of the two streams, were infected. However,

we cannot be certain that *M. emscheri* n. sp. is absent from the upper stretch of the Boye catchment.

This ciliate might be present in lower abundances and evaded detection due to a relatively small

sample size (~ 30 individuals). Furthermore, it is possible that seasonal dynamics may play a role.

Accordingly, the prevalence of *P. branchiarum* in *G. pulex* collected in a locality in proximity to

Prague varied between 1% in January and 20% in August (De Puytorac and Lom, 1962). The close

proximity of BOYuhHA, the most downstream site, with the River Emscher (2.7 km), suggests

that *M. emscheri* n. sp. might also be present in the latter.

Assuming that the observed replacement of the muscle mass in infected amphipods may

hinder the host's ability to withstand water flow, it is plausible that a good portion of infected

individuals passively drift downstream and end up in the River Emscher. Furthermore, all infected

amphipods had relatively large sizes, which alone might influence amphipod drift. Accordingly,

both large-sized and microsporidian-infected *G. pulex* in a nearby watercourse, the Rotbach,

drifted more than smaller and uninfected individuals (Prati et al., 2023). The exclusive presence

of *M. emscheri* n. sp. in relatively large-sized amphipods is not surprising. Generally, parasite

prevalence in aquatic organisms, including amphipods, is positively correlated with host size, as

older individuals are more likely to be exposed to infections over their lifespan (Poulin, 2000; Prati

et al., 2023; Soldánová et al., 2010). Furthermore, size-asymmetric cannibalism, which is common

in many aquatic taxa, including *G. pulex*, may potentially favor the transmission of parasites via

ingestion of infected individuals (MacNeil et al., 2003; McGrath et al., 2007).

4.2 Transmission of *Metacollinia emscheri* n. sp.

The transmission of *M*. *emscheri* n. sp., like that of other members of the Collinidae and

Pseudocollinidae family, remains unknown (Chantangsi et al., 2013; Gómez-Gutiérrez et al., 2012;

Lynn, 2008; Metz and Hechinger, 2021). Thus, we don´t know if the parasite can be orally

transmitted. Poisson (1921) described *M. luciensis* from the digestive tract of *Orchestia*

gammarellus but mentioned that the ciliate might be able to pierce through the intestine wall and

penetrate the coelom, where it can also be found. On the other hand, Balbiani (in Cépède, 1910)

hypothesized that host injury is required for ciliate transmission. Such a pathway might also be

possible for *G. pulex* as injuries often occur in adult individuals, particularly during mating, when

males compete to assert their dominance over females (Dick and Elwood, 1990).

Alternatively, preexisting parasitic infections might favor secondary infections by

hindering the host immune system (Cox, 2001). This does not seem to be the case in the present

study, as the vast majority of infected amphipods did not show coinfections. A more plausible

theory has been proposed by Metz and Hechinger (2021). After failing multiple attempts to

experimentally infect healthy and injured crabs with *L. grapsolytica* by exposure to swimming

tomites, infected conspecifics, or direct injection of life cells, they proposed that the parasite has a

complex life cycle with development outside the host, possibly involving an intermediate host.

Such a possibility might also apply to *M. emscheri* n. sp.; however, more extensive studies are

required to shed light on its life cycle.

4.3 Ecological importance of *Metacollinia emscheri* n. sp.

Irrespective of its elusive life cycle, *M. emscheri* n. sp. is of high ecological importance. With

prevalences up to 21% in the infected populations, this parasite is likely to strongly affect host

population dynamics. Due to reduced muscular mass, host movement and activities such as feeding

and mating might be affected in individuals with high infection levels. As amphipods are keystone

species in freshwater aquatic environments, *M. emscheri* n. sp. may indirectly impact ecosystem

functioning. For instance, infected individuals may experience reduced shredding activity and,

consequently, a slowdown of leaf-litter breakdown decreases the amount of nutrients and
processed material available for microorganisms and other invertebrates. Furthermore, if *M.*
emscheri n. sp. directly or indirectly increases the mortality rate in amphipods, species that rely on
such prey may face reduced food availability.

4.4 Taxonomy and nomenclature of the family Collinidae

Jankowski (2007) undertook a comprehensive revision of the nomenclature and taxonomy of
Collinidae, suggesting that the genus *Collinia* should be split into three distinct genera: *Collinia*,
Paracollinia, and *Metacollinia*. The first, with the type species *Collinia circulans*, was originally
observed by Balbiani 1885 (in Cépède, 1910) in the freshwater isopod *Asellus aquaticus*. *Collinia*
circulans is characterized by a large tomont (70-80 µm long and 30-35 µm wide) with ~ 13 spiraled
bipolar kineties and a single row of ~ 12 contractile vacuoles. The tomite is small, showing short
x, y, and z kineties reminiscent of those of the foettingeriids (De Puytorac and Lom, 1962; Lynn,
2008; Lynn and Strüder-Kypke, 2019). The genus *Paracollinia*, whose type species *Paracollinia*
branchiarum was observed in 1852 by Stein infecting the freshwater amphipod *G. pulex* has large
tomonts with 55 to 60 straight evenly spaced kineties without a non-ciliated cortical band and a
smaller flattened tomite with 9 kineties and short x, y, and z kineties, reminiscent of those of the
foettingeriids (Jankowski, 2007; Lynn and Strüder-Kypke, 2019). The genus *Metacollinia*, whose
type species *Metacollinia luciensis* was observed in the marine amphipod *O. gammarellus* by
Poisson (1921), has large tomonts with 65 to 70 slightly spiraling kineties, a large non-ciliated
cortical band and two lateral rows of 8-10 contractile vacuoles. Like the previous species, the
tomite has 9 kineties and short x, y, and z kineties, reminiscent of those of the foettingeriids
(Jankowski, 2007; Lynn and Strüder-Kypke, 2019). More recently, a novel genus, *Lynnia*, with
the type species *L. grapsolytica* found in the marine crab *Pachygrapsus crassipes* has been
included in the Collinidae (Metz and Hechinger, 2021). The tomont is large with ~ 11 equidistant
spiraled bipolar kineties, lacks a non-ciliated cortical band, and possesses randomly scattered
subpellicular contractile vacuoles. The tomite is small, showing short x, y, and z kineties
reminiscent of those of the foettingeriids; however, unique compared to any known apostome, the
eighth and ninth kineties are both interrupted, and once impregnated with silver, the tomite
presents a hook-shaped arrangement of ventral somatic kineties (Metz and Hechinger, 2021).

4.5 Differences between *M. emscheri* n. sp. and other Collinids

Compared to *L. grapsolytica*, the tomite of *M. emscheri* n. sp. bears more resemblance with that of *C. ciruclans*, *P. branchiarum,* and *M. luciensis*. Only the ninth somatic kinety in *M. emscheri* n. sp. is interrupted, and no hook-shaped arrangement of ventral somatic kineties is present. Like *M. luciensis*, the tomont of *M. emscheri* n. sp. displayed slightly spiraled kineties and a non-ciliated cortical band, suggesting the placement of the ciliate in this genus. The average length of tomont in *M. emscheri* n. sp. (~67 µm) was similar to that reported in *M. luciensis* (~70 µm) by De Puytorac and Grain (1975). However, the kineties, numbering 40-47, were lower than the 65-70 reported in *M. luciensis.* It is possible that the smaller size and lower number of kineties observed in *M. emscheri* n. sp. might be due to regression processes occurring during the tomitogenesis, as reported for *M. luciensis* collected by Lynn and Strüder-Kypke (2019). Interestingly, the contractile vacuoles in the tomont of *M. emscheri* n. sp., which numbers 6-10 in contrast to 8-10 found in *M. luciensis* (De Puytorac and Grain, 1975)*,* are arranged either in two parallel rows or scattered across the cytoplasm.

Phylogenetic analyses indicate that *M. emscheri* n. sp. is more closely related to *M. lucensis* than *L. grapsolytica*, the only other Collinidae for which DNA sequences are available. In the absence of DNA sequences of species belonging to the genera *Collinia* and *Paracollinia*, and based on morphological similarity with *M. lucensis* we provisionally place *M. emscheri* n. sp. in the genus *Metacollina*. Given the gap existing in DNA sequence references, we cannot exclude that the proposed placement might change once more sequences of Collinidae species become available.

4.6 Taxonomic summary

Kingdom: Protozoa

Superphylum: Alveolata

Phylum: Ciliophora Doflein, 1901

Subphylum: Intramacronucleata Lynn, 1996

Class: Oligohymenophorea de Puytorac et al. (1974)

Order: Apostomatida Chatton and Lwoff, 1928

Family: Colliniidae Cepede, 1910

Species: *Metacollinia emscheri* n. sp. Prati, Rückert, Grabner, Sures, Bojko, 2024

Formal taxonomic summary: The tomonts are ~67 µm long with 40-47 (possibly more) slightly spiraled kineties, a non-ciliated cortical band, and 6-10 contractile vacuoles arranged either in two parallel rows or scattered across the cytoplasm. At the end of palintomy, during which the number of kineties decreases to nine, oral fragments (buccal kineties x, y, and z) appear.

Type locality: Boye catchment, North-Rhine Westphalia, Germany (Boye 51.553107; 6.948059 and 51.534515; 6.997740, Haarbach 51.570306; 6.960891 and 51.562638; 6.955581)

Type host: *Gammarus pulex* Linnaeus, 1758 (Crustacea, Malacostraca, Amphipoda, Gammaridae)

Type habitat: Endoparasitic, found in the hemolymph and muscle tissues of hosts collected in lowland streams at the type locality.

Holotype: The name-bearing type of this species is the specimen illustrated in Figures 2 and 3 (ICZN 1999, Articles 73.1.4).

Etymology: The species epithet emscheri refers to the Emscher River catchment, which comprises the type localities found in its tributary, the Boye and the Haarbach.

Authors contribution

Sebastian Prati, Daniel S. Grabner, Bernd Sures, and Jamie Bojko conceived the study. Sebastian Prati carried out the sampling. Sebastian Prati, Sonja Rückert, and Jamie Bojko performed laboratory analyses. Jamie Bojko supervised the study. Sebastian Prati wrote the first draft of the manuscript, and all authors contributed critically to the final draft. All authors read and approved the final manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This study was performed within the Collaborative Research Center (CRC) RESIST (A09) and funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - CRC 1439/1 - project number: 426547801. We are grateful to the Developmental Biology group of the University of Duisburg-Essen and, in particular, to Sabine Schneider for providing access and assistance with the histological instruments. We thank the Imaging Center Essen (IMCES) at the Faculty of Medicine of the University of Duisburg-Essen, in particular Bernd Walkenfort for providing access and assistance with the SEM equipment.

Appendix A. Supplementary data

Supplementary material 1

References

Adlard, R.D., Lester, R.J.G., 1995. Development of a diagnostic test for *Mikrocytos roughleyi*, the aetiological agent of Australian winter mortality of the commercial rock oyster, *Saccostrea commercialis* (Iredale & Roughley). J. Fish Dis. 18, 609–614. https://doi.org/10.1111/j.1365-2761.1995.tb00365.x

Bojko, J., Ovcharenko, M., 2019. Pathogens and other symbionts of the Amphipoda: taxonomic diversity and pathological significance. Dis. Aquat. Org. 136, 3–36. https://doi.org/10.3354/dao03321

Capriulo, G., Small, E., 1985. Discovery of an apostome ciliate (*Collinia beringensis* n. sp.) endoparasitic in the Bering Sea euphausiid *Thysanoessa inermis*. Dis. Aquat. Org. 1, 141–146. https://doi.org/10.3354/dao001141

Cépède, C., 1910. Recherches sur les infusoires astomes: anatomie, biologie, ethologie parasitaire, systeématique. Librairie A. Schulz, Paris.

Chantangsi, C., Lynn, D.H., Rueckert, S., Prokopowicz, A.J., Panha, S., Leander, B.S., 2013. *Fusiforma themisticola* n. gen., n. sp., a new genus and species of apostome ciliate infecting the Hyperiid amphipod *Themisto libellula* in the Canadian Beaufort Sea (Arctic Ocean), and establishment of the Pseudocolliniidae (Ciliophora, Apostomatia). Protist 164, 793–810. https://doi.org/10.1016/j.protis.2013.09.001

Cox, F.E.G., 2001. Concomitant infections, parasites and immune responses. Parasitology 122,
S23–S38. https://doi.org/10.1017/S003118200001698X

De Puytorac, P., Grain, J., 1975. Etude de la tomitogenèse et de l'ultrastructure de *Collinia*
orchestiae, cilié apostome sanguicole, endoparasite du crustacé *Orchestia gammarella*
pallas. Protistologica 11, 61–74.

De Puytorac, P., Lom, J., 1962. La tomitogénèse des *Collinia* ciliés apostomes sanguicoles
endoparasites des crustacés. Ann. Parasitol. Hum. Comp. 37, 195–209.
https://doi.org/10.1051/parasite/1962373195

Dick, J.T.A., Elwood, R.W., 1990. Symmetrical assessment of female quality by male
Gammarus pulex (Amphipoda) during struggles over precopula females. Anim. Behav.
40, 877–883. https://doi.org/10.1016/S0003-3472(05)80989-3

Giari, L., Fano, E.A., Castaldelli, G., Grabner, D., Sures, B., 2020. The ecological importance of
amphipod–parasite associations for aquatic ecosystems. Water 12, 2429.
https://doi.org/10.3390/w12092429

Gómez-Gutiérrez, J., Kawaguchi, S., 2017. *Pseudocollinia* histophagous ciliates infect krill in
the Pacific and Atlantic Oceans and possibly worldwide. CICIMAR Oceánides 32, 15–
24. https://doi.org/10.37543/oceanides.v32i2.204

Gómez-Gutiérrez, J., Strüder-Kypke, M.C., Lynn, D.H., Shaw, T.C., Aguilar-Méndez, M.J.,
López-Cortés, A., Martínez-Gómez, S., Robinson, C.J., 2012. *Pseudocollinia brintoni*
gen. nov., sp. nov. (Apostomatida: Colliniidae), a parasitoid ciliate infecting the
euphausiid *Nyctiphanes simplex*. Dis. Aquat. Org. 99, 57–78.
https://doi.org/10.3354/dao02450

Grabner, D.S., Weigand, A.M., Leese, F., Winking, C., Hering, D., Tollrian, R., Sures, B., 2015.
Invaders, natives and their enemies: distribution patterns of amphipods and their
microsporidian parasites in the Ruhr Metropolis, Germany. Parasit. Vectors 8, 419.
https://doi.org/10.1186/s13071-015-1036-6

Jankowski, A., 2007. Phylum Ciliophora Doflein, 1901, in: Alimov, A. F. (Ed.), Protista, Part 2,
Handbook of Zoology. Russian Academy of Science, Zoological Institute, St. Petersburg,
pp. 415–993.

Ji, D., Wang, Y., 2018. An optimized protocol of protargol staining for ciliated protozoa. J.
Eukaryot. Microbiol. 65, 705–708. https://doi.org/10.1111/jeu.12515

Katoh, K., Rozewicki, J., Yamada, K.D., 2019. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief. Bioinformatics. 20,
1160–1166. https://doi.org/10.1093/bib/bbx108

Kurilov, A.V., 2017. Improvement of silver impregnation technique using in situ synthesized
protargol. Acta Protozool. 56.

Lynn, D.H. (Ed.), 2008. The ciliated protozoa: characterization, classification, and guide to the
literature., 3rd ed. Springer Netherlands, Dordrecht. https://doi.org/10.1007/978-1-4020-
8239-9_17

Lynn, D.H., Gómez-Gutiérrez, J., Strüder-Kypke, M.C., Shaw, C.T., 2014. Ciliate species
diversity and host-parasitoid codiversification in *Pseudocollinia* infecting krill, with
description of *Pseudocollinia similis* sp. nov. Dis. Aquat. Org. 112, 89–102.
https://doi.org/10.3354/dao02796

Lynn, D.H., Strüder-Kypke, M.C., 2019. The sanguicolous apostome *Metacollinia luciensis*
Jankowski 1980 (Colliniidae, Apostomatia, Ciliophora) is not closely related to other
sanguicolous apostomes. J. Eukaryot. Microbiol. 66, 140–146.
https://doi.org/10.1111/jeu.12638

Macneil, C., Dick, J.T.A., Elwood, R.W., 1997. The trophic ecology of freshwater *Gammarus*
spp. (Crustacea: Amphipoda): problems and perspectives concerning the functional
feeding group concept. Biol. Rev. Cambridge Philos. Soc. 72, 349–364.
https://doi.org/10.1017/S0006323196005038

MacNeil, C., Dick, J.T.A., Hatcher, M.J., Fielding, N.J., Hume, K.D., Dunn, A.M., 2003.
Parasite transmission and cannibalism in an amphipod (Crustacea). Int. J. Parasitol. 33,
795–798. https://doi.org/10.1016/S0020-7519(03)00110-3

McGrath, K.E., Peeters, E.T.H.M., Beijer, J.A.J., Scheffer, M., 2007. Habitat-mediated
cannibalism and microhabitat restriction in the stream invertebrate *Gammarus pulex*.
Hydrobiologia 589, 155–164. https://doi.org/10.1007/s10750-007-0731-5

Metz, D.C.G., Hechinger, R.F., 2021. *Lynnia grapsolytica* n. gen, n. sp. (Ciliophora:
Apostomatida: Colliniidae), a deadly blood parasite of crabs with a novel
pseudocytopharynx. J. Eukaryot. Microbiol. 68, e12847.
https://doi.org/10.1111/jeu.12847

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,
Lanfear, R., 2020. IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. Mol. Biol. Evol. 37, 1530–1534.
https://doi.org/10.1093/molbev/msaa015

Poisson, R., 1921. Sur un infusoire du genre *Balantidium*, parasite du tube digestif d'*Orchestia*
littorea Mont. Compt. Rend. Seances Soc. Biol. Fil 84, 333–335.

Poulin, R., 2000. Variation in the intraspecific relationship between fish length and intensity of
parasitic infection: biological and statistical causes. J. Fish Biol. 56, 123–137.
https://doi.org/10.1111/j.1095-8649.2000.tb02090.x

Prati, S., Enß, J., Grabner, D.S., Huesken, A., Feld, C.K., Doliwa, A., Sures, B., 2023. Possible
seasonal and diurnal modulation of *Gammarus pulex* (Crustacea, Amphipoda) drift by
microsporidian parasites. Sci Rep 13, 9474. https://doi.org/10.1038/s41598-023-36630-2

Prati, S., Grabner, D.S., Pfeifer, S.M., Lorenz, A.W., Sures, B., 2022. Generalist parasites persist
in degraded environments: a lesson learned from microsporidian diversity in amphipods.
Parasitology 149, 973–982. https://doi.org/10.1017/S0031182022000452

Prokopowicz, A.J., Rueckert, S., Leander, B.S., Michaud, J., Fortier, L., 2010. Parasitic infection
of the hyperiid amphipod *Themisto libellula* in the Canadian Beaufort Sea (Arctic
Ocean), with a description of *Ganymedes themistos* sp. n. (Apicomplexa,
Eugregarinorida). Polar Biol 33, 1339–1350. https://doi.org/10.1007/s00300-010-0821-0

Skibbe, O., 1994. An improved quantitative protargol stain for ciliates and other planktonic
protists. Arch. Hydrobiol. 339–347. https://doi.org/10.1127/archiv-
hydrobiol/130/1994/339

Soldánová, M., Selbach, C., Sures, B., Kostadinova, A., Pérez-del-Olmo, A., 2010. Larval
trematode communities in *Radix auricularia* and *Lymnaea stagnalis* in a reservoir system
of the Ruhr River. Parasit Vectors 3, 56. https://doi.org/10.1186/1756-3305-3-56

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Mol. Biol. Evol. 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

Table 1 Sampling locations, sampling date, amphipod species, average size of the 4[th] coaxal plate

in mm with SD, and prevalence in % of *Metacollinia emscheri* n. sp.

Location	Coordinates	Date	N	Species	Size in mm	Prevalence in %
BOYohBR	51.569135; 6.927698	27/04/2023	4	*Gammarus pulex* clade C	2.88 ± 0.21	0
			26	*Gammarus pulex* clade E	2.95 ± 0.46	0
BOYohKI	51.553107; 6.948059	07/12/2022	8	*Gammarus pulex* clade C	2.57 ± 0.31	12.50 (1/8)
			4	*Gammarus pulex* clade E	2.49 ± 0.37	0
		13/05/2023	18	*Gammarus pulex* clade C	2.63 ± 0.43	0
			12	*Gammarus pulex* clade E	2.71 ± 0.41	0
BOYohSP	51.564297; 6.930277	27/04/2023	10	*Gammarus fossarum* clade 2	2.82 ± 0.56	0
			8	*Gammarus pulex* clade C	2.99 ± 0.74	0
			12	*Gammarus pulex* clade E	2.96 ± 0.57	0
BOYuhHA	51.534515; 6.997740	02/05/2023	26	*Gammarus fossarum* clade 2	3.08 ± 0.51	11.53 (3/26)
			4	*Gammarus pulex* clade C	3.28 ± 0.66	0
BRA1000	51.577238; 6.932752	27/04/2023	1	*Gammarus pulex* clade C	3.36	0
			25	*Gammarus pulex* clade E	2.95 ± 0.38	0
HAAob	51.570306; 6.960891	13/05/2023	11	*Gammarus pulex* clade C	2.87 ± 0.30	9.09 (1/11)
			19	*Gammarus pulex* clade E	2.81 ± 0.36	5.26 (1/19)
HAAun	51.562638; 6.955581	13/05/2023	9	*Gammarus pulex* clade C	3.02 ± 0.31	22.22 (2/9)
			21	*Gammarus pulex* clade E	2.92 ± 0.49	19.05 (4/21)
		02/03/2024	42	Not determined	3.07 ± 0.47	21.43 (9/42)
VORohBOY	51.554753; 6.932435	07/12/2022	8	*Gammarus pulex* clade C	2.84 ± 0.29	0
		13/05/2023	25	*Gammarus pulex* clade C	3.15 ± 0.63	0
			5	*Gammarus pulex* clade E	3.07 ± 0.71	0

List of Figures

Figure 1 Histopathological preparations of the ciliate infection in amphipod tissue. A) Ciliated protozoans (white triangle) with basophilic staining nuclei (black triangle) are present in the hemocoel of the infected amphipod, alongside hemocytes (orange arrow). The image includes hemocoel ('Heam') surrounding the hepatopancreas ('HP') and muscle. B) Ciliated protozoans (white triangle) passing through the gill lamellae ('Gill'), alongside the outer host cuticle. C) Ciliate-like features are observable (white arrow) on the outside of the protozoans. D) Ciliated protozoans (white triangle) seen to travel through the hemocoel ('Heam') adjacent to the testis, gut, and hepatopancreas ('HP'). An individual in section highlights the common U-shaped nucleus often noted in Ciliophora members.

Figure 2 A) tomont with 47 kineties, B) internal morphology of tomont showing macronucleus and contractile vacuoles, C) regressing tomont displaying a non-ciliated cortical band, D) dividing protomite showing 9 kineties and the x, y, and z kineties, and E) conjugant.

Figure 3 A) dividing tomont, B) regressing tomont displaying a non-ciliated cortical band, C) tomont showing kineties merging toward the poles but not touching them, D) dividing tomite and conjugant, E) light microscopy of freshly isolated tomont with kineties and a large macronucleous visible, and F) QPS staining of tomont showing contractile vacuoles arranged in parallel lines and a large macronucleous.

Figure 4 Maximum-likelihood phylogeny of Apostome ciliates incorporating 18S rDNA sequences. Gray dots indicate bootstrap support values above 90 (1,000 replicates).

Figure 5 Maximum-likelihood phylogeny of Apostome ciliates incorporating COI sequences. Gray dots indicate bootstrap support values above 90 (1,000 replicates).

561