Metacollinia emscheri **n. sp., a novel sanguicolous apostome ciliate of freshwater**

amphipods (*Gammarus* spp.)

Sebastian Prati[1,2]*, Sonja Rückert[2,3], Daniel S. Grabner[1,2], Bernd Sures[1,2,4], Jamie Bojko[5,6]

[1]Aquatic Ecology, University of Duisburg-Essen, Universitaetsstr. 5, 45141 Essen, Germany.

[2]Centre for Water and Environmental Research, University of Duisburg-Essen, Universitaetsstr.

5, 45141 Essen, Germany. [3]Eukaryotic Microbiology, University of Duisburg-Essen,

Universitaetsstr. 5, 45141 Essen, Germany. [4]Research Center One Health Ruhr of the University

Alliance Ruhr, University of Duisburg-Essen, Universitätsstraße 5, 45141 Essen, Germany.

[5]National Horizons Centre, Teesside University, Darlington, DL1 1HG, United Kingdom. [6]School

of Health and Life Sciences, Teesside University, Middlesbrough, TS1 3BX, United Kingdom

*Corresponding author: sebastian.prati@uni-due.de

Keywords: Collinidae, endoparasitic ciliate, amphipod pathogen, crustacea, mitogenome

Abstract

We describe a novel sanguicolous parasitic ciliate, *Metacollinia emscheri* n. sp., found in the

freshwater amphipods *Gammarus pulex* and *G. fossarum*. This ciliate infected up to 20% of the

amphipods collected in a German stream catchment, the Boye, a tributary of the river Emscher.

The ciliate showed morphological characteristics fitting the genus *Metacollinia*. Different life

stages of variable size occurred simultaneously in the hemocoel throughout the hosts' body. The

tomont had 40-47 slightly spiraled kineties, a non-ciliated cortical band, a large macronucleus, and

contractile vacuoles arranged in rows or scattered throughout the cytoplasm. The

protomites/tomites with nine somatic kineties presented evidence of the buccal kineties x, y, and

z reminiscent of those of the order Foettingeriida. Phylogenetic analyses of the 18S rRNA and COI

regions confirm the ciliate placement in the Collinidae and a close relatedness to the type species

of the genus *Metacollinia*, *Metacollinia lucensis*.

We formally describe this new parasite as *Metacollinia emscheri* n. sp. using pathological,

morphological, and nuclear/mitochondrial genetic data. The systemic infections observed in

histological preparations and the pathogenicity of *Metacollinia emscheri* n. sp. suggest that this

parasite might influence host population dynamics. Given the ecological importance of amphipods

as keystone species in freshwater ecosystems, an outbreak of this parasite might indirectly impact
ecosystem functioning.

1. Introduction

Macroinvertebrates, including amphipods, are essential components of stream ecosystem food webs. Amphipods are highly diversified and ubiquitous keystone organisms that often represent the dominant element of invertebrate communities (Giari et al., 2020; Macneil et al., 1997). Their shredding activity can account for up to 75% of overall leaf-litter breakdown in some ecosystems, and they are an important food source for a wide range of predators, linking several trophic levels within riverine food webs (Giari et al., 2020). Their ecological niche makes them ideal hosts for a hyper-diverse range of pathogens ranging from viruses to trophically transmitted metazoans (Bojko and Ovcharenko, 2019; Warren et al., 2023, 2022). Yet most studies focus on metazoan parasites, gregarines (McKinley et al. 2024), or microsporidians due to their abundance, diversity, and ecological relevance (Prati et al., 2023).

Protists, such as ciliates, are commonly found as commensal epibionts on the carapace, gills, and extremities of amphipods, where they benefit from the host movement to collect particulate matter (Bojko et al., 2013). However, few apostome ciliates have specialized symbiotic and parasitic relationships with crustaceans, including amphipods (Lynn, 2008). Among them, sanguicolous parasites are able to penetrate the host cuticle and feed on the cells and fluid of the hemocoel (Lynn et al., 2014; Lynn and Strüder-Kypke, 2019). Representatives of the Collinidae and the Pseudocollinidae families reproduce continuously in the host hemolymph, leading to systemic infections, which might ultimately result in host death (Chantangsi et al., 2013; Lynn et al., 2014; Lynn and Strüder-Kypke, 2019; Prokopowicz et al., 2010). For instance, Pseudocollinidae representatives have been linked to mass mortality events in krill in marine environments (Gómez-Gutiérrez et al., 2012; Gómez-Gutiérrez and Kawaguchi, 2017) and the Collinidae member *Lynnia grapsolytica* strongly enhances the mortality rate of infected crabs under laboratory conditions (Metz and Hechinger, 2021).

The majority of Collinidae are parasites of freshwater amphipods and isopods (De Puytorac and Lom, 1962). One of the first observations in freshwater amphipods dates back to 1852 when an apostome ciliate, later described as *Paracollinia branchiarum* was found in the hemolymph of *Gammarus pulex* (De Puytorac and Lom, 1962). Successive investigations revealed the presence

of Collinidae in freshwater amphipods from Lake Baikal and subterranean environments in Europe

and the USA (Capriulo and Small, 1985). More recently, an unidentified ciliate has been

histologically observed in the hemolymph of *Gammarus roeselii* (Bojko et al., 2017). These

findings indicate that these parasites might be ubiquitous but poorly investigated.

Given the ecological importance of amphipods and the possible high pathogenicity of

ciliates belonging to the Collinidae family, deepening our knowledge of these sanguicolous

parasites is paramount. If highly pathogenic, these parasites might strongly influence host

population dynamics and indirectly affect ecosystem functioning. We document the pathology,

morphological characteristics, and genetic detail of a novel member of the Collinidae, in the

hemocoel of the amphipods *Gammarus pulex* and *Gammarus fossarum,* and formally describe it

as *Metacollinia emscheri* n. sp.

74 2. Materials and methods

75 2.1 Sampling

Amphipods were collected between December 2022 and March 2024 from eight sampling

locations across the Boye catchment in North Rhine-Westphalia, Germany (Table 1). The first

collection of amphipods (December 2022) initially aimed at describing novel microsporidian

parasites comprised 72 individuals (n = 52 from BOYohKI and n = 20 from VORohBOY, Table

1). Thus, only microsporidian-infected amphipods (n = 20; for molecular identification method,

see Prati et al. 2023) were processed for histological analyses. After finding an individual infected

with an unknown sanguicolous ciliate, we decided to carry out a non-targeted sampling in April

2023.

All amphipods were kept alive and dissected within 6h post collection. The head and

anterior part of the thorax were fixed in 96% ethanol for molecular analyses. The proximate

segments of the thorax (2 mm) were placed in 2.5% glutaraldehyde 0.1 M sodium cacodylate

buffer (pH 7.4) for electron microscopy. The rest of the body was fixed in Davidson´s freshwater

fixative for 24 hours, then transferred to 70% ethanol for histological examination. Additionally,

given the high morphological variability observed in the ciliates, we collected 42 amphipods in

March 2024. We dissected them and separately stored the freshly isolated ciliates by life stages in

96% ethanol for molecular analyses to confirm their taxonomic placement.

2.2 Histological analyses

Amphipod tissues were infiltrated with paraffin wax in an automated tissue processor (ethanol-
xylene substitute-wax), embedded into wax blocks, and sectioned to 4 µm thickness through the
animal center. Each section was mounted on a glass slide and stained with hematoxylin and eosin,
following a standard de-wax, stain, and rehydration protocol. The slides were screened for
pathogens using an Olympus CX 33 trinocular light microscope and photographed with a Sony
Alpha 6600 mounted via extension tubes on the camera port.

2.3 Scanning Electron Microscopy

Haemocoel from glutaraldehyde-fixed or fresh amphipods was transferred into self-made filter
baskets with a 10 μm polycarbonate membrane filter (Millipore Corp.) attached to a cut-off
BEEM® conical tip capsule. These capsules were placed in 0.9% NaCl in a multi-well plate. A
piece of filter paper was mounted on the inside of the multi-well plate lid and saturated with 4%
(w/v) OsO_4. The ciliates were fixed with OsO_4 vapor for 30 min in the dark. Ten drops of 4% (w/v)
OsO_4 were added directly to the filter baskets, and the samples were fixed again for 30 min. The
filters with the ciliates were washed with distilled water, dehydrated with a graded series of ethyl
alcohol, and critical point dried with CO_2 (Leica EM CPD 300 critical point drier). Filters were
mounted on stubs, sputter coated with 5 nm platinum (Leica ACE 600 sputter coater), and viewed
using the Zeiss Crossbeam 540 scanning electron microscope.

2.4 Quantitative protargol staining

We employed a rapid Quantitative protargol staining (QPS) on ciliates isolated from
glutaraldehyde-fixed amphipods to better observe the internal structures of ciliates. The protocol
(see details in Supplementary material 1) is a modified version of those proposed by Skibbe (1994),
Kurilov (2017), and Ji and Wang (2018). Briefly, the steps employed involved in-house production
of liquid protargol (Kurilov, 2017), the concentration of fixed ciliates on cellulose filters (Skibbe,
1994), coverage of ciliates with a thin layer of agarose, hardening of agarose with formaldehyde,
fast impregnation in warm protargol solution (Skibbe, 1994), dehydration and mounting of the
filter on permanent slides. The process included steps to correct overbleaching and overstaining
(Ji and Wang, 2018).

2.5 Molecular analyses

DNA was isolated from muscular tissue following a modified salt precipitation protocol described by Grabner et al. (2015). DNA extracted from the only amphipod infected with an unknown ciliate in December 2022 was sequenced using shotgun next-generation sequencing (NGS) to assemble the ciliate mitogenome. Based on the obtained sequences, we then conducted a molecular screening of sanguicolous ciliates in amphipods collected in April 2023 using DNA barcoding (Sanger sequencing). DNA barcoding was also used to confirm that all ciliate life stages isolated in March 2024 belonged to the same species. Furthermore, we molecularly identified infected hosts using DNA barcodes.

2.6 Genomic library preparation, sequencing, and bioinformatics

A DNA sample from an animal with histological confirmation of the infection was used for shotgun next-generation sequencing (NGS). The DNA extract was submitted to Novogene, where a library was prepared (NEBNext® Ultra™ DNA Library Preparation Kit), which results in paired-end (150 bp) sequence data. The library was sequenced on an Illumina NovaSeq. Data were returned in fastq format (raw forward reads: 34.6 M; raw reverse reads: 34.6 M) and trimmed and quality checked using Trimmomatic (Bolger et al., 2014). Assembly of the data was conducted in SPAdes v.3.15.3 (Bankevich et al., 2012), resulting in a contiguous sequence file containing 107,638 contigs (>1000 bp) and assembly statistics of: N50: 830; N90: 545; L50: 181,716; L90: 460,369; determined using quast v.5.2.0 (Gurevich et al., 2013).

The contigs were screened for mitochondrial sequences using a combination of metaxa2 v.2.1.3 (Bengtsson-Palme et al., 2015) and blastp (based on a bespoke protein database developed from available Ciliophora mitogenomes). This isolated two contigs that contributed to the genome of the suspected ciliate observed in histological section. These contigs were mapped to confirm contiguity (using CLC v.12) and annotated using Mfannot (Lang et al., 2023; translation table 4 - mold and protozoan).

2.7 DNA barcoding

For the molecular screening of sanguicolous ciliates in amphipods collected in April 2023 and for the ciliates isolated in March 2024, we used the primer pair: SB2F (5-TGTAAAACGACGGCCAGTGTTCCCCTTGAACGAGGAATTC-3) and ITS2.2R (5-

CAGGAAACAGCTATGACCCTGGTTAGTTTCTTTTCCTCCGC-3) in conjunction with the

pair cil18SspannerF (5-TGTAAAACGACGGCCAGT-AATCAGAGTAATGATTAATAGG-3)

and cil18SspannerR (5-CAGGAAACAGCTATGAC-TGGTTCACCGGACCACTCGA-3);

which targets the 18S rRNA and LSU rRNA region (Adlard and Lester, 1995; Metz and Hechinger,

2021) as these primers bound well to the mitogenome sequences obtained from the ciliate collected

in December 2022 and did not amplify amphipods DNA. All infected amphipods were identified

using the universal eukaryotic primers LCO1490 and HCO2198, which amplify the COI region.

Each PCR reaction volume used for the SB2F-ITS2.2R and cil18SspannerF-

cil18SspannerR primer pairs consisted of 20 μL composed of 10 μL of 2 × AccuStart II PCR

ToughMix (Quantabio), 1 μL of each primer (0.5 μM), 0.35 μL of 50 × GelTrack Loading Dye

(Quantabio), 6.65 μL MilliQ water and 1 μL of DNA template. For the LCO1490 and HCO2198

primer pair, the PCR reaction consisted of 20 μL assay with 10 μL of OneTaq Quick-Load 2x

Master mix (New England Biolabs), 1 μL of each primer (0.5 μM), 7 μL of nuclease-free water

and 1 μL of DNA template. The PCR settings for the SB2F- ITS2.2R and cil18SspannerF-

cil18SspannerR primers were as follows: initial denaturation at 94°C for 5 min, 35 cycles of

denaturation at 94°C for 45 sec, annealing at 55°C for 75 sec, and extension at 72°C for 90 sec,

with a final extension at 72°C for 7 min. PCR setting for LCO1490 and HCO2198 primers were

as follows: initial denaturation at 94 °C for 60 sec, five cycles of denaturation at 94 °C for 60 sec,

annealing at 45 °C for 90 sec, and extension at 68 °C for 90 sec, followed by 40 cycles at 94 °C

for 60 sec, 50 °C for 90 sec, 68 °C for 60 sec, with a final extension at 72 °C for 5 min. PCR

products were sent to Microsynth Seqlab (Germany) for sequencing using the forward primers

SB2F, cil18SspannerF, and LCO1490.

2.8 Genetic distance and phylogenetic analyses

Raw 18S rRNA sequences (infected amphipods collected in April 2023 and ciliate isolated in

March 2024) and a COI sequence (extracted from the mitogenome assembly) were quality-

checked and edited using Geneious Prime v2024.0.5 (Biomatters). The obtained sequences were

separately aligned with apostome ciliate sequences available on GenBank

(www.ncbi.nlm.nih.gov) using the MAFFT v7.490 algorithm with standard settings (Katoh et al.,

2019). All available 18S rRNA sequences with a minimum length of 1500 bp were used for genetic

distance analyses, whereas 18S rRNA and COI sequences with a minimum length of 750 bp were

used for phylogenetic analyses. Genetic distance was calculated in Mega11 v11.0.13 (Tamura et

al., 2021) from 1450 unambiguously aligned nucleotide positions using the Kimura two-parameter

(K2P) model with 1000 bootstrap replicates. Maximum likelihood phylogenetic trees with

bootstrap support values (1000 replicates) were produced in IQ-Tree 2.2.2.6 (Minh et al., 2020).

The HKY+F+I+R2 (18s) and TPM2u+F+I+G4 (COI) substitution models were selected based on

Bayesian information criterion scores. The ciliates *Cinetochilum ovale* (Genbank accession

number FJ870103) and *Paramecium bursaria* (JX082062) were used respectively as outgroups for

18S rRNA and COI sequences. Ciliates and their hosts were identified by comparing obtained

sequences against GenBank records using Blastn.

3. Results

A total of 256 amphipods collected between December 2022 and April 2023 from eight

sampling sites were screened for sanguicolous ciliates molecularly and histologically (Table 1).

All known amphipod species occurring in the Boye system (Prati et al., 2022) were represented in

the sample. More specifically, *G. fossarum* clade 2 (n = 36, 100% similarity, 100% coverage, e-

value = 0 to ON093817), *G. pulex* clade C (n = 96, 99.9-100% similarity, 100% coverage, e-value

= 0 to ON093813), and clade E (n = 124, 99.9-100% similarity, 100% coverage, e-value = 0 to

ON093815). The most common amphipods in the investigated sites were *G. pulex* clade E and C.

3.1 Prevalence

Half of the sampling sites had ciliate-infected amphipod populations, with overall prevalences

ranging from 6.67 to 21.43% (Table 1). Among the 12 infected amphipods collected between

December 2022 and April 2023, three were *G. fossarum* clade 2, four *G. pulex* clade C, and 5 *G.*

pulex clade E. Two ciliate-infected individuals presented simultaneous infections with

Microsporidium sp. 515. One of them was also infected with a trematode. The 9 infected

amphipods collected in March 2024 and used to isolate fresh ciliates were neither molecularly

identified nor screened for co-infections. Overall, the average size of infected amphipods (4[th]

coaxal plate length 3.07 mm ± 0.44 SD, n = 21) was similar to that of uninfected ones (2.93 mm ±

0.49 SD, n = 277). No individual below 2.57 mm was infected.

3.2 Histology and pathology

Histological analysis indicated that the ciliate is a parasite that travels through the hemolymph and
causes a systemic infection. Freely dividing tomonts and protomites-tomites were visible in the
host hemocoel, surrounding the testis, muscle, hepatopancreas, and gill lamellae (Fig. 1). We found
no evidence that the ciliate infiltrated these organs or any sign of visible immune reactions in the
form of melanization. However, compared to uninfected individuals, the muscular mass
progressively reduced with increasingly high ciliate densities in infected individuals.

3.3 Morphology

Similar to the histological sections, we observed tomonts and protomites-tomites in the
glutaraldehyde-fixed gammarids. The tomonts were subelliptical and had a dense ciliature. Their
length ranged between 34.6 and 90.8 µm (66.8 µm ± 15.3 SD), and the width between 15.4 and
59.1 µm (32.5 µm ± 8 SD). Tomonts had a large central macronucleus (width 48.5 µm ± 13.2 SD
and length 15.9 µm ± 4.8 SD) and contractile vacuoles either arranged in rows parallel or slightly
diagonal to kineties or rather scattered throughout the cytoplasm (Fig. 2 and 3). The largest tomonts
displayed 40 to 47 irregularly spaced kineties converging toward the extremities but ending shortly
before reaching the poles (Fig. 2 and 3). However, the vast majority of tomonts were along the
path of regressive division; thus, it is possible that non-regressing cells might attain a larger size
and kineties number than what is reported here.

Following palintomy, a process during which the parental cell freely undergoes a sequence
of repeated divisions without intervening growth in the host hemocoel, the size and the number of
kineties of the daughter cells are progressively reduced. Regressing tomonts showed a non-ciliated
band likely resulting from the reabsorption of kineties (Fig. 2 and 3). The protomites-tomites shape
varied from pyriform to subelliptical, and their size ranged between 9.9 and 27.8 µm (20.7 ± 4.6)
in length and 9.3 to 23 µm (14 ± 2.9) in width. They had a large macronucleus (length 11.1 ± 3.1
µm and width 6.73 ± 1.8 µm), nine somatic kineties, and in some cases, presented tree buccal
kineties indicated as x, y, and z (Fig. 2 and 3).

3.4 Molecular phylogeny and genetics

Molecular identification of the collected sanguicolous ciliates using aligned 18S rRNA sequences
indicated that these were closely related to *Metacollinia luciensis* (MH200618, 95.90% similarity,
100% coverage, e-value = 0.0), and all belonged to the same haplotype (Fig. 4). Similar results

were obtained for the COI region (MH182621, 75.50% similarity, 99% coverage, e-value = 1e-
140); however, the availability of sequences for this region was very limited (Fig. 4). Molecular
identification of all ciliate life stages isolated from amphipods collected in March 2024 confirmed
that these belonged to the same species and haplotype. The genetic divergence (K2P) between the
sequences obtained in the current study (0 differing nucleotides) and those of the closest related
species *M. lucensis* (69 SNPs) and *Lynnia grapsolytica* (74 SNPs), both belonging to the
Collinidae family, were respectively 4% and 4.7%, suggesting a closer genetic relationship with
M. luciensis. Apostome ciliates belonging to other families diverged by 6.4-6.7% (92-100 SNPs),
supporting the placement of the ciliate in the Collinidae.

3.5 Mitochondrial genetics

The mitochondrial genome of the new ciliated protozoan was identified in two separate contigs,
the larger consisting of 39,710 bp and encoding 28 protein-coding genes (PCGs); and the second
was composed of 20,221 bp and encoded 4 PCGs, as well as both the SSU and LSU rRNA
sequences. tRNA sequences remain unannotated and require more detailed exploration since
predictive methods were unable to clearly identify putative coding regions. The most similar
known relatives are *Metacollina*, *Tetrahymena*, and *Cryptocaryon* where proteins are compared
from both contigs using blastp (Table 2).

Data available from across the Ciliophora provided 23 potential comparators, which
together included four directly comparable proteins (ATP9-COB-COX2-COX1) and were
prepared into a maximum-likelihood phylogenetic tree (Fig. 5). The tree revealed that the novel
species groups most closely with *M. luciensis* isolates MH182621 and AWT08281 (Fig. 5). Within
this same group, we determined that a mitochondrial genome available from the host *Idotea baltica*
(Isopoda; JAACYD010007607) also showed similarity and phylogenetic positioning, highlighting
another putative ciliate of this group that may have gone undiagnosed.

Nuclear genes were not assessed from the metagenomic data during this study; however,
information was gathered from the data to support the PCR diagnostic used to reproduce the 18S
sequence for this species, as described above.

4. Taxonomic summary

Kingdom: Protozoa

Superphylum: Alveolata

Phylum: Ciliophora Doflein, 1901

Subphylum: Intramacronucleata Lynn, 1996

Class: Oligohymenophorea de Puytorac et al. (1974)

Order: Apostomatida Chatton and Lwoff, 1928

Family: Colliniidae Cepede, 1910

Species: *Metacollinia emscheri* n. sp. Prati, Rückert, Grabner, Sures, Bojko, 2024

Formal taxonomic summary: The tomonts are ~67 µm long with 40-47 (possibly more) slightly

spiraled kineties, a non-ciliated cortical band, and 6-10 contractile vacuoles arranged either in two

parallel rows or scattered across the cytoplasm. At the end of palintomy, during which the number

of kineties decreases to nine, oral fragments (buccal kineties x, y, and z) appear.

Type locality: Boye catchment, North-Rhine Westphalia, Germany (Boye 51.553107; 6.948059

and 51.534515; 6.997740, Haarbach 51.570306; 6.960891 and 51.562638; 6.955581)

Type host: *Gammarus pulex* Linnaeus, 1758 (Crustacea, Malacostraca, Amphipoda, Gammaridae)

Type habitat: Endoparasitic, found in the hemolymph and muscle tissues of hosts collected in

lowland streams at the type locality.

Holotype: The name-bearing type of this species is the specimen illustrated in Fig. 2 and 3 (ICZN

1999, Articles 73.1.4).

DNA sequences: 18S rRNA and the draft mitogenome are publicly available via GenBank (18S

PQ013206 - PQ013208, mitogenome: …………….)

Etymology: The species epithet emscheri refers to the Emscher River catchment, which comprises

the type localities found in its tributary, the Boye and the Haarbach.

5. Discussion

This study describes a novel sanguicolous apostome ciliate that infects the amphipods *G. pulex*

and *G. fossarum*, from four populations of the Boye (North Rhine-Westphalia, Germany). Our

phylogenetic analyses place *M. emscheri* n. sp. within the Collinidae family. Due to morphological

characteristics, phylogenetic similarity with the type species, *M. lucensis*, and the absence of DNA

sequences of less molecularly divergent species, we provisionally place this ciliate in the genus

Metacollinia. This concept is supported by *18s, cox1,* and a concatenated multi-protein
phylogenetic approach (Fig. 4-5).

5.1 Prevalence of *Metacollinia emscheri* n. sp.

The presence of *M. emscheri* n. sp. in *G. pulex* clade E, *G. pulex* clade C, and *G. fossarum* clade
C2 indicate lower host specificity than other collinids, as these have mostly been reported from
single hosts. The prevalence of *M. emscheri* n. sp. across the four infected amphipod populations
indicates medium infection levels and, in the case of the population collected from the lower stretch
of the Haarbach (HAAun) temporal stability. This sanguicolous ciliate appears to be localized in
the Haarbach and the lower part of the Boye, since none of the amphipods collected upstream of
BOYohKI (located just above the junction of the two streams) were infected. However, we cannot
be certain that *M. emscheri* n. sp. is absent from the upper stretch of the Boye catchment. This
ciliate might be present in lower abundances and have evaded detection, due to a relatively small
sample size (~ 30 individuals). It is plausible that seasonal dynamics may also play a role - for
example, the prevalence of *P. branchiarum* in *G. pulex* collected in a locality in proximity to
Prague varied between 1% in January and 20% in August (De Puytorac and Lom, 1962). The close
proximity of BOYuhHA, the most downstream site, with the River Emscher (2.7 km), suggests
that *M. emscheri* n. sp. might also be present in the latter.

Assuming that the observed replacement of the muscle mass in infected amphipods may
hinder the host's ability to withstand water flow, it is plausible that a good portion of infected
individuals passively drift downstream and end up in the River Emscher. Furthermore, all infected
amphipods had relatively large sizes, which alone might influence amphipod drift. Accordingly,
both large-sized and microsporidian-infected *G. pulex* in a nearby watercourse, the Rotbach,
drifted more than smaller and uninfected individuals (Prati et al., 2023). The exclusive presence
of *M. emscheri* n. sp. in relatively large-sized amphipods is not surprising. Generally, parasite
prevalence in aquatic organisms, including amphipods, is positively correlated with host size, as
older individuals are more likely to be exposed to infections over their lifespan (Poulin, 2000; Prati
et al., 2023; Soldánová et al., 2010). Furthermore, size-asymmetric cannibalism, which is common
in many aquatic taxa, including *G. pulex*, may potentially favor the transmission of parasites via
ingestion of infected individuals (Arundell et al., 2019a, 2019b; MacNeil et al., 2003; McGrath et
al., 2007).

5.2 Transmission of *Metacollinia emscheri* n. sp.

The transmission of *M. emscheri* n. sp., like that of other members of the Collinidae and Pseudocollinidae families, remains unknown (Chantangsi et al., 2013; Gómez-Gutiérrez et al., 2012; Lynn, 2008; Metz and Hechinger, 2021). Poisson (1921) described *M. luciensis* from the digestive tract of *Orchestia gammarellus*, but mentioned that the ciliate might be able to pierce through the intestine wall and penetrate the coelom. Balbiani (in Cépède, 1910) hypothesized that host injury is required for ciliate transmission. Such a pathway might also be possible for *G. pulex* as injuries often occur in adult individuals, particularly during mating, when males compete to assert their dominance over females (Dick and Elwood, 1990).

Alternatively, pre-existing parasitic infections might favor secondary infections by hindering the host immune system (Cox, 2001). This does not seem to be the case in the present study since most infected amphipods did not show co-infections; similarly, Bojko et al. (2017) noted a few additional parasites present alongside their ciliate in *G. roeselii*, which was sampled in Poland and observed at a prevalence of <1%. A more plausible theory has been proposed by Metz and Hechinger (2021), who, after failing multiple attempts to experimentally infect healthy and injured crabs with *L. grapsolytica* (via exposure to swimming tomites, infected conspecifics, or direct injection of life cells) proposed that the parasite has a complex life cycle with development outside the host. This might possibly involve an intermediate host. Such a possibility might also apply to *M. emscheri* n. sp.; however, more extensive studies are required to shed light on its life cycle. Given that we have identified proteins from the mitochondrial genome with similarity to a protozoan symbiont of *Idotea baltica*, it may be that isopods can also host this group of parasitic ciliates.

5.3 Ecological importance of *Metacollinia emscheri* n. sp.

Irrespective of its elusive life cycle, *M. emscheri* n. sp. is of high ecological importance. With prevalences up to 21% in the infected populations, this parasite is likely to strongly affect host population dynamics. Due to reduced muscular mass, host movement and activities such as feeding and mating might be affected in individuals with high infection levels. As amphipods are keystone species in freshwater aquatic environments, *M. emscheri* n. sp. may indirectly impact ecosystem functioning. For instance, infected individuals may experience reduced shredding activity and,

consequently, a slowdown of leaf-litter breakdown decreases the amount of nutrients and processed material available for microorganisms and other invertebrates. Such a slowdown might be particularly detrimental to small stream food webs as the decomposition of terrestrial organic matter often constitutes the main source of energy (Gessner et al., 1999; Wallace et al., 1997). Furthermore, if *M. emscheri* n. sp. directly or indirectly increases the mortality rate in amphipods, species that rely on such prey may face reduced food availability. Gammarus spp. are an especially important food source for fish and, unlike other macroinvertebrates, are available year-round (Macneil et al., 1999).

5.4 Taxonomy and nomenclature of the family Collinidae

Jankowski (2007) undertook a comprehensive revision of the nomenclature and taxonomy of Collinidae, suggesting that the genus *Collinia* should be split into three distinct genera: *Collinia*, *Paracollinia*, and *Metacollinia*. *Collinia* (type species: *Collinia circulans*) was originally observed by Balbiani 1885 (in Cépède, 1910) in the freshwater isopod, *Asellus aquaticus*. *Collinia circulans* is characterized by a large tomont (70-80 µm long and 30-35 µm wide) with ~ 13 spiraled bipolar kineties and a single row of ~ 12 contractile vacuoles. The tomite is small, showing short x, y, and z kineties reminiscent of those of the foettingeriids (De Puytorac and Lom, 1962; Lynn, 2008; Lynn and Strüder-Kypke, 2019).

The genus *Paracollinia* (type species: *Paracollinia branchiarum*) was observed in 1852 by Stein, to infect the freshwater amphipod *G. pulex*. It has large tomonts with 55 to 60 straight evenly spaced kineties without a non-ciliated cortical band and a smaller flattened tomite with 9 kineties and short x, y, and z kineties, reminiscent of those of the foettingeriids (Jankowski, 2007; Lynn and Strüder-Kypke, 2019).

The genus *Metacollinia* (type species: *Metacollinia luciensis*) was observed in the marine amphipod *O. gammarellus* by Poisson (1921). It has large tomonts with 65 to 70 slightly spiraling kineties, a large non-ciliated cortical band and two lateral rows of 8-10 contractile vacuoles. Like the previous species, the tomite has 9 kineties and short x, y, and z kineties, reminiscent of those of the foettingeriids (Jankowski, 2007; Lynn and Strüder-Kypke, 2019).

More recently, a novel genus, *Lynnia*, (type species: *L. Grapsolytica*) found in the marine crab *Pachygrapsus crassipes* has been included in the Collinidae (Metz and Hechinger, 2021). The tomont is large with ~ 11 equidistant spiraled bipolar kineties, lacks a non-ciliated cortical band,

and possesses randomly scattered subpellicular contractile vacuoles. The tomite is small, showing

short x, y, and z kineties reminiscent of those of the foettingeriids; however, unique compared to

any known apostome, the eighth and ninth kineties are both interrupted, and once impregnated

with silver, the tomite presents a hook-shaped arrangement of ventral somatic kineties (Metz and

Hechinger, 2021).

Compared to *L. grapsolytica*, the tomite of *M. emscheri* n. sp. bears more resemblance with

that of *C. ciruclans*, *P. branchiarum,* and *M. luciensis*. Only the ninth somatic kinety in *M.*

emscheri n. sp. is interrupted, and no hook-shaped arrangement of ventral somatic kineties is

present. Like *M. luciensis*, the tomont of *M. emscheri* n. sp. displayed slightly spiraled kineties and

a non-ciliated cortical band, suggesting the placement of the ciliate in this genus. The average

length of tomont in *M. emscheri* n. sp. (~67 µm) was similar to that reported in *M. luciensis* (~70

µm) by De Puytorac and Grain (1975). However, the kineties, numbering 40-47, were lower than

the 65-70 reported in *M. luciensis.* It is possible that the smaller size and lower number of kineties

observed in *M. emscheri* n. sp. might be due to regression processes occurring during the

tomitogenesis, as reported for *M. luciensis* collected by Lynn and Strüder-Kypke (2019).

Interestingly, the contractile vacuoles in the tomont of *M. emscheri* n. sp., which numbers 6-10 in

contrast to 8-10 found in *M. luciensis* (De Puytorac and Grain, 1975)*,* are arranged either in two

parallel rows or scattered across the cytoplasm.

Phylogenetic analyses supports these morphological deductions, and indicates that *M.*

emscheri n. sp. is more closely related to *M. lucensis* than *L. grapsolytica*, the only other Collinidae

for which DNA sequences are available. In the absence of DNA sequences of species belonging

to the genera *Collinia* and *Paracollinia*, and based on morphological similarity with *M. lucensis*,

we provisionally place *M. emscheri* n. sp. in the genus *Metacollina*. Given the gap in DNA

sequence reference data, we cannot exclude that the proposed placement might change once more

sequences of Collinidae species become available.

Authors contribution

429 Sebastian Prati, Daniel S. Grabner, Bernd Sures, and Jamie Bojko conceived the study. Sebastian

430 Prati carried out the sampling. Sebastian Prati, Sonja Rückert, and Jamie Bojko performed

431 laboratory analyses. Jamie Bojko conducted the metagenomic and bioinformatic analysis and

supervised the study. Sebastian Prati wrote the first draft of the manuscript, and all authors contributed critically to the final draft. All authors read and approved the final manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This study was performed within the Collaborative Research Center (CRC) RESIST (A09) and funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – CRC 1439/1 – project number: 426547801. We are grateful to the Developmental Biology group of the University of Duisburg-Essen and, in particular, to Sabine Schneider for providing access and assistance with the histological instruments. We thank the Imaging Center Essen (IMCES) at the Faculty of Medicine of the University of Duisburg-Essen, in particular Bernd Walkenfortfor for providing access and assistance with the SEM equipment.

Appendix A. Supplementary data

Supplementary material 1

References

Adlard, R.D., Lester, R.J.G., 1995. Development of a diagnostic test for *Mikrocytos roughleyi*, the aetiological agent of Australian winter mortality of the commercial rock oyster, *Saccostrea commercialis* (Iredale & Roughley). J. Fish Dis. 18, 609–614. https://doi.org/10.1111/j.1365-2761.1995.tb00365.x

Arundell, K.L., Bojko, J., Wedell, N., Dunn, A.M., 2019a. Fluctuating asymmetry, parasitism and reproductive fitness in two species of gammarid crustacean. Dis. Aquat. Org. 136, 37–49. https://doi.org/10.3354/dao03395

Arundell, K.L., Dubuffet, A., Wedell, N., Bojko, J., Rogers, M.S.J., Dunn, A.M., 2019b. *Podocotyle atomon* (Trematoda: Digenea) impacts reproductive behaviour, survival and

461   physiology in *Gammarus zaddachi* (Amphipoda). Dis. Aquat. Org. 136, 51–62.

462   https://doi.org/10.3354/dao03416

463 Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,

464   Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N.,

465   Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: a new genome assembly

466   algorithm and its applications to single-cell sequencing. J. Comput. Biol. 19, 455–477.

467   https://doi.org/10.1089/cmb.2012.0021

468 Bengtsson-Palme, J., Hartmann, M., Eriksson, K.M., Pal, C., Thorell, K., Larsson, D.G.J.,

469   Nilsson, R.H., 2015. metaxa2: improved identification and taxonomic classification of

470   small and large subunit rRNA in metagenomic data. Mol. Ecol. Resour. 15, 1403–1414.

471   https://doi.org/10.1111/1755-0998.12399

472 Bojko, J., Bącela-Spychalska, K., Stebbing, P.D., Dunn, A.M., Grabowski, M., Rachalewski, M.,

473   Stentiford, G.D., 2017. Parasites, pathogens and commensals in the "low-impact" non-

474   native amphipod host *Gammarus roeselii*. Parasites Vectors 10, 193.

475   https://doi.org/10.1186/s13071-017-2108-6

476 Bojko, J., Ovcharenko, M., 2019. Pathogens and other symbionts of the Amphipoda: taxonomic

477   diversity and pathological significance. Dis. Aquat. Org. 136, 3–36.

478   https://doi.org/10.3354/dao03321

479 Bojko, J., Stebbing, P., Bateman, K., Meatyard, J., Bacela-Spychalska, K., Dunn, A., Stentiford,

480   G., 2013. Baseline histopathological survey of a recently invading island population of

481   'killer shrimp', *Dikerogammarus villosus*. Dis. Aquat. Org. 106, 241–253.

482   https://doi.org/10.3354/dao02658

483 Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina

484   sequence data. Bioinformatics 30, 2114–2120.

485   https://doi.org/10.1093/bioinformatics/btu170

486 Capriulo, G., Small, E., 1985. Discovery of an apostome ciliate (*Collinia beringensis* n. sp.)

487   endoparasitic in the Bering Sea euphausiid *Thysanoessa inermis*. Dis. Aquat. Org. 1,

488   141–146. https://doi.org/10.3354/dao001141

489 Cépède, C., 1910. Recherches sur les infusoires astomes: anatomie, biologie, ethologie

490   parasitaire, systeématique. Librairie A. Schulz, Paris.

Chantangsi, C., Lynn, D.H., Rueckert, S., Prokopowicz, A.J., Panha, S., Leander, B.S., 2013.
Fusiforma themisticola n. gen., n. sp., a new genus and species of apostome ciliate
infecting the Hyperiid amphipod *Themisto libellula* in the Canadian Beaufort Sea (Arctic
Ocean), and establishment of the Pseudocolliniidae (Ciliophora, Apostomatia). Protist
164, 793–810. https://doi.org/10.1016/j.protis.2013.09.001

Cox, F.E.G., 2001. Concomitant infections, parasites and immune responses. Parasitology 122,
S23–S38. https://doi.org/10.1017/S003118200001698X

De Puytorac, P., Grain, J., 1975. Etude de la tomitogenèse et de l'ultrastructure de *Collinia*
orchestiae, cilié apostome sanguicole, endoparasite du crustacé *Orchestia gammarella*
pallas. Protistologica 11, 61–74.

De Puytorac, P., Lom, J., 1962. La tomitogénèse des *Collinia* ciliés apostomes sanguicoles
endoparasites des crustacés. Ann. Parasitol. Hum. Comp. 37, 195–209.
https://doi.org/10.1051/parasite/1962373195

Dick, J.T.A., Elwood, R.W., 1990. Symmetrical assessment of female quality by male
Gammarus pulex (Amphipoda) during struggles over precopula females. Anim. Behav.
40, 877–883. https://doi.org/10.1016/S0003-3472(05)80989-3

Gessner, M.O., Chauvet, E., Dobson, M., 1999. A perspective on leaf litter breakdown in
streams. Oikos 85, 377–384. https://doi.org/10.2307/3546505

Giari, L., Fano, E.A., Castaldelli, G., Grabner, D., Sures, B., 2020. The ecological importance of
amphipod–parasite associations for aquatic ecosystems. Water 12, 2429.
https://doi.org/10.3390/w12092429

Gómez-Gutiérrez, J., Kawaguchi, S., 2017. *Pseudocollinia* histophagous ciliates infect krill in
the Pacific and Atlantic Oceans and possibly worldwide. CICIMAR Oceánides 32, 15–
24. https://doi.org/10.37543/oceanides.v32i2.204

Gómez-Gutiérrez, J., Strüder-Kypke, M.C., Lynn, D.H., Shaw, T.C., Aguilar-Méndez, M.J.,
López-Cortés, A., Martínez-Gómez, S., Robinson, C.J., 2012. *Pseudocollinia brintoni*
gen. nov., sp. nov. (Apostomatida: Colliniidae), a parasitoid ciliate infecting the
euphausiid *Nyctiphanes simplex*. Dis. Aquat. Org. 99, 57–78.
https://doi.org/10.3354/dao02450

Grabner, D.S., Weigand, A.M., Leese, F., Winking, C., Hering, D., Tollrian, R., Sures, B., 2015.
Invaders, natives and their enemies: distribution patterns of amphipods and their

microsporidian parasites in the Ruhr Metropolis, Germany. Parasit. Vectors 8, 419.

https://doi.org/10.1186/s13071-015-1036-6

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: quality assessment tool for

genome assemblies. Bioinformatics 29, 1072–1075.

https://doi.org/10.1093/bioinformatics/btt086

Jankowski, A., 2007. Phylum Ciliophora Doflein, 1901, in: Alimov, A. F. (Ed.), Protista, Part 2,

Handbook of Zoology. Russian Academy of Science, Zoological Institute, St. Petersburg,

pp. 415–993.

Ji, D., Wang, Y., 2018. An optimized protocol of protargol staining for ciliated protozoa. J.

Eukaryot. Microbiol. 65, 705–708. https://doi.org/10.1111/jeu.12515

Katoh, K., Rozewicki, J., Yamada, K.D., 2019. MAFFT online service: multiple sequence

alignment, interactive sequence choice and visualization. Brief. Bioinformatics. 20,

1160–1166. https://doi.org/10.1093/bib/bbx108

Kurilov, A.V., 2017. Improvement of silver impregnation technique using in situ synthesized

protargol. Acta Protozool. 56.

Lang, B.F., Beck, N., Prince, S., Sarrasin, M., Rioux, P., Burger, G., 2023. Mitochondrial

genome annotation with MFannot: a critical analysis of gene identification and gene

model prediction. Front. Plant Sci. 14. https://doi.org/10.3389/fpls.2023.1222186

Lynn, D.H. (Ed.), 2008. The ciliated protozoa: characterization, classification, and guide to the

literature., 3rd ed. Springer Netherlands, Dordrecht. https://doi.org/10.1007/978-1-4020-

8239-9_17

Lynn, D.H., Gómez-Gutiérrez, J., Strüder-Kypke, M.C., Shaw, C.T., 2014. Ciliate species

diversity and host-parasitoid codiversification in *Pseudocollinia* infecting krill, with

description of *Pseudocollinia similis* sp. nov. Dis. Aquat. Org. 112, 89–102.

https://doi.org/10.3354/dao02796

Lynn, D.H., Strüder-Kypke, M.C., 2019. The sanguicolous apostome *Metacollinia luciensis*

Jankowski 1980 (Colliniidae, Apostomatia, Ciliophora) is not closely related to other

sanguicolous apostomes. J. Eukaryot. Microbiol. 66, 140–146.

https://doi.org/10.1111/jeu.12638

Macneil, C., Dick, J.T.A., Elwood, R.W., 1999. The dynamics of predation on *Gammarus* spp.

(Crustacea: Amphipoda). Biol. Rev. 74, 375–395.

https://doi.org/10.1017/S0006323199005368

Macneil, C., Dick, J.T.A., Elwood, R.W., 1997. The trophic ecology of freshwater *Gammarus*

spp. (Crustacea: Amphipoda): problems and perspectives concerning the functional

feeding group concept. Biol. Rev. Cambridge Philos. Soc. 72, 349–364.

https://doi.org/10.1017/S0006323196005038

MacNeil, C., Dick, J.T.A., Hatcher, M.J., Fielding, N.J., Hume, K.D., Dunn, A.M., 2003.

Parasite transmission and cannibalism in an amphipod (Crustacea). Int. J. Parasitol. 33,

795–798. https://doi.org/10.1016/S0020-7519(03)00110-3

McGrath, K.E., Peeters, E.T.H.M., Beijer, J.A.J., Scheffer, M., 2007. Habitat-mediated

cannibalism and microhabitat restriction in the stream invertebrate *Gammarus pulex*.

Hydrobiologia 589, 155–164. https://doi.org/10.1007/s10750-007-0731-5

Metz, D.C.G., Hechinger, R.F., 2021. *Lynnia grapsolytica* n. gen, n. sp. (Ciliophora:

Apostomatida: Colliniidae), a deadly blood parasite of crabs with a novel

pseudocytopharynx. J. Eukaryot. Microbiol. 68, e12847.

https://doi.org/10.1111/jeu.12847

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,

Lanfear, R., 2020. IQ-TREE 2: New models and efficient methods for phylogenetic

inference in the genomic era. Mol. Biol. Evol. 37, 1530–1534.

https://doi.org/10.1093/molbev/msaa015

Poisson, R., 1921. Sur un infusoire du genre *Balantidium*, parasite du tube digestif d'*Orchestia*

littorea Mont. Compt. Rend. Seances Soc. Biol. Fil 84, 333–335.

Poulin, R., 2000. Variation in the intraspecific relationship between fish length and intensity of

parasitic infection: biological and statistical causes. J. Fish Biol. 56, 123–137.

https://doi.org/10.1111/j.1095-8649.2000.tb02090.x

Prati, S., Enß, J., Grabner, D.S., Huesken, A., Feld, C.K., Doliwa, A., Sures, B., 2023. Possible

seasonal and diurnal modulation of *Gammarus pulex* (Crustacea, Amphipoda) drift by

microsporidian parasites. Sci Rep 13, 9474. https://doi.org/10.1038/s41598-023-36630-2

580 Prati, S., Grabner, D.S., Pfeifer, S.M., Lorenz, A.W., Sures, B., 2022. Generalist parasites persist

581   in degraded environments: a lesson learned from microsporidian diversity in amphipods.

582   Parasitology 149, 973–982. https://doi.org/10.1017/S0031182022000452

583 Prokopowicz, A.J., Rueckert, S., Leander, B.S., Michaud, J., Fortier, L., 2010. Parasitic infection

584   of the hyperiid amphipod *Themisto libellula* in the Canadian Beaufort Sea (Arctic

585   Ocean), with a description of *Ganymedes themistos* sp. n. (Apicomplexa,

586   Eugregarinorida). Polar Biol 33, 1339–1350. https://doi.org/10.1007/s00300-010-0821-0

587 Skibbe, O., 1994. An improved quantitative protargol stain for ciliates and other planktonic

588   protists. Arch. Hydrobiol. 339–347. https://doi.org/10.1127/archiv-

589   hydrobiol/130/1994/339

590 Soldánová, M., Selbach, C., Sures, B., Kostadinova, A., Pérez-del-Olmo, A., 2010. Larval

591   trematode communities in *Radix auricularia* and *Lymnaea stagnalis* in a reservoir system

592   of the Ruhr River. Parasit Vectors 3, 56. https://doi.org/10.1186/1756-3305-3-56

593 Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis

594   Version 11. Mol. Biol. Evol. 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

595 Wallace, J.B., Eggert, S.L., Meyer, J.L., Webster, J.R., 1997. Multiple trophic levels of a forest

596   stream linked to terrestrial litter inputs. Science 277, 102–104.

597   https://doi.org/10.1126/science.277.5322.102

598 Warren, D.A., Burgess, A.L., Karemera, F., Bacela-Spychalska, K., Stentiford, G.D., Bojko, J.,

599   2022. Histopathological survey for parasite groups in *Gammarus varsoviensis*

600   (Amphipoda). Diseases of Aquatic Organisms 149, 47–51.

601   https://doi.org/10.3354/dao03658

602 Warren, D.A., Burgess, A.L., Prati, S., Bacela-Spychalska, K., S. J. Rogers, M., Bojko, J., 2023.

603   Histopathological screening of *Pontogammarus robustoides* (Amphipoda), an invader on

604   route to the United Kingdom. Journal of Invertebrate Pathology 200, 107970.

605   https://doi.org/10.1016/j.jip.2023.107970

607 **Table 1.** Sampling locations, sampling date, amphipod species, average size of the 4th coaxal plate

608 in mm with SD, and prevalence in % of *Metacollinia emscheri* n. sp.

Location	Coordinates	Date	N	Species	Size in mm	Prevalence in %
BOYohBR	51.569135; 6.927698	27/04/2023	4	*Gammarus pulex* clade C	2.88 ± 0.21	0
			26	*Gammarus pulex* clade E	2.95 ± 0.46	0
BOYohKI	51.553107; 6.948059	07/12/2022	8	*Gammarus pulex* clade C	2.57 ± 0.31	12.50 (1/8)
			4	*Gammarus pulex* clade E	2.49 ± 0.37	0
		13/05/2023	18	*Gammarus pulex* clade C	2.63 ± 0.43	0
			12	*Gammarus pulex* clade E	2.71 ± 0.41	0
BOYohSP	51.564297; 6.930277	27/04/2023	10	*Gammarus fossarum* clade 2	2.82 ± 0.56	0
			8	*Gammarus pulex* clade C	2.99 ± 0.74	0
			12	*Gammarus pulex* clade E	2.96 ± 0.57	0
BOYuhHA	51.534515; 6.997740	02/05/2023	26	*Gammarus fossarum* clade 2	3.08 ± 0.51	11.53 (3/26)
			4	*Gammarus pulex* clade C	3.28 ± 0.66	0
BRA1000	51.577238; 6.932752	27/04/2023	1	*Gammarus pulex* clade C	3.36	0
			25	*Gammarus pulex* clade E	2.95 ± 0.38	0
HAAob	51.570306; 6.960891	13/05/2023	11	*Gammarus pulex* clade C	2.87 ± 0.30	9.09 (1/11)
			19	*Gammarus pulex* clade E	2.81 ± 0.36	5.26 (1/19)
HAAun	51.562638; 6.955581	13/05/2023	9	*Gammarus pulex* clade C	3.02 ± 0.31	22.22 (2/9)
			21	*Gammarus pulex* clade E	2.92 ± 0.49	19.05 (4/21)
		02/03/2024	42	Not determined	3.07 ± 0.47	21.43 (9/42)
VORohBOY	51.554753; 6.932435	07/12/2022	8	*Gammarus pulex* clade C	2.84 ± 0.29	0
		13/05/2023	25	*Gammarus pulex* clade C	3.15 ± 0.63	0
			5	*Gammarus pulex* clade E	3.07 ± 0.71	0

610 **Table 2.** A blast table comparing the protein coding genes (as determined by MFannot) from both

611 segments of the mitochondrial genome to available sequences held by NCBI.

Mitogenome segment	Gene	Coding region (bp)	BlastP hit (accession)	Coverage, Similarity, e-value	Function/Protein
1	*ccmF/yejr*	732-2111	*Tetrahymena rostrata* (YP010411968)	32%, 32.52%, 1e^{-12}	Heme maturase
	nad7	2148-3722	*Tetrahymena paravorax* (YP740772)	77%, **61.07**%, 3e^{-127}	NADH7
	rp114	4539-4898	*Tetrahymena pyriformis* (NP049608)	98%, 53.85%, 1e^{-34}	Ribosomal protein
	nad2	5252-5725	-	-	-
	cob	6026-7306	Paramecium caudatum (YP003734430)	96%, 42.03%, 2e^{-100}	COB
	cox1	8120-10108	*Metacollinia luciensis* (AWT08281)	40%, **75.19**%, 2e^{-133}	COX1
	orf254	10123-10887	-	-	-
	nad5	1095-12718	Protozoan of *Idotea baltica* (MCL4129281)	42%, 52.78%, 9e^{-79}	NADH5
	orf525	12734-14311	-	-	-
	nad1	14326-15138	Paramecium gigas (YP010183551)	85%, 55.41%, 1e^{-80}	NADH1
	orf196	16186-16776	-	-	-
	nad4	17260-18777	Tetrahymena rostrata (YP010411987)	73%, 45.99%, 1e^{-87}	NADH4
	nad3	19343-19618	Protozoan of *Idotea baltica* (MCL4129282)	95%, 56.32%, 1e^{-24}	NADH3
	orf334	19717-20721	-	-	-
	orf237	20883-21596	-	-	-
	ymf64	21779-24178	*Pseudocohnilembus persalinus* (QLI42455)	18%, 35.10%, 2e^{-07}	YMF64
	nad10	24187-24663	*Cryptocaryon irritans* (YP010632201)	99%, **73.89**%, 6e^{-90}	NADH10
	orf198	24672-25268	-	-	-
	rpl2	25343-26119	*Cryptocaryon irritans* (YP010632200)	95%, 36.11%, 7e^{-23}	Ribosomal protein
	rps19	26139-26441	-	-	Ribosomal protein
	rps13	26453-27424	Tetrahymena rostrata (QBI37874)	68%, 38.76%, 3e^{-30}	Ribosomal protein
	orf171	28449-28964	-	-	-
	nad9	28984-29610	-	-	-
	rps12	29954-30349	Protozoan of *Idotea baltica* (MCL4137936)	75%, **66.67**%, 2e^{-37}	Ribosomal protein
	orf169	30374-30883	-	-	-
	cox2	30978-34934	Paramecium caudatum (YP003734428)	29%, 43.62%, 6e^{-37}	COX2
	orf265	35066-35863	-	-	-
	orf474	37121-38545	-	-	-
2	*orf699*	4260-2161	-	-	-
	orf525	10199-8622	-	-	-
	orf549	11884-10235	-	-	-
	atp-9	13388-13158	Tetrahymena paravorax (YP740778)	89%, **67.65**%, 1e^{-24}	ATP-9

Fig. 1. Histopathological preparations of the ciliate infection in amphipod tissue. A) Ciliated protozoans (white triangle) with basophilic staining nuclei (black triangle) are present in the hemocoel of the infected amphipod, alongside hemocytes (orange arrow). The image includes hemocoel ('Heam') surrounding the hepatopancreas ('HP') and muscle. B) Ciliated protozoans (white triangle) passing through the gill lamellae ('Gill'), alongside the outer host cuticle. C)

619   Ciliate-like features are observable (white arrow) on the outside of the protozoans. D) Ciliated

620   protozoans (white triangle) seen to travel through the hemocoel ('Heam') adjacent to the testis, gut,

621   and hepatopancreas ('HP'). An individual in section highlights the common U-shaped nucleus often

622   noted in Ciliophora members.

Fig. 2. A) tomont with 47 kineties, B) internal morphology of tomont showing macronucleus and contractile vacuoles, C) regressing tomont displaying a non-ciliated cortical band, D) dividing protomite showing 9 kineties and the x, y, and z kineties, and E) conjugant.

Fig. 3. A) dividing tomont, B) regressing tomont displaying a non-ciliated cortical band, C) tomont showing kineties merging toward the poles but not touching them, D) dividing tomite and conjugant, E) light microscopy of freshly isolated tomont with kineties and a large macronucleous visible, and F) QPS staining of tomont showing contractile vacuoles arranged in parallel lines and a large macronucleous.

Fig. 4. A) maximum-likelihood phylogeny of Apostome ciliates incorporating 18S rDNA and B) mitochondrial COI sequences. The tree includes the ciliates *Cinetochilum ovale* (18s) and *Paramecium bursaria* (COI) as outgroup. Best-fit model: HKY+F+I+R2 (18s) and TPM2u+F+I+G4 (COI) chosen according to Bayesian Information Criterion. Gray dots indicate bootstrap support values above 90 (1,000 replicates).

ATP9-COB-COX1-COX2

Fig. 5. A maximum-likelihood tree based on four proteins (ATP9, COB, COX1, COX2), using
mitochondrial sequences from across the Ciliophora. The tree includes 25 sequences (*Saprolegnia*
is used as an outgroup) and includes 3509 columns, 2357 distinct patterns, 1548 parsimony-
informative, 595 singleton sites, 1366 constant sites. Best-fit model: mtZOA+F+I+G4 chosen
according to Bayesian Information Criterion. Log-likelihood of consensus tree: -46022.229.

650