Changing feeding levels reveal plasticity in elasmobranch life history strategies

Sol Lucas*, Per Berggren, Ellen Barrowclift, Isabel M. Smallegange*

School of Natural and Environmental Sciences, Newcastle University,

Newcastle upon Tyne, NE1 7RU, UK

*Author contact: s.a.lucas2@newcastle.ac.uk, Isabel.smallegange@newcastle.ac.uk

ORCID ID's: Smallegange Orcid: 0000-0001-6218-7358. Lucas Orcid: 0000-0003-2497-7089. Berggren Orcid: 0000-0001-8204-7256. Barrowclift Orcid: 0000-0003-0625-0919.

Abstract

Life history strategies are shaped by phylogeny, environmental conditions and individual energy budgets, and have implications for population performance. Here, we used an approach that merges demography with energy budget theory to structure life history traits of 151 elasmobranch species into life history strategies for two contrasting feeding levels. We assessed how phylogeny and habitat impacted life history strategies, and tested if these strategies predict population performance and conservation status. Elasmobranch life history strategies are structured along the fast-slow continuum and reproductive strategy axes. However, species' positions in this life history space were not fixed, but instead moved in an anticlockwise 'whirlpool' manner along the two axes in response to an increase in feeding level. We also found that population growth rate does not necessarily inform on a species demographic resilience. Finally, only at the higher feeding level does the fast-slow continuum predict IUCN conservation status, with the slowest species at the highest risk of extinction. Our analyses reveal plasticity in species life history strategies, and warn against extrapolating the fast-slow continuum and reproductive strategy framework from one environment to another when predicting a species' response to (climate) change, perturbations, and, particularly in case of elasmobranchs, (over)exploitation.

1. Introduction

Life history strategies of many animals and plants reflect "life history speed" or "pace of life" along a fast-slow continuum, and reproductive biology along a reproductive strategy axis[1–3]. Fast species mature early, grow fast and are short-lived and slow species are slow-growing and long-lived with low fecundity[4]. The reproductive strategy axis is characterised by high lifetime reproductive output and high mortality at one end, and low reproductive output and low mortality at the other[5] (in some studies this second axis relates to development instead[6]). These axes are shaped by trade-offs[4]. For

example, the fast-slow continuum represents the trade-off between reproduction versus survival[6],

often irrespective of body size or phylogenetic relatedness[7–9]. Yet, trade-offs that operate within

individuals, such as trading off growth versus reproduction in the energy budget[7,8], should be

considered, as different dynamics can exist between populations or species with similar traits[10,11].

Specifically, the classical association between life history strategies and population responses to

environmental change breaks down when accounting for individual-level trade-offs and energy

allocation in ray-finned fish[12]. There is also limited understanding of how ecological and evolutionary

factors, like phylogeny or habitat type, shape variation in life history strategies across species[1,2,4,13].

Identifying patterns in life history strategies across the tree of life is essential to predicting

population growth rates and demographic resilience (herein 'population performance')[2]. However,

to do so requires in-depth understanding of how energy budgets, phylogeny and habitat structure

life history strategies across species[1,13,14].

Elasmobranchs (sharks, skates and rays) encompass a vast amount of life history variation.

Longevity, for example, can range from five (e.g. *Carcharhinus sealei* and *Carcharhinus sorrah*[15]) to

almost 400 years (*Somniosus microcephalus*[16]). Reproduction is highly variable, including oviparity

(skates and three families of shark), aplacental viviparity (in various forms; sharks and rays) and

placental viviparity (some sharks)[17,18]. Elasmobranch population performance typically varies with

body size, reproductive strategy and environmental temperature[19–21], yet little is known about the

impact of energy budgets. Here, we investigate how individual energy budgets, phylogeny and

habitat structure elasmobranch life history strategies, and how the resulting fast-slow continuum

and reproductive strategy framework links to their population performance, demographic resilience

and conservation biology[2,13]. We parameterise Dynamic Energy Budget Integral Projection Models

(DEB-IPMs) for 157 elasmobranch species[22–26]. In DEB-IPMs, an energy budget model describes a

species' growth and reproduction, where energy intake is determined by setting an experienced

feeding level[27,28]. To identify life history strategies, we (objective 1) calculate, using the

parameterised DEB-IPMs, a set of representative life history traits based on schedules of survival,

growth and reproduction for a low and a high feeding level. We (objective 2) evaluate the variation

in these traits along major axes using a phylogenetically-corrected principal components analysis

(PCA). We then (objective 3) assess to what extent phylogenetic ancestry and habitat (water

temperature) determine species position along these major axes. Finally, (objective 4) we test

whether species position along these axes predicts population growth rate, speed of recovery from

perturbations (demographic resilience) and IUCN conservation status.

2. Methods

2.1 Brief description of the DEB-IPM

A DEB-IPM describes the dynamics of a population comprising cohorts of females of different sizes

as a result of their survival, growth and reproduction[22,26]. Specifically, denoting the number of

females at year t by $N(L,t)$ gives the dynamics of the body length number distribution from year t

to $t+1$ as:

$$N(L',t+1) = \int_\Omega \left[D(L',L(t))R(L(t)) + G(L',L(t))S(L(t)) \right] N(L,t)dL \qquad \text{eqn 1}$$

where the closed interval Ω denotes the length domain. The survival function $S(L(t))$ in equation (1)

is the probability that an individual of length L survives from time t to $t + 1$:

$$S(L(t)) = \begin{cases} e^{-\mu_j} & \text{for } L_b \leq L < L_p \ \& \ L \leq L_m E(Y)/\kappa, \\ e^{-\mu_a} & \text{for } L_p \leq L \leq L_m \ \& \ L \leq L_m E(Y)/\kappa, \\ 0 & \text{otherwise} \end{cases} \qquad \text{eqn 2}$$

where $E(Y)$ can range from zero (empty gut) to one (full gut), L_b is length at birth, L_p length at puberty

and L_m maximum attainable length (Fig. 1). Individuals die from starvation at a body length at which

maintenance requirements exceeds the total amount of assimilated energy, which occurs when $L >$

$L_m \cdot E(Y)/\kappa$ and hence, $S(L(t)) = 0$ (e.g. an individual of length L_m will die of starvation if $E(Y) < \kappa$,

where κ is the fraction of assimilated energy allocated to respiration, with $1 - \kappa$ allocated to

reproduction [Fig. 1]). Juveniles and adults often have different mortality rates, and thus, juveniles

$(L_b \leq L < L_p)$ that do not die of starvation (i.e. $L \leq L_m \cdot E(Y)/\kappa$) have a mortality rate of μ_j and

adults $(L_p \leq L \leq L_m)$ that do not die of starvation (i.e. $L \leq L_m \cdot E(Y)/\kappa$) have a mortality rate of μ_a

(Fig. 1).

In equation (1), the function $G(L',L(t))$ is the probability that an individual of body length L at

time t grows to length L' at $t + 1$, conditional on survival, and follows a Gaussian distribution:

$$G(L',L(t)) = \frac{1}{\sqrt{2\pi\sigma_L^2(L(t+1))}} e^{\frac{-(L'-E(L(t+1))^2}{2\sigma_L^2(L(t+1))}} \qquad \text{eqn 3}$$

with the growth realised by a cohort of individuals with length $L(t)$ equalling:

$$E(L(t+1)) = \begin{cases} L(t)e^{-r_B} + (1 - e^{-r_B})L_m E(Y) & \text{for } L \leq L_m E(Y) \\ L(t) & \text{otherwise} \end{cases}, \qquad \text{eqn 4}$$

and the variance in length at time $t + 1$ for a cohort of individuals of length L as:

$$\sigma^2(L(t+1)) = \begin{cases} (1 - e^{-r_B})^2 L_m^2 \sigma^2(Y) & \text{for } L \leq L_m E(Y) \\ 0 & \text{otherwise} \end{cases} \qquad \text{eqn 5}$$

where $\sigma(Y)$ is the standard deviation of the expected feeding level, and where $\dot{r}_B$ is the von

Bertalanffy growth rate (Fig. 1).

The reproduction function $R(L(t))$ in equation (1) gives the number of offspring produced

between time t and $t + 1$ by an individual of length L at time t:

$$R(L(t)) = \begin{cases} 0 & \text{for } L_b \leq L < L_p \\ E(Y)R_m L(t)^2/L_m^2 & \text{for } L_p \leq L \leq L_m E(Y) \\ \frac{R_m}{1-\kappa}\left[E(Y)L(t)^2 - \frac{\kappa L(t)^3}{L_m}\right] & \text{for } L_m E(Y) < L \leq L_m E(Y)/\kappa \end{cases} \qquad \text{eqn 6}$$

Individuals are mature when they reach puberty at body length L_p and only surviving adults

reproduce; thus, only individuals within a cohort of length $L_p \leq L \leq L_m Y/\kappa$ reproduce.

Finally, the probability density function $D(L', L(t))$ gives the probability that the offspring of

an individual of body length L are of length L' at time $t + 1$, and hence describes the association

between parent and offspring character values:

$$D(L', L(t)) = \begin{cases} 0 & \text{for } L < L_p \\ \frac{1}{\sqrt{2\pi\sigma_{L_b}^2(L(t))}} e^{\frac{-(L' - E_{L_b}(L(t)))^2}{2\sigma_{L_b}^2(L(t))}} & \text{otherwise} \end{cases} \qquad \text{eqn 7}$$

where $E_{L_b}(L(t))$ is the expected size of offspring produced by a cohort of individuals with length

$L(t)$, and $\sigma_{L_b}^2(L(t))$ the associated variance. For simplicity, $E_{L_b}(L(t))$ is set to be constant and its

associated variance, $\sigma_{L_b}^2(L(t))$ is assumed to be very small.

DEB-IPMs were parameterised (**Error! Reference source not found.**) using the DEBBIES database of e

ctotherms for 157 elasmobranch species, from 11 orders and 37 families[26].

2.2 Life history strategies

Nine representative life history traits were calculated based on schedules of survival, growth and

reproduction (Table 1)[1,2,13] in MATLAB Version 9.12[29] using the code in Smallegange and Lucas[26]

(objective 1). For each species, this was done for a low and high feeding level, $E(Y)$= 0.6 and $E(Y)$=

0.9, respectively. Feeding levels lower than $E(Y)$= 0.6 were not explored because these were too low

for individual growth and reproduction to occur for many species. To identify life history strategies

along major axes (objective 2), we performed a varimax-rotated, phylogenetically corrected principal

component analysis (pPCA)[30], using the *phy.pca* function in the R library *phytools*[2,30–33]. Traits were

log-transformed and scaled to adhere to PCA assumptions ($\mu = 0$ and SD$= 1$).

To correct for phylogenetic relatedness among species, a pool of 10,000 possible phylogenetic trees

were obtained from Stein et al.[34], available at Vertlife.org. These trees represented 500 root node

ages combined with 20 scenarios for infilling taxon with no genetic data. We randomly selected

twenty trees and ran the pPCA for each, on both the high and low feeding levels ($E(Y)= 0.6$ and $E(Y)=$

0.9). Each rooted phylogenetic tree had its branch lengths scaled proportionally based on time of

separation of clades and species, using the *rotl* package[35]. In the tree, branch length informs on

phylogenetic relatedness. The pPCA then links the phylogeny to the life-history traits via a modified

covariance matrix and estimated Pagel's λ, a scaling parameter for the phylogenetic correlation

between species, expected under Brownian motion[36]. Pagel's λ was estimated using the *ape*

package[37] and varies between 0 (phylogenetic independence) and 1 (species traits covary

proportionally to their shared evolutionary history)[38]. The loadings for each axis were consistent

across each of the trees (Appendix: File S1) and Pagel's λ was either consistently above or below our

cut-off value of $\lambda > 0.25$ (Appendix: File S2), above which we assumed the phylogenetic signal is

strong for both feeding levels. Therefore, we report results from one single tree, selected at random

from the subsample of twenty trees (Appendix: Fig. S1). Three species (*S. hawaiiensis*, *A. narutobiei*

and *M. ambigua*) were not in the Vertlife.org database and thus excluded from the phylogenetic tree

and analyses. Two species, *E. granulosus* and *A. bovinus*, were removed because they did not have

an associated polygon shapefile on the IUCN database (see below). After visual inspection of the

PCAs, *S. microcephalus* (Greenland shark) was identified as an outlier and removed, because of its

extreme life history trait values compared to all other species in the dataset.

We checked for the influence of phylogeny and body mass on the final structuring of life history

strategies (see Jeschke & Kokko for a detailed discussion[39]) by running PCA analyses with and

without a correction for phylogeny, and with and without a correction for body mass. Body mass

was corrected for by computing the residuals for linear models between the log_{10}-transformed life

history traits (Table 1) and the body mass for each species for both the PCA and the pPCA[3,40,41]. Body

mass values were extracted from Fishbase[42] using the *rfishbase* package [43]. Where available, we used

the maximum mass given on Fishbase. For species without mass records, we employed length-

weight relationships to infer the maximum body mass. If the length-weight relationship used a

length that was not present in the database, length-length conversions were conducted prior to the

length-weight conversion (e.g. converting fork length to pre-caudal length). Data were not available

for all species. The total number of species included in each PCA are summarised in Table S1

(Appendix). We assessed the significance of PC axes using Kaiser's criterion, retaining those PC axes

with eigenvalues greater than unity[44].

To compare if and how species position changed between the low and high feeding levels, we took

the standardised score (z-score) of each principal components for all species in each analysis (n= 63

for *E(Y)*= 0.6, n= 151 for *E(Y)*= 0.9). The standardised scores were plotted only for the 63 species that

were in both analyses.

2.3 Effect of phylogenetic ancestry and habitat (water temperature) on life history patterns

For each feeding level, we assessed if habitat and phylogenetic clade elasmobranch life history

variation (objective 3). First, we sourced habitat information (water depth and environmental

temperature) for each species. For depth, continuous variable 'commonly used depth' was sourced

for each species from the Fishbase database[42,43]. If commonly used depth was unavailable, median

depth was sourced from the IUCN database[45]. For the continuous variable temperature, we used

'temperature-at-depth' following the methodology of Barrowclift et al.[19]. To this end, shape files

containing polygons of the known depth range for each species were taken from the IUCN

database[45]. These shapefiles were overlaid onto the International Pacific Research Center's mean

annual ocean temperatures across 27 depth levels (0-2000m), based on a dataset from the Argo

Project (available at

http://apdrc.soest.hawaii.edu/projects/Argo/data/statistics/On_standard_levels/Ensemble_mean/1

x1/m00/index.html). Median temperature-at-depth was calculated for each species using values at

the nearest depth level to each species' commonly used depth (or the IUCN median depth where

commonly used depth was unavailable). For mesotherm species (family *Lamnidae*) a correction

factor of 3.5°C was applied to their median temperature.

We tested collinearity between depth, temperature and clade using visual inflation factor (threshold

value VIF>10) in the *faraway* R package[46]. Depth, temperature and clade were all colinear.

Therefore, we separately assessed the effect of temperature at depth (as the representative variable

for habitat) and clade on the PC scores of the first and second axis of the best fitting (p)PCA for each

feeding level. We used four general linear models (LMs) for each feeding level, where the predictor

variable was either temperature at depth or phylogenetic clade, and the response variable was

either PC1 or PC2 score. For each LM, the model assumptions of Gaussian errors, homoscedasticity

and collinearity were confirmed by inspecting the probability plots and error structures in R[32,46,47].

2.4 Life history strategies as predictors of population growth rate, demographic resilience and

conservation status

We used LMs to test, for each feeding level, if the PC scores (continuous variables) of the first and

second axis of the best fitting PCA, and their interaction, predict population growth rate (λ) and

demographic resilience across species (objective 4). To calculate λ, we discretised each DEB-IPM

(Eqn 1) by dividing the length domain Ω into 200 very small-width discrete bins, resulting in a matrix

A of size $m \times n$, where $m = n = 200$, and which dominant eigenvalue equals λ (Table 1). Demographic

resilience was calculated as the damping ratio ξ, with $\xi = \lambda / ||\lambda_2||$, where λ_2 is the highest

subdominant eigenvalue of matrix **A**[48,49]. To assess predictive links between life history strategies

and conservation status, global conservation statuses for each species were sourced from the IUCN

database[45], where LC indicates Least Concern, NT indicates Near Threatened, VU indicates

Vulnerable, EN indicates Endangered, CR indicates Critically Endangered and DD indicates Data

Deficient. We used an ordinal regression to assess if PC scores predict species IUCN conservation

status (excluding Data Deficient species).

All LMs, ordinal regression analyses and plots[50–53] were performed in R version 4.3.1[32] in Rstudio[54],

using the *dplyr* and *tibble* packages for data manipulation[55,56]. For each LM, the model assumptions

of Gaussian errors, homoscedasticity and collinearity were confirmed by inspecting the probability

plots and error structures in R[32,46,47].

205   3   Results

3.1 Elasmobranch life history strategies structure along three axes (objective 1 & 2)

At the low feeding level ($E(Y) = 0.6$), we found that elasmobranch life history traits structure along

three separate PC axes that cumulatively explain 88.2% of the total variance in life histories (Table 1:

PC1: 42.1%, PC2: 31.2%, PC3: 14.9%). Pagel's λ was lower than our cut-off value of 0.25 (Pagel's $\lambda <$

0.001) (Table 1), so the phylogenetic signal did not impact the results. Correcting for body mass did

not qualitatively change the results either (Appendix: Table S1) and we thus continued with the PCA

results. Generation time (T), progressive growth (γ) and mature life expectancy (L_ω) loaded positively

onto PC1 (Fig. 2A) (Table 1). As PC1 scores move from negative to positive, individuals increase their

allocation to progressive growth and mature life expectancy (longevity) at the expense of population

turnover (i.e., greater generation time), suggesting an axis analogous to the fast-slow continuum[4]

(Fig. 2A). Degree of iteroparity (S), mean recruitment success (φ) and net reproductive rate (R_0) were

loaded positively onto PC2 and retrogressive growth (ρ) negatively (Fig. 2A) (Table 1). These life

history traits are all associated with reproductive strategies. Thus, as PC2 scores move from negative

to positive, elasmobranchs attain greater lifetime reproductive success while their retrogressive

growth decreases (Fig. 2A). Finally, survivorship curve type (H) and mean age at maturity (L_α) were

positively and negatively, respectively, loaded onto PC3 (Table 1). This means that as age at maturity

increases (high L_α), senescence rates increase (mortality rate increases with age, corresponding to

low H).

Like the low feeding level, at the high feeding level ($E(Y)$= 0.9), we also found that elasmobranch life

history traits structure along three separate pPC axes that cumulatively explain 86.1% of the total

variation in life histories (Table 1: pPC1: 43.1%, pPC2: 31.0%, pPC3: 12.0%). Pagel's λ was higher than

our cut-off value of 0.25 (Pagel's λ= 0.31), indicating the phylogenetic signal was significant.

Correcting for body mass did not qualitatively change the results (Appendix: Table S1) and thus we

continued with the pPCA results. Generation time (T), progressive growth (γ), mature life expectancy

(L_ω) and age at maturity (L_α) were loaded positively onto PC1 and mean recruitment success (φ)

negatively (Fig. 2B) (Table 1). Like the low feeding level, as PC1 score increases, individuals increase

their allocation to progressive growth and mature life expectancy at the expense of longer

generation times and later maturity. Therefore, again, PC1 most closely represents the fast-slow

continuum (Fig. 2B). Degree of iteroparity (S), mean recruitment success (φ) and net reproductive

rate (R_0) were loaded positively onto PC2 and retrogressive growth (ρ) negatively, respectively (Fig.

2B) (Table 1). As PC2 score increases, elasmobranchs attain greater lifetime reproductive success

(iteroparity and net reproductive rate) and an increased number of breeding females (recruitment

success), again representing the reproductive strategy axis (Fig. 2B). Only survivorship curve type (H)

was loaded positively onto PC3 (Table 1), so for higher PC3 scores, senescence rates decrease

(decreasing mortality with age).

Comparing the position of each species at each feeding level showed that species move along each

PC axis in particular ways (Fig. 2C). For example, at the low feeding level, species that are fast paced

but have a low life reproductive strategy score (e.g. *Mobula birostris*), increase mostly in life history

speed with increasing feeding level (Fig. 2C: bottom left-hand corner). But, at low feeding level,

species that are fast paced and also have a high reproductive strategy (e.g. *Anoxypristis cuspidata*),

move to a lower reproductive strategy with an increase in feeding level (Fig. 2C: top left-hand

corner). Likewise, species that are slow paced with a low reproductive strategy at low feeding level

(e.g. *Beringraja binoculata*), show a higher reproductive strategy as feeding level increases (Fig. 2C:

bottom right-hand corner). But species that are slow paced with a high reproductive strategy (e.g.

Prionace glauca) tend to show a faster life history speed with increasing feeding level (Fig. 2C: top

right-hand corner). Together, these shifts create an anticlockwise 'whirlpool' effect as species move

across the life history strategy space in response to an increase in feeding level (Fig. 2C).

3.2 Habitat type shapes life history strategies differently for different feeding levels and clades
(objective 3)

For the low feeding level $E(Y)$= 0.6, neither temperature at depth (p= 0.350) nor clade (p= 0.320)
significantly affected species position on the PC1 (fast-slow) axis. Similarly, neither temperature at
depth (p= 0.924) nor clade (p= 0.336) significantly affected species position on the PC2 (reproductive
strategy) axis.

At the high feeding level $E(Y)$= 0.9, phylogenetic clade did not significantly affect PC1 score (fast-slow
continuum) (p= 0.445). There was a significant effect of temperature at depth on PC1 score ($F_{1,149}$=
23.93, adjusted-R^2= 0.13, p= <0.001). PC1 scores decreased in value (and pace of life increased) with
increasing temperature at depth, indicating faster pace of life in warmer waters (Fig. 3A). PC2 scores
(reproductive strategy) decreased with increasing temperature at depth ($F_{1,149}$= 4.86, adjusted-R^2=
0.03, p= 0.029: Fig. 3B). This suggests that across all elasmobranchs, species with lower reproductive
outputs occur in warmer waters. PC2 scores were higher for skates than sharks and rays, indicating
that skates have a high reproductive output ($F_{2,148}$= 6.97, adjusted-R^2= 0.11, p= <0.001, Fig. 2B).

3.3 Life history strategies as predictors of population performance and IUCN status depends on
feeding level (objective 4)

At the low feeding level $E(Y)$= 0.6, population growth rate significantly increased with increasing PC1
scores (fast-slow axis) and increasing PC2 scores (reproductive strategy axis) (Table 2), showing that
slower paced species and species with higher reproductive outputs have higher population growth
rates (Fig. 4A: e.g. *Raja montagui, Leucoraja ocellata, Squalus acanthias*). Resilience to perturbations
(damping ratio) was significantly affected by the interaction between PC1 and PC2 scores (Table 2).
Damping ratio significantly increased with both decreasing PC1 scores and increasing PC2 scores,
showing that faster paced species with higher reproductive outputs have increased resilience to
perturbations (Fig. 4C: e.g. *Scoliodon laticaudus, Urolophus paucimaculatus, Rhizoprionodon acutus*).
There was no significant association of PC1 scores, PC2 scores, or their interaction on the IUCN
category of species in the model (Table 2, Fig. 4E).

At the high feeding level $E(Y)$= 0.9, population growth rate was significantly affected by the
interaction between PC1 scores (fast-slow axis) and PC2 scores (reproductive strategy) (Table 2).
Population growth rate increased with increasing PC2 scores, but this increase was steeper for
species with high, negative PC1 scores (faster species) than for species with high, positive PC1 scores
(slower species) (Fig. 4B: e.g. *Rostroraja eglanteria, Raja microocellata, Raja montagui*). Resilience to
perturbations (damping ratio) was significantly affected by the interaction between PC1 scores and
PC2 scores (Table 2). Resilience to perturbations increased with decreasing PC1 (increasing pace of

life) at a higher rate at low PC2 scores than at higher PC2 scores (Fig. 4D). Overall, faster species with

intermediate reproductive outputs show the highest resilience to perturbations (Fig. 4D: e.g.

Scoliodon laticaudus, Rhizoprionodon acutus, Anoxypristis cuspidata). There was a significant

positive association between PC1 scores (fast-slow axis) and IUCN category, but not with PC2 scores

or their interaction (Table 2). Species with a higher PC1 score (slower pace of life) were more likely

to be in a higher threat category than species with a lower PC1 score (faster pace of life) (Fig. 4F).

4   Discussion

Our aim was to investigate how individual energy budgets, phylogeny and habitat type structure

elasmobranch life history variation, and if this structured variation can predict population

performance, resilience and conservation biology. Elasmobranch life histories are primarily

structured along a fast-slow continuum and reproductive strategy axes at both a low and high

feeding level. Interestingly, we found that as feeding level increased, species moved along the life

history axes in an anticlockwise manner. Regardless, the two life history strategies (the fast-slow

continuum and reproductive strategy) predicted species' population growth rate and demographic

resilience for both feeding levels, and IUCN conservation status in the higher feeding level.

4.1 The impact of feeding level on quantifying and applying life history strategies

Typically, most of the variation in animal life history strategies, including the elasmobranchs studied

here, is captured with two axes: the fast-slow continuum and reproductive strategy axis[2–4]. For many

taxa, including elasmobranchs, these life history strategies are influenced by phylogeny and habitat[1],

likely due to distinct environmental selective pressures. In this study, we also captured the

phenomenon that species move in life history space in an anticlockwise 'whirlpool' manner as their

feeding level changes. Specifically, as feeding levels increase, faster-paced species (on the fast end of

the fast-slow continuum) with low reproductive output speed up, whilst those with high

reproductive output shift to lower reproductive output. Slower-paced species with low reproductive

output increase their reproductive output whilst those with high reproductive output increase their

pace of life. The exact mechanism that underpins this 'whirlpool' effect is not immediately obvious

but possibly related to each species-specific energy budget structure. Such plasticity in life history

strategies can be adaptive when developing organisms adjust their allocation decisions in response

to their local (feeding) environment, following evolved rules that maximise expected fitness in

different ecological conditions[57]. Crucially, the consequences of such feeding level effects can be far-

reaching. Like other studies[2,13], we found that the two life history strategy axes predict species

population performance, resilience and even conservation status. But the plasticity in life history

strategies highlights how predictions on performance, resilience and conservation status *specific to a*

species will depend on its feeding level, and thus, the environment it lives in. One should therefore

be cautious when extrapolating how species-specific life history strategies predict their response to

(climate) change and perturbations in one environment, to another.

Although we find qualitatively the same two life history strategy axes as in other studies (Appendix:

Table S2)[2,5,13,49], there is one notable difference in the life history traits that comprise these life

history strategies. One would expect that investment into development, i.e. high progressive

growth, is characteristic of species with a faster life history speed. But in our analyses, progressive

growth was higher for species with slower life history speeds, not faster (e.g.[2,13]) (Appendix: Table

S2). We surmise that this result is due to high variation in the von Bertalanffy growth rate between

species that drives a positive correlation between generation time and progressive growth (cf.[58]). For

example, the values of progressive growth and generation time change with different values of von

Bertalanffy growth rate and feeding level. The relationship between generation time and progressive

growth for the reef manta ray, *Mobula alfredi,* can be plotted for a range of feeding levels and values

of von Bertalanffy growth rate (Appendix: Fig. S2). For the same von Bertalanffy growth rate value,

generation time increases with decreasing feeding level and progressive growth decreases. At the

same time, generation time decreases with increasing von Bertalanffy growth rate values. The result

is that, at the same feeding level (black solid lines in Appendix: Fig. S2), progressive growth increases

with decreasing values of the von Bertalanffy growth rate, and at the same time, generation time

increases. Because our dataset covers a wide range of von Bertalanffy growth rate values, this would

explain why we found that the slower a species' life history speed, the higher its apparent

progressive growth. These findings highlight the need for careful interpretation of correlations

between life history traits calculated from demographic datasets (our objective 1 and e.g.[1,2,5,13,31]),

and how these are used to predict population characteristics[58].

4.2 Consequences for elasmobranch conservation

For many oceanic species, demographic data are sparce or lacking[59–61]. This often means that the

only measure available to evaluate a species relative risk to perturbations like fishing or

environmental change, is an estimate of its maximum intrinsic rate of population increase, r_{max}[61–64].

Typically, species with higher (or lower) r_{max} values are assumed to have higher (or lower) resilience

to perturbations (cf.[61,65,66]). The intrinsic rate of population increase, r, is related to the population

growth rate ($\lambda = e^{r}$). We found that species with the highest population growth rates do not

necessarily show the highest resilience. In fact, particularly at high feeding level, those species where

populations grow at a relatively low rate, show the highest resilience. This highlights how it is

important to exercise caution when using summary metrics such as r_{max} to predict a species'

resilience [cf.[67]]. Our approach presents a more mechanistic methodology that can be applied to

data-sparce elasmobranchs to understand and predict their resilience and conservation status. For

example, we found that faster-paced elasmobranchs show higher resilience. We also found that

these faster-paced species generally live in warmer waters. Faster paced elasmobranchs might thus

be better able to deal with the current rise in sea temperatures than slower species[68,69]. However,

some species living in warmer, shallower waters are more intrinsically vulnerable to exploitation

(based on their r_{max} values)[19], and may experience higher mortality due to their proximity to densely

populated areas with fisheries[70,71] compared to those living in colder and deeper waters[72]. What is

more, global (sea water) warming is linked to poorer hunting performance in sharks[73]. Warming can

thus reduce the sharks' experienced feeding levels, their population growth rate and ultimately their

resilience. Understanding how elasmobranch life histories, their habitats and feeding translate into

differences in resilience is crucial to improve management decisions and predict elasmobranch

conservation status. Given that the most immediate threat to sharks, skates and rays is

overfishing[72,74,75], further study using an energy budget, demographic approach like ours could

examine the effects of different fisheries mortalities (either by empirical measures or by derived

estimates[76]) on population performance of threatened or highly fished species.

4.3 Conclusion

Many studies have shown that population responses to future environmental change and

perturbations depend on the species-specific life-history strategy. Our analyses reveal that feeding

level can alter a population's life history strategy; such plasticity in life history strategies will impact

how the fast-slow continuum and reproductive strategy framework can be used to predict

population responses to change and perturbations. For elasmobranchs, we also provide strong

support for the expansion of the classical use of maximum intrinsic rate of population increase, r_{max},

to the fast-slow continuum and reproductive strategy framework, and highlight how our approach

can be used to explore different scenarios of (over)fishing to quantify sustainable levels of

exploitation. Further research should explore if variations in environmental conditions fuel plasticity

of life history strategies in a range of taxa, and how these ultimately impact population performance

and responses to change.

Author contributions

Sol Lucas: Conceptualisation, Methodology, Formal analysis, Investigation, Validation, Data curation,
Writing- Original draft preparation, Writing- Review & Editing, Visualisation; Per Berggren: Writing-
Reviewing and Editing, Supervision; Ellen Barrowclift: Methodology, Formal analysis, Writing-
Reviewing and Editing; Isabel M. Smallegange: Conceptualisation, Methodology, Formal analysis,

Investigation, Validation, Data curation, Writing- Original draft preparation, Writing- Review &
Editing, Visualisation, Supervision.

Declaration of competing interest

389 The authors declare no competing interests.

Additional Information

Acknowledgements

396 SL was funded by the Leverhulme Doctoral Scholarship in Behaviour Informatics (DS-2017-015).

References

1. Capdevila, P. *et al.* Longevity, body dimension and reproductive mode drive differences in
aquatic versus terrestrial life-history strategies. *Funct Ecol* **34**, 1613–1625 (2020).

2. Salguero-Gómez, R. *et al.* Fast–slow continuum and reproductive strategies structure plant
life-history variation worldwide. *Proc Natl Acad Sci U S A* **113**, 230–235 (2016).

3. Gaillard, A. J. *et al.* An Analysis of Demographic Tactics in Birds and Mammals. **56**, 59–76
(1989).

4. Gaillard, J. M. *et al.* Life Histories, Axes of Variation in. *Encyclopedia of Evolutionary Biology*
312–323 (2016) doi:10.1016/B978-0-12-800049-6.00085-8.

5. Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal life history
is shaped by the pace of life and the distribution of age-specific mortality and reproduction.
Nat Ecol Evol **3**, 1217–1224 (2019).

6. Stearns, S. C. The Influence of Size and Phylogeny on Patterns of Covariation among Life-
History Traits in the Mammals. *Oikos* **41**, 173–187 (1983).

7. Gadgil, M. *et al.* Life Historical Consequences of Natural Selection. *Am Nat* **104**, 1–24 (1970).

8. Reznick, D. The Structure of Guppy Life Histories: The Tradeoff between Growth and
Reproduction. *Ecology* **64**, 862–873 (1983).

9.   Williams, G. C. Natural Selection, the Costs of Reproduction, and a Refinement of Lack 's
Principle. *Am Nat* **100**, 687–690 (1966).

10.   Gamelon, M. *et al.* Effects of pulsed resources on the dynamics of seed consumer
populations: a comparative demographic study in wild boar. *Ecosphere* **12**, (2021).

11.   Nilsen, E. B. *et al.* A slow life in hell or a fast life in heaven: Demographic analyses of
contrasting roe deer populations. *Journal of Animal Ecology* **78**, 585–594 (2009).

12.   Rademaker, M., van Leeuwen, A. & Smallegange, I. M. Why we cannot always expect life
history strategies to directly inform on sensitivity to environmental change. *Journal of Animal*
Ecology 348–366 (2024) doi:10.1111/1365-2656.14050.

13.   Salguero-Gómez, R. Applications of the fast–slow continuum and reproductive strategy
framework of plant life histories. *New Phytologist* **213**, 1618–1624 (2017).

14.   Romeijn, J. & Smallegange, I. M. Exploring how the fast-slow pace of life continuum and
reproductive strategies structure microorganism life history variation. *bioRxiv* (2022).

15.   Ebert, D. A., Dando, M. & Fowler, S. *Sharks of the World: A Complete Guide*. (Princeton
University Press, 2021).

16.   Nielsen, J. *et al.* Eye lens radiocarbon reveals centuries of longevity in the Greenland shark
(Somniosus microcephalus). *Science (1979)* **353**, 702–704 (2016).

17.   Carrier, J. C., Pratt, H. L. & Castro, J. I. Reproductive biology of elasmobranchs. in *Biology of*
Sharks and Their Relatives (eds. Carrier, J. C., Musick, J. A. & Heithaus, M. R.) 269–286 (CRC
Press, 2004). doi:10.1201/9780203491317.ch10.

18.   Miller, E., Wails, C. N. & Sulikowski, J. It's a shark-eat-shark world, but does that make for
bigger pups? A comparison between oophagous and non-oophagous viviparous sharks. *Rev*
Fish Biol Fish **32**, 1019–1033 (2022).

19.   Barrowclift, E. *et al.* Tropical rays are intrinsically more sensitive to overfishing than the
temperate skates. *Biol Conserv* **281**, (2023).

20.   Pardo, S. A. & Dulvy, N. K. Body mass, temperature, and depth shape the maximum intrinsic
rate of population increase in sharks and rays. *Ecol Evol* **12**, 1–11 (2022).

21.   Gravel, S., Bigman, J. S., Pardo, S. A., Wong, S. & Dulvy, N. K. Metabolism, population growth,
and the fast-slow life history continuum of marine fishes. *Fish and Fisheries* **25**, 349–361
(2024).

22.   Smallegange, I. M., Caswell, H., Toorians, M. E. M. & de Roos, A. M. Mechanistic description
446   of population dynamics using dynamic energy budget theory incorporated into integral
447   projection models. *Methods Ecol Evol* **8**, 146–154 (2017).

448   23.   Sousa, T., Domingos, T., Poggiale, J. C. & Kooijman, S. A. L. M. Dynamic energy budget theory
449   restores coherence in biology. *Philosophical Transactions of the Royal Society B: Biological*
450   *Sciences* **365**, 3413–3428 (2010).

451   24.   Kooijman, S. A. L. M. & Metz, J. A. J. On the dynamics of chemically stressed populations: The
452   deduction of population consequences from effects on individuals. *Ecotoxicol Environ Saf* **8**,
453   254–274 (1984).

454   25.   Ellner, S. P., Childs, D. Z. & Rees, M. *Data-Driven Modelling of Structured Populations: A*
455   *Practical Guide to the Integral Projection Model*. (Springer, Cham, Switzerland, 2016).

456   26.   Smallegange, I. M. & Lucas, S. DEBBIES Dataset to study Life Histories across Ectotherms. *Sci*
457   *Data* **11**, 1–10 (2024).

458   27.   Kooijman, S. A. L. M., Sousa, T., Pecquerie, L., van der Meer, J. & Jager, T. From food-
459   dependent statistics to metabolic parameters, a practical guide to the use of dynamic energy
460   budget theory. *Biological Reviews* **83**, 533–552 (2008).

461   28.   Kooijman, S. A. L. M. *Dynamic Energy and Mass Budgets in Biological Systems*. (Cambridge
462   University Press, 2001). doi:10.1086/420555.

463   29.   The MathWorks Inc. MATLAB. Preprint at https://www.mathworks.com (2022).

464   30.   Revell, L. J. phytools: An R package for phylogenetic comparative biology (and other things).
465   *Methods Ecol Evol* **3**, 217–223 (2012).

466   31.   Paniw, M., Ozgul, A. & Salguero-Gómez, R. Interactive life-history traits predict sensitivity of
467   plants and animals to temporal autocorrelation. *Ecol Lett* **21**, 275–286 (2018).

468   32.   R Core Team. R: A language and environment for statistical computing. *R Foundation for*
469   *Statistical Computing, Vienna, Austria* https://www.r-project.org/ (2023).

470   33.   Chamberlain, S. A. & Szöcs, E. Taxize: Taxonomic search and retrieval in R. *F1000Res* **2**, 1–30
471   (2013).

472   34.   Stein, R. W. *et al.* Global priorities for conserving the evolutionary history of sharks, rays and
473   chimaeras. *Nat Ecol Evol* **2**, 288–298 (2018).

35. Michonneau, F., Brown, J. W. & Winter, D. J. rotl: an R package to interact with the Open Tree
of Life data. *Methods Ecol Evol* **7**, 1476–1481 (2016).

36. Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative data: A test
and review of evidence. *American Naturalist* **160**, 712–726 (2002).

37. Paradis, E. & Schliep, K. Ape 5.0: An environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics* **35**, 526–528 (2019).

38. Revell, L. J. Phylogenetic signal and linear regression on species data. *Methods Ecol Evol* **1**,
319–329 (2010).

39. Jeschke, J. M. & Kokko, H. The roles of body size and phylogeny in fast and slow life histories.
Evol Ecol **23**, 867–878 (2009).

40. Jolicoeur, P., Pirlot, P., Baron, G. & Stephan, H. Brain structure and correlation patterns in
insectivora, chiroptera, and primates. *Syst Biol* **33**, 14–29 (1984).

41. Revell, L. J. Size-correction and principal components for interspecific comparative studies.
Evolution (N Y) **63**, 3258–3268 (2009).

42. Froese, R. & Pauly, D. FishBase. *World Wide Web electronic publication* www.fishbase.org
(2023).

43. Boettiger, C., Lang, D. T. & Wainwright, P. C. Rfishbase: Exploring, manipulating and
visualizing FishBase data from R. *J Fish Biol* **81**, 2030–2039 (2012).

44. Kaiser, H. F. The Application of Electronic Computers to Factor Analysis. *Educ Psychol Meas*
20, 141–151 (1960).

45. IUCN. The IUCN Red List of Threatened Species. *Version 2022-2* https://www.iucnredlist.org
(2022).

46. Faraway, J. J. faraway: Functions and Datasets for Books by Julian Faraway. Preprint at
https://cran.r-project.org/package=faraway (2022).

47. Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *J Open Source Softw*
6, 3139 (2021).

48. Caswell, H. *Matrix Population Models: Construction, Analysis and Interpretation*. (Sinauer
Associates, Sunderland, MA, USA, 2001).

49. Capdevila, P., Stott, I., Beger, M. & Salguero-Gómez, R. Towards a Comparative Framework of
Demographic Resilience. *Trends Ecol Evol* **35**, 776–786 (2020).

50. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York, 2016).

51. Slowikowski, K. ggrepel: Automatically Position Non-Overlapping Text Labels with 'ggplot2'. *R*
package version 0.9.3 https://github.com/slowkow/ggrepel (2023).

52. Attali, D. & Baker, C. ggExtra: Add Marginal Histograms to 'ggplot2', and More 'ggplot2'
Enhancements. (2023).

53. Hijmans, R. J. raster: Geographic Data Analysis and Modeling. R package version 3.6-21
https://rspatial.org/raster (2023).

54. RStudio Team. RStudio: Integrated Development Environment for R. *RStudio, PBC, Boston,*
MA http://www.rstudio.com/ (2023).

55. Wickham, H., François, R., Henry, L. & Müller, K. dplyr: A Grammar of Data Manipulation.
https://dplyr.tidyverse.org (2022).

56. Müller, K. & Wickham, H. tibble: Simple Data Frames. https://tibble.tidyverse.org/ (2023).

57. Mcnamara, J. M. & Houston, A. I. State-dependent life histories. *Nature* **380**, 215–221 (1996).

58. van Noordwijk, A. J. & de Jong, G. Acquisition and Allocation of Resources: Their Influence on
Variation in Life History Tactics. *Am Nat* **128**, 137–142 (1986).

59. Bradshaw, C. J. A., Mollet, H. F. & Meekan, M. G. Inferring population trends for the world's
largest fish from mark-recapture estimates of survival. *Journal of Animal Ecology* **76**, 480–489
(2007).

60. Temple, A. J. *et al.* Life-history, exploitation and extinction risk of the data-poor Baraka's
whipray (Maculabatis ambigua) in small-scale tropical fisheries. *J Fish Biol* **97**, 708–719
(2020).

61. Pardo, S. A. *et al.* Growth, productivity, and relative extinction risk of a data-sparse devil ray.
Sci Rep **6**, 1–10 (2016).

62. Myers, R. A. & Fowlow, P. S. Maximum population growth rates and recovery times for
Atlantic cod, Gadus morhua. *Fishery bulletin* **95**, 762–772 (1997).

63. Myers, R. A., Bowen, K. G. & Barrowman, N. J. Maximum reproductive rate of fish at low
population sizes. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 2404–2419 (1999).

64.   Dulvy, N. K. *et al.* Methods of assessing extinction risk in marine fishes. *Fish and Fisheries* **5**,
533   255–276 (2004).

534   65.   Pardo, S. A., Kindsvater, H. K., Reynolds, J. D. & Dulvy, N. K. Maximum intrinsic rate of
535   population increase in sharks, rays, and chimaeras: The importance of survival to maturity.
536   *Canadian Journal of Fisheries and Aquatic Sciences* **73**, 1159–1163 (2016).

537   66.   Cortés, E. Perspectives on the intrinsic rate of population growth. *Methods Ecol Evol* **7**, 1136–
538   1145 (2016).

539   67.   Smallegange, I. M., Flotats Avilés, M. & Eustache, K. Unusually Paced Life History Strategies of
540   Marine Megafauna Drive Atypical Sensitivities to Environmental Variability. *Front Mar Sci* **7**,
541   1–18 (2020).

542   68.   Rosa, R. *et al.* Early-life exposure to climate change impairs tropical shark survival.
543   *Proceedings of the Royal Society B: Biological Sciences* **281**, (2014).

544   69.   Osgood, G. J., White, E. R. & Baum, J. K. Effects of climate-change-driven gradual and acute
545   temperature changes on shark. *Journal of Animal Ecology* **90**, 2547–2559 (2021).

546   70.   Letessier, T. B. *et al.* Remote reefs and seamounts are the last refuges for marine predators
547   across the Indo-Pacific. *PLoS Biol* **17**, 1–20 (2019).

548   71.   Letessier, T. B. *et al.* Divergent responses of pelagic and benthic fish body-size structure to
549   remoteness and protection from humans. *Science (1979)* **383**, 976–982 (2024).

550   72.   Dulvy, N. K. *et al.* Overfishing drives over one-third of all sharks and rays toward a global
551   extinction crisis. *Current Biology* **31**, 4773-4787.e8 (2021).

552   73.   Pistevos, J. C. A., Nagelkerken, I., Rossi, T., Olmos, M. & Connell, S. D. Ocean acidification and
553   global warming impair shark hunting behaviour and growth. *Sci Rep* **5**, 1–10 (2015).

554   74.   Pacoureau, N. *et al.* Half a century of global decline in oceanic sharks and rays. *Nature* **589**,
555   567–571 (2021).

556   75.   Worm, B. *et al.* Global shark fishing mortality still rising despite widespread regulatory
557   change. *Science (1979)* **383**, 225–230 (2024).

558   76.   Smith, S. E., Au, D. W. & Show, C. Intrinsic rebound potentials of 26 species of Pacific sharks.
559   *Mar Freshw Res* **49**, 663–678 (1998).

77.   Caswell, H. Sensitivity Analysis: Matrix Methods in Demography and Ecology. in *Demographic*

Research Monographs Chapter 10 (Springer Open, 2019).

78.   Tuljapurkar, S., Gaillard, J. M. & Coulson, T. From stochastic environments to life histories and

back. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**, 1499–1509

(2009).

Figure 1. *Workflow of parameterising a DEB-IPM using the DEBBIES database[26], using the derived life history traits to plot*
elasmobranch life history strategies in a Principal Components Analysis (PCA). Species biogeography and phylogeny were
used to estimate position on PCA axes, and the PC scores were used to predict conservation status, population growth rates
and demographic resilience. White arrows indicate parameterisation of models or calculation of traits. Blue arrows
represent metrics used to predict life history strategies and green arrows indicate the use of life history strategies to make
predictions of conservation status and demography.

Table 1. Demographic quantities and their loadings onto the Principal Components axes. To calculate life history traits, we discretised each DEB-IPM (Eqn 1) by dividing the length domain Ω into 200 very small-width discrete bins, resulting in a matrix **A** of size $m \times n$, where $m = n = 200$, and which dominant eigenvalue equals λ. Mean lifetime reproductive success R_0 is the dominant eigenvalue of the matrix $\mathbf{F} = \mathbf{V}(\mathbf{I} - \mathbf{GS})^{-1}$, where **I** is the identity matrix and $\mathbf{V} = \mathbf{DR}$, with **D** as the parent-offspring association, **R** the reproduction, **G** the growth and **S** the survival matrix[48]; this gives generation time $T = \log(R_0)/\log(\lambda)$ (Caswell 2001[48]). The mean life expectancy, η_e, is calculated as $\eta_e = \mathbf{1}^T\mathbf{N}$, where **1** is a vector of ones of length m and **N** is the fundamental matrix $\mathbf{N} = (\mathbf{I} - \mathbf{S})^{-1}$. The longevity of an individual of length L is η_L, which means we can calculate age at sexual maturity $L_\alpha = \eta_{L_b}$ and mature life expectancy $L_\varpi = \eta_{L_p}$ so that $\eta_e = L_\alpha + L_\varpi$ (Caswell 2018[77], eqn 4.21). l_x is the probability of surviving to age at least x, and m_x is the average fertility of age class x (cf. Tuljapurkar et al. 2009[78]), $\overline{\mathbf{G}}$ is the mean of G, $\overline{\mathbf{V}}$ is the mean of V, and i and j are the row and column entries of the matrix, respectively. The vital rates included in the studied set of demographic quantities (progressive growth γ, retrogressive growth ρ, and sexual reproduction φ) were averaged across the columns j (the length bins), weighted by the relative contributions of each stage at stationary equilibrium. For example, to calculate mean sexual reproduction φ, we summed the values in the columns j of the **V** matrix and multiplied each φ_{ij} by the corresponding jth element w_j of the stable stage distribution w, calculated as the right eigenvector of **A**.

Demographic quantity	Symbol	Definition	Equation	E(Y)= 0.6			E(Y)= 0.9		
		Feeding level							
		Number of species in analysis		63			151		
				PC1	PC2	PC3	p-PC1	p-PC2	p-PC3
Generation time	T	Number of days required for the individuals of a population to be fully replaced by new ones	$T = \dfrac{\log(R_0)}{\log(\lambda)}$	**0.532**	-0.215	-0.007	**0.942**	-0.234	0.147
Survivorship curve	H	Keyfitz' entropy ($H < 1$ denotes increasing mortality rate with age, $H > 1$ denotes decreasing mortality rate with age, $(H)= 1$ denotes a constant mortality rate with age) (Keyfitz & Caswell 2005).	$H = -\dfrac{\sum_{x=0}^{x=\eta_e} \log(l_x)l_x}{\sum_{x=0}^{x=\eta_e} l_x}$	0.281	0.110	**0.772**	0.086	0.083	**0.987**
Age at maturity	L_α	Number of days that it takes an average individual in the population to become reproductive	$L_\alpha = \eta_{L_b}$	0.274	0.048	**-0.538**	**0.828**	0.384	-0.228
Progressive growth	γ	Mean probability of growing to a larger length across the length domain Ω.	$\gamma = \sum_i^m \overline{\mathbf{G}}_{i,j}\vert_{i<j}$	**0.511**	0.006	-0.007	**0.869**	-0.004	0.043
Retrogressive growth	ρ	Mean probability of growing to a smaller length across the length domain Ω.	$\rho = \sum_i^m \overline{\mathbf{G}}_{i,j}\vert_{i>j}$	-0.161	**-0.462**	0.042	-0.138	**-0.898**	-0.066
Mean recruitment success	φ	Mean per-capita number of recruits across the length domain Ω.	$\varphi = \sum_i^m \overline{\mathbf{V}}_{i,j}$	-0.210	**0.521**	0.043	**-0.720**	**0.643**	0.092
Degree of iteroparity	S	Coefficient of variation in age at reproduction	$S = \overline{m}_x/\sigma(m_x)$	-0.045	**0.451**	0.169	-0.228	**0.734**	0.059
Net reproductive rate	R_0	Mean number of recruits produced during the mean life expectancy of an individual in the population.	$R_0 = \sum_{x=0}^{x=\eta_e} l_x m_x$	0.097	**0.474**	-0.287	0.344	**0.870**	-0.028
Mature life expectancy	L_ϖ	Number of days from the mean age at maturity (L_α) until the mean life expectancy (η_e) of an individual in the population.	$L_\varpi = \eta_{L_p}$	**0.469**	0.162	-0.023	**0.899**	0.264	0.187
Axis				Fast-slow	Repro	Longevity	Fast-slow	Repro	Longevity
Proportion of variance explained				42.1%	31.2%	14.9%	43.1%	31.0%	12.0%
Cumulative proportion of variance explained				42.1%	73.3%	88.2%	43.1%	74.1%	86.1%
Kaiser criterion				3.79	2.80	1.34	15.05	7.78	1.17
Pagel's lambda (for the pPCA)					-			0.303	

Figure 2. *(p)PCA plots for the 0.6 (A) and 0.9 (B) feeding levels, representing 63 and 151 species respectively. Marginal boxplots of the split by clade (Shark, Skate and Ray) are included for each principal component. Species present in A are highlighted in yellow in B. Labelled arrows correspond to the variables used in the PCA analysis and show the magnitude and direction of the loadings. Red arrows align with the fast-slow continuum, blue arrows align with the reproductive axis and the orange arrow does not align with either of the primary two principal components. The dashed blue/red arrow aligns with both the Slow-Fast continuum and the reproductive axis in plot B. (C) the change standardised (z-scored) PC scores plotted for the 63 species in both low (white points) and high (black points) feeding level. The change in life history strategies between low and high feeding level creates a 'whirlpool' effect.*

Figure 3. *For the high feeding level E(Y)= 0.9. (A): The relationship between temperature-at-depth and PC1 (fast slow axis). Warmer temperatures relate to negative PC1 scores (faster paces of life). (B) PC2 score decreased for increasing temperature-at-depth. Reproductive output was greater for species inhabiting cooler waters. F-statistics, p-values and R^2 for each analysis are in each plot window. The grey band around the fitted models shows the 95% confidence intervals.*

Table 2. General linear models for population growth rate (λ) and demographic resilience (damping ratio, ξ), as well as ordinal regression models for IUCN conservation status, as a response to PC1 (fast-slow axis) and PC2 (reproductive strategy) scores, for both the low (*E(Y)*=0.6) and high (*E(Y)*=0.9) feeding levels.

Variable	Axes	Estimate	df	Std. Error	p	R²
		Feeding Level E(Y)= 0.6				
λ ~ *PC1 * PC2*	Intercept	1.010		0.008	**<0.001**	0.674
	PC1	0.042	1	0.004	**<0.001**	
	PC2	0.031	1	0.005	**<0.001**	
	PC1:PC2	0.002	1	0.003	0.402	
	Error term	0.067	59			
ξ ~ *PC1 * PC2*	Intercept	1.164		0.011	**<0.001**	0.590
	PC1	-0.034	1	0.006	**<0.001**	
	PC2	0.053	1	0.007	**<0.001**	
	PC1:PC2	-0.009	1	0.004	**0.029**	
	Error term	0.092	59			
as.factor(IUCN) ~ PC1 * PC2	PC1	0.019		0.116	0.869	0.012
	PC2	-0.128		0.133	0.336	
	PC1:PC2	0.088		0.073	0.227	
		Feeding Level E(Y)= 0.9				
λ ~ *PC1 * PC2*	Intercept	1.215		0.007	**<0.001**	0.826
	PC1	-0.068	1	0.007	**<0.001**	
	PC2	0.167	1	0.007	**<0.001**	
	PC1:PC2	-0.082	1	0.007	**<0.001**	
	Error term	0.086	147			
ξ ~ *PC1 * PC2*	Intercept	1.237		0.012	**<0.001**	0.544
	PC1	-0.151	1	0.013	**<0.001**	
	PC2	-0.041	1	0.013	**0.001**	
	PC1:PC2	0.037	1	0.013	**0.004**	
	Error term	0.151	147			
as.factor(IUCN) ~ PC1 * PC2	PC1	0.483		0.151	**0.001**	0.030
	PC2	-0.095		0.150	0.526	
	PC1:PC2	-0.160		0.150	0.287	

R² values are adjusted-R² for the linear models and pseudo-R² for the ordinal regression on conservation status. McFadden's pseudo R² was calculated for the ordinal regression models using the equation $R^2 = 1 - \frac{residual\ deviance}{null\ deviance}$.

Figure 4. *Overlays on the PCA figures for both feeding levels: E(Y)= 0.6 (A,C,E) and E(Y)= 0.9 (B,D,F). Population growth rate (A,B), damping ratio (C,D), IUCN conservation status (E,F).*