1 Bacterial Protein Signatures Identified in *Porphyromonas gingivalis* Containing-Autophagic
2 Vacuoles Reveal Evolutionary History Between Oral Red/Orange Complex Bacteria and Gut
3 Resident Bacteria
4

Ayana Paul[a], Bridgette Wellslager[a], Maddie Williamson[a], Özlem Yilmaz [a, b, c]

[a] Department of Oral Health Sciences, Medical University of South Carolina, Charleston, South

Carolina 29425, USA

[b] Department of Microbiology and Immunology, Medical University of South Carolina,

Charleston, South Carolina 29425, USA

[c] Hollings Cancer Center, Medical University of South Carolina, 86 Jonathan Lucas Street,

Charleston, SC 29425, USA

Running Head: Oral and Gut Bacteria Share Evolutionary History for Virulence

*Corresponding Author(s): Özlem Yilmaz (yilmaz@musc.edu) and Ayana Paul

(ayana.paul@emory.edu)

Lead Contact: Özlem Yilmaz (yilmaz@musc.edu)

Abstract

Oral bacterial species have evolved in humans to enable the colonization of a diversity of

commensals and opportunistic pathogens like *Porphyromonas gingivalis*. Due to the

communalistic survival mechanism shared amongst oral bacteria and gut bacteria, these bacteria

have likely evolved together to establish in the human oral cavity and transfer genetic

information. Our liquid-chromatography with tandem-mass-spectrometry (LC-MS-MS) analysis

has revealed protein signatures, Elongation Factor Tu, RagB/SusD nutrient uptake outer

membrane protein and DnaK, specifically from *P. gingivalis*-containing autophagic vacuoles

isolated from the infected primary gingival epithelial cells. Interestingly, our Mass-Spectrometry

analysis reported similar proteins from closely related oral bacteria, *Tannerella forsythia* and

Prevotella intermedia. In our phylogenetic study of these key protein signatures, we have

established that pathogenic oral bacteria share extensive relatedness to each other and gut

resident bacteria. We have shown that virulence factors, Elongation Factor Tu and DnaK, there

are several structural similarities and conservation with proteins from pathogenic oral bacteria.

We have also shown that the RagB/SusD nutrient uptake These findings may highlight the

shared virulence mechanisms not only amongst oral bacterial pathogens but gut bacteria and

their evolution in the human host.

Keywords: *P. gingivalis:* red/orange complex bacteria; oral and gut microbiome; mucosal

colonizers; virulence, adaptation

1. Introduction

A microbiome can be defined as the ecological community of pathogenic and commensal microorganisms that inhabit a particular area of the body. The oral microbiome is likely one of the most influential microbiomes of the human body as the oral cavity is a point of entry of microorganisms into other areas of the body and thus may play a role in the pathologies outside of this area (Willis and Gabaldón, 2020, Dewhirst et al., 2010). This microbiome is substantially diverse and can be divided into several niches including the buccal mucosa, throat, tongue, saliva, gingiva, and sub-gingiva (Lee et al., 2021, Faran Ali and Tanwir, 2012). The modern human oral microbiota is believed to have evolved from changes diet, environment, hygiene, health, genetics, lifestyle over human evolution and industrialization (Shaw et al., 2017). Characterization of the modern oral microbiome has resulted in the organization of microbial complexes of varying association with healthy and diseased states of the oral cavity (Avila et al., 2009). Blue, purple, green and yellow complex bacteria are understood to colonize and proliferate in the oral cavity in the early stages of oral microbiome development. A number of bacteria are commonly found in healthy oral microbiomes such as, *Streptococcus, Staphylococcus, Veillonella* and *Eubacteria* (Avila et al., 2009, Wilson, 2004). Conversely, bacteria that have been characterized as pathogenic and associated with diseased states are *Prevotella* species, *Tannerella forsythia, Treponema denticola* and *Porphyromonas gingivalis* (Haffajee and Socransky, 1994, Spooner et al., 2016, Haffajee and Socransky, 2005). These bacteria, commonly characterized as orange and red complex bacteria, colonize the oral microbiome last and enable the proliferation of other pathogenic bacteria and induction of host proinflammatory responses. Figure 1 briefly demonstrates how the oral bacterial complexes are established and how these oral complexes can lead to disease pathologies and translocation in the

human body. Poor oral hygiene and antibiotic use are important contributing factors to oral

dysbiosis (Dewhirst et al., 2010, Jiao et al., 2014). Thus, dysbiosis of the oral microbiome can

result in numerous oral diseases such as gingivitis, dental carries and periodontitis (Jiao et al.,

2014, Simón-Soro et al., 2013). The number of people with oral conditions in general have also

increased globally from 1990 to 2015 with disability adjusted life years (DALYS) increasing by

64% (Kassebaum et al., 2017). The organization, pathogenicity and dysbiosis of the oral

microbiome are pertinent to our understanding of oral disease outcomes.

The oral microbiome is a unique ecological space in that is physically connected to other areas of

the body, especially the gut microbiome (Figure 1) (Narengaowa et al., 2021). Due to this

accessibility, it is likely that the oral microbiota shares similar evolutionary histories with the gut

microbiome. In fact, in a study conducted by The Human Microbiome Project Consortium, it

was found 125 bacterial species consistently present in both saliva and stool from 470 people

(Human Microbiome Project, 2012). Despite the consistencies in disease associations and

physical connections in the microbiomes, the relationship between oral and gut bacteria is

incredibly understudied. Studying the co-evolution of these bacteria and their molecular

mechanisms could potentially shine a light on novel holistic therapeutic strategies for non-

communicable microbially associated diseases.

P. gingivalis, is a Gram (-), obligate anaerobic bacterium and is a member of the oral red

complex within the oral cavity. This bacterium demonstrates a unique ability to persist and

spread within gingival epithelial cells (GECs), which serve as a microbial protective barrier in

the oral mucosa (Lee et al., 2020a, Nakayama and Ohara, 2017, Mohanty et al., 2019, Jenkinson

and Lamont, 2005, Atanasova et al., 2016, Yilmaz et al., 2006). *P. gingivalis* can avoid the host

immune system and induce dysbiosis of the oral microbiome and possesses a number of

virulence factors to aid in its pathogenesis such as lipopolysaccharides, gingipains, out

membrane vacuoles as well as pili (Jia et al., 2019, Chen et al., 2001a, Zhang et al., 2021). Our

lab has begun to characterize the intracellular life of *P. gingivalis* and has revealed that the

microorganism can evade a variety of cellular mechanisms designed to eliminate pathogens

(Roberts et al., 2017a, Lee et al., 2020b, Yao et al., 2010) (Spooner et al., 2014). One key

mechanism that *P. gingivalis* modulates is xenophagy in order to avoid phagolysosomal fusion

while inside the Endoplasmic Reticulum (ER)/LC3 rich double-membrane vacuoles for which

P. gingivalis can replicate and survive in (Lee et al., 2018b, Wellslager et al., 2024). In fact, *P.*

gingivalis is most strongly associated with chronic periodontitis, extensive inflammation of the

periodontal tissue leading to tooth loss, and demonstrates unique virulence mechanisms for

subduing host immune response, positioning as a major pathogen within the oral microbiome

(Hajishengallis et al., 2012, Lee et al., 2018a, Lee et al., 2020c, Lee et al., 2020d, Roberts et al.,

2017b, Yilmaz, 2008, Lee and Yilmaz, 2020, Atanasova and Yilmaz, 2015, Nakhjiri et al., 2001)

Our recent study into the bacterial proteomics involved in the vacoular life of this bacterium

using Mass Spectrometry has prompted study of several bacterial protein signatures (Lee et al.,

2018b, Wellslager et al., 2024, Chowdhury et al., 2024). Elongation Factor Tu, DnaK and SusD

nutrient uptake outer membrane protein signatures, belonging to pathogenic oral bacteria

Prevotella intermedia and *T. forsythia* were identified inside of *P. gingivalis* containing

autophagic vacuoles which were selectively isolated for *P. gingivalis* from the infected primary

GECs as we recently described (Wellslager et al., 2024). These bacteria are both closely related

P. gingivalis and of the identical or proximal oral complex (Mohanty et al., 2019). It is not fully

understood what functional role these proteins play in the survival of *P. gingivalis* in the

autophagic vacuolesvacoules. Further, the evolution of established pathogenic oral complexes

and their shared survival mechanisms in host niches is not well understood. Lastly, the extent of

relatedness of these protein signatures from *P. gingivalis*, related pathogenic oral and gut

bacteria is not known as well. In this study, our objective was to determine the extent of

relatedness of proteins identified in *P. gingivalis* contained vacoules across oral complex bacteria

to understand the evolution, maintenance, and establishment of this microbiome and their

established bacterial pathogens. We have found that proteins identified from red and orange

complex bacteria *T. forsythia* and *P. intermedia* are evolutionary related with bacteria from not

only the oral microbiome but the gut microbiome based on phylogenetic analysis. Many of these

bacteria have well characterized virulence mechanisms involving the identified protein

signatures. The predictive secondary structures of these proteins belonging to these oral

pathogens offer additional support for homology and conserved structure. Understanding the

evolutionary dynamics of these bacterial species may aid in our understanding of the virulence

and survival mechanisms amongst oral bacteria as well as gut bacteria. Extensive study into the

protein homology shared across these two microbiomes may contribute to the growing interest in

the oral-gut axis and its connection to systemic diseases.

2. Materials and Methods

The Magnetic Isolation of *P. gingivalis*-Containing Autophagosomes

P. gingivalis ATCC 33277 strain was incubated in 5 μg/mL Lipobiotin (LB) in PBS overnight at

4°C. As we recently described, labelled *P. gingivalis* was then incubated with MagCellect

Streptavidin Ferrofluid (R&D Systems) at a concentration of $7–14 \times 10^8$ particles per

2×10^8 bacteria for 20 min at 4°C (Wellslager et al., 2024, Chowdhury et al., 2024). To

determine the importance of HSp27 on *P. gingivalis*-containing autophagosomal integrity, GECs

were plated at a density of 1.4×10^6 in 100 mM dishes and were cultured until they reached

~80% confluence. Following siRNA transfection, redox-related treatments, and *P. gingivalis*

infection, *P. gingivalis*-containing autophagosomes were then isolated as described in (*5*). In

brief, GECs were washed with Buffer A (15 mm HEPES buffer with 20 mm sucrose,

50 mm MgCl$_2$, and protease inhibitors containing 0.02% ethylenediaminetetraacetic acid

(EDTA)) to remove non-ingested bacteria, were trypsinized, and resuspended in Buffer A

combined with Benzonase Nuclease (25 U/mL; Novagen) and Cytochalasin D (5 µg/mL;

Sigma). Following resuspension, cell membranes were disrupted and *P. gingivalis*-

containing autophagosomes were released via sonicating GECs on ice via a ThermoFisher

Qsonica Sonicator Q500 (FisherSci) at an amplitude of 75% for 10 s. GECs were then

centrifuged at 300 g for 2 min between sonication steps, after which the supernatant was

removed and replaced with 500 µL of Buffer A. After the third sonication, the samples were

left on ice for 10 min. The supernatants were then removed and incubated at 37°C for

5 min. Following incubation, the samples were placed in a Magnetic Rack (Bio-Rad) for 30 min

prior to repeated manual rinsing of the magnetic fraction with Buffer A at a flow rate of 250

µL/min for 15 min to ensure purity. Finally, the autophagosome-containing fractions were

collected and centrifuged at 15,000 g for 12 min, before the supernatant was decanted and the

autophagosomes were resuspended in 100 µL of HEPES buffer combined with protease

inhibitors.

Mass Spectrometry Sample preparation

Cells were lysed in 9M urea, 50 mM Tris pH 8, and 100 units/mL Pierce Universal Nuclease

(ThermoScientific cat. # 88700) and the concentration of protein was measured using a BCA

assay (ThermoScientific cat. # 23225). Protein was LysC and trypsin (Sigma cat. # T6567)

digested at 37°C for 18 hours, and the resulting peptides were desalted using C18 Stage tips

(Pierce cat. # 84850)

Liquid Chromatography and Mass Spectrometry Data Acquisition Parameters

Peptides were separated and analyzed on an EASY nLC 1200 System (ThermoFisher) in-line

with the Orbitrap Fusion Lumos Tribrid mass spectrometer (ThermoFisher) with instrument

control software v. 4.2.28.14. 2 ug of tryptic peptides were pressure loaded at 1,180 bar and

peptides were separated on a C18 reversed phase column (Acclaim PepMap RSLC, 75 µm x 50

cm (C18, 2 µm, 100 Å) ThermoFisher) using a gradient of 5% to 40% B in 180 min (Solvent A:

5% acetonitrile/0.1% formic acid; Solvent B: 80% acetonitrile/ 0.1% formic acid) at a flow rate

of 300 nL/min with a column heater set to 50C.

Mass spectra were acquired in data-dependent mode with a high resolution (60,000)

FTMS survey scan, mass range of m/z 375-1575, followed by tandem mass spectra (MS/MS) of

the most intense precursors with a cycle time of 3 s. The automatic gain control target value was

4.0e5 for the survey MS scan. Fragmentation was performed with a precursor isolation window

of 1.6 m/z, a maximum injection time of 22 ms, and HCD collision energy of 35%.

Monoisotopic-precursor selection was set to "peptide". Apex detection was not enabled.

Precursors were dynamically excluded from resequencing for 20 sec and a mass tolerance of 10

ppm. Precursor ions with charge states that were undetermined, 1, or >5 were excluded. The

assay was done with at least 6-replicates pooled together during LC-MS/MS analysis.

Mass Spectrometry Data Processing

Protein identification and quantification were extracted from raw LC-MS/MS data using the

MaxQuant platform v.1.6.3.3 with the Andromeda database searching algorithm and label free

quantification (LFQ) algorithm (Cox J, Mann M. MaxQuant enables high peptide identification

rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat

Biotechnol. 2008 Dec;26(12):1367-72; Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann

M. Accurate proteome-wide label-free quantification by delayed normalization and maximal

peptide ratio extraction, termed MaxLFQ. Mol Cell Proteomics. 2014 Sep;13(9):2513-26;

Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based

shotgun proteomics. Nat Protoc. 2016 Dec;11(12):2301-2319). Data were searched against a

human Uniprot reference database UP0000005640 with 74,468 proteins (August, 2019), a

Porphyromonas gingivalis database, and a database with other common bacterial species. The

false discovery rate (FDR), determined using a reversed database strategy, was set at 1% at the

protein and peptide level. Fully tryptic peptides with a minimum of 7 residues were required

including cleavage between lysine and proline. Two missed cleavages were permitted. The LFQ

feature was on with "Match between runs" enabled for those features that had spectra in at least

one of the runs. The "stabilize large ratios" feature was enabled. The first search was performed

with a 20 ppm mass tolerance; after recalibration, a 4.5 ppm tolerance was used for the main

search. A minimum ratio counts of 2 was required for LFQ protein quantification with at least

one unique peptide. Parameters included static modification of cysteine with carbamidomethyl

and variable N-terminal acetylation and oxidation of methionine.

The protein groups text file from the MaxQuant search results was processed in Perseus v.

1.6.7.0 (Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J. The

Perseus computational platform for comprehensive analysis of proteomics data. Nat Methods.

2016 Sep;13(9):731-40). Identified proteins were filtered to remove proteins only identified by a

modified peptide, matches to the reversed database, and potential contaminants.   The MQ

normalized LFQ intensities for each biological replicate were log2 transformed. Gene ontology

annotations were added from a human protein database downloaded from Uniprot with

associated GO terms and protein properties of interest.

Homologous Protein Search and Phylogenetic Tree Construction

All protein signatures identified via Mass Spectrometry were subjected to a NCBI Protein

BLAST search using their associated NCBI Protein ID. Each BLAST search was performed

under the Position-specific iterated blast algorithm. A total of 20 homologous bacterial proteins

ranging from Firmicutes, Bacteroidetes, Actinobacteria and Proteobacteria (including 1

eukaryotic protein if possible) for each protein search were recorded for analysis based on high

percent identities (>50%) and low E-value (closest to 0) (Sayers et al., 2022). However, a

homologous (significant protein identity) *P. gingivalis* protein could not be identified for *T.*

forsythia's Ragb/SusD protein. Instead *P.gingivalis'* RagB (Protein ID: U2JPV9) was included

in the phylogenetic analysis of *T*. forsythia's Ragb/SusD protein. Phylogenetic analysis of each

protein was performed by first constructing multiple sequence alignments using the ClustalW

program within the MEGA X application. Each ClustalW alignment was run through the

MEGAX Find Best DNA/Protein Model program in MEGA X in order to apply the best

evolutionary history model. The Le Gascuel 2008 model was used to construct phylogenetic tree

from *P. intermedia'*s and *T. forsythia*'s EF-Tu and *P. intermedia*'s DnaK (Le and Gascuel,

2008). Whelan And Goldman model was used to construct phylogenetic tree from *T. forsythia*'s

SusD/RagB protein (Whelan and Goldman, 2001). The maximum likelihood trees were finally

generated and exported from 500 bootstrap replicates in the MEGA X application (Tamura et al.,

2021, Kumar et al., 2018, Pattengale et al., 2010).

Functional Regions and Conserved Site Identification

To search for functional conserved sites within homologous proteins, the EXPASY-PROSITE

tool and InterPROscan database was utilized (de Castro et al., 2006, Blum et al., 2021).

Homologous proteins sequences were uploaded to a CLUSTAL Omega program available

through UniProt to generate alignments for annotation (Sievers et al., 2011). Additional known

and putative binding substrates for each protein were found from literature review (Sun et al.,

2022, Harvey et al., 2019, Tomoyasu et al., 2012, Candela et al., 2010).

Protein Functional Interaction Prediction

In order to identify known functional interactions within the context of *P. gingivalis*, the

STRING database was accessed under confidence mode. *P. gingivalis'* EF-Tu and DnaK were

searched within the STRING database. *B. fragilis'* NanU protein and *T. forsythia's* RagB/SuSD

protein was searched by amino acid sequence within the STRING database. A protein of 98.8%

identity (BF1797) was used to construct a STRING network for *T. forsythia's* RagB/SuSD

protein. Only functional interaction networks with a confidence score above 0.7 were displayed

and exported (Szklarczyk et al., 2019).

Assessing the structural homology of select proteins

Published data for the crystallized structures of *P. gingivalis's EF-Tu and T. forsythia's*

RagB/SuSD nutrient uptake outer membrane protein) and thus are no established models

available in public depositories. The secondary structure of these proteins was predicted using a

predictive structural biology program RosettaTTAFold under Robetta (Yang et al., 2020). Out of

the five models generated, the model with the least angstrom error estimates across the protein's

sequence was selected for analysis. Model 3 for *P. gingivalis'* predicted EF-Tu structure and

Model 2 for *T. forsythia's* predicted RagB/SusD family protein was selected from the Robetta

provided models. Characterized models for *E.coli*'s EF-Tu (PDB ID:1EFC) and its EF-Tu in

complex with *A. baumanni* DsbA (PDB ID: 4P3Y), *P. gingivalis'* RagB (PDB ID: 5CX8) and *B.*

fragilis NanU (PBD ID: 4L7T) were acquired from Protein Data Bank (PDB). All models were

visualized, annotated and superimposed in UCSF ChimeraX (Goddard et al., 2018, Pettersen et

al., 2004, Pettersen et al., 2021).

3. Results

3.1 Bacterial protein quantities of Elongation Factor Tu (EF-Tu), chaperone protein dnaK and a

RagB/SusD increase over a 24 hour P. gingivalis infection

Primary GECs were infected for up to 24 hours with *P. gingivalis* ATCC 33277. After *P.*

gingivalis containing vacoules were isolated from infected cells at 3 hour, 6 hour and 24 hour

post infection time points, numerous protein signatures belonging to *P. gingivalis, T. forsythia*

and *P. intermedia* were identified via mass spectrometry. Label free-quantification (LFQ) was

used to measure the amount of specific proteins present from within the vacoules. According to

the reported LFQ values of these proteins, the synthesis of homologous intracellular *P. gingivalis*

proteins increased over the course of the 24 hour infection (Figure 2). This study will discuss

bacterial proteins Elongation Factor Tu (EF-Tu), chaperone protein dnaK and a RagB/SusD

nutrient uptake outer membrane protein and their shared properties amongst oral and gut bacteria

(Table 1). Mass Spectrometry analysis specifically identified protein signatures residing in

Domain 2 of EF-Tu, a nucleotide binding site superfamily domain of dnaK and a SusD-like N

terminal domain of the RagB/SusD nutrient uptake outer membrane protein.

3.2 Pathogenic oral bacteria, T. forsythia's Elongation Factor Tu shares conserved properties

with neighboring pathogenic bacteria within the oral microbiome

According to mass spectrometry results, EF-Tu protein signatures belonging to oral bacteria *P.*

gingivalis, *T. forsythia*, and *P. intermedia* were identified in the vacuoles isolated from infected

primary GEC's. EF-Tu is a conserved GTPase protein that is intracellularly abundant and

essential for protein synthesis across both eukaryotes and prokaryotes (Yikilmaz et al., 2014,

Stark et al., 1997). Thus, a number of homologous EF-Tu proteins were identified ranging from

bacteria that reside in both oral and gut microbiomes but primarily from bacteria resident in the

oral cavity (Supplementary Table 1). In order to assess the extent of relatedness of *T. forsythia*'s

EF-Tu with other oral and gut bacteria, phylogenetic analysis was performed. *T. forsythia*'s EF-

Tu protein was found to be most closely related to neighboring red oral complex bacteria,

P. gingivalis from the maximum likelihood tree. Additionally, several homologous EF-Tu

proteins from orange oral complex bacteria such as *P. intermedia* and *Fusobacterium nucleatum*

had close relation to members of the red oral complex. Red and orange complex bacteria also

appear to be extensively related to gut bacteria such as *E. coli, Citrobacter feundii* and

Parabacteriotes distasonis (Figure 3A). In order to further analyze the relatedness of the *T.*

forsythia's EF-Tu with other bacteria, protein sequences from the same clade were aligned for

annotation and SNP detection. A conserved binding site for the ligand GTP was identified via

InterPro and denoted in an alignment of these homologous EF-Tu proteins from this closely

related gut and oral bacteria (Figure 3A). After extensive literature review, several accessory

proteins beyond oral or gut bacterial EF-Tu's main ligand, GTP, were found to bind to EF-Tu to

serve accessory functions (Harvey et al., 2019). These proteins include bacterial DsbA, human

holotransferrin, human mucin and human nucleolin (Figure 3B). In summary, these results

provide support for the putative and ancestral functional properties of EF-Tu across pathogenic

oral bacteria and related bacterial species that reside in different regions of a human host.

3.3 Orange complex bacteria, P. intermedia's EF-Tu is most related to human gut resident

bacteria

A total of 20 homologous proteins were identified via BLAST search ranging from both oral and

gut bacteria, most of which were gut resident bacteria. The most homologous proteins to *P.*

intermedia's EF-Tu protein signature were from other orange complex bacteria and bacteria

residing in the gut microbiome (e-value closest to 0) (Supplementary Table 2). In concordance

with *T. forsythia*'s EF-Tu, EF-Tu in pathogenic oral bacteria are often extensively related to each

other (Figure 4). However, in contrast to *T. forsythia*, EF-Tu from *P. intermedia* is primarily

related to commensal gut bacteria based on phylogenetic analysis. As in Figure 3A,

P. intermedia's and the bacteria that reside within the same clade demonstrate a conserved

binding site for GTP indicating a conserved primary function within these closely related

bacteria. In conclusion, oral bacteria such as *P. intermedia* may share more proteomic

similarities with members of different microbiomes which may serve as evidence for genetic

transfer between microbiomes.

3.4 P. gingivalis' EF-Tu structurally homologous to moonlighting EF-Tu from E. coli to

moonlighting which may enable pro-survival functions

In order to further assess the relatedness of oral bacterial EF-Tu proteins to members of the gut

microbiome, we sought to analysis these proteins structurally. Both *P. gingivalis* and *T.*

forsythia's EF-Tu structure were predicted using a predictive modeling program Robetta to

generate possible models with 88% confidence. After these predicted structure were

superimposed with *E.coli*'s EF-Tu in its GDP bound state, the two proteins appear to have

extensive similarity with a percent homology of 78% (Figure 5A). In addition to there being

evidence that these species are related based phylogenetic analysis of these proteins, these

proteins shared conserved domains as established in Figure 5B. In a study investigating the

biochemical interactions of *Acinetobacter baumannii*'s DsbA enzyme, researchers found that *E.*

coli's EF-Tu can be isolated in complex with *A. baumannii*'s DsbA. To investigate the accessory

binding similarities between these two bacteria, we sought to predict if *P. gingivalis'* EF-Tu

could potentially bind to *A. baumannii*'s DsbA. After superimposing the predicted structure of *P.*

gingivalis with *E.coli*'s EF-Tu bound to *A. baumanni*'s DsbA, it appears that *P. gingivalis* EF-Tu

may share the same accessory binding (Figure 5C). To summarize, the moonlighting potential of

EF-Tu protein from pathogenic oral bacteria can be predicted via structural homology and this

shared moonlighting potential may enable pro-survival functions yet to be studied.

3.5 T. forsythia's RagB/SusD family nutrient uptake outer membrane protein is mostly

homologous to gut Parabacteriotes

In addition to EF-Tu, a protein signature from the RagB/SusD family was identified from red

oral complex bacteria, *T. forsythia*. Rag/SusD proteins are starch binding proteins that allow

bacteria to uptake nutrients in their microbial environment and are localized on the outer

membrane of *Bacteroidetes* (Glenwright et al., 2017). Phylogenetic analysis revealed that *T.*

forsythia's Rag/SusD protein is most similar to proteins from other *Parabacteriotes* species from

the gut. However, this protein was found to be related to other oral bacteria such as orange oral

complex bacteria, *P. intermedia* (Supplementary Figure 3, Figure 6). Amongst the closely related

proteins, there is a conserved starch binding protein and a lipoprotein lipid attachment domain

(Figure 6). This includes a putative haloacid dehalogenase (HAD) protein from P. *gingivalis*

which is understood to be involved in metabolism and homeostasis (Tribble et al., 2006). Taken

together, this protein derived from oral bacteria, *T. forsythia*, appears to share functional

characteristics to gut resident bacteria which may highlight additional conserved or inherited

nutrient uptake pathways.

3.6 Structural similarities in T. forsythia's RagB/SusD family protein may enable starch

utilization and antigenicity

In order to predict the function of the SusD family protein signature identified from *T, forsythia*,

we sought to compare the structures of SusD proteins and RagB protein to the protein of interest.

After superimposing *P. gingivalis'* RagB and *T. forsythia's* RagB/SusD family protein predicted

structure with *B. fragilis'* NanU, all three homologous proteins appeared to have similar

structures as well. However, despite *P. gingivalis'* RagB protein and *T. forsythia*'s SusD/RagB

protein sharing similar structures they exhibit low percent shared identity (Figure 7A-C). Lastly,

the predicted structure of RagB/SusD from *T. forsythia* was generated with 72% confidence and

appeared to have a wide range of electrostatic potential and a positively charged region outside

the putative binding pocket (Figure 7D). After literature review, it was found that the structure of

P. gingivalis' RagB is extensively similar to SusD of gut bacteria, *Bacteroides thetaiotaomicron*

and *Tannerella forsythia's* NanU (Goulas et al., 2016). In summary, this data indicates that *P.*

gingivalis' associated antigen and the nutrient uptake protein exhibits a substantial structural and

functional similarity as well as conservation in both oral and gut bacteria.

3.7 STRING analysis confirms conserved protein functions and predicts putative functions in

oral bacteria

In order to determine the known functional interactions of oral bacterial SusD family protein, the

STRING database tool was utilized. The known functional interactions of this RagB/SusD

protein from *T. forsythia* was found to include connections with a number of putative,

hypothetical proteins, a PBS lyase HEAT-like repeat protein and a TonB linked outer membrane

protein (Figure 8A). *Bacteroides fragilis*, a gut bacterium that utilizes a SusD like protein called

NanU, exhibits similar functional interactions to *T. forsythia*'s RagB/SusD protein

(Supplementary Table 6, Figure 8B). In summary, T. *forsythia's* SusD family protein may enable

survival and nutrient uptake functions similar to the gut bacteria, B. fragilis.

3.8 Pathogenic Oral bacteria and Gut Bacterial share similar properties in chaperone protein

dnaK

A chaperone dnaK protein signature from orange oral complex bacteria *P. intermedia* was also

amongst the proteins identified within the *P. gingivalis* contained vacoules. Similar to bacterial

EF-Tu, this chaperone protein is essential, and several homologous proteins were identified

across oral and gut bacteria (Supplementary Table 4). The most closely related proteins to *P.*

*gingivali*s' dnaK were from oral bacteria, *P. intermedia* and *P. nigrescens* as well as gut bacteria

such as *E. coli*. In addition, red oral complex bacteria *T. denticola* and a number of gut bacteria

were shown to have extensive relation to this protein (Figure 9A). Of the homologous proteins

most closely related to *P. gingivalis'* dnaK, there are numerous conserved. There are conserved

domains shared amongst orange and red complex bacteria such as the conserved Heat Shock

protein 70 family signature 1. However, known functional interactions for *P. gingivalis'* dnaK

involve other heat shock proteins, chaperone proteins and proteases (Figure 9B). After literature

review of chaperone DnaK utility in oral and gut bacteria, it was found that this protein has

functional interactions with proteins such as human laminin and plasminogen (Figure 9C). Taken

together, these results indicate that oral and gut bacteria have extensive conservation in structure

and sequence which may allow for additional pathogenic functions observed in bacteria like *B.*

lactis (Candela et al., 2010).

4. Discussion

There is little known on the extent of shared survival and pathogenic functions amongst these

neighboring bacteria. Understanding the proteomics of one bacterium within the context of one

microbiome may aid in our understanding of a related species. Further, assessing the homology

of shared proteins may offer potential therapeutic strategies for various clinically relevant oral

and gut associated chronic illnesses and pathogens.

Based on phylogenetic analysis of oral bacterial EF-Tu protein signatures identified in *P.*

gingivalis contained vacoules, *P. gingivalis'* EF-Tu is closely related to neighboring orange and

red oral complex bacteria. As shown in Figure 3A and Figure 4, EF-Tu proteins from oral

pathogens, *T. forsythia* or *P. intermedia* are extensively related to pathogenic and commensal

oral bacteria. In addition, this study has shown that the evolution of host-adapted bacterial

proteomics is not limited to the oral microbiota but extends into the gut bacterial community. For

example, EF-Tu proteins from commensal bacteria, *Porphyromonas catoniae*, and commensal

gut bacteria, *Bifidobacterium pseudocatenulatum,* appear to be related based on their EF-Tu

(Gonzalez-Vazquez et al., 2022) . This may suggest genetic transfer between not only oral

bacteria but gut resident bacteria which is likely due to the physical connection between the oral

cavity and human intestinal tract (Roberts and Kreth, 2014, Guilloux et al., 2021). Moreover,

some of these related bacteria can be isolated from not only humans but other animal

microbiomes which may suggest potential zoonotic transfer of opportunistic bacteria (Wylensek

et al., 2020, Companys et al., 2021, Lagkouvardos et al., 2016) Interestingly, this cytosolic

enzyme has enabled similar extracellular virulence and survival functions in various

microbiomes of the human host and the potential accessory substrates of bacterial EF-Tu utilized

in pathogens are numerous and can be appreciated from Figure 3B (Harvey et al., 2019, Widjaja

et al., 2017, Premkumar et al., 2014, Archambaud et al., 2005, Torres et al., 2020). The

evolutionary relatedness between bacteria that use EF-Tu as a moonlighting protein and other

bacteria may imply that there are uncharacterized extracellular functions for EF-Tu in other oral

bacterial pathogens.

In addition to bacterial EF-Tu, the conservation of SusD/RagB outer membrane protein in both

oral and gut microbiomes is likely that expression of this protein was transferred between

bacteria. In a study examining the evolution of *P. gingivalis*' RagB protein, they found evidence

of long term selection pressure through likelihood ratio tests which is likely as a result of

horizontal gene transfer from other Bacteroides bacteria (Su et al., 2010). Our structural analysis

as seen in Figure 6B further revealed homology between the RagB/SusD family protein in *T.*

forsythia with Chain A of *B. fragilis* NanU protein, a glycan utilization protein. Furthermore,

string analysis of *T. forsythia's* RagB/SusD family proteins and *B. fragilis* NanU protein

confirms its utilization in *T. forsythia* as a starch outer membrane binding protein. It was

similarly suggested by Hanley et al that the RagAB operon of *P. gingivalis* arose via horizontal

gene transfer from a foreign source, potentially from one comprising another component of the

periodontal microflora (Hanley et al., 1999). It has also been shown that PgRagB exhibits

structural similarity with the *B. thetaiomicron* SusD protein and *T. forsythia* NanU which are

both involved in starch and sialic acid uptake (Goulas et al., 2016). Therefore, it is possible that

P. gingivalis utilizes this conserved starch binding domain in its RagB to acquire sialic acid in a

similar manner to that of other species in both the gut and oral microbiomes for later use in

several virulence mechanisms. Moreover, *P. gingivalis* utilizes a metabolic protein, HAD

phospahatse SerB653, to enable entry into GECs. Given that *T. forsythia's* SusD/RagB protein is

homologous with a putative HAD protein from *P. gingivalis* it possible that these metabolic and

nutritional similarities shared amongst oral bacteria may enable virulence and invasiveness

(Chen et al., 2001b, Tribble et al., 2006). Overall, oral and gut bacteria appear to utilize similar

nutrient uptake systems which further supports a shared evolutionary history.

Furthermore, phylogenetic analysis of DnaK in *P. gingivalis* contained vacoules revealed close

relationships to DnaK of other orange complex oral bacteria such as *Prevotella intermedia* and

Prevotella nigrescens. As previously mentioned, one of the exhibited conserved sites among

460 these homologous proteins was the Heat Shock Protein (HSP) 70 family signature 1. It was

461 hypothesized that the secretion of DnaK may occur due to the ability of *P. intermedia* to escape

462 the oral cavity and colonize in distant parts of the body (Karched et al., 2022), which may be a

463 strong indicator of potential horizontal gene transfer of the DnaK protein between the oral and

464 gut microbiome. Despite possessing several unique characteristics from other known bacterial

465 HSP70s, *P. gingivalis* DnaK was found to behave similarly to *P. intermedia* DnaK in its ability

466 to cooperate with DnaJ and GrpE of the *E. coli* chaperone system (Yoshida et al., 1999). The

467 ability of this protein to cooperate with the DnaJ and GrpE proteins was exhibited further via

468 string network analysis as seen in Figure 8B. Given the homology between these bacteria and the

469 analogous chaperone functioning it is possible that this role and response of DnaK is conserved

470 between species of pathogenic oral and gut bacteria.

472 Despite there being evidence that oral and gut bacterial protein may share moonlighting

473 functions as a moonlighting factor and mode of adhesion extracellularly, it is important to note

474 that there are limitations to making novel mechanistic conclusions based solely on

475 bioinformatics. Further experimentation is required to conclude whether the binding activities

476 and resulting function of these binding events with EF-Tu can be applied to other bacteria. It is

477 possible that the identification of EF-Tu outside *P. gingivalis* may be an artifact due to

478 upregulated protein synthesis genes while inside the host autophagic vacuoles. In addition, it is

479 understood that *P. gingivalis* can adapt to oral bacterial species communities including *F.*

480 *nucleatum* and *S. gordonii* by up regulating protein synthesis genes (Kuboniwa et al., 2009).

481 Based on the string network generated in Figure 3B, there is only evidence of *P. gingivalis*' EF-

482 Tu interacting with ribosomal subunits and the EF-Tu protein which is also required for protein

synthesis. The caution must be applied to *P. gingivalis'* RagB protein that is homologous to the

SusD family protein shared amongst. Although it may have structural homology with SusD

proteins the known function of this protein remains associated with immunogenicity (Potempa et

al., 2021).

5. Conclusions

Overall, these findings suggest conserved survival and potentially pathogenic mechanisms

between oral bacteria, and that this conservation may extend outside the oral microbiome into the

gut microbiome. The protein signatures identified within the *P. gingivalis* infected GECs not

only characterize the intracellular survival of *P. gingivalis* but also highlight the broad

similarities it shares with other pathogenic bacteria. The extent of relatedness between

pathogenic oral complex bacteria and pathogenic gut bacteria may provide evidence for the

evolution of shared pathogenicity between the two microbiomes. *P. gingivalis* and closely related

bacteria use a multitude of virulence mechanisms to enable persistence and invasion within host

(Yilmaz, 2008, Lee et al., 2020c, Lee et al., 2018b, Harvey et al., 2019, Yilmaz et al., 2008, Han,

2015, Dashper et al., 2011, Chiu et al., 2017). Further study into the shared survival and

virulence pathways amongst oral and gut bacteria may support identification of effective

therapeutics and our understanding of established bacteria within microbiomes.

Funding Source This work was primarily supported by funding from the NIH grants

R01DE030313, F31DE032273, R25GM113278 and T32DE017551.

Declaration of Competing Interests: Authors declare no conflicts of interest

Acknowledgments

The authors would like to thank Dr. Lauren Ball, MUSC proteomics core director and MUSC

core facilities were supported in part by Hollings Cancer Center Support Grant (P30 CA138313)

and the MUSC Digestive Disease Research Cores Center (P30 DK123704).

Figures and Tables

FIGURE 1 The Human oral-gut axis can be organized into different core complexes that are

linked via physical connections. A schematic diagram demonstrating the organization of key

bacteria within the oral microbiome and its physical connection to external microbiomes such as

the gut microbiome. Bacteria are not limited to their prospective niches and have physical access

to different areas of the body.

FIGURE 2 Specific protein signatures for A) Elongation Factor Tu, B) RagB/SusD Family

Protein, and C) DnaK, which were originally characterized in other bacterial species and were

persistently present in *P. gingivalis*-specific autophagosomes. These autophagosomes were

isolated from human primary gingival epithelial cells (GECs) infected with wild-type *P.*

gingivalis. GECs were infected with *P. gingivalis* for 3 h, 6 h, or 24 h. The assay was done with

at least 6-replicates pooled together during LC-MS/MS analysis.

TABLE 1 List of protein signatures of interest identified from vacuoles containing *P. gingivalis*.

Protein IDs were reported from Mass Spectrometry analysis after identifying specific protein

signatures belonging in the biological samples. Homologous *P. gingivalis* proteins containing the

protein signatures are displayed as well.

FIGURE 3 Pathogenic oral complex bacteria share putative accessory binding sites in bacterial

EF-TU. (A) Maximum likelihood tree based on *T. forsythia*'s EF-Tu and 20 homologous

proteins from oral/gut bacterial species and a control eukaryotic species (n=21). A sequence

alignment of *T. forsythia* EF-Tu and closely related species is displayed to the right of the clade.

A conserved region shared amongst these species is highlighted in yellow. Bootstrap percentages

are hidden if under 50. Red complex bacteria are denoted by a red circle, orange complex

bacteria are denoted by a orange circle and yellow complex bacteria are denoted by a yellow

circle. Bacteria that can colonize the human gut are denoted by a triangle and bacteria that

colonize other parts of the human host are denoted by a square. The black star specifies

P. gingivalis. (B) Schematic diagram of EF-Tu locus and potential ligands binding to bacterial

EF-Tu. GTP is displayed above the sequence alignment binding to the Guanine Nucleotide

Binding Domain.

FIGURE 4 *P. intermedia's* EF-Tu shares a conserved GTP binding site with gut and oral

bacteria. (A) Maximum likelihood tree based on *P. intermedia's* 20 homologous proteins from

oral/gut bacterial species and a control eukaryotic species (n=21). A sequence alignment of *P.*

intermedia and closely related species is shown from one clade to the right. A conserved region

shared amongst these species is highlighted in yellow. Bootstrap percentages are hidden if under

50. Red complex bacteria are denoted by a red circle, orange complex bacteria are denoted by an

orange circle and yellow complex bacteria are denoted by a yellow circle. Bacteria that can

colonize the human gut are denoted by a triangle and bacteria that colonize other parts of the

human host are denoted by a square. The black star specifies interactions of *P. gingivalis'* EF-Tu.

(B) Representative string networks displaying the known functional interactions of *P. gingivalis'*

EF-Tu.

FIGURE 5 *P. gingivalis'* EF-Tu is structurally similar to *E.coli's* EF-Tu. (A) Superimposed

predicted models of *P. gingivalis'* EF-Tu (Orange Ribbon) with *E.coli's* EF-TU (Purple Ribbon).

(B) A sequence alignment of P. gingivalis', T. forsythia's and *E. coli's* EF-Tu protein. A

conserved region shared amongst these species is highlighted in yellow. (C) *P. gingivalis'*

predicted EFTu (orange ribbon) bound to *A. baumanni's* DsbA (red ribbon).

FIGURE 6 *T. forsythia's* SusD/RagB like protein share conserved regions with gut bacteria and

P. gingivalis' RagB. (A) Maximum likelihood tree based on *T. forsythia's* SusD/RagB protein

EF-Tu and 19 homologous proteins from oral/gut bacterial species and a control eukaryotic

species (n=20). A sequence alignment of *T. forsythia* and closely related species is shown from

one clade is shown to the right. A conserved region shared amongst these species is highlighted

in yellow. Bootstrap percentages are hidden if under 50. Red complex bacteria are denoted by a

red circle, orange complex bacteria are denoted by an orange circle and yellow complex bacteria

are denoted by a yellow circle. Bacteria that can colonize the human gut are denoted by a triangle

and bacteria that colonize other parts of the human host are denoted by a square. The black star

specifies *P. gingivalis'* RagB protein.

FIGURE 7 *P. gingivalis'* RagB and *T. forsythia's* SusD family protein is structurally similar to

Bacillus fragilis' NanU protein**.** (A) Superimposed models of *P. gingivalis'* Chain A of RagB

(Sky Blue Ribbon) and *T. forsythia's* predicted Rag/SusD like protein (Beige Ribbon). The

starch binding domain is highlighted in dark blue. (B) Superimposed models of *P. gingivalis'*

Chain A of RagB (Beige Ribbon) with *B. fragilis* Chain A of NanU (Sky Blue Ribbon). The

starch binding domain is highlighted in dark blue. (C) *T. forsythia's* predicted Rag/SusD like

protein (Beige Ribbon) superimposed with *B. fragilis* Chain A of NanU (Sky Blue Ribbon). The

conserved starch binding domain is highlighted in dark blue. (D) A ribbon structure and three-

dimensional predicted secondary structure image of *T. forsythia's* EF-Tu (Orange ribbon)

depicting the predicted range of surface hydrophobicity. The range in electrostatic potential was

calculated using Coulomb's law. The putative starch binding domain is highlighted in dark blue.

FIGURE 8 *T. forsythia's* SusD family protein and *B. fragilis* NanU share similar functional

interactions**.** (A) A string network generated from *T.* forsythia's Rag/SusD family protein with

the Rag/SusD like protein at the center. (B) A string network generated from *B. fragilis'* NanU

protein with the NanU protein at the center. The thickness of the "edges" between each node

indicates the level of confidence of the interaction between each protein.

FIGURE 9 *P. intermedia* DnaK is conserved in pathogenic oral bacteria and gut bacteria. (A)

Maximum likelihood tree based on *P. intermedia's* EF-Tu and 20 homologous proteins from

oral/gut bacterial species and a control eukaryotic species (n=21). A sequence alignment of

T. forsythia and closely related species is shown from one clade is shown to the right. A

conserved region shared amongst these species is highlighted in yellow. Bootstrap percentages

are hidden if under 50. Red Complex bacteria are denoted in red circle and orange complex are

denoted in orange circle. Bacteria that can colonize the human gut are denoted by a triangle and

bacteria that colonize other parts of the human host are denoted by a square. The black star

specifies *P. gingivalis'*. (B) Schematic diagram of DnaK locus and potential and known ligands.

Ligands highlighted in gray indicate there is no literature available for the exact binding on

DnaK.

References

ARCHAMBAUD, C., GOUIN, E., PIZARRO-CERDA, J., COSSART, P. & DUSSURGET, O. 2005. Translation elongation factor EF-Tu is a target for Stp, a serine-threonine phosphatase involved in virulence of Listeria monocytogenes. *Mol Microbiol,* 56**,** 383-96.

ATANASOVA, K., LEE, J., ROBERTS, J., LEE, K., OJCIUS, D. M. & YILMAZ, Ö. 2016. Nucleoside-Diphosphate-Kinase of P. gingivalis is Secreted from Epithelial Cells In the Absence of a Leader Sequence Through a Pannexin-1 Interactome. *Scientific Reports,* 6**,** 37643.

ATANASOVA, K. R. & YILMAZ, Ö. 2015. Prelude to oral microbes and chronic diseases: past, present and future. *Microbes and Infection,* 17**,** 473-483.

AVILA, M., OJCIUS, D. M. & YILMAZ, O. 2009. The oral microbiota: living with a permanent guest. *DNA Cell Biol,* 28**,** 405-11.

BLUM, M., CHANG, H. Y., CHUGURANSKY, S., GREGO, T., KANDASAAMY, S., MITCHELL, A., NUKA, G., PAYSAN-LAFOSSE, T., QURESHI, M., RAJ, S., RICHARDSON, L., SALAZAR, G. A., WILLIAMS, L., BORK, P., BRIDGE, A., GOUGH, J., HAFT, D. H., LETUNIC, I., MARCHLER-BAUER, A., MI, H., NATALE, D. A., NECCI, M., ORENGO, C. A., PANDURANGAN, A. P., RIVOIRE, C., SIGRIST, C. J. A., SILLITOE, I., THANKI, N., THOMAS, P. D., TOSATTO, S. C. E., WU, C. H., BATEMAN, A. & FINN, R. D. 2021. The InterPro protein families and domains database: 20 years on. *Nucleic Acids Res,* 49**,** D344-D354.

CANDELA, M., CENTANNI, M., FIORI, J., BIAGI, E., TURRONI, S., ORRICO, C., BERGMANN, S., HAMMERSCHMIDT, S. & BRIGIDI, P. 2010. DnaK from Bifidobacterium animalis subsp. lactis is a surface-exposed human plasminogen receptor upregulated in response to bile salts. *Microbiology (Reading),* 156**,** 1609-1618.

CHEN, T., NAKAYAMA, K., BELLIVEAU, L. & DUNCAN, M. J. 2001a. Porphyromonas gingivalis gingipains and adhesion to epithelial cells. *Infection and immunity,* 69**,** 3048-3056.

CHEN, W., LAIDIG, K. E., PARK, Y., PARK, K., YATES, J. R., 3RD, LAMONT, R. J. & HACKETT, M. 2001b. Searching the Porphyromonas gingivalis genome with peptide fragmentation mass spectra. *Analyst,* 126**,** 52-7.

CHIU, K.-H., WANG, L.-H., TSAI, T.-T., LEI, H.-Y. & LIAO, P.-C. 2017. Secretomic Analysis of Host–Pathogen Interactions Reveals That Elongation Factor-Tu Is a Potential Adherence Factor of Helicobacter pylori during Pathogenesis. *Journal of Proteome Research,* 16**,** 264-273.

CHOWDHURY, N., WELLSLAGER, B., LEE, H., GILBERT, J. L. & YILMAZ, Ö. 2024. Glutamate is a key nutrient for *Porphyromonas gingivalis* growth and survival during intracellular autophagic life under nutritionally limited conditions. *bioRxiv,* 2024.07.08.602514.

COMPANYS, J., GOSALBES, M. J., PLA-PAGA, L., CALDERON-PEREZ, L., LLAURADO, E., PEDRET, A., VALLS, R. M., JIMENEZ-HERNANDEZ, N., SANDOVAL-RAMIREZ, B. A., DEL BAS, J. M., CAIMARI, A., RUBIO, L. & SOLA, R. 2021. Gut Microbiota Profile and Its Association with Clinical Variables and Dietary Intake in Overweight/Obese and Lean Subjects: A Cross-Sectional Study. *Nutrients,* 13.

DASHPER, S. G., SEERS, C. A., TAN, K. H. & REYNOLDS, E. C. 2011. Virulence factors of the oral spirochete Treponema denticola. *Journal of dental research,* 90**,** 691-703.

DE CASTRO, E., SIGRIST, C. J., GATTIKER, A., BULLIARD, V., LANGENDIJK-GENEVAUX, P. S., GASTEIGER, E., BAIROCH, A. & HULO, N. 2006. ScanProsite: detection of PROSITE signature matches and ProRule-associated functional and structural residues in proteins. *Nucleic Acids Res,* 34**,** W362-5.

DEWHIRST, F. E., CHEN, T., IZARD, J., PASTER, B. J., TANNER, A. C., YU, W. H., LAKSHMANAN, A. & WADE, W. G. 2010. The human oral microbiome. *J Bacteriol,* 192**,** 5002-17.

FARAN ALI, S. M. & TANWIR, F. 2012. Oral microbial habitat a dynamic entity. *Journal of Oral Biology and Craniofacial Research,* 2**,** 181-187.

GLENWRIGHT, A. J., POTHULA, K. R., BHAMIDIMARRI, S. P., CHOREV, D. S., BASLE, A., FIRBANK, S. J., ZHENG, H., ROBINSON, C. V., WINTERHALTER, M., KLEINEKATHOFER, U., BOLAM, D. N. & VAN DEN BERG, B. 2017. Structural basis for nutrient acquisition by dominant members of the human gut microbiota. *Nature,* 541**,** 407-411.

GODDARD, T. D., HUANG, C. C., MENG, E. C., PETTERSEN, E. F., COUCH, G. S., MORRIS, J. H. & FERRIN, T. E. 2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci,* 27**,** 14-25.

GONZALEZ-VAZQUEZ, R., ZUNIGA-LEON, E., TORRES-MARAVILLA, E., LEYTE-LUGO, M., MENDOZA-PEREZ, F., HERNANDEZ-DELGADO, N. C., PEREZ-PASTEN-BORJA, R., AZAOLA-ESPINOSA, A. & MAYORGA-REYES, L. 2022. Genomic and Biochemical Characterization of Bifidobacterium pseudocatenulatum JCLA3 Isolated from Human Intestine. *Microorganisms,* 10.

GOULAS, T., GARCIA-FERRER, I., HUTCHERSON, J. A., POTEMPA, B. A., POTEMPA, J., SCOTT, D. A. & GOMIS-RUTH, F. X. 2016. Structure of RagB, a major immunodominant outer-membrane surface receptor antigen of Porphyromonas gingivalis. *Mol Oral Microbiol,* 31**,** 472-485.

GUILLOUX, C. A., LAMOUREUX, C., BEAURUELLE, C. & HERY-ARNAUD, G. 2021. Porphyromonas: A neglected potential key genus in human microbiomes. *Anaerobe,* 68**,** 102230.

HAFFAJEE, A. D. & SOCRANSKY, S. S. 1994. Microbial etiological agents of destructive periodontal diseases. *Periodontol 2000,* 5**,** 78-111.

HAFFAJEE, A. D. & SOCRANSKY, S. S. 2005. Microbiology of periodontal diseases: introduction. *Periodontol 2000,* 38**,** 9-12.

HAJISHENGALLIS, G., DARVEAU, R. P. & CURTIS, M. A. 2012. The keystone-pathogen hypothesis. *Nat Rev Microbiol,* 10**,** 717-25.

HAN, Y. W. 2015. Fusobacterium nucleatum: a commensal-turned pathogen. *Current opinion in microbiology,* 23**,** 141-147.

HANLEY, S. A., ADUSE-OPOKU, J. & CURTIS, M. A. 1999. A 55-kilodalton immunodominant antigen of Porphyromonas gingivalis W50 has arisen via horizontal gene transfer. *Infect Immun,* 67**,** 1157-71.

HARVEY, K. L., JAROCKI, V. M., CHARLES, I. G. & DJORDJEVIC, S. P. 2019. The Diverse Functional Roles of Elongation Factor Tu (EF-Tu) in Microbial Pathogenesis. *Front Microbiol,* 10**,** 2351.

HUMAN MICROBIOME PROJECT, C. 2012. Structure, function and diversity of the healthy human microbiome. *Nature,* 486**,** 207-214.

JENKINSON, H. F. & LAMONT, R. J. 2005. Oral microbial communities in sickness and in health. *Trends Microbiol,* 13**,** 589-95.

JIA, L., HAN, N., DU, J., GUO, L., LUO, Z. & LIU, Y. 2019. Pathogenesis of Important Virulence Factors of Porphyromonas gingivalis via Toll-Like Receptors. *Frontiers in cellular and infection microbiology,* 9**,** 262-262.

JIAO, Y., HASEGAWA, M. & INOHARA, N. 2014. The Role of Oral Pathobionts in Dysbiosis during Periodontitis Development. *Journal of Dental Research,* 93**,** 539-546.

KARCHED, M., BHARDWAJ, R. G., QUDEIMAT, M., AL-KHABBAZ, A. & ELLEPOLA, A. 2022. Proteomic analysis of the periodontal pathogen Prevotella intermedia secretomes in biofilm and planktonic lifestyles. *Sci Rep,* 12**,** 5636.

KASSEBAUM, N. J., SMITH, A. G. C., BERNABÉ, E., FLEMING, T. D., REYNOLDS, A. E., VOS, T., MURRAY, C. J. L. & MARCENES, W. 2017. Global, Regional, and National Prevalence, Incidence, and Disability-Adjusted Life Years for Oral Conditions for 195 Countries, 1990–2015: A Systematic Analysis for the Global Burden of Diseases, Injuries, and Risk Factors. *Journal of Dental Research,* 96**,** 380-387.

KUBONIWA, M., HENDRICKSON, E. L., XIA, Q., WANG, T., XIE, H., HACKETT, M. & LAMONT,
R. J. 2009. Proteomics of Porphyromonas gingivalis within a model oral microbial community.
BMC Microbiol, 9**,** 98.
KUMAR, S., STECHER, G., LI, M., KNYAZ, C. & TAMURA, K. 2018. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol,* 35**,** 1547-1549.
LAGKOUVARDOS, I., PUKALL, R., ABT, B., FOESEL, B. U., MEIER-KOLTHOFF, J. P., KUMAR,
N., BRESCIANI, A., MARTINEZ, I., JUST, S., ZIEGLER, C., BRUGIROUX, S., GARZETTI,
D., WENNING, M., BUI, T. P., WANG, J., HUGENHOLTZ, F., PLUGGE, C. M., PETERSON,
D. A., HORNEF, M. W., BAINES, J. F., SMIDT, H., WALTER, J., KRISTIANSEN, K.,
NIELSEN, H. B., HALLER, D., OVERMANN, J., STECHER, B. & CLAVEL, T. 2016. The
Mouse Intestinal Bacterial Collection (miBC) provides host-specific insight into cultured
diversity and functional potential of the gut microbiota. *Nat Microbiol,* 1**,** 16131.
LE, S. Q. & GASCUEL, O. 2008. An improved general amino acid replacement matrix. *Mol Biol Evol,*
25**,** 1307-20.
LEE, J., ROBERTS, J. S., ATANASOVA, K. R., CHOWDHURY, N. & YILMAZ, Ö. 2018a. A novel
kinase function of a nucleoside-diphosphate-kinase homologue in Porphyromonas gingivalis is
critical in subversion of host cell apoptosis by targeting heat-shock protein 27. *Cell Microbiol,* 20**,**
e12825.
LEE, J., SPOONER, R., CHOWDHURY, N., PANDEY, V., WELLSLAGER, B., ATANASOVA, K.,
EVANS, Z. & YILMAZ, O. 2020a. In Situ Intraepithelial Localizations of Opportunistic
Pathogens, Porphyromonas gingivalis and Filifactor alocis, in Human Gingiva. Current Research
in Microbial Sciences. *Current Research in Microbial Sciences,* 1**,** 7-17.
LEE, J. S., CHOWDHURY, N., ROBERTS, J. S. & YILMAZ, O. 2020b. Host surface ectonucleotidase-
CD73 and the opportunistic pathogen, Porphyromonas gingivalis, cross-modulation underlies a
new homeostatic mechanism for chronic bacterial survival in human epithelial cells. *Virulence,*
11**,** 414-429.
LEE, J. S., CHOWDHURY, N., ROBERTS, J. S. & YILMAZ, Ö. 2020c. Host surface ectonucleotidase-
CD73 and the opportunistic pathogen, Porphyromonas gingivalis, cross-modulation underlies a
new homeostatic mechanism for chronic bacterial survival in human epithelial cells. *Virulence,*
11**,** 414-429.
LEE, J. S., SPOONER, R., CHOWDHURY, N., PANDEY, V., WELLSLAGER, B., ATANASOVA, K.
R., EVANS, Z. & YILMAZ, Ö. 2020d. In Situ Intraepithelial Localizations of Opportunistic
Pathogens, Porphyromonas gingivalis and Filifactor alocis, in Human Gingiva. *Current Research
in Microbial Sciences,* 1**,** 7-17.
LEE, J. S. & YILMAZ, Ö. 2020. Key Elements of Gingival Epithelial Homeostasis upon Bacterial
Interaction. *Journal of Dental Research,* 100**,** 333-340.
LEE, K., ROBERTS, J. S., CHOI, C. H., ATANASOVA, K. R. & YILMAZ, O. 2018b. Porphyromonas
gingivalis traffics into endoplasmic reticulum-rich-autophagosomes for successful survival in
human gingival epithelial cells. *Virulence,* 9**,** 845-859.
LEE, Y.-H., CHUNG, S. W., AUH, Q. S., HONG, S.-J., LEE, Y.-A., JUNG, J., LEE, G.-J., PARK, H. J.,
SHIN, S.-I. & HONG, J.-Y. 2021. Progress in Oral Microbiome Related to Oral and Systemic
Diseases: An Update. *Diagnostics (Basel, Switzerland),* 11**,** 1283.
MOHANTY, R., ASOPA, S. J., JOSEPH, M. D., SINGH, B., RAJGURU, J. P., SAIDATH, K. &
SHARMA, U. 2019. Red complex: Polymicrobial conglomerate in oral flora: A review. *Journal
of family medicine and primary care,* 8**,** 3480-3486.
NAKAYAMA, M. & OHARA, N. 2017. Molecular mechanisms of Porphyromonas gingivalis-host cell
interaction on periodontal diseases. *The Japanese dental science review,* 53**,** 134-140.
NAKHJIRI, S. F., PARK, Y., YILMAZ, O., CHUNG, W. O., WATANABE, K., EL-SABAENY, A.,
PARK, K. & LAMONT, R. J. 2001. Inhibition of epithelial cell apoptosis by Porphyromonas
gingivalis. *FEMS Microbiol Lett,* 200**,** 145-9.

NARENGAOWA, KONG, W., LAN, F., AWAN, U. F., QING, H. & NI, J. 2021. The Oral-Gut-Brain
AXIS: The Influence of Microbes in Alzheimer's Disease. *Frontiers in Cellular Neuroscience,*
15.
PATTENGALE, N. D., ALIPOUR, M., BININDA-EMONDS, O. R., MORET, B. M. & STAMATAKIS,
A. 2010. How many bootstrap replicates are necessary? *J Comput Biol,* 17**,** 337-54.
PETTERSEN, E. F., GODDARD, T. D., HUANG, C. C., COUCH, G. S., GREENBLATT, D. M.,
MENG, E. C. & FERRIN, T. E. 2004. UCSF Chimera--a visualization system for exploratory
research and analysis. *J Comput Chem,* 25**,** 1605-12.
PETTERSEN, E. F., GODDARD, T. D., HUANG, C. C., MENG, E. C., COUCH, G. S., CROLL, T. I.,
MORRIS, J. H. & FERRIN, T. E. 2021. UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci,* 30**,** 70-82.
POTEMPA, J., MADEJ, M. & SCOTT, D. A. 2021. The RagA and RagB proteins of Porphyromonas
gingivalis. *Mol Oral Microbiol,* 36**,** 225-232.
PREMKUMAR, L., KURTH, F., DUPREZ, W., GROFTEHAUGE, M. K., KING, G. J., HALILI, M. A.,
HERAS, B. & MARTIN, J. L. 2014. Structure of the Acinetobacter baumannii dithiol oxidase
DsbA bound to elongation factor EF-Tu reveals a novel protein interaction site. *J Biol Chem,* 289**,**
19869-80.
ROBERTS, A. P. & KRETH, J. 2014. The impact of horizontal gene transfer on the adaptive ability of
the human oral microbiome. *Front Cell Infect Microbiol,* 4**,** 124.
ROBERTS, J. S., ATANASOVA, K. R., LEE, J., DIAMOND, G., DEGUZMAN, J., HEE CHOI, C. &
YILMAZ, O. 2017a. Opportunistic Pathogen Porphyromonas gingivalis Modulates Danger
Signal ATP-Mediated Antibacterial NOX2 Pathways in Primary Epithelial Cells. *Front Cell*
Infect Microbiol, 7**,** 291.
ROBERTS, J. S., ATANASOVA, K. R., LEE, J., DIAMOND, G., DEGUZMAN, J., HEE CHOI, C. &
YILMAZ, Ö. 2017b. Opportunistic Pathogen Porphyromonas gingivalis Modulates Danger
Signal ATP-Mediated Antibacterial NOX2 Pathways in Primary Epithelial Cells. *Frontiers in*
Cellular and Infection Microbiology, 7.
SAYERS, E. W., BOLTON, E. E., BRISTER, J. R., CANESE, K., CHAN, J., COMEAU, D. C.,
CONNOR, R., FUNK, K., KELLY, C., KIM, S., MADEJ, T., MARCHLER-BAUER, A.,
LANCZYCKI, C., LATHROP, S., LU, Z., THIBAUD-NISSEN, F., MURPHY, T., PHAN, L.,
SKRIPCHENKO, Y., TSE, T., WANG, J., WILLIAMS, R., TRAWICK, B. W., PRUITT, K. D.
& SHERRY, S. T. 2022. Database resources of the national center for biotechnology information.
Nucleic Acids Res, 50**,** D20-D26.
SHAW, L. P., SMITH, A. M. & ROBERTS, A. P. 2017. The oral microbiome. *Emerging Topics in Life*
Sciences, 1**,** 287-296.
SIEVERS, F., WILM, A., DINEEN, D., GIBSON, T. J., KARPLUS, K., LI, W., LOPEZ, R.,
MCWILLIAM, H., REMMERT, M., SODING, J., THOMPSON, J. D. & HIGGINS, D. G. 2011.
Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal
Omega. *Mol Syst Biol,* 7**,** 539.
SIMÓN-SORO, A., TOMÁS, I., CABRERA-RUBIO, R., CATALAN, M. D., NYVAD, B. & MIRA, A.
2013. Microbial geography of the oral cavity. *J Dent Res,* 92**,** 616-21.
SPOONER, R., DEGUZMAN, J., LEE, K. L. & YILMAZ, O. 2014. Danger signal adenosine via
adenosine 2a receptor stimulates growth of Porphyromonas gingivalis in primary gingival
epithelial cells. *Mol Oral Microbiol,* 29**,** 67-78.
SPOONER, R., WEIGEL, K. M., HARRISON, P. L., LEE, K., CANGELOSI, G. A. & YILMAZ, O.
2016. In Situ Anabolic Activity of Periodontal Pathogens Porphyromonas gingivalis and
Filifactor alocis in Chronic Periodontitis. *Sci Rep,* 6**,** 33638.
STARK, H., RODNINA, M. V., RINKE-APPEL, J., BRIMACOMBE, R., WINTERMEYER, W. &
VAN HEEL, M. 1997. Visualization of elongation factor Tu on the Escherichia coli ribosome.
Nature, 389**,** 403-6.

SU, Z., KONG, F., WANG, S., CHEN, J., YIN, R., ZHOU, C., ZHANG, Y., HE, Z., SHI, Y., XUE, Y.,
SHI, X., LU, L., SHAO, Q. & XU, H. 2010. The rag locus of Porphyromonas gingivalis might
arise from Bacteroides via horizontal gene transfer. *Eur J Clin Microbiol Infect Dis,* 29**,** 429-37.
SUN, Y., WANG, X., LI, J., XUE, F., TANG, F. & DAI, J. 2022. Extraintestinal pathogenic Escherichia
coli utilizes the surface-expressed elongation factor Tu to bind and acquire iron from holo-
transferrin. *Virulence,* 13**,** 698-713.
SZKLARCZYK, D., GABLE, A. L., LYON, D., JUNGE, A., WYDER, S., HUERTA-CEPAS, J.,
SIMONOVIC, M., DONCHEVA, N. T., MORRIS, J. H., BORK, P., JENSEN, L. J. & MERING,
C. V. 2019. STRING v11: protein-protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res,* 47**,**
D607-D613.
TAMURA, K., STECHER, G. & KUMAR, S. 2021. MEGA11: Molecular Evolutionary Genetics
Analysis Version 11. *Mol Biol Evol,* 38**,** 3022-3027.
TOMOYASU, T., TABATA, A., IMAKI, H., TSURUNO, K., MIYAZAKI, A., SONOMOTO, K.,
WHILEY, R. A. & NAGAMUNE, H. 2012. Role of Streptococcus intermedius DnaK chaperone
system in stress tolerance and pathogenicity. *Cell Stress Chaperones,* 17**,** 41-55.
TORRES, A. N., CHAMORRO-VELOSO, N., COSTA, P., CADIZ, L., DEL CANTO, F., VENEGAS, S.
A., LOPEZ NITSCHE, M., COLOMA-RIVERO, R. F., MONTERO, D. A. & VIDAL, R. M.
2020. Deciphering Additional Roles for the EF-Tu, l-Asparaginase II and OmpT Proteins of
Shiga Toxin-Producing Escherichia coli. *Microorganisms,* 8.
TRIBBLE, G. D., MAO, S., JAMES, C. E. & LAMONT, R. J. 2006. A Porphyromonas gingivalis
haloacid dehalogenase family phosphatase interacts with human phosphoproteins and is important
for invasion. *Proc Natl Acad Sci U S A,* 103**,** 11027-32.
WELLSLAGER, B., ROBERTS, J., CHOWDHURY, N., MADAN, L., ORELLANA, E. & YILMAZ, Ö.
2024. *Porphyromonas gingivalis* activates Heat-Shock-Protein 27 to drive a LC3C-specific
probacterial form of select autophagy that is redox sensitive for intracellular bacterial survival in
human gingival mucosa. *bioRxiv***,** 2024.07.01.601539.
WHELAN, S. & GOLDMAN, N. 2001. A general empirical model of protein evolution derived from
multiple protein families using a maximum-likelihood approach. *Mol Biol Evol,* 18**,** 691-9.
WIDJAJA, M., HARVEY, K. L., HAGEMANN, L., BERRY, I. J., JAROCKI, V. M., RAYMOND, B. B.
A., TACCHI, J. L., GRÜNDEL, A., STEELE, J. R., PADULA, M. P., CHARLES, I. G.,
DUMKE, R. & DJORDJEVIC, S. P. 2017. Elongation factor Tu is a multifunctional and
processed moonlighting protein. *Scientific Reports,* 7**,** 11227.
WILLIS, J. R. & GABALDÓN, T. 2020. The Human Oral Microbiome in Health and Disease: From
Sequences to Ecosystems. *Microorganisms,* 8**,** 308.
WILSON, M. 2004. *Microbial Inhabitants of Humans: Their Ecology and Role in Health and Disease,*
Cambridge, Cambridge University Press.
WYLENSEK, D., HITCH, T. C. A., RIEDEL, T., AFRIZAL, A., KUMAR, N., WORTMANN, E., LIU,
T., DEVENDRAN, S., LESKER, T. R., HERNANDEZ, S. B., HEINE, V., BUHL, E. M., P, M.
D. A., CUMBO, F., FISCHODER, T., WYSCHKON, M., LOOFT, T., PARREIRA, V. R., ABT,
B., DODEN, H. L., LY, L., ALVES, J. M. P., REICHLIN, M., FLISIKOWSKI, K., SUAREZ, L.
N., NEUMANN, A. P., SUEN, G., DE WOUTERS, T., ROHN, S., LAGKOUVARDOS, I.,
ALLEN-VERCOE, E., SPROER, C., BUNK, B., TAVERNE-THIELE, A. J., GIESBERS, M.,
WELLS, J. M., NEUHAUS, K., SCHNIEKE, A., CAVA, F., SEGATA, N., ELLING, L.,
STROWIG, T., RIDLON, J. M., GULDER, T. A. M., OVERMANN, J. & CLAVEL, T. 2020. A
collection of bacterial isolates from the pig intestine reveals functional and taxonomic diversity.
Nat Commun, 11**,** 6389.
YANG, J., ANISHCHENKO, I., PARK, H., PENG, Z., OVCHINNIKOV, S. & BAKER, D. 2020.
Improved protein structure prediction using predicted interresidue orientations. *Proc Natl Acad*
Sci U S A, 117**,** 1496-1503.

YAO, L., JERMANUS, C., BARBETTA, B., CHOI, C., VERBEKE, P., OJCIUS, D. M. & YILMAZ, O.
2010. Porphyromonas gingivalis infection sequesters pro-apoptotic Bad through Akt in primary
gingival epithelial cells. *Mol Oral Microbiol,* 25**,** 89-101.
YIKILMAZ, E., CHAPMAN, S. J., SCHRADER, J. M. & UHLENBECK, O. C. 2014. The interface
between Escherichia coli elongation factor Tu and aminoacyl-tRNA. *Biochemistry,* 53**,** 5710-20.
YILMAZ, Ö. 2008. The chronicles of Porphyromonas gingivalis: the microbium, the human oral
epithelium and their interplay. *Microbiology (Reading),* 154**,** 2897-2903.
YILMAZ, O., VERBEKE, P., LAMONT, R. J. & OJCIUS, D. M. 2006. Intercellular spreading of
Porphyromonas gingivalis infection in primary gingival epithelial cells. *Infect Immun,* 74**,** 703-10.
YILMAZ, O., YAO, L., MAEDA, K., ROSE, T. M., LEWIS, E. L., DUMAN, M., LAMONT, R. J. &
OJCIUS, D. M. 2008. ATP scavenging by the intracellular pathogen Porphyromonas gingivalis
inhibits P2X7-mediated host-cell apoptosis. *Cellular microbiology,* 10**,** 863-875.
YOSHIDA, A., NAKANO, Y., YAMASHITA, Y., OHO, T., SHIBATA, Y., OHISHI, M. & KOGA, T.
1999. A novel dnaK operon from Porphyromonas gingivalis. *FEBS Lett,* 446**,** 287-91.
ZHANG, Z., LIU, D., LIU, S., ZHANG, S. & PAN, Y. 2021. The Role of Porphyromonas gingivalis
Outer Membrane Vesicles in Periodontal Disease and Related Systemic Diseases. *Frontiers in*
Cellular and Infection Microbiology, 10.
862

2

The early colonizers allow
for the establishment of
orange complex bacteria

Prevotella intermedia
Prevotella nigrescens
Fusobacterium nucleatum
Streptococcus constellatus

3

Porphyromonas gingivalis
Tannerella forsythia
Treponema denticola

The orange complex
bacteria permit the
colonization of invasive
pathogenic bacteria that
cause inflammation in the
oral mucosa

The Human
Oral-Gut Axis

1

Early colonizers, belonging to
the blue, green and purple oral
complex, are established from
birth and contribute to the initial
development of the oral
microbiome.

4

A physical connection between the oral
cavity and digestive tract permits
translocation of oral bacteria to the gut

Escherichia coli
Helicobacter pylori
Bacteroidetes fragilis
Citrobacter freundii
Porphyromonas gingivalis
Fusobacterium nucleatum

Mucosal Epithelium

FIGURE 1

PROTEIN	PROTEIN SIGNATURE SEQUENCES	PROTEIN ID	SPECIES
Elongation Factor Tu	LLDEGQAGDNVGLLLR	B2RL52; A0A1D3UQ89; A0A2A6EK37	*Porphyromonas gingivalis, Tannerella forsythia, Prevotella intermedia*
RagB/SusD family OM Protein	MPYIDRVIMPNDKM	A0A2A6EAR0, A0A2D2N8A1	*Tannerella forsythia, Porphyromonas gingivalis*
DnaK	IVNEPTAAALAYGIDKIVN EPTAAALAYGIDK	AAQ66298; A0A2A6ENH2	*Porphyromonas gingivalis, Prevotella intermedia,*

Table 1

FIGURE 2

FIGURE 3

A

Legend:
□ Other
○ Oral
△ Gut
★ *P. gingivalis*
● Red Complex
● Orange Complex
● Yellow Complex

P. gingivalis	MAKEHFNRSKPHVNVGTIGHVDHGKTTLTAAITTVLAKRGL--SELRSFDSIDNAPEEKE	58
P. salivae	MAKEEFVRTKPHVNIGTIGHVDHGKTTLTAAISKVLHEKGFGSEDVKSFDQIDNAPEEKE	60
P. nigrescens	MAKETFQRTKPHVNIGTIGHVDHGKTTLTAAISKVLHDKGFGSEDVKSFDQIDNAPEEKE	60
P. intermedia	MAKETFQRTKPHVNIGTIGHVDHGKTTLTAAISKTLHDKGFGGEEAKSFDQIDNAPEEKE	60
P. corporis	MAKETFQRTKPHVNIGTIGHVDHGKTTLTAAISKTLHDKGFGGEDAKSFDQIDNAPEEKE	60

**** * *:.***** :***************:.. .:*: .: :***.********

P. gingivalis	RGITINTSHVEYQTANRHYAHVDCPGHADYVKNMVTGAAQMDGAIIVVAATDGPMPQTRE	118
P. salivae	RGITINSAHIEYETAKRHYAHVDCPGHADYVKNMVTGAAQMDGAILVVAATDGPMPQTRE	120
P. nigrescens	RGITINSAHIEYETAKRHYAHVDCPGHADYVKNMVTGAAQMDGAILVVAATDGPMPQTRE	120
P. intermedia	RGITINSSHIEYETAKRHYAHVDCPGHADYVKNMVTGAAQMDGAILVCAATDGPMPQTRE	120
P. corporis	RGITINSSHIEYETAKRHYAHVDCPGHADYVKNMVTGAAQMDGAILVVAATDGPMPQTRE	120

******:.:.**:.**:.************************** :. *********** *

Conserved Translational type guanine nucleotide binding domain

B

P. gingivalis' EF-TU

FIGURE 4

A

P.gingivalis EF-TU
T. forsythia f EF-TU
E. coli EF-TU

Percent Identity 78%

B

```
T. forsythia   MAKETFQRTKPHVNIGTIGHVDHGKTTLTAAISKTLHDKGFGGEEAKSFDQIDNAPEEKE     60
E. coli        MAKEKFDRSKPHVNIGTIGHVDHGKTTLTAAITTVLAKKGLS--ELRSFDSIDNAPEEKE     58
P. gingivalis  MAKEHFNRSKPHVNVGTIGHVDHGKTTLTAAITTVLAKRGLS--ELRSFDSIDNAPEEKE     58
               **** *.*.*****:***************...* .:*:.  * ;***.********

T. forsythia   RGITINSSHIEYETAKRHYAHVDCPGHADYVKNMVTGAAQMDGAILVCAATDGPMPQTRE    120
E. coli        RGITINTSHVEYQTANRHYAHVDCPGHADYVKNMVTGAAQMDGAIIVVAATDGPMPQTRE    118
P. gingivalis  RGITINTSHVEYQTANRHYAHVDCPGHADYVKNMVAGAAQMDGAIIVVAATDGPMPQTRE    118
               ******.**:**.**.**********************:.*********.********* **********

T. forsythia   HVLLARQVNVPRLVVFLNKCDMVEDEEMLELVEMELGEILTQYEYEE-DTPIIRGSALGA    179
E. coli        HILLARQVNVPRLVVFMNKCDMVDDEEMLELVEMEMRELLSFYQFDGDNTPIIRGSALGA    178
P. gingivalis  HILLARQVNVPRLVVFMNKCDMVDDEEMLELVEMDMRELLSFYDFDGDNTPIIRGSALGA    178
               *:*************.*******:.**********:: *:*: *:::  :***********
```

Conserved EF-TU Domain 1

C

EF-TU

DsbA

EF-TU

DsbA

FIGURE 5

A

Conserved Starch binding associating with outer membrane and lipoprotein lipid attachment domain

FIGURE 6

A

P. gingivalis RagB
T. forsythia RagB/SuSD
Low Percent Identity

B

P. gingivalis RagB
B. fragilis NanU
Percent Identity: 45.45%

C

T. forsythia RagB/SuSD
B. fragilis NanU
Percent Identity: 30.65%

D

FIGURE 7

A

T. forsythia Rag/SusD family protein

B

B. fragilis NanU

FIGURE 8

FIGURE 9

ACCESSION #	SPECIES	Percent Identities	Amino Acid #	Region
WP_014225269.1	*Tannerella forsythia*		395	Oral
WP_005468460	*Porphyromonas catoniae*	92.41	395	Oral
WP_121736507.1	*Parabacteroides distasonis*	91.86	395	Gut
MSK95816.1	*Escherichia coli*	91.86	395	Gut
WP_195341511	*Bacteroides fragilis*	91.14	394	Gut
MZM33066	*Bifidobacterium pseudocatenulatum*	91.14	394	Gut
CUO27731	*Catenibacterium mitsuokai*	90.38	394	Gut
WP_010955998	*Porphyromonas gingivalis*	90.13	395	Oral
WP_004366570.1	*Prevotella nigrescens*	80.86	396	Oral, urogenital
WP_028905370.1	*Prevotella intermedia*	80.6	396	Oral, urogenital
PZR28947	*Citrobacter freundii*	73.86	395	Gut, Oral
WP_096015169	*Campylobacter lanienae*	71.93	399	Gut
WP_001040716	*Streptococcus oralis*	71.72	398	Oral, Upper respiratory
WP_049480505.1	*Streptococcus constellatus*	71.72	398	Oral, Urogenital, Gut
WP_024962200	*Campylobacter ureolyticus*	71.5	399	Gut
WP_001040579.1	*Helicobacter pylori*	71.07	399	Oral, Gut
WP_032887443.1	*Fusobacterium nucleatum*	70.56	394	Oral
WP_040304280.1	*Campylobacter gracilis*	69.92	399	Oral
WP_004894320.1	*Lactobacillus johnsonii*	69.47	396	Gut
WP_002669993.1	*Treponema denticola*	62.72	395	Oral
AAB00499	*Homo sapiens*	54.68	452	Control

Supplementary Table 1.

ACCESSION #	SPECIES	Percent Identities	Amino Acid #	Region
WP_028905370	*Prevotella intermedia*		396	Oral, Urogenital
WP_025875819	*Prevotella corporis*	96.21	396	Oral, Urogenital
WP_211791898	*Prevotella nigrescens*	93.94	396	Oral
MBF1523572	*Prevotella salivae*	87.37	398	Oral, Gut
WP_007763555	*Bacteroides finegoldii*	85.61	394	Gut
WP_065539685	*Bacteroides caecimuris*	85.1	394	Gut
WP_234097660	*Bacteroides faecis*	84.6	394	Gut
WP_005656566	*Bacteroides stercoris*	84.34	394	Gut
CUO27731	*Catenibacterium mitsuokai*	83.84	394	Gut
MZU51045	*Escherichia coli*	83.33	394	Gut
WP_004335339	*Porphyromonas endodontalis*	79.9	396	Oral
WP_010955998	*Porphyromonas gingivalis*	78.34	395	Oral
VDR34100	*Faecalibacterium prausnitzii*	76.96	395	Gut
MBF1213542	*Fusobacterium periodonticum*	70.56	395	Oral
WP_032887443	*Fusobacterium nucleatum*	70.05	394	Oral
WP_118358359	*Dorea formicigenerans*	69.6	397	Gut
WP_071441959	*Traorella massiliensis*	69.54	394	Gut
WP_009527606	*Peptoanaerobacter stomatis*	68.84	397	Oral
WP_040304280	*Campylobacter gracilis*	67.92	399	Oral
KXT83818	*Streptococcus oralis*	66.92	398	Oral
AAB00499	*Homo sapiens*	54.20	452	Control

Supplementary Table 2.

ACCESSION #	SPECIES	Percent Identities	Amino Acid #	Region
WP_097530727	*Tannerella forsythia*		631	Oral
WP_106831261	*Parabacteroides pacaensis*	74.96	630	Gut
MBS6574981	*Parabacteroides goldsteinii*	69.15	627	Gut
WP_183668678	*Parabacteroides faecis*	68.76	627	Gut
WP_195292649	*Parabacteroides merdae*	67.19	634	Gut
GAP81744.1	*Porphyromonas gingivalis*	64.29	503	Oral
WP_007573186	*Prevotella multisaccharivorax*	52.66	645	Oral
WP_004351005	*Prevotella buccalis*	52.45	642	Oral
MBF1544323	*Prevotella salivae*	52.14	642	Oral
WP_240062496	*Bacteroides xylanisolvens*	52.02	636	Gut
OKZ23327	*Bacteroides finegoldii*	51.93	642	Gut
WP_022136618	*Bacteroides congonensis*	51.91	642	Gut
WP_074637268	*Bacteroides ovatus*	51.85	642	Gut
MTT48421	*Proteus mirabilis*	43.88	370	Gut
MZM33202	*Bifidobacterium pseudocatenulatum*	29.41	240	Gut
WP_195207990	*Bacteroides uniformis*	28.73	563	Gut
PZR29998	*Citrobacter freundii*	27.00	579	Gut
WP_135958631	*Parabacteroides distasonis*	24.88	535	Gut
PFX22450	*Stylophora pistillata*	24.66	3185	Control
MBL1007498	*Escherichia coli*	24.5	654	Gut
WP_188807090	*Porphyromonas pasteri*	23.6	548	Oral
WP_028906323	*Prevotella intermedia*	22.39	544	Oral

Supplementary Table 3.

ACCESSION #	SPECIES	Percent Identities	Amino Acid #	Region
WP_097646862.1	*Prevotella intermedia*		633	Oral
WP_211795957	*Prevotella nigrescens*	95.58	633	Oral
WP_004349103	*Prevotella buccalis*	88.65	643	Oral
HAW5551234	*Escherichia coli*	80.03	639	Gut
VDR35622	*Faecalibacterium prausnitzii*	74.29	648	Gut
WP_004583505	*Porphyromonas gingivalis*	73.52	640	Oral
WP_204898322	*Pseudoflavonifractor capillosus*	61.55	612	Gut
WP_002669872	*Treponema denticola*	61.28	646	Oral
WP_050622774	*Clostridium phoceensis*	60.95	618	Gut
WP_058962768	*Fournierella massiliensis*	60.51	609	Gut
WP_204741179	*Intestinimonas butyriciproducens*	60.28	621	Gut
WP_034438123	*Clostridium ihumii*	60	607	Gut
WP_039997401	*Flavonifractor plautii*	59.91	617	Gut
WP_204560969	*Collinsella tanakaei*	59.81	632	Gut
WP_022348576	*Enorma massiliensis*	59.7	631	Gut
WP_204793018	*Olsenella profusa*	59.12	633	Oral
WP_142414544	*Hathewaya massiliensis*	58.87	619	Gut
WP_204584696	*Collinsella intestinalis*	58.64	633	Gut
WP_035432935	*Lancefieldella parvula*	58.45	634	Oral
AAH00478	*Homo sapiens*	57.97	679	Control
WP_005869203	*Campylobacter gracilis*	56.69	628	Oral

Supplementary Table 4.

Node	Protein	Confidence Score
CAH07419.1	NanU Protein	0.706-0.929
CAH07421.1	Similar to Bacteroides fragilis outer membrane protein omp121	0.769
CAH07497.1	Putative outer membrane protein	0.764
CAH07502.1	Starch-binding outer membrane protein	0.765
CAH09392.1	Putative export protein	0.706
CAH09730.1	Putative outer membrane protein	0.756

Supplementary Table 5.

Node	Protein	Confidence Score
BFO_1797	Rag/SusD Family Protein (Vesicle Isolation)	0.707-0.912
BFO_1796	TonB linked outer membrane protein	0.912
BFO_1798	Tat pathway signal sequence domain protein	0.734-0.925
BFO_1799	Putative lipoprotein	0.707-0.925

Supplementary Table 6