Title: Direct piriform-to-auditory cortical projections shape auditory-olfactory integration

Abbreviated title: Auditory-olfactory integration in auditory cortex

Nathan W. Vogler[1], Ruoyi Chen[1], Alister Virkler[2], Violet Y. Tu[1], Jay A. Gottfried[2], and Maria N.

Geffen[1,*]

[1]Department of Otorhinolaryngology, Perelman School of Medicine, University of Pennsylvania

[2]Department of Neurology, Perelman School of Medicine, University of Pennsylvania

*Corresponding Author: mgeffen@pennmedicine.upenn.edu

3450 Hamilton Walk, Stemmler Hall G10, Philadelphia, PA 19104

Number of pages: 46

Number of figures: 6

Number of tables: 1

Number of words: Abstract (169), Introduction (649), Discussion (1283)

Keywords: auditory cortex, piriform cortex, auditory, olfactory, multisensory integration

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: Funding: NIH F32MH127788 (to N.W.V.), Kaufman Foundation KA2020-

114796 (to M.N.G. and J.A.G.), NIH R01DC015527, NIH R01DC014479, and NIH R01NS113241

(to M.N.G.). The authors would like to thank Tyler Ling for technical assistance.

Abstract

In a real-world environment, the brain must integrate information from multiple sensory modalities,

including the auditory and olfactory systems. However, little is known about the neuronal circuits

governing how odors influence and modulate sound processing. Here, we investigated the

mechanisms underlying auditory-olfactory integration using anatomical, electrophysiological, and

optogenetic approaches, focusing on the auditory cortex as a key locus for cross-modal

integration. First, retrograde and anterograde viral tracing strategies revealed a direct projection

from the piriform cortex to the auditory cortex. Next, using *in vivo* electrophysiological recordings

of neuronal activity in the auditory cortex of awake male or female mice, we found that odors

modulate auditory cortical responses to sound. Finally, we used *in vivo* optogenetic manipulations

during electrophysiology to demonstrate that olfactory modulation in auditory cortex, specifically,

odor-driven enhancement of sound responses, depends on direct input from the piriform cortex.

Together, our results identify a novel role of piriform-to-auditory cortical circuitry in shaping

olfactory modulation in the auditory cortex, shedding new light on the neuronal mechanisms

underlying auditory-olfactory integration.

Significance Statement

All living organisms exist within multisensory environments, yet there is a lack in our

understanding of how the brain integrates multisensory information. This work highlights a direct

piriform-to-auditory cortical circuit governing auditory-olfactory integration. Our results shed new

light on a relatively understudied area of multisensory research, promising a more robust

understanding of how animals and humans perceive and interact within complex environments.

Introduction

A fundamental role of the central nervous system is to integrate information across multiple

sensory modalities to optimize decision-making and behavior. Indeed, rarely does any individual

sensory modality operate alone and unabetted: humans and animals must integrate combinations

of sounds, smells, visual input, tactile and proprioceptive information (Gottfried and Dolan, 2003;

Deroy et al., 2016; Angeloni and Geffen, 2018; Choi et al., 2018; Wallace et al., 2020). When

combining sounds and smells, the brain must integrate information from both the auditory and

olfactory systems. Auditory-olfactory integration is an ethologically important domain of

multisensory integration. When a cat approaches a mouse, the mouse may use the combination

of sounds and smell of the cat to identify it as a predator (Halene et al., 2009). Moreover,

combinations of auditory and olfactory stimuli are crucial for pup retrieval behavior in mice, and

play critical roles during development — bonds between parents and offspring form based on

body odors and vocalization sounds (Marlin and Froemke, 2017; Tasaka et al., 2020; McRae et

al., 2023). In humans, the sounds of eating food, such as crunchy potato chips, affect perception

of the pleasantness of the food odor, such as potato chip odor/staleness (Seo and Hummel, 2011;

Seo et al., 2014). However, the circuits and mechanisms underlying auditory-olfactory integration

have remained elusive.

Previous studies examining interactions between audition and olfaction demonstrated cross-

modal activation of piriform and auditory cortical areas (Smith et al., 2009; Wesson and Wilson,

2010; Cohen et al., 2011; Varga and Wesson, 2013; Cohen and Mizrahi, 2015; Gnaedinger et al.,

2019; Olofsson et al., 2019; Tasaka et al., 2020; Gilday and Mizrahi, 2023; Wu et al., 2023). In

the auditory cortex (ACx) of lactating mother mice, pup odors reduce spontaneous activity and

increase evoked activity in response to pup ultrasonic vocalizations (Cohen et al., 2011; Cohen

and Mizrahi, 2015). Furthermore, learned associations between an odor and a subsequent sound

modulate single-neuron responses in the auditory cortex (Gilday and Mizrahi, 2023). Together,

these studies suggest that the ACx serves as a locus for auditory-olfactory cross-modal

integration, specifically regarding the olfactory modulation of sound processing.

While these studies point to an involvement of the ACx in auditory-olfactory integration, much

remains unknown regarding where and how these cross-modal inputs converge. By comparison,

previous research on audio-visual integration has found that concurrent visual stimuli improve

sound detection and encoding in ACx, facilitated by cross-cortical connections between visual

cortex and ACx (Bizley et al., 2016; Atilgan et al., 2018; Meijer et al., 2018; Zhou et al., 2020; Han

et al., 2021). Moreover, audio-visual integration crucially depends on the temporal binding of the

combined audio-visual stimuli (Meijer et al., 2017, 2019). Similar studies on auditory-olfactory

integration have been limited by experimental challenges in delivering combinations of odors and

sounds with temporal precision, as well as a lack of knowledge of the brain circuitry underlying

auditory-olfactory interactions. Notably, reports of potential cross-modal anatomical connections

linking the piriform cortex (Pir) and ACx have been inconsistent. Some have found that neurons

in Pir display sound-evoked responses (Varga and Wesson, 2013), and that ACx projects to Pir

but not vice versa (Budinger et al., 2006; Budinger and Scheich, 2009). Others have reported a

sparse but detectable direct projection from Pir to ACx (Costa et al., 2017; Tasaka et al., 2020),

yet its potential role in odor-driven modulation of ACx is unclear. Identifying and characterizing

these pathways are important first steps in deciphering how the brain integrates auditory and

olfactory stimuli.

Here, we confirm the direct Pir-to-ACx projection using multiple circuit tracing strategies and

examine its functional role in auditory-olfactory integration in ACx. We explore the impact of this

pathway on olfactory modulation of auditory processing, using a novel experimental setup for

delivering odors and sounds with temporal precision to awake mice during *in vivo*

electrophysiological recordings in ACx. Our findings reveal that the Pir-to-ACx pathway

contributes specifically to odor-mediated enhancement of sound responses in ACx.

Materials and Methods

Mice

For anatomical circuit tracing experiments, we used 5 male C57BL/6J mice (The Jackson

Laboratory; strain #000664; RRID:IMSR_JAX:000664; aged 8-16 weeks), 2 male Cdh23 mice

(B6N(Cg)-Cdh23$^{tm2.1Kjn}$/Kjn; The Jackson Laboratory, strain #:018399; RRID:IMSR_JAX:018399;

aged 22 weeks), and 13 Ai14 Cre reporter mice (progeny of Cdh23 mice crossed with Ai14 mice

[B6.Cg-Gt(ROSA)$^{26Sortm14(CAG-tdTomato)Hze}$/J; The Jackson Laboratory, strain #:007914;

RRID:IMSR_JAX:007914], 7 male, 6 female, aged 18-26 weeks). For electrophysiological

recordings, we used 18 mice: 4 C57BL/6J mice (2 male, 2 female; aged 12-17 weeks at time of

recording); and 14 Cdh23 mice (5 male, 9 female; aged 12-26 weeks). Cdh23 mice were used for

most experiments because they do not exhibit age-related hearing loss which is common in

C57BL/6J mice (Johnson et al., 2017). Any C57BL/6J mice were used by 17 weeks of age. Mice

were housed at 28 °C with a 12-h reversed light cycle and food and water provided *ad libitum*.

Experiments were performed during the animals' dark cycle. All experimental procedures were

approved by the Institutional Care and Use Committee at the University of Pennsylvania and in

accordance with National Institutes of Health guidelines.

Stereotaxic surgery and virus injections

Mice were anesthetized under isoflurane (1-3%). Prior to surgery, mice were administered

subcutaneous doses of buprenorphine (0.05-0.1 mg/kg Buprenex or 1.3 mg/kg Ethiqa),

dexamethasone (0.2 mg/kg), and bupivicane (local; 2 mg/kg). A small craniotomy (~1 mm

diameter) was made above the respective ACx or Pir injection sites. To label ACx-projecting

neurons in Pir in Cdh23 mice, we injected AAVrg-hSyn-eGFP (Addgene #50465-AAVrg;

RRID:Addgene_50465; 1.7x10^{13} GC/mL; 250 nL) into left ACx (4.3 mm ML and -2.65 mm AP

[relative to Bregma], -0.7 mm DV [relative to brain surface]). To label ACx-projecting neurons in

Pir in Ai14 mice, we injected AAVrg-hSyn-Cre-WPRE-hGH (Addgene #105553-AAVrg;

RRID:Addgene_105553; 1.9 x10^{13} GC/mL; 150-250 nL) into left ACx. To label Pir-recipient

neurons in ACx in Ai14 mice, we injected AAV1-hSyn-Cre-WPRE-hGH (Addgene #105553-AAV1;

RRID:Addgene_105553; 1.9 x10^{13} GC/mL; 150-180 nL) into left Pir (3.9 mm ML, -1.5 mm AP, -

4.00 DV). Viruses were delivered through 30-40 μm diameter glass micropipettes connected to a

syringe pump (Harvard Apparatus Pump 11 Elite; 60 nL/min infusion rate). For optogenetic

experiments, in Cdh23 or C57BL/6J mice we injected AAVrg-hSyn-Cre-WPRE-hGH (300 nL) into

left ACx and AAV5-hSyn1-SIO-stGtACR1-FusionRed (Addgene #105678-AAV5;

RRID:Addgene_105678; 2.1 x10^{13} GC/mL; 300 nL) or AAV5-FLEX-tdTomato (Addgene #28306-

AAV5; RRID:Addgene_28306; 2.1 x10^{13} GC/mL; 300 nL) into left Pir (3.9 mm ML, -1.0 mm AP, -

4.00 mm DV). A fiber optic cannula (ThorLabs CFML52L05; Ø200 μm Core, 0.50 NA, L=5mm)

was implanted over left Pir (3.9 mm ML, -1.0 mm AP, -3.6 mm DV) and secured with dental cement

(C&B Metabond). For all electrophysiological recordings, a ground pin was inserted via small

craniotomy over the left frontal cortex, and a custom-made stainless steel headplate (eMachine

Shop) was secured to the skull using dental cement and acrylic (Lang Dental). After all surgeries,

we subcutaneously administered meloxicam (5 mg/kg) and an antibiotic (Baytril 5 mg/kg) for 4

days during postoperative care. After virus injections, to allow adequate time for virus expression,

we waited at least two weeks (for retrograde tracing/optogenetics) or three weeks (for anterograde

tracing) before subsequent recording and/or transcardial perfusion.

Histology and cell counting

Mice were anesthetized via intraperitoneal injection of ketamine (120 mg/kg) and

dexmedetomidine (1 mg/kg), then perfused with phosphate-buffered saline (PBS) followed by 4%

paraformaldehyde (PFA) in PBS. Brains were extracted and immersed in 4% PFA overnight then

in 30% sucrose in PBS for at least 36 hours at 4°C. Coronal brain sections (60 μm thickness)

were cut on a cryostat (Leica CM1860) at -25°C and mounted on slides using Invitrogen ProLong

Antifade Mountant with DAPI. Fluorescent images were acquired using a slide scanning

microscope (Zeiss Axio Scan.Z1) under 20x magnification and ZEN 2.3 software. Whole brain

fluorescent images were manually aligned with the reference mouse brain atlas (Franklin &

Paxinos) to confirm the accuracy of injection sites and extent of viral spread, and to determine

regions of interest (ROIs) for cell counting. For the anterior Pir, four coronal sections between 1.0-

2.0 mm anterior to Bregma were counted in each mouse. For the posterior Pir, four coronal

sections between 1.5-2.5 mm posterior to bregma were counted in each mouse. Fluorescent-

labeled cells within ROIs were manually counted using ImageJ software. For ACx subregions,

eight coronal sections between 2.2-3.6 mm posterior to bregma were counted in each mouse. A

custom-written MATLAB script was used to register the normalized depth of each cell from the

pia during manual counting. Specifically, for each ACx subregion, a line segment spanning the

depth of the cortex was drawn perpendicular to the pia in the middle of the subregion. The

normalized depth for each cell was defined as the relative length of the projection of the cell's

location onto that line segment.

Electrophysiological recordings

Recordings were performed in a custom-built acoustic isolation booth. At least one week following

headplate surgery, mice were habituated to the recording setup for increasing durations (5 – 60

min) over the course of 3 days. On the day of recording, mice were anesthetized with 1.5-2.5%

isoflurane, and a small craniotomy was made over the left auditory cortex using a dental drill,

centered ~1.5mm anterior to the lambdoid suture along the caudal end of the squamosal suture.

Following the craniotomy, mice were allowed to recover from anesthesia for at least 30 min. Acute

extracellular recordings in ACx were performed using a 32-channel silicon probe (A1x32-Poly2-

5mm-50s-177, NeuroNexus). Prior to insertion, the probe was coated in a dye (CM-DiI or SP-DiO;

Invitrogen) for post hoc visualization of recording sites. The probe was positioned perpendicularly

to the brain surface and lowered at 1-2 μm/s to a final depth of 900-1200 μm or until all channels

were within the brain. During lowering of the probe, we monitored responses to brief broadband

noise clicks to confirm localization within auditory cortex. Neuronal signals were amplified and

digitized with an Intan headstage (RHD 32ch) and recorded by an OpenEphys acquisition board

(Siegle et al., 2017) at a rate of 30 kHz. Signals were filtered between 500 and 6000 Hz, offset

corrected, and re-referenced to the median across all channels. Spiking data were sorted using

Kilosort2 (Pachitariu et al., 2016) and manually curated in phy2 to identify putative single- and

multi-units.

Stimuli

Stimuli during electrophysiological recordings consisted of combinations of auditory and olfactory

stimuli. Odors consisted of acetophenone (CAS 98-86-2; Sigma-Aldrich), eucalyptus (100% pure

eucalyptus oil; NOW essential oils), or an acetophenone + eucalyptus mixture (1:1 vol/vol),

delivered using a respiration-triggered flow dilution olfactometer (Shusterman et al., 2011; Dewan

et al., 2018; Williams and Dewan, 2020). We chose these odorants due to their use in previous

studies (acetophenone in Cohen et al., 2011), and initially to compare the effects of a

monomolecular odorant (acetophenone) versus a more complex mixture (eucalyptus). Because

preliminary experiments showed no odor-dependent difference in effects, we combined the

groups for subsequent experiments and analysis. This approach allowed us to vary odors

between successive recordings in the same mouse, to minimize potential habituation to a single

odor. Odorants were liquid diluted 1:1 vol/vol in mineral oil (CAS 8042-47-5; Sigma-Aldrich) in

amber vials, then further air diluted (2% N_2/Air) in the olfactometer, for a final dilution of 1%. The

control (non-odorant) vial contained pure mineral oil. The olfactometer and stimulus delivery were

controlled by custom Python software and Arduino MEGA 2560. Throughout the recording, we

monitored the respiratory cycle of the mouse. Respiration (sniffing) was measured non-invasively

using a pressure sensor connected via tubing to the odor port around the mouse's nose. The

pressure was transduced using the sensor and a preamplifier, and the analog signal from the

preamplifier was recorded along with electrophysiological data in OpenEphys. Between each trial,

the olfactometer tubing was cleared and primed with the next odor for at least 3 s. At the onset of

a trial, the mouse's inhale triggered the 'final valve' to switch from delivering continuous clean

background air to odorized air, which rapidly delivered the odorized air to the mouse's nose with

no change in total air flow (valve opening duration 1.25 s). Opening of the final valve also triggered

playing of the sound stimulus with a delay of 250 ms to allow adequate time for the odor to reach

the mouse's nose and subsequent inhale. The onset timing and kinetics of odors were verified

using a photo-ionization detector (miniPID; Aurora Scientific). Sounds consisted of 1 s duration

broadband gaussian white noise filtered between 3-60 kHz, either unmodulated or amplitude-

modulated (10 Hz, 100% modulation depth). Sounds were generated in Python and converted to

analog signals via National Instruments card (NI PCIe-6353, 200 kHz sample rate) and delivered

through a speaker (MF-1; Tucker-Davis Technologies) pointed toward the mouse's right ear, at 0,

45, 55, 65, or 75 dB SPL. Background noise from the olfactometer air flow was measured to be

~50 dB SPL (1/4-inch condenser microphone; Brüel &Kjær). During recordings, on half of trials,

we delivered either odorized air or control air. For recordings with optogenetics, on half of trials,

we delivered 470 nm LED light (M470F4; ThorLabs) through a fiber optic cannula (ThorLabs

CFML52L05; power 2.8-3.7 mW at cannula tip). LED stimulus onset and duration (1.25 s) were

the same as final valve opening. For all recordings, trial order was randomized and different for

each recording.

Electrophysiological data analysis

Neuronal spiking data from each recorded unit were organized by trial type and aligned to the trial

onset (i.e. the time of final valve opening). For each unit, peristimulus time histograms (PSTHs)

of neuronal spikes were generated using 10 ms time bins, then averaged across trials for each

condition. We included all units with baseline firing rate > 1 Hz. For responses to unmodulated

noise stimuli, we analyzed spiking during a response window of 0-250 ms after sound onset; for

responses to 10 Hz AM stimuli, we analyzed spiking during a response window of 0-1 s. To classify

each unit as sound-responsive or odor-modulated, we input the number of spikes during each

trial's response window into a generalized linear model (GLM; predictor variables: sound level

[continuous variable normalized from 0-1], odor [0 or 1]; response variable: number of spikes

during response window; Poisson distribution, log link function). This allowed the classification of

each unit's responses as having a main effect ($p<0.05$) of sound, odor, and/or a sound-odor

interaction. Sound-responsive units that were also responsive to odor or had a significant sound-

odor interaction were classified as 'odor-modulated' units. For each odor-modulated unit, to

determine the direction and magnitude of olfactory modulation, we calculated an Odor Modulation

Index (OMI) according to:

$$OMI = \frac{FR_{Odor} - FR_{Air}}{FR_{Odor} + FR_{Air}}$$

Where FR_{Odor} and FR_{Air} are the mean firing rates in response to sound, with or without odor,

respectively. For each unit we calculated the OMI using the mean firing rates for all responses to

5, 15, and 25 dB SNR sounds. Odor-modulated units with OMI>0 were classified as 'enhanced'

units; units with OMI<0 were classified as 'suppressed' units. In addition, for both enhanced and

suppressed units, we calculated the OMI for each sound level using the responses to each sound

stimulus separately. To calculate z-scored firing rates, we used the mean and standard deviation

across trials of firing rate during the 1 s baseline period prior to final valve opening. For 'enhanced'

units in optogenetics experiments (stGtACR1 and control mice), we calculated an LED Modulation

Index according to:

$$LED\ Modulation\ Index = \frac{FR_{Odor\ (LED\ ON)} - FR_{Odor\ (LED\ off)}}{FR_{Odor\ (LED\ ON)} + FR_{Odor\ (LED\ off)}}$$

Where $FR_{Odor\ (LED\ ON)}$ and $FR_{Odor\ (LED\ off)}$ are the mean firing rates in response to Odor+Sound, with

or without LED light, respectively.

Sniff data analysis

Raw analog sniff data from each recording were bandpass filtered between 0.5 and 20 Hz, then

subsequent analyses were performed on the filtered sniff traces for each trial during the 1.25 s

trial duration, as described above. To calculate sniff frequency for each trial, we measured the

average time between peaks (exhales) and between troughs (inhales) in the sniff trace. To

calculate sniff amplitude for each trial, we measured the difference between the average peak

amplitude minus the average trough amplitude.

Statistics

Data analysis and figure generation were performed using Matlab or Python. We used the

Shapiro-Wilk test to assess normality. For parametric data we used two-tailed paired or unpaired

t-tests between groups; for non-parametric data we used two-tailed Wilcoxon signed-rank or

Mann-Whitney tests between groups, or Friedman tests for multiple groups. We performed

Bonferroni corrections for multiple comparisons. Significance levels are defined as $p<0.05$. Group

data in figures are presented as mean $\pm$ SEM, unless otherwise noted. The statistical tests

performed are indicated in the text and figure legends. Table 1 contains a detailed summary of

statistical tests and comparisons.

Data and code availability

Source data is available on Dryad: https://doi.org/10.5061/dryad.dncjsxm7c. Original code is

available on GitHub: https://github.com/geffenlab/Vogler_2024.

Results

The auditory cortex receives direct input from the piriform cortex.

To examine potential anatomical substrates for auditory-olfactory integration in the auditory cortex

(ACx), we first sought to identify whether ACx receives direct projections from olfactory brain

areas such as the piriform cortex (Pir). We started by injecting a retrograde virus expressing GFP

(AAVrg-hSyn-eGFP) into the left ACx of C57BL/6J or Cdh23 mice to selectively label the cell

bodies of ACx-projecting neurons (Fig. 1A). We observed fluorescently labeled cells in the

ipsilateral Pir (GFP, Fig. 1B). Because the anterior and posterior regions of the Pir are known to

exhibit differential long-range connectivity patterns (Bekkers and Suzuki, 2013), we quantified and

compared retrograde GFP expression in the anterior Pir (aPir) versus posterior Pir (pPir) (Fig. 1B-

C). In most mice, we observed a greater number of labeled cells in pPir compared to aPir. (Fig.

1C), suggesting that the Pir-to-ACx projection predominantly arises from pPir. To confirm the

accuracy of injection sites, we examined the extent of viral spread in each mouse (Fig. 1D-E),

ensuring localization of the virus injection to ACx with minimal spread into surrounding non-

auditory regions.

To further confirm the existence of this projection and validate our approach, we performed a

similar retrograde tracing experiment in Ai14 cre-reporter mice (progeny of Cdh23 mice crossed

with Ai14 mice), injecting retroAAV expressing Cre (AAVrg-hSyn-Cre) into the ACx (Fig. 1E).

Again, we found retrogradely labeled cell bodies in ipsilateral Pir (tdTomato, Fig. 1F), and we

observed a significantly greater number of labeled cells in pPir compared to aPir (Fig. 1G,

*p=0.020, paired t-test). These retrograde tracing experiments reveal a direct projection to ACx

arising from Pir, predominantly from pPir.

Next, to further characterize Pir input to ACx, we performed anterograde viral tracing to reveal the

targets of Pir neurons within ACx. We injected a transsynaptic anterograde virus (Zingg et al.,

2020) expressing Cre (AAV1-hSyn-Cre) into the left Pir of Ai14 cre-reporter mice (progeny of

Cdh23 mice crossed with Ai14 mice) to selectively label neurons in ACx that receive Pir input (Fig.

2A). Because our retrograde tracing experiments (Fig. 1) showed that the Pir-to-ACx projection

arises mostly from pPir, we specifically targeted pPir for the anterograde virus injections.

Consistent with our retrograde tracing experiments, we observed fluorescently labeled cell bodies

in the ipsilateral ACx, including the primary ACx (Au1) and the dorsal and ventral secondary ACx

(AuD, AuV) (Fig. 2B-C). We additionally observed Pir-recipient cells in the temporal association

cortex (TeA) (Fig. 2B-C), consistent with previous studies (Tasaka et al., 2020). We quantified and

compared anterograde tdTomato expression across these cortical subregions and found an

increase in the number of labeled cells from dorsal-to-ventral auditory subregions (Fig. 2C,

*p=0.014, Friedman test). Furthermore, we quantified the cortical depth of the labeled cells and

found that Pir-recipient cells span all layers of ACx, but predominantly in layers 2/3 and 6 (Fig.

2D). To again confirm the accuracy of injection sites, we examined the extent of viral spread in

each mouse (Fig. 2E), ensuring localization of the virus injection to Pir with minimal spread into

surrounding regions. These anterograde tracing experiments further confirm the direct projection

from Pir to ACx and reveal that Pir-recipient ACx neurons span auditory cortical and layers and

subregions.

Taken together, our tracing experiments confirm a direct projection from pPir to ACx using multiple

retrograde and anterograde viral approaches. Our results are consistent with previous studies

showing a sparse but detectable Pir-to-ACx projection (Costa et al., 2017; Tasaka et al., 2020).

These results suggest an anatomical substrate by which olfactory sensory information may affect

auditory processing in ACx, pointing toward a possible mechanism for auditory-olfactory

integration.

***In vivo* electrophysiology for assessing auditory-olfactory integration in auditory cortex.**

Next, we sought to determine whether and how odors modulate auditory processing in the ACx

of awake mice. While previous studies have demonstrated olfactory modulation of sound

responses in ACx, most of these studies were performed in anesthetized animals using complex

odorants and sound stimuli, or in awake animals but influenced by a behavioral task (Cohen et

al., 2011; Cohen and Mizrahi, 2015; Gnaedinger et al., 2019; Gilday and Mizrahi, 2023). As such,

it remains unclear how auditory-olfactory integration may depend on brain state, stimulus complexity, and/or experience-dependent learning. We therefore sought to answer the basic question of whether and how a temporally coupled odor modulates ACx responses to simple sounds in untrained, awake mice. To answer this question, we designed a custom-made system combining sound stimuli with a respiration-triggered olfactometer to deliver sounds and odors to awake head-fixed mice with high temporal precision during *in vivo* electrophysiology (Fig. 3A). We recorded the activity of ACx neurons in response to combinations of odor (acetophenone or eucalyptus, 1%) and sound stimuli (unmodulated or amplitude-modulated [AM, 10 Hz] noise, 0 – 75 dB). We used the respiration-triggered olfactometer to control the timing of odor/sound stimulus delivery with the mouse's inhale (Fig. 3B). To examine auditory-olfactory integration in ACx, we targeted recordings to the ACx (Fig. 3C) and recorded the activity of single- and multi-units in response to sounds, with or without odor (Fig. 3D).

Because our anatomical experiments pointed toward the Pir-to-ACx pathway as a possible substrate for auditory-olfactory integration in ACx, we tested whether and how this pathway contributes to odor modulation of sound responses using an *in vivo* optogenetic approach. To selectively inactivate ACx-projecting Pir neurons in C57BL/6J or Cdh23 mice, we injected AAVrg-hSyn-Cre into ACx, and the cre-dependent soma-targeted inhibitory opsin stGtACR1 (AAV5-SIO-stGtACR1) (Mahn et al., 2018) into pPir (Fig. 3E). During *in vivo* recordings, we delivered 470 nm LED light through a fiber optic cannula implanted over pPir (Fig. 3E-F). Fig. 3G-H shows sound responses in an example ACx unit under the various trial conditions: odor enhances the sound responses, but this enhancement is eliminated by LED inactivation of ACx-projecting Pir neurons.

Odor modulates sound responses in auditory cortex.

First, we tested whether and how odors modulate responses to sound stimuli in ACx. We identified 'sound-responsive' and 'odor-modulated' units using a generalized linear model (GLM). In

362 response to unmodulated broadband noise, ~20% of sound-responsive units were significantly

363 modulated by odor (Fig. 4A). However, the effects of the odor on sound responses were

364 heterogenous: in response to unmodulated noise, we observed both enhancement (Fig. 4B-D,

365 left) and suppression (Fig. 4B-D, right) of sound responses by odor. To quantify the amount of

366 enhancement or suppression in odor-modulated units, we calculated an Odor Modulation Index

367 (OMI) for each unit for each sound level (Fig. 4C). We found that OMIs vary significantly across

368 sound level for enhanced units (Fig. 4C, left; *p=0.039, Friedman test), but not suppressed units

369 (Fig. 4C, right; n.s. p=0.326, Friedman test). Because we used acetophenone and eucalyptus as

370 odorants in our experiments and varied the odor between recordings, we tested for an odor-

371 dependent difference in enhancement or suppression. We compared the amount of odor

372 modulation (OMI) using acetophenone versus eucalyptus (Fig. 4E) and found that odor

373 modulation does not differ between odorants (Fig. 4E, left: n.s. p=0.93, Mann-Whitney test; Fig

374 4E, right: n.s. p=0.63, Mann-Whitney test). Therefore, we combined the groups here and in

375 subsequent experiments and analyses.

377 In addition, we tested the effect of odor on responses to 10 Hz amplitude-modulated (AM) noise

378 and performed the same analyses (Fig. 4F-J). In response to AM noise, we found a greater

379 proportion of odor-modulated sound-responsive units (~33%, Fig. 4F). We again observed both

380 enhancement and suppression of sound responses by odor (Fig. 4G-I), and that OMIs vary

381 significantly across sound level for enhanced units (Fig. 4H, left; *p=0.028, Friedman test).

382 Moreover, we again found that odor modulation does not differ between odorants (Fig. 4J, left:

383 n.s. p=0.59, Mann-Whitney test; Fig 4J, right: n.s. p=0.88, Mann-Whitney test).

385 In sound-responsive odor-modulated units, we tested the effect of odor on neuronal firing in the

386 absence of sound. We found an overall slight but significant enhancement of neuronal firing rate

387 in response to odor compared to control (Fig. 4K, **p=0.0042, Wilcoxon signed-rank test). This

result suggests that odors may impact ACx processing in the absence of sound, as well as in

response to sound stimuli. Specifically, odors enhance baseline neuronal firing, but either

enhance or suppress sound-evoked responses.

In response to odors, rodents actively change their respiratory patterns (sniffing) (Wesson et al.,

2009; Wachowiak, 2011). To evaluate the potential effects of respiration on the observed changes

in neuronal firing in our experiments, we compared the sniff frequency and amplitude during odor

versus control trials in each recording (Fig. 4L). We observed that while odor did not significantly

affect sniff frequency (Fig. 4L, left; n.s. p=0.28, paired t-test), odor reduced sniff amplitude (Fig.

4L, right; ***p=6.7e-7). This may reflect shallower breathing in response to the odor in our

experiments. Nevertheless, our results demonstrate odor-evoked changes in ACx activity,

enabling further dissection of the underlying mechanisms.

Together, these results demonstrate that: 1) olfactory stimuli modulate auditory processing in

awake mice, including enhancement of baseline firing as well as bidirectional modulation of

sound-evoked responses; and 2) olfactory stimuli enhance or suppress neuronal responses to

multiple types of sound stimuli, including both unmodulated and AM noise.

Piriform input to auditory cortex contributes to auditory-olfactory integration.

To test the role of the newly characterized Pir-to-ACx pathway (Fig. 1-2) in auditory-olfactory

olfactory integration, we combined *in vivo* electrophysiology in awake mice with optogenetics

specifically targeting ACx-projecting Pir neurons (Fig. 3). We again recorded sound responses

with or without odor, but now we also optogenetically inactivated ACx-projecting Pir neurons by

shining LED light through a cannula implanted over ipsilateral pPir (Fig. 3E-H). Consistent with

our previous experiments (Fig. 4), we observed both enhancement and suppression of sound

responses by odor in a subpopulation of ACx units (Fig. 5A). For both the enhanced and

suppressed units, we compared the amount of odor modulation, using the OMI, with or without

LED-inactivation of Pir neurons. Strikingly, we observed that for the enhanced units, the LED

significantly reduced odor modulation (**p=0.001, Wilcoxon signed-rank test), whereas odor

modulation in suppressed units was unaffected (n.s. p=0.96, Wilcoxon signed-rank test) (Fig. 5A).

We then examined the effect of the LED on odor modulation for each sound level (Fig. 5B-C) and

found that in enhanced units, the LED significantly reduced the amount of odor modulation (OMI)

at the highest sound level (Fig. 5C, left; 25 dB SNR: *p=0.042, Wilcoxon signed-rank test),

whereas in suppressed units the odor modulation was unaffected (Fig. 5C, right; 25 dB SNR: n.s.

p=0.73, Wilcoxon signed-rank test). We again repeated the same analyses for responses to AM

noise and observed similar results: the LED significantly reduced the amount of odor modulation

in enhanced units (**p=0.004, Wilcoxon signed-rank test), but not in suppressed units (n.s.

p=0.63, Wilcoxon signed-rank test) (Fig. 5E). For enhanced units, the LED significantly reduced

the amount of odor modulation at 15 dB SNR and 25 dB SNR (Fig. 5G, left; 15 dB SNR: *p=0.013;

25 dB SNR: *p=0.021; Wilcoxon signed-rank tests), whereas suppressed units were unaffected

(Fig. 5G, right; 15 dB SNR: n.s. p=0.10, Wilcoxon signed-rank test; 25 dB SNR: n.s. p=0.23,

paired t-test). These results demonstrate that odor *enhancement* of sound responses, but not

odor suppression, requires the Pir-to-ACx pathway, suggesting that this pathway provides an

excitatory drive.

In addition, we examined odor-driven enhancement of ACx neuronal firing in the absence of

sound, with or without *in vivo* optogenetic inactivation of Pir neurons. Consistent with our previous

experiments (Fig. 4K), in odor-modulated ACx units we observed that odor significantly enhanced

neuronal firing rate, compared to control (Fig. 5I, left; LED off: ***p=3.07e-7, paired t-test).

Strikingly, however, this enhancement was eliminated by the LED (Fig. 5I, left; LED ON, n.s. p>1,

Wilcoxon signed-rank test). When comparing the amount of odor modulation (OMI) with versus

without the LED, we found that the LED significantly reduced the odor-driven enhancement in the

absence of sound (Fig. 5I, middle; ***p=2.3e-7, Wilcoxon signed-rank test). This result suggests

that optogenetic inactivation of ACx-projecting Pir neurons eliminates odor-driven enhancement

of baseline ACx neuronal firing, and therefore that this enhancement requires the direct Pir-to-

ACx pathway.

Finally, we examined potential effects of respiration on the changes in neuronal firing we observed

in our optogenetic experiments (Fig. 5J). Consistent with our previous results (Fig. 4L), we

observed an odor-mediated reduction of sniff amplitude (Fig. 5J, right; LED off, ***p=6.2e-5,

Wilcoxon signed-rank test) but not sniff frequency (Fig. 5J, left; LED off, n.s. p=0.087, paired t-

test). Importantly, during trials with the LED, we observed similar effects (Fig. 5J, right: LED ON,

***p=0.0004, Wilcoxon signed-rank test; Fig. 5J, left: LED ON, n.s. p=0.076, paired t-test).

Moreover, when comparing respiration during odor trials with versus without the LED, we

observed that the LED did not affect sniff frequency (Fig. 5J, left; n.s. p=0.78, paired t-test) or

amplitude (Fig. 5J, right, n.s. p=0.35, Wilcoxon signed-rank test). These results critically

demonstrate that the optogenetic reduction of odor-driven enhancement is not due to any effects

of the LED on sniffing, and is likely specifically due to inactivation of the Pir-to-ACx pathway.

To further confirm that the optogenetic effects we observed (Fig. 5) were specifically due to

inactivation of Pir-to-ACx input via light activation of stGtACR1, and not potential other non-

specific effects of the LED light, we performed additional control experiments. Control Cdh23 mice

underwent an identical experimental procedure as the optogenetic mice, except we injected a

control virus into Pir (AAV5-Flex-tdTomato) instead of opsin (Fig. 6A). In control mice, LED light

did not significantly reduce odor modulation in enhanced units, for both unmodulated noise (Fig.

6B-C, left; n.s. p=0.11, Wilcoxon signed-rank test) and AM noise (Fig. 6B-C, right; n.s. 0.115,

Wilcoxon signed-rank test). Next, we examined the effect of the LED on enhanced units in the

controls. To quantify the effect of the LED on olfactory modulation, we calculated an LED

Modulation Index (Fig. 6E). For both unmodulated noise (Fig. 6E, left; **p=0.01, Mann-Whitney

test) and AM noise (Fig. 6E, right; *p=0.037, Mann-Whitney test), the LED significantly reduced

odor modulation in mice expressing stGtACR1 as compared to controls.

In control mice we again observed odor-driven enhancement of baseline neuronal firing in the

absence of sound (Fig. 6F, left; LED off, ***p=8.7e-9, paired t-test); however, this enhancement

persisted with the LED (Fig. 6F, left; LED ON, ***p=1.64e-6, Wilcoxon signed-rank test). When

comparing the amount of odor modulation (OMI) with versus without the LED, we found that the

LED did not affect the odor-driven enhancement in the absence of sound (Fig. 6F, right; n.s.

p=0.31, Wilcoxon signed-rank test). Additionally, when comparing respiration during the various

stimulation conditions, we again confirmed that the LED did not affect sniff frequency (Fig. 6G,

left; n.s. p>1, Wilcoxon signed-rank test) or amplitude (Fig. 6G, right; n.s. p=0.40, Wilcoxon

signed-rank test). These results confirm that the reduction of odor-driven enhancement we

observed in mice expressing stGtACR1 (Fig. 5) was specifically due to inactivation of the Pir-to-

ACx pathway.

Together, our results identify the role of the Pir-to-ACx pathway in auditory-olfactory integration.

Optogenetic inactivation of ACx-projecting Pir neurons selectively reduced odor-driven

enhancement of baseline ACx firing as well as enhancement of sound-evoked responses.

Therefore, the Pir-to-ACx pathway contributes to odor-mediated enhancement of sound

responses, but not suppression.

Discussion

In this study we tested the functional role of a direct projection from the piriform cortex to the

auditory cortex. We identified the Pir-to-ACx pathway using retrograde and anterograde viral

tracing, highlighting that this pathway arises primarily from posterior Pir (Fig. 1), and projects to

multiple ACx subregions and laminae (Fig. 2). Next, using *in vivo* electrophysiology in ACx, we

demonstrated that olfactory stimuli enhance baseline activity in ACx neurons and bidirectionally

modulate sound-evoked activity, either enhancing or suppressing sound responses (Figs. 3-4).

Finally, using *in vivo* optogenetics during electrophysiology, we found that the Pir-to-ACx

projection is necessary for odor-mediated enhancement of ACx activity and sound responses, but

is minimally engaged during suppression (Figs. 5-6). Our results reveal that the Pir-to-ACx

pathway shapes auditory-olfactory integration in ACx, contributing to odor-mediated

enhancement of sound processing.

Our anatomical results demonstrate that the Pir-to-ACx projection primarily arises from posterior

Pir (pPir), and projects across multiple ACx subregions and cortical layers. Its origination in pPir

is consistent with previous studies showing that pPir contains more associational connections

with other higher-order brain areas compared to anterior Pir (Bekkers and Suzuki, 2013). For

example, pPir is known to have interconnections with the prefrontal cortex, amygdala, and

entorhinal and perirhinal cortices, and is thought to encode odor quality (Johnson et al., 2000;

Zelano et al., 2011). Our results add ACx to the list of Pir-recipient regions and expand knowledge

of cross-modal cortical circuitry. Moreover, the distribution of Pir-recipient neurons spanning ACx

is consistent with previous studies showing that cortico-cortical intra-telencephalic neurons

project to/from multiple cortical layers including L2/3 and L6 (Shepherd, 2013; Harris and

Shepherd, 2015).

Notably, while previous studies have shown evidence for the Pir-to-ACx projection (Costa et al.,

2017; Tasaka et al., 2020), such findings have gone relatively underexplored. Our results are

consistent with these previous studies and provide further evidence for the existence and

functional impact of the Pir-to-ACx cross-cortical circuit. Although this projection is sparse, we

revealed this circuity using a rigorous approach comprised of four different viral strategies:

retroAAV-GFP (Fig. 1), retroAAV-Cre (Fig. 1), AAV1-Cre (Fig. 2), and selective labeling of ACx-

projecting Pir neurons for optogenetics (retroAAV-Cre and AAV5-stGtACR1, Fig. 3). It is possible

that a few Pir cells labeled as ACx-projecting (Fig. 1) or Pir-recipient ACx cells (Fig. 2) may instead

result from minor viral spread outside the target region (Fig. 1D-E, 2E); however, our redundant

approach increases our confidence in our results revealing the Pir-to-ACx pathway, which are

also consistent with some previous studies (Costa et al., 2017; Tasaka et al., 2020). Yet while our

experiments yielded robust and reproducible results using these strategies, the question remains

why some other research groups previously did not observe this projection. One explanation could

simply relate to differences in the tracer used (retroAAVs here versus fluorescein-labeled dextran

(Budinger et al., 2006; Budinger and Scheich, 2009)) and/or the animal model (mice here versus

gerbils (Budinger et al., 2006; Budinger and Scheich, 2009)). In addition, due to the sparseness

of the Pir-to-ACx projection, it is possible that this pathway may have gone unnoticed in some

previous studies in which it was not the target of investigation.

Some previous rodent studies examined auditory-olfactory integration in ACx, providing an

important foundation for understanding where and how the brain integrates auditory and olfactory

stimuli (Cohen et al., 2011; Cohen and Mizrahi, 2015; Gilday and Mizrahi, 2023). Our data build

upon this foundation and underscore modulation of sound responses by odors in naïve, passively

exposed, awake mice. Previous experiments using more complex stimuli, such as modulation of

ACx responses to pup vocalizations by pup odors, used acetophenone as a control odor stimulus,

showing no effect on sound responses (Cohen et al., 2011). That said, the Cohen et al. study was

performed in anesthetized animals, and it has since been well-established that neuronal

responses vary greatly in anesthetized versus awake states (Fontanini and Katz, 2008; Lesicko

et al., 2022). Our results suggest that olfactory modulation by simple odorants such as

acetophenone may depend on brain state. Furthermore, the combined olfactory and auditory

stimuli in previous studies were not bound precisely in time, and often involved other factors such

as behavioral training and associative learning that could have altered cross-modal coding

(Cohen et al., 2011; Cohen and Mizrahi, 2015; Gilday and Mizrahi, 2023). Our results in passively

exposed, untrained mice suggest that auditory-olfactory integration in ACx does not require

behavioral training. Future studies will be needed to further disentangle the complex relationships

between brain state, stimulus complexity, and experience-dependent learning involved in

auditory-olfactory integration and its neurophysiological underpinnings. Interestingly, an important

recent study has suggested that auditory-olfactory integration occurs in higher-order areas of

entorhinal cortex due to shared auditory and olfactory inputs (Wu et al., 2023). Our results add to

this growing body of literature and suggest that auditory-olfactory integration occurs at even

earlier stages in the sensory processing hierarchy than previously realized.

In sound-responsive odor-modulated units, we found that odor alone produces a slight

enhancement of firing rate even in the absence of sound (Figs. 4K, 5I, 6F). Thus, olfactory input

may affect both baseline and sound-evoked ACx activity. Our findings of both enhanced and

suppressed sound-evoked responses are consistent with previous studies in other modalities

such as in audio-visual integration, showing that visual stimuli evoke both excitation and

modulation of ACx (Bizley et al., 2007; Kayser et al., 2008; Atilgan et al., 2018). However, the

relatively modest degree of the odor-mediated effects on ACx activity observed here (Figs. 3-6),

coupled with the sparse nature of the Pir-to-ACx projection (Figs. 1-2), overall suggest that this

olfactory input to ACx likely sculpts auditory processing in ACx, rather than providing a strong

excitatory drive.

We found that Pir input to ACx contributes to odor-mediated enhancement of baseline and sound-

evoked responses, but not suppression of sound-evoked responses. It is intriguing to consider

what mechanism might account for this dichotomy. Because most long-range intra-telencephalic

projections are excitatory (Shepherd, 2013; Harris and Shepherd, 2015), odor enhancement may

be due to activation of ACx by direct Pir input, whereas odor suppression may arise from indirect

input via local circuitry or via other brain regions. In our anterograde tracing experiments

identifying Pir-recipient neurons, we observed that the TeA receives a large amount of Pir input

(Fig. 2), a finding that is consistent with previous studies (Tasaka et al., 2020, 2023) and suggests

that TeA is a region that might relay Pir input to ACx, ultimately affecting ACx activity. Together

these results pave the way for future studies to disentangle the relative contributions of direct Pir

input versus indirect pathways to auditory-olfactory integration in ACx, and how the Pir-to-ACx

pathway interacts with other multimodal and higher-order brain circuitry.

Finally, it is worth noting that previous studies of auditory-olfactory integration have largely

focused on parental behavior, including the effects of pup odors on auditory processing in

maternal mice, as well as pup retrieval (Cohen and Mizrahi, 2015; Marlin et al., 2015; Marlin and

Froemke, 2017; McRae et al., 2023). Moving forward, it will be interesting to determine the extent

to which the Pir-to-ACx pathway may contribute to parental behavior, pup retrieval, and other

complex behaviors. Although our present study has focused on the effects of simple neutral

odorants, future studies are poised to investigate the additional effects of ethologically relevant

odors, such as pup odors, social odors, food odors, or predator odors. Furthermore, auditory-

olfactory integration in ACx is known to undergo plasticity during motherhood as well as during

formation of odor-sound associations (Cohen and Mizrahi, 2015; Gnaedinger et al., 2019; Gilday

and Mizrahi, 2023). Drawing from these findings, we suggest that the Pir-to-ACx pathway serves

as a novel contributor to multimodal processing that may play a role in multisensory plasticity and

learning.

References

Angeloni C, Geffen MN (2018) Contextual modulation of sound processing in the auditory

cortex. Curr Opin Neurobiol 49:8–15 Available at:

www.sciencedirect.comwww.sciencedirect.com [Accessed November 18, 2020].

Atilgan H, Town SM, Wood KC, Jones GP, Maddox RK, Lee AKC, Bizley JK (2018) Integration

of Visual Information in Auditory Cortex Promotes Auditory Scene Analysis through

Multisensory Binding. Neuron 97:640-655.e4 Available at:

https://doi.org/10.1016/j.neuron.2017.12.034 [Accessed December 7, 2020].

Bekkers JM, Suzuki N (2013) Neurons and circuits for odor processing in the piriform cortex.

Trends Neurosci 36:429–438.

Bizley JK, Jones GP, Town SM (2016) Where are multisensory signals combined for perceptual

decision-making? Curr Opin Neurobiol 40:31–37 Available at:

http://dx.doi.org/10.1016/j.conb.2016.06.003 [Accessed November 18, 2020].

Bizley JK, Nodal FR, Bajo VM, Nelken I, King AJ (2007) Physiological and anatomical evidence

for multisensory interactions in auditory cortex. Cereb Cortex 17:2172–2189.

Budinger E, Heil P, Hess A, Scheich H (2006) Multisensory processing via early cortical stages:

Connections of the primary auditory cortical field with other sensory systems. Neuroscience

143:1065–1083.

Budinger E, Scheich H (2009) Anatomical connections suitable for the direct processing of

neuronal information of different modalities via the rodent primary auditory cortex. Hear

Res 258:16–27.

Choi I, Lee JY, Lee SH (2018) Bottom-up and top-down modulation of multisensory integration.

Curr Opin Neurobiol 52:115–122 Available at: https://doi.org/10.1016/j.conb.2018.05.002.

Cohen L, Mizrahi A (2015) Plasticity during motherhood: Changes in excitatory and inhibitory

layer 2/3 neurons in auditory cortex. J Neurosci 35:1806–1815.

Cohen L, Rothschild G, Mizrahi A (2011) Multisensory integration of natural odors and sounds in

the auditory cortex. Neuron 72:357–369.

Costa NM da, Martin KAC, Sägesser FD (2017) A weighted graph of the projections to mouse

auditory cortex. bioRxiv:228726.

Deroy O, Faivre N, Lunghi C, Spence C, Aller M, Noppeney U (2016) The Complex Interplay

between Multisensory Integration and Perceptual Awareness. Multisens Res 29:585–606.

Dewan A, Cichy A, Zhang J, Miguel K, Feinstein P, Rinberg D, Bozza T (2018) Single olfactory

receptors set odor detection thresholds. Nat Commun 9:1–12 Available at:

http://dx.doi.org/10.1038/s41467-018-05129-0.

Fontanini A, Katz DB (2008) Behavioral states, network states, and sensory response variability.

J Neurophysiol 100:1160–1168.

Gilday OD, Mizrahi A (2023) Learning-Induced Odor Modulation of Neuronal Activity in Auditory

Cortex. J Neurosci 43:1375–1386.

Gnaedinger A, Gurden H, Gourévitch B, Martin C (2019) Multisensory learning between odor

and sound enhances beta oscillations. Sci Rep 9:11236.

Gottfried JA, Dolan RJ (2003) The nose smells what the eye sees: Crossmodal visual facilitation

of human olfactory perception. Neuron 39:375–386.

Halene TB, Talmud J, Jonak GJ, Schneider F, Siegel SJ (2009) Predator odor modulates

auditory event-related potentials in mice. Neuroreport 20:1260–1264.

Han X, Xu J, Chang S, Keniston L, Yu L (2021) Multisensory-Guided Associative Learning

Enhances Multisensory Representation in Primary Auditory Cortex. Cereb Cortex:1–15.

Harris KD, Shepherd GMG (2015) The neocortical circuit: Themes and variations. Nat Neurosci

18:170–181.

Johnson DMG, Illig KR, Behan M, Haberly LB (2000) New features of connectivity in piriform

cortex visualized by intracellular injection of pyramidal cells suggest that "primary" olfactory

cortex functions like "association" cortex in other sensory systems. J Neurosci 20:6974–

6982.

Johnson KR, Tian C, Gagnon LH, Jiang H, Ding D, Salvi R (2017) Effects of Cdh23 single

nucleotide substitutions on age-related hearing loss in C57BL/6 and 129S1/Sv mice and

comparisons with congenic strains. Sci Rep 7:1–13.

Kayser C, Petkov CI, Logothetis NK (2008) Visual modulation of neurons in auditory cortex.

Cereb Cortex 18:1560–1574.

Lesicko AMH, Angeloni CF, Blackwell JM, De Biasi M, Geffen MN (2022) Cortico-Fugal

Regulation of Predictive Coding. Elife 11:1–29.

Mahn M, Gibor L, Patil P, Cohen-Kashi Malina K, Oring S, Printz Y, Levy R, Lampl I, Yizhar O

(2018) High-efficiency optogenetic silencing with soma-targeted anion-conducting

channelrhodopsins. Nat Commun 9 Available at: http://dx.doi.org/10.1038/s41467-018-

06511-8.

Marlin BJ, Froemke RC (2017) Oxytocin modulation of neural circuits for social behavior. Dev

Neurobiol 77:169–189.

Marlin BJ, Mitre M, D'Amour JA, Chao M V, Froemke RC (2015) Oxytocin enables maternal

behaviour by balancing cortical inhibition. Nature 520:499–504.

McRae BR, Andreu V, Marlin BJ (2023) Integration of olfactory and auditory cues eliciting

parental behavior. J Neuroendocrinol:1–11.

Meijer GT, Mertens PEC, Pennartz CMA, Olcese U, Lansink CS (2019) The circuit architecture

of cortical multisensory processing: Distinct functions jointly operating within a common

anatomical network. Prog Neurobiol 174:1–15 Available at:

https://doi.org/10.1016/j.pneurobio.2019.01.004.

Meijer GT, Montijn JS, Pennartz CMA, Lansink CS (2017) Audiovisual modulation in mouse

primary visual cortex depends on cross-modal stimulus configuration and congruency. J

Neurosci 37:8783–8796.

Meijer GT, Pie JL, Dolman TL, Pennartz CMA, Lansink CS (2018) Audiovisual integration

enhances stimulus detection performance in mice. Front Behav Neurosci 12:231 Available

at: https://www.frontiersin.org/article/10.3389/fnbeh.2018.00231/full [Accessed November

18, 2020].

Olofsson JK, Zhou G, East BS, Zelano C, Wilson DA (2019) Odor identification in rats:

Behavioral and electrophysiological evidence of learned olfactory-auditory associations.

eNeuro 6 Available at: https://doi.org/10.1523/ENEURO.0102-19.2019 [Accessed July 1,

2020].

Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris KD (2016) Kilosort: realtime spike-

sorting for extracellular electrophysiology with hundreds of channels. bioRxiv:061481

Available at: http://biorxiv.org/lookup/doi/10.1101/061481.

Seo H-S, Hummel T (2011) Auditory-Olfactory Integration: Congruent or Pleasant Sounds

Amplify Odor Pleasantness. Chem Senses 36:301–309 Available at:

https://academic.oup.com/chemse/article/36/3/301/506704 [Accessed November 24,

2020].

Seo H-S, Lohse F, Luckett CR, Hummel T (2014) Congruent Sound Can Modulate Odor

Pleasantness. Chem Senses 39:215–228 Available at:

https://academic.oup.com/chemse/article/39/3/215/502704 [Accessed November 24,

2020].

Shepherd GMG (2013) Corticostriatal connectivity and its role in disease. Nat Rev Neurosci

14:278–291 Available at: www.nature.com/reviews/neuro [Accessed October 15, 2020].

Shusterman R, Smear MC, Koulakov AA, Rinberg D (2011) Precise olfactory responses tile the

sniff cycle. Nat Neurosci 14:1039–1044.

Siegle JH, López AC, Patel YA, Abramov K, Ohayon S, Voigts J (2017) Open Ephys: An open-

source, plugin-based platform for multichannel electrophysiology. J Neural Eng 14.

Smith JJ, Shionoya K, Sullivan RM, Wilson DA (2009) Auditory stimulation dishabituates

olfactory responses via noradrenergic cortical modulation. Neural Plast 2009.

Tasaka G ichi, Feigin L, Maor I, Groysman M, DeNardo LA, Schiavo JK, Froemke RC, Luo L,

Mizrahi A (2020) The Temporal Association Cortex Plays a Key Role in Auditory-Driven

Maternal Plasticity. Neuron 107:566-579.e7 Available at:

https://doi.org/10.1016/j.neuron.2020.05.004.

Tasaka G ichi, Maggi C, Taha E, Mizrahi A (2023) The local and long-range input landscape of

inhibitory neurons in mouse auditory cortex. J Comp Neurol 531:502–514.

Varga AG, Wesson DW (2013) Distributed auditory sensory input within the mouse olfactory

cortex. Eur J Neurosci 37:564–571.

Wachowiak M (2011) All in a Sniff: Olfaction as a Model for Active Sensing. Neuron 71:962–973

Available at: http://dx.doi.org/10.1016/j.neuron.2011.08.030.

Wallace MT, Woynaroski TG, Stevenson RA (2020) Multisensory integration as a window into

orderly and disrupted cognition and communication. Annu Rev Psychol 71:193–219

Available at: https://doi.org/10.1146/annurev-psych-010419- [Accessed November 20,

2020].

Wesson DW, Verhagen J V., Wachowiak M (2009) Why sniff fast? The relationship between

sniff frequency, odor discrimination, and receptor neuron activation in the rat. J

Neurophysiol 101:1089–1102.

Wesson DW, Wilson DA (2010) Smelling sounds: Olfactory-auditory sensory convergence in the

olfactory tubercle. J Neurosci 30:3013–3021.

Williams E, Dewan A (2020) Olfactory detection thresholds for primary aliphatic alcohols in

mice. Chem Senses 45:513–521.

Wu T, Li S, Du D, Li R, Liu P, Yin Z, Zhang H, Qiao Y, Li A (2023) Olfactory–auditory sensory

integration in the lateral entorhinal cortex. Prog Neurobiol 221:102399 Available at:

https://doi.org/10.1016/j.pneurobio.2022.102399.

Zelano C, Mohanty A, Gottfried JA (2011) Olfactory Predictive Codes and Stimulus Templates

in Piriform Cortex. Neuron 72:178–187 Available at:

http://dx.doi.org/10.1016/j.neuron.2011.08.010.

Zhou H yu, Cheung EFC, Chan RCK (2020) Audiovisual temporal integration: Cognitive

processing, neural mechanisms, developmental trajectory and potential interventions.

Neuropsychologia 140 Available at:

https://doi.org/10.1016/j.neuropsychologia.2020.107396 [Accessed November 20, 2020].

Zingg B, Peng B, Huang J, Tao HW, Zhang LI (2020) Synaptic specificity and application of

anterograde transsynaptic AAV for probing neural circuitry. J Neurosci 40:3250–3267

Available at: https://doi.org/10.1523/JNEUROSCI.2158-19.2020 [Accessed October 15,

2020].

 Figures and Legends

Figure 1. Neurons in the ipsilateral piriform cortex directly project to the auditory cortex.

A) Injection of AAVrg-hSyn-eGFP into the left auditory cortex (ACx) of C57BL/6J mice (n=5) or Cdh23 mice (n=2). **B)** Representative fluorescent micrographs of ACx injection site (left), posterior piriform (pPir) ROI (middle), anterior piriform (aPir) ROI (right). Scale bars = 500 µm. **C)** Number of cells labeled with GFP in aPir and pPir (n=7 mice; n.s. Wilcoxon signed-rank test). **D)** Viral spread around ACx injection site. Colored outlines indicate extent of viral spread for each mouse (n=7 mice). From left to right, atlas overlays are bregma -3.08 mm, -2.54 mm, and -2.06 mm. **E)** Injection of AAVrg-hSyn-Cre into the left ACx of Ai14 mice (n=7). Colored outlines indicate extent of viral spread around each ACx injection site (n=7 mice). **F)** Representative fluorescent micrographs of ACx injection site (left), pPir ROI (middle), aPir ROI (right). Scale bars = 500 µm. **G)** Number of cells labeled with tdTomato in aPir and pPir (n=7 mice; *p<0.05, paired t-test).

Figure 2. Neurons that receive piriform cortical input span auditory cortical regions. A) Injection of AAV1-hSyn-Cre into the left piriform cortex (Pir) of Ai14 mice (n=6). B) Representative fluorescent micrographs of the Pir injection site (left) and auditory cortical ROIs (right). Scale bars = 500 μm. C) Number of cells labeled with tdTomato across auditory cortical regions (n=6 mice; *$p<0.05$, Friedman test). D) Histograms of the depth distribution (normalized cortical depth) of labeled cells in each auditory cortical region (all cells from 6 mice). E) Viral spread around Pir injection site. Colored outlines indicate extent of viral spread for each mouse (n=6 mice). From top to bottom, atlas overlays are bregma -1.94 mm, -1.46 mm, and -0.94 mm.

Figure 3. *In vivo* electrophysiology for assessing auditory-olfactory integration in auditory

cortex. A) Experimental setup for delivering combinations of odor/sounds to awake mice during

recordings. **B)** Stimulus timing using respiration-triggered olfactometer. **C)** Example post-hoc

visualization of probe location using DiI. **D)** Example raster plot from sound-responsive ACx unit.

Trials sorted by sound intensity and odor/control**. E)** Virus injections for selectively expressing

stGtACR1 in ACx-projecting Pir neurons in C57BL/6J or Cdh23 mice. Right: example opsin

expression in Pir and cannula placement for in vivo optogenetics using 470 nm LED. Scale bar =

500 μm. **F)** Stimulus timing during optogenetic experiments and example raster plot from ACx

unit. Trials sorted by sound intensity and odor/LED combination. **G)** Average firing rates (Hz) of

764   example unit shown in F, in response to various sound levels (left: 0 dB [no sound]; right: dB SNR

765   [relative to background noise of olfactometer air flow]) in control and odor, with and without LED

766   inactivation of PCx-to-ACx projections. **H)** Peristimulus time histogram (PSTH) of example unit

767   shown in F and G, in response to 15 dB SNR sound.

Figure 4. Odor modulates sound responses in auditory cortex. A) Left: Percentage of significantly odor-modulated units in ACx in response to unmodulated broadband noise, according to GLM. Right: percentage of Enhanced vs Suppressed odor-modulated units. n=5 mice (1 C57BL/6J, 4 Cdh23), 368 sound-responsive units, 293 'Sound only' units, 75 'Odor mod.' units, 33 Enhanced units, 42 Suppressed units. **B)** Average responses to sound levels (dB SNR relative to noise of background air flow), with Odor versus control, for Enhanced units (left) and Suppressed units (right). *p<0.05, **p<0.01, ***p<0.001, Wilcoxon signed-rank tests. **C)** Average Odor Modulation Index (OMI) for each sound level, for Enhanced units (left) and Suppressed units (right). *p<0.05, Friedman test. **D)** Average PSTHs in response to 5 dB SNR (left) and 15 dB SNR (right) sound. Dashed magenta line indicates odor valve opening time, gray lines indicate sound stimulus start and stop times. **E)** Average OMI for experiments with acetophenone (Acet.) versus eucalyptus (Euc.), for enhanced units (left; Acet. n=21 units, Euc. n=12 units) and suppressed units (right; Acet. n=17 units, Euc. n=25 units). n.s. Mann-Whitney tests. **F-J)** Same as A-E, for 10 Hz amplitude modulated noise. n= 5 mice, 397 sound-responsive units, 262 'Sound only' units, 135 'Odor mod.' units, 65 Enhanced units, 70 Suppressed units. **G)** *p<0.05, **p<0.01, ***p<0.001, Wilcoxon signed-rank tests. **H)** *p<0.05, Friedman test. **J)** Left: Acet. n=31 units, Euc. n=34 units. Right: Acet. n=33 units, Euc. n=37 units. n.s. Mann-Whitney tests. **K)** Left and middle: Average firing rate (z-scored) for odor-modulated units (10 Hz AM; units in F, n=135 units), in response to Odor versus control, in the absence of sound (0 dB). **p<0.01, Wilcoxon signed-rank test. Right: PSTH from an example odor-modulated unit, in the absence of sound (0 dB). **L)** Left: Average sniff frequency during Odor versus control trials (n=13 recordings). n.s. paired t-test. Right: Average sniff amplitude during Odor versus control trials (n=13 recordings). ***p<0.001, paired t-test.

Figure 5. *In vivo* optogenetic inactivation of ACx-projecting piriform neurons reduces odor-

mediated enhancement but not suppression. A) Left: Odor Modulation Index (OMI) with vs

without LED, for Enhanced and Suppressed ACx units in response to unmodulated broadband

noise. n= 6 mice (3 C57BL/6J, 3 Cdh23), 16 Enhanced units, 29 Suppressed units. Wilcoxon

signed-rank tests LED off vs LED ON. Right: Average OMI for Enhanced and Suppressed units,

LED off vs LED ON. **p<0.01, Wilcoxon signed-rank test. **B)** Average responses to various sound

levels (dB SNR relative to noise of background air flow): sound alone versus odor with or without

LED, for Enhanced units (left) and Suppressed units (right). *p<0.05, **p<0.01, Wilcoxon signed-

rank tests vs. 'Sound (control)'. **C)** Average OMI for each sound level, for Enhanced units (left)

and Suppressed units (right). *p<0.05, Wilcoxon signed-rank test. **D)** Average PSTHs in response

to 25 dB SNR sound stimulus. Dashed magenta and cyan lines indicate odor valve opening and

LED ON time, gray lines indicate sound stimulus start and stop times. **E-H)** Same as A-D, for 10

Hz amplitude modulated noise. n= 6 mice, 37 Enhanced units, 54 Suppressed units. E) **p<0.01,

Wilcoxon signed-rank test. **F)** *p<0.05, ***p<0.001, Wilcoxon signed-rank tests vs. 'Sound

(control)' **G)** *p<0.05, Wilcoxon signed-rank tests. **I)** Left: Average firing rate (z-scored) for odor

modulated units (10 Hz AM; units in E, n=91 units) in the absence of sound (0 dB), in response

to Odor versus control with LED off and LED ON. LED off: ***p<0.001, paired t-test. LED ON: n.s.

p>1, Wilcoxon signed-rank test. Middle: Average OMI in the absence of sound (0 dB), LED off vs

LED ON, ***p<0.001, Wilcoxon signed-rank test. Right: Average PSTH for odor-modulated units,

in the absence of sound (0 dB). **J)** Left: Average sniff frequency during Odor versus control trials,

with LED off and LED ON (n=16 recordings). LED off: n.s. LED ON: n.s. paired t-tests. LED off vs

LED ON: n.s. paired t-test. Right: Average sniff amplitude during Odor versus control trials (n=13

recordings). LED off: ***p<0.001, LED ON: ***p<0.001, Wilcoxon signed-rank tests. Odor (LED

off) vs Odor (LED ON): n.s. Wilcoxon signed-rank test.

Figure 6. Optogenetic inhibition of odor-mediated enhancement is specifically due to inactivation of ACx-projecting Pir neurons. A) Virus injections for optogenetic control experiments expressing control virus in ACx-projecting Pir neurons in Cdh23 mice. **B)** Left: OMI with vs without LED, for Enhanced and Suppressed ACx units in response to unmodulated broadband noise. n= 7 Cdh23 mice, 29 Enhanced units, 37 Suppressed units. Wilcoxon signed-rank tests LED off vs LED ON. Right: Same as Left, but in response to 10 Hz AM noise. n= 7 mice, 60 Enhanced units, 72 Suppressed units. Wilcoxon signed-rank tests LED off vs LED ON. **C)** Average OMI for Enhanced and Suppressed units, LED off vs LED ON., in response to unmodulated noise (left) and 10 Hz AM noise (right). ***$p<0.001$, Wilcoxon signed-rank test. **D)** Average PSTHs in response to 25 dB SNR sound, for unmodulated (left) and 10 Hz AM noise (right). **E)** LED Modulation Index for Enhanced units in response to unmodulated (left) and 10 Hz AM (right) noise, in stGtACR1 mice versus Control mice. Left: stGtACR1: n=16 units; Control: n=29 units; *$p<0.05$, Mann-Whitney test. Right: stGtACR1: n=37 units; Control: n=60 units; *$p<0.05$, Mann-Whitney test. **F)** Left: Average firing rate (z-scored) for odor-modulated units (10 Hz AM; units in B, right, n=132 units) in the absence of sound (0 dB), in response to Odor versus control with LED off and LED ON. LED off: ***$p<0.001$, paired t-test. LED ON: ***$p<0.001$, Wilcoxon signed-rank test. Right: Average OMI in the absence of sound (0 dB), LED off vs LED ON. n.s. Wilcoxon signed-rank test. **G)** Left: Average sniff frequency during Odor versus control trials, with LED off and LED ON (n=22 recordings). LED off: n.s. paired t-test, LED ON: n.s. Wilcoxon signed-rank test. LED off vs LED ON: n.s. Wilcoxon signed-rank test. Right: Average sniff amplitude during Odor versus control trials (n=22 recordings). LED off: ***$p<0.001$, LED ON: ***$p<0.001$, Wilcoxon signed-rank tests. Odor (LED off) vs Odor (LED ON): n.s. Wilcoxon signed-rank test.

841 **Table 1. Statistical comparisons.** Detailed values and statistical tests for all data figures.

Figure	Comparison	Test	n	Test statistic	p-value	Bonferroni corrected p-value
1C	Cell count, aPir vs. pPir	Wilcoxon signed-rank test	7 mice	W = 8	0.38	
1G	Cell count, aPir vs. pPir	Paired t-test	7 mice	t = -3.14	0.020	
2C	Cell count, by Region	Friedman test	6 mice	Q = 10.6	0.014	
4B, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound	Wilcoxon signed-rank test	33 units	W = 235	0.59	
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound			W = 130.5	0.013	
	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound			W = 154	0.023	
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound			W = 127	0.0093	
4B, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound	Wilcoxon signed-rank test	42 units	W = 357	0.65	
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound			W = 219	0.0061	
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound			W = 128	0.00015	
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound			W = 159	0.00043	
4C, left	Enhanced: OMI, by Sound Level	Friedman test	33 units	Q = 8.36	0.039	
4C, right	Suppressed: OMI, by Sound Level	Friedman test	42 units	Q = 3.46	0.33	
4E, left	Enhanced: OMI, Acet. vs. Euc.	Mann-Whitney test	21 units; 12 units	U = 129	0.93	
4E, right	Suppressed: OMI, Acet. vs. Euc.	Mann-Whitney test	17 units; 25 units	U = 232	0.63	
4G, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound	Wilcoxon signed-rank test	65 units	W = 983	0.56	
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound			W = 373	8.2e-6	

	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound			W = 531	0.0004	
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound			W = 789	0.064	
4G, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound	Wilcoxon signed-rank test	70 units	W = 819	0.013	
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound			W = 642	0.0012	
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound			W = 661	0.0007	
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound			W = 354	2.0e-7	
4H, left	Enhanced: OMI, by Sound Level	Friedman test	65 units	Q = 9.13	0.028	
4H, right	Suppressed: OMI, by Sound Level	Friedman test	70 units	Q = 3.88	0.27	
4J, left	Enhanced: OMI, Acet. vs. Euc.	Mann-Whitney test	31 units; 34 units	U = 568	0.59	
4J, right	Suppressed: OMI, Acet. vs. Euc.	Mann-Whitney test	33 units; 37 units	U = 624	0.88	
4K	Firing Rate, control vs. Odor	Wilcoxon signed-rank test	135 units	W = 3286	0.0042	
4L, left	Sniff frequency: control vs. Odor	Paired t-test	13 rec.	t = -1.12	0.28	
4L, right	Sniff amplitude: control vs. Odor	Paired t-test	13 rec.	t = 9.43	6.7e-7	
5A	Enhanced: OMI (LED off) vs. OMI (LED ON)	Wilcoxon signed-rank test	16 units	W = 9	0.001	
	Suppressed: OMI (LED off) vs. OMI (LED ON)		29 units	W = 201	0.96	
5B, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound (LED off)	Paired t-test	16 units	t = -1.14	0.27	0.54
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound (LED off)			t = -1.85	0.085	0.17
	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound (LED off)	Wilcoxon signed-rank test		W = 28	0.069	0.14
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 15	0.0042	0.0084

5B, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)	Wilcoxon signed-rank test	16 units	W = 61	0.74	1.49
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 62	0.78	1.56
	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 52	0.43	0.87
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 52	0.43	0.87
5B, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound (LED off)	Wilcoxon signed-rank test	29 units	W = 155	0.18	0.37
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 53	0.0006	0.0012
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 47	0.0006	0.0012
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 119	0.093	0.19
5B, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)	Wilcoxon signed-rank test	29 units	W = 206	0.81	1.63
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 54	0.0007	0.0014
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 57	0.0009	0.0018
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 92	0.0056	0.011
5C, left	Enhanced: OMI -5 dB SNR: LED off vs. LED ON	Paired t-test	16 units	t = 1.30	0.21	
	Enhanced: OMI 5 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test		W = 47	0.30	
	Enhanced: OMI 15 dB SNR: LED off vs. LED ON			t = 1.31	0.21	
	Enhanced: OMI 25 dB SNR: LED off vs. LED ON	Paired t-test		t = 2.22	0.042	
5C, right	Suppressed: OMI -5 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test	29 units	W = 105	0.044	

	Comparison	Test	Units	Statistic		
	Suppressed: OMI 5 dB SNR: LED off vs. LED ON	Paired t-test		t = 0.49	0.63	
	Suppressed: OMI 15 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test		W = 171	0.33	
	Suppressed: OMI 25 dB SNR: LED off vs. LED ON			W = 188	0.73	
5E	Enhanced: OMI (LED off) vs. OMI (LED ON)	Wilcoxon signed-rank test	37 units	W = 165	0.0041	
	Suppressed: OMI (LED off) vs. OMI (LED ON)		54 units	W = 636	0.63	
5F, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound (LED off)	Wilcoxon signed-rank test	37 units	W = 292	0.38	0.76
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 210	0.053	0.11
	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 150	0.0069	0.014
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 95	0.0002	0.0004
5F, left	Enhanced: -5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)	Wilcoxon signed-rank test	37 units	W = 222	0.08	0.16
	Enhanced: 5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 308	0.52	1.04
	Enhanced: 15 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 332	0.78	1.55
	Enhanced: 25 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 294	0.39	0.79
5F, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound (LED off)	Wilcoxon signed-rank test	54 units	W = 514	0.11	0.22
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 490	0.03	0.06
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 229	1.7e-5	3.3e-5
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound (LED off)			W = 243	1.7e-5	3.4e-5

Figure	Comparison	Test	Units	Statistic	p	p adj
5F, right	Suppressed: -5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)	Wilcoxon signed-rank test	54 units	W = 490	0.03	0.059
	Suppressed: 5 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 484	0.026	0.052
	Suppressed: 15 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 525.5	0.06	0.12
	Suppressed: 25 dB SNR: Sound (control) vs. Odor+Sound (LED ON)			W = 220	6.8e-6	1.4e-5
5G, left	Enhanced: OMI -5 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test	37 units	W = 249	0.19	
	Enhanced: OMI 5 dB SNR: LED off vs. LED ON	Paired t-test		t = 1.69	0.11	
	Enhanced: OMI 15 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test		W = 188	0.013	
	Enhanced: OMI 25 dB SNR: LED off vs. LED ON			W = 199	0.021	
5G, right	Suppressed: OMI -5 dB SNR: LED off vs. LED ON	Wilcoxon signed-rank test	54 units	W = 726	0.89	
	Suppressed: OMI 5 dB SNR: LED off vs. LED ON			W = 710	0.96	
	Suppressed: OMI 15 dB SNR: LED off vs. LED ON			W = 553	0.10	
	Suppressed: OMI 25 dB SNR: LED off vs. LED ON	Paired t-test		t = 1.23	0.26	
5I, left	Firing Rate, control vs. Odor (LED off)	Paired t-test	91 units	t = -5.70	1.5e-7	3.1e-7
	Firing Rate, control vs. Odor (LED ON)	Wilcoxon signed-rank test		W = 1957	0.59	1.18
5I, middle	OMI, LED off vs. LED ON	Wilcoxon signed-rank test	91 units	W = 786	2.3e-7	
5J, left	Freq.: control vs. Odor (LED off)	Paired t-test	16 rec.	t = -2.21	0.043	0.087
	Freq.: control vs. Odor (LED ON)			t = -2.28	0.038	0.076
	Freq.: Odor (LED off) vs. Odor (LED ON)			t = -0.88	0.39	0.78
5J, right	Amp.: control vs. Odor (LED off)	Wilcoxon signed-rank test	16 rec.	W = 0	3.1e-5	6.2e-5
	Amp.: control vs. Odor (LED ON)			W = 4	0.0002	0.0004

Figure	Comparison	Test	Units	Statistic	p	p adj.
	Amp.: Odor (LED off) vs. Odor (LED ON)			W = 41	0.18	0.35
6B-C, left	Enhanced: OMI (LED off) vs. OMI (LED ON)	Wilcoxon signed-rank test	29 units	W = 142	0.11	
	Suppressed: OMI (LED off) vs. OMI (LED ON)		37 units	W = 239	0.14	
6B-C, right	Enhanced: OMI (LED off) vs. OMI (LED ON)	Wilcoxon signed-rank test	60 units	W = 676	0.11	
	Suppressed: OMI (LED off) vs. OMI (LED ON)		72 units	W = 497	7.6e-6	
6E, left	LED Modulation Index, stGtACR1 vs. Control	Mann-Whitney test	16 units; 29 units	U = 125	0.012	
6E, right	LED Modulation Index, stGtACR1 vs. Control	Mann-Whitney test	37 units; 60 units	U = 828	0.037	
6F, left	Firing Rate, control vs. Odor (LED off)	Paired t-test		t = -6.29	4.3e-9	8.7e-9
	Firing Rate, control vs. Odor (LED ON)	Wilcoxon signed-rank test	132 units	W = 2218	8.2e-7	1.6e-6
6F, right	OMI, LED off vs. LED ON	Wilcoxon signed-rank test	132 units	W = 3882	0.31	
6G, left	Freq.: control vs. Odor (LED off)	Paired t-test		t = 0.85	0.40	0.81
	Freq.: control vs. Odor (LED ON)	Wilcoxon signed-rank test	22 rec.	W = 63	0.04	0.08
	Freq.: Odor (LED off) vs. Odor (LED ON)			W = 110	0.61	1.2
6G, right	Amp.: control vs. Odor (LED off)	Wilcoxon signed-rank test	22 rec.	W = 1	9.5e-7	1.9e-6
	Amp.: control vs. Odor (LED ON)			W = 18	0.0001	0.0002
	Amp.: Odor (LED off) vs. Odor (LED ON)			W = 86	0.20	0.40