Olfactory basis for essential amino acid perception during foraging in *Caenorhabditis elegans*

Ritika Siddiqui [1, ¥], Nikita Mehta [1, ¥], Gopika Ranjith[2, ¥], Marie-Anne Felix[3], Changchun Chen[4] and Varsha Singh[1, 2] [*]

[1] Division of Molecular Microbiology, School of Life Sciences, University of Dundee, UK

[2] Department of Developmental Biology and Genetics, Indian Institute of Science, Bengaluru, India

[3] Institute of Biology of the Ecole Normale Supérieure, Paris, France

[4] Department of Molecular Biology, Umeå University, Sweden

[¥] Equal authorship

[*] Correspondence: vsingh001@dundee.ac.uk

Abstract

Acquisition of essential nutrients through diet is crucial for the survival of animals. Dietary odors might enable foraging in animals for nutrient-rich diets. We asked if *Caenorhabditis elegans*, a bacterivorous nematode, uses olfactory cues to forage for essential amino acid-rich (EAA) diets. Using the native microbiome of *C. elegans,* we show that worms rely on olfaction to select leucine (EAA)-enriched bacteria. Using gas chromatography, we find that leucine-enriched bacteria produce isoamyl alcohol (IAA) odor in the highest abundance. Prior adaptation of worms to IAA diminishes the diet preference of worms. Several wild isolates of *C. elegans* display robust responses to IAA emphasizing its ecological relevance. We find that foraging for a leucine-enriched diet is mediated via the AWC olfactory neurons. Finally, we identify SRD-12, in AWC neurons, as a receptor for IAA and a mediator of dietary decisions in worms. Our study identifies a receptor-ligand module underpinning foraging behavior in *C. elegans*.

Keywords: *C. elegans*; microbiota; olfaction; odors; isoamyl alcohol; odor sensory neurons; GPCR

Introduction

Amino acids are central to metabolism, not only as building blocks for proteins but also as signaling molecules. During the course of evolution, animals have lost the ability to synthesize half of the proteogenic amino acids, making dietary intake of such amino acids indispensable [1]. Leucine (Leu), isoleucine (Ile), valine (Val), histidine (His), lysine (Lys), tryptophan (Trp), phenylalanine (Phe), methionine (Met), threonine (Thr), and arginine (Arg) constitute essential amino acids (EAAs) [2]. Several studies have investigated the mechanism of amino acid surveillance by cells and tissues. Leucine, glutamine, and arginine have been shown to activate the mTOR pathway and control protein translation and cell growth [3-9]. Altered EAA levels influence the longevity, behavior, and health of animals [10-12]. Increased levels of circulating branched-chain amino acids (BCAAs) are associated with an increased risk of obesity and diabetes in humans [13,14]. Restricted intake of tryptophan and methionine extends lifespan in rats, while a tryptophan-deficient diet reduces the weight of organs [15,16]. Methionine restriction is reported to extend the lifespan of *Drosophila* [17]. Central administration of a BCAA, leucine, reduces food intake in rats [18]. Extended lifespan upon diet restriction in silkworms is linked to enhanced utilization of EAAs [19]. However, it is not clear if animals have a way to modulate EAA-specific dietary intake. Do animals use specific sensing mechanisms to find an EAA-enriched diet? Several studies on feeding behaviors suggest that both invertebrates and vertebrates select nutrient-rich diets. For instance, rats prefer a protein-rich diet over a carbohydrate-rich diet [20]. Many animals display mechanisms to choose for or reject food with specific EAAs. Sparrows consume combination of different diets to get adequate levels of various EAAs, while gastropods reject diets deficient in methionine [21-23]. When maintained on an EAA-restricted diet, flies compensate by increasing feeding [24]. Kittens prefer diets rich in methionine and threonine, while rats select for a leucine-enriched diet [25]. Despite evidence that organisms seek EAA, specific molecular mechanisms for search behavior are not well understood.

Chemoperception, taste and olfaction, allow animals to quickly assess their diet utilizing G-protein coupled receptors (GPCRs). EAAs evoke taste perceptions such as

sweet (Thr, His, Leu, Phe, Trp) and bitter (Arg, Ile, Lys, Met), while non-EAAs can elicit umami (Glu) and sour (Asp) taste [26-28]. Amino acids that elicit sweet responses in humans, like threonine, are found to be attractive in rodents [29,30]. GPCRs are implicated in sensing different classes of amino acids and regulate foraging behavior in animals [31-35]. Metabotropic glutamate receptors (mGlu) sense glutamate, while the extracellular calcium-sensing receptor senses aliphatic, aromatic, and polar amino acids [34,36]. Invertebrates like *Caenorhabditis elegans* and *Drosophila melanogaster* also possess GPCRs-based amino acid sensing mechanisms [37]. *C. elegans* senses glutamate using mGlu type receptors, Mgl-1/2/3, while *D. melanogaster* utilizes a cationic amino-acid transporter, Dm *slif,* to sense arginine [38,39]. The goldfish 5.24, an odorant receptor, responds to arginine, while its human homolog, GPRC6A, responds to basic as well as aliphatic amino acids [40,41]. Olfactory GPCRs are known to sense various odors, including alcohol, esters, ketones, etc. and many of these are products of EAA catabolism. It is not known if animals rely on EAA-derived odors to find an EAA-rich diets.

The presence of an elaborate chemosensory system in soil-dwelling animals suggests the usage of olfactory cues for foraging. *C. elegans,* a bacterivorous nematode, relies heavily on its simple but effective chemosensory system composed of 302 neurons for foraging, although specific cues regulating this phenomenon are unknown [42,43]. Olfactory neurons of *C. elegans* express as many as 335 GPCRs [44], however, the odors stimulating them remain largely unknown. Like other invertebrates, *C. elegans* requires ten EAAs, of these, leucine, tryptophan, valine, arginine, and lysine extend their lifespan when supplemented at low concentration [45]. Does *C. elegans* rely on olfactory cues to sense diets with high(er) levels of EAAs? It is conceivable since some bacteria possess biosynthetic pathways to convert amino acids into odors, such as dimethyl sulfide, indole, isoamyl acetate, phenylethyl alcohol, and benzyl alcohol [46]. *C. elegans* can sense over 120 odors representing extensive structural diversity [47]. Do worms utilize some of them as foraging signals to find nutrient-enriched bacteria?

We hypothesized that certain odors produced by the microbiome of *C. elegans* are olfactory signals for EAA-rich bacteria and drive worms' foraging behavior. To test this hypothesis, we analyzed the preference of worms for natural microbiota species supplemented with individual EAAs. We found that odors drive worms' feeding

preference in favor of leucine supplemented diets. By carefully analyzing the odor bouquet produced by natural microbiota upon leucine supplementation, we identified isoamyl alcohol (IAA) as the most abundant odor in preferred diets. We show that IAA is produced from leucine using the Ehrlich degradation pathway in bacteria. IAA is an ecologically relevant odor and regulates the preference of worms for a leucine supplemented diet. Finally, we identified SRD-12, a GPCR expressed in AWC neurons, as the receptor for IAA. Taken together, SRD-12 regulates the dietary preference of worms to IAA-producing bacteria and thereby mediates the foraging behavior of *C. elegans* to leucine-enriched diets. Thus, IAA produced by bacteria is a diet quality code for leucine-enriched bacteria.

Results

Microbial odors drive the preference of *C. elegans* for leucine-enriched diets

To understand if *C. elegans* used olfactory signals to distinguish between standard and EAA-enriched diets, we performed 'odor-only' diet preference assays using natural microbiota of *C. elegans*, CeMbio, and laboratory food, *Escherichia coli* OP50. CeMbio is a collection of 12 bacteria that best reflects the metabolic complexity of the microbes in the natural habitat of worms (see Table S1). N2 or wild type (WT) worms were allowed to choose between odors from each strain grown on separate sections, EAA supplemented (+ EAA) or unsupplemented (- EAA), of a tripartite plate (schematic in Figure 1A). Of the ten EAAs tested, WT worms showed a preference only for bacteria on leucine supplemented (+ LEU) over unsupplemented (- LEU) for three microbial strains, namely, *Enterobacter hormaechei* (CEent1), *Lelliottia amnigena* (JUb66)*,* and *Sphingobacterium multivorum* (BIGb0170) (Figure 1B). However, worms were not able to distinguish between supplemented and unsupplemented bacteria for the remaining nine EAAs (Figures 1C, 1D and S1A-S1G). We found that worms did not respond to leucine at three different concentrations in a chemotaxis assay, suggesting that worms are incapable of sensing leucine directly (Figure S1H). These results suggested that worms can forage for diets enriched in a specific EAA, leucine, using odors produced by these bacteria.

We next asked whether worms prefer their native microbiome over laboratory food *E. coli* OP50. We performed 'odor-only' diet preference assays to test if worms can distinguish between CeMbio and *E. coli* OP50 using odors (schematic in Figure S2A). We found that the preference indices (PI) of worms for CEent1, JUb66, and BIGb0170 were positive, making them the preferred diets, while neutral or negative for other strains such as *Pseudomonas lurida* MYb11 (Figure S2B). These findings suggested that worms not only rely on odors to choose between two bacteria, but also to find leucine-enriched bacteria.

Figure 1. *C. elegans* relies on odors to select leucine-enriched bacteria
(A) Schematic representation of 'odor-only' diet preference assay.
Preference index (PI) of WT worms in an 'odor-only' diet preference assay (indicated as ☺) for individual diet supplemented with (B) 5 mM leucine (+ LEU), (C) 5 mM isoleucine (+ ILE), and (D) 5 mM valine (+ VAL). Error bars indicate SEM (n≥15). Also see Figure S1.

Isoamyl alcohol odor is a signature for a leucine-enriched diet

We hypothesized that bacterial odor(s) enriched upon leucine supplementation are likely to be drivers of foraging behavior in worms. We identified all the odors in the headspace of individual bacteria using solid-phase microextraction followed by gas chromatography-mass spectrometry (GC-MS/MS). By examining levels of individual odors in leucine supplemented and unsupplemented bacteria, we found that several odors were present in the headspace of bacteria (Figures 2A, 2B, and S3A-S3D). However, isoamyl alcohol (IAA) was the most abundant and the only shared odor in the worms' preferred diets, CEent1, JUb66, and BIGb0170 (Figures 2A-2C and S3A-S3D). We found that under standard conditions, IAA constituted 40-70% of the headspace of preferred bacteria and remarkably increased up to 90% upon leucine supplementation (Figure 2C). We also quantified leucine-dependent increase in IAA levels in the headspace of preferred bacteria (Figure S3E, see methods). We found that abundance of IAA significantly increased in CEent1, JUb66, and BIGb0170 by 1.2 - 4 folds upon leucine supplementation (Figures 2D, S3F and S3G).

Microbes can produce short-chain alcohols from carbon sources as well as branched-chain amino acids [46]. Leucine can be catabolized to IAA via the Ehrlich degradation pathway in three enzymatic steps. The first step is catalyzed by IlvE, a transaminase, which converts leucine to α-Ketoisocaproate (schematic in Figure 2E). The *ilvE* gene is present in the genome of CEent1, JUb66, and BIGb0170 [48]. We predicted that leucine supplementation would result in the upregulation of *ilvE* in a substrate-dependent manner. Using qRT-PCR, we found that transcript levels of *ilvE* were 2-fold higher in leucine supplemented CEent1 over unsupplemented (Figure 2F). This indicated that the Ehrlich degradation pathway is functional in IAA-producing bacteria and further stimulated in response to leucine. Altogether, these findings show that IAA produced via Ehrlich degradation is a signature for a leucine-enriched bacteria.

Figure 2. Leucine supplementation boosts isoamyl alcohol levels via Ehrlich degradation pathway

GC-MS/MS profile of odors produced by CEent1 under (A) leucine supplemented and (B) leucine unsupplemented conditions. Unmarked peaks represent masses contributed by fiber or media alone (n≥3).

(C) Heat map representing the relative abundance of various odors in the headspace of CEent1, JUb66, and BIGb0170 with and without leucine supplementation.

(D) Absolute abundance of IAA (in moles) produced by CEent1 under leucine supplemented and unsupplemented conditions. * P ≤ 0.05 as determined by a two-tailed unpaired *t*-test.

(E) Schematic of Ehrlich degradation pathway.

(F) Fold change of transcript levels of *ilvE* under leucine supplemented over unsupplemented conditions for CEent1.

Error bars indicate SEM (n≥3).

AWC odor sensory neurons facilitate the diet preference of *C. elegans* for leucine-enriched diets

Worms use two pairs of odor sensory neurons, AWA and AWC, to sense attractive olfactory cues [47,49,50]. To identify the odor sensory neurons that mediate preference for leucine-enriched diets, we studied the response of *odr-7* worms, which do not have functional AWA neurons, and AWC ablation (-) worms [51,52]. We found that the preference of worms for leucine supplemented diets was dramatically lost in AWC(-) worms but was retained in *odr-7* worms (Figures 3A-3C). This suggested that AWC neurons play a crucial role in sensing odors from leucine-enriched diets. We also tested the role of these neurons in mediating diet preference between preferred CeMbio bacteria and *E. coli* OP50. We found that AWC(-) worms showed reduced preference for CEent1, JUb66 and BIGb0170 over *E. coli*, while *odr-7* mutants had slightly diminished preference for CEent1 and BIGb0170 over *E. coli* (Figures S4A-S4C). This suggested AWC, predominantly, and AWA odor sensory neurons contribute to the diet preference of worms for specific microbes in their natural habitat (Figure S4D). The finding that diet preference entails two pairs of neurons suggests that multiple odors constitute foraging signal enabling worms to find diet bacteria.

To identify the constituents of foraging signal, we sampled the headspace of each CeMbio bacteria. Each bacterium produced a unique bouquet of odors composed of chemically diverse molecules, including alcohols, aldehydes, ketones, esters, and carboxylic acids (Table S1). To identify foraging cues amongst odors in the headspace of preferred bacteria, we examined the response of WT worms to ten different odors in chemotaxis assays (schematic in Figure S5A). We found that WT worms were attracted to IAA, acetoin (ACE), isovalerate (ISV), isobutanol (ISB), and phenylethyl alcohol (PEA) in a dose-dependent manner (Figures S5B-S5F). Worms showed attraction to butyl acetate (BA) and methyl isovalerate (MIV) only at specific concentrations but did not display a clear dose-response (Figures S5G and S5H). Worms showed no response to 2,3-butanediol (BDL), isoamyl acetate (ISA), and indole (IND) (Figures S5I-S5K). This is the first report of acetoin and methyl isovalerate as attractants for *C. elegans*. These results show that the headspace of preferred bacteria contains multiple attractive odors for *C. elegans*.

To identify neurons responsible for sensing attractive odors produced by preferred bacteria, we used the *odr-7* and AWC(-) worms in chemotaxis assays for each of the seven attractive odors. We found that AWC(-) worms had severely diminished responses to IAA, ISV, ISB, PEA, BA, and MIV (Figures 3D and 3F-3J). Acetoin, on the other hand, was sensed by AWA neurons as *odr-7* showed reduced response to acetoin (Figure 3E). Taken together, our findings revealed that worms sense an extensive repertoire of chemically diverse odors produced by preferred bacteria as attractants, predominantly using AWC neurons (summary in Figure 3K).

Figure 3. Odors sensed through AWC neurons mediate the diet preference of *C. elegans*
Preference index (PI) of WT, *odr-7*, and AWC(-) worms for (A) CEent1, (B) JUb66, and (C) BIGb0170 supplemented with leucine over unsupplemented conditions. ☹ symbol indicates 'odor-only' preference assays.
Chemotaxis index (CI) of WT, *odr-7*, and AWC(-) worms for (D) isoamyl alcohol (IAA), (E) acetoin (ACE), (F) isovalerate (ISV), (G) isobutanol (ISB), (H) phenylethyl alcohol (PEA), (I) butyl acetate (BA), and (J) methyl isovalerate (MIV)
(K) Summary schematic representing the role of AWA and AWC odor sensory neurons in 'odor-only' diet preference for leucine-enriched diets and chemotaxis assays.
Significant differences are indicated as * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, and **** $P \leq 0.0001$ determined by one-way ANOVA followed by post hoc Dunnett's multiple comparison test. Error bars indicate SEM (n=15).

Isoamyl alcohol drives foraging behavior in *C. elegans*

To identify the most relevant foraging signal for *C. elegans*, we took advantage of the well-established odor adaptation assay using each of the attractive odors in its diets (schematic in Figure 4A). Prolonged exposure of *C. elegans* to an odorant is known to result in the loss or reduction of the ability of worms to sense the same odorant in a subsequent exposure [53]. We performed a modified adaptation assay where worms were exposed to a bouquet of odors from a bacterial lawn followed by testing their chemotaxis response to individual attractive odors. Worms adapted to odors from CEent1 lawns had dramatically reduced responses to IAA, isobutanol, and phenylethyl alcohol compared to naïve worms (Figures 4B, 4E, and 4F). Worms adapted to odors from JUb66 and BIGb0170 lawns also had dramatically reduced response to IAA (Figure 4B). While adaptation to bacterial odors did not influence worms' response to acetoin, isovalerate, butyl acetate and methyl isovalerate (Figure 4C, 4D, 4G and 4H). These findings suggested that the odor bouquets of preferred bacteria have levels of IAA high enough to modulate chemoperception in *C. elegans.*

If IAA is the most relevant foraging signal, we predicted that prior adaptation to IAA will diminish the diet preference of *C. elegans*. To test this, we adapted worms to IAA odor and checked their diet preference (schematic in Figure S6A). IAA-adapted worms showed a reduced chemotaxis response to IAA, indicating that the adaptation regimen is robust (Figure S6B). We found that IAA-adapted worms displayed a reduced preference for CEent1, JUb66, or BIGb0170 over *E. coli* OP50 compared to naïve worms (Figures S6C-S6E). These results indicated that IAA is the predominant olfactory cue that determines the diet preference of *C. elegans.*

Figure 4. Isoamyl alcohol regulates the diet preference of worms

(A) Schematic representation of the adaptation regimen followed by chemotaxis assays.

Chemotaxis index (CI) for naïve worms and worms adapted with CEent1, JUb66, or BIGb0170 odors to (B) isoamyl alcohol (IAA), (C) acetoin (ACE), (D) isovalerate (ISV), (E) isobutanol (ISB) (F) phenylethyl alcohol (PEA), (G) butyl acetate (BA), and (H) methyl isovalerate (MIV). Dark gray bars indicate naïve worms and light gray bars indicate worms adapted to bacterial odors.

Significant differences are indicated as * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, and **** P ≤ 0.0001 determined by one-way ANOVA followed by post hoc Dunnett's multiple comparison test. Error bars indicate SEM (n=15).

IAA is an ecologically relevant odor for *C. elegans*

If an odor is relevant as a foraging cue in nature, the ability to sense it must be under positive selection in wild *C. elegans* populations. We anticipated that most wild isolates of *C. elegans* will have a robust response to IAA but not to other attractants reported in this study. To test this hypothesis, we studied the response to IAA in nine wild isolates of *C. elegans* collected from geographically distinct regions (Figure 5A and Table S1) [54,55]. Additionally, we tested their chemotaxis response to diacetyl, a known food cue, and two other attractants (PEA and ACE) identified in this study. We found that all the strains displayed robust response to IAA comparable to WT worms (Figure 5B). The chemotaxis response of these strains to diacetyl was also quite robust (Figure 5C). However, the chemotaxis response to PEA and acetoin were variable within and across wild isolates suggesting they are not crucial for foraging (Figures 5D and 5E). The robust response of wild isolates to IAA supports the notion that the ability of *C. elegans* to sense IAA has been selected during evolution. These findings suggest that IAA is an ecologically relevant foraging cue for *C. elegans*.

Figure 5. Robust chemoperception of isoamyl alcohol in wild isolates of *C. elegans*

(A) World map representing the distinct geographical locations (source) of wild isolates of *C. elegans* used in this study.

Chemotaxis index (CI) of WT and wild isolates of worms for (B) IAA, (C) diacetyl (DA), (D) phenylethyl alcohol (PEA), and (E) acetoin (ACE).

Significant differences are indicated as * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, and **** P ≤ 0.0001 determined by one-way ANOVA followed by post hoc Dunnett's multiple comparison test. Error bars indicate SEM (n=15).

SRD-12 senses isoamyl alcohol and facilitates foraging in *C. elegans*

C. elegans genome encodes for ~1300 putative GPCRs, many of which are predicted to sense soluble and olfactory cues [56]. Since the preference for IAA and leucine-enriched

diets is mediated by AWC odor sensory neurons (Figures 3A-3D), we hypothesized that the receptor for IAA is a GPCR expressed in AWC neurons. Indeed, calcium response to IAA has been reported in AWC neurons [57,58]. Using single-cell RNAseq data for *C. elegans* neurons, available at CeNGEN [44], we listed 18 putative GPCRs expressed in AWC^{on} and AWC^{off} for further analysis (Figure 6A). We tested response to IAA in 15 strains harboring mutants in one or more of these GPCRs (see Table S1) [59]. One of these strains, CHS1169, had a significantly reduced response to IAA compared to WT worms (Figure 6A). Of the six GPCRs edited in the strain, *srd-12* had the highest expression in AWC^{off} odor sensory neuron [44]. Using CRISPR/Cas9-based approach, we generated two *srd-12* deletion lines, VSL2401, and VSL2402, carrying mutation in *srd-12* gene (see methods and Table S1). We recorded the chemotaxis response of WT and mutant worms to IAA (schematic in Figure 6B) and found they were indeed defective in their response to IAA (Figures 6C-6F). By analyzing the locomotion of WT, VSL2401, and VSL2402 worms, in the chemotaxis arena, we determined quantitative differences in their response to IAA (Figure S7). We observed that VSL2401 and VSL2402 strains had speeds comparable to WT worms and traveled distances similar to WT worms suggesting that *srd-12* mutants do not have locomotory defects (Figures S7A and S7B). We found that both *srd-12* mutant strains had significantly reduced chemotaxis response to IAA compared to WT worms (Figure 6F). However, *srd-12* mutants had a comparable response to WT worms for acetoin and phenylethyl alcohol (Figures 6G and 6H). As expected, *srd-12* mutant strains had diminished preference for CEent1 over *E. coli* OP50 in the diet preference assay consistent with the idea that IAA is the most relevant foraging signal (Figure 6I). Taken together, our findings show that *C. elegans* senses foraging signal, IAA, using newly identified GPCR, SRD-12.

Figure 6. SRD-12 GPCR mediates isoamyl alcohol sensing and diet preference in *C. elegans*

(A) List of GPCRs that are highly expressed in AWC neurons. Chemotaxis index (CI) of WT and GPCR edited strains to 1:1000 IAA (n≥3).

(B) Schematic for chemotaxis assay plate used for movement track analysis.

Movement tracks of 5 animals in a chemotaxis arena with 1:1000 IAA for 15 minutes, for (C) WT, (D) VSL2401, and (E) VSL2402. Also, refer to the video S2. Individual color represents the track for a single worm.

Chemotaxis index (CI) of WT, VSL2401, and VSL2402 worms for (F) 1:1000 IAA, (G) 500 mM acetoin (ACE), and (H) absolute phenylethyl alcohol (PEA).

(I) Preference index (PI) of WT, VSL2401, and VSL2402 worms for a preferred diet, CEent1, over *E. coli* OP50. ⊘ symbol indicates 'odor-only' preference assays.

Significant differences are indicated as * $P \leq 0.05$, ** $P \leq 0.01$, and **** $P \leq 0.0001$ determined by one-way ANOVA followed by post hoc Dunnett's multiple comparison test. Error bars indicate SEM (n≥15).

Based on our study, we propose a model describing an odor-dependent mechanism for sensing EAA-enriched bacteria in *C. elegans*. We show that IAA is an olfactory signal for leucine-enriched diet. AWC olfactory neurons of *C. elegans* allow worms to choose for IAA producing bacteria. Bacteria preferred by worms have the metabolic capacity to produce IAA from leucine via the Ehrlich degradation pathway. SRD-12, a GPCR in AWC neurons, mediates foraging for bacteria in *C. elegans* by facilitating chemoperception of IAA (Figure 7).

Figure 7. Model depicting odor-based foraging strategy used by *C. elegans*
C. elegans prefers leucine-enriched diets in an odor-dependent manner. Preferred bacteria catabolize leucine, an EAA, to produce IAA via the Ehrlich degradation pathway. SRD-12 GPCR expressed in AWC odor sensory neurons mediates the foraging behavior of *C. elegans* by sensing IAA.

Discussion

Our study provides evidence for an odor-dependent mechanism for foraging in *C. elegans*. Using CeMbio, a model microbiome for *C. elegans*, we show that worms prefer

leucine supplemented bacteria, namely *Enterobacter hormaechei, Lelliottia amnigena*, and *Sphingobacterium multivorum*, in an odor-dependent manner. Leucine-enriched diets produce significantly higher levels of IAA odor, making up to 90% of their headspace. These bacteria produce IAA from leucine using the Ehrlich degradation pathway. Preference for leucine-enriched bacteria solely depends on the AWC odor sensory neurons of *C. elegans*. Although the preferred bacteria produce several AWC-sensed odors, IAA exclusively influences the diet preference of worms. Wild isolates of *C. elegans*, representing nine geographically distinct locations on Earth, respond robustly to IAA underscoring its relevance as a foraging signal in nature. *C. elegans* utilizes SRD-12, a GPCR primarily expressed in AWC neurons, to sense IAA and forage for the preferred bacteria. Thus, IAA-SRD-12 represents a ligand-receptor module used by *C. elegans* to forage for bacteria in its natural environment.

Is olfaction a major contributor to foraging behavior in animals in general? Artificial flowers, containing nectars with high amino acid content, are better at attracting butterflies [60]. Honeybees prefer to consume artificial nectar rich in proline [61]. Hummingbirds, moths, and ants are attracted to flowers emitting benzyl acetone, an odor synthesized from an EAA, phenylalanine [62-65]. For odors to serve as an attractant, the said odor must serve as a proxy or code for a needed primary metabolite, abundant in carbon, nitrogen, phosphorus, or micronutrients. In such a scenario, it is easy to envisage that each attractive cue might be a code for a specific essential nutrient.

Is isoamyl alcohol a signal for essential nutrient for *C. elegans* and other foraging animals. IAA was one of 121 odors reported to be sensed by *C. elegans* several decades ago [47], but its ecological relevance was unclear to date. Our study suggests that IAA is the primary cue that drives foraging in favor of leucine-enriched diets in *C. elegans*. Does IAA reflect the metabolic state of bacteria producing it? IAA is indeed derived from a primary metabolite leucine, an essential branched chain amino acid for animals [66]. When bacteria were provided with additional leucine, the absolute abundance of IAA increases significantly in three different bacteria examined (Figures 2C and S3). The arrival of feeding insects to ripening fruits such as bananas coincides with for production of IAA and isoamyl acetate in the fruit via Ehrlich degradation pathway [67]. The presence of Ehrlich degradation pathway in specific microbes and many insect attracting plants underscores the importance of odors in mediating

interkingdom interactions. IAA, in combination with other odors, has been reported to attract *Drosophila melanogaster* and help in establishing a fly-fruit-yeast relationship [68-74]. Why do animals need BCAA? Not only is leucine a major component of *C. elegans* proteome, intracellular accumulation of BCAA including leucine prolongs the lifespan of *C. elegans* [75]. The lack of direct leucine perception (Figure S1H in this study) but presence of IAA perception ability, via SRD-12, suggests that worms rely on IAA sensing to identify bacteria with leucine. Additional natural microbiotas of *C. elegans* (not part of CeMbio) which produce IAA have also been reported to attract worms [76,77]. Worms also prefer odors derived from other BCAA such as isobutanol derived from valine via Ehrlich degradation (Figure 4E) [78-80]. Phenylethyl alcohol (PEA in this study) can be produced by microbial degradation of phenyl alanine, another essential amino acid that animals require for their proteins as well as for the synthesis of neurotransmitter dopamine. All these reports support the idea that animals use olfactory cues to identify diets rich in essential amino acids.

Animals use GPCRs to sense soluble and olfactory cues present in their environment. Rats utilize taste receptors T1R1/T1R3 for sensing alanine [81]. *C. elegans* can also sense lysine and histidine [82] but not leucine (this study). Several studies have implicated olfactory GPCRs in regulating animal behavior. In termites, olfactory co-receptor OCRO influences foraging by enabling pheromone sensing [83]. Or83b, an odorant receptor in *D. melanogaster* regulates lifespan [84]. Six odor sensory neurons of *C. elegans* express hundreds of GPCRs, most of them remain orphan [44]. ODR-10 expressed in AWA odor sensory neurons regulates foraging behavior in *C. elegans* for lactic acid bacteria which produce diacetyl odor [85]. AWC odor sensory neurons express as many as 171 GPCRs, of which 18 GPCRs display strong expression. These receptors likely mediate the foraging behavior of worms. One of these, STR-2, senses 2-heptanone odor produced by *Bacillus nematocidal* [86]. STR-2 also regulates lipid accumulation and lifespan in a temperature-dependent manner [87]. The ligand(s) for the remaining 17 receptors were unknown. The discovery of one of these, SRD-12, as receptor for sensing IAA (this study) suggests that the ligand for other receptors may also be found in the headspace of the *C. elegans* microbiome. Our study provides a framework for the identification of ligands for orphan GPCR, by searching an ecologically relevant repertoire, the microbiome of an animal.

Methods

Experimental model and subject details

Strains and growth media

C. elegans strains used in this study are listed in see Table S1. All strains were maintained as hermaphrodites at 20°C on nematode growth media (NGM) plates seeded with *E. coli* OP50, as originally described [88]. All the CeMbio strains used in this study (see Table S1) were grown on Difco Luria-Bertani (LB) media and maintained at 25°C, as described by Dirksen *et al.* [48]. A list of all the reagents used in this study is provided in Table S1.

CRISPR approach for gene editing

Homology-directed integration of the ssDNA oligo was done in the *srd-12* gene using a CRISPR/Cas9-based approach as described previously [89]. ssDNA oligo was designed to contain two stop codons along with a restriction site between two 35 bp long homology arms flanking the PAM sites of the targeted gene. ssDNA oligo was incubated with optimized RNP complexes of Cas9, crRNA, and tracrRNA. The mix was injected in the gonad of *C. elegans* along with *rol-6* [pRF4::rol-6(su1006)] marker plasmid. The F1 progenies were segregated and genotyped using PCR, followed by restriction digestion for the introduction of mutation. *srd-12* CRISPR mutants were back-crossed three times with N2 Bristol WT strain to generate VSL2401 and VSL2402.

Diet preference assay

For diet preference assays, individual CeMbio bacterium (test bacterium) and *E. coli* (control bacterium) were grown in LB broth overnight at 25°C and adjusted to an OD_{600} of 1. Standard diet choice assays were conducted on 90 mm NGM plates, wherein 25 µl of cultures of the test and control bacteria were spotted 1 cm away from the periphery of the plate on diametrically opposite points of the dish. These plates were incubated at 25°C for 12 hours. Gravid adult worms were washed thrice with S-basal buffer (100 mM NaCl, 5.75 mM K_2HPO_4, 44 mM KH_2PO_4). 50-70 worms were placed in the center of the assay plate and incubated at 25°C. The preference index (PI) was calculated after 3 hours using the following equation:

Preference Index (PI) = (Worms on the test lawn – Worms on the control lawn) / Total worms

'Odor-only' diet preference assays were performed in tripartitioned plates, wherein one compartment was filled with buffered agar or BA (1 mM $CaCl_2$, 1 mM $MgSO_4$, 5 µg/ml cholesterol, 25 mM KPO_4 buffer, and 2% agar) and other two compartments were filled with NGM. All the assay plates were air-dried for 70-90 minutes. For EAA supplementation, one of the NGM-filled compartments was supplemented with individual EAA at 5 mM concentration (+ EAA), while the other was left unsupplemented (- EAA). All bacterial strains were grown in LB broth overnight at 25°C and adjusted to an OD_{600} of 1. A volume of 25 µl of the test bacterial culture was spotted on the two compartments with NGM. The plates were incubated at 25°C for 12 hours. At the time of assay, 1 µl of sodium azide was spotted near the edges of the BA-containing compartment. Gravid adult worms were washed thrice with S-basal buffer, and 50-70 worms were placed at the center of the BA-containing compartment and incubated at 25°C (schematic in Figure 1A). The preference index (PI) was calculated after 3 hours using the following equation:

Preference index (PI) = [(Worms near diet+EAA) – (Worms near diet-EAA)] / Total worms

To understand worms' preference for individual CeMbio bacterium (test bacterium) over *E. coli* OP50 (control bacterium), we performed an 'odor-only' diet preference assay as described above without any supplementations. The test bacterium culture was spotted in one of the NGM-filled compartments, while *E. coli* OP50 was spotted in the other compartment (schematic in Figure S2A). The remaining steps of the assay were performed in a manner similar to the one mentioned above. The PI was calculated using the following equation:

Preference index (PI) = (Worms near CeMbio – Worms near OP50) / Total worms

All diet choice experiments were conducted as standard preference assays unless otherwise specified.

Identification of volatiles produced by microbiota of *C. elegans*

The volatile profiles of CeMbio strains and *E. coli* OP50 were determined using gas chromatography-mass spectrometry (GC-MS/MS) [90]. Briefly, the individual bacterium was

inoculated in 3 ml of LB broth and grown overnight at 25°C. NGM dishes (60 mm) were seeded with seven spots of 50 µl bacterial cultures adjusted to an OD_{600} of 1. Seeded plates were then incubated at 25°C for 23 hours. Two plates of the same conditions were sealed together using parafilm and incubated at 25°C for 1 hour to build odor concentration in the headspace for sampling. The volatiles were collected using a solid-phase microextraction fiber (SPME DVB: Divinylbenzene – Carboxen-WR – polydimethyl siloxane, 80 µm: Agilent Technologies, Part no. 5191-5874). Odors were subjected to thermal desorption and were identified using gas chromatography-mass spectrometry in an 8890C gas chromatography machine coupled with a 7000D GC/TQ, which used a capillary column HP-5MS ultra inert (30 m x 0.25 mm and 0.25 m, Agilent 19091S-433UI: 0245625H) with Helium as the carrier gas at a constant flow rate of 1.5 ml/min. The injection of the sample was done at 40°C. The GC program included a 40°C hold for 1 minute followed by a temperature ramp to 170°C at a rate of 5°C/min, then by a ramp to 270°C at a rate of 100°C/min, and finally, hold for 2 minutes. The inlet, M.S. source, and M.S. quadrupole temperatures were maintained at 225°C, 230°C, and 150°C, respectively. Odors were then identified using NIST (National Institute of Standards and Technology) 2019 V2.3 Mass Spectral Library and Agilent MassHunter Workstation version 10.0. The m/z peaks detected in the odor profile of NGM plates seeded with LB broth were subtracted from the odor profile of each tested bacterium. For leucine supplementation, preferred diets (CEent1, JUb66, and BIGb0170) were seeded on NGM supplemented with 5 mM leucine (+ LEU) and without supplementation (- LEU). GC-MS/MS was performed as described above, and a heat map was generated by analyzing the area under the curve of individual volatile peaks (Figure 2C).

For quantification of IAA, a standard curve was prepared by measuring the area under the curve for five concentrations (0.18 µM, 0.36 µM, 2.25 µM, 4.50 µM, and 8.99 µM) of IAA (Figure S3E). The preferred diets were seeded with two spots of 50 µl culture on NGM dishes supplemented with (+ LEU) and without (- LEU) leucine. The amount of IAA produced by each bacterium was determined by comparing the area under the curve in the GC-MS/MS plot of each bacterium against the standard curve for IAA.

Bacterial RNA extraction and qRT-PCR relative expression analysis

For RNA isolation, bacterial culture was grown in 20 ml of NGM broth with 5 mM leucine (+ LEU) and without leucine (- LEU) for 24 hours at 25°C. Bacterial cells were harvested by centrifugation at 5000 rpm for 20 min. The bacterial pellet was treated with RNAprotect Bacteria Reagent (Qiagen, Cat. No. 76506). Total RNA was extracted from the samples using the RNeasy Mini kit (Qiagen, Cat. No. 74104), and DNase I (NEB Cat. No. M0303S) treatment was done to remove genomic DNA. cDNA was synthesized using the iScript cDNA synthesis kit (Bio-Rad, Cat. No. 170-8891) and used for qRT-PCR to analyze the relative gene expression of *ilvE* using SYBR Green detection mix (Bio-Rad Cat. No. 1725124) on StepOnePlus (Applied Biosystems) machine. *rboB* was used as a housekeeping gene. The comparative ΔΔCt method was used to determine the fold change of the *ilvE* target gene.

Chemotaxis assay

Chemotaxis assays were performed as described previously [90]. Briefly, BA plates (90 mm) air-dried for 90 minutes were used for the assays. Sodium azide (2 µl) was spotted on the two diametrically opposite ends of the plate, followed by 2 µl of the test chemical to one side and solvent to the opposite side (schematic in Figure S4A). Test chemicals were diluted using suitable solvents (Table S1). Gravid adult worms were washed thrice with S-basal buffer. 50 to 80 gravid adult worms were placed in the center of the plate and incubated at 25°C for 3 hours. The chemotaxis index (CI) was calculated using the following equation:

Chemotaxis index (CI) = (Worms towards test – Worms towards control) / Total worms

Odor adaptation

Gravid adult worms were washed thrice with S-basal buffer and transferred to 60 mm NGM plates. For odor adaption, worms were exposed to bacterial odors by sealing an NGM plate seeded with bacteria (7 spots of 50 µL bacterial culture) with the plate containing washed worms. These plates were incubated at 25°C for 90 minutes to desensitize the worms to the odor. After exposure, worms were washed once with S basal buffer and used to perform chemotaxis assays (schematic in Figure 4A). For the modified odor adaptation assays, the NGM plate containing worms was exposed to IAA (1:10 dilution) by sealing it with another NGM agar plate containing 4 spots of 3 µl IAA or

without IAA (schematic in Figure S6A). The adapted worms were then used for diet preference or chemotaxis assays as required.

Worm movement track analysis

We recorded the motion of five worms on a chemotaxis assay plate using a simple imaging setup. We used 1800 frames (1 fps) to track the behavior of worms. Using MATLAB, we extracted the set of binary images from the recorded frames and processed them using the Trackmate plugin of Fiji software to track worms' motion. We used the Advanced Kalman algorithm to create tracks and interpolate for missing frames. The trajectory information was exported in the motilitylab spreadsheet format. This data was analyzed using MATLAB to measure the average speed of worms and the total distance traveled by worms [91-94].

Statistical analyses

All statistical analyses were done using GraphPad Prism version 8. Statistical analyses were performed either by a two-tailed unpaired *t*-test between two groups or one-way ANOVA, followed by post hoc Dunnett's multiple comparison test across multiple groups. The significant differences were denoted according to P-values: $*P \leq 0.1$; $**P \leq 0.01$; $***P \leq 0.001$; and $****P \leq 0.0001$. Data are presented as means ± SEM. All experiments were performed with 15 biological replicates done over three days.

Acknowledgments

Some *C. elegans* strains were provided by CGC, which is funded by the NIH Office of Infrastructure Programs. We thank Ms. Aatira and Mr. Karthick for helping with chemotaxis assays and Ms. Navjot Kaur for some GC-MS analysis. This work was supported by funds from the University of Dundee and from the Royal Society.

Author Contributions

GR, NM, RS, M-AF, CC and VS conceptualized the study. GR, NM, RS and CC performed the experiments. GR, NM, RS, and VS analyzed and interpreted the data. GR, NM, RS, M-AF, and VS wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

1.   Gietzen, D.W., and Rogers, Q.R. (2006). Nutritional homeostasis and indispensable amino acid sensing: a new solution to an old puzzle. Trends in Neurosciences *29*, 91-99. https://doi.org/10.1016/j.tins.2005.12.007.

2.   Tomlinson, C., Rafii, M., Ball, R.O., and Pencharz, P.B. (2011). Arginine can be synthesized from enteral proline in healthy adult humans. J Nutr *141*, 1432-1436. 10.3945/jn.110.137224.

3.   Sancak, Y., Peterson, T.R., Shaul, Y.D., Lindquist, R.A., Thoreen, C.C., Bar-Peled, L., and Sabatini, D.M. (2008). The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. Science *320*, 1496-1501. 10.1126/science.1157535.

4.   Hara, K., Yonezawa, K., Weng, Q.P., Kozlowski, M.T., Belham, C., and Avruch, J. (1998). Amino acid sufficiency and mTOR regulate p70 S6 kinase and eIF-4E BP1 through a common effector mechanism. The Journal of biological chemistry *273*, 14484-14494. 10.1074/jbc.273.23.14484.

5.   van der Vos, K.E., and Coffer, P.J. (2012). Glutamine metabolism links growth factor signaling to the regulation of autophagy. Autophagy *8*, 1862-1864. 10.4161/auto.22152.

6.   Bauchart-Thevret, C., Cui, L., Wu, G., and Burrin, D.G. (2010). Arginine-induced stimulation of protein synthesis and survival in IPEC-J2 cells is mediated by mTOR but not nitric oxide. American journal of physiology. Endocrinology and metabolism *299*, E899-909. 10.1152/ajpendo.00068.2010.

7.   Gauthier-Coles, G., Vennitti, J., Zhang, Z., Comb, W.C., Xing, S., Javed, K., Bröer, A., and Bröer, S. (2021). Quantitative modelling of amino acid transport and homeostasis in mammalian cells. Nat Commun *12*, 5282. 10.1038/s41467-021-25563-x.

8.   Bar-Peled, L., and Sabatini, D.M. (2014). Regulation of mTORC1 by amino acids. Trends in Cell Biology *24*, 400-406. https://doi.org/10.1016/j.tcb.2014.03.003.

9.   González, A., and Hall, M.N. (2017). Nutrient sensing and TOR signaling in yeast and mammals. The EMBO journal *36*, 397-408. 10.15252/embj.201696010.

10.   Austad, S.N., Smith, J.R., and Hoffman, J.M. (2024). Amino acid restriction, aging, and longevity: an update. Front Aging *5*, 1393216. 10.3389/fragi.2024.1393216.

11.   Minussi, I., Gerrits, W.J.J., Jansman, A.J.M., Gerritsen, R., Lambert, W., Zonderland, J.J., and Bolhuis, J.E. (2023). Amino acid supplementation counteracts negative effects of low protein diets on tail biting in pigs more than extra environmental enrichment. Scientific Reports *13*, 19268. 10.1038/s41598-023-45704-0.

12.   Enomoto, H., Sakai, Y., Aizawa, N., Iwata, Y., Tanaka, H., Ikeda, N., Hasegawa, K., Yoh, K., Ishii, A., Takashima, T., et al. (2013). Association of amino acid imbalance with the severity of liver fibrosis and esophageal varices. Ann Hepatol *12*, 471-478.

13.   Tai, E.S., Tan, M.L., Stevens, R.D., Low, Y.L., Muehlbauer, M.J., Goh, D.L., Ilkayeva, O.R., Wenner, B.R., Bain, J.R., Lee, J.J., et al. (2010). Insulin resistance is associated with a metabolic profile of altered protein metabolism in Chinese and Asian-Indian men. Diabetologia *53*, 757-767. 10.1007/s00125-009-1637-8.

14.   Wang, T.J., Larson, M.G., Vasan, R.S., Cheng, S., Rhee, E.P., McCabe, E., Lewis, G.D., Fox, C.S., Jacques, P.F., Fernandez, C., et al. (2011). Metabolite profiles and the risk of developing diabetes. Nature medicine *17*, 448-453. 10.1038/nm.2307.

15.   De Marte, M.L., and Enesco, H.E. (1986). Influence of low tryptophan diet on survival and organ growth in mice. Mechanisms of ageing and development *36*, 161-171. 10.1016/0047-6374(86)90017-5.

16.   Miller, R.A., Buehner, G., Chang, Y., Harper, J.M., Sigler, R., and Smith-Wheelock, M. (2005). Methionine-deficient diet extends mouse lifespan, slows immune and lens aging, alters glucose, T4, IGF-I and insulin levels, and increases hepatocyte MIF levels and stress resistance. *4*, 119-125. https://doi.org/10.1111/j.1474-9726.2005.00152.x.

17.   Lee, B.C., Kaya, A., Ma, S., Kim, G., Gerashchenko, M.V., Yim, S.H., Hu, Z., Harshman, L.G., and Gladyshev, V.N. (2014). Methionine restriction extends lifespan of Drosophila melanogaster under conditions of low amino-acid status. Nature Communications *5*, 3592. 10.1038/ncomms4592.

18.   Cota, D., Proulx, K., Smith, K.A.B., Kozma, S.C., Thomas, G., Woods, S.C., and Seeley, R.J. (2006). Hypothalamic mTOR Signaling Regulates Food Intake. *312*, 927-930. doi:10.1126/science.1124147.

19.   Wang, M., Shen, Y., Tan, Z., Yasen, A., Fan, B., and Shen, X. (2023). Metabolomics analysis of dietary restriction results in a longer lifespan due to alters of amino acid levels in larval hemolymph of Bombyx mori. Scientific Reports *13*, 6828. 10.1038/s41598-023-34132-9.

20.   Makarios-Lahham, L., Roseau, S.M., Fromentin, G., Tome, D., and Even, P.C. (2004). Rats free to select between pure protein and a fat-carbohydrate mix ingest high-protein mixed meals during the dark period and protein meals during the light period. J Nutr *134*, 618-624. 10.1093/jn/134.3.618.

21. Delaney, K., and Gelperin, A. (1986). Post-ingestive food-aversion learning to amino acid deficient diets by the terrestrial slugLimax maximus. Journal of Comparative Physiology A *159*, 281-295. 10.1007/BF00603975.

22. Gietzen, D.W. (1993). Neural Mechanisms in the Responses to Amino Acid Deficiency1. The Journal of Nutrition *123*, 610-625. https://doi.org/10.1093/jn/123.4.610.

23. Murphy, M.E., and Pearcy, S.D. (1993). Dietary amino acid complementation as a foraging strategy for wild birds. Physiology & Behavior *53*, 689-698. https://doi.org/10.1016/0031-9384(93)90175-F.

24. Juricic, P., Grönke, S., and Partridge, L. (2019). Branched-Chain Amino Acids Have Equivalent Effects to Other Essential Amino Acids on Lifespan and Aging-Related Traits in Drosophila. The Journals of Gerontology: Series A *75*, 24-31. 10.1093/gerona/glz080 %J The Journals of Gerontology: Series A.

25. Rogers, Q.R., Wigle, A.R., Laufer, A., Castellanos, V.H., and Morris, J.G. (2004). Cats Select for Adequate Methionine but Not Threonine. The Journal of Nutrition *134*, 2046S-2049S. https://doi.org/10.1093/jn/134.8.2046S.

26. Shallenberger, R.S. (1993). Taste Chemistry Principles. In Taste Chemistry, R.S. Shallenberger, ed. (Springer US), pp. 47-109. 10.1007/978-1-4615-2666-7_3.

27. Schiffman, S.S., Sennewald, K., and Gagnon, J. (1981). Comparison of taste qualities and thresholds of D- and L-amino acids. Physiology and Behavior *27*, 51-59. 10.1016/0031-9384(81)90298-5.

28. Zhao, C.J., Schieber, A., and Gänzle, M.G. (2016). Formation of taste-active amino acids, amino acid derivatives and peptides in food fermentations – A review. Food Research International *89*, 39-47. https://doi.org/10.1016/j.foodres.2016.08.042.

29. Bachmanov, A.A., Bosak, N.P., Glendinning, J.I., Inoue, M., Li, X., Manita, S., McCaughey, S.A., Murata, Y., Reed, D.R., Tordoff, M.G., and Beauchamp, G.K. (2016). Genetics of Amino Acid Taste and Appetite. Advances in Nutrition 7, 806S-822S. https://doi.org/10.3945/an.115.011270.

30. Yoshida, M., and Saito, S. (1969). MULTIDIMENSIONAL SCALING OF THE TASTE OF AMINO ACIDS. Japanese Psychological Research *11*, 149-166. 10.4992/psycholres1954.11.149.

31. Wellendorph, P., Hansen, K.B., Balsgaard, A., Greenwood, J.R., Egebjerg, J., and Bräuner-Osborne, H. (2005). Deorphanization of GPRC6A: A promiscuous L-α-amino acid receptor with preference for basic amino acids. Molecular Pharmacology *67*, 589-597. 10.1124/mol.104.007559.

32. Kuang, D., Yao, Y., Wang, M., Pattabiraman, N., Kotra, L.P., and Hampson, D.R. (2003). Molecular Similarities in the Ligand Binding Pockets of an Odorant Receptor and the Metabotropic Glutamate Receptors. Journal of Biological Chemistry *278*, 42551-42559. 10.1074/jbc.M307120200.

33. Nelson, G., Chandrashekar, J., Hoon, M.A., Feng, L., Zhao, G., Ryba, N.J.P., and Zuker, C.S. (2002). An amino-acid taste receptor. Nature *416*, 199-202. 10.1038/nature726.

34. Conigrave, A.D., Quinn, S.J., and Brown, E.M. (2000). L-amino acid sensing by the extracellular Ca2+-sensing receptor. Proceedings of the National Academy of Sciences of the United States of America *97*, 4814-4819. 10.1073/pnas.97.9.4814.

35. Conigrave, A.D., and Hampson, D.R. (2006). Broad-spectrum l-amino acid sensing by class 3 G-protein-coupled receptors. Trends in Endocrinology & Metabolism *17*, 398-407. https://doi.org/10.1016/j.tem.2006.10.012.

36. Nakanishi, S. (1992). Molecular Diversity of Glutamate Receptors and Implications for Brain Function. *258*, 597-603. doi:10.1126/science.1329206.

37. Miguel-Aliaga, I. (2012). Nerveless and gutsy: intestinal nutrient sensing from invertebrates to humans. Seminars in cell & developmental biology *23*, 614-620. 10.1016/j.semcdb.2012.01.002.

38. Dillon, J., Franks, C.J., Murray, C., Edwards, R.J., Calahorro, F., Ishihara, T., Katsura, I., Holden-Dye, L., and O'Connor, V. (2015). Metabotropic Glutamate Receptors: MODULATORS OF CONTEXT-DEPENDENT FEEDING BEHAVIOUR IN C. ELEGANS. The Journal of biological chemistry *290*, 15052-15065. 10.1074/jbc.M114.606608.

39. Colombani, J., Raisin, S., Pantalacci, S., Radimerski, T., Montagne, J., and Léopold, P. (2003). A Nutrient Sensor Mechanism Controls Drosophila Growth. Cell *114*, 739-749. https://doi.org/10.1016/S0092-8674(03)00713-X.

40. Speca, D.J., Lin, D.M., Sorensen, P.W., Isacoff, E.Y., Ngai, J., and Dittman, A.H. (1999). Functional Identification of a Goldfish Odorant Receptor. Neuron *23*, 487-498. https://doi.org/10.1016/S0896-6273(00)80802-8.

41. Wellendorph, P., Hansen, K.B., Balsgaard, A., Greenwood, J.R., Egebjerg, J., and Bräuner-Osborne, H. (2005). Deorphanization of GPRC6A: a promiscuous L-alpha-amino acid receptor with preference for basic amino acids. Mol Pharmacol *67*, 589-597. 10.1124/mol.104.007559.

42. Sawin, E.R., Ranganathan, R., and Horvitz, H.R. (2000). C. elegans Locomotory Rate Is Modulated by the Environment through a Dopaminergic Pathway and by Experience through a Serotonergic Pathway. Neuron *26*, 619-631. https://doi.org/10.1016/S0896-6273(00)81199-X.

43. Rhoades, J.L., Nelson, J.C., Nwabudike, I., Yu, S.K., McLachlan, I.G., Madan, G.K., Abebe, E., Powers, J.R., Colón-Ramos, D.A., and Flavell, S.W. (2019). ASICs Mediate Food Responses in an Enteric Serotonergic Neuron that Controls Foraging Behaviors. Cell *176*, 85-97.e14. 10.1016/j.cell.2018.11.023.

44. Hammarlund, M., Hobert, O., Miller, D.M., 3rd, and Sestan, N. (2018). The CeNGEN Project: The Complete Gene Expression Map of an Entire Nervous System. Neuron *99*, 430-433. 10.1016/j.neuron.2018.07.042.

45. Edwards, C., Canfield, J., Copes, N., Brito, A., Rehan, M., Lipps, D., Brunquell, J., Westerheide, S.D., and Bradshaw, P.C. (2015). Mechanisms of amino acid-mediated lifespan extension in Caenorhabditis elegans. BMC genetics *16*, 8. 10.1186/s12863-015-0167-2.

46. Weisskopf, L., Schulz, S., and Garbeva, P. (2021). Microbial volatile organic compounds in intra-kingdom and inter-kingdom interactions. Nature Reviews Microbiology *19*, 391-404. 10.1038/s41579-020-00508-1.

47. Bargmann, C.I., Hartwieg, E., and Horvitz, H.R. (1993). Odorant-selective genes and neurons mediate olfaction in C. elegans. Cell *74*, 515-527. 10.1016/0092-8674(93)80053-h.

48. Dirksen, P., Assie, A., Zimmermann, J., Zhang, F., Tietje, A.M., Marsh, S.A., Felix, M.A., Shapira, M., Kaleta, C., Schulenburg, H., and Samuel, B.S. (2020). CeMbio - The Caenorhabditis elegans Microbiome Resource. G3 (Bethesda) *10*, 3025-3039. 10.1534/g3.120.401309.

49. Bargmann, C.I. (2006). Chemosensation in C. elegans. WormBook : the online review of C. elegans biology, 1-29. 10.1895/wormbook.1.123.1.

50. Troemel, E.R., Kimmel, B.E., and Bargmann, C.I. (1997). Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in C. elegans. Cell *91*, 161-169. 10.1016/s0092-8674(00)80399-2.

51. Beverly, M., Anbil, S., and Sengupta, P. (2011). Degeneracy and neuromodulation among thermosensory neurons contribute to robust thermosensory behaviors in Caenorhabditis elegans. The Journal of neuroscience : the official journal of the Society for Neuroscience *31*, 11718-11727. 10.1523/jneurosci.1098-11.2011.

52. Sengupta, P., Colbert, H.A., and Bargmann, C.I. (1994). The C. elegans gene odr-7 encodes an olfactory-specific member of the nuclear receptor superfamily. Cell *79*, 971-980. 10.1016/0092-8674(94)90028-0.

53. Colbert, H.A., and Bargmann, C.I. (1995). Odorant-specific adaptation pathways generate olfactory plasticity in C. elegans. Neuron *14*, 803-812. 10.1016/0896-6273(95)90224-4.

54. Andersen, E.C., Gerke, J.P., Shapiro, J.A., Crissman, J.R., Ghosh, R., Bloom, J.S., Félix, M.A., and Kruglyak, L. (2012). Chromosome-scale selective sweeps shape Caenorhabditis elegans genomic diversity. Nature genetics *44*, 285-290. 10.1038/ng.1050.

55. Cook, D.E., Zdraljevic, S., Roberts, J.P., and Andersen, E.C. (2017). CeNDR, the Caenorhabditis elegans natural diversity resource. Nucleic Acids Res *45*, D650-d657. 10.1093/nar/gkw893.

56. Thomas, J.H., and Robertson, H.M. (2008). The Caenorhabditis chemoreceptor gene families. BMC Biol *6*, 42. 10.1186/1741-7007-6-42.

57. Zaslaver, A., Liani, I., Shtangel, O., Ginzburg, S., Yee, L., and Sternberg, P.W. (2015). Hierarchical sparse coding in the sensory system of *Caenorhabditis elegans*. *112*, 1185-1189. doi:10.1073/pnas.1423656112.

58. Lin, A., Qin, S., Casademunt, H., Wu, M., Hung, W., Cain, G., Tan, N.Z., Valenzuela, R., Lesanpezeshki, L., Venkatachalam, V., et al. (2023). Functional imaging and quantification of multineuronal olfactory responses in *C. elegans*. *9*, eade1249. doi:10.1126/sciadv.ade1249.

59. Pu, L., Wang, J., Lu, Q., Nilsson, L., Philbrook, A., Pandey, A., Zhao, L., Schendel, R.V., Koh, A., Peres, T.V., et al. (2023). Dissecting the genetic landscape of GPCR signaling through phenotypic profiling in C. elegans. Nat Commun *14*, 8410. 10.1038/s41467-023-44177-z.

60. Alm, J., Ohnmeiss, T.E., Lanza, J., and Vriesenga, L. (1990). Preference of cabbage white butterflies and honey bees for nectar that contains amino acids. Oecologia *84*, 53-57. 10.1007/bf00665594.

61. Bertazzini, M., Medrzycki, P., Bortolotti, L., Maistrello, L., and Forlani, G. (2010). Amino acid content and nectar choice by forager honeybees (Apis mellifera L.). Amino Acids *39*, 315-318. 10.1007/s00726-010-0474-x.

62. Haverkamp, A., Yon, F., Keesey, I.W., Mißbach, C., Koenig, C., Hansson, B.S., Baldwin, I.T., Knaden, M., and Kessler, D. (2016). Hawkmoths evaluate scenting flowers with the tip of their proboscis. eLife *5*, e15039. 10.7554/eLife.15039.

63. Yon, F., Joo, Y., Cortés Llorca, L., Rothe, E., Baldwin, I.T., and Kim, S.-G. (2016). Silencing Nicotiana attenuata LHY and ZTL alters circadian rhythms in flowers. *209*, 1058-1066. https://doi.org/10.1111/nph.13681.

64. Bhattacharya, S., and Baldwin, I.T. (2012). The post-pollination ethylene burst and the continuation of floral advertisement are harbingers of non-random mate selection in Nicotiana attenuata. *71*, 587-601. https://doi.org/10.1111/j.1365-313X.2012.05011.x.

65. Kessler, D., and Baldwin, I.T. (2007). Making sense of nectar scents: the effects of nectar secondary metabolites on floral visitors of Nicotiana attenuata. *49*, 840-854. https://doi.org/10.1111/j.1365-313X.2006.02995.x.

66. Hazelwood, L.A., Daran, J.-M., Maris, A.J.A.v., Pronk, J.T., and Dickinson, J.R. (2008). The Ehrlich Pathway for Fusel Alcohol Production: a Century of Research on *Saccharomyces cerevisiae* Metabolism. Applied and Environmental Microbiology *74*, 2259-2266. doi:10.1128/AEM.02625-07.

67. Dou, T., Hu, C., Zhao, S., Gao, H., He, W., Deng, G., Sheng, O., Bi, F., Yang, Q., Li, C., et al. (2022). RNA-Seq Reveals the Effect of Ethylene on the Volatile Organic Components (VOCs) of Cavendish Banana at Different Post-harvesting

Stages. Journal of Plant Growth Regulation *41*, 3061-3074. 10.1007/s00344-021-10496-y.

68. Becher, P.G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S., Larsson, M.C., Hansson, B.S., Piškur, J., Witzgall, P., and Bengtsson, M. (2012). Yeast, not fruit volatiles mediate Drosophila melanogaster attraction, oviposition and development. Functional Ecology *26*, 822-828. https://doi.org/10.1111/j.1365-2435.2012.02006.x.

69. Dotterl, S., and Gershenzon, J. (2023). Chemistry, biosynthesis and biology of floral volatiles: roles in pollination and other functions. Nat Prod Rep. 10.1039/d3np00024a.

70. He, J., Tuo, W., Zhang, X., Dai, Y., Fang, M., Zhou, T., Xiu, M., and Liu, Y. (2022). Olfactory Senses Modulate Food Consumption and Physiology in Drosophila melanogaster. Front Behav Neurosci *16*, 788633. 10.3389/fnbeh.2022.788633.

71. Liu, J., Sun, L., Fu, D., Zhu, J., Liu, M., Xiao, F., and Xiao, R. (2022). Herbivore-Induced Rice Volatiles Attract and Affect the Predation Ability of the Wolf Spiders, Pirata subpiraticus and Pardosa pseudoannulata. Insects *13*. 10.3390/insects13010090.

72. Zjacic, N., and Scholz, M. (2022). The role of food odor in invertebrate foraging. Genes Brain Behav *21*, e12793. 10.1111/gbb.12793.

73. Haupt, S.S., Sakurai, T., Namiki, S., Kazawa, T., and Kanzaki, R. (2010). Olfactory Information Processing in Moths. In The Neurobiology of Olfaction, A. Menini, ed.

74. Pham-Delegue, M.H., Bailez, O., Blight, M.M., Masson, C., Picard-Nizou, A.L., and Wadhams, L.J. (1993). Behavioural discrimination of oilseed rape volatiles by the honeybee Apis mellifera L. Chemical Senses *18*, 483-494. 10.1093/chemse/18.5.483.

75. Mansfeld, J., Urban, N., Priebe, S., Groth, M., Frahm, C., Hartmann, N., Gebauer, J., Ravichandran, M., Dommaschk, A., Schmeisser, S., et al. (2015). Branched-chain amino acid catabolism is a conserved regulator of physiological ageing. Nat Commun *6*, 10043. 10.1038/ncomms10043.

76. Worthy, S.E., Haynes, L., Chambers, M., Bethune, D., Kan, E., Chung, K., Ota, R., Taylor, C.J., and Glater, E.E. (2018). Identification of attractive odorants released by preferred bacterial food found in the natural habitats of C. elegans. PloS one *13*, e0201158. 10.1371/journal.pone.0201158.

77. Worthy, S.E., Rojas, G.L., Taylor, C.J., and Glater, E.E. (2018). Identification of Odor Blend Used by Caenorhabditis elegans for Pathogen Recognition. Chem Senses *43*, 169-180. 10.1093/chemse/bjy001.

78. Bizzio, L.N., Tieman, D., and Munoz, P.R. (2021). Branched-Chain Volatiles in Fruit: A Molecular Perspective. Front Plant Sci *12*, 814138. 10.3389/fpls.2021.814138.

79.   Maoz, I., Lewinsohn, E., and Gonda, I. (2022). Amino acids metabolism as a source for aroma volatiles biosynthesis. Curr Opin Plant Biol *67*, 102221. 10.1016/j.pbi.2022.102221.

80.   Yan, Y., Sun, L., Xing, X., Wu, H., Lu, X., Zhang, W., Xu, J., and Ren, Q. (2022). Microbial succession and exploration of higher alcohols-producing core bacteria in northern Huangjiu fermentation. AMB Express *12*, 79. 10.1186/s13568-022-01418-6.

81.   Taylor-Burds, C.C., Westburg, Ä.M., Wifall, T.C., and Delay, E.R. (2004). Behavioral Comparisons of the Tastes of l-Alanine and Monosodium Glutamate in Rats. Chem Senses *29*, 807-814. 10.1093/chemse/bjh246 %J Chemical Senses.

82.   Ward, S. (1973). Chemotaxis by the nematode Caenorhabditis elegans: identification of attractants and analysis of the response by use of mutants. Proceedings of the National Academy of Sciences of the United States of America *70*, 817-821. 10.1073/pnas.70.3.817.

83.   Xu, H., Gao, Y., Hassan, A., Liu, Y., Zhao, X., and Huang, Q. (2024). Neuroregulation of foraging behavior mediated by the olfactory co-receptor Orco in termites. International Journal of Biological Macromolecules *262*, 129639. https://doi.org/10.1016/j.ijbiomac.2024.129639.

84.   Libert, S., Zwiener, J., Chu, X., VanVoorhies, W., Roman, G., and Pletcher, S.D. (2007). Regulation of <i>Drosophila</i> Life Span by Olfaction and Food-Derived Odors. *315*, 1133-1137. doi:10.1126/science.1136610.

85.   Choi, J.I., Yoon, K.H., Subbammal Kalichamy, S., Yoon, S.S., and Il Lee, J. (2016). A natural odor attraction between lactic acid bacteria and the nematode Caenorhabditis elegans. The ISME journal *10*, 558-567. 10.1038/ismej.2015.134.

86.   Zhang, C., Zhao, N., Chen, Y., Zhang, D., Yan, J., Zou, W., Zhang, K., and Huang, X. (2016). The Signaling Pathway of Caenorhabditis elegans Mediates Chemotaxis Response to the Attractant 2-Heptanone in a Trojan Horse-like Pathogenesis. The Journal of biological chemistry *291*, 23618-23627. 10.1074/jbc.M116.741132.

87.   Dixit, A., Sandhu, A., Modi, S., Shashikanth, M., Koushika, S.P., Watts, J.L., and Singh, V. (2020). Neuronal control of lipid metabolism by STR-2 G protein-coupled receptor promotes longevity in Caenorhabditis elegans. *19*, e13160. https://doi.org/10.1111/acel.13160.

88.   Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94. 10.1093/genetics/77.1.71.

89.   Pu, L., Nilsson, L., Chen, C., and Wang, J. (2023). Iterative editing of multiple genes using CRISPR/Cas9 in C. elegans. microPublication biology *2023*. 10.17912/micropub.biology.000898.

90. Prakash, D., Ms, A., Radhika, B., Venkatesan, R., Chalasani, S.H., and Singh, V. (2021). 1-Undecene from Pseudomonas aeruginosa is an olfactory signal for flight-or-fight response in Caenorhabditis elegans. The EMBO journal *40*, e106938. 10.15252/embj.2020106938.

91. Saxton, M.J., and Jacobson, K. (1997). SINGLE-PARTICLE TRACKING:Applications to Membrane Dynamics. *26*, 373-399. https://doi.org/10.1146/annurev.biophys.26.1.373.

92. Codling, E.A., Plank, M.J., and Benhamou, S. (2008). Random walk models in biology. Journal of the Royal Society, Interface *5*, 813-834. 10.1098/rsif.2008.0014.

93. Li, L., Nørrelykke, S.F., and Cox, E.C. (2008). Persistent cell motion in the absence of external signals: a search strategy for eukaryotic cells. PloS one *3*, e2093. 10.1371/journal.pone.0002093.

94. Berg, H.C. (1993). Random Walks in Biology

New and Expanded Edition, REV - Revised Edition (Princeton University Press). 10.2307/j.ctv7r40w6.