APP β-CTF triggers cell-autonomous synaptic toxicity independent of Aβ

Mengxun Luo[1,2], Jia Zhou[1,2], Cailu Sun[1,2], Wanjia Chen[1,2], Chaoying Fu[1], Chenfang Si[1,2], Yaoyang Zhang[1], Yang Geng[1], Yelin Chen[1]*

Affiliations:

[1]Interdisciplinary Research Center on Biology and Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 100 Haike Rd., Pudong, Shanghai, 201210, China

[2]University of Chinese Academy of Sciences, Beijing, 100049, China.

*Correspondence: Yelin Chen (chenyelin@sioc.ac.cn)

Abstract

Aβ is believed to play a significant role in synaptic degeneration observed in Alzheimer's disease (AD) and is primarily investigated as a secreted peptide. However, the contribution of intracellular Aβ or other cleavage products of its precursor protein (APP) to synaptic loss remains uncertain. In this study, we conducted a systematic examination of their cell-autonomous impact using a sparse expression system. Here, these proteins/peptides were overexpressed in a single neuron, surrounded by thousands of untransfected neurons. Surprisingly, we found that APP induced dendritic spine loss only when co-expressed with BACE1. This effect was mediated by β-CTF, a β-cleavage product of APP, through an endosome-related pathway independent of Aβ. Neuronal expression of β-CTF in mouse brains resulted in defective synaptic transmission and cognitive impairments, even in the absence of amyloid plaques. These findings unveil a β-CTF-initiated mechanism driving synaptic toxicity irrespective of amyloid plaque formation and suggest a potential intervention by inhibiting the endosomal GTPase Rab5.

Keywords: Alzheimer's disease, Amyloid precursor protein, synapse, β-CTF, endosome

Introduction

Alzheimer's disease (AD) stands as the most prevalent form of neurodegenerative disease and a leading cause of dementia. It is characterized by extracellular amyloid plaque deposition, intracellular neurofibrillary tangles, synaptic and neuronal loss, and neuroinflammation (Blennow et al., 2006; Gómez-Isla et al., 1996; JohnHardy and Selkoe, 2002; Leyns and Holtzman, 2017; Selkoe, 2002). The amyloid plaque primarily consists of aggregated Aβ, a cleavage product of amyloid precursor protein (APP) (Glenner and Wong, 1984; Masters et al., 1985). Notably, two Aβ antibodies have demonstrated efficacy in removing amyloid plaques and slowing AD progression in phase III clinical trials (Sims et al., 2023; van Dyck et al., 2023). Furthermore, numerous naturally occurring mutations in genes encoding APP or its catalytic enzyme γ-secretase result in early onset familial AD or reduce AD risk (Jonsson et al., 2012; Liu et al., 2017; Mullan et al., 1992; Nilsberth et al., 2001). This collective evidence, spanning AD pathology, human genetics, and intervention trials, strongly supports a causal role of Aβ and amyloid plaque in AD pathogenesis. However, despite clinical trials employing Aβ antibodies targeting Aβ oligomers, protofibrils, or deposited plaque, AD progression has been slowed down by only ~30% (Mintun et al., 2021; Pleen and Townley, 2022; Sims et al., 2023; van Dyck et al., 2023). Notably, antibodies against monomeric soluble Aβ failed to yield clinical benefits (Sperling et al., 2023). It is possible that these Aβ antibodies may overlook certain pathogenic factors crucial for AD pathogenesis.

APP, a type I transmembrane protein, undergoes cleavage primarily by α-secretase on the cytoplasmic membrane, producing soluble α-cleavage N-terminal fragment (sAPPα) and α-cleavage C-terminal fragment (α-CTF). Some APP molecules bypass α-cleavage and undergo endocytosis into endocytic compartments, where they are subsequently cleaved by β-secretase, generating soluble β-cleavage N-terminal fragment (sAPPβ) and β-cleavage C-terminal fragment (β-CTF). β-CTF is further cleaved by γ-secretase to produce Aβ and APP intracellular domain (AICD) (Golde et

al., 1992; Zhang and Song, 2013; Zhang et al., 2011). While extracellular Aβ, targeted by Aβ antibodies, is widely studied, the potential contribution of intracellular Aβ, APP, and other APP cleavage products to AD pathogenesis remains uncertain (Konietzko, 2012; Kwart et al., 2019; Nikolaev et al., 2009; Oddo et al., 2003; Vohra et al., 2010; Willem et al., 2015). For instance, β-CTF has been implicated in endosomal dysfunction(Israel et al., 2012; Jiang et al., 2010; Kim et al., 2016; Kwart et al., 2019; Xu et al., 2016), yet its downstream functional impacts remain unclear.

Synapse loss represents an early feature of AD neurodegeneration and is closely associated with cognitive dysfunction (de Wilde et al., 2016; DeKosky et al., 1996; Terry et al., 1991). Secreted Aβ induces synaptic dysfunction by interacting with its receptors on the neuronal plasma membrane (Kamenetz et al., 2003; Kessels et al., 2013; Wei et al., 2010). Other studies have reported γ-secretase inhibition reduced spine density *in vivo* via an APP-dependent pathway (Bittner et al., 2009). Additionally, deletion of APP in mice has been shown to decrease dendritic spine density (Tyan et al., 2012). The diverse outcomes of APP on synapse suggest a complex impact of APP and metabolites. It remains unclear whether APP or other APP fragments can also induce synaptic toxicity in a cell-autonomous manner.

To address these inquiries, we employed a sparse transfection system utilizing the Helios gene gun to explore the potential role of intracellular Aβ or other APP fragments. These molecules were expressed in a single neuron surrounded by untransfected wild type neurons. Surprisingly, full-length APP did not induce synaptic toxicity. However, co-expression of APP with BACE1 resulted in significant loss of dendritic spines, indicating a crucial role of APP β-cleavage in synaptic damage. Further investigations unveiled that this detrimental effect was mediated by β-CTF in a cell-autonomous manner, independent of Aβ. Additionally, *in vivo* expression of β-CTF was adequate to induce synaptic dysfunction and cognitive impairments in mice, even in the absence of amyloid plaques. In summary, our study delineates a

82　mechanism initiated by β-CTF that can induce synaptic degeneration in a

83　cell-autonomous manner, thus extending beyond the scope of Aβ antibody-based

84　therapies.

Results

APP only led to spine loss when co-expressed with BACE1

To investigate the cell-autonomous impact of APP on neurons, we utilized a Helios gene gun transfection system to sparsely express APP in rat organotypic hippocampal slice cultures (Chen et al., 2014). In hippocampal slice cultures at DIV9, transient expression of a familial AD APP mutation (APP$_{Swedish}$) (Mullan et al., 1992) with GFP in CA1 pyramidal neurons for six days did not reduce their dendritic spine densities compared with neurons expressing GFP alone (Figure 1A-B), suggesting that APP alone does not significantly contribute to synaptic loss.

BACE1-mediated β-cleavage of APP is crucial for amyloidogenesis, and BACE1 activity is notably elevated in AD patient brains (Cheng et al., 2014; Vassar et al., 1999). In human brains, APP and BACE1 are expressed at a ratio of about 15:1 (Uhlén et al., 2015). Endogenous BACE1 levels may not suffice to cleave the overexpressed APP. To assess whether insufficient β-cleavage of APP underlies the lack of synaptic toxicity caused by overexpressed APP, we co-expressed APP with BACE1 (in a 15:1 ratio) in organotypic hippocampal slice cultures and observed a significant ~40% reduction in spine density compared to neurons expressing GFP alone (Figure 1C-D). Transient expression of BACE1 alone did not affect spine density (Figure 1E-F). Co-expression of APP and BACE1 with different ratios or using an internal ribosome entry site (IRES) also resulted in significant spine loss (Supplementary figure 1A-B). These findings support the requirement of BACE1 for APP to induce synaptic loss.

When expressed alone in HEK293T cells, APP is predominantly cleaved at its α site to produce sAPPα and α-CTF (C-terminal fragment after APP α-cleavage) (Figure 1G, Supplementary figure 1C). Co-expression with BACE1 led to increased cleavage of APP at its β and β' sites, resulting in elevated β-CTF (more than 4-fold) and β'-CTF

as the major metabolic products and a substantial reduction in α-CTF levels (Figure 1G-H, Supplementary figure 1C).

APP$_{MV}$ (M596V) cannot be cleaved by BACE1 to produce β-CTF and Aβ but has no impact on β'-cleavage (Figure 1I-K) (Citron et al., 1995). When co-expressed with BACE1, APP$_{MV}$ failed to induce spine loss (Figure 1L-M), supporting the requirement of β-cleavage of APP to induce spine loss.

In summary, these findings suggest that certain β-cleavage products of APP, rather than APP or BACE1 alone, could lead to spine loss in a cell-autonomous manner.

β-CTF induced spine loss independent of Aβ.

APP undergoes cleavage by secretases, generating soluble N-terminal fragments (sAPPs) and C-terminal fragment (CTFs). We then explored the impact of different sAPPs and APP-CTFs on dendritic spines. Transient expression of β-CTF significantly reduced the spine density of CA1 pyramidal neurons in organotypic rat hippocampal cultures, whereas expression of sAPPα, sAPPβ, α-CTF or β'-CTF did not produce such an effect (Figure 2A-B, Supplementary figure 2A-B). APP CTFs exhibited similar expression levels in HEK293T cells (Figure 2C).

β-CTF undergoes processing by γ-secretase to generate Aβ and AICD (APP intracellular domain) (Thinakaran and Koo, 2008). Subsequently, we investigated whether Aβ generated from β-CTF was responsible for the β-CTF-induced spine loss. Treatment with a γ-secretase inhibitor, PF03084014 (1μM), effectively reduced Aβ to baseline levels in cells expressing β-CTF without altering the expression levels of β-CTF itself (Figure 2D-E). PF03084014 treatment did not affect the spine density in neurons expressing GFP alone (Supplementary figure 2C-D). Notably, PF03084014 treatment failed to reverse the spine loss induced by β-CTF expression (Figure 2F-G), or by co-expression of APP and BACE1 (Figure 2H-K), suggesting that Aβ might not be the causative factor.

Formatted: Space After: 1 line

To delve deeper into the impact of Aβ on dendritic spines, we engineered two APP

mutants (APP$_{\Delta 59}$ and APP$_{\Delta 57}$) lacking the AICD domain, which are known to generate

significant amounts of Aβ40 and Aβ42, respectively (Figure 2L-M). Co-expression of

either APP$_{\Delta 59}$ or APP$_{\Delta 57}$ with BACE1, did not alter spine density (Figure 2N-O),

further bolstering the idea that Aβ is not responsible for the β-CTF-induced spine loss.

Subsequently, we investigated the involvement of another β-CTF cleavage product,

AICD, which has been reported to interact with the transcription factor forkhead box

O (FoxO) and promote FoxO-induced transcription of proapoptotic genes, leading to

cell death (Wang et al., 2014). However, transient expression of AICD failed to alter

the density of spines (Supplementary figure 2E-F).

In conclusion, these results support the idea that APP can induce spine loss in a

cell-autonomous manner through β-CTF, independent of Aβ and AICD.

Expression of β-CTF damaged synapses in mice.

To ascertain whether β-CTF-induced spine loss could manifest *in vivo*, we examined

the density of dendritic spines from CA1 pyramidal neurons infected with lentivirus

encoding β-CTF and GFP, or GFP alone, in adult mice. Sparse expression of GFP and

β-CTF was observed in CA1 pyramidal neurons (Figure 3A). The spine density of

CA1 pyramidal neurons expressing β-CTF and GFP was significantly lower than that

of neurons expressing GFP alone (Figure 3B-C), indicating that β-CTF could induce

spine loss in a cell-autonomous manner *in vivo*.

Next, we evaluated excitatory synaptic transmission through whole-cell patch-clamp

recording of hippocampal pyramidal neurons infected with lentivirus encoding β-CTF

or GFP. Neurons expressing β-CTF exhibited a ~65% lower frequency of mEPSCs

compared to neighboring uninfected neurons, whereas there was no significant change

in mEPSC frequency in neurons expressing GFP alone (Figure 3D-E). However,

mEPSC amplitude remained unaltered in neurons expressing either β-CTF or GFP

Formatted: Space After: 1 line

(Figure 3D, F). These findings collectively suggest that β-CTF leads to reduced excitatory synapse density without affecting the strength of the remaining synapses.

Expression of β-CTF damaged cognitive function in mice in the absence of plaque formation.

We proceeded to investigate whether β-CTF affects cognitive functions. Adeno-associated viruses (AAV) encoding GFP or β-CTF were bilaterally injected into the hippocampus of 1-month-old mice (Figure 4A-B). After 4 months of expression, animal behaviors were examined. Immunostaining revealed widespread expression of GFP or β-CTF throughout the entire hippocampus (Figure 4B), with no detectable amyloid plaque formation observed using ThS staining (Figure 4C). Side-by-side staining demonstrated robust amyloid plaque deposition in the brain of two-month-old 5XFAD mice (Figure 4D), validating the efficacy of the staining method. The body weight of mice expressing β-CTF in the hippocampus was approximately 17% lower than that of GFP controls (Figure 4E). In the Y-maze test, the mean alternations were significantly reduced in mice expressing β-CTF compared to GFP controls, suggesting abnormal working memory in mice expressing β-CTF (Figure 4F).

The fear conditioning test assesses associative fear learning and memory (Figure 4G) (Xiao et al., 2018). In this test, mice expressing β-CTF exhibited similar baseline levels of freezing time as GFP controls (Figure 4H). However, in the hippocampus-dependent contextual fear conditioning test, mice expressing β-CTF showed a significantly shorter freezing time (approximately 40% less) than those expressing GFP after training (Figure 4H). In the amygdala-dependent cued fear conditioning test, these two groups performed similarly (Figure 4H). In the water T maze test (Figure 4I), mice expressing β-CTF displayed slower learning curves than GFP controls, indicating impairments in acquisition learning (Figure 4J) but not in reversal learning (Figure 4K).

In the open field test, mice expressing β-CTF traveled a longer distance
(Supplementary figure 3A-C) and spent more time in the center area with a higher
frequency of center entrances (Supplementary figure 3D-E) compared to mice
expressing GFP. However, there were no differences in rearing frequency and
duration between the two groups (Supplementary figure 3F-G). These results suggest
that mice expressing β-CTF exhibited increased locomotion and reduced anxiety-like
behaviors. Consistently, mice injected with AAV encoding β-CTF spent significantly
more time exploring and traveled a greater distance in the open arms compared to the
GFP controls in the EPM (Supplementary figure 3H-J). In tail suspension tests, these
two groups exhibited similar levels of immobility (Supplementary figure 3K),
indicating that β-CTF expression in the hippocampus did not alter depression-like
behaviors.

In conclusion, these findings support that β-CTF expression is sufficient to disrupt
hippocampus-dependent cognitive functions.

The C-terminal YENPTY motif was necessary for β-CTF to induce endosomal
dysfunction and synapse loss.
Endosome abnormalities mediated by APP β-CTFs have been reported across various
cell types, including human iPSC-induced neurons, PC12M cells, and N2a cells (Kim
et al., 2016; Kwart et al., 2019; Xu et al., 2016). Next, we investigated whether β-CTF
impacted endosomes in hippocampal neurons. Dissociated cultured hippocampal
neurons were co-transfected with APP-CTFs and Rab5-GFP, an endosomal marker
fused with GFP, at DIV7. In neurons expressing either Rab5-GFP alone or Rab5-GFP
with α-CTF, Rab5 puncta appeared uniform and smoothly rounded (Figure 5A-B).
However, in neurons co-transfected with β-CTF, Rab5 puncta were larger and
exhibited less uniform shapes, often appearing lobular, and showed robust
co-localization with β-CTF (Figure 5C). Notably, the morphology of lysosomes, as
observed by Lamp1 staining, remained unaffected by the expression of α- or β-CTF in

Formatted: Space After: 1 line

neurons (Supplementary figure 4A-C), thereby suggesting a specific interaction of β-CTF with endosomes.

The C-terminal YENPTY motif of APP was found to be crucial for its interaction with endosomes (Lai et al., 1995). To elucidate the significance of this interaction, we expressed a mutant form of β-CTF (β-CTF$_{mut}$), where the YENPTY motif was substituted with AENATA. Interestingly, β-CTF$_{mut}$ was expressed at similar levels to wildtype β-CTF and exhibited comparable Aβ production (Figure 5D-G). However, unlike wildtype β-CTF, β-CTF$_{mut}$ showed a more diffuse distribution throughout the neurons and failed to induce enlarged Rab5 puncta (Figure 5H). Notably, pyramidal neurons transiently expressing β-CTF$_{mut}$ displayed a higher spine density compared to those expressing wildtype β-CTF (Figure 5I-J). These findings underscore the critical role of the YENPTY motif-mediated interaction with endosomes in β-CTF-induced spine loss.

Spine loss induced by β-CTF was prevented by Rab5 inhibition.

To investigate the downstream mechanism responsible for β-CTF-induced synaptic loss, we analyzed the proteomic alterations triggered by β-CTF in dissociated cultured hippocampal neurons (Figure 6A). Our findings revealed significant changes in protein levels upon exposure to β-CTF (Figure 6B). Notably, the expression of Synapsin 1, a presynaptic protein associated with synaptic vesicles, and GluR1, an AMPA receptor subunit, were both diminished in neurons expressing β-CTF (Figure 6C-D). Gene Ontology analysis further elucidated that β-CTF expression downregulated proteins involved in membrane trafficking, synaptic vesicle cycle, pre-synapse, and post-synapse (Figure 6E-F), aligning with previous observations indicating that β-CTF induces synaptic dysfunction and endosomal abnormalities(Zhou et al., 2019). Conversely, upregulated proteins were primarily associated with peptide metabolic processes and translation (Supplementary figure 5A-B).

Overexpression of β-CTF resulted in the enlargement of Rab5-positive endosomes,

similar to the effects observed with a constitutively active mutant of Rab5, $Rab5_{Q79L}$

(Supplementary figure 5C) (Kim et al., 2016). Notably, the introduction of a dominant

negative mutant of Rab5, $Rab5_{S34N}$, attenuated the endosome enlargement induced by

β-CTF (Figure 6G-H). In neurons expressing $Rab5_{S34N}$, there was reduced

co-localization of β-CTF with Rab5 (Figure 6G-H). Critically, co-expression of

$Rab5_{S34N}$ with β-CTF effectively mitigated the spine loss induced by β-CTF in

hippocampal slice cultures (Figure 6I-J). These findings underscored that Rab5

overactivation-induced endosomal dysfunction contributed to β-CTF-induced spine

loss.

Discussion

Treatment with Lecanemab or Donanemab, two Aβ antibodies, significantly slowed down AD progression by approximately 30% in phase III clinical trials (Sims et al., 2023; van Dyck et al., 2023). The success of these antibodies in modifying AD progression validates Aβ as a cause of AD pathogenesis. However, the relatively modest benefits have raised additional questions. Despite effectively reducing amyloid plaques to near baseline levels after 18 months of treatment in some patients, functional benefits were limited to only ~30%. This raises the question of whether amyloid plaque is the primary source of Aβ-related toxicity or if these patients were treated too late to reverse their disease progression more effectively. This study aims to investigate whether Aβ-associated pathways could lead to pathogenesis beyond secreted Aβ and amyloid plaques, which falls outside the scope of Aβ antibody-based therapies due to their inability to penetrate cytoplasmic membranes.

AD is characterized by significant synaptic degeneration. Conventionally, Aβ is considered detrimental to synapses, thus synaptic loss in AD is attributed to various forms of Aβ, including amyloid plaques (Kamenetz et al., 2003; Oakley et al., 2006). However, conflicting studies have suggested that Aβ may actually promote synaptogenesis (Abramov et al., 2009; Bittner et al., 2009; Puzzo et al., 2008; Zhou et al., 2022). The exact role of Aβ in AD-related synaptic degeneration remains elusive. Most investigations have focused on the non-cell-autonomous function of Aβ after its secretion from neurons (Reinders et al., 2016). The potential contribution of intracellular Aβ or its precursors to synaptic toxicity has been largely unexplored. Utilizing a sparse neuron transfection system, we systematically examined this question. Intriguingly, among Aβ, APP, and major APP cleavage products, only β-CTF induced synaptic toxicity in a cell-autonomous manner (Figure 1, 2, Suppplementary figure 2), aligning with predictions based on indirect evidence (Kwart et al., 2019; Lee et al., 2022; Tamayev et al., 2012; Vaillant-Beuchot et al., 2021; Xu et al., 2016). In mouse brains, β-CTF also triggered substantial synaptic loss

and cognitive deficits in the absence of amyloid plaques (Figure 3C), further
supporting the notion that synaptic loss and amyloid plaque formation are mediated
by distinct mechanisms.

It is noteworthy that β-CTF induces spine loss independent of Aβ, implying that while
synaptic degeneration and amyloidogenesis both occur downstream of β-CTF, they
may represent two parallel processes independent of each other following APP
β-cleavage. Aβ antibodies effectively facilitate the clearance of secreted Aβ, including
that deposited in amyloid plaques. However, β-CTF localizes within the intracellular
membrane of endo/lysosomal vesicles, rendering it inaccessible to Aβ antibodies.
Consequently, Aβ antibody-based therapies cannot mitigate the neuronal toxicities
initiated by β-CTF. This limitation could partially explain the restricted clinical
benefits observed with Aβ antibodies. Addressing β-CTF-mediated synaptic toxicity
should be prioritized in the future development of improved AD therapies, potentially
in conjunction with Aβ antibodies.

Endosomal dysfunction is an early neuropathological signature of AD and Down
syndrome (DS), its downstream functional impacts remain unclear. One of the
impacts is deficits of NGF signaling in basal forebrain cholinergic neurons (BFCNs)
(Salehi et al., 2006; Xu et al., 2016). Here, we report endosomal dysfunction-induced
by β-CTF also contributes to synaptic dysfunction. Synaptic abnormalities are
corrected by inhibiting Rab5 activity to restore endosomal function (Figure 6G-J),
indicating that endosomal abnormalities may serve as a potential target for AD
treatment.

β-CTF elicited significant spine loss, whereas α-CTF or β'-CTF, which are only 16 or
10 amino acids shorter at their N-terminal ends compared to β-CTF, did not exhibit
this detrimental effect. Since the C-terminal **YENPTY** motif of β-CTF is essential for
this activity, likely through its interaction with endosomal proteins, β-CTF represents

the minimum size of the APP fragment capable of inducing cell-autonomous synaptic

toxicity. However, the specific contribution of the N-terminal domain of β-CTF to

synaptic loss remains unclear. It remains to be investigated whether longer fragments

containing β-CTF could also induce synaptic loss. If confirmed, strategies aimed at

reducing APP β-cleavage while promoting α- or β'-cleavage to decrease both Aβ and

β-CTF may offer a more effective treatment for AD in the future.

Materials and Methods

Dissociated hippocampal neuron culture and transfection

We prepared hippocampal neurons from embryonic day 17/18 (E17/18) SD rats. Initially, whole hippocampi were dissected under stereomicroscopes and then digested into single cells using papain (Worthington) for 25 minutes at 37 ℃. The dissociated cells were seeded at a density of 180,000 cells per well in a 24-well plate, using Neurobasal media (Thermo Fisher Scientific) supplemented with 2% B27 (Life technology), 0.25% Glutamax (Thermo Fisher Scientific), and 100 units/mL Penicillin/Streptomycin (Life technology). Neurons were cultured in incubators maintained at 37 ℃ with 5% CO_2, with fresh media half replenished every 5 days. For virus infection, dissociated hippocampal neurons at day *in vitro* (DIV) 14 were exposed to purified lentivirus. After 5 days, the infected neurons were lysed in RIPA buffer (Solarbio) supplemented with a protease inhibitor cocktail (Selleck) for further analysis. For plasmid transfection, dissociated hippocampal neurons at DIV7 were transfected with Lipofectamine 2000 (Thermo Fisher Scientific) following the manufacturer's instructions.

Western blotting and immunofluorescent staining

For western blotting, we first removed the culture medium, and then immediately placed neurons or HEK293T cells on ice. Subsequently, they were lysed using 2× loading buffer (100 mM Tris-HCl, 150 mM NaCl, 4% SDS, 20% glycerol). The lysates were agitated for 10 minutes at room temperature, followed by boiling at 98 ℃ for 10 minutes and centrifugation at 14,000 × g for 10 minutes. The proteins were separated using 8%-12% Tris-glycine SDS-PAGE and transferred to a PVDF membrane (Millipore, 0.22 μm). Primary antibodies were then incubated in blocking buffer overnight at 4 °C, followed by secondary antibodies at room temperature for 2 hours. The signal was detected using an Amersham Imager 600 (GE Healthcare), and densitometry was measured using ImageJ.

For immunofluorescent staining, we first washed neurons on cover glasses with PBS

(supplemented with 1 mM $CaCl_2$ and 5 mM $MgCl_2$) once. Then, we permeabilized

them with 0.15% Triton-X in PBS for 15 minutes. After blocked with 5% bovine

serum albumin in PBS for 30 minutes at room temperature, the samples were

incubated with primary antibodies in blocking buffer at 4 °C overnight. The following

day, the samples were washed three times with PBS and then incubated with

secondary antibodies at room temperature for 1 hour in the dark. Subsequently, the

samples were washed three times with PBS and mounted in Mounting Medium with

DAPI (Solarbio). Finally, images were acquired using an Andor spinning disk or

Nikon confocal microscopy system. All images were captured and analyzed in a

blinded manner.

Organotypic hippocampal slice culture, transfection and imaging

We prepared organotypic hippocampal slice cultures following previously established

protocols (Chen et al., 2014). Briefly, slices were obtained from P7-8

Sprague-Dawley rats. At DIV 3, slices were biolistically transfected using a gene gun

(Bio-Rad). Gold particles (1.6 μm in diameter; Bio-Rad) were coated with 25 μg of

cDNA along with 5 μg of GFP. Live cell imaging was conducted at DIV9/10 using a

Nikon confocal system equipped with a water-immersion 60× objective. Protrusions

from dendrites longer than 0.4 μm were counted as spines. For spine density analysis,

secondary basal dendrites were selected, and 2-5 different dendrites were imaged

from each pyramidal cell. Both image acquisition and spine counting were carried out

in a blinded manner.

Aβ40 and Aβ42 ELISA

We analyzed all cell sample sets using Aβ40 and Aβ42 ELISA kits (Invitrogen,

KHB3481 and KHB3441) according to the manufacturer's instructions. The cells were

sonicated in RIPA buffer (Solarbio) containing a protease inhibitor cocktail (Selleck),

followed by centrifugation at 12,000 × g for 10 minutes to prepare the samples.

Lentivirus production

We inserted β-CTF-flag and GFP into the pFHTrePW vector (Ni et al., 2023) backbone under the control of the Tet Response Element (TRE 3G). The rtTA for the tet-off system was expressed under the control of the human synapsin I promoter. The pFHTrePW plasmid containing β-CTF-flag or GFP, along with PSPAX2 and PMD2g, was transfected into HEK293T cells. After 48 hours, the cell culture medium containing the virus was collected and filtered through a 0.45μm syringe filter (Millipore). Subsequently, it was concentrated using ultracentrifugation, aliquoted, and stored at -80 ℃.

Adeno-Associated virus production

We performed AAV purification using triple-transfected HEK293T cells. Briefly, we transfected pHelper, pAAV2-8, and pAAV-MCS containing GFP or β-CTF into HEK293T cells at a ratio of 1:1:1. After 72 hours of incubation, the culture medium was collected in a 50 mL centrifuge tube, and the cell pellet was collected in another tube with lysis buffer (50 mM Tris-HCl, 150 mM NaCl, 2 mM $MgCl_2$, pH 8.0). The cell pellet was subjected to three cycles of freezing and thawing, followed by mixed with the culture medium. PEG8000 (10%) and NaCl (1 M/L) were added to the mixture. After incubating for 1 hour on ice, the mixture was centrifuged for 15 minutes at $12,000 \times g$ and 4 ℃. The precipitate was gently resuspended in lysis buffer containing DNase I (Roche) using a 1000-ml pipette. After a 30-minute incubation at 37 ℃, chloroform in a 1:1 (v:v) ratio was added to each tube, shaken for 1 hour, and then centrifuged for 15 minutes at $12,000 \times g$ and 4℃. To harvest the concentrated AAV, the supernatant was processed using an ultrafiltration column (Millipore).

Electrophysiology

We performed patch-clamp recordings from hippocampal pyramidal cells in acute brain slices. Adult mice approximately 8 weeks old were anesthetized with isoflurane and then decapitated. The hippocampus was harvested intact and placed into cold

choline cutting buffer (110 mM Choline Cl, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 25 mM

$NaHCO_3$, 7 mM $MgSO_4$, 25 mM D-glucose, 3.1 mM Na Pyruvate, 11.6 mM Na

Ascorbate, 0.5 mM $CaCl_2$ at pH 7.25). Using a vibrating microtome (Leica VT1200S),

the hippocampus was sliced into 400 μm sections, followed by a 30-minute recovery

period at 34 °C and another 30-minute recovery period at room temperature in

artificial cerebrospinal fluid (ACSF).

The recording external solution consisted of ACSF containing (in mM): 127 NaCl, 2.5

KCl, 12.5 NaH_2PO_4, 25 $NaHCO_3$, 25 D-glucose, 2.5 $CaCl_2$, and 1.3 $MgCl_2$, aerated

with 95% O_2/5% CO_2 to maintain a pH around 7.25. Pipettes (TW150F-4, World

Precision Instruments) with a resistance of 4–6 mΩ were filled with an internal

solution containing (in mM): 115 cesium methanesulfonate, 20 CsCl, 10 HEPES, 2.5

$MgCl_2$, 4 Na_2ATP, 0.4 Na_3GTP, 10 Na phosphocreatine, and 0.6 EGTA (pH 7.25).

Miniature excitatory postsynaptic currents (mEPSCs) were recorded by whole-cell

voltage clamp. The mEPSCs were recorded at a holding potential of -70 mV in the

presence of 1 μM tetrodotoxin (TTX, MedChem express, HY-12526A) and 100 μM

picrotoxin (PTX, Sigma, P1675). EPSCs were analyzed using MiniAnalysis software

(Synaptosoft) with an amplitude threshold of 5 pA.

Antibodies and drugs

The following antibodies were used in this study: Y188 antibody (Abcam, ab32136),

D10E5 antibody (Cell Signaling, 5606), Flag antibody (Sigma-Aldrich, F3165),

β-Tubulin III antibody (Sigma, T 2200), GAPDH antibody (Proteintech, 60004-1-Ig),

HA antibody (Cell Signaling, 3724S), GFP antibody (Abcam, ab13970), SYN

antibody (Millipore, AB1543), GluR1 antibody (Millipore, MAB2263), GluR2

antibody (Millipore, AB1768-1). Peroxidase-Conjugated secondary antibodies: Goat

Anti-Rabbit IgG (YESEN, 33101ES60), Goat Anti-Mouse IgG (YESEN, 33201ES60).

Fluorescent secondary antibodies: Alexa 488 Goat anti-chicken IgG (Thermo Fisher,

A11039), Alexa 568 Goat anti-chicken IgY (Thermo Fisher, A11041), Alexa 568

Goat anti-rabbit IgG (Thermo Fisher, A11036), Alexa 647 Goat anti-Mouse IgG

(Thermo Fisher, A21236). Inhibitors: PF-03084014 (Selleck, S7731).

Plasmids construction

We sub-cloned the expression of GFP, BACE1-GFP, GFP-Rab5, APP,

APP-IRES-BACE1 or related mutations into the pCAGGS mammalian expression

vector. The AAV constructs of GFP and β-CTF-HA were sub-cloned into the

pAAV-MCS vector. Lentivirus constructs of GFP, β-CTF, and β-CTF-IRES-GFP

were sub-cloned into the pFHTrePW vector backbone under the control of the TRE

3G.

Behavioral Analysis

432   Animals underwent a 1-hour habituation period in the room before all behavioral tests.

433   All tests were conducted and analyzed in a blinded manner.

Open field test (OFT)

The mice were introduced into a 40×40 cm box without roofs and permitted to

explore the apparatus freely for 1 hour. Their activities were monitored using the

Ethovision video tracking system (Noldus Information Technology Inc., Leesburg,

VA, USA). The field was divided into a center area (24×24 cm) and the whole arena.

Parameters such as total travel distance, duration in the center area, velocity, and

rearing frequency were automatically recorded and analyzed.

Y maze

442   Each mouse was placed in one arm of a Y-maze (with arms measuring 30 cm in

443   length) and allowed to explore the maze freely for 8 minutes. The movements of the

444   mice were captured by a video camera positioned above the arena. The number of

445   alternations and entries were analyzed using the Ethovision video tracking system.

Elevated plus maze (EPM)

Anxiety-like behavior was assessed using an elevated plus maze (EPM), which is an elevated apparatus shaped like a plus sign (+). It consists of two open arms ($35 \times 5 \times 0.3$ cm), two closed arms ($36 \times 5 \times 18$ cm) with 15 cm walls and open roofs, and a 5×5 cm central area. At the beginning of the test, each mouse was placed in the center of the maze facing an open arm and allowed to explore the maze freely for 5 minutes. The distance explored and time spent in the apparatus were recorded and analyzed using the SMART software (Panlab, Barcelona, Spain).

Water T maze (WTM)

Spatial learning and memory were evaluated using the Water T Maze (WTM) behavioral paradigm. In this task, mice were trained to utilize spatial cues within a room to locate a concealed platform and escape from water. In the reversal test, the hidden platform was relocated to the opposite arm to assess cognitive flexibility. The testing apparatus consisted of a plus maze, with each arm measuring 45 cm in length and 10 cm in width, constructed from clear Plexiglas. Each arm was designated as N, S, E, or W. A divider was positioned to block off an arm, allowing the mice to choose only the E or W arm for escape. The water temperature was maintained at 25-26 °C and made opaque by adding white, nontoxic powder. An escape platform was submerged 1 cm below the water's surface on the E side of the maze, rendering it invisible to the mice.

At the beginning of each trial, the divider was inserted to block access to the appropriate arm, and mice were placed at the starting point. The experimenter recorded whether each response was correct or incorrect, and mice were permitted to remain on the platform for 10 seconds before being removed. Each day, mice underwent 5 trials with starting points semi-randomized between the N and S positions. The criterion for acquisition was achieving 80% or more correct responses averaged across the 5 trials for 2 consecutive days. Following acquisition by both groups, reversal training commenced. The hidden platform was relocated to the

opposite side (W), and the same procedure was repeated until the mice successfully

learned the new platform position. Mice that achieved 80% or more correct responses

across the 5 trials for 2 consecutive days were deemed to have passed the test.

Fear conditioning (FC)

The fear conditioning test comprised two components: contextual fear conditioning

and cued fear conditioning. During the training stage, mice were introduced into

conditioning boxes and allowed to freely explore the environment for 3 minutes. The

baseline freezing behavior was recorded using a visual camera and analyzed with the

Ethovision video tracking system. Subsequently, mice were exposed to an 80 dB,

2 kHz tone for 30 seconds (conditioned stimulus, CS), during which an inescapable

0.3 mA foot shock (unconditioned stimulus, US) was delivered in the last 2 seconds of

the tone. This procedure was repeated three times with a 30-second

inter-stimulus-interval.

For the contextual fear conditioning test, conducted on the second day, mice were

placed in the same conditioning box used on the previous day for 5 minutes, and their

freezing behavior was recorded for further analysis. In the cued fear conditioning test,

conducted on the third day after training and contextual fear conditioning, mice were

placed in a different chamber from the previous one. After a 3-minute free exploration

period, mice were exposed to a 3-minute tone stimulus (2 kHz, 80 dB). Freezing

behavior during the 3-minute tone stimulus was measured and analyzed using the

Ethovision video tracking system.

Tail suspension test (TST)

Each mouse was suspended by its tail using tape affixed to the ceiling of a

three-walled rectangular compartment measuring 30 cm in height, 15 cm in width, and

15 cm in depth. The mice dangled downward, with ample space provided for

movement. Video recording was conducted from the side, capturing the animals'

immobile time (defined as cessation of limb movements for more than 2 seconds) during the final 4 minutes of a 6-minute session.

Stereotactic virus injection

Mice aged 4 weeks or 7-8 weeks received injections of lentivirus or AAV in the hippocampus at coordinates ML:±1.5 mm, AP: -2 mm, DV: -1.5 mm. For spine density analysis, mice were harvested after 2 months; for behavioral tests, mice were evaluated after 4 months.

Liquid chromatography tandem mass spectrometry (LC-MS/MS) sample preparation and analysis

For protein sample preparation, neurons seeded in 24-well plates underwent lysis using ultrasonic waves in RIPA buffer. After centrifugation at $18,000 \times g$ for 10 minutes at 4 ℃, the supernatants were collected and quantified using a BCA assay (Thermo Fisher). Protein extracts were subjected to an in-solution digest protocol, involving reduction with 5 mM Tris(2-carboxyethyl) phosphine hydrochloride (TCEP) (Aldrich, USA) followed by alkylation with 10 mM iodoacetamide (IAA) (Sigma, USA). Trypsin was added at a 1:100 ratio, and the mixture was incubated at 37 ℃ overnight in the dark. Digested peptides were then collected by centrifugation and desalted using C18 tips (Pierce, USA) for subsequent analysis.

For LC-MS/MS analysis, the peptide mixture was analyzed using an online EASY-nLC 1000 HPLC system coupled with an Orbitrap Fusion mass spectrometer (Thermo Fisher Scientific). The sample was loaded directly onto a 15-cm homemade capillary column (100 μm I.D., C18-AQ 1.9 μm resin, Dr. Maisch). Mobile phase A consisted of 0.1% formic acid (FA), 2% acetonitrile (ACN), and 98% H_2O, while mobile phase B comprised 0.1% FA, 2% H2O, and 98% ACN. A 180-minute gradient (mobile phase B: 2% at 0 min, 5% at 7 min, 20% at 127 min, 35% at 167 min, 95% at 173 min, 95% at 180 min) was employed at a static flow rate of 300 nl/min.

We acquired data for proteomic analysis in a data-dependent mode, starting with one

full MS1 scan in the Orbitrap (m/z: 300-1800; resolution: 120,000; AGC target value:

500,000; maximal injection time: 50 ms), followed by an MS2 scan in the linear ion

trap (32% normalized collision energy; maximal injection time: 250 ms). The

isolation window was set at 1.6 m/z.

Statistics

Data were processed using Microsoft Excel, and statistical analysis was performed

with GraphPad Prism 7. Figures were made using Adobe Illustrator V26.3.1. Sample

sizes and the statistical analyses performed are described in the respective figure

legends. For all analyses, $P<0.05$ was considered statistically significant.

Acknowledgements

We thank the staff members of the Animal Facility at the National Facility for Protein Science in Shanghai (NFPS), Shanghai Advanced Research Institute, Chinese Academy of Sciences, China for providing assistance in mouse breeding and maintenance. This study was supported by Shanghai Basic Research Pioneer Project. This project received support from National Natural Science Foundation of China to Y.C. (31671044, 91849204); National Program on Key Research Project of China to Y.C. (2016YFA0501901); Shanghai Municipal Science and Technology Major Project to Y.C. (Grant No. 2019SHZDZX02).

Author contributions

Conceptualization: Yelin Chen, Yang Geng.; Methodology: Yelin Chen, Yang Geng.,

Mengxun Luo; Investigation and Validation: Mengxun Luo., Jia Zhou., Cailu Sun,

Chaoying Fu, Wanjia Chen, Chenfang Si; Visualization: Mengxun Luo; Formal

analysis: Mengxun Luo, Jia Zhou, Cailu Sun, Chenfang Si, Data curation: Mengxun

Luo, Chenfang Si, Yaoyang Zhang; Writing – original draft preparation: Mengxun

Luo; Writing – review & editing: Yelin Chen, Yang Geng; Supervision and Project

administration: Yelin Chen; Funding acquisition: Yelin Chen.

Conflict of Interest

The authors declare no competing interests.

References

Abramov, E., Dolev, I., Fogel, H., Ciccotosto, G.D., Ruff, E., and Slutsky, I. (2009). Amyloid-beta as a positive endogenous regulator of release probability at hippocampal synapses. Nature neuroscience *12*, 1567-1576.

Bittner, T., Fuhrmann, M., Burgold, S., Jung, C.K., Volbracht, C., Steiner, H., Mitteregger, G., Kretzschmar, H.A., Haass, C., and Herms, J. (2009). Gamma-secretase inhibition reduces spine density in vivo via an amyloid precursor protein-dependent pathway. The Journal of neuroscience : the official journal of the Society for Neuroscience *29*, 10405-10409.

Blennow, K., de Leon, M.J., and Zetterberg, H. (2006). Alzheimer's disease. Lancet (London, England) *368*, 387-403.

Chen, Y., Wang, Y., Ertürk, A., Kallop, D., Jiang, Z., Weimer, R.M., Kaminker, J., and Sheng, M. (2014). Activity-induced Nr4a1 regulates spine density and distribution pattern of excitatory synapses in pyramidal neurons. Neuron *83*, 431-443.

Cheng, X., He, P., Lee, T., Yao, H., Li, R., and Shen, Y. (2014). High activities of BACE1 in brains with mild cognitive impairment. The American journal of pathology *184*, 141-147.

Citron, M., Teplow, D.B., and Selkoe, D.J. (1995). Generation of amyloid beta protein from its precursor is sequence specific. Neuron *14*, 661-670.

de Wilde, M.C., Overk, C.R., Sijben, J.W., and Masliah, E. (2016). Meta-analysis of synaptic pathology in Alzheimer's disease reveals selective molecular vesicular machinery vulnerability. Alzheimer's & dementia : the journal of the Alzheimer's Association *12*, 633-644.

DeKosky, S.T., Scheff, S.W., and Styren, S.D. (1996). Structural correlates of cognition in dementia: quantification and assessment of synapse change. Neurodegeneration : a journal for neurodegenerative disorders, neuroprotection, and neuroregeneration *5*, 417-421.

Glenner, G.G., and Wong, C.W. (1984). Alzheimer's disease: initial report of the

purification and characterization of a novel cerebrovascular amyloid protein.

Biochemical and biophysical research communications *120*, 885-890.

Golde, T.E., Estus, S., Younkin, L.H., Selkoe, D.J., and Younkin, S.G. (1992).

Processing of the amyloid protein precursor to potentially amyloidogenic derivatives.

Science *255*, 728-730.

Gómez-Isla, T., Price, J.L., McKeel, D.W., Jr., Morris, J.C., Growdon, J.H., and

Hyman, B.T. (1996). Profound loss of layer II entorhinal cortex neurons occurs in

very mild Alzheimer's disease. The Journal of neuroscience : the official journal of the

Society for Neuroscience *16*, 4491-4500.

Israel, M.A., Yuan, S.H., Bardy, C., Reyna, S.M., Mu, Y., Herrera, C., Hefferan, M.P.,

Van Gorp, S., Nazor, K.L., Boscolo, F.S.*, et al.* (2012). Probing sporadic and familial

Alzheimer's disease using induced pluripotent stem cells. Nature *482*, 216-220.

Jiang, Y., Mullaney, K.A., Peterhoff, C.M., Che, S., Schmidt, S.D., Boyer-Boiteau, A.,

Ginsberg, S.D., Cataldo, A.M., Mathews, P.M., and Nixon, R.A. (2010).

Alzheimer's-related endosome dysfunction in Down syndrome is Abeta-independent

but requires APP and is reversed by BACE-1 inhibition. Proceedings of the National

Academy of Sciences of the United States of America *107*, 1630-1635.

JohnHardy, and Selkoe, D.J. (2002). The Amyloid Hypothesis of Alzheimer's Disease:

Progress and Problems on the Road to Therapeutics. Science *297*, 353-356.

Jonsson, T., Atwal, J.K., Steinberg, S., Snaedal, J., Jonsson, P.V., Bjornsson, S.,

Stefansson, H., Sulem, P., Gudbjartsson, D., Maloney, J.*, et al.* (2012). A mutation in

APP protects against Alzheimer's disease and age-related cognitive decline. Nature

488, 96-99.

Kamenetz, F., Tomita, T., Hsieh, H., Seabrook, G., Borchelt, D., Iwatsubo, T., Sisodia,

S., and Malinow, R. (2003). APP processing and synaptic function. Neuron *37*,

925-937.

Kessels, H.W., Nabavi, S., and Malinow, R. (2013). Metabotropic NMDA receptor

function is required for β-amyloid-induced synaptic depression. Proceedings of the

National Academy of Sciences of the United States of America *110*, 4033-4038.

Kim, S., Sato, Y., Mohan, P.S., Peterhoff, C., Pensalfini, A., Rigoglioso, A., Jiang, Y.,

and Nixon, R.A. (2016). Evidence that the rab5 effector APPL1 mediates

APP-βCTF-induced dysfunction of endosomes in Down syndrome and Alzheimer's

disease. Molecular psychiatry *21*, 707-716.

Konietzko, U. (2012). AICD nuclear signaling and its possible contribution to

Alzheimer's disease. Current Alzheimer research *9*, 200-216.

Kwart, D., Gregg, A., Scheckel, C., Murphy, E.A., Paquet, D., Duffield, M., Fak, J.,

Olsen, O., Darnell, R.B., and Tessier-Lavigne, M. (2019). A Large Panel of Isogenic

APP and PSEN1 Mutant Human iPSC Neurons Reveals Shared Endosomal

Abnormalities Mediated by APP β-CTFs, Not Aβ. Neuron *104*, 256-270.e255.

Lai, A., Sisodia, S.S., and Trowbridge, I.S. (1995). Characterization of sorting signals

in the beta-amyloid precursor protein cytoplasmic domain. The Journal of biological

chemistry *270*, 3565-3573.

Lee, S.E., Kwon, D., Shin, N., Kong, D., Kim, N.G., Kim, H.Y., Kim, M.J., Choi,

S.W., and Kang, K.S. (2022). Accumulation of APP-CTF induces mitophagy

dysfunction in the iNSCs model of Alzheimer's disease. Cell death discovery *8*, 1.

Leyns, C.E.G., and Holtzman, D.M. (2017). Glial contributions to neurodegeneration

in tauopathies. Molecular neurodegeneration *12*, 50.

Liu, C.Y., Ohki, Y., Tomita, T., Osawa, S., Reed, B.R., Jagust, W., Van Berlo, V., Jin,

L.W., Chui, H.C., Coppola, G.*, et al.* (2017). Two Novel Mutations in the First

Transmembrane Domain of Presenilin1 Cause Young-Onset Alzheimer's Disease.

Journal of Alzheimer's disease : JAD *58*, 1035-1041.

Masters, C.L., Simms, G., Weinman, N.A., Multhaup, G., McDonald, B.L., and

Beyreuther, K. (1985). Amyloid plaque core protein in Alzheimer disease and Down

syndrome. Proceedings of the National Academy of Sciences of the United States of

America *82*, 4245-4249.

Mintun, M.A., Lo, A.C., Duggan Evans, C., Wessels, A.M., Ardayfio, P.A., Andersen,

S.W., Shcherbinin, S., Sparks, J., Sims, J.R., Brys, M.*, et al.* (2021). Donanemab in

Early Alzheimer's Disease. The New England journal of medicine *384*, 1691-1704.

Mullan, M., Crawford, F., Axelman, K., Houlden, H., Lilius, L., Winblad, B., and

Lannfelt, L. (1992). A pathogenic mutation for probable Alzheimer's disease in the

APP gene at the N-terminus of beta-amyloid. Nature genetics *1*, 345-347.

Ni, J., Ren, Y., Su, T., Zhou, J., Fu, C., Lu, Y., Li, D., Zhao, J., Li, Y., Zhang, Y.*, et*

al. (2023). Loss of TDP-43 function underlies hippocampal and cortical synaptic

deficits in TDP-43 proteinopathies. Molecular psychiatry *28*, 931-945.

Nikolaev, A., McLaughlin, T., O'Leary, D.D., and Tessier-Lavigne, M. (2009). APP

binds DR6 to trigger axon pruning and neuron death via distinct caspases. Nature *457*,

981-989.

Nilsberth, C., Westlind-Danielsson, A., Eckman, C.B., Condron, M.M., Axelman, K.,

Forsell, C., Stenh, C., Luthman, J., Teplow, D.B., Younkin, S.G.*, et al.* (2001). The

'Arctic' APP mutation (E693G) causes Alzheimer's disease by enhanced Abeta

protofibril formation. Nature neuroscience *4*, 887-893.

Oakley, H., Cole, S.L., Logan, S., Maus, E., Shao, P., Craft, J., Guillozet-Bongaarts,

A., Ohno, M., Disterhoft, J., Van Eldik, L.*, et al.* (2006). Intraneuronal beta-amyloid

aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial

Alzheimer's disease mutations: potential factors in amyloid plaque formation. The

Journal of neuroscience : the official journal of the Society for Neuroscience *26*,

10129-10140.

Oddo, S., Caccamo, A., Shepherd, J.D., Murphy, M.P., Golde, T.E., Kayed, R.,

Metherate, R., Mattson, M.P., Akbari, Y., and LaFerla, F.M. (2003). Triple-transgenic

model of Alzheimer's disease with plaques and tangles: intracellular Abeta and

synaptic dysfunction. Neuron *39*, 409-421.

Pleen, J., and Townley, R. (2022). Alzheimer's disease clinical trial update 2019-2021.

Journal of neurology *269*, 1038-1051.

Puzzo, D., Privitera, L., Leznik, E., Fà, M., Staniszewski, A., Palmeri, A., and

Arancio, O. (2008). Picomolar amyloid-beta positively modulates synaptic plasticity

and memory in hippocampus. The Journal of neuroscience : the official journal of the

Society for Neuroscience *28*, 14537-14545.

Reinders, N.R., Pao, Y., Renner, M.C., da Silva-Matos, C.M., Lodder, T.R., Malinow,

R., and Kessels, H.W. (2016). Amyloid-β effects on synapses and memory require

AMPA receptor subunit GluA3. Proceedings of the National Academy of Sciences of

the United States of America *113*, E6526-e6534.

Salehi, A., Delcroix, J.D., Belichenko, P.V., Zhan, K., Wu, C., Valletta, J.S.,

Takimoto-Kimura, R., Kleschevnikov, A.M., Sambamurti, K., Chung, P.P.*, et al.*

(2006). Increased App expression in a mouse model of Down's syndrome disrupts

NGF transport and causes cholinergic neuron degeneration. Neuron *51*, 29-42.

Selkoe, D.J. (2002). Alzheimer's disease is a synaptic failure. Science *298*, 789-791.

Sims, J.R., Zimmer, J.A., Evans, C.D., Lu, M., Ardayfio, P., Sparks, J., Wessels,

A.M., Shcherbinin, S., Wang, H., Monkul Nery, E.S.*, et al.* (2023). Donanemab in

Early Symptomatic Alzheimer Disease: The TRAILBLAZER-ALZ 2 Randomized

Clinical Trial. Jama *330*, 512-527.

Sperling, R.A., Donohue, M.C., Raman, R., Rafii, M.S., Johnson, K., Masters, C.L.,

van Dyck, C.H., Iwatsubo, T., Marshall, G.A., Yaari, R.*, et al.* (2023). Trial of

Solanezumab in Preclinical Alzheimer's Disease. The New England journal of

medicine *389*, 1096-1107.

Tamayev, R., Matsuda, S., Arancio, O., and D'Adamio, L. (2012). β- but not

γ-secretase proteolysis of APP causes synaptic and memory deficits in a mouse model

of dementia. EMBO molecular medicine *4*, 171-179.

Terry, R.D., Masliah, E., Salmon, D.P., Butters, N., DeTeresa, R., Hill, R., Hansen,

L.A., and Katzman, R. (1991). Physical basis of cognitive alterations in Alzheimer's

disease: synapse loss is the major correlate of cognitive impairment. Annals of
neurology *30*, 572-580.

Thinakaran, G., and Koo, E.H. (2008). Amyloid precursor protein trafficking,
processing, and function. The Journal of biological chemistry *283*, 29615-29619.

Tyan, S.H., Shih, A.Y., Walsh, J.J., Maruyama, H., Sarsoza, F., Ku, L., Eggert, S.,
Hof, P.R., Koo, E.H., and Dickstein, D.L. (2012). Amyloid precursor protein (APP)
regulates synaptic structure and function. Mol Cell Neurosci *51*, 43-52.

Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu,
A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A.*, et al.* (2015). Proteomics.
Tissue-based map of the human proteome. Science *347*, 1260419.

Vaillant-Beuchot, L., Mary, A., Pardossi-Piquard, R., Bourgeois, A., Lauritzen, I.,
Eysert, F., Kinoshita, P.F., Cazareth, J., Badot, C., Fragaki, K.*, et al.* (2021).
Accumulation of amyloid precursor protein C-terminal fragments triggers
mitochondrial structure, function, and mitophagy defects in Alzheimer's disease
models and human brains. Acta neuropathologica *141*, 39-65.

van Dyck, C.H., Swanson, C.J., Aisen, P., Bateman, R.J., Chen, C., Gee, M.,
Kanekiyo, M., Li, D., Reyderman, L., Cohen, S.*, et al.* (2023). Lecanemab in Early
Alzheimer's Disease. The New England journal of medicine *388*, 9-21.

Vassar, R., Bennett, B.D., Babu-Khan, S., Kahn, S., Mendiaz, E.A., Denis, P., Teplow,
D.B., Ross, S., Amarante, P., Loeloff, R.*, et al.* (1999). Beta-secretase cleavage of
Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE.
Science *286*, 735-741.

Vohra, B.P., Sasaki, Y., Miller, B.R., Chang, J., DiAntonio, A., and Milbrandt, J.
(2010). Amyloid precursor protein cleavage-dependent and -independent axonal
degeneration programs share a common nicotinamide mononucleotide
adenylyltransferase 1-sensitive pathway. The Journal of neuroscience : the official
journal of the Society for Neuroscience *30*, 13729-13738.

Wang, X., Wang, Z., Chen, Y., Huang, X., Hu, Y., Zhang, R., Ho, M.S., and Xue, L.

(2014). FoxO mediates APP-induced AICD-dependent cell death. Cell death &

disease *5*, e1233.

Wei, W., Nguyen, L.N., Kessels, H.W., Hagiwara, H., Sisodia, S., and Malinow, R.

(2010). Amyloid beta from axons and dendrites reduces local spine number and

plasticity. Nature neuroscience *13*, 190-196.

Willem, M., Tahirovic, S., Busche, M.A., Ovsepian, S.V., Chafai, M., Kootar, S.,

Hornburg, D., Evans, L.D., Moore, S., Daria, A.*, et al.* (2015). η-Secretase processing

of APP inhibits neuronal activity in the hippocampus. Nature *526*, 443-447.

Xiao, C., Liu, Y., Xu, J., Gan, X., and Xiao, Z. (2018). Septal and Hippocampal

Neurons Contribute to Auditory Relay and Fear Conditioning. Frontiers in cellular

neuroscience *12*, 102.

Xu, W., Weissmiller, A.M., White, J.A., 2nd, Fang, F., Wang, X., Wu, Y., Pearn,

M.L., Zhao, X., Sawa, M., Chen, S.*, et al.* (2016). Amyloid precursor

protein-mediated endocytic pathway disruption induces axonal dysfunction and

neurodegeneration. The Journal of clinical investigation *126*, 1815-1833.

Zhang, X., and Song, W. (2013). The role of APP and BACE1 trafficking in APP

processing and amyloid-β generation. Alzheimer's research & therapy *5*, 46.

Zhang, Y.W., Thompson, R., Zhang, H., and Xu, H. (2011). APP processing in

Alzheimer's disease. Molecular brain *4*, 3.

Zhou, B., Lu, J.G., Siddu, A., Wernig, M., and Südhof, T.C. (2022). Synaptogenic

effect of APP-Swedish mutation in familial Alzheimer's disease. Science translational

medicine *14*, eabn9380.

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O.,

Benner, C., and Chanda, S.K. (2019). Metascape provides a biologist-oriented

resource for the analysis of systems-level datasets. Nature communications *10*, 1523.

Figure 1. APP only led to spine loss when co-expressed with BACE1.

Figure 1. APP only led to spine loss when co-expressed with BACE1.

A-B Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with APP. (filler, n =21; APP, n=16).

C-D Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with APP and BACE1 with a ratio of 15:1. (filler, n=8; APP plus BACE1, n=5)

E-F Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with BACE1. (filler, n=15; BACE1, n=15)

G Western blot and corresponding statistical analysis of APP (Y188), BACE1 and GAPDH from HEK293T cells expressing APP or APP plus BACE1 with a ratio of 15:1. GAPDH was measured as a loading control.

H Measurement of APP β-CTF (β C-Terminal Fragment) level from HEK293T cells expressing APP alone or APP and BACE1 with a ratio of 15:1. n=4.

I Western blot and corresponding statistical analysis of APP fragments (Y188) from HEK293T cells expressing APP plus BACE1 or APP_{MV} plus BACE1.

J Measurement of APP β-CTF levels from HEK293T cells expressing APP plus BACE1 or APP_{MV} plus BACE1. n=4.

K Measurement of Aβ40 levels from HEK293T cells expressing APP plus BACE1 or APP_{MV} plus BACE1. n=4.

L-M Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with APP and BACE1 or APP_{MV} and BACE1. (filler, n=12; APP plus BACE1, n=14; APP_{MV} plus BACE1, n=10)

All dendritic images were acquired from rat organotypic hippocampal slice cultures after transfection for 6-7 days. Statistics: One-way ANOVA or Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM. Scale bars, 10 μm.

Figure 2. β-CTF induced spine loss independent of Aβ.

Figure 2. β-CTF induced spine loss independent of Aβ.

A-B Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with α-CTF, β'-CTF or β-CTF. (filler, n=19; α-CTF, n=7; β'-CTF, n=8; β-CTF, n=7)

C Western blot of APP fragments from HEK293T cells expressing α-CTF-flag, β'-CTF-flag and β-CTF-flag (flag antibody).

D Western blot of APP fragments from HEK293T cells expressing β-CTF-flag treatment with vehicle or PF (flag antibody).

E Measurement of Aβ40 secreted from HEK293T cells expressing β-CTF treatment with vehicle or PF. n=4.

F-G Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with β-CTF after treated with vehicle or PF. (filler, n=11; β-CTF, n=7, β-CTF plus PF, n=14)

H-I Representative images and their spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP together with APP and BACE1 after treated with vehicle or PF. (APP plus BACE1 with vehicle, n=9; APP plus BACE1 with PF, n=14)

J Western blot of APP fragments (Y188) from HEK293T cells co-expressing APP and BACE1 after treated with vehicle or PF.

K Measurements of Aβ40 from HEK 293T cells expressing APP and BACE1 after treated with vehicle or PF. n=4.

L Measurements of Aβ40 from HEK293T cells expressing APP and BACE1 or APP$_{\Delta 59}$ and BACE1. n=4.

M Measurements of Aβ42 from HEK293T cells expressing APP and BACE1 or APP$_{\Delta 57}$ and BACE1. n=4.

N-O Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP alone or together with APP$_{\Delta 59/\Delta 57}$ and BACE1. (filler, n=9; APP$_{\Delta 59}$ and BACE1, n=7; APP$_{\Delta 57}$ and BACE1, n=6)

All dendritic images were acquired from rat organotypic hippocampal slice cultures after transfection for 6-7 days. PF, PF03084014, a γ secretase inhibitor. Statistics: One-way ANOVA or Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM. Scale bars, 10 μm.

Figure 3. Expression of β-CTF damaged synapses in mouse brains.

Figure 3. Expression of β-CTF damaged synapses in mouse brains.

A Representative images of mouse CA1 pyramidal neurons infected with lentivirus expressing GFP and β-CTF. Scale bar, 100 μm.

B-C Representative images from basal dendrites and quantitation of their spine density in CA1 pyramidal neurons infected with lentivirus expressing GFP alone or GFP with β-CTF in mouse cortices. Scale bar, 10 μm. GFP only, n=10; GFP plus β-CTF, n=13.

D-F Representative recording traces and quantitations of mEPSCs from adult mouse hippocampal neurons infected with lentivirus (identified with GFP signal) and neighboring uninfected neurons. n=10.

Statistics: Two-way ANOVA or Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM.

Figure 4. Expression of β-CTF damaged cognitive function in mice in the absence of plaque formation.

Figure 4. Expression of β-CTF damaged cognitive function in mice in the absence of plaque formation.

A Schematic diagram showing the time line for stereotactic injection of AAV, behavioral training and tests.

B Representative images showing immunofluorescence staining of expressed GFP or β-CTF-HA by AAV in mouse hippocampi (WT). Scare bars represent 1000 μm (left) and 500 μm (right).

C Images of mouse hippocampus (WT) stained with ThS and HA antibody after infected with AAV encoding β-CTF-HA. Scale bar, 100 μm.

D Representative image of ThS staining of a 5XFAD mice hippocampal slice. Scale bar, 100 μm.

E Weight of mice infected with AAV encoding GFP or β-CTF in their hippocampi. GFP, n=14; β-CTF, n=15.

F Quantitations of spontaneous alternation in Y-maze from mice infected with AAV encoding GFP or β-CTF. GFP, n=14; β-CTF, n=15.

G Schematic diagram showing fear conditioning test design.

H Quantitation of freezing in contextual and cued fear conditioning tests from mice infected with AAV encoding GFP and β-CTF in their hippocampi. GFP, n=13; β-CTF, n=15.

I Schematic diagram showing water T maze test design.

J-K The percentage of correct responses across the 5 trials of water T maze test was quantified in each day of Acquisition (J) or Reversal (K). GFP, n=14; β-CTF, n=15.

Statistics: Two-way ANOVA or Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM.

Figure 5. The C-terminal YENPTY motif was necessary for β-CTF to induce endosomal dysfunction and synapse loss.

Figure 5. The C-terminal YENPTY motif was necessary for β-CTF to induce endosomal dysfunction and synapse loss.

A Immunofluorescent staining of GFP (green) from dissociated rat hippocampal neurons expressing Rab5-GFP.

B Immunofluorescent staining of GFP (green) and flag (Red) from dissociated rat hippocampal neurons expressing Rab5-GFP and α-CTF-flag.

C Immunofluorescent staining of GFP (green) and flag (Red) from dissociated rat hippocampal neurons expressing Rab5-GFP and β-CTF-flag.

D Schematic diagram of the β-CTF and β-CTFmut.

E Western blot of APP fragments from HEK293T cells expressing β-CTF-flag and β-CTFmut-flag with a flag antibody.

F Quantitations of APP fragments from HEK293T cells expressing β-CTF-flag and β-CTFmut-flag. n=4.

G Measurement of Aβ40 from HEK 293T cells expressing β-CTF or β-CTFmut. n=5.

H Immunofluorescent staining of GFP (green) and flag (Red) from dissociated rat hippocampal neurons expressing Rab5-GFP and β-CTFmut-flag.

I-J Representative images and spine density of basal dendrites from CA1 pyramidal neurons transiently expressing GFP together with β-CTF or β-CTFmut for 6-7 days in rat organotypic hippocampal slice cultures. β-CTF, n=12; β-CTFmut, n=15.

Scale bar, 10 μm. Statistics: Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM.

Figure 6. Spine loss induced by β-CTF was prevented by Rab5 inhibition.

Figure 6. Spine loss induced by β-CTF was prevented by Rab5 inhibition.

A Representative images of dissociated rat hippocampal neurons infected with lenti virus encoding GFP or β-CTF. Scale bar, 100 μm.

B Volcano plot of quantitative mass spectrometry analysis showing protein omics changes in neurons infected with lenti virus encoding β-CTF v.s. GFP. Proteins with fold change >1.2 or <0.83 and p value < 0.05 are colored.

C-D Western blot and corresponding statistic analysis of Synapsin 1, GluR 1, GluR 2 in dissociated hippocampal neurons infected with lentivirus expressing GFP or β-CTF. Tuj 1 was used as an internal control.

E The Gene Ontology (GO) processes analysis of decreased proteins by Metascape.

F The Reactome Gene Sets processes analysis of decreased protein by Metascape.

G-H Immunofluorescent staining of GFP (green), flag (Red) and HA (magenta) from dissociated rat hippocampal neurons expressing Rab5-GFP and β-CTF-flag or together with $Rab5_{S34N}$-HA. Scale bar, 10 μm.

I-J Representative images and measurements of spine density from basal dendrites or CA1 pyramidal neurons transiently expressing GFP and β-CTF with or without $Rab5_{S34N-HA}$ for 6-7 days in rat organotypic hippocampal slice cultures. Scale bar, 10 μm. (β-CTF, n=17; β-CTF plus $Rab5_{S34N}$, n=11)

Statistics: Two-way ANOVA or Student's t test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Error bars show SEM.