The penetration ring is a novel infection structure formed by the penetration peg for

invading plant cell membrane in rice blast fungus

Wenqin Fang[1,2,3], Xiaoyu Zai[1,2], Jia Chen[1,2], Yakubu Saddeeq Abubakar[1,2,4], Qiuqiu Wu[1,2], Zhenyu

Fang[1,2], Xiuwei Huang[1,2], Xiang Gan[1,2], Daniel J. Ebbole[5], Zonghua Wang[1,2,3,*], Wenhui

Zheng[1,2,*]

[1] State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, College of Plant

Protection, Fujian Agriculture and Forestry University, Fuzhou, Fujian, 350002, China.

[2] Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant

Protection, Fujian Agriculture and Forestry University, Fuzhou, Fujian, 350002, China.

[3] Fuzhou Institute of Oceanography, Minjiang University, Fuzhou 350108, China

[4] Department of Biochemistry, Faculty of Life Sciences, Ahmadu Bello University, Zaria, 810281,

Nigeria

[5] Department of Plant Pathology & Microbiology, Texas A&M University, College Station, TX,

77843, USA

* Corresponding author:

Wenhui Zheng (wenhuiz@fafu.edu.cn)

Zonghua Wang (wangzh@fafu.edu.cn)

Author Contributions: D.J.E., Z.W., and W.Z. designed research; W.F., X.Z., J.C., Q.W., Z.F.,

X.H., X.G., and W.Z. performed research; W.F., Z.W., and W.Z. analyzed data; and W.F., Y.S.A.,

Z.W., and W.Z. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: penetration ring, secreted protein, *Magnaporthe oryzae*, peg, appressorium

Abstract

Many fungal pathogens develop specialized infection structures such as appressoria to penetrate plant cells. However, it is not clear whether special structures are formed after cell wall penetration before invading host cytoplasm membrane in hemibiotrophic pathogens. Here, we showed that a penetration ring consisting of Ppe1 secreted proteins is formed after appressorium-mediated cell wall penetration and remained at the base of penetration site after invading plant cytoplasm membrane in the rice blast fungus *Magnaporthe oryzae*. The same persistent Ppe1 ring is formed after the penetration of neighboring cells by transpressoria. *PPE1* is specifically expressed during plant infection and the Δ*ppe1* mutant is defective in penetration and invasive growth. Blockage of penetration peg formation impedes the development of the Ppe1 ring. Close examinations showed that the penetration ring is formed at the collar of penetration pegs between plant cell wall and cytoplasm membrane and it is persistent as a fixed ring even after invasive hyphae invaded neighboring cells. Furthermore, Ppe1 is a member of an expanded family of secreted proteins that are unique to fungal pathogens using extreme appressorium turgor for plant penetration. Other members of the Ppe1 family also localize to the penetration ring for anchoring on cytoplasm membrane during plant infection. Taken together, a penetration ring consisting of a family of secreted proteins is formed between plant cell wall and cytoplasm membrane, which may function as a novel physical structure at the interface between the tip of penetration pegs and plant cytoplasm membrane before the differentiation of invasive hyphae.

Significance Statement

Like many other plant pathogens, the rice blast fungus forms melanized appressoria (specialized infection structures) to penetrate plant cells. In this study, we showed that a penetration ring is formed by penetration pegs after appressorium- or transpressorium-mediated penetration of plant cell wall. This ring of secreted proteins is persistent at the point of penetration pegs invading plant cytoplasm membrane to form invasive hyphae. Therefore, after the penetration of plant cell wall, the rice blast fungus forms a penetration ring that consists of a family of secreted proteins unique to pathogens using extreme appressorium turgor for penetration and may function as a physical

structure for anchoring onto plant cytoplasm membrane and developing invasive hyphae in penetrated cells.

Introduction

Magnaporthe oryzae, a filamentous ascomycete, is the causal agent of the devastating rice blast disease, which poses a significant threat to global rice production (1, 2). In addition to rice, *M. oryzae* is capable of infecting a wide range of grass species, including major cereal crops such as wheat, barley, oat, and millet, thus compromising global food security (3, 4). As one of the most important plant diseases, it is mainly controlled by planting resistant cultivars. Nevertheless, outbreaks of rice blast still occur throughout rice growing regions worldwide because of breakdown of resistance by new races of the pathogen. In addition, wheat blast caused by *M. oryzae* pathovar *Triticum* has recently spread from South America to Asia, threatening wheat production (5, 6).

Like many other foliar fungal pathogens, penetration of the cuticle and cell wall of host plants is the most critical infection step (7, 8). The infection cycle of *M. oryzae* starts with attachment of dispersing conidia to plant surface by releasing sport tip mucilage. Germ tubes emerging from attached conidia can sense surface hydrophobicity and hardness and develop highly specialized infection structures known as appressoria at the tip (1, 9). Mature dome-shaped appressoria are anchored on rice surface with appressorium mucilage (10) and heavily melanized by depositing a melanin layer between the cell wall and membrane as an impermeable barrier for turgor generation (11). Subsequently, the appressorium uses the extremely high turgor pressure to physically penetrate the cuticle and cell wall with a narrow penetration peg (7, 12, 13). As an hemibiotrophic pathogen, after the penetration of plant cell wall, the penetration peg does not penetrate plant cytoplasm membrane but differentiates into primary invasive hyphae and invasive hyphae that are enveloped by plant-derived extra-invasive hyphal membrane (EIHM) in penetrated cells (7, 14). To date, it is not clear whether there is any special structure at the contact point between the penetration peg and plant cytoplasm membrane before the differentiation of primary invasive hyphae. However, a biotrophic interfacial complex (BIC), a plant membrane-rich structure, is formed at the tip of the primary invasive hyphae for effector delivery (15, 16). Following

proliferation within the first colonized cell, the tips of invasive hyphae differentiate into
transpressoria, which subsequently form narrow infection pegs to traverse the plasmodesmata
(PD)-rich cell junctions, allowing entry into adjacent host cells (17, 18). The fungus continues this
colonization strategy until disease symptoms manifest on infected rice plants.

To suppress plant defense responses and facilitate invasive growth, *M. oryzae* deploys effectors
with various functions that can be delivered into host tissues or cells with two secretion systems
during infection (19). Apoplastic effectors, such as Bas4 and Slp1, accumulate in the extracellular
space between the fungal invasive hyphal cell wall and EIHM, outlining the entire invasive hyphae,
are secreted through the conventional endoplasmic reticulum (ER)-Golgi-dependent pathway
which is sensitive to brefeldin A (BFA) (19). Conversely, cytoplasmic effectors, including Pwl2,
AvrPiz-t, and Bas1, accumulate in the BIC before being delivered into plant cells using a Golgi-
independent pathway involving the exocyst complex and t-SNAREs, which is insensitive to BFA
(19). To date, all the known proteinaceous effectors in *M. oryzae* belong to these two categories
and localize to the extracellular space outlining invasive hyphae or the BICs and intracellular
targets inside plant cells.

In a previous study, we identified a family of small secreted proteins that may function as effectors
in *M. oryzae* (20). In this study, we found that one of them named as *PPE1* (for peg periphery-
localized effector 1, MGG_16603) is specifically expressed during plant infection and is a crucial
virulence factor. Surprisingly, Ppe1 preferentially localizes to the penetration site in a ring
conformation that is described as the penetration ring below. The penetration ring consisting of
Ppe1 proteins is formed after appressorium- and transpressorium- mediated plant cell wall
penetration and remained at the point where penetration pegs transitioning into primary invasive
hyphae. Ppe1 homologs are only present in fungal pathogens using extreme appressorium turgor
for plant penetration. Other members of the Ppe1 family also localize to the penetration ring that
is at the base of primary invasive hyphae outside plant cytoplasm membrane. Taken together, a
penetration ring consisting of secreted Ppe1 family proteins is formed between plant cell wall and
cytoplasm membrane, which may function as a novel physical structure at the interface between

the tip of penetration pegs and plant cytoplasm membrane before the differentiation of invasive

hyphae.

Results

Ppe1 is important for full virulence of *M. oryzae*

We previously identified 21 host-adapted effector genes in *M. oryzae* that may play critical roles

during rice infection based on *in silico* analyses (20). To explore the functions of these effectors,

we selected *PPE1* (MGG_16603) for further functional characterization because its high

transcription level during early infection stages as reported (20, 21). We first assayed the

expression profile of *PPE1* during infection by qRT-PCR. The result showed that its expression

increased during the early stages of plant infection (24-36 hpi) and peaked at 36 hpi (Fig. 1A).

Moreover, fluorescence signals of mCherry driven by the *PPE1* promoter indicated that its

expression was only observed in matured appressoria and during infection (Fig. 1B). These results

were consistent with the transcriptome data, suggesting that Ppe1 is functional during

appressorium penetration and plant infection. Ppe1 encodes a 116-aa protein with typical

characteristics of fungal effectors, including an N-terminal signal peptide and eight cysteine

residues (Fig. S1A). The signal peptide of Ppe1 is functional in directing the secretion of yeast

Suc2 invertase in a yeast signal trap system (Fig. S1B), suggesting that Ppe1 is a secreted protein.

To determine the biological functions of Ppe1 in *M. oryzae,* we generated a *PPE1* gene deletion

mutant in the wild-type strain 70-15 (Fig. S1C). Deletion of *PPE1* had no obvious effect on the

fungal growth, conidiation, and appressorium formation (Fig. S1D-G). However, in spray infection

assays with rice leaves, the Δ*ppe1* mutant was significantly reduced in the number of disease

lesions compared to the wild type (Fig. 1C). In wound infection assays, the Δ*ppe1* mutant also was

significantly reduced in virulence and formed smaller lesions than the wild type (Fig. 1D). To

confirm the role of *PPE1* in virulence, we generated a complemented strain Δ*ppe1*-com that had a

copy of the *PPE1* gene ectopically integrated into its genome. In spray or wound infection assays,

Δ*ppe1*-com was similar with the wild type in virulence (Fig. 1C-D), indicating a complete recovery

in phenotypes. We then conducted rice leaf sheath penetration assays to observe the infection

processes. At 32 hpi, over 91% of the appressoria formed by the wild type and the complemented

strain had penetrated and developed invasive hyphae in rice cells. Invasive hyphae in most of the

penetrated cells (approximately 60%) had more than two branches, and approximately 10% of
them had spread into the neighboring cells (Fig. 1E). In contrast, approximately 25% of the
appressoria formed by the Δppe1 mutant had no penetration peg. Although 53.5% of these mutant
appressoria successfully penetrated into host cells, only 23% of Δppe1 invasive hyphae had more
than two branches and less than 4% had spread into adjacent cells (Fig. 1E). These results indicate
that the Δppe1 mutant has defects in appressorium penetration, invasive hyphal growth and
subsequent spread of invasive hyphae into neighboring plant cells.

Ppe1 forms a ring at the base of appressoria during host penetration
To determine the localization of Ppe1 during infection, we expressed a full-length Ppe1-mCherry
fusion protein under the control of *PPE1* promoter in both the wild type (70-15) and the Δppe1
mutant. Rice leaf sheaths were then infected with these strains and the localization of Ppe1 was
analyzed by confocal microscopy. In these transformants, Ppe1-mCherry signals were observed as
a ring underneath the appressoria (likely at the site of the emergence of penetration pegs) at 22 hpi
(Fig. 2A, Video S1). The Ppe1-mCherry ring persisted after appressorial penetration and the ring-
shaped fluorescence signal of the Ppe1-mCherry was still observed even after the formation of
primary and secondary invasive hyphae (Fig. 2B). We further measured the size of this ring and
found that the average diameter of the inner cycle of the Ppe1-mCherry ring was 1.53 μm. The
width of the Ppe1-mCherry ring varied from 1.79 to 4.78 μm, which is smaller than the width of
appressoria (Fig. 2C) and known septin ring (5.9 μm) at the appressorium pore (13). Furthermore,
the Ppe1-mCherry fusion protein could be detected by western blot analysis with proteins isolated
from infected rice leave (Fig. S2). Because the Ppe1 ring always appears at the point of emergence
of the penetration peg, we named this ring structure as penetration ring.

To confirm the localization of Ppe1, we also generated Ppe1-GFP transformants. Like Ppe1-
mCherry, Ppe1-GFP also formed a ring at the penetration site underneath the appressorium (Fig.
2D). These results indicate that, irrespective of being fused with mCherry or GFP, Ppe1 localizes
to the penetration ring. To avoid possible overexpression or unexpected effects due to ectopic
integration of the fusion constructs, we generated a transformant from 70-15 strain in which
mCherry was inserted right before the TAA stop codon of the endogenous *PPE1* gene. In the

resulting transformant expressing the in situ *PPE1*-mCherry fusion construct, the penetration ring
was also observed (Fig. 2E). These results further verify the localization of Ppe1 to the penetration
ring during appressorial penetration.
175
Penetration ring is also formed during the penetration of neighboring cells by transpressoria
In *M. oryzae*, appressoria-like structures called transpressoria are formed by invasive hyphae on
the wall boundary of the invaded cell in order to penetrate into neighboring cells (17, 18).
Therefore, we observed localization of Ppe1 during transpressorium-mediated penetration of plant
cells. In the *PPE1*-mCherry transformant, like during appressorial penetration, the penetration ring
was observed during transpressoria-mediated penetration of neighboring cells (Fig. 3A). Similarly,
we observed the penetration ring in the neighboring cells penetrated by invasive hyphae of the
PPE1-GFP transformant (Fig. 3B). The Ppe1-mCherry or Ppe1-GFP ring (Fig. 3A and B – zoomed
view) appeared to be at the transpressorial penetration point. These results indicate that the
penetration ring is also formed during invasive hyphal cell-to-cell penetration. Furthermore,
similar localization of Ppe1 to the penetration ring was observed in barley epidermal cells (Fig.
3C). Taken together, we conclude that Ppe1 localizes to the penetration ring during appressorium-
mediated and transpressorium-mediated plant cell penetrations in *M. oryzae.*
189
Penetration ring is formed on the periphery of the penetration peg
In *M. oryzae*, appressorium-mediated penetration is a septin-dependent process in which septins
aggregate and generate a hetero-oligomeric 5.9 μm ring structure at the appressorial pore to re-
organize the F-actin cytoskeleton network to promote the emergence of penetration pegs (13).
Following the penetration peg formation, septin ring undergoes further constriction into a smaller
ring (22). To clarify the precise site of the penetration ring, we expressed Ppe1-mCherry (under
the control of the *PPE1* promoter) in a strain expressing Sep3-GFP (Guy11 background) and then
observed their spatiotemporal dynamics during rice leaf sheath infection. At 11 hpi, Sep3-GFP big
ring was formed at the appressorium pore but the Ppe1-mCherry ring was not observed (Fig. S3A,
Video S2). The Ppe1-mCherry ring signal was observed at 16 hpi, after the constriction of Sep3-
GFP ring and the former was completely surrounding the latter (Fig. S3B). Subsequently (at 19
hpi), the Ppe1-mCherry ring (still surrounding the Sep3-GFP ring) became constricted (Fig. 4A,

Fig. S3C, Video S3). However, on impenetrable hydrophobic coverslips, the Sep3-GFP ring was

formed normally at the appressorium pore, but the Ppe1-mCherry ring and constriction of Sep3-

GFP were not observed (Fig. S4). These results suggest that Ppe1-mCherry ring is formed after

the formation of penetration pegs, outside the constricted Sep3-GFP ring.

Penetration pegs are rich in microfilaments, filasomes and microtubules (23), therefore, it can be

labeled by Lifeact-GFP. Hence, we co-expressed Ppe1-mCherry with Lifeact-GFP in 70-15 and

observed their positioning in penetration assays with rice leaf sheaths. The Ppe1-mCherry ring was

also observed to surround the Lifeact-GFP signal (Fig. 4B, Video S4), indicating that the

penetration ring is localized at the periphery of penetration pegs during the initial infection stage.

To confirm this observation, we co-expressed the Ppe1-mCherry with cytosolic GFP in 70-15.

Using a three-dimensional (3D) volume rendering image constructed by confocal optical z-series

images, we found that the Ppe1-mCherry ring was situated around the narrow site between the

appressorium and invasive hyphae labeled with GFP (Fig. 4C).

A previous study showed that callose deposition is dynamic during *M. oryzae* infection and has

three different forms: callose-rich papillae underneath appressorium penetration sites, callose

collars formed at cell wall crossing sites around the invasive hyphae and callose deposition at

plasmodesmata pit fields (24). To investigate whether the penetration ring locates outside the peg,

we stained callose deposits with aniline blue to examine the possible positioning of the callose ring

underneath the appressorial penetration site and Ppe1-mCherry. We discovered that the Ppe1-

mCherry ring was surrounded by the callose ring, exhibiting partial colocalization and close

attachment (Fig. 4D, Video S5). Taken together, we conclude that the penetration ring locates at

the periphery of penetration peg during the initial appressorium-mediated host penetration.

Formation of the penetration ring at the base of penetration pegs after transpressorium

penetration

To determine the precise localization of the penetration ring during the transpressoria-mediated

penetration, we expressed Ppe1-mCherry (under the control of the *PPE1* promoter) in a Sep5-GFP

strain and then observed their spatiotemporal dynamics during rice leaf sheath infection. The

results showed that Sep5-GFP first displayed a ring structure at the swollen part of the invasive

hyphal tip (Fig. S5A), similar to the septin big ring at the appressorium pore. It then exhibited a

collar structure at the constricted infection peg site (Fig. S5B-C), which is consistent with a

previous report by Sakulkoo (25). Meanwhile, the Ppe1-mCherry ring or collar-like ring was

formed outside the Sep5-GFP collar (Fig. S5B-C), indicating that the penetration ring formed

during transpressoria-mediated cell-to-cell penetration is positioned outside the penetration peg.

Taken together, we conclude that the penetration ring of *M. oryzae* is localized at the periphery of

penetration pegs both during appressoria- and transpressoria-mediated host penetrations.

Penetration ring formation requires the initiation of penetration peg formation

To further characterize the relationship between the penetration ring and penetration peg, we

observed the expression and localization of Ppe1-mCherry in Δ*mst12,* Δ*nox2* and Δ*nox1* mutant

strains that are defective in plant penetration (26, 27). In the Δ*mst12* mutant that is defective in

both penetration initiation and peg formation, the Ppe1-mCherry fluorescence signal could not be

detected. In the Δ*nox2* mutant that is unable to initiate penetration peg formation, the Ppe1-

mCherry ring could not be formed but displayed a disorganized mass underneath the appressorium.

Interestingly, in the Δ*nox1* mutant that forms normal penetration pegs but is defective in invasive

hyphal elongation, the Ppe1-mCherry ring was formed during rice leaf sheath infection (Fig. 5A),

indicating that penetration peg formation is a prerequisite for the emergence of the penetration ring.

To confirm this observation, the localization of Ppe1-mCherry fusion proteins was observed after

inducing appressorium formation on hydrophobic coverslips and cellophane membranes.

Although melanized appressoria were formed, we failed to observed the Ppe1-mCherry ring on

the impenetrable coverslips (Fig. 5B). However, the penetration ring is formed after the penetration

of cellophane membranes (Fig. 5C). These results indicate that initiation of penetration peg

formation is necessary for the formation of the penetration ring.

Other Ppe proteins also localize to the penetration ring

To determine whether other secreted proteins homologous to Ppe1 also localize to the penetration

ring, we used the amino acid sequence of Ppe1 to carry out BLASTp search at the NCBI database

to uncover the homolog proteins of Ppe1 and further analyzed their phylogenetic relationships.

The Ppe1 homologs only exist in *Magnaporthe* and *Colletotrichum* species, such as the most

common cereal phytopathogens *M. oryzae* and *C. graminicola* (Fig. 6A). Remarkably, these fungi

are known to form melanized appressoria and deploy extremely high turgor pressure for plant

penetration (8). To understand whether other members of the Ppe1 family (Ppe2, Ppe3, Ppe3, Ppe4,

Ppe5, and Ppe6) from *M. oryzae* have similar localization as Ppe1, we tagged each of them with

mCherry and expressed them individually in the 70-15 background under the control of their native

promoters. Examination of their localizations during infection revealed that Ppe2-mCherry, Ppe3-

mCherry and Ppe5-mCherry also exhibit a ring structure during appressorium-mediated host

penetration, while the fluorescence signals of Ppe4-mCherry and Ppe6-mCherry were hardly

observed, possibly due to their low expression (Fig. 6B). Moreover, when MoT-Ppe1 from *M.*

oryzae pathovar *Triticum* was expressed in *M. oryzae* under the control of its native promoter, the

characteristic penetration ring was also observed (Fig. 6B). All these results suggest that the

secreted Ppe proteins accumulate at the penetration ring.

As controls, we also examined the localizations of Bas4, Slp1, Pwl2, and AvrPiz-t. Under the same

experimental conditions, Bas4-GFP and Slp1-GFP were observed to uniformly accumulate within

the extra-invasive hyphal membrane (EIHM), outlining the entire invasive hyphae, while Pwl2-

mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC) (Fig. 6C),

which is consistent with previous reports (16, 28, 29). EIHM and BIC are specifically associated

with invasive hyphae during biotrophic growth. In contrast, Ppe1 localized to the penetration ring

at the periphery of penetration pegs (Fig. 6C). Therefore, unlike subcellular localizations of other

effectors to the extracellular space outlining invasive hyphae, BICs, or intracellular targets in plant

cells, Ppe1 and its homologs localize to the penetration ring that may function as a novel physical

structure at the interface between the tip of penetration pegs and plant cytoplasm membrane before

the differentiation of invasive hyphae.

Discussion

As the model plant pathogenic fungus, appressorium formation and appressorium-mediated

penetration have been well-characterized in *M. oryzae* (18). The fungus uses extremely high turgor

pressure generated in appressoria cemented to the surface with appressorium mucilage to penetrate

underlying plant cells (12). As an hemibiotrophic pathogen, after piercing plant cell wall, the
penetration peg does not penetrate the cytoplasm membrane but differentiates into invasive hyphae
enveloped in a layer of EIHM (16). In this study, we showed that a penetration ring of Ppe1
proteins is formed by the penetration peg on the surface of the plant cytoplasm membrane before
invading and differentiating into primary invasive hyphae (Fig. 2 and Fig. 7). Other Ppe proteins
including Ppe2, Ppe3, and Ppe5 proteins also localize to the penetration ring (Fig. 6B). Because
the penetration ring is persistent at the point penetration pegs transitioning into invasive hyphae
outside the cytoplasm membrane even after the penetrated cells are filled with invasive hyphae, it
may function as a novel physical structure for anchoring penetration pegs on the surface of plant
cytoplasm membrane after cell wall penetration (Fig. 7). It is also possible that this penetration
ring functions as a collar or landmark that is associated with the differentiation of penetration pegs
(on the surface of cytoplasm membrane) into primary invasive hyphae enveloped in the EIHM
cytoplasm membrane (Fig. 7). Although this transition is not well characterized, it likely involves
changes in cell polarity and cytoskeleton as well as cell wall modifications. Ppe1 and its homologs
may be able to assemble into a protein complex in the penetration ring, which likely involves other
proteins or components as facilitating factors. In *U. maydis*, the Stp secreted effectors are known
to form a protein complex during plant infection (30).

The penetration ring is also formed after transpressoria-mediated penetration of neighboring plant
cells (Fig. 7). Similarly, Ppe1 and its homologs form a ring after the penetration of cell wall but
before differentiating into invasive hyphae enveloped with the EIHM. The penetration ring formed
by transpressoria-mediated penetration is also persistent at the penetration site on the surface of
plant cytoplasm membrane in penetrated cells (Fig. 3 and Fig. S5), suggesting that Ppe proteins
are assembled into a complex that is not diffusible. In fact, similar penetration ring is observed
after penetration of artificial membrane (Fig. 5C), which has no barrier to restrict the diffusion of
secreted proteins, further indicating that the penetration ring of Ppe1 and other Ppe proteins is
likely a structural feature or component for signaling the transition of penetration pegs to invasive
hyphae. Although appressoria and transpressoria are different, the processes after the penetration
of cell wall are the same in appressorium- and transpressorium-mediated plant penetration.

Therefore, formation of the penetration ring in both appressorium- and transpressorium-mediated

penetration is another evidence for it to be a structural element.

M. oryzae is known to secrete a repertoire of effectors that can be categorized into cytoplasmic

and intracellular effectors (14). Ppe1 and its homologs are putative effectors that are described as

HAG (host adaptation genes) effectors in an earlier study (20). Although it remains possible that

Ppe1 and its homologs function as effectors during plant infection. However, in comparison with

the localizations of known effectors in *M. oryzae*, we found that the localization of Ppe1 to the

penetration ring is unique and Ppe1 homologs have the same localization pattern (Fig. 6). Because

of the persistent localization of Ppe1 and its homologs to the penetration ring, we think the

penetration ring is likely a physical structure as described above. If Ppe1 and its homologs are

effector proteins being secreted at this specific location, fluorescent signals of Ppe1 and its

homologs will diffuse and disappear or dimmish when effectors being translocated into plant cells.

Furthermore, it is unlikely that secreted proteins localized to the penetration ring that has limited

contact with the plant cytoplasm membrane will function as effector proteins to interact with plant

targets. Nevertheless, although unlikely, it remains possible that the penetration ring discovered in

this study is a secretion site for effectors involved in the transition of penetration pegs to invasive

hyphae. After the penetration of plant cell wall, the penetration ring may not only function as a

physical structure but also serve as an initial effector secretion site for the release of specific

effectors to overcome plant immunity in early infection stages.

Taken together, in this study we observed the formation of the penetration ring that appears to be

a structural element for signaling or triggering the differentiation of penetration pegs after the

penetration of plant cell wall into invasive hyphae enveloped by the EIHM in *M. oryzae*. Based on

the limited distribution of Ppe1 homologs, it is likely that only fungal pathogens using high

appressorium turgor for plant penetration, such as *M. oryzae* and *Colletotrichum* species,

developed the penetration ring. Therefore, it will be important to further characterize the functional

relationship between the penetration ring and appressorium turgor as well as the transition of

penetration pegs driven by turgor pressure into primary invasive hyphae (lower turgor pressure).

Materials and Methods

Strains and growth conditions

The *M. oryzae* strains 70-15 and Guy11 were used as the wild types in this study; all other fungal strains generated from the wild types are listed in Table S1 in the supplemental materials. All the fungal strains were grown on complete medium CM (CM: 10 g D-glucose, 2 g peptone, 1 g yeast extract, 1 g casamino acids, 50 mL 20 × nitrate salts (120 g/L NaNO$_3$, 10.4 g/L KCl, 30 g/L KH$_2$PO$_4$, and 10.4 g/L MgSO$_4$), 1 mL trace elements (0.1 g/L biotin, 0.1 g/L pyridoxine, 0.1 g/L thiamine, 0.1 g/L riboflavin, 0.1 g/L p-aminobenzonic acid and 0.1 g/L nicotinic acid), adjusted to pH 6.5 with NaOH, 15 g agar and distilled water added to 1 L) (31). Agar cultures were incubated at 26 °C with a 12-h light/12-h dark photoperiod for mycelial growth and conidiation. Liquid CM medium was used to harvest mycelia for extraction of genomic DNA and protoplast preparation. The bacterial strains *Escherichia coli* DH5α and *Agrobacterium tumefaciens* AGL1 were grown in Luria-Bertani (LB) medium at 37 °C and 28 °C, respectively.

Plant and growth conditions

Rice cultivar CO-39 (*Oryza sativa* cv. CO-39) was grown in a greenhouse at 26 °C and 70% relative humidity with a 14-h light/10-h dark photoperiod. Barley (*Hordeum vulgare* cv. Golden Promise) was grown at 23 °C with a 12-h light /12-h dark photoperiod.

Bioinformatic and phylogenetic analyses of Ppe1 homolog proteins

The Ppe1 was previously sequenced and the amino acid sequence established from our lab (32). Ppe1 signal peptide was predicted using SignalP 5.0 software. BLASTp was used to find Ppe1 homolog proteins at the NCBI database (https://www.ncbi.nlm.nih.gov/) and the representative homolog protein sequences were downloaded from the database. Phylogenetic tree was generated using MEGA7.0 software by neighbor-joining method.

Construction of recombinant DNA and *M. oryzae* transformation

The plasmids used in the study were generated by infusion cloning method based on homologous recombination using ClonExpress II one-step cloning kit or ClonExpress MultiS kit (Cat. No. C112-01 or C113-01, Vazyme, Nanjing, China). The primers used (Table S2 in the supplemental material) were specially designed to contain 18-22-bp overlap with adjoining sequences to allow

the fragments assemble by homologous recombination. To observe the expression pattern and
localization of *PPE1*, the *PPE1* promoter (1485 bp) fragment amplified from the genomic DNA
of the wild type (70-15) was seamlessly cloned into the *Kpn*I/*Hin*dIII-digested pKNT-
mCherry/GFP fusion vectors (33) using the ClonExpress II one-step cloning kit (Cat. No. C112-
01, Vazyme, Nanjing, China). For Ppe1-mCherry/GFP plasmid, the entire Ppe1 coding sequence
with its native promoter but without stop codon was amplified and cloned into the *Kpn*I/*Hin*dIII-
digested vector pKNT-mCherry/GFP. The pKNT-AvrPizt-GFP and other plasmids used were was
constructed using the same method. The constructed plasmids were transformed into the fungal
protoplasts via PEG-mediated transformation (34) and further verified by PCR screening and
confocal microscopic examination of fluorescence signals. For situ fusion of Ppe1 with mCherry,
the mCherry was inserted just before the Ppe1 stop codon and the construct was transformed by
Agrobacterium tumefaciens-mediated (ATMT) transformation for homologous recombination
(34).
393
Assays for the secretory function of the SP of Ppe1
To confirm the secretory function of the predicted signal peptide (SP), yeast signal trap system
was performed as described previously (35). The predicted SP nucleotide sequences of Ppe1 (60
bp) and Pwl2 (63 bp) were respectively cloned into pSUC2 vector that carries a truncated invertase
gene *SUC2*. The generated pSUC2-SP constructs and the empty pSUC2 vector were transformed
into the yeast *suc2* mutant YTK12, respectively, and then their growth conditions analyzed on
CMD-W (minus Trp) plates (0.67% yeast nitrogen base without amino acids, 0.075% tryptophan
dropout supplement, 2% sucrose, 0.1% glucose, and 2% agar) and YPRAA medium plate (1%
yeast extract, 2% peptone, 2% raffinose, and 2 μg/mL antimicyn A) containing raffinose and
lacking glucose. Transformants of YTK12 carrying the empty pSUC vector or pSUC2-Pwl2SP
were used as negative and positive controls, respectively. Reduction of 2,3,5-triphenyltetrazolium
chloride (TTC) to insoluble red-colored 1,3,5-triphenylformazan (TPF) was used as an indicator
of the invertase enzymatic activity (36). Each of the above YTK12 transformants strain was
inoculated in 5 mL of CMD-W liquid medium and incubated for 24 h at 30 ℃. The cells were then
collected, washed and resuspended in 0.1% TTC in new centrifuge tubes and further incubated at

35 °C for 35 min. The color change was observed after 10 min of incubation at room temperature

to detect the signal peptide secretion function.

Targeted gene deletion and complementation

Deletions of *NOX1*, *NOX2* and *MST12* genes were performed using Split-marker recombination

method (37). The open reading frames (ORFs) of the various genes were replaced with a

hygromycin phosphotransferase (*HPH*) resistance cassette, respectively. Briefly, about 1-kb

upstream and downstream sequences of the targeted genes were amplified from the genomic DNA

of the wild type (70-15) using specially designed primer pairs AF+AR and BF+BR, respectively.

At the same time, the upstream and downstream sequences of the *HPH* with overlapping segments

were amplified from pCB1003 plasmid using the primer pairs HYG/F + HY/R and YG/F + HYG/R,

respectively. The corresponding upstream and downstream fragments were respectively ligated by

splicing for overlap extension (SOE) PCR using the primer pairs AF+HY/R and YG/F+BR. The

PCR products were purified and transformed into the Ppe1-mCherry strain protoplasts via a PEG-

mediated transformation (34). Putative transformants were selected in the presence of hygromycin

B and then confirmed by PCR using the primers OF+OR and UA+H853. PCR confirmations were

repeated three times for consistency. All the primer sequences used are listed in Table S2 of the

supplemental material.

For *PPE1* gene deletion, *HPH* sequence was inserted between the upstream and downstream

sequences of the *PPE1* gene, replacing the ORF of the *PPE1* gene, in an ATMT1 plasmid using a

ClonExpress MultiS One Step Cloning Kit (Cat. no. C113-01, Vazyme, Nanjing, China). The

constructed plasmid was introduced into *Agrobacterium tumefaciens* AGL1 using freeze-thaw

method (38). *Agrobacterium tumefaciens*-mediated transformation (ATMT) of the construct into

the wild-type spores was then carried out. The transformants were selected on hygromycin B plates

and confirmed by PCR. The mutants were further confirmed by Southern blot assay using DIG-

High Prime DNA labeling and detection starter kit I (Cat. no. 11745832910, Roche, Mannheim,

Germany). For complementation, the constructed plasmid pKNT-Ppe1-mCherry was transformed

into the Δ*ppe1* mutant protoplast and further verified by PCR screening and confocal microscopic

examination for fluorescence signal detection.

Live cell imaging

To monitor the expression and subcellular localization of Ppe1 during plant infection, rice leaf sheath inoculation and live cell imaging of fluorescently labeled proteins were performed as described (15) with slight modifications. In brief, 5-7 cm long leaf sheath segments were cut from 4-week-old CO-39 rice seedlings, and the whole hollow spaces of the sheaths were filled with conidia suspensions (5×10^4 conidia/mL) using a 200 μL pipette. The inoculated sheaths were placed horizontally (fixed with wet filter paper fragments) on an inverted 96-well PCR plate covered by one layer of wet filter paper and incubated in a sealed box whose inner bottom was lined with two layers of wet filter paper to keep the environment humid at 26 °C. The epidermal parts of the inoculated sheaths were trimmed into thin and transparent samples and immediately imaged using an inverted Nikon A1 laser scanning confocal microscope (Nikon, Tokyo, Japan) at the indicated time points post inoculation. The 60×/1.42 numerical aperture (NA) oil immersion objective lens was used for the imaging. The fluorescence and differential interference contrast (DIC) images were collected by confocal optical z-series acquired at 0.28 μm intervals spanning the entire fluorescence signal depth between the top plane of the appressorium formed and the bottom plane of the rice leaf sheath cross sections. The fluorescence excitation/emission wavelengths were 488 nm/500-550 nm for GFP, 561 nm/570-620 nm for mCherry and 405 nm/425-475 nm for callose staining. The images were processed using NIS-Elements AR analysis software (Nikon, Tokyo, Japan). Unless otherwise stated in the figure legend, micrograph fluorescence represents maximum intensity projection of z-series (with a 0.28 μm intervals) after removing extraneous z-planes that do not contain relevant information for a particular fluorescence. Three-dimensional (3D) volume rendering images and videos were generated from z-series in maximum intensity projection.

To observe the subcellular localizations of Ppe1 in conidia incubated on artificial hydrophobic surfaces and barley leaves, conidia suspensions (5×10^4 conidia/mL) were dropped on the surfaces of hydrophobic coverslips or on the back sides of barley leaves (5-day old plants) and incubated under humid and dark conditions at a 26 °C. For observation, the hydrophobic coverslips containing the appressoria were inverted, placed on 24mm×50 mm microscope glass slides and

then observed under a scanning confocal microscope. Epidermal parts were removed from the

infected barley leaves for observation. For penetration assay, cellophane was cut into round sheets

of 4 cm in diameter and sterilized by autoclaving at 115 ℃ for 15 min. The sterile cellophane

sheets were placed flat on CM plates and then the conidia suspensions ($5×10^4$ conidia/mL) were

dropped on the surface of the cellophane and incubated at 26 ℃. Small pieces of the cellophane

containing mycelia were cut for observation under a confocal microscope at 36-48 hpi.

Callose staining

For visualization of deposited callose, aniline blue (Cat. no. 28631-66-5, Sangon, Shanghai, China)

staining was adopted (25). Rice leaf sheaths were inoculated with conidia suspensions ($3×10^4$

conidia/mL). The inoculated sheaths were trimmed to thin and transparent samples at the indicated

time points and stained with aniline blue (0.1mg/mL) solution prepared in 150 mM KH_2PO_4 (pH

9.5) for up to 30 min. The stained samples were imaged by using a laser scanning confocal

microscope (Nikon, Japan).

Phenotypic analysis

For vegetative growth and conidiation assays, small mycelial blocks from each strain were cut

from the edges of 4-day-old cultures and placed at the center of freshly prepared CM plates (90

mm) and incubated for ten days at 26 ℃. The colony diameters of the cultures were measured

using vernier calipers and then photographs of the cultures were taken. The cultures were then

washed with 5 mL sterilized double-distilled water, filtered through three layers of lens paper and

the conidia counted using a hemocytometer under a light microscope. For conidia germination and

appressoria formation, spore suspensions ($1×10^5$ conidia/mL) were incubated on hydrophobic

coverslips (Cat. no. 12-545-101P, Fisherbrand™ Cover Glasses, USA) under humid and dark

conditions at 26 ℃ for 24 h. The conidia were then observed under a light microscope for

germination and formation of appressoria.

To investigate the pathogenicity of each strain, conidia spray, punch and leaf sheath inoculations

were performed. Conidia suspensions ($5×10^4$ conidia/mL containing 0.02% Tween-20) were used

to spray the leaf surfaces of 3-week-old CO-39 rice seedlings. The inoculated seedlings were

incubated in a growth chamber at 25 °C with 90% humidity. The seedlings were first kept in the dark for 24 h, and then subjected to 12 h light / 12 h dark photoperiod for 7 days, after which photographs were taken and the number of disease lesions on the leaf surfaces counted. Punch inoculation assay was carried out according to a method previously described (29). Briefly, 5-week-old CO-39 rice leaves were slightly wounded using mouse ear punch and 10 μL of conidia suspensions (5×10^5 conidia/mL containing 0.02% Tween-20) from the various strains were respectively dropped at the wounded regions. Then, both sides of each inoculated spot were sealed with scotch tape to sustain the spore suspension. The inoculated plants were kept in a growth chamber under the same conditions described above for 10 days. The disease lesions formed were scanned, and later analyzed using Adobe Photoshop software to measure the lesion areas. For leaf sheaths inoculation, conidia suspensions (1×10^5 conidia/mL) from the various strains were respectively injected into 4-5 weeks old CO-39 rice leaf sheaths. The epidermal parts of the infected leaves were trimmed to thin and transparent samples for observation under a light microscope at 32 hpi. At least 100 infected sites were observed using a Nikon A1 laser confocal microscope (Nikon, Tokyo, Japan).

qRT-PCR analysis

To analyze the expression level of *PPE1* during *M. oryzae* infection, rice leaf sheaths were cut from 4- to 5-week-old CO-39 rice seedings and inoculated with conidia suspensions (2×10^5 conidia/mL) from the wild type (70-15). The epidermal parts of the inoculated sheaths were cut and collected at 12-, 36- and 48 hours post inoculation (hpi). At least 15 segments of the inoculated rice sheath samples were collected at each time point. Total RNAs were extracted from the leaf sheaths or rice seeding leaves using an Eastep Super Total RNA Extraction Kit (Cat. no. LS1040, Promega, Shanghai, China). Reverse transcription was used to generate cDNA using a reverse transcription kit (Cat. no. RR037B, Takara, Dalian, China). Quantitative PCR was finally conducted using a SYBR kit (Cat. no. RR067A, Takara, Dalian, China) using specific primers (Table S2). Fugal *β-TUBULIN* gene used as endogenous reference genes, and the expression levels were calculated using $2^{-\Delta\Delta Ct}$ method as previously reported (39). All qRT-PCR assays were repeated three times for each sample.

Statistical analyses

Statistical analyses were performed using GraphPad Prism7 (GraphPad Software, Inc., San Diego, CA). Statistically significant differences were determined by one-way ANOVA with two-sided least significant difference. **** indicated significant difference at $p < 0.0001$，ns indicated no significant difference.

Data Availability

The authors declare that all the data supporting the findings of this study are available within the paper and SI Appendix. Raw data and details of experimental procedures are available upon request from the corresponding authors.

Acknowledgments

We thank Prof. Nicholas J. Talbot and Prof. Barbara Valent for providing us with *Δsec5* mutant and pBV591 plasmid respectively. We also thank Prof. Min Guo for providing us with Sep3/5-GFP strains and Δsep3/4/5 mutants. We also appreciate Profs. Guodong Lu, Jie Zhou, Huawei Zheng, Wei Tang, Ya Li, Xiaofeng Chen, and Huakun Zheng, as well as Drs Yijuan Hang, Meilian Chen and Yonghe Hong for their helpful suggestions. Thanks to Jin Zhang, Chunling Li and Yuping Fan's help. This research was supported by the National Key Research and Development Program of China (2023YFD1400200), the National Natural Science Foundation of China (32122071, 31601596), and the Natural Science Foundation of Fujian Province (2021J06015).

References

1. Talbot NJ (2003) On the trail of a cereal killer: Exploring the biology of Magnaporthe grisea. *Annual review of microbiology* 57:177-202.
2. Dean R*, et al.* (2012) The Top 10 fungal pathogens in molecular plant pathology. *Molecular plant pathology* 13(4):414-430.
3. Couch BC*, et al.* (2005) Origins of host-specific populations of the blast pathogen Magnaporthe oryzae in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice. *Genetics* 170(2):613-630.
4. Langner T, Białas A, & Kamoun S (2018) The Blast Fungus Decoded: Genomes in Flux. *mBio* 9(2).
5. Ceresini PC*, et al.* (2018) Wheat Blast: Past, Present, and Future. *Annual review of phytopathology* 56:427-456.

6.   Malaker PK*, et al.* (2016) First Report of Wheat Blast Caused by Magnaporthe oryzae Pathotype
triticum in Bangladesh. 100(11):2330-2330.
7.   Howard RJ & Valent B (1996) Breaking and entering: host penetration by the fungal rice blast
pathogen Magnaporthe grisea. *Annual review of microbiology* 50:491-512.
8.   Talbot NJ (2019) Appressoria. *Current biology : CB* 29(5):R144-r146.
9.   Hamer JE, Howard RJ, Chumley FG, & Valent B (1988) A mechanism for surface attachment in
spores of a plant pathogenic fungus. *Science (New York, N.Y.)* 239(4837):288-290.
10.   Rocha RO, Elowsky C, Pham NTT, & Wilson RA (2020) Spermine-mediated tight sealing of the
Magnaporthe oryzae appressorial pore-rice leaf surface interface. *Nature microbiology*
5(12):1472-1480.
11.   Chumley FG & Valent BJMP-mI (1990) Genetic analysis of melanin-deficient, nonpathogenic
mutants of Magnaporthe grisea. 3:135-143.
12.   de Jong JC, McCormack BJ, Smirnoff N, & Talbot NJ (1997) Glycerol generates turgor in rice blast.
Nature 389(6648):244-244.
13.   Dagdas YF*, et al.* (2012) Septin-mediated plant cell invasion by the rice blast fungus,
Magnaporthe oryzae. *Science (New York, N.Y.)* 336(6088):1590-1595.
14.   Fernandez J & Orth K (2018) Rise of a Cereal Killer: The Biology of Magnaporthe oryzae
Biotrophic Growth. *Trends in microbiology* 26(7):582-597.
15.   Kankanala P, Czymmek K, & Valent B (2007) Roles for rice membrane dynamics and
plasmodesmata during biotrophic invasion by the blast fungus. *The Plant cell* 19(2):706-724.
16.   Khang CH*, et al.* (2010) Translocation of Magnaporthe oryzae effectors into rice cells and their
subsequent cell-to-cell movement. *The Plant cell* 22(4):1388-1403.
17.   Cruz-Mireles N, Eseola AB, Osés-Ruiz M, Ryder LS, & Talbot NJ (2021) From appressorium to
transpressorium-Defining the morphogenetic basis of host cell invasion by the rice blast fungus.
PLoS pathogens 17(7):e1009779.
18.   Ryder LS*, et al.* (2022) The appressorium at a glance. *Journal of cell science* 135(14).
19.   Giraldo MC*, et al.* (2013) Two distinct secretion systems facilitate tissue invasion by the rice blast
fungus Magnaporthe oryzae. *Nature communications* 4:1996.
20.   Ebbole DJ*, et al.* (2021) Evolution and Regulation of a Large Effector Family of Pyricularia oryzae.
Molecular plant-microbe interactions : MPMI 34(3):255-269.
21.   Yan X*, et al.* (2023) The transcriptional landscape of plant infection by the rice blast fungus
Magnaporthe oryzae reveals distinct families of temporally co-regulated and structurally
conserved effectors. *The Plant cell* 35(5):1360-1385.
22.   Eisermann I & Talbot NJ (2024) Septin-dependent invasive growth by the rice blast fungus
Magnaporthe oryzae. *Journal of Plant Diseases and Protection*.
23.   Bourett TM & Howard RJ (1992) Actin in penetration pegs of the fungal rice blast
pathogen,Magnaporthe grisea. *Protoplasma* 168(1-2):20-26.
24.   Osés-Ruiz M*, et al.* (2021) Appressorium-mediated plant infection by Magnaporthe oryzae is
regulated by a Pmk1-dependent hierarchical transcriptional network. *Nature microbiology*
6(11):1383-1397.
25.   Sakulkoo W*, et al.* (2018) A single fungal MAP kinase controls plant cell-to-cell invasion by the
rice blast fungus. *Science (New York, N.Y.)* 359(6382):1399-1403.
26.   Park G, Xue C, Zheng L, Lam S, & Xu JR (2002) MST12 regulates infectious growth but not
appressorium formation in the rice blast fungus Magnaporthe grisea. *Molecular plant-microbe
interactions : MPMI* 15(3):183-192.

27. Ryder LS*, et al.* (2013) NADPH oxidases regulate septin-mediated cytoskeletal remodeling during
plant infection by the rice blast fungus. *Proceedings of the National Academy of Sciences of the*
United States of America 110(8):3179-3184.
28. Mentlak TA*, et al.* (2012) Effector-mediated suppression of chitin-triggered immunity by
magnaporthe oryzae is necessary for rice blast disease. *The Plant cell* 24(1):322-335.
29. Park CH*, et al.* (2012) The Magnaporthe oryzae effector AvrPiz-t targets the RING E3 ubiquitin
ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *The*
Plant cell 24(11):4748-4762.
30. Ludwig N*, et al.* (2021) A cell surface-exposed protein complex with an essential virulence
function in Ustilago maydis. *Nature microbiology* 6(6):722-730.
31. Talbot NJ, Ebbole DJ, & Hamer JE (1993) Identification and characterization of MPG1, a gene
involved in pathogenicity from the rice blast fungus Magnaporthe grisea. *The Plant cell*
5(11):1575-1590.
32. Zhong Z*, et al.* (2018) Population genomic analysis of the rice blast fungus reveals specific events
associated with expansion of three main clades. *The ISME journal* 12(8):1867-1878.
33. Chen X*, et al.* (2021) Trehalose Phosphate Synthase Complex-Mediated Regulation of Trehalose
6-Phosphate Homeostasis Is Critical for Development and Pathogenesis in Magnaporthe oryzae.
mSystems 6(5):e0046221.
34. Khang CH, Park SY, Rho HS, Lee YH, & Kang S (2006) Filamentous Fungi (Magnaporthe grisea and
Fusarium oxysporum). *Methods in molecular biology (Clifton, N.J.)* 344:403-420.
35. Jiang C*, et al.* (2020) An orphan protein of Fusarium graminearum modulates host immunity by
mediating proteasomal degradation of TaSnRK1α. *Nature communications* 11(1):4382.
36. Oh SK*, et al.* (2009) In planta expression screens of Phytophthora infestans RXLR effectors reveal
diverse phenotypes, including activation of the Solanum bulbocastanum disease resistance
protein Rpi-blb2. *The Plant cell* 21(9):2928-2947.
37. Kershaw MJ & Talbot NJ (2009) Genome-wide functional analysis reveals that infection-
associated fungal autophagy is necessary for rice blast disease. *Proceedings of the National*
Academy of Sciences of the United States of America 106(37):15967-15972.
38. Holsters M*, et al.* (1978) Transfection and transformation of Agrobacterium tumefaciens.
Molecular & general genetics : MGG 163(2):181-187.
39. Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)* 25(4):402-
408.
40. Dean RA*, et al.* (2005) The genome sequence of the rice blast fungus Magnaporthe grisea.
Nature 434(7036):980-986.

Figures and Tables

Fig. 1 Ppe1 is important for full virulence in *M. oryzae*. (A) The relative expression level of Ppe1 during infection of rice leaf sheaths assayed by qRT-PCR. (B) Fluorescence signals of the P*PPE1*-mCherry transformant in penetration assays with rice leaf sheaths. Scale bars = 10 μm. (C) Typical leaves of CO-39 seedlings sprayed with conidia of the wild type (70-15), *PPE1* gene deletion mutants (Δ*ppe1-45*, Δ*ppe1-101*) and complement strains (Δ*ppe1-com-3*, Δ*ppe1-com-8*) examined at 7 days post inoculation (dpi). Boxplot below shows the number of disease lesions per 5 cm of individual inoculated leaf. n = 22, 23, 19, 19, and 22, respectively. (D) Wounded leaves of CO-39 drop-inoculated with the marked strains were examined at 10 dpi (upper panel) and measured for

the lesion area (lower panel). n = 10, 9, 7, 5, and 4 respectively. (E) Rice leaf sheath penetration

assays. Conidia suspensions from each strain were respectively injected into 4-5 weeks old rice

leaf sheaths. Infection levels were observed at 32 hpi (hours post inoculation). The infection levels

were grouped into four types: Type 1, appressorium formation without penetration; Type 2, having

penetration peg or primary invasive hyphae; Type 3, having more than two branched invasive

hyphae restricted in the first infected rice cell; Type 4, having invasive hyphae crossing to

neighboring host cells. Scale bar = 10 μm. Statistical analyses of the infection types and

percentages are also shown. Significant differences were determined by one-way ANOVA.

Asterisks indicated significant differences (p<0.0001). ns indicated no significant difference

(p>0.05).

Fig. 2 Ppe1-mCherry forms a ring during appressorial penetration. (A) A ring of Ppe1-mCherry

signals at the penetration site was observed at 22 hpi in leaf sheath penetration assays with the

PPE1-mCherry transformant. Upper panel indicates the Ppe1-mCherry expressed in wild-type

strain (70-15). Lower panel indicates the Ppe1-mCherry expressed in Δ*ppe1* mutant. Scale bars =

2 μm. (B) The Ppe1-mCherry penetration ring was persist even after the growth of primary and

secondary invasive hyphae. Scale bars = 5 μm. (C) The diameters of appressoria, Ppe1-mCherry

penetration ring (ring), and its inner cycle. n= 21, 65 and 65 respectively. (D) The formation of

Ppe1-GFP rings at the penetration site formed by the *PPE1*-GFP transformant. Scale bars = 2 μm.

(E) The in-situ *PPE1*-mCherry transformant also formed a fluorescent ring at the penetration site

in leaf sheath penetration assays. Scale bar = 2 μm. AP, Appressorium. Triangle arrows pointed to

penetration rings.

Fig. 3 Penetration ring formation during invasive hyphal penetration into neighboring cells. (A)

Ppe1-mCherry forms rings at about 36 hpi. (B) Ppe1-GFP also forms rings similar to Ppe1-

mCherry during penetration to neighboring cells. (C) Ppe1 ring formation is also observed during

barley infection. Scale bars = 10 μm. Arrows heads point to penetration rings at the penetration

sites of invasive hyphae. White squares indicate zoomed regions.

Fig. 4 The penetration ring localizes on the periphery of penetration peg. (A) Ppe1 ring surrounds the Sep3 constricted ring. Scale bar = 2 μm. (B) Ppe1 ring also surrounds the Lifeact ring. Scale bar = 1 μm. (C) Ppe1 ring is formed around the narrow site between an appressorium and invasive hypha. Three-dimensional (3D) volume rendering images were constructed by confocal optical z-series images captured at 0.35 μm intervals, with a total of 22 steps. (D) Ppe1 ring partly colocalized with callose ring at the apoplastic papilla. Scale bar = 1 μm. Hollow triangles indicate the site of penetration peg. Line graphs were generated at the directions pointed by the white arrows.

Fig. 5 Penetration ring formation requires the emergence of penetration peg. (A) Ppe1 ring expression and localization in 70-15, Δ*mst12*, Δ*nox2* and Δ*nox1* strains at 24 hpi. Conidia suspensions from each strain were inoculated on rice leaf sheaths and observed at 24 hpi. (B) Ppe1 ring could not be formed on impenetrable coverslips. Conidia from Ppe1-mCherry strain were inoculated on hydrophobic coverslips and observed at 24 hpi. (C) Ppe1 ring was observed at 38 hpi on penetrable cellophane membrane. Scale bars = 2 μm.

Fig. 6 Other secreted Ppe proteins also localize to the penetration ring. (A) Phylogenetic tree of Ppe1 constructed using neighbor-joining method. (B) Paralogs of Ppe1 in *M. oryzae* localize to the penetration ring. Ppe2-mCherry, Ppe3-mCherry and Ppe5-mCherry were driven by their respective native promoters. MoTPpe1-mCherry from *Magnaporthe oryzae Triticum* was expressed in *M. oryzae* (70-15), driven by its native promoter. Scale bars = 2 μm. (C) Comparison of the localization of penetration ring with those of some previously reported fungal effectors. Bas4-GFP and Slp1-GFP uniformly outlined the extra-invasive hyphal membrane (EIHM), while Pwl2-mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC). Scale bars = 10 μm.

Fig. 7 A proposed working model for the discovery of penetration ring in *M. oryzae*. Penetration

ring forms at the periphery of peg and accumulates fungal secreted protein such as Ppe1 to promote

M. oryzae invasion.

Supplementary figures

Fig. S1 Ppe1 is dispensable for mycelial growth, conidiation and appressorium formation in *M.*

oryzae. (A) Amino acid sequence of *M. oryzae* Ppe1. Signal peptides (green) were predicted by

Signal-5. Cysteine residues are shown in red. (B) Detection of Ppe1 signal peptide secretion by

yeast signal trap assay. The recombinant pSUC2-Ppe1SP and pSUC2-Pwl2SP constructs were

transformed into the yeast strain YTK12 respectively. The pSUC2-Pwl2SP-expressing strain

served as a positive control while pSUC2 empty vector was used as a negative control. (C)

Targeted gene deletion strategy for *PPE1* in *M. oryzae* 70-15. Schematic map showing targeted

disruption of *PPE1* gene via Agrobacterium tumefaciens-mediated transformation. The Southern

blot result shows a successful replacement of *PPE1* by single insertion of hygromycin

phosphotransferase (*HPH*) resistance cassette at *PPE1* ORF locus. (D) Colonies of the wild type

(70-15), two independent *Δppe1* transformants, and complemented strains. (E) Statistical analyses

of the colony diameters of each strain after being cultured on complete medium (CM) at 25 °C for

10 days. n = 9. (F) Statistical analyses of the number of conidia produced by each strain after being

cultured on CM at 25 °C for 10 days. n = 17. (G) Appressorium formation by the indicated strain

observed at 24 hours on artificial hydrophobic surfaces. Scale bar = 50 μm. Significant differences

were determined by one-way ANOVA. ns stands for no significant difference (p>0.05).

Fig. S2 Detection of Ppe1-mCherry fusion protein by western blot. M, protein marker. CK, rice

leaf sheaths injected with sterile double distilled water.

Fig. S3 Ppe1 ring is formed simultaneously with peg penetration. (A-C) Spatiotemporal dynamics of Ppe1 ring and Sep3 ring during appressorium-mediated peg penetration into rice leaf sheaths. Scale bars = 2 μm. Hollow triangles show the sites of penetration peg. Line graphs were generated at the directions pointed by the white arrows.

Fig. S4 Spatiotemporal dynamics of Ppe1-mCherry and Sep3-GFP on hydrophobic coverslips.

Conidia were harvested from a strain co-expressing Ppe1-mCherry and Sep3-GFP and inoculated

on hydrophobic coverslips and observed at each time point. Scale bars = 1 μm.

Fig. S5 Ppe1 ring was also observed during transpressorium-mediated peg penetration. (A-D) The

spatiotemporal dynamics of Ppe1-mCherry and Sep5-GFP during transpressorium-mediated cell-

to-cell penetration. Red squares indicate the zoomed regions, respectively. Scale bars = 10 μm.

Supplementary Videos

Video S1. 3D visualization of Ppe1-mCherry ring formation at 22 hpi.

Video S2. 3D visualization of Ppe1-mCherry and Sep3-GFP localizations at 11 hpi.

Video S3. 3D visualization of Ppe1-mCherry ring surrounding the Sep3-GFP small ring at 19
hpi.

Video S4. 3D visualization of Ppe1-mCherry ring surrounding the penetration peg at 22 hpi.

Video S5. 3D visualization of Ppe1-mCherry ring and papilla callose ring at the penetration site
at 22 hpi.

Supplementary Table S1. Fungal strains used in this study

Strain	Description	Reference
70-15	Wild-type strain	(40)
mCherry$_{PPE1\ promoter}$	pKNT-mCherry$_{PPE1\ promoter}$ transformant of 70-15	this study
Ppe1-mCherry	pKNT-MoPpe1-mCherry and pCB1532 co-transformant of 70-15	this study
Ppe1-GFP	pKNT-MoPpe1-GFP and pCB1532 co-transformant of 70-15	this study
Ppe1-mCherry $_{situ\ insertion}$	ATMT1-Ppe1-mCherry $_{situ\ insertion}$ transformant of 70-15	this study
Sep3-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Sep5-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Ppe1-mCherry + Sep3-GFP	pKNT-Ppe1-mCherry transformant of Sep3-GFP strain	this study
Ppe1-mCherry + Sep5-GFP	pKNT-Ppe1-mCherry transformant of Sep5-GFP strain	this study
Ppe1-mCherry + Lifeact-GFP	pXY198-Lifeact-GFP transformant of Ppe1-mCherry strain	this study
Ppe1-mCherry + GFP	pKNT-RP27-GFP transformant of Ppe1-mCherry strain	this study
Δppe1-45/101	*PPE1* deletion mutant of 70-15	this study
Δppe1-com-3/8	pKNT-Ppe1-mCherry transformant of Δppe1-45 mutant	this study
Δmst12	*MST12* deletion mutant of Guy11	(26)
Δnox1	*NOX1* deletion mutant of Ppe1-mCherry strain	this study
Δnox2	*NOX2* deletion mutant of Ppe1-mCherry strain	this study
Ppe2-mCherry	pKNT-MGG_10732-mCherry transformant of 70-15	this study
Ppe3-mCherry	pKNT-MGG_16041-mCherry transformant of 70-15	this study
Ppe5-mCherry	pKNT-MGG_16379-mCherry transformant of 70-15	this study
MoTPpe1-mCherry	pKNT-MoTPpe1-mCherry transformant of 70-15	this study

PBV591	pBV591(Co-expression of Pwl2-mCherry and Bas4-GFP) transformant of 70-15	(16)
Slp1-GFP	pKNT-Slp1-mCherry transformant of 70-15	this study
AvrPiz-t-GFP	pKNT-AvrPiz-t-GFP transformant of 70-15	this study

783
784 **References**
785

Dean, R. A., Talbot, N. J., Ebbole, D. J., Farman, M. L., Mitchell, T. K., Orbach, M. J., Thon, M., Kulkarni,
R., Xu, J. R., Pan, H., Read, N. D., Lee, Y. H., Carbone, I., Brown, D., Oh, Y. Y., Donofrio, N., Jeong, J. S.,
Soanes, D. M., Djonovic, S., Kolomiets, E., Rehmeyer, C., Li, W., Harding, M., Kim, S., Lebrun, M. H.,
Bohnert, H., Coughlan, S., Butler, J., Calvo, S., Ma, L. J., Nicol, R., Purcell, S., Nusbaum, C., Galagan, J. E.,
& Birren, B. W., 2005, Nature 434, 980.

Khang, C. H., Berruyer, R., Giraldo, M. C., Kankanala, P., Park, S. Y., Czymmek, K., Kang, S., & Valent, B.,
2010, The Plant cell 22, 1388.

Park, G., Xue, C., Zheng, L., Lam, S., & Xu, J. R., 2002, Molecular plant-microbe interactions : MPMI 15,
183.

Supplementary Table S2 List of Primers used in this study
797

Primer Name	Primer Sequence (5'→3')
Construction of pSUC2 plasmid for confirming the secretory function of signal peptide (SP)	
pSUC2-Ppe1SP-F	TTTTAATTAAGAATTCATGCGTTCATCTACGATTATTCTG
pSUC2-Ppe1SP-R	CTATAGGGAGAACCTCGAGGGCGGCAACCAGGCCAGTG
pSUC2-Pwl2SP-F	TTTTAATTAAGAATTCATGAAATGCAACAACATCATCCTC
pSUC2-Pwl2SP-R	CTATAGGGAGAACCTCGAGGGCGGTTACAGTGGTCG
pSUC2-yF	ACAGATGTCGTTGTTCCAGA
Construction of plasmid for studying subcellular localization	
pKNT-Ppe1-promoter-F	AGGGAACAAAAGCTGGGTACCTCTTTCTCCCAGTCTCATC
pKNT-Ppe1-promoter-R	GCCCTTGCTCACCATAAGCTTGTTGACGCCAGGTCGTGAGA
pKNT-Ppe1-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAATCCAGCATGC
ATMT-mCherry$_{situ\ instertion}$-AF	GAATTCGAGCTCGGTACCGGCAACGACGACAAGAA

ATMT-mCherry$_{situ\ insertion}$-AR	GTCGACTCTAGAGGATCCATTGCACATCCAATAAATCCAGC
mCherry$_{situ\ instertion}$-F	ATGGTGAGCAAGGGCGAGGAG
mCherry$_{situ\ instertion}$-R	TTGACCTCCACTAGCTCCAGCCAAGCCCTACTTGTACAGCTCGTCCATGCC
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HYG/R	AACCCGCGGTCGGCATCTACTCTATTC
ATMT-mCherry$_{situ\ insertion}$-BF	GAATAGAGTAGATGCCGACCGCGGGTTGTTGAGCGGAATTGCCTGTTAT
ATMT-mCherry$_{situ\ insertion}$-BR	GTCGACTCTAGAGGATCCTTCGCTTGTTGTTCTTCG
pKNT-AvrPiz-t-M-F	AGGGAACAAAAGCTGGGTACCCCAACGTAGAGCGTATGCTGT
pKNT-AvrPiz-t-M-R	GCCCTTGCTCACCATAAGCTTTTGGCGCTGAGCCTGAG
pCB1532-Slp1-G-F	ACTATAGGGCGAATTGGGTACCATTGAACAGTGCGATGCC
pCB1532-Slp1-G-R	CGCCCTTGCTCACCATAAGCTTGTTCTTGCAGATGGGGATG
GFP-sR	AGTTCACCTTGATGCCGTTC
mCherry-sR	TGGTCTTCTTCTGCATTACGG
pKNT-MoTPpe1-M-F	AGGGAACAAAAGCTGGGTACCTCTCTCAGTCTCATTCTACAGC
pKNT-MoTPpe1-M-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAATCCAGC
pKNT-MGG_10732-M-F	AGGGAACAAAAGCTGGGTACCGCACGTACCTTTGGTGGTTGC
pKNT-MGG_10732-M-R	GCCCTTGCTCACCATAAGCTTAGCACATCCAACATGTCCAGG
pFGL1010M-MGG_16041-M-F	CATGATGATGCTCGAGAATTCACGCCGCTTTCTGGACTA
pFGL1010M-MGG_16041-M-R	CTCGCCCTTGCTCACGGTACCGACGCATCCAACAAACCCAGGAA
pFGL1010M-MGG_16379-M-F	CATGATGATGCTCGAGAATTCTATCGTTTCGCGGGTAGATTG
pFGL1010M-MGG_16379-M-R	CTCGCCCTTGCTCACGGTACCCTTTGCGCACCCAATGAATGC

Construction of gene deletion

ATMT-Ppe1-AF	GAATTCGAGCTCGGTACCCCACCTTTTCGTCCTCC
ATMT-Ppe1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGACGCCAGGTCGTGAG

ATMT-Ppe1-BF	GAATAGAGTAGATGCCGACCGCGGGTTAGCGGAATTGCCTGTTA
ATMT-Ppe1-BR	GTCGACTCTAGAGGATCCACTGCCGACCGTGACTA
Ppe1-OF	ATGCGTTCATCTACGATTATTCTG
Ppe1-OR	TGCACATCCAATAAATCCAG
Ppe1-UA	CATCATCGCCGTCCTCT
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HY/R	GTATTGACCGATTCCTTGCGGTCCGAA
YG/F	GATGTAGGAGGGCGTGGATATGTCCT
HYG/R	AACCCGCGGTCGGCATCTACTCTATTC
H853	GACAGACGTCGCGGTGAGTT
Nox1-AF	CACCAGATGCAGTAGTCAATT
Nox1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTCGTTAAAGAGCGTAT GTC
Nox1-BF	GAATAGAGTAGATGCCGACCGCGGGTTGCCGAAATCTATTACGA CG
Nox1-BR	CCTGAATACTTCCAAGACACC
Nox1-OF	TCACCACCTGTTTATCCCG
Nox1-OR	GACACCAATACCAGTTCCAAT
Nox1-UA	CAGAGCCACGATGCGAAAA
Nox2-AF	CGATTGTGCCTGGATGA
Nox2-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGCTTGGTCGGTTGTC
Nox2-BF	GAATAGAGTAGATGCCGACCGCGGGTTGAAGCAAGCCTGGCAAT A
Nox2-BR	CAAAAGCCTGTTTACGAGAT
Nox2-OF	AAGATGGACGCCGCTAA
Nox2-OR	AAACGATGAAGAAGTGGTGA
Nox2-UA	ACCTCCTTGACTCTTCCACC
qRT-PCR analysis	
Ppe1-qF	TTTGCTGCGACGCTGGTA
Ppe1-qR	GGAAAATCTGGGAATGGGT
Tublin-qF	TGACAACGAGGCTCTGTACG
Tublin-qR	CCATGAAGAAGTGCAGACGA