The penetration ring is a novel infection structure formed by the penetration peg for

invading plant cell membrane in rice blast fungus

Wenqin Fang[1,2,3], Xiaoyu Zai[1,2], Jia Chen[1,2], Yakubu Saddeeq Abubakar[1,2,4], Qiuqiu Wu[1,2],

Zhenyu Fang[1,2], Xiuwei Huang[1,2], Xiang Gan[1,2], Daniel J. Ebbole[5], Zonghua Wang[1,2,3,]*, Wenhui

Zheng[1,2,]*

[1] State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection,

Fujian Agriculture and Forestry University, Fuzhou, Fujian, 350002, China.

[2] Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant

Protection, Fujian Agriculture and Forestry University, Fuzhou, Fujian, 350002, China.

[3] Fuzhou Institute of Oceanography, Minjiang University, Fuzhou 350108, China

[4] Department of Biochemistry, Faculty of Life Sciences, Ahmadu Bello University, Zaria,

810281, Nigeria

[5] Department of Plant Pathology & Microbiology, Texas A&M University, College Station, TX,

77843, USA

* Corresponding author:

Wenhui Zheng (wenhuiz@fafu.edu.cn)

Zonghua Wang (wangzh@fafu.edu.cn)

Author Contributions: D.J.E., Z.W., and W.Z. designed research; W.F., X.Z., J.C., Q.W., Z.F.,

X.H., X.G., and W.Z. performed research; W.F., Z.W., and W.Z. analyzed data; and W.F.,

Y.S.A., Z.W., and W.Z. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: penetration ring, secreted protein, *Magnaporthe oryzae*, peg, appressorium

Abstract

Many fungal pathogens develop specialized infection structures such as appressoria to penetrate plant cells. However, it is not clear whether special structures are formed after cell wall penetration before invading host plasma membrane in hemibiotrophic pathogens. Here, we showed that a penetration ring consisting of Ppe1 secreted proteins is formed after appressorium-mediated cell wall penetration and remained at the base of penetration site after invading plant plasma membrane by the rice blast fungus *Magnaporthe oryzae*. The same persistent Ppe1 ring is formed after the penetration of neighboring cells by transpressoria. *PPE1* is specifically expressed during plant infection and the Δ*ppe1* mutant is defective in penetration and invasive growth. Blockage of penetration peg formation impedes the development of the Ppe1 ring. Close examinations showed that the penetration ring is formed at the collar of penetration pegs between the plant cell wall and plasma membrane and it is persistent as a fixed ring even after invasive hyphae invaded neighboring cells. Furthermore, Ppe1 is a member of an expanded family of secreted proteins that are unique to fungal pathogens using extreme appressorium turgor for plant penetration. Other members of the Ppe1 family also localize to the penetration ring for anchoring on plasma membrane during plant infection. Taken together, a penetration ring consisting of a family of secreted proteins is formed between plant cell wall and plasma membrane, which may function as a novel physical structure at the interface between the tip of penetration pegs and plant plasma membrane before the differentiation of invasive hyphae.

Significance Statement

Like many other plant pathogens, the rice blast fungus forms melanized appressoria (specialized infection structures) to penetrate plant cells. In this study, we showed that a penetration ring is formed by penetration pegs after appressorium -mediated penetration of plant cell wall. This ring of secreted proteins is persistent at the point of penetration pegs invading plant plasma membrane to form invasive hyphae. Therefore, after the penetration of plant cell wall, the rice blast fungus forms a penetration ring that consists of a family of secreted proteins unique to pathogens using extreme appressorium turgor for penetration and may function as a physical

structure for anchoring onto plant plasma membrane and developing invasive hyphae in

penetrated cells.

Introduction

Magnaporthe oryzae, a filamentous ascomycete, is the causal agent of the devastating rice blast

disease, which poses a significant threat to global rice production (1, 2). In addition to rice, *M.*

oryzae is capable of infecting a wide range of grass species, including major cereal crops such as

wheat, barley, oat, and millet, thus compromising global food security (3, 4). As one of the most

important plant diseases, it is mainly controlled by planting resistant cultivars. Nevertheless,

outbreaks of rice blast still occur throughout rice growing regions worldwide because of

breakdown of resistance by new races of the pathogen. In addition, wheat blast caused by *M.*

oryzae pathovar *Triticum* has recently spread from South America to Asia, threatening wheat

production (5, 6).

Like many other foliar fungal pathogens, penetration of the cuticle and cell wall of host plants is

the most critical infection step (7, 8). The infection cycle of *M. oryzae* starts with attachment of

dispersing conidia to plant surface by releasing sport tip mucilage. Germ tubes emerging from

attached conidia can sense surface hydrophobicity and hardness and develop highly specialized

infection structures known as appressoria at the tip (1, 9). Mature dome-shaped appressoria are

anchored on rice surface with appressorium mucilage (10) and heavily melanized by depositing a

melanin layer between the cell wall and membrane as an impermeable barrier for turgor

generation (11). Subsequently, the appressorium uses the extremely high turgor pressure to

physically penetrate the cuticle and cell wall with a narrow penetration peg (7, 12, 13). As an

hemibiotrophic pathogen, after the penetration of plant cell wall, the penetration peg does not

penetrate plant plasma membrane but differentiates into primary invasive hyphae and invasive

hyphae that are enveloped by plant-derived extra-invasive hyphal membrane (EIHM) in

penetrated cells (7, 14). To date, it is not clear whether there is any special structure at the

contact point between the penetration peg and plant plasma membrane before the differentiation

of primary invasive hyphae. However, a biotrophic interfacial complex (BIC), a plant

membrane-rich structure, is formed at the tip of the primary invasive hyphae for effector delivery

(15, 16). Following proliferation within the first colonized cell, the tips of invasive hyphae

differentiate into transpressoria, which subsequently form narrow infection pegs to traverse the

plasmodesmata (PD)-rich cell junctions, allowing entry into adjacent host cells (17, 18). The

fungus continues this colonization strategy until disease symptoms manifest on infected rice

plants.

To suppress plant defense responses and facilitate invasive growth, *M. oryzae* deploys effectors

with various functions that can be delivered into host tissues or cells with two secretion systems

during infection (19). Apoplastic effectors, such as Bas4 and Slp1, accumulate in the

extracellular space between the fungal invasive hyphal cell wall and EIHM, outlining the entire

invasive hyphae, are secreted through the conventional endoplasmic reticulum (ER)-Golgi-

dependent pathway which is sensitive to brefeldin A (BFA) (19). Conversely, cytoplasmic

effectors, including Pwl2, AvrPiz-t, and Bas1, accumulate in the BIC before being delivered into

plant cells using a Golgi-independent pathway involving the exocyst complex and t-SNAREs,

which is insensitive to BFA (19). To date, all the known proteinaceous effectors in *M. oryzae*

belong to these two categories and localize to the extracellular space outlining invasive hyphae

or the BICs and intracellular targets inside plant cells.

In a previous study, we identified a family of small secreted proteins that may function as

effectors in *M. oryzae* (20). In this study, we found that one of them named as *PPE1* (for peg

periphery-localized effector 1, MGG_16603) is specifically expressed during plant infection and

is a crucial virulence factor. Surprisingly, Ppe1 preferentially localizes to the penetration site in a

ring conformation that is described as the penetration ring below. The penetration ring consisting

of Ppe1 proteins is formed after appressorium- and transpressorium- mediated plant penetration

and remained at the point where penetration pegs transitioning into primary invasive hyphae.

Ppe1 homologs are only present in fungal pathogens using extreme appressorium turgor for plant

penetration. Other members of the Ppe1 family also localize to the penetration ring that is at the

base of primary invasive hyphae outside plant plasma membrane. Taken together, a penetration

ring consisting of secreted Ppe family proteins is formed between plant cell wall and plasma

membrane, which may function as a novel physical structure at the interface between the tip of

penetration pegs and plant plasma membrane before differentiation into invasive hyphae.

Results

Ppe1 is important for full virulence of *M. oryzae*

We previously identified 21 host-adapted effector genes in *M. oryzae* that may play critical roles

during rice infection based on *in silico* analyses (20). To explore the functions of these effectors,

we selected *PPE1* (MGG_16603) for further functional characterization because its high

transcription level during early infection stages as reported (20, 21). We first assayed the

expression profile of *PPE1* during infection by qRT-PCR. The result showed that its expression

increased during the early stages of plant infection (24-36 hpi) and peaked at 36 hpi (Fig. 1A).

Moreover, fluorescence signals of mCherry driven by the *PPE1* promoter indicated that its

expression was only observed in matured appressoria and during infection (Fig. 1B). These

results were consistent with the transcriptome data, suggesting that Ppe1 is functional during

appressorium penetration and plant infection. Ppe1 encodes a 116-aa protein with typical

characteristics of fungal effectors, including an N-terminal signal peptide and eight cysteine

residues (Fig. S1A). The signal peptide of Ppe1 is functional in directing the secretion of yeast

Suc2 invertase in a yeast signal trap system (Fig. S1B), suggesting that Ppe1 is a secreted protein.

To determine the biological functions of Ppe1 in *M. oryzae,* we generated a *PPE1* gene deletion

mutant in the wild-type strain 70-15 (Fig. S1C). Deletion of *PPE1* had no obvious effect on the

fungal growth, conidiation, and appressorium formation (Fig. S1D-G). However, in spray

infection assays with rice leaves, the Δ*ppe1* mutant was significantly reduced in the number of

disease lesions compared to the wild type (Fig. 1C). In wound infection assays, the Δ*ppe1*

mutant also was significantly reduced in virulence and formed smaller lesions than the wild type

(Fig. 1D). To confirm the role of *PPE1* in virulence, we generated a complemented strain Δ*ppe1*-

com that had a copy of the *PPE1* gene ectopically integrated into its genome. In spray or wound

infection assays, Δ*ppe1*-com was similar with the wild type in virulence (Fig. 1C-D), indicating a

complete recovery in phenotypes. We then conducted rice leaf sheath penetration assays to

observe the infection processes. At 32 hpi, over 91% of the appressoria formed by the wild type

and the complemented strain had penetrated and developed invasive hyphae in rice cells.

Invasive hyphae in most of the penetrated cells (approximately 60%) had more than two

branches, and approximately 10% of them had spread into the neighboring cells (Fig. 1E). In

contrast, approximately 25% of the appressoria formed by the Δppe1 mutant had no penetration.

Although 53.5% of these mutant appressoria successfully penetrated into host cells, only 23% of

Δppe1 invasive hyphae had more than two branches and less than 4% had spread into adjacent

cells (Fig. 1E). To further validate these observations, we generated a *PPE1* knockout in the

wild-type reference strain Guy11. Similarly, we observed a significant decrease in the

pathogenicity of the Δppe1 mutants compared to Guy11 in the spray assay (Fig. S2). These

results indicate that the Δppe1 mutant has defects in appressorium penetration, invasive hyphal

growth and subsequent spread of invasive hyphae into neighboring plant cells.

Ppe1 forms a ring at the base of appressoria during host penetration

To determine the localization of Ppe1 during infection, we expressed a full-length Ppe1-mCherry

fusion protein under the control of *PPE1* promoter in both the wild type (70-15) and the Δppe1

mutant. Rice leaf sheaths were then infected with these strains and the localization of Ppe1 was

analyzed by confocal microscopy. In these transformants, Ppe1-mCherry signals were observed

as a ring underneath the appressoria (likely at the site of the emergence of penetration pegs) at 22

hpi (Fig. 2A, Video S1). The Ppe1-mCherry ring persisted after appressorial penetration and the

ring-shaped fluorescence signal of the Ppe1-mCherry was still observed even after the formation

of primary and secondary invasive hyphae (Fig. 2B). We further measured the size of this ring

and found that the average diameter of the inner circle of the Ppe1-mCherry ring was 1.53 μm.

The width of the Ppe1-mCherry ring varied from 1.79 to 4.78 μm, which is smaller than the

width of appressoria (Fig. 2C) and known septin ring (5.9 μm) at the appressorium pore (13).

Furthermore, the Ppe1-mCherry fusion protein could be detected by western blot analysis with

proteins isolated from infected rice leave (Fig. S3). Because the Ppe1 ring always appears at the

point of emergence of the penetration peg, we named this ring structure as penetration ring.

To confirm the localization of Ppe1, we also generated Ppe1-GFP transformants. Like Ppe1-

mCherry, Ppe1-GFP also formed a ring at the penetration site underneath the appressorium (Fig.

2D). These results indicate that, irrespective of being fused with mCherry or GFP, Ppe1 localizes

to the penetration ring. To avoid possible overexpression or unexpected effects due to ectopic

integration of the fusion constructs, we generated a transformant from 70-15 strain in which

mCherry was inserted right before the TAA stop codon of the endogenous *PPE1* gene. In the

resulting transformant expressing the in situ *PPE1*-mCherry fusion construct, the penetration

ring was also observed (Fig. 2E). These results further verify the localization of Ppe1 to the

penetration ring during appressorial penetration.

Penetration ring is also formed during the penetration of neighboring cells by
transpressoria

In *M. oryzae*, appressoria-like structures called transpressoria are formed by invasive hyphae on

the wall boundary of the invaded cell in order to penetrate into neighboring cells (17, 18).

Therefore, we observed localization of Ppe1 during transpressorium-mediated penetration of

plant cells. In the *PPE1*-mCherry transformant, like during appressorial penetration, the

penetration ring was observed during transpressoria-mediated penetration of neighboring cells

(Fig. 3A). Similarly, we observed the penetration ring in the neighboring cells penetrated by

invasive hyphae of the *PPE1*-GFP transformant (Fig. 3B). The Ppe1-mCherry or Ppe1-GFP ring

(Fig. 3A and B – zoomed view) appeared to be at the transpressorial penetration point. These

results indicate that the penetration ring is also formed during invasive hyphal cell-to-cell

penetration. Furthermore, similar localization of Ppe1 to the penetration ring was observed in

barley epidermal cells (Fig. 3C). Taken together, we conclude that Ppe1 localizes to the

penetration ring during appressorium-mediated and transpressorium-mediated plant cell

penetrations in *M. oryzae*.

Penetration ring is formed on the periphery of the penetration peg

In *M. oryzae*, appressorium-mediated penetration is a septin-dependent process in which septins

aggregate and generate a hetero-oligomeric 5.9 μm ring structure at the appressorial pore to re-

organize the F-actin cytoskeleton network to promote the emergence of penetration pegs (13).

Following the penetration peg formation, septin ring undergoes further constriction into a smaller

ring (22). To clarify the precise site of the penetration ring, we expressed Ppe1-mCherry (under

the control of the *PPE1* promoter) in a strain expressing Sep3-GFP (Guy11 background) and

then observed their spatiotemporal dynamics during rice leaf sheath infection. At 11 hpi, Sep3-GFP big ring was formed at the appressorium pore but the Ppe1-mCherry ring was not observed (Fig. S4A, Video S2). The Ppe1-mCherry ring signal was observed at 16 hpi, after the constriction of Sep3-GFP ring and the former was completely surrounding the latter (Fig. S4B). Subsequently (at 19 hpi), the Ppe1-mCherry ring (still surrounding the Sep3-GFP ring) became constricted (Fig. 4A, Fig. S4C, Video S3). However, on impenetrable hydrophobic coverslips, the Sep3-GFP ring was formed normally at the appressorium pore, but the Ppe1-mCherry ring and constriction of Sep3-GFP were not observed (Fig. S5). These results suggest that Ppe1-mCherry ring is formed after the formation of penetration pegs, outside the constricted Sep3-GFP ring.

Penetration pegs are rich in microfilaments, filasomes and microtubules (23), therefore, it can be labeled by Lifeact-GFP. Hence, we co-expressed Ppe1-mCherry with Lifeact-GFP in 70-15 and observed their positioning in penetration assays with rice leaf sheaths. The Ppe1-mCherry ring was also observed to surround the Lifeact-GFP signal (Fig. 4B, Fig. S6, Video S4), indicating that the penetration ring is localized at the periphery of penetration pegs during the initial infection stage and primary invasive hyphal development. To confirm this observation, we co-expressed the Ppe1-mCherry with cytosolic GFP in 70-15. Using a three-dimensional (3D) volume rendering image constructed by confocal optical z-series images, we found that the Ppe1-mCherry ring was situated around the narrow site between the appressorium and invasive hyphae labeled with GFP (Fig. 4C, Fig. S7).

A previous study showed that callose deposition is dynamic during *M. oryzae* infection and has three different forms: callose-rich papillae underneath appressorium penetration sites, callose collars formed at cell wall crossing sites around the invasive hyphae and callose deposition at plasmodesmata pit fields (24). To investigate whether the penetration ring locates outside the peg, we stained callose deposits with aniline blue to examine the possible positioning of the callose ring underneath the appressorial penetration site and Ppe1-mCherry. We discovered that the Ppe1-mCherry ring was surrounded by the callose ring, exhibiting partial colocalization and close attachment (Fig. 4D, Video S5). Furthermore, we examined the localization of Ppe1-

mCherry in transgenic rice expressing GFP-OsPIP2 as a plasma membrane marker (25). Our data demonstrate that the penetration ring is tightly associated with the host plasma membrane (Fig. S8). Taken together, we conclude that the penetration ring locates at the periphery of the penetration peg and anchors the peg to the host plasma membrane during the initial appressorium-mediated host penetration.

Formation of the penetration ring at the base of penetration pegs after transpressorium penetration

To determine the precise localization of the penetration ring during the transpressoria-mediated penetration, we expressed Ppe1-mCherry (under the control of the *PPE1* promoter) in a Sep5-GFP strain and then observed their spatiotemporal dynamics during rice leaf sheath infection. The results showed that Sep5-GFP first displayed a ring structure at the swollen part of the invasive hyphal tip (Fig. S9A), similar to the septin big ring at the appressorium pore. It then exhibited a collar structure at the constricted infection peg site (Fig. S9B-C), which is consistent with a previous report by Sakulkoo (26). Meanwhile, the Ppe1-mCherry ring or collar-like ring was formed outside the Sep5-GFP collar (Fig. S9B-C), indicating that the penetration ring formed during transpressoria-mediated cell-to-cell penetration is positioned outside the penetration peg. Taken together, we conclude that the penetration ring of *M. oryzae* is localized at the periphery of penetration pegs both during appressoria- and transpressoria-mediated host penetrations.

Penetration ring formation requires the initiation of penetration peg formation

To further characterize the relationship between the penetration ring and penetration peg, we observed the expression and localization of Ppe1-mCherry in Δ*mst12, Δnox2* and Δ*nox1* mutant strains that are defective in plant penetration (27, 28). In the Δ*mst12* mutant that is defective in both penetration initiation and peg formation, the Ppe1-mCherry fluorescence signal could not be detected. In the Δ*nox2* mutant that is unable to initiate penetration peg formation, the Ppe1-mCherry ring could not be formed but displayed a disorganized mass underneath the appressorium. Interestingly, in the Δ*nox1* mutant that forms normal penetration pegs but is defective in invasive hyphal elongation, the Ppe1-mCherry ring was formed during rice leaf

sheath infection (Fig. 5A), indicating that penetration peg formation is a prerequisite for the
emergence of the penetration ring. To confirm this observation, the localization of Ppe1-mCherry
fusion proteins was observed after inducing appressorium formation on hydrophobic coverslips
and cellophane membranes. Although melanized appressoria were formed, we failed to observed
the Ppe1-mCherry ring on the impenetrable coverslips (Fig. 5B). However, the penetration ring
is formed after the penetration of cellophane membranes (Fig. 5C). These results indicate that
penetration is necessary for the formation of the penetration ring.

Other Ppe proteins also localize to the penetration ring
To determine whether other secreted proteins homologous to Ppe1 also localize to the
penetration ring, we used the amino acid sequence of Ppe1 to carry out BLASTp search at the
NCBI database to uncover the homolog proteins of Ppe1 and further analyzed their phylogenetic
relationships. The Ppe1 homologs only exist in *Magnaporthe* and *Colletotrichum* species, such
as the most common cereal phytopathogens *M. oryzae* and *C. graminicola* (Fig. 6A).
Remarkably, these fungi are known to form melanized appressoria and deploy extremely high
turgor pressure for plant penetration (8). To understand whether other members of the Ppe1
family (Ppe2, Ppe3, Ppe3, Ppe4, Ppe5, and Ppe6) from *M. oryzae* have similar localization as
Ppe1, we tagged each of them with mCherry and expressed them individually in the 70-15
background under the control of their native promoters. Examination of their localizations during
infection revealed that Ppe2-mCherry, Ppe3-mCherry and Ppe5-mCherry also exhibit a ring
structure during appressorium-mediated host penetration, while the fluorescence signals of Ppe4-
mCherry and Ppe6-mCherry were hardly observed, possibly due to their low expression (Fig.
6B). Moreover, when MoT-Ppe1 from *M. oryzae* pathovar *Triticum* was expressed in *M. oryzae*
under the control of its native promoter, the characteristic penetration ring was also observed
(Fig. 6B). All these results suggest that the secreted Ppe proteins accumulate at the penetration
ring.

As controls, we also examined the localizations of Bas4, Slp1, Pwl2, and AvrPiz-t. Under the
same experimental conditions, Bas4-GFP and Slp1-GFP were observed to uniformly accumulate
within the extra-invasive hyphal membrane (EIHM), outlining the entire invasive hyphae, while

Pwl2-mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC) (Fig.

6C), which is consistent with previous reports (16, 29, 30). EIHM and BIC are specifically

associated with invasive hyphae during biotrophic growth. In contrast, Ppe1 localized to the

penetration ring at the periphery of penetration pegs (Fig. 6C). Therefore, unlike subcellular

localizations of Bas4, Slp1, Pwl2, and AvrPiz-t to the extracellular space outlining invasive

hyphae, BICs, or intracellular targets in plant cells, Ppe1 and its homologs localize to the

penetration ring that may function as a novel physical structure at the interface between the tip of

penetration pegs and plant plasma membrane before the differentiation of invasive hyphae.

Discussion

As the model plant pathogenic fungus, appressorium formation and appressorium-mediated

penetration have been well-characterized in *M. oryzae* (18). The fungus uses extremely high

turgor pressure generated in appressoria cemented to the surface with appressorium mucilage to

penetrate underlying plant cells (12). As an hemibiotrophic pathogen, after piercing plant cell

wall, the penetration peg does not penetrate the plasma membrane but differentiates into invasive

hyphae enveloped in a layer of EIHM (16). In this study, we showed that a penetration ring of

Ppe1 proteins is formed by the penetration peg on the surface of the plant plasma membrane

before invading and differentiating into primary invasive hyphae (Fig. 2 and Fig. 7). Other Ppe

proteins including Ppe2, Ppe3, and Ppe5 proteins also localize to the penetration ring (Fig. 6B).

Because the penetration ring is persistent at the point penetration pegs transitioning into invasive

hyphae outside the plasma membrane even after the penetrated cells are filled with invasive

hyphae, it may function as a novel physical structure for anchoring penetration pegs on the

surface of plant plasma membrane after cell wall penetration (Fig. 7). It is also possible that this

penetration ring functions as a collar that is associated with the differentiation of penetration

pegs (on the surface of plasma membrane) into primary invasive hyphae enveloped in the EIHM

(Fig. 7). Although this transition is not well characterized, it likely involves changes in cell

polarity and cytoskeleton as well as cell wall modifications. Ppe1 and its homologs may be able

to assemble into a protein complex in the penetration ring, which likely involves other proteins

or components as facilitating factors. In *U. maydis*, the Stp secreted effectors are known to form

a protein complex during plant infection (31).

The penetration ring is also formed after transpressoria-mediated penetration of neighboring plant cells (Fig. 7). Similarly, Ppe1 and its homologs form a ring after the penetration of cell wall but before differentiating into invasive hyphae enveloped with the EIHM. The penetration ring formed by transpressoria-mediated penetration is also persistent at the penetration site on the surface of plant plasma membrane in penetrated cells (Fig. 3 and Fig. S5), suggesting that Ppe proteins are assembled into a complex that is not diffusible. In fact, similar penetration ring is observed after penetration of artificial membrane (Fig. 5C), which has no barrier to restrict the diffusion of secreted proteins, further indicating that the penetration ring of Ppe1 and other Ppe proteins is likely a structural feature or component for signaling the transition of penetration pegs to invasive hyphae. Although appressoria and transpressoria are different, the processes after the penetration of cell wall are the same in appressorium- and transpressorium-mediated plant penetration. Therefore, formation of the penetration ring in both appressorium- and transpressorium-mediated penetration is another evidence for it to be a structural element.

M. oryzae is known to secrete a repertoire of effectors that can be categorized into cytoplasmic and intracellular effectors (14). Ppe1 and its homologs are putative effectors that are described as HAG (host adaptation genes) effectors in an earlier study (20). Although it remains possible that Ppe1 and its homologs function as effectors during plant infection. However, in comparison with the localizations of known effectors in *M. oryzae*, we found that the localization of Ppe1 to the penetration ring is unique and Ppe1 homologs have the same localization pattern (Fig. 6). Because of the persistent localization of Ppe1 and its homologs to the penetration ring, we think the penetration ring is likely a physical structure as described above. If Ppe1 and its homologs are effector proteins being secreted at this specific location, fluorescent signals of Ppe1 and its homologs will diffuse and disappear or dimmish when effectors being translocated into plant cells. Furthermore, it is unlikely that secreted proteins localized to the penetration ring that has limited contact with the plant plasma membrane will function as effector proteins to interact with plant targets. Nevertheless, although unlikely, it remains possible that the penetration ring discovered in this study is a secretion site for secreted proteins involved in the transition of penetration pegs to invasive hyphae. After the penetration of plant cell wall, the penetration ring

may not only function as a physical structure but also serve as an secretion site for the release of
specific proteins to overcome plant immunity in early infection stages.

Taken together, in this study we observed the formation of the penetration ring that appears to be
a structural element for signaling or triggering the differentiation of penetration pegs after the
penetration of plant cell wall into invasive hyphae enveloped by the EIHM in *M. oryzae*. Based
on the limited distribution of Ppe1 homologs, it is likely that only fungal pathogens using high
appressorium turgor for plant penetration, such as *M. oryzae* and *Colletotrichum* species,
developed the penetration ring. Therefore, it will be important to further characterize the
functional relationship between the penetration ring and appressorium turgor as well as the
transition of penetration pegs driven by turgor pressure into primary invasive hyphae (lower
turgor pressure).

In summary, our findings have several significant implications. First, we believe that the
discovery of the penetration ring as a novel physical structure associated with the differentiation
of invasive hyphae represents a breakthrough in plant-pathogen interactions. Secondly, our study
presents new role of the peg as a specialized platform for the deployment of secreted protein, in
addition to its commonly known role as a physical penetration tool for the pathogen. Thirdly, we
identified Ppe1 as a potential molecular target for controlling the devastating rice blast disease,
as Ppe homologs are absent in plants and mammals. The primary challenge for future research
lies in elucidating the pathogenesis associated with penetration rings. Achieving this will further
our understanding of the mechanism of rice blast fungus-host interaction.

371
372

Materials and Methods
Strains and growth conditions
The *M. oryzae* strains 70-15 and Guy11 were used as the wild types in this study; all other fungal
strains generated from the wild types are listed in Table S1 in the supplemental materials. All the
fungal strains were grown on complete medium CM (CM: 10 g D-glucose, 2 g peptone, 1 g yeast
extract, 1 g casamino acids, 50 mL 20 × nitrate salts (120 g/L $NaNO_3$, 10.4 g/L KCl, 30 g/L
KH_2PO_4, and 10.4 g/L $MgSO_4$), 1 mL trace elements (0.1 g/L biotin, 0.1 g/L pyridoxine, 0.1 g/L

thiamine, 0.1 g/L riboflavin, 0.1 g/L p-aminobenzonic acid and 0.1 g/L nicotinic acid), adjusted

to pH 6.5 with NaOH, 15 g agar and distilled water added to 1 L) (32). Agar cultures were

incubated at 26 °C with a 12-h light/12-h dark photoperiod for mycelial growth and conidiation.

Liquid CM medium was used to harvest mycelia for extraction of genomic DNA and protoplast

preparation. The bacterial strains *Escherichia coli* DH5α and *Agrobacterium tumefaciens* AGL1

were grown in Luria-Bertani (LB) medium at 37 °C and 28 °C, respectively.

Plant and growth conditions

388   Rice cultivar CO-39 (*Oryza sativa* cv. CO-39) was grown in a greenhouse at 26 □ and 70%

389   relative humidity with a 14-h light/10-h dark photoperiod. Barley (*Hordeum vulgare* cv. Golden

390   Promise) was grown at 23 □ with a 12-h light /12-h dark photoperiod.

Bioinformatic and phylogenetic analyses of Ppe1 homolog proteins

393   The Ppe1 was previously sequenced and the amino acid sequence established from our lab (33).

394   Ppe1 signal peptide was predicted using SignalP 5.0 software. BLASTp was used to find Ppe1

395   homolog proteins at the NCBI database (https://www.ncbi.nlm.nih.gov/) and the representative

396   homolog protein sequences were downloaded from the database. Phylogenetic tree was

397   generated using MEGA7.0 software by neighbor-joining method.

Construction of recombinant DNA and *M. oryzae* transformation

The plasmids used in the study were generated by infusion cloning method based on homologous

recombination using ClonExpress II one-step cloning kit or ClonExpress MultiS kit (Cat. No.

C112-01 or C113-01, Vazyme, Nanjing, China). The primers used (Table S2 in the supplemental

material) were specially designed to contain 18-22-bp overlap with adjoining sequences to allow

the fragments assemble by homologous recombination. To observe the expression pattern and

localization of *PPE1*, the *PPE1* promoter (1485 bp) fragment amplified from the genomic DNA

of the wild type (70-15) was seamlessly cloned into the *Kpn*I/*Hin*dIII-digested pKNT-

mCherry/GFP fusion vectors (34) using the ClonExpress II one-step cloning kit (Cat. No. C112-

01, Vazyme, Nanjing, China). For Ppe1-mCherry/GFP plasmid, the entire Ppe1 coding sequence

with its native promoter but without stop codon was amplified and cloned into the *Kpn*I/*Hin*dIII-

digested vector pKNT-mCherry/GFP. The pKNT-AvrPizt-GFP and other plasmids used were

was constructed using the same method. The constructed plasmids were transformed into the

fungal protoplasts via PEG-mediated transformation (35) and further verified by PCR screening

and confocal microscopic examination of fluorescence signals. For situ fusion of Ppe1 with

mCherry, the mCherry was inserted just before the Ppe1 stop codon and the construct was

transformed by *Agrobacterium tumefaciens*-mediated (ATMT) transformation for homologous

recombination (35).

Assays for the secretory function of the SP of Ppe1

To confirm the secretory function of the predicted signal peptide (SP), yeast signal trap system

was performed as described previously (36). The predicted SP nucleotide sequences of Ppe1 (60

bp) and Pwl2 (63 bp) were respectively cloned into pSUC2 vector that carries a truncated

invertase gene *SUC2*. The generated pSUC2-SP constructs and the empty pSUC2 vector were

transformed into the yeast *suc2* mutant YTK12, respectively, and then their growth conditions

analyzed on CMD-W (minus Trp) plates (0.67% yeast nitrogen base without amino acids,

0.075% tryptophan dropout supplement, 2% sucrose, 0.1% glucose, and 2% agar) and YPRAA

medium plate (1% yeast extract, 2% peptone, 2% raffinose, and 2 μg/mL antimicyn A)

containing raffinose and lacking glucose. Transformants of YTK12 carrying the empty pSUC

vector or pSUC2-Pwl2SP were used as negative and positive controls, respectively. Reduction of

2,3,5-triphenyltetrazolium chloride (TTC) to insoluble red-colored 1,3,5-triphenylformazan

(TPF) was used as an indicator of the invertase enzymatic activity (37). Each of the above

YTK12 transformants strain was inoculated in 5 mL of CMD-W liquid medium and incubated

for 24 h at 30 □. The cells were then collected, washed and resuspended in 0.1% TTC in new

centrifuge tubes and further incubated at 35 □ for 35 min. The color change was observed after

10 min of incubation at room temperature to detect the signal peptide secretion function.

Targeted gene deletion and complementation

Deletions of *NOX1*, *NOX2* and *MST12* genes were performed using Split-marker recombination

method (38). The open reading frames (ORFs) of the various genes were replaced with a

hygromycin phosphotransferase (*HPH*) resistance cassette, respectively. Briefly, about 1-kb

upstream and downstream sequences of the targeted genes were amplified from the genomic

DNA of the wild type (70-15) using specially designed primer pairs AF+AR and BF+BR,

respectively. At the same time, the upstream and downstream sequences of the *HPH* with

overlapping segments were amplified from pCB1003 plasmid using the primer pairs HYG/F +

HY/R and YG/F + HYG/R, respectively. The corresponding upstream and downstream

fragments were respectively ligated by splicing for overlap extension (SOE) PCR using the

primer pairs AF+HY/R and YG/F+BR. The PCR products were purified and transformed into

the Ppe1-mCherry strain protoplasts via a PEG-mediated transformation (35). Putative

transformants were selected in the presence of hygromycin B and then confirmed by PCR using

the primers OF+OR and UA+H853. PCR confirmations were repeated three times for

consistency. All the primer sequences used are listed in Table S2 of the supplemental material.

For *PPE1* gene deletion, *HPH* sequence was inserted between the upstream and downstream

sequences of the *PPE1* gene, replacing the ORF of the *PPE1* gene, in an ATMT1 plasmid using

a ClonExpress MultiS One Step Cloning Kit (Cat. no. C113-01, Vazyme, Nanjing, China). The

constructed plasmid was introduced into *Agrobacterium tumefaciens* AGL1 using freeze-thaw

method (39). *Agrobacterium tumefaciens*-mediated transformation (ATMT) of the construct into

the wild-type spores was then carried out. The transformants were selected on hygromycin B

plates and confirmed by PCR. The mutants were further confirmed by Southern blot assay using

DIG-High Prime DNA labeling and detection starter kit I (Cat. no. 11745832910, Roche,

Mannheim, Germany). For complementation, the constructed plasmid pKNT-Ppe1-mCherry was

transformed into the Δ*ppe1* mutant protoplast and further verified by PCR screening and

confocal microscopic examination for fluorescence signal detection.

Live cell imaging

To monitor the expression and subcellular localization of Ppe1 during plant infection, rice leaf

sheath inoculation and live cell imaging of fluorescently labeled proteins were performed as

described (15) with slight modifications. In brief, 5-7 cm long leaf sheath segments were cut

from 4-week-old CO-39 rice seedlings, and the whole hollow spaces of the sheaths were filled

with conidia suspensions (5×10^4 conidia/mL) using a 200 μL pipette. The inoculated sheaths

were placed horizontally (fixed with wet filter paper fragments) on an inverted 96-well PCR

plate covered by one layer of wet filter paper and incubated in a sealed box whose inner bottom

was lined with two layers of wet filter paper to keep the environment humid at 26 □. The

epidermal parts of the inoculated sheaths were trimmed into thin and transparent samples and

immediately imaged using an inverted Nikon A1 laser scanning confocal microscope (Nikon,

Tokyo, Japan) at the indicated time points post inoculation. The 60×/1.42 numerical aperture

(NA) oil immersion objective lens was used for the imaging. The fluorescence and differential

interference contrast (DIC) images were collected by confocal optical z-series acquired at 0.28

µm intervals spanning the entire fluorescence signal depth between the top plane of the

appressorium formed and the bottom plane of the rice leaf sheath cross sections. The

fluorescence excitation/emission wavelengths were 488 nm/500-550 nm for GFP, 561 nm/570-

620 nm for mCherry and 405 nm/425-475 nm for callose staining. The images were processed

using NIS-Elements AR analysis software (Nikon, Tokyo, Japan). Unless otherwise stated in the

figure legend, micrograph fluorescence represents maximum intensity projection of z-series

(with a 0.28 µm intervals) after removing extraneous z-planes that do not contain relevant

information for a particular fluorescence. Three-dimensional (3D) volume rendering images and

videos were generated from z-series in maximum intensity projection.

To observe the subcellular localizations of Ppe1 in conidia incubated on artificial hydrophobic

surfaces and barley leaves, conidia suspensions ($5×10^4$ conidia/mL) were dropped on the surfaces

of hydrophobic coverslips or on the back sides of barley leaves (5-day old plants) and incubated

under humid and dark conditions at a 26 □. For observation, the hydrophobic coverslips

containing the appressoria were inverted, placed on 24mm×50 mm microscope glass slides and

then observed under a scanning confocal microscope. Epidermal parts were removed from the

infected barley leaves for observation. For penetration assay, cellophane was cut into round

sheets of 4 cm in diameter and sterilized by autoclaving at 115 □ for 15 min. The sterile

cellophane sheets were placed flat on CM plates and then the conidia suspensions ($5×10^4$

conidia/mL) were dropped on the surface of the cellophane and incubated at 26 □. Small pieces

of the cellophane containing mycelia were cut for observation under a confocal microscope at

36-48 hpi.

Callose staining

For visualization of deposited callose, aniline blue (Cat. no. 28631-66-5, Sangon, Shanghai, China) staining was adopted (26). Rice leaf sheaths were inoculated with conidia suspensions (3×10^4 conidia/mL). The inoculated sheaths were trimmed to thin and transparent samples at the indicated time points and stained with aniline blue (0.1mg/mL) solution prepared in 150 mM KH_2PO_4 (pH 9.5) for up to 30 min. The stained samples were imaged by using a laser scanning confocal microscope (Nikon, Japan).

Phenotypic analysis

For vegetative growth and conidiation assays, small mycelial blocks from each strain were cut from the edges of 4-day-old cultures and placed at the center of freshly prepared CM plates (90 mm) and incubated for ten days at 26 □. The colony diameters of the cultures were measured using vernier calipers and then photographs of the cultures were taken. The cultures were then washed with 5 mL sterilized double-distilled water, filtered through three layers of lens paper and the conidia counted using a hemocytometer under a light microscope. For conidia germination and appressoria formation, spore suspensions (1×10^5 conidia/mL) were incubated on hydrophobic coverslips (Cat. no. 12-545-101P, Fisherbrand™ Cover Glasses, USA) under humid and dark conditions at 26 □ for 24 h. The conidia were then observed under a light microscope for germination and formation of appressoria.

To investigate the pathogenicity of each strain, conidia spray, punch and leaf sheath inoculations were performed. Conidia suspensions (5×10^4 conidia/mL containing 0.02% Tween-20) were used to spray the leaf surfaces of 3-week-old CO-39 rice seedlings. The inoculated seedlings were incubated in a growth chamber at 25 □ with 90% humidity. The seedlings were first kept in the dark for 24 h, and then subjected to 12 h light / 12 h dark photoperiod for 7 days, after which photographs were taken and the number of disease lesions on the leaf surfaces counted. Punch inoculation assay was carried out according to a method previously described (30). Briefly, 5-week-old CO-39 rice leaves were slightly wounded using mouse ear punch and 10 μL of conidia suspensions (5×10^5 conidia/mL containing 0.02% Tween-20) from the various strains were

respectively dropped at the wounded regions. Then, both sides of each inoculated spot were

sealed with scotch tape to sustain the spore suspension. The inoculated plants were kept in a

growth chamber under the same conditions described above for 10 days. The disease lesions

formed were scanned, and later analyzed using Adobe Photoshop software to measure the lesion

areas. For leaf sheaths inoculation, conidia suspensions (1×10^5 conidia/mL) from the various

strains were respectively injected into 4-5 weeks old CO-39 rice leaf sheaths. The epidermal

parts of the infected leaves were trimmed to thin and transparent samples for observation under a

light microscope at 32 hpi. At least 100 infected sites were observed using a Nikon A1 laser

confocal microscope (Nikon, Tokyo, Japan).

qRT-PCR analysis

To analyze the expression level of *PPE1* during *M. oryzae* infection, rice leaf sheaths were cut

from 4- to 5-week-old CO-39 rice seedings and inoculated with conidia suspensions (2×10^5

conidia/mL) from the wild type (70-15). The epidermal parts of the inoculated sheaths were cut

and collected at 12-, 36- and 48 hours post inoculation (hpi). At least 15 segments of the

inoculated rice sheath samples were collected at each time point. Total RNAs were extracted

from the leaf sheaths or rice seeding leaves using an Eastep Super Total RNA Extraction Kit

(Cat. no. LS1040, Promega, Shanghai, China). Reverse transcription was used to generate cDNA

using a reverse transcription kit (Cat. no. RR037B, Takara, Dalian, China). Quantitative PCR

was finally conducted using a SYBR kit (Cat. no. RR067A, Takara, Dalian, China) using

specific primers (Table S2). Fugal *β-TUBULIN* gene used as endogenous reference genes, and

the expression levels were calculated using $2^{-\Delta\Delta Ct}$ method as previously reported (40). All qRT-

PCR assays were repeated three times for each sample.

Statistical analyses

Statistical analyses were performed using GraphPad Prism7 (GraphPad Software, Inc., San

Diego, CA). Statistically significant differences were determined by one-way ANOVA with two-

sided least significant difference. **** indicated significant difference at $p < 0.0001$, ns

indicated no significant difference.

Data Availability

The authors declare that all the data supporting the findings of this study are available within the paper and SI Appendix. Raw data and details of experimental procedures are available upon request from the corresponding authors.

Acknowledgments

We thank Prof. Nicholas J. Talbot and Prof. Barbara Valent for providing us with Δ*sec5* mutant and pBV591 plasmid respectively. We also thank Prof. Min Guo for providing us with Sep3/5-GFP strains and Δsep3/4/5 mutants. We also appreciate Profs. Guodong Lu, Jie Zhou, Huawei Zheng, Wei Tang, Ya Li, Xiaofeng Chen, and Huakun Zheng, as well as Drs Yijuan Hang, Meilian Chen and Yonghe Hong for their helpful suggestions. Thanks to Jin Zhang, Chunling Li and Yuping Fan's help. This research was supported by the National Key Research and Development Program of China (2023YFD1400200), the National Natural Science Foundation of China (32122071, 31601596), and the Natural Science Foundation of Fujian Province (2021J06015).

References

1. Talbot NJ (2003) On the trail of a cereal killer: Exploring the biology of Magnaporthe grisea. *Annual review of microbiology* 57:177-202.
2. Dean R, *et al.* (2012) The Top 10 fungal pathogens in molecular plant pathology. *Molecular plant pathology* 13(4):414-430.
3. Couch BC, *et al.* (2005) Origins of host-specific populations of the blast pathogen Magnaporthe oryzae in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice. *Genetics* 170(2):613-630.
4. Langner T, Białas A, & Kamoun S (2018) The Blast Fungus Decoded: Genomes in Flux. *mBio* 9(2).
5. Ceresini PC, *et al.* (2018) Wheat Blast: Past, Present, and Future. *Annual review of phytopathology* 56:427-456.
6. Malaker PK, *et al.* (2016) First Report of Wheat Blast Caused by Magnaporthe oryzae Pathotype triticum in Bangladesh. 100(11):2330-2330.
7. Howard RJ & Valent B (1996) Breaking and entering: host penetration by the fungal rice blast pathogen Magnaporthe grisea. *Annual review of microbiology* 50:491-512.
8. Talbot NJ (2019) Appressoria. *Current biology : CB* 29(5):R144-r146.
9. Hamer JE, Howard RJ, Chumley FG, & Valent B (1988) A mechanism for surface attachment in spores of a plant pathogenic fungus. *Science (New York, N.Y.)* 239(4837):288-290.

595   10.   Rocha RO, Elowsky C, Pham NTT, & Wilson RA (2020) Spermine-mediated tight sealing of the
596   Magnaporthe oryzae appressorial pore-rice leaf surface interface. *Nature microbiology*
597   5(12):1472-1480.

598   11.   Chumley FG & Valent BJMP-ml (1990) Genetic analysis of melanin-deficient, nonpathogenic
599   mutants of Magnaporthe grisea. 3:135-143.

600   12.   de Jong JC, McCormack BJ, Smirnoff N, & Talbot NJ (1997) Glycerol generates turgor in rice blast.
601   *Nature* 389(6648):244-244.

602   13.   Dagdas YF, *et al.* (2012) Septin-mediated plant cell invasion by the rice blast fungus,
603   Magnaporthe oryzae. *Science (New York, N.Y.)* 336(6088):1590-1595.

604   14.   Fernandez J & Orth K (2018) Rise of a Cereal Killer: The Biology of Magnaporthe oryzae
605   Biotrophic Growth. *Trends in microbiology* 26(7):582-597.

606   15.   Kankanala P, Czymmek K, & Valent B (2007) Roles for rice membrane dynamics and
607   plasmodesmata during biotrophic invasion by the blast fungus. *The Plant cell* 19(2):706-724.

608   16.   Khang CH, *et al.* (2010) Translocation of Magnaporthe oryzae effectors into rice cells and their
609   subsequent cell-to-cell movement. *The Plant cell* 22(4):1388-1403.

610   17.   Cruz-Mireles N, Eseola AB, Osés-Ruiz M, Ryder LS, & Talbot NJ (2021) From appressorium to
611   transpressorium-Defining the morphogenetic basis of host cell invasion by the rice blast fungus.
612   *PLoS pathogens* 17(7):e1009779.

613   18.   Ryder LS, *et al.* (2022) The appressorium at a glance. *Journal of cell science* 135(14).

614   19.   Giraldo MC, *et al.* (2013) Two distinct secretion systems facilitate tissue invasion by the rice blast
615   fungus Magnaporthe oryzae. *Nature communications* 4:1996.

616   20.   Ebbole DJ, *et al.* (2021) Evolution and Regulation of a Large Effector Family of Pyricularia oryzae.
617   *Molecular plant-microbe interactions : MPMI* 34(3):255-269.

618   21.   Yan X, *et al.* (2023) The transcriptional landscape of plant infection by the rice blast fungus
619   Magnaporthe oryzae reveals distinct families of temporally co-regulated and structurally
620   conserved effectors. *The Plant cell* 35(5):1360-1385.

621   22.   Eisermann I & Talbot NJ (2024) Septin-dependent invasive growth by the rice blast fungus
622   Magnaporthe oryzae. *Journal of Plant Diseases and Protection*.

623   23.   Bourett TM & Howard RJ (1992) Actin in penetration pegs of the fungal rice blast
624   pathogen,Magnaporthe grisea. *Protoplasma* 168(1-2):20-26.

625   24.   Osés-Ruiz M, *et al.* (2021) Appressorium-mediated plant infection by Magnaporthe oryzae is
626   regulated by a Pmk1-dependent hierarchical transcriptional network. *Nature microbiology*
627   6(11):1383-1397.

628   25.   Chen Z, *et al.* (2019) Green Fluorescent Protein- and Discosoma sp. Red Fluorescent Protein-
629   Tagged Organelle Marker Lines for Protein Subcellular Localization in Rice. *Frontiers in plant*
630   *science* 10:1421.

631   26.   Sakulkoo W, *et al.* (2018) A single fungal MAP kinase controls plant cell-to-cell invasion by the
632   rice blast fungus. *Science (New York, N.Y.)* 359(6382):1399-1403.

633   27.   Park G, Xue C, Zheng L, Lam S, & Xu JR (2002) MST12 regulates infectious growth but not
634   appressorium formation in the rice blast fungus Magnaporthe grisea. *Molecular plant-microbe*
635   *interactions : MPMI* 15(3):183-192.

636   28.   Ryder LS, *et al.* (2013) NADPH oxidases regulate septin-mediated cytoskeletal remodeling during
637   plant infection by the rice blast fungus. *Proceedings of the National Academy of Sciences of the*
638   *United States of America* 110(8):3179-3184.

639   29.   Mentlak TA, *et al.* (2012) Effector-mediated suppression of chitin-triggered immunity by
640   magnaporthe oryzae is necessary for rice blast disease. *The Plant cell* 24(1):322-335.

30. Park CH, *et al.* (2012) The Magnaporthe oryzae effector AvrPiz-t targets the RING E3 ubiquitin
ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *The*
Plant cell 24(11):4748-4762.
31. Ludwig N, *et al.* (2021) A cell surface-exposed protein complex with an essential virulence
function in Ustilago maydis. *Nature microbiology* 6(6):722-730.
32. Talbot NJ, Ebbole DJ, & Hamer JE (1993) Identification and characterization of MPG1, a gene
involved in pathogenicity from the rice blast fungus Magnaporthe grisea. *The Plant cell*
5(11):1575-1590.
33. Zhong Z, *et al.* (2018) Population genomic analysis of the rice blast fungus reveals specific events
associated with expansion of three main clades. *The ISME journal* 12(8):1867-1878.
34. Chen X, *et al.* (2021) Trehalose Phosphate Synthase Complex-Mediated Regulation of Trehalose
6-Phosphate Homeostasis Is Critical for Development and Pathogenesis in Magnaporthe oryzae.
mSystems 6(5):e0046221.
35. Khang CH, Park SY, Rho HS, Lee YH, & Kang S (2006) Filamentous Fungi (Magnaporthe grisea and
Fusarium oxysporum). *Methods in molecular biology (Clifton, N.J.)* 344:403-420.
36. Jiang C, *et al.* (2020) An orphan protein of Fusarium graminearum modulates host immunity by
mediating proteasomal degradation of TaSnRK1α. *Nature communications* 11(1):4382.
37. Oh SK, *et al.* (2009) In planta expression screens of Phytophthora infestans RXLR effectors reveal
diverse phenotypes, including activation of the Solanum bulbocastanum disease resistance
protein Rpi-blb2. *The Plant cell* 21(9):2928-2947.
38. Kershaw MJ & Talbot NJ (2009) Genome-wide functional analysis reveals that infection-
associated fungal autophagy is necessary for rice blast disease. *Proceedings of the National*
Academy of Sciences of the United States of America 106(37):15967-15972.
39. Holsters M, *et al.* (1978) Transfection and transformation of Agrobacterium tumefaciens.
Molecular & general genetics : MGG 163(2):181-187.
40. Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)* 25(4):402-
408.
41. Dean RA, *et al.* (2005) The genome sequence of the rice blast fungus Magnaporthe grisea.
Nature 434(7036):980-986.

676 **Figures and Tables**

Fig. 1 Ppe1 is important for full virulence in *M. oryzae*. (A) The relative expression level of Ppe1 during infection of rice leaf sheaths assayed by qRT-PCR. (B) Fluorescence signals of the P*PPE1*-mCherry transformant in penetration assays with rice leaf sheaths. Scale bars = 10 μm. (C) Typical leaves of CO-39 seedlings sprayed with conidia of the wild type (70-15), *PPE1* gene deletion mutants (Δ*ppe1-45*, Δ*ppe1-101*) and complement strains (Δ*ppe1-com-3*, Δ*ppe1-com-8*) examined at 7 days post inoculation (dpi). Boxplot below shows the number of disease lesions per 5 cm of individual inoculated leaf. n = 22, 23, 19, 19, and 22, respectively. (D) Wounded leaves of CO-39 drop-inoculated with the marked strains were examined at 10 dpi (upper panel) and measured for the lesion area (lower panel). n = 10, 9, 7, 5, and 4 respectively. (E) Rice leaf sheath penetration assays. Conidia suspensions from each strain were respectively injected into 4-5 weeks old rice leaf sheaths. Infection levels were observed at 32 hpi (hours post inoculation).

The infection levels were grouped into four types: Type 1, appressorium formation without

penetration; Type 2, having penetration peg or primary invasive hyphae; Type 3, having more

than two branched invasive hyphae restricted in the first infected rice cell; Type 4, having

invasive hyphae crossing to neighboring host cells. Scale bar = 10 μm. Statistical analyses of the

infection types and percentages are also shown. Significant differences were determined by one-

way ANOVA. Asterisks indicated significant differences (p<0.0001). ns indicated no significant

difference (p>0.05).

Fig. 2 Ppe1-mCherry forms a ring during appressorial penetration. (A) A ring of Ppe1-mCherry

signals at the penetration site was observed at 22 hpi in leaf sheath penetration assays with the

PPE1-mCherry transformant. Upper panel indicates the Ppe1-mCherry expressed in wild-type

strain (70-15). Lower panel indicates the Ppe1-mCherry expressed in Δ*ppe1* mutant. Scale bars =

2 μm. (B) The Ppe1-mCherry penetration ring was persist even after the growth of primary and

secondary invasive hyphae. Scale bars = 5 μm. (C) The diameters of appressoria, Ppe1-mCherry

penetration ring (ring), and its inner circle. n= 21, 65 and 65 respectively. (D) The formation of

Ppe1-GFP rings at the penetration site formed by the *PPE1*-GFP transformant. Scale bars = 2

μm. (E) The in-situ *PPE1*-mCherry transformant also formed a fluorescent ring at the penetration

site in leaf sheath penetration assays. Scale bar = 2 μm. AP, Appressorium. Triangle arrows

pointed to penetration rings.

Fig. 3 Penetration ring formation during invasive hyphal penetration into neighboring cells. (A)

Ppe1-mCherry forms rings at about 36 hpi. (B) Ppe1-GFP also forms rings similar to Ppe1-

mCherry during penetration to neighboring cells. (C) Ppe1 ring formation is also observed during

barley infection. Scale bars = 10 μm. Arrows heads point to penetration rings at the penetration

sites of invasive hyphae. White squares indicate zoomed regions.

Fig. 4 The penetration ring localizes on the periphery of penetration peg. (A) Ppe1 ring surrounds the Sep3 constricted ring. Scale bar = 2 μm. (B) Ppe1 ring also surrounds the Lifeact ring. Scale bar = 1 μm. (C) Ppe1 ring is formed around the narrow site between an appressorium and invasive hypha. Three-dimensional (3D) volume rendering images were constructed by confocal optical z-series images captured at 0.35 μm intervals, with a total of 22 steps. (D) Ppe1 ring partly colocalized with callose ring at the apoplastic papilla. Scale bar = 1 μm. Hollow triangles indicate the site of penetration peg. Line graphs were generated at the directions pointed by the white arrows.

Fig. 5 Penetration ring formation requires the emergence of penetration peg. (A) Ppe1 ring expression and localization in 70-15, Δ*mst12*, Δ*nox2* and Δ*nox1* strains at 24 hpi. Conidia suspensions from each strain were inoculated on rice leaf sheaths and observed at 24 hpi. (B) Ppe1 ring could not be formed on impenetrable coverslips. Conidia from Ppe1-mCherry strain were inoculated on hydrophobic coverslips and observed at 24 hpi. (C) Ppe1 ring was observed at 38 hpi on penetrable cellophane membrane. Scale bars = 2 μm.

Fig. 6 Other secreted Ppe proteins also localize to the penetration ring. (A) Phylogenetic tree of Ppe1 constructed using neighbor-joining method. (B) Paralogs of Ppe1 in *M. oryzae* localize to the penetration ring. Ppe2-mCherry, Ppe3-mCherry and Ppe5-mCherry were driven by their respective native promoters. MoTPpe1-mCherry from *Magnaporthe oryzae Triticum* was expressed in *M. oryzae* (70-15), driven by its native promoter. Scale bars = 2 μm. (C) Comparison of the localization of penetration ring with those of some previously reported fungal effectors. Bas4-GFP and Slp1-GFP uniformly outlined the extra-invasive hyphal membrane (EIHM), while Pwl2-mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC). Scale bars = 10 μm.

Fig. 7 A proposed working model for the discovery of penetration ring in *M. oryzae*. Penetration ring forms at the periphery of peg and accumulates fungal secreted protein such as Ppe1 to promote *M. oryzae* invasion.

762   **Supplementary figures**

Fig. S1 Ppe1 is dispensable for mycelial growth, conidiation and appressorium formation in *M. oryzae*. (A) Amino acid sequence of *M. oryzae* Ppe1. Signal peptides (green) were predicted by Signal-5. Cysteine residues are shown in red. (B) Detection of Ppe1 signal peptide secretion by yeast signal trap assay. The recombinant pSUC2-Ppe1SP and pSUC2-Pwl2SP constructs were transformed into the yeast strain YTK12 respectively. The pSUC2-Pwl2SP-expressing strain served as a positive control while pSUC2 empty vector was used as a negative control. (C) Targeted gene deletion strategy for *PPE1* in *M. oryzae* 70-15. Schematic map showing targeted disruption of *PPE1* gene via Agrobacterium tumefaciens-mediated transformation. The Southern blot result shows a successful replacement of *PPE1* by single insertion of hygromycin phosphotransferase (*HPH*) resistance cassette at *PPE1* ORF locus. (D) Colonies of the wild type (70-15), two independent *Δppe1* transformants, and complemented strains. (E) Statistical analyses of the colony diameters of each strain after being cultured on complete medium (CM) at 25 °C for 10 days. n = 9. (F) Statistical analyses of the number of conidia produced by each strain after being cultured on CM at 25 °C for 10 days. n = 17. (G) Appressorium formation by the indicated strain observed at 24 hours on artificial hydrophobic surfaces. Scale bar = 50 μm. Significant differences were determined by one-way ANOVA. ns stands for no significant difference (p>0.05).

Fig. S2 Ppe1 is essential for virulence in *M. oryzae* strain Guy11. (A) Disease symptoms on leaves of CO-39 rice seeding inoculated with conidial suspension (5×10^{4} spores/ml) from

786   Guy11, Δ*ppe1-21* and Δ*ppe1-24*. Photographs were taken 7 days after inoculation. (B)

787   Quantification of the lesion number per 5 cm length of rice leaf. Asterisk represents significant

788   difference. **, $p < 0.01$. ***, $p < 0.001$.

791   Fig. S3 Detection of Ppe1-mCherry fusion protein by western blot. M, protein marker. CK, rice

792   leaf sheaths injected with sterile double distilled water.

Fig. S4 Ppe1 ring is formed simultaneously with peg penetration. (A-C) Spatiotemporal dynamics of Ppe1 ring and Sep3 ring during appressorium-mediated peg penetration into rice leaf sheaths. Scale bars = 2 μm. Hollow triangles show the sites of penetration peg. Line graphs were generated at the directions pointed by the white arrows.

Fig. S5 Spatiotemporal dynamics of Ppe1-mCherry and Sep3-GFP on hydrophobic coverslips.

Conidia were harvested from a strain co-expressing Ppe1-mCherry and Sep3-GFP and inoculated

on hydrophobic coverslips and observed at each time point. Scale bars = 1 μm.

Fig. S6 3D images of the co-expression of Ppe1-mCherry and Lifeact-GFP during primary

invasive hyphal development. AP, Appressorium. Arrow shows penetration ring.

Fig. S7 Co-expression of Ppe1-mCherry with cytoplasmic GFP at the early infection stage of *M.*

oryzae. (A) Ppe1-mCherry signal formed a tight ring-like structure closely associated with the

base of the appressorium at 16 hpi. Scale bar = 5 µm. (B) At 23 hpi, Ppe1-mCherry ring signal

was still beneath the appressorium, and around the penetration peg which differentiated into the

primary invasive hyphae. Scale bar = 10 µm. AP, Appressorium. IH, invasive hyphae. hpi, hours

post-inoculation. Arrows show penetration ring.

Fig. S8 The spatial positioning of Ppe1-mCherry ring and rice plasma membrane (marked by

GFP-OsPIP2). The zoomed images represent the sites pointed by the white arrows, respectively.

Scale bar = 10 µm; 2 µm for the zoomed images.

Fig. S9 Ppe1 ring was also observed during transpressorium-mediated peg penetration. (A-D)

The spatiotemporal dynamics of Ppe1-mCherry and Sep5-GFP during transpressorium-mediated

cell-to-cell penetration. Red squares indicate the zoomed regions, respectively. Scale bars = 10

μm.

Supplementary Videos

Video S1. 3D visualization of Ppe1-mCherry ring formation at 22 hpi.

Video S2. 3D visualization of Ppe1-mCherry and Sep3-GFP localizations at 11 hpi.

Video S3. 3D visualization of Ppe1-mCherry ring surrounding the Sep3-GFP small ring at 19

hpi.

Video S4. 3D visualization of Ppe1-mCherry ring surrounding the penetration peg at 22 hpi.

Video S5. 3D visualization of Ppe1-mCherry ring and papilla callose ring at the penetration site

at 22 hpi.

Supplementary Table S1. Fungal strains used in this study

Strain	Description	Reference
70-15	Wild-type strain	(41)
mCherry$_{PPE1\ promoter}$	pKNT-mCherry$_{PPE1\ promoter}$ transformant of 70-15	this study
Ppe1-mCherry	pKNT-MoPpe1-mCherry and pCB1532 co-transformant of 70-15	this study
Ppe1-GFP	pKNT-MoPpe1-GFP and pCB1532 co-transformant of 70-15	this study
Ppe1-mCherry $_{situ\ insertion}$	ATMT1-Ppe1-mCherry $_{situ\ insertion}$ transformant of 70-15	this study
Sep3-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Sep5-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Ppe1-mCherry + Sep3-GFP	pKNT-Ppe1-mCherry transformant of Sep3-GFP strain	this study
Ppe1-mCherry + Sep5-GFP	pKNT-Ppe1-mCherry transformant of Sep5-GFP strain	this study
Ppe1-mCherry + Lifeact-GFP	pXY198-Lifeact-GFP transformant of Ppe1-mCherry strain	this study
Ppe1-mCherry + GFP	pKNT-RP27-GFP transformant of Ppe1-mCherry strain	this study
Δppe1-45/101	PPE1 deletion mutant of 70-15	this study
Δppe1-com-3/8	pKNT-Ppe1-mCherry transformant of Δppe1-45 mutant	this study
Δmst12	MST12 deletion mutant of Guy11	(27)
Δnox1	NOX1 deletion mutant of Ppe1-mCherry strain	this study
Δnox2	NOX2 deletion mutant of Ppe1-mCherry strain	this study
Ppe2-mCherry	pKNT-MGG_10732-mCherry transformant of 70-15	this study
Ppe3-mCherry	pKNT-MGG_16041-mCherry transformant of 70-15	this study
Ppe5-mCherry	pKNT-MGG_16379-mCherry transformant of	this study

	70-15	
MoTPpe1-mCherry	pKNT-MoTPpe1-mCherry transformant of 70-15	this study
PBV591	pBV591(Co-expression of Pwl2-mCherry and Bas4-GFP) transformant of 70-15	(16)
Slp1-GFP	pKNT-Slp1-mCherry transformant of 70-15	this study
AvrPiz-t-GFP	pKNT-AvrPiz-t-GFP transformant of 70-15	this study

References

Dean, R. A., Talbot, N. J., Ebbole, D. J., Farman, M. L., Mitchell, T. K., Orbach, M. J., Thon, M., Kulkarni, R.,

Xu, J. R., Pan, H., Read, N. D., Lee, Y. H., Carbone, I., Brown, D., Oh, Y. Y., Donofrio, N., Jeong, J. S.,

Soanes, D. M., Djonovic, S., Kolomiets, E., Rehmeyer, C., Li, W., Harding, M., Kim, S., Lebrun, M. H.,

Bohnert, H., Coughlan, S., Butler, J., Calvo, S., Ma, L. J., Nicol, R., Purcell, S., Nusbaum, C., Galagan, J. E.,

& Birren, B. W., 2005, Nature 434, 980.

Khang, C. H., Berruyer, R., Giraldo, M. C., Kankanala, P., Park, S. Y., Czymmek, K., Kang, S., & Valent, B.,

2010, The Plant cell 22, 1388.

Park, G., Xue, C., Zheng, L., Lam, S., & Xu, J. R., 2002, Molecular plant-microbe interactions : MPMI 15,

183.

Supplementary Table S2 List of Primers used in this study

Primer Name	Primer Sequence (5'→3')
Construction of pSUC2 plasmid for confirming the secretory function of signal peptide (SP)	
pSUC2-Ppe1SP-F	TTTTAATTAAGAATTCATGCGTTCATCTACGATTATTCTG
pSUC2-Ppe1SP-R	CTATAGGGAGAACCTCGAGGGCGGCAACCAGGCCAGTG
pSUC2-Pwl2SP-F	TTTTAATTAAGAATTCATGAAATGCAACAACATCATCCTC
pSUC2-Pwl2SP-R	CTATAGGGAGAACCTCGAGGGCGGTTACAGTGGTCG
pSUC2-yF	ACAGATGTCGTTGTTCCAGA
Construction of plasmid for studying subcellular localization	
pKNT-Ppe1-promoter-F	AGGGAACAAAGCTGGGTACCTCTTTCTCCCAGTCTCATC
pKNT-Ppe1-promoter-R	GCCCTTGCTCACCATAAGCTTGTTGACGCCAGGTCGTGAGA
pKNT-Ppe1-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAATCCAGCATGC
ATMT-mCherry$_{situ\ instertion}$ -	GAATTCGAGCTCGGTACCGGCAACGACGACAAGAA

AF	
ATMT-mCherry_{situ insertion}-AR	GTCGACTCTAGAGGATCCATTGCACATCCAATAAATCCAGC
mCherry_{situ instertion}-F	ATGGTGAGCAAGGGCGAGGAG
mCherry_{situ instertion}-R	TTGACCTCCACTAGCTCCAGCCAAGCCCTACTTGTACAGCTCGTCCATGCC
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HYG/R	AACCCGCGGTCGGCATCTACTCTATTC
ATMT-mCherry_{situ instertion}-BF	GAATAGAGTAGATGCCGACCGCGGGTTGTTGAGCGGAATTGCCTGTTAT
ATMT-mCherry_{situ instertion}-BR	GTCGACTCTAGAGGATCCTTCGCTTGTTGTTCTTCG
pKNT-AvrPiz-t-M-F	AGGGAACAAAAGCTGGGTACCCCAACGTAGAGCGTATGCTGT
pKNT-AvrPiz-t-M-R	GCCCTTGCTCACCATAAGCTTTTGGCGCTGAGCCTGAG
pCB1532-Slp1-G-F	ACTATAGGGCGAATTGGGTACCATTGAACAGTGCGATGCC
pCB1532-Slp1-G-R	CGCCCTTGCTCACCATAAGCTTGTTCTTGCAGATGGGGATG
GFP-sR	AGTTCACCTTGATGCCGTTC
mCherry-sR	TGGTCTTCTTCTGCATTACGG
pKNT-MoTPpe1-M-F	AGGGAACAAAAGCTGGGTACCTCTCTCAGTCTCATTCTACAGC
pKNT-MoTPpe1-M-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAATCCAGC
pKNT-MGG_10732-M-F	AGGGAACAAAAGCTGGGTACCGCACGTACCTTTGGTGGTTGC
pKNT-MGG_10732-M-R	GCCCTTGCTCACCATAAGCTTAGCACATCCAACATGTCCAGG
pFGL1010M-MGG_16041-M-F	CATGATGATGCTCGAGAATTCACGCCGCTTTCTGGACTA
pFGL1010M-MGG_16041-M-R	CTCGCCCTTGCTCACGGTACCGACGCATCCAACAAACCCAGGAA
pFGL1010M-MGG_16379-M-F	CATGATGATGCTCGAGAATTCTATCGTTTCGCGGGTAGATTG
pFGL1010M-MGG_16379-M-R	CTCGCCCTTGCTCACGGTACCCTTTGCGCACCCAATGAATGC

Construction of gene deletion

ATMT-Ppe1-AF	GAATTCGAGCTCGGTACCCCACCTTTTCGTCCTCC
ATMT-Ppe1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGACGCCAGGTCGTGAG

ATMT-Ppe1-BF	GAATAGAGTAGATGCCGACCGCGGGTTAGCGGAATTGCCTGTTA
ATMT-Ppe1-BR	GTCGACTCTAGAGGATCCACTGCCGACCGTGACTA
Ppe1-OF	ATGCGTTCATCTACGATTATTCTG
Ppe1-OR	TGCACATCCAATAAATCCAG
Ppe1-UA	CATCATCGCCGTCCTCT
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HY/R	GTATTGACCGATTCCTTGCGGTCCGAA
YG/F	GATGTAGGAGGGCGTGGATATGTCCT
HYG/R	AACCCGCGGTCGGCATCTACTCTATTC
H853	GACAGACGTCGCGGTGAGTT
Nox1-AF	CACCAGATGCAGTAGTCAATT
Nox1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTCGTTAAAGAGCGTATGTC
Nox1-BF	GAATAGAGTAGATGCCGACCGCGGGTTGCCGAAATCTATTACGACG
Nox1-BR	CCTGAATACTTCCAAGACACC
Nox1-OF	TCACCACCTGTTTATCCCG
Nox1-OR	GACACCAATACCAGTTCCAAT
Nox1-UA	CAGAGCCACGATGCGAAAA
Nox2-AF	CGATTGTGCCTGGATGA
Nox2-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGCTTGGTCGGTTGTC
Nox2-BF	GAATAGAGTAGATGCCGACCGCGGGTTGAAGCAAGCCTGGCAATA
Nox2-BR	CAAAAGCCTGTTTACGAGAT
Nox2-OF	AAGATGGACGCCGCTAA
Nox2-OR	AAACGATGAAGAAGTGGTGA
Nox2-UA	ACCTCCTTGACTCTTCCACC
qRT-PCR analysis	
Ppe1-qF	TTTGCTGCGACGCTGGTA
Ppe1-qR	GGAAAATCTGGGAATGGGT
Tublin-qF	TGACAACGAGGCTCTGTACG
Tublin-qR	CCATGAAGAAGTGCAGACGA