The replication principle revisited: a shared functional organization between pulvinar-cortical and cortico-cortical connectivity and its structural and molecular imaging correlates

Gianpaolo Antonio Basile[1], Augusto Ielo[2], Lilla Bonanno[2], Antonio Cerasa[3,4,5], Giuseppe Santoro[1], Demetrio Milardi[1], Giuseppe Pio Anastasi[1], Ambra Torre[1], Sergio Baldari[6], Riccardo Laudicella[6], Michele Gaeta[7], Marina Quartu[8], Maria Pina Serra[8], Marcello Trucas[8], Angelo Quartarone[2], Manojkumar Saranathan[9], Alberto Cacciola[1,*]

[1] Brain Mapping Lab, Department of Biomedical, Dental Sciences and Morphological and Functional Imaging, University of Messina, Messina, Italy

[2] IRCCS Centro Neurolesi "Bonino Pulejo", Messina, Italy

[3] Institute for Biomedical Research and Innovation (IRIB), National Research Council of Italy (CNR), Messina, Italy

[4] S. Anna Institute, 1680067 Crotone, Italy

[5] Pharmacotechnology Documentation and Transfer Unit, Preclinical and Translational Pharmacology, Department of Pharmacy, Health Science and Nutrition, University of Calabria, 87036 Arcavacata, Italy

[6] Nuclear Medicine Unit, Department of Biomedical, Dental Sciences and Morphological and Functional Imaging, University of Messina, Messina, Italy

[7] Radiology Unit, Department of Biomedical, Dental Sciences and Morphological and Functional Imaging, University of Messina, Messina, Italy

[8] Section of Cytomorphology, Department of Biomedical Sciences, University of Cagliari, Cittadella Universitaria di Monserrato, 09042 Monserrato CA, Italy

[9] Department of Radiology, University of Massachusetts Chan Medical School, Worcester, MA, USA

* Correspondence

Alberto Cacciola, MD

Brain Mapping Lab, Department of Biomedical, Dental Sciences and Morphological and Functional Imaging, University of Messina, Messina, Italy

Phone: +39 0902217143

alberto.cacciola0@gmail.com

Abstract

The pulvinar, the largest nucleus in the human thalamus, is a complex, highly interconnected

structure. Through a dense, organized network of cortical and subcortical areas, it provides adequate

cooperation between neural systems, which is crucial for multiple high-order functions such as

perception, visuospatial attention, and emotional processing. Such a central role is made possible by

a precise internal topographical organization, which is mirrored by anatomical connections as well

as by the expression of neurochemical markers. While being traditionally subdivided into sub-

nuclei, each characterized by distinct connectional and morphological features, recent studies in

both primate and human brains have highlighted that this topographical organization only

marginally aligns with the conventional histological subdivision. Instead, it has been delineated in

the context of continuous gradients of cortical connections along the dorsoventral and mediolateral

axes. While this multi-gradient organization has been extensively documented in primate models, it

remains relatively underexplored in the human brain. The present work combines high-quality,

multi-modal structural and functional imaging data with a recently published whole-brain, large-

scale, positron emission tomography (PET) atlas detailing 19 neurotransmitters and receptors

distributed across the human brain. By applying diffusion embedding analysis to tractography,

functional connectivity, and receptor coexpression data, we identify and characterize multiple

topographically organized gradients of structural connections, functional coactivation, and

molecular binding patterns. We demonstrate that such gradients converge on a shared representation

along the dorsoventral and mediolateral axes of the human pulvinar. This representation aligns with

transitions in both structural and functional connectivity, spanning from lower-level to higher-order

cortical regions. Moreover, it is paralleled by gradual changes in the expression of molecular

markers associated with key neuromodulator systems, including serotoninergic, noradrenergic,

dopaminergic, and opioid systems. We contend that our findings mark a significant stride towards a

more comprehensive understanding of pulvinar anatomy and function, providing a nuanced

characterization of its role in health and disease.

Keywords

Connectivity, dMRI, fMRI, gradients, PET

Introduction

The pulvinar complex stands out as the largest nucleus within the human thalamus, serving as a pivotal hub in a myriad of cortico-subcortical networks that intricately interconnect various cortical regions of the brain (Benarroch 2015). Anatomical studies in primate brains have highlighted its primary sources of afferent and efferent connections, primarily stemming from the primary and secondary visual areas. However, its connectivity also extends to other crucial regions such as the temporal lobe, primary sensory, prefrontal, and cingulate cortices (Pons and Kaas 1985; Romanski et al. 1997b; Lyon and Kaas 2002; Fang et al. 2006). Through its extensive interplay with cerebral cortical areas, the pulvinar is believed to play a pivotal role in orchestrating integrative processes crucial for context-dependent modulation of visuospatial attention (Jaramillo et al. 2019; Fiebelkorn and Kastner 2020). It has been postulated that the pulvinar mediates the selection of relevant information from the environment by generating alpha oscillations thereby potentially modulating neuronal gain in thalamocortical circuits (Bourgeois et al. 2020). Nonetheless, despite the acknowledged significance of cortico-pulvinar connectivity in these theoretical frameworks, the functional and anatomical organization of this network in the human brain remains relatively underexplored.

Traditionally, the primate pulvinar complex has been anatomically divided into four sub-nuclei: the anterior (oral), the inferior, the lateral, and the medial pulvinar nuclei (Olszewski and Baxter 1982). While this subdivision has long served as a foundational framework for understanding pulvinar organization, anatomical investigations in non-human primates have revealed that connectivity patterns extend beyond discrete cytoarchitectonic subdivisions (Lysakowski et al. 1988; Webster et al. 1993; Adams et al. 2000). Instead, accumulating evidence suggests a spatially continuous organization in space, wherein multiple representations of the cortical sheet and visual fields coexist (Shipp 2001). These representations, often termed "maps", adhere to the "replication principle," wherein densely connected cortical regions exhibit overlapping representations within the pulvinar

(Shipp 2003). Moreover, these representations are loosely associated with chemoarchitectural

domains (Gutierrez and Cusick 1997; Stepniewska and Kaas 1997; Gutierrez et al. 2000).

Translating our understanding of the functional and anatomical organization of the pulvinar

complex to the human brain has proven challenging due to the small size of the underlying

structures and the inherent limitations of imaging techniques. Consequently, there has been a

relative scarcity of experimental works investigating pulvinar structure and function in vivo in the

human brain. While functional and structural neuroimaging techniques have been successfully

employed to explore the anatomy and connectivity of the human pulvinar (Leh et al. 2008; Tamietto

et al. 2012; Arcaro et al. 2015, 2018; Mai and Majtanik 2019; Basile et al. 2021), only a few studies

have delved into the connectional topography of this structure. Functional MRI-based parcellation

studies, employing task-based activation or resting-state connectivity fingerprinting, have provided

insights into functional specialization within sub-regions of the human pulvinar (Barron et al. 2015;

Arcaro et al. 2018; Guedj and Vuilleumier 2020). These investigations confirm that, akin to

findings in non-human primates, functional regions within the human pulvinar are loosely

associated with the anatomical subdivision into nuclei. However, relying on voxel-wise clustering

methods, these studies share the common assumption of discrete connectivity units and therefore

may not fully capture the spatially continuous and transient nature of pulvinar connectivity patterns.

Recently, gradient mapping techniques have emerged as a valuable tool for investigating the

structure-function relationship in the human brain. These techniques utilize dimensional

decomposition algorithms, such as diffusion embedding (Coifman and Lafon 2006), to map high-

dimensional to manifold low-dimensional brain features, known as "gradients". These gradients are

commonly interpreted as representing spatial patterns of smooth transitions in biological features of

interest, either within or across brain structures (Haak et al. 2018; Hong et al. 2020). Over the past

decade, gradient mapping methods have been extensively applied to various biological features,

including structural and functional connectivity, structural covariance, MRI-based microstructural

measures, receptor, and gene expression (Bajada et al. 2017; Paquola et al. 2019b, a, 2020; Valk et

al. 2020; Vos De Wael et al. 2021; Hansen et al. 2022b). These methods have been employed to explore the functional anatomy of the cerebral cortex (Margulies et al. 2016), as well as specific cortical or subcortical areas, such as the striatum, thalamus, hippocampus, and cerebellum (Guell et al. 2018; Przeździk et al. 2019; Tian et al. 2020; Yang et al. 2020; Müller et al. 2020; Katsumi et al. 2023).

The present study aims to investigate the gradient organization of the human pulvinar using a multimodal approach. By leveraging high-quality structural, diffusion-weighted, and resting state functional MRI from two independent datasets (Human Connectome Project; Leipzig Mind-Brain-Body dataset) (Van Essen et al. 2013; Babayan et al. 2019), along with a recently published multi-tracer receptor expression atlas derived from ~1000 positron emission tomography (PET) scans (Hansen et al. 2022b), we aim to provide insights on the spatial organization of connectivity within the human pulvinar and its relation to molecular expression.

Results

1. The multiscale gradient organization of the human pulvinar goes beyond discrete anatomical subdivisions

We collected functional and diffusion-weighted imaging data from two high-quality, independent datasets of healthy subjects: a primary dataset from the Human Connectome Project (HCP) (Van Essen et al. 2012), including 210 healthy subjects (males=92, females=118, age range 22-36 years); and a validation dataset from the Leipzig Mind-Brain-Body dataset (LEMON) (Babayan et al. 2019), consisting of 213 healthy subjects (males=138, females=75, age range 20-70 years). The pulvinar complex, along with its constituent subnuclei, was delineated bilaterally in the left and right hemispheres according to a whole brain labeling atlas (Rolls et al. 2020). For each participant, we generated voxel-wise estimates of pulvinar functional and structural connectivity to a 400-label parcellation of the cerebral cortex (Schaefer et al. 2018), respectively from preprocessed resting-state functional MRI and from whole brain probabilistic tractograms derived from diffusion data.

Individual connectivity maps were then normalized and aggregated to group-level, dense functional

and structural connectomes.

From the normative neurotransmitter atlas (Hansen et al. 2022b), we extracted the voxel-wise

density profiles of 19 receptors and transporters across nine different neurotransmitter systems

(serotonin, noradrenaline, dopamine, glutamate, gamma-aminobutyric acid, acetylcholine,

histamine, opiates, endocannabinoids), and we calculated the pairwise correlation for each voxel

(coexpression). The functional and structural dense connectomes, as well as the coexpression

matrix, were converted to modality-specific affinity matrices quantifying the similarity of each

voxel profile to each other.

By applying diffusion embedding (Coifman and Lafon 2006) on modality-specific affinity matrices

sampled from left and right pulvinar, we derived distinct functional connectivity, structural

connectivity, and receptor expression embeddings (Figure 1).

Figure 1. Schematic overview of the gradient mapping protocol. For each different imaging modality (DWI; BOLD;
PET) a feature matrix was built by extracting voxel-wise features in the left and right pulvinar separately. Structural and
functional connectivity to 400 cortical areas, as well as the expression profiles of 19 receptors and transporters, were
extracted for each pulvinar voxel. Symmetric, affinity matrices comparing the feature profile similarity of each voxel
against each other were estimated using cosine similarity and fed into the diffusion embedding algorithm. Additional
details can be found in the main text.

These embeddings, which we henceforth refer to as "gradients", represent unitless entities in which

each value identifies the position of pulvinar voxels along the respective embedding axis that

encodes dominant differences in voxel connectivity or coexpression patterns (Margulies et al. 2016;

Haak et al. 2018).

The first three functional connectivity gradients ($G_{FC}1$-$G_{FC}3$) collectively explained ~80% of the

total variance in functional connectivity for both the left and right pulvinar, as evidenced by the

elbow observed in the scree plot (Figure 2A). The principal gradient ($G_{FC}1$, ~50% of total variance

explained for both left and right pulvinar) reflected a dorsomedial-to-ventrolateral topographical

organization. Of the two secondary gradients, explaining each approximately ~10/15% of the total

variance for both left and the right pulvinar, one ($G_{FC}2$) ran to the posteromedial part of the pulvinar

to the most anterior dorsal region, the other ($G_{FC}3$) was aligned to the posterolateral to the

anteromedial axis (Figure 2A).

Nearly 90% of the global variance in structural connectivity values was explained by the first two

connectivity gradients ($G_{SC}1$-$G_{SC}2$). The first structural gradient ($G_{SC}1$, ~70% of total variance

explained, bilaterally) was arranged on a medio-lateral axis, while the second ($G_{SC}2$), explaining

approximately 20% of global explained variance, delineated a dorsal-to-ventral topography (Figure

2B).

Finally, the first three gradient embeddings ($G_{RC}1$-$G_{RC}3$) explained globally ~80% of the variance

in receptor coexpression data both for the left and right pulvinar, although with minor differences

between the left and right hemispheres for individual gradients. The main gradient ($G_{RC}1$),

explaining 60% of the total variance in the left hemisphere and 70% in the right, depicted a dorso-

medial to ventro-lateral axis organization. The second gradient ($G_{RC}2$), accounting for ~20% of the

explained variance on the left hemisphere and ~15% on the right hemisphere, displayed a medio-

lateral topography. Finally, the third gradient ($G_{RC}3$), capturing approximately 10% of the total

variance on the left and less than 10% on the right, ran along the anterior-posterior axis (Figure 2C).

Figure 2. Functional connectivity, structural connectivity, and receptor coexpression pulvinar gradients. For each panel, the top row shows the scree plots of explained variance for each gradient. The gradients that have been considered for subsequent analyses are marked with red circles. The bottom row shows 3D reconstruction of the normalized gradient images overlaid on the thalamic outline (gray-shaded area). The thalamic volume has been obtained from the AAL3 atlas. Gradients are shown in posterior (left), superior (center), and anterior (right) views. A: anterior; P: posterior; L: left; R: right.

To characterize the relationship between these pulvinar gradients and the conventional anatomical subdivision of the pulvinar complex into discrete nuclei, we analyzed the distribution of gradient values across the four traditional anatomical nuclei (medial, lateral, anterior, inferior) as defined by an atlas-based parcellation based on post-mortem histology (Iglesias et al. 2018). First, by plotting gradient values grouped by each pulvinar nucleus, we observed limited correspondence to discrete

pulvinar nuclei, regardless of the imaging modality. Although some gradients showed a progressing trend along different pulvinar nuclei, most gradient values were evenly distributed across nuclei. To assess if discrete pulvinar nuclear structure could be inferred from the continuous gradient values derived from diffusion embedding, k-means data-driven clustering in gradient space was applied, either by considering the most relevant gradients for each single imaging modality (functional, structural, or receptor coexpression) or by concatenating gradient values from all the modalities. The appropriate number of clusters (4 for functional gradients, 3 for structural gradients, 4 for coexpression gradients, and 8 for combined clustering) was selected based on silhouette score plots for each hemisphere. Clustering of gradient values resulted in generally poor overlap to anatomical nuclei either for single modality (maximum Dice coefficient left/right pulvinar: 0.52/0.49 to medial pulvinar for functional gradients, 0.72/0.66 to medial pulvinar for structural gradients; 0.47/0.46 to medial pulvinar for coexpression gradients) or concatenated modalities (maximum Dice coefficient left/right pulvinar: 0.48/0.46 to anterior pulvinar) (Supplementary Figure 1).

213
214 **Supplementary Figure 1. K-means clustering of pulvinar gradients and their correspondence with histological**
215 **pulvinar nuclei.** Panels to the left show the silhouette plots for left and right pulvinar clustering solutions; error bars are
216 standard errors calculated across 50 resamples. Panels to the right show matrix plots of Dice similarity coefficients of
217 pulvinar clusters against histological nuclei (AAL atlas). INF: inferior; ANT: anterior; LAT: lateral; MED: medial.
218

2. Pulvinar-cortical functional connectivity replicates patterns of cortico-cortical functional

connectivity

Connectivity gradients serve as low-dimensional representations of the topographical organization

patterns of connectivity, whether functional or structural connectivity. With this concept in mind,

we sought to investigate the extent to which pulvinar-cortical connectivity reflects cortico-cortical

connectivity. To achieve this, we generated pulvinar-cortical gradient-weighted connectivity maps

for the most relevant structural and functional gradients. This involved calculating the average of

the dot products of gradient values in each voxel and their connectivity to each cortical ROI.

Subsequently, we correlated these maps with gradients derived from the embedding of cortico-

cortical connectivity. To address the issue of spatial autocorrelation (SA), which commonly affects

brain maps, we computed SA-corrected permutational p-values using a method based on SA-

preserving surrogates (Burt et al. 2020). Furthermore, these SA-corrected p-values were adjusted

for false discovery rate (FDR) using the Benjamini-Hochberg correction method.

Diffusion embedding of cortico-cortical functional connectivity gradients revealed an elbow in the

explained variance graph at the fifth gradient, collectively explaining ~55% of the total variance.

Similarly, the first four gradients of cortico-cortical structural connectivity accounted for ~60% of

the total variance (Supplementary Figure 2-3).

236
237 **Supplementary Figure 2. Cortico-cortical functional connectivity gradients.** The left panel shows the scree plot of
238 explained variance for each gradient, where gradients that have been considered for subsequent analyses (1-5) are
239 marked with red circles. The right panel shows normalized gradient values for gradients 4-5 plotted on the cortical
240 surface; Gradients 1-3 are shown in Figure 3A.

241
We found that pulvinar-cortical connectivity patterns associated with pulvinar gradients exhibited

high correlations with the first three cortico-cortical functional connectivity gradients. Specifically,

$G_{FC}1$-weighted connectivity maps demonstrated the highest correlation with the first cortico-cortical

functional gradient (left pulvinar: r=0.89, p < 0.01; right pulvinar: r=0.83, p < 0.01); in line with the

available evidence, this gradient delineates a unimodal-to-transmodal cortical hierarchy, spanning

from low-level sensory and motor regions to higher-order associative and limbic cortices

(Margulies et al. 2016). $G_{FC}2$-weighted connectivity maps strongly correlated with the third cortico-

cortical functional gradient (left pulvinar: r=0.78, p < 0.01; right pulvinar: r=0.78, p < 0.01); as in

previous investigations, the cortical topography of this gradient spans across cortical regions

involved in low-versus-high complexity cognitive task, thus highlighting a cortical organization

depending on cognitive demand (Turnbull et al. 2020). Finally, $G_{FC}3$-weighted connectivity was

found strongly correlated with the second cortico-cortical functional gradient (left pulvinar: r=0.71,

p < 0.01; right pulvinar: r=0.67, p < 0.01): in line with previous works, this cortical gradient is

anchored on one end on visual processing areas, and on the opposite end on sensorimotor

processing networks, reflecting a secondary cortico-cortical hierarchy that is orthogonal to the

principal cortical gradient (Margulies et al. 2016). Additionally, weaker yet significant correlations

were found between $G_{FC}1$-weighted connectivity maps and the second cortico-cortical functional

gradient (left pulvinar: r=0.27, p=0.01; right pulvinar: r=0.38, p < 0.01) (Figure 3A).

260
Figure 3. A) The relationship between pulvinar-cortical and cortico-cortical functional gradients. The left panels
show gradient-weighted connectivity values for the first three cortico-pulvinar functional connectivity gradients, each

paired with the most correlated cortico-cortical functional connectivity gradient. Normalized values of each cortical
ROI (Schaefer atlas, 400 parcels) are plotted on the cortical surface. The right panels show matrix plots of Pearson's
correlation values between cortico-cortical gradients (y-axis) and pulvinar-cortical gradient-weighted connectivity
maps. Only values showing statistical significance (SA-corrected, FDR-adjusted $p < 0.01$) are shown. **B) The**
relationship between pulvinar-cortical functional gradients and cortical connectivity networks. Violin plots of
normalized gradient-weighted connectivity values grouped by seven intrinsic connectivity networks (as in Yeo et al.,
2011); VN: visual network; SMN: sensorimotor network; DAN: dorsal attention network; VAN: ventral attention
network; LN: limbic network; FPN: frontoparietal network, DMN: default-mode network.
271
On the contrary, gradient-weighted structural connectivity maps did not exhibit the same correlation

pattern with cortico-cortical structural gradient maps, displaying only weaker correlations to cortical

gradients. In both left and right pulvinar, $G_{SC}2$-weighted structural connectivity maps showed weak

correlations with the principal structural connectivity gradient (left pulvinar: r=0.31, p=0.02; right

pulvinar: r=0.28, p < 0.01). Additionally, there was a trend towards significance observed for the

correlation between left $G_{SC}1$-weighted connectivity and the third cortico-cortical gradient (r=0.17,

p=0.02) and between right $G_{SC}2$-weighted connectivity and the third cortico-cortical gradient

(r=0.19, p=0.01) (Supplementary Figure 3).

Supplementary Figure 3. Cortico-cortical structural connectivity gradients and their relationship to pulvinar-cortical structural gradients. The left panel shows cortico-cortical structural gradients; normalized gradient values are plotted on the cortical surface (Schaefer atlas, 400 parcels). The right panel shows scree plots of explained variance, where gradients that have been considered for subsequent analyses (1-4) are marked with red circles. Bottom right: heatmaps of Pearson's correlation values between pulvinar-cortical gradient-weighted connectivity (x-axis) and the corresponding four cortico-cortical structural connectivity gradients (y-axis).

3. The unimodal-to-transmodal gradient ($G_{FC}1$) aligns with receptor expression on the dorso-ventral pulvinar axis

We discovered that the main gradient of cortico-cortical functional connectivity is mirrored on the pulvinar by $G_{FC}1$. The cortical connectivity of this gradient progresses from unimodal cortical areas of the sensorimotor and visual network to multi-modal, higher-order associative areas of the limbic,

frontoparietal, and default mode network (Figure 3B). Similarly, in the pulvinar, this trend

manifests as a progression of gradient values in a ventral-to-dorsal fashion, from the anterior and

inferior to the lateral and medial pulvinar nuclei (Figure 4B).

We further delved into the relationship between multi-scale organization features of the pulvinar by

correlating gradients obtained from embedding data obtained from different imaging modalities. For

this gradient, we found the strongest association with the main gradient of receptor expression $G_{RC}1$

(Figure 4A; left pulvinar: r=-0.71, p < 0.01; right pulvinar: r=-0.78, p < 0.01). It is worth noting

that, due to the intrinsic sign indeterminacy of gradient values, absolute correlation values were

considered. Similarly to $G_{FC}1$, $G_{RC}1$ was also organized along the dorso-ventral axis of the pulvinar

complex and displayed a similar distribution of values across pulvinar nuclei, albeit inverted with

respect to $G_{FC}1$, with loadings progressing from lateral and medial to anterior and inferior pulvinar

nuclei.

To identify the receptor coexpression patterns driving this specific mode of organization, we

examined the values of the five tracers with the strongest correlation to gradient values. We

observed that expression of the serotonin and noradrenaline reuptake transporters (5HTT and NAT)

positively correlated with gradient values. Conversely, the expression of markers of dopaminergic

activity, such as the D2 receptor or the reuptake transporter DAT, as well as NMDA glutamatergic

receptors decreased with increasing gradient values (Figure 4C).

Furthermore, we observed a significant correlation between $G_{FC}1$ and the secondary gradient of

structural connectivity $G_{SC}2$ (left pulvinar: r=0.70, p < 0.01; right pulvinar: r=0.72, p < 0.01), which

explained ~20% of the variance in structural connectivity. Similar to $G_{FC}1$, $G_{SC}2$ follows the dorso-

ventral axis of the pulvinar complex, progressing from lower values in the anterior and lateral nuclei

to higher values in the inferior and medial nuclei (Figure 4C). Consequently, this gradient also

displayed a weaker negative correlation with $G_{RC}1$ (left pulvinar: r= -0.44, p < 0.01; right pulvinar:

r=-0.46, p < 0.01). However, the cortical structural connectivity pattern associated with $G_{FC}1$ does

not mirror that of its functional connectivity counterpart. Instead, gradient-weighted structural

319   connectivity progresses from lower values in modality-independent networks (e.g., default mode,

320   frontoparietal) to intermediate values in unimodal, low-level sensory networks (e.g., visual and

321   sensorimotor) with maximal values in networks involved in higher-order sensory processing (e.g.,

322   limbic, ventral attention, dorsal attention).

323
Figure 4. A) Pulvinar correlates of the unimodal-transmodal cortical gradient. Scatter plots illustrate the
relationship between the first pulvinar-cortical connectivity gradient, corresponding to the unimodal-transmodal
hierarchy of cortico-cortical connectivity, and its most correlated gradient values across other modalities. Due to the
intrinsic sign indeterminacy of gradient values, absolute correlation values are considered. **B) The relationship
between correlated pulvinar gradients and discrete histological nuclei.** Violin plots illustrating normalized gradient
values grouped by histological nuclei (AAL atlas). MED: medial pulvinar; LAT: lateral pulvinar; ANT: anterior
pulvinar; INF: inferior pulvinar. **C) Structural connectivity and receptor coexpression patterns correlating with
the unimodal-transmodal hierarchy.** Left panels: gradient-weighted structural connectivity. Normalized values for
each cortical ROI (Schaefer atlas, 400 parcels) are plotted on the cortical surface. Violin plots show values grouped by
seven intrinsic connectivity networks (as in Yeo et al., 2011). VN: visual network; SMN: sensorimotor network; DAN:
dorsal attention network; VAN: ventral attention network; LN: limbic network; FPN: frontoparietal network; DMN:
default-mode network. Right panels: gradient-correlated receptor density values for the top 5 most correlated receptors.
Details can be found in the main text.

In summary, our findings demonstrated that $G_{FC}1$, $G_{RC}1$, and $G_{SC}2$ substantially delineate

multiscale differences between the ventral and dorsal aspects of the pulvinar. Moving along the

ventral-dorsal axis of the pulvinar complex, more ventral regions showed higher functional

connectivity to unimodal sensory processing networks, higher levels of 5HTT and NAT expression,

and preferentially higher structural connectivity to modality-independent or low-level sensory

processing cortices. Conversely, dorsal regions displayed increasingly higher functional

connectivity to multi-modal processing networks, higher levels of D2, NMDA receptors, and

greater structural connectivity to higher-order sensory processing cortices (Figure 4).

4. The visual-to-sensorimotor gradient ($G_{FC}3$) aligns with structural connectivity on the

medio-lateral pulvinar axis

The secondary gradient of cortico-cortical organization was found to be correlated with $G_{FC}3$,

which explained ~10% of the variance in pulvinar-cortical functional connectivity. $G_{FC}3$ exhibited a

well-delineated medial-lateral and antero-posterior arrangement, corresponding to lower values in

the anterior and lateral pulvinar divisions, and higher values in the inferior and medial nuclei. The

cortical functional connectivity patterns associated with $G_{FC}3$ spanned from the lower extreme of

the sensorimotor network to the higher extreme of the visual network. This gradient reflected

transitions from increasingly visual-related processing networks (such as dorsal attention and limbic

networks) to increasingly action-related processing networks (including the default mode,

frontoparietal, and ventral attention networks) (Figure 5B).

We found that $G_{FC}3$ is the most correlated, in absolute values, to the principal gradient $G_{SC}1$, which

explained ~70% of variance in pulvinar-cortical structural connectivity, showing a strong negative

correlation (Figure 5A; left pulvinar: r= -0.61, p < 0.01; right pulvinar: r=-0.77, p < 0.01). $G_{SC}1$ also

delineates a clear progression along the medio-lateral axis of the pulvinar, with lower values in the

inferior and medial pulvinar nuclei transitioning to higher values in the lateral and anterior pulvinar

nuclei (Figure 5B). On the cortical mantle, the structural gradient-weighted connectivity patterns

exhibited a heterogeneous distribution of values within functional networks, rather than delineating

a specific pattern of progression across functional networks (Figure 5C).

Additionally, we found a significant negative correlation to $G_{RC}2$, which accounts for ~15% of the

variance in receptor expression (left pulvinar: r= -0.47, p < 0.01; right pulvinar: r=-0.72, p < 0.01).

This gradient consistently showed an association in both hemispheres with the expression of mu-

opioid receptors (MOR), which was positively correlated to the gradient axis, and of the 5HTT,

which was negatively correlated to the gradient axis (Figure 5C). Similar to $G_{FC}3$, $G_{RC}2$ was also

found to be correlated with the main gradient of structural connectivity $G_{SC}1$ (Figure 5A; left

pulvinar: r= 0.59, p < 0.01; right pulvinar: r= 0.77, p < 0.01). Like $G_{SC}1$, $G_{RC}2$ exhibited a

progression of values from inferior and medial pulvinar nuclei to the lateral and anterior nuclei

(Figure 5B).

375
Figure 5 A) Pulvinar correlates of the visual-to-sensorimotor cortical gradient. Scatter plots illustrate the
relationship between the third pulvinar-cortical connectivity gradient, corresponding to the visual-to-sensorimotor
hierarchy of cortico-cortical connectivity, and its most correlated gradient values across other modalities. Due to the
intrinsic sign indeterminacy of gradient values, absolute correlation values are considered. **B) The relationship
between correlated pulvinar gradients and discrete histological nuclei**. Violin plots illustrating normalized gradient
values grouped by histological nuclei (AAL atlas). MED: medial pulvinar; LAT: lateral pulvinar; ANT: anterior
pulvinar; INF: inferior pulvinar. **C) Structural connectivity and receptor coexpression patterns correlating with
the visual-to-sensorimotor hierarchy.** Left panels: gradient-weighted structural connectivity. Normalized values for
each cortical ROI (Schaefer atlas, 400 parcels) are plotted on the cortical surface. Violin plots show values grouped by
seven intrinsic connectivity networks (as in Yeo et al., 2011). VN: visual network; SMN: sensorimotor network; DAN:
dorsal attention network; VAN: ventral attention network; LN: limbic network; FPN: frontoparietal network, DMN:
default-mode network. Right panels: gradient-correlated receptor density values for the top 5 most correlated receptors.
Details can be found in the main text.

To summarize, our findings reveal a strong association between the medio-lateral axis of pulvinar

organization, which reflects the progression of functional connectivity from sensorimotor to

associative and visual areas, and the principal gradient of structural connectivity. The cortical

connectivity profiles of this gradient delineate within-network patterns of progression, suggesting

that multiple functional networks host separate representations of the pulvinar complex.

Additionally, this structural and functional connectivity pattern is paralleled by a secondary gradient

of receptor expression. Specifically, there is increasing expression of MOR and decreasing

expression of 5HTT in the medial end of the gradient, which corresponds to regions functionally

connected to sensorimotor-associative regions. Conversely, there is a decreasing expression of

MOR and an increasing expression of 5HTT in the lateral end of the gradient, which corresponds to

regions functionally connected to associative-visual regions (Figure 5).

5. Reliability and reproducibility

In general, diffusion embedding of cortico-pulvinar connectivity data demonstrated high stability

and reproducibility, with structural connectivity gradients exhibiting superior performance

compared to functional connectivity gradients.

The structural connectivity gradients $G_{SC}1$ and $G_{SC}2$ showed excellent stability across 100

resampling of random halves of the main dataset in both hemispheres (all median $r > 0.99$ for both

left and right pulvinar, with IQR < 0.01). Additionally, group-level structural gradients

demonstrated the highest repeatability when compared between test and retest samples of the main

dataset (all $r > 0.99$ for both left and right pulvinar). Finally, structural gradients showed high

reproducibility across both the main and validation datasets (left pulvinar, $G_{SC}1$: $r = 0.87$; $G_{SC}2$: $r =$

0.88; right pulvinar, $G_{SC}1$: $r = 0.87$; $G_{SC}2$: $r = 0.85$).

Functional connectivity gradients showed overall very high or moderate-to-high stability across

split-half resamples of the main dataset in both hemispheres. However, stability decreased with

decreasing variance explained (left pulvinar, $G_{FC}1$: median $r = 0.85$, IQR$= 0.021$; $G_{FC}2$: median $r =$

0.70, IQR $= 0.042$; $G_{FC}3$: median $r = 0.59$, IQR $= 0.061$; right pulvinar, $G_{FC}1$: median $r = 0.87$,

IQR$= 0.023$; $G_{FC}2$: median $r = 0.75$, IQR $= 0.039$; $G_{FC}3$: median $r = 0.72$, IQR $= 0.039$).

Repeatability on test-retest samples of the main dataset also showed a decreasing trend following

the decrease of explained variance, with moderate to good repeatability for $G_{FC}1$ and $G_{FC}2$, and low

repeatability for $G_{FC}3$. In contrast, reproducibility of the group-level gradients across the main and

validation sample, while showing a similar trend with higher values for $G_{FC}1$ compared to the other

secondary gradients, was found high both for left (all r > 0.75) and right pulvinar (all r > 0.80)

(Supplementary Figure 4).

424
425   **Supplementary Figure 4. Reliability and reproducibility of diffusion embedding of pulvinar-cortical connectivity**
426   **data. A) Split-half stability analysis.** Violin plots show the distribution of correlation values across 100 split-half
427   resamples (bars on the median and extreme values)**. B) Test-retest repeatability. C) Reproducibility on the**
428   **supplementary dataset (LEMON).**
429

Discussion

In the present study, we employed a data-driven dimensionality reduction analysis on a

comprehensive array of high-resolution, large-scale multimodal datasets encompassing structural,

functional connectivity, and receptor expression data. Our objective was to delve into the

organizational principles underlying pulvinar-cortical connectivity and its relationship to cortico-

cortical connectivity. Through the application of diffusion embedding, a widely utilized gradient

analysis technique, we successfully collapsed the high-dimensional connectivity and coexpression

data into a reduced set of low-level representations, reflecting the spatial patterns of organization of

multimodal features on the pulvinar complex.

Previous reports on functional or structural connectivity of the human pulvinar complex have been

predominantly either ROI-based and hypothesis-driven (Leh et al. 2008; Tamietto et al. 2012;

Arcaro et al. 2015; Basile et al. 2021) and thus unable to capture finer details regarding the

topographical organization patterns of pulvinar connectivity in its entirety, or data-driven and

focused on hard parcellations from data clustering algorithms (Barron et al.2015; Guedj and

Vuilleumier 2020). The later methodological approach tends to force the topographical

heterogeneity of pulvinar connectivity patterns into discrete, spherical units rather than providing

insights into their continuous transitions in space.

Over the past five decades, the connectivity of the pulvinar complex has undergone through

investigations in non-human primates, utilizing anatomical tract-tracing or electrophysiological

methods (Asanuma et al. 1985; Baleydier and Mauguiere 1985; Pons and Kaas 1985; Cusick and

Gould 1990; Baleydier and Morel 1992; Romanski et al. 1997a; Shipp 2001). A prevailing principle

suggests that connectivity to cortical regions does not adhere to clear-cut distinctions between

histological subdivisions of the pulvinar complex but instead follows a topographical organization

spanning across anatomical nuclei (Shipp 2003). Accordingly, existing connectivity-based

parcellation models of the pulvinar complex have generally shown moderate-to-low correspondence

between functional connectivity or coactivation clusters and histological delineations of pulvinar

nuclei (Barron et al. 2015; Ji et al. 2016; Guedj and Vuilleumier 2020). Our results align with this

perspective by demonstrating that while structural and functional connectional topography often

follows recognizable patterns of progression across anatomical nuclei, hard clustering of either

connectivity or coexpression data failed to reveal the underlying boundaries of nuclear anatomy. In

other words, the topographical pattern of connection and coexpression only partially reflects the

anatomical boundaries outlined by the connectivity or coexpression differences between nuclei.

It is worth noting, however, that connectivity profiles of individual pulvinar nuclei can be properly

inferred from gradient values and their relative gradient-weighted connectivity maps. For instance,

$G_{FC}1$, which showed progression of gradient-weighted connectivity from low-level sensorimotor

and visual processing regions to higher-level multimodal, limbic, and default-mode regions,

demonstrated a corresponding progression of values from anterior (mostly sensorimotor) and

inferior (low-level visual) nuclei to lateral (higher-level visual) and medial (associative and limbic)

pulvinar nuclei. This observation is consistent with existing literature in non-human primates

(Benarroch 2015; Homman-Ludiye and Bourne 2019). In synthesis, the notion of connectivity

gradients reconciles the concept of continuous and graded cortical representations on the pulvinar

complex with the hypothesis of functional and anatomical specialization of discrete pulvinar

subregions.

Drawing from anatomical literature in primates, Shipp (2003) proposed a model of pulvinar-cortical

connectivity, positing a clear dissociation between dorsal and ventral pulvinar, with each region

hosting distinct and parallel representations of cortical regions along their latero-medial axis.

According to this model, connections to the dorsal pulvinar are organized along a dorsal parietal-

superior temporal axis, while connections to the ventral pulvinar follow an occipital-inferior

temporal axis. Subsequent investigations using resting-state fMRI and diffusion tractography in the

human brain have further substantiated this specific organizational pattern (Arcaro et al. 2015,

2018). Findings from connectivity-based parcellation studies substantially agree with this model.

Anterior (and ventral) clusters predominantly exhibit connections to precentral and postcentral gyri;

dorsomedial clusters show connectivity to the cingulum, precuneus and inferior parietal lobules;

dorsal-lateral clusters connected to prefrontal and parietal regions and a ventromedial cluster

displaying connectivity to sensorimotor and occipitotemporal cortical regions (Guedj and

Vuilleumier 2020; Basile et al 2024, under review).

Consistent with these findings, our study reveals that the two major axes of both structural and

functional connectivity converge on the dorso-medial and ventro-lateral axes of the pulvinar

complex, confirming the double dissociation pattern described therein. Ventral regions of the pulvinar (low on $G_{FC}1$) exhibited stronger connections to occipital and inferior temporal regions, while dorsal regions (high on $G_{FC}1$) were predominantly connected to frontoparietal and cingulate regions. Consequently, the secondary gradient of pulvinar-cortical functional connectivity progressed from lower extreme values in the dorsolateral pulvinar (associated with connectivity to frontoparietal regions) to dorsomedial pulvinar (mostly connected to superior temporal cortex), passing through ventromedial and ventrolateral pulvinar (respectively connected to occipital and inferior temporal lobes). Lastly, medial regions of the pulvinar (low on $G_{FC}3$) exhibited stronger connections to the occipital and parietal lobe compared to lateral regions (high on $G_{FC}3$), which were instead more correlated with superior and inferior temporal cortices.

Our findings provide a nuanced perspective on Shipp's model of pulvinar-cortical connectivity, shedding light on the intricate relationships between pulvinar-cortical and cortico-cortical connection patterns. While anatomical studies in non-human primates have suggested that highly connected cortical areas may share overlapping representations on the pulvinar complex, known as the "replication principle" (Shipp 2003), our work offers robust formal evidence of this organizational principle in the human brain. Although we could not replicate the same results for structural connectivity, likely due to the inherent disproportion between pulvinar-cortical connections and the whole brain connectome, we demonstrated a one-to-one correspondence between the first three cortico-cortical functional connectivity gradients and the three main functional connectivity gradients of the pulvinar.

Beyond merely confirming that cortical functional networks converge on shared connectional domains within the pulvinar complex, our findings underscore how pulvinar-cortical connectivity closely mirrors the organizational principles observed in the cerebral cortex. These principles, initially elucidated in the seminal work of Margulies and colleagues (2016) and grounded in earlier anatomical models based on tract-tracing studies in non-human primates (Mesulam 1998), reflect a

hierarchical patterning of information transfer from unimodal sensory or motor representation towards increasing levels of integration and abstraction in multi-modal associative cortices.

Specifically, the principal gradient of cortical connectivity, extending from primary visual, somatosensory, motor, and auditory cortices to associative regions of the limbic, frontoparietal and default-mode network, not only accounted for the largest amount of variance of pulvinar functional connectivity, but also delineated the ventro-dorsal pulvinar axis. Likewise, the gradient capturing the second-most cortical connectivity variance, which spans from visual to motor-auditory unimodal cortices with modality-independent regions situated at various levels, corresponds to the third pulvinar-cortical gradient in terms of explained variance, mapped along the medio-lateral axis. Interestingly, the third gradient of cortico-cortical connectivity, in terms of pulvinar-cortical connectivity appears slightly more represented than this secondary, cross-modal gradient, as indicated by the higher explained variance. This cortical gradient, extending from dorso-lateral to dorso-medial pulvinar regions, progresses from task-negative to task-positive networks of the cerebral cortex and has been linked to spontaneous attention during naturalistic tasks (Samara et al. 2023).

Together, these results reinforce the concept of pulvinar involvement in multimodal sensory integration and top-down attentional processing, consistent with recent literature (Saalmann et al. 2012; Benarroch 2015; Fiebelkorn and Kastner 2020; Froesel et al. 2021). They also align with the proposed role of pulvinar-cortical connectivity in enhancing communication and synchronization within cortical processing modules in a context-dependent and competitive manner (Rockland et al. 1999; Cortes et al. 2021, 2024; Aussel et al. 2023; Bastuji et al. 2024).

Neurochemical correlates of pulvinar-cortical topographical organization

In addition to elucidating the intricate pulvinar-cortical connectivity patterns, our study unveils unprecedented insights into the correlates of this complex topographical organization at multiple organizational scales, including structural connectivity and neurotransmission. Notably, we found

that the hierarchical organization of pulvinar-cortical connectivity aligns with the principal axis of receptor expression, indicating a substantial influence of neuromodulator receptor expression on this level of organization. This principal axis of receptor expression reflects diverging expression patterns of markers of catecholaminergic neurotransmission, encompassing dopaminergic, noradrenergic, and serotonergic systems. While previous ex-vivo and in-vivo investigations in the human and non-human primate brain confirm that pulvinar is a recipient of dopaminergic, noradrenergic, and serotonergic projections (Lavoie and Parent 1991; Oke et al. 1997; Rieck et al. 2004; Sánchez-González et al. 2005; Pérez-Santos et al. 2021), our work provides information on their specific distribution pattern, representing a novel contribution to the field. Moreover, studies in non-human primates have highlighted many receptor markers whose topographic distribution mirrors connectivity patterns in the pulvinar complex, including acetylcholine-esterase (AChE), parvalbumin, calbindin, or vesicular glutamate transporters 1 and 2 (vGLUT1, vGLUT2) (Gutierrez et al. 1995, 2000; Stepniewska and Kaas 1997; Balaram et al. 2013, 2015). Since some of these markers are known to colocalize with catecholaminergic receptors in different brain areas (Liang et al. 1996; Graham et al. 2015), and have been recently correlated to functional topography in the cortex and subcortex, including the thalamus (Anderson et al. 2018, 2020; Müller et al. 2020), the macroscale pattern of receptor coexpression observed by PET may reflect broader chemoarchitectural features at the cellular level. Furthermore, serotonergic neurotransmission in the thalamus plays a crucial role in higher-order sensory integration, especially within the visual system. In this regard, alterations in thalamo-cortical functional connectivity, influenced by powerful serotonergic psychedelics, are known to mediate this process (Preller et al. 2018; Gaddis et al. 2022; Onofrj et al. 2023). This perspective sheds light on the increasing 5HTT expression from high to low-level somatosensory and visual regions of the pulvinar. Conversely, the increasing expression of dopamine receptors and transporters observed in the dorsal, higher-order regions of the pulvinar is consistent with the role of dopamine neuromodulation in mediating attentional processes, as supported by the findings of increased functional connectivity between thalamus and

higher-order cortical regions after administration of the selective DAT-blocker modafinil (Pizzi et al. 2023). Finally, recent multi-modal investigations have reported higher levels of dopamine receptors and transporters in cortical and subcortical regions structurally and functionally connected with the dorsal attention network, including the dorsomedial pulvinar (Alves et al. 2022). This further supports our results and underscores the pivotal role of dopamine neuromodulation in attentional processes.

A multimodal pulvinar gradient architecture as a possible benchmark for neurological disorders

By integrating functional and structural connectivity data with neuroreceptor expression patterns, our proposed model of multi-level gradient architecture of the pulvinar complex holds promise for elucidating alterations in pulvinar structure and connectivity observed in various pathological conditions. For instance, reduced availability of D2/3 receptors in the pulvinar correlated with functional connectivity to the superior temporal sulcus and medial occipital lobe and autistic social communication symptoms in individuals with autistic spectrum disorders (ASD) (Murayama et al. 2022). Furthermore, the right anterior pulvinar (PuA), situated at the extreme of our principal coexpression gradient characterized by high expression of 5HTT and low expression of D2 receptors, exhibited reduced volume in patients with Parkinson's disease and comorbid depression. Notably, treatment with serotonergic antidepressants was associated with increased volume of this region, suggesting a protective effect of serotonergic transmission against dopamine depletion-driven degeneration (Bhome et al. 2022). In another study involving first-episode psychotic patients, heightened functional connectivity to nodes of the default mode network and central executive network was observed in the dorsal pulvinar. This finding aligns with our multimodal model, wherein increased expression of D2 receptors corresponds to this region (Kwak et al. 2021). We suggest that our integrative, gradient-based model of the organizational features of the pulvinar complex may provide an explorative framework to understand the role of this structure in

neuropsychiatric disorders such as Lewy Body Dementia (LBD), Alzheimer's Disease,

frontotemporal dementia (FTD), medial temporal lobe epilepsy (MTLE), or schizophrenia and other

psychotic disorders (Guye 2006; Anticevic et al. 2015; Erskine et al. 2018; Capecchi et al. 2020;

Perez-Rando et al. 2022; Velioglu et al. 2023; Zhang et al. 2023), as well as the factors influencing

pharmacological treatment response.

Finally, the potential to modulate activity in specific cortical functional networks based on the

localization on the main pulvinar axes, as well as its relationships to structural connectivity, offers

valuable insights for therapeutic interventions.

This includes deep brain stimulation of the pulvinar, an emerging area of interest in functional

neurosurgery for epilepsy, which could benefit from a deeper understanding of pulvinar

connectivity and its implications for treatment outcomes (Kalamatianos et al. 2023; Vakilna et al.

2023; Wong et al. 2023). Specifically, the medial (as opposed to the lateral) pulvinar is the desired

target due to its connectivity to the medial temporal lobe. Overall, our model opens avenues for

both understanding the pathophysiology of neurological and psychiatric disorders and developing

targeted therapeutic interventions.

While our study offers valuable insights into the organizational principles of the pulvinar complex,

it is not without limitations. One significant limitation concerns the relatively small size of the

pulvinar complex, if compared to other larger cortical or subcortical structures. Although the

generally high quality of both the main and the validation dataset, including rs-fMRI data (Uğurbil

et al. 2013; Babayan et al. 2019), finer resolution imaging techniques, such as ultra-high field fMRI,

may provide more detailed information about pulvinar topographical organization at a finer scale.

The limitation of small region size is also particularly relevant for the normative PET data, that

have been resampled to the voxel size of $2mm^2$ to match the resolution of BOLD data and suffer

from all the inherent limitations of PET imaging in small brain volumes, such as partial volume

effects or tracer spillout effects (Kanel et al. 2023). In addition, the neurotransmitter normative atlas

is derived from group-level averages of data acquired across different centers. As a consequence,

the receptor density profiles employed to quantify receptor "coexpression" are measured across spatially aligned and group-averaged PET scans of different individuals (Hansen et al. 2022b), since the radioactive nature of PET and SPECT tracers prevents the study of multiple neurotransmitter systems in the same individual for technical and clinical reasons, yet. On the other hand, this neurotransmitter atlas has been extensively validated both by correlation to an independent autoradiography dataset and to ex-vivo gene expression levels measured with microarray techniques (Hansen et al. 2022a). As such, it therefore represents, to the best of our knowledge, the most advanced tool available to investigate the relations between neurotransmitter expression patterns in the human brain in vivo. Indeed, as mentioned above, "gold-standard" reference ex-vivo studies on molecular expression levels of the receptors and transporters analyzed in our study, relative to the pulvinar complex, are substantially sparse, and nearly absent for the human brain. Therefore, further investigations, possibly by employing immunohistochemical or immunofluorescence microscopy in the human brain coupled with highly sampled gene expression probes, are warranted to validate and enhance our understanding of molecular expression in the human pulvinar.

Conclusion

This study offers advanced insights into the interplay between pulvinar-cortical and cortico-cortical connectivity in the human brain, along with its structural and molecular correlates. Our proposed model confirms the hypothesis of a "replication principle", illustrating that the pulvinar complex harbors multiple representations of cortico-cortical functional connectivity organized hierarchically by their information processing significance across its ventro-dorsal and medio-lateral axes. These shared representations traverse pulvinar nuclei continuously, reconciling discrete, individual nuclear connectivity and coactivation patterns with the idea of a gradient-like organization of pulvinar connectivity.

By systematically assessing the relationship between functional, structural and molecular aspects of pulvinar organization, we delineate the topographically organized features of the human pulvinar

and reveal substantial convergence among the main directives of functional connectivity,

anatomical connectivity and receptor coexpression. Our study underscores the pivotal role of the

pulvinar complex in mediating information integration and communication across brain networks at

multiple levels of brain organization. We propose that this comprehensive understanding may offer

a unified explanatory framework to enhance comprehension of its involvement in higher-order

perceptual and cognitive functions both in healthy patients and in various neuropsychiatric diseases.

Materials and Methods

1. *Data acquisition and preprocessing*

1.1. Primary dataset (HCP)

Structural, diffusion-weighted, and resting-state functional MRI data of 210 healthy subjects

(males=92, females=118, age range 22-36 years) were retrieved from the HCP repository

(https://humanconnectome.org). The study protocol was approved by the Washington University in

St. Louis Institutional Review Board (IRB) (Van Essen et al. 2012).

MRI data were acquired on a 3T custom-made Siemens "Connectome Skyra" scanner (Siemens,

Erlangen, Germany), equipped with a Siemens SC72 gradient coil.

High-resolution T1-weighted scans (isotropic voxel size = 0.7 mm) were acquired with the

following parameters: MP-RAGE sequence, TR = 2400 ms, TE = 2.14 ms (Van Essen et al. 2012).

Skull-stripped T1 weighted images were segmented into cortical and subcortical gray matter (GM),

white matter (WM), and cerebrospinal fluid (CSF) using FAST and FIRST FSL's tools (Smith et al.

2004; Patenaude et al. 2011). The MNI-space transformations available as part of the minimally

preprocessed data were employed (FLIRT 12 degrees of freedom affine; FNIRT nonlinear

registration) (Glasser et al. 2013a).

Resting-state functional MRI data (rs-fMRI) (isotropic voxel size = 2mm) were acquired with a

gradient-echo EPI sequence, using the following parameters: TR = 720 ms, TE = 33.1 ms, 1200

frames, ~15 min/run. Data were acquired separately on different days along two different sessions,

each session consisting of a left-to-right (LR) and a right-to-left (RL) phase encoding acquisition (Van Essen et al. 2012; Uğurbil et al. 2013; Smith et al. 2013a). The LR and RL acquisitions of the first session only have been employed in the present work.

Data were retrieved in a minimally preprocessed form which includes artifact and motion correction, registration to 2-mm resolution MNI 152 standard space, high pass temporal filtering (> 2000 s full width at half maximum) (Glasser et al. 2013b), ICA-based denoising with ICA-FIX (Salimi-Khorshidi et al. 2014), and regression of artifacts and motion-related parameters (Smith et al. 2013b). Additionally, the global WM and CSF signal was regressed out to further improve ICA-based denoising results (Plachti et al. 2019).

Multi-shell DWI data (isotropic voxel size = 1.25 mm) were acquired using a single-shot 2D spin-echo multiband Echo Planar Imaging (EPI) sequence (Sotiropoulos et al. 2013). The following acquisition parameters were employed: b-values: 1000, 2000, 3000 mm/s^2; 90 directions per shell; spatial isotropic resolution 1.25 mm. DWI scans were retrieved in a minimally preprocessed form including eddy currents, EPI distortion and motion correction, and cross-modal linear registration of structural and DWI images (Glasser et al. 2013b).

A multi-shell multi-tissue (MSMT) CSD signal modeling was performed to estimate individual response functions in WM, GM, and CSF (Jeurissen et al. 2014). Whole brain, probabilistic tractography (10 million streamlines, iFOD2 algorithm; anatomically constrained tractography) was generated with MRtrix3 default parameters (Tournier et al. 2010; Smith et al. 2012).

1.2. Validation dataset (LEMON)

Structural, diffusion, and rs-fMRI data of 213 healthy subjects (males=138, females=75, age range 20-70 years) were retrieved from the Leipzig Study for Mind-Body-Emotion Interactions (LEMON) dataset (http://fcon_1000.projects.nitrc.org/indi/retro/MPI_LEMON.html). The study protocol was approved by the Ethics Committee of the medical faculty of the University of Leipzig.

MRI data were acquired on a 3T scanner (MAGNETOM Verio, Siemens Healthcare GmbH,

Erlangen, Germany) equipped with a 32-channel head coil was employed for MRI data acquisition.

High-resolution T1-weighted (isotropic voxel size = 1 mm) scans were acquired with the following

parameters: MP-RAGE sequence, TR = 5000 ms, TE = 2.92 ms. Skull-stripped T1 weighted images

were segmented into cortical and subcortical GM, WM, and CSF using FAST and FIRST FSL's

tools (Smith et al. 2004; Patenaude et al. 2011). T1-weighted volumes were also non-linearly

registered to the 1-mm resolution MNI 152 asymmetric template using a FLIRT 12 degrees of

freedom affine transform and FNIRT non-linear registration (Jenkinson and Smith 2001; Jenkinson

et al. 2002; Andersson et al. 2007).

Single-shell DWI data (isotropic voxel size = 1.7 mm) were acquired using a multi-band accelerated

sequence with b-value of 1000, and 60 unique diffusion-encoding directions. Data were

preprocessed according to the dedicated pipeline implemented via the MRtrix3 software (Tournier

et al. 2019), which features denoising using Marchenko-Pastur principal component analysis (MP-

PCA) (Veraart et al. 2016), removal of Gibbs ringing artifacts (Kellner et al. 2016); 3) eddy

currents, distortion, and motion correction (Andersson et al. 2003; Smith et al. 2004; Andersson and

Sotiropoulos 2016) and bias field correction using the N4 algorithm (Tustison et al. 2010). A single-

shell 3-tissue (SS3T) CSD signal modeling was performed using MRtrix3Tissue (Dhollander et al.

2016), a fork of MRtrix3 software. Whole brain, probabilistic tractography (5 million streamlines,

iFOD2 algorithm; anatomically constrained tractography) was generated with default parameters

(Tournier et al. 2010; Smith et al. 2012).

For rs-fMRI data (isotropic voxel size = 2.3 mm), a gradient-echo EPI was acquired with the

following parameters: TR = 1400 ms, TE = 30 ms, 15.30 min/run (Babayan et al. 2019). Data were

obtained in a minimally-preprocessed form, consisting of the following steps: 1) removal of the first

5 volumes to allow for signal equilibration; 2) motion and distortion correction; 3) outlier and

artifact detection (rapidart) and denoising using component-based noise correction (aCompCor); 4)

mean-centering and variance normalization of the time series; 5) spatial normalization to 2-mm

resolution MNI 152 standard space (Babayan et al. 2019; Mendes et al. 2019).

1.3. Positron emission tomography (PET) data

Volumetric 3D PET images were retrieved from the Network Neurosciences Lab (NetNeuroLab)

github page (https:github.com/netneurolab/hansen_receptors/tree/main/data/PET_nifti_images).

Images were collected for 19 distinct neurotransmitter receptors and transporters in a multicentric

data acquisition approach resulting in multiple studies across the world. Images were acquired using

optimized imaging preprocessing strategies for each radioligand (Nørgaard et al. 2019). In all

studies, only healthy participants were scanned, for a total of 1,238 healthy individuals (males=718,

females=520). Full methodologic details about each study, the associated receptor/transporter,

tracers, PET cameras, and modeling methods can be found in (Hansen et al. 2022b). In general, the

measured estimates for each image (binding potential and/or tracer distribution volume) are

proportional to receptor density and then, we will refer to them as "receptor density maps" for

simplicity.

Receptor density maps, originally available at different resolutions, were resliced to the 2mm

MNI152 standard template resolution and converted to z-scores. Four receptor density maps

(5HT1b, D2, mGluR5, vAChT) were acquired using the same tracer in different studies and were

converted to a weighted average as in the reference paper. Ten receptor/transmitters (5HT1a,

5HT1b, 5HTT, CB1, D2, DAT, GABA-A, MOR and NET) were acquired with different tracers in

different studies; these receptor maps were not averaged but kept as separated, resulting in a total of

29 receptor density maps. Supplementary Material 1 summarizes details about the neurotransmitter,

tracers, and number of subjects for each study.

2. Diffusion map embedding

We derived functional connectivity, structural connectivity, and receptor expression gradients for the left and right pulvinar separately. All the analyses were performed in standard space (MNI152 2006 template, 2 mm voxel size). Voxels belonging to the left or right pulvinar were defined according to the Automated Anatomical Labeling v3 Atlas (AAL3, 2mm version) (Rolls et al. 2020), while cortical regions of interest (ROIs) were derived from the 400-areas version of the local-global cortical parcellation proposed by Schaefer and colleagues (Schaefer et al. 2018).

Individual functional connectomes were obtained from Pearson's correlation of BOLD signal in each pulvinar voxel to the average signal within each cortical ROI. As in similar work, negative correlation values were zeroed and sparsity sampling was applied by row-wise thresholding of top 10% connectivity values, with all the values below the 90[th] percentile set to 0 (Margulies et al. 2016; Katsumi et al. 2023). Functional connectivity matrices were then normalized with Fisher's r-to-z transformation and averaged across all subjects (and across LR and RL sessions, for the HCP dataset) to obtain a group-level dense functional connectome.

Structural connectomes were obtained after registration of each whole-brain tractogram to the MNI space. Streamlines connecting the pulvinar to each cortical ROI were extracted from the whole brain tractogram and mapped back to their pulvinar endpoint voxels using the *tckmap* command in MRtrix3, to obtain individual, voxel-wise pulvinar tract-density distribution maps. Such individual maps were concatenated for all cortical ROIs, converted to voxel-to-cortical ROI tract-density matrices, and averaged across all subjects to obtain a group-level dense structural connectome.

Finally, the normalized receptor density was sampled for each voxel within the pulvinar by concatenating This process resulted in a final pulvinar voxel-by-receptor density matrix, representing the expression levels of 19 neurotransmitters or receptors for each voxel within the pulvinar. This matrix will be referred to as the "coexpression matrix".

We applied diffusion embedding (Coifman and Lafon 2006), an unsupervised learning algorithm widely employed in the field of gradient mapping (Guell et al. 2018; Tian et al. 2020; Hong et al. 2020; Katsumi et al. 2023). Following previous work, each group's asymmetric feature matrix

(functional dense connectome, structural dense connectome, coexpression matrix) was converted to a symmetric voxel-by-voxel cosine distance similarity matrix (1-minus-cosine distance). For diffusion embedding, an alpha value of 0.5 was employed to maximize robustness to noise. The source code for this method is available at https://github.com/satra/mapalign. Eigenvalues were utilized to quantify the variance explained by each gradient, and scree plots of explained variance against the number of gradients were employed to select the most appropriate number of gradients for each side (left or right) and modality. To address the intrinsic sign indeterminacy of gradient representations, right-sided gradients were sign-flipped to align with their left-sided counterparts if needed.

Functional and structural connectivity gradients were also obtained from cortico-cortical connectivity matrices and employed for further analysis. Functional connectivity was computed as the pairwise Pearson's correlation between the average time series of the 400 cortical ROIs. For structural connectivity gradient embedding, the raw number of streamlines connecting each pair of cortical ROIs was extracted from the whole brain tractogram separately for the left and right hemispheres, to emphasize intra-hemispheric patterns of connectivity. Right hemisphere gradients were sign-flipped to align with their left hemisphere counterparts if necessary. The diffusion embedding analysis was performed using the same methods and parameters as for the pulvinar gradients. A graphical overview of the gradient mapping protocol is presented in Figure 1.

3. Gradient analysis and statistics

To facilitate the interpretation of their significance, we performed post-hoc characterization of each pulvinar gradient at various levels.

Firstly, we assessed the involvement of discrete pulvinar nuclei in gradient organization by calculating the distribution of gradient values for each nucleus. Histological nuclei were identified using the AAL atlas, which incorporates digitalized parcellation of thalamic nuclei derived from post-mortem histology (Iglesias et al. 2018). Furthermore, to explore whether the continuous

gradient architecture derived from diffusion embedding could be mapped to discrete units overlapping with pulvinar nuclei, we employed a k-means clustering strategy on gradient values, as described in previous work (Guell et al. 2020). The relevant gradients to be included in the clustering analysis were determined by identifying the elbow in their explained variance percentage graph (as in the paragraph below). For both the left and right pulvinar, the relevant gradients were then normalized within a range from 0 to 1 and concatenated. Various iterations of k-means clustering were then applied in the resulting gradient space, ranging from k=2 to k=30, and the silhouette coefficient (Rousseeuw 1987) was utilized to identify the optimal number of clusters.

Gradient clustering was performed separately for left and right pulvinar, both after concatenating relevant gradients from each single imaging modality alone (functional connectivity, structural connectivity, receptor coexpression) or by concatenating together the most relevant gradients from all the imaging modalities (combined clustering). For each single-modality and combined cluster, the similarity to anatomical pulvinar nuclei was quantified using the Dice similarity coefficient (Dice 1945).

To analyze the functional or structural connectivity gradients, we obtained gradient-weighted cortical connectivity maps by multiplying each voxel's connectivity to the 400 cortical ROIs by the corresponding gradient value. These voxel-wise, gradient-weighted pulvinar-cortical connectivity maps were then averaged to derive a single cortical projection for each pulvinar connectivity gradient.

To further understand the involvement of cortical networks in each of the cortical projections of structural and functional cortical gradients, we examined the distribution of gradient-weighted connectivity values across the seven major functional cortical networks, as defined in Thomas Yeo et al. 2011).

Additionally, we explored the relationship between pulvinar-cortical and cortico-cortical connectivity gradients by calculating Pearson's correlation for each gradient-weighted functional connectivity map with each cortico-cortical gradient map.

Receptor coexpression gradients were characterized by correlating voxel-wise gradient values to normalized receptor density distributions.

Finally, to investigate the relation between pulvinar gradients obtained from different modalities, we computed the pairwise Pearson's correlation between gradients obtained with different modalities.

To account for the spatial autocorrelation (SA) properties of gradient maps, for all the correlations described, statistical significance was assessed using the permutational approach described in Burt et al. (2020). Briefly, this method takes as input geometric distance matrices for SA estimation and involves the generation of a given number of SA-preserving permuted surrogate maps, which are then employed as nulls to estimate a permutational null distribution of the test statistic (Burt et al. 2020). Pairwise Euclidean distances between left or right pulvinar voxel coordinates were employed for pulvinar null models, while for cortical parcellated connectivity data Euclidean distances were estimated between centroids of each cortical ROI. In both cases, 1000 surrogates were generated to estimate the null distribution. Statistical tests were controlled for false discovery rate (FDR) using Benjamini and Hochberg's correction.

4. Reliability and reproducibility assessment

We conducted several analyses to assess the robustness and consistency of our main findings. Specifically, both functional and structural connectivity gradients underwent split-half and test-retest replicability analysis, as well as reproducibility analysis.

For split-half replicability analysis, the main dataset was segmented into two halves, with 105 subjects each. Subjects were randomly assigned to each half, and 100 instances of the shuffling algorithm were performed. In each iteration, individual connectivity matrices of each half were averaged into dense connectomes and underwent diffusion embedding. The average Pearson's correlation for each gradient across all the iterations was used as a reliability measure.

Test-retest replicability was based on an extension of the HCP dataset, consisting of 44 subjects

(males = 13; females = 31; age range: 22–36 years) with available test-retest structural and diffusion

acquisition. The dense structural and functional connectomes obtained from the test and retest

datasets were compared using Pearson's correlation as a measure of similarity.

Finally, reproducibility was determined by recomputing results in the validation dataset and

correlating pulvinar gradients between the primary and replication datasets.

In all the mentioned cases, gradients were realigned before correlation by using Procrustes' analysis

to obtain more reliable estimates of gradient similarity (Hong et al. 2020). This approach ensured

that the gradients from different datasets were aligned properly before assessing their similarity,

enhancing the robustness of our analyses.

References

Adams MM, Hof PR, Gattass R, et al (2000) Visual cortical projections and chemoarchitecture of
macaque monkey pulvinar. J Comp Neurol 419:377–393. https://doi.org/10.1002/(SICI)1096-
9861(20000410)419:3<377::AID-CNE9>3.0.CO;2-E
Alves PN, Forkel SJ, Corbetta M, Thiebaut de Schotten M (2022) The subcortical and
neurochemical organization of the ventral and dorsal attention networks. Commun Biol
5:1343. https://doi.org/10.1038/s42003-022-04281-0
Anderson KM, Collins MA, Chin R, et al (2020) Transcriptional and imaging-genetic association of
cortical interneurons, brain function, and schizophrenia risk. Nat Commun 11:2889.
https://doi.org/10.1038/s41467-020-16710-x
Anderson KM, Krienen FM, Choi EY, et al (2018) Gene expression links functional networks across
cortex and striatum. Nat Commun 9:1428. https://doi.org/10.1038/s41467-018-03811-x
Andersson JLR, Jenkinson M, Smith S, Andersson J (2007) FNIRT — FMRIB' non-linear image
registration tool
Andersson JLR, Skare S, Ashburner J (2003) How to correct susceptibility distortions in spin-echo
echo-planar images: application to diffusion tensor imaging. Neuroimage 20:870–888.
https://doi.org/10.1016/S1053-8119(03)00336-7
Andersson JLR, Sotiropoulos SN (2016) An integrated approach to correction for off-resonance
effects and subject movement in diffusion MR imaging. Neuroimage 125:1063–1078.
https://doi.org/10.1016/j.neuroimage.2015.10.019
Anticevic A, Haut K, Murray JD, et al (2015) Association of thalamic dysconnectivity and conversion
to psychosis in youth and young adults at elevated clinical risk. JAMA Psychiatry.
https://doi.org/10.1001/jamapsychiatry.2015.0566
Arcaro MJ, Pinsk MA, Chen J, Kastner S (2018) Organizing principles of pulvino-cortical functional
coupling in humans. Nat Commun. https://doi.org/10.1038/s41467-018-07725-6
Arcaro MJ, Pinsk MA, Kastner S (2015) The anatomical and functional organization of the human
visual pulvinar. Journal of Neuroscience. https://doi.org/10.1523/JNEUROSCI.1575-14.2015

Asanuma C, Andersen RA, Cowan WM (1985) The thalamic relations of the caudal inferior parietal
lobule and the lateral prefrontal cortex in monkeys: Divergent cortical projections from cell
clusters in the medial pulvinar nucleus. Journal of Comparative Neurology 241:357–381.
https://doi.org/10.1002/cne.902410309
Aussel A, Fiebelkorn IC, Kastner S, et al (2023) Interacting rhythms enhance sensitivity of target
detection in a fronto-parietal computational model of visual attention. Elife 12:.
https://doi.org/10.7554/eLife.67684
Babayan A, Erbey M, Kumral D, et al (2019) A mind-brain-body dataset of MRI, EEG, cognition,
emotion, and peripheral physiology in young and old adults. Sci Data 6:180308.
https://doi.org/10.1038/sdata.2018.308
Bajada CJ, Jackson RL, Haroon HA, et al (2017) A graded tractographic parcellation of the temporal
lobe. Neuroimage 155:503–512. https://doi.org/10.1016/J.NEUROIMAGE.2017.04.016
Balaram P, Hackett TA, Kaas JH (2013) Differential expression of vesicular glutamate transporters 1
and 2 may identify distinct modes of glutamatergic transmission in the macaque visual
system. J Chem Neuroanat 50–51:21–38. https://doi.org/10.1016/j.jchemneu.2013.02.007
Balaram P, Isaamullah M, Petry HM, et al (2015) Distributions of vesicular glutamate transporters
1 and 2 in the visual system of tree shrews (Tupaia belangeri). Journal of Comparative
Neurology 523:1792–1808. https://doi.org/10.1002/cne.23727
Baleydier C, Mauguiere F (1985) Anatomical evidence for medial pulvinar connections with the
posterior cingulate cortex, the retrosplenial area, and the posterior parahippocampal gyrus in
monkeys. Journal of Comparative Neurology. https://doi.org/10.1002/cne.902320207
Baleydier C, Morel A (1992) Segregated thalamocortical pathways to inferior parietal and
inferotemporal cortex in macaque monkey. Vis Neurosci 8:391–405.
https://doi.org/10.1017/S0952523800004922
Barron DS, Eickhoff SB, Clos M, Fox PT (2015) Human pulvinar functional organization and
connectivity. Hum Brain Mapp 36:2417–2431. https://doi.org/10.1002/hbm.22781
Basile GA, Bertino S, Bramanti A, et al (2021) In Vivo Super-Resolution Track-Density Imaging for
Thalamic Nuclei Identification. Cerebral Cortex. https://doi.org/10.1093/cercor/bhab184
Bastuji H, Cadic-Melchior A, Ruelle-Le Glaunec L, et al (2024) Functional connectivity between
medial pulvinar and cortical networks as a predictor of arousal to noxious stimuli during
sleep. European Journal of Neuroscience 59:570–583. https://doi.org/10.1111/ejn.15958
Benarroch EE (2015) Pulvinar: Associative role in cortical function and clinical correlations.
Neurology 84:738–747. https://doi.org/10.1212/WNL.0000000000001276
Bhome R, Zarkali A, Thomas GEC, et al (2022) Thalamic white matter macrostructure and subnuclei
volumes in Parkinson's disease depression. NPJ Parkinsons Dis 8:2.
https://doi.org/10.1038/s41531-021-00270-y
Bourgeois A, Guedj C, Carrera E, Vuilleumier P (2020) Pulvino-cortical interaction: An integrative
role in the control of attention. Neurosci Biobehav Rev 111:104–113.
https://doi.org/10.1016/j.neubiorev.2020.01.005
Burt JB, Helmer M, Shinn M, et al (2020) Generative modeling of brain maps with spatial
autocorrelation. Neuroimage 220:117038.
https://doi.org/10.1016/j.neuroimage.2020.117038
Capecchi F, Mothersill I, Imbach LL (2020) The medial pulvinar as a subcortical relay in temporal
lobe status epilepticus. Seizure 81:276–279. https://doi.org/10.1016/j.seizure.2020.08.016
Coifman RR, Lafon S (2006) Diffusion maps. Appl Comput Harmon Anal 21:5–30.
https://doi.org/10.1016/j.acha.2006.04.006
Cortes N, Abbas Farishta R, Ladret HJ, Casanova C (2021) Corticothalamic Projections Gate Alpha
Rhythms in the Pulvinar. Front Cell Neurosci 15:. https://doi.org/10.3389/fncel.2021.787170

Cortes N, Ladret HJ, Abbas-Farishta R, Casanova C (2024) The pulvinar as a hub of visual processing
and cortical integration. Trends Neurosci 47:120–134.
https://doi.org/10.1016/j.tins.2023.11.008
Cusick CG, Gould HJ (1990) Connections between area 3b of the somatosensory cortex and
subdivisions of the ventroposterior nuclear complex and the anterior pulvinar nucleus in
squirrel monkeys. Journal of Comparative Neurology 292:83–102.
https://doi.org/10.1002/cne.902920106
Dhollander T, Raffelt D, Connelly A (2016) Unsupervised 3-tissue response function estimation
from single-shell or multi-shell diffusion MR data without a co-registered T1 image. ISMRM
Workshop on Breaking the Barriers of Diffusion MRI
Dice LR (1945) Measures of the Amount of Ecologic Association Between Species. Ecology.
https://doi.org/10.2307/1932409
Erskine D, Ding J, Thomas AJ, et al (2018) Molecular changes in the absence of severe pathology in
the pulvinar in dementia with Lewy bodies. Movement Disorders 33:982–991.
https://doi.org/10.1002/mds.27333
Fang P-C, Stepniewska I, Kaas JH (2006) The thalamic connections of motor, premotor, and
prefrontal areas of cortex in a prosimian primate (Otolemur garnetti). Neuroscience 143:987–
1020. https://doi.org/10.1016/j.neuroscience.2006.08.053
Fiebelkorn IC, Kastner S (2020) Functional Specialization in the Attention Network. Annu Rev
Psychol 71:221–249. https://doi.org/10.1146/annurev-psych-010418-103429
Froesel M, Cappe C, Ben Hamed S (2021) A multisensory perspective onto primate pulvinar
functions. Neurosci Biobehav Rev 125:231–243.
https://doi.org/10.1016/j.neubiorev.2021.02.043
Gaddis A, Lidstone DE, Nebel MB, et al (2022) Psilocybin induces spatially constrained alterations
in thalamic functional organizaton and connectivity. Neuroimage 260:119434.
https://doi.org/10.1016/j.neuroimage.2022.119434
Glasser MF, Sotiropoulos SN, Wilson JA, et al (2013a) The minimal preprocessing pipelines for the
Human Connectome Project. Neuroimage 80:105–124.
https://doi.org/10.1016/j.neuroimage.2013.04.127
Glasser MF, Sotiropoulos SN, Wilson JA, et al (2013b) The minimal preprocessing pipelines for the
Human Connectome Project. Neuroimage 80:105–124.
https://doi.org/10.1016/j.neuroimage.2013.04.127
Graham DL, Durai HH, Garden JD, et al (2015) Loss of Dopamine D2 Receptors Increases
Parvalbumin-Positive Interneurons in the Anterior Cingulate Cortex. ACS Chem Neurosci
6:297–305. https://doi.org/10.1021/cn500235m
Guedj C, Vuilleumier P (2020) Functional connectivity fingerprints of the human pulvinar: Decoding
its role in cognition. Neuroimage. https://doi.org/10.1016/j.neuroimage.2020.117162
Guell X, D'Mello AM, Hubbard NA, et al (2020) Functional Territories of Human Dentate Nucleus.
Cerebral Cortex 30:2401–2417. https://doi.org/10.1093/cercor/bhz247
Guell X, Schmahmann JD, Gabrieli J DE, Ghosh SS (2018) Functional gradients of the cerebellum.
Elife 7:. https://doi.org/10.7554/eLife.36652
Gutierrez C, Cola MG, Seltzer B, Cusick C (2000) Neurochemical and connectional organization of
the dorsal pulvinar complex in monkeys. J Comp Neurol 419:61–86.
https://doi.org/10.1002/(SICI)1096-9861(20000327)419:1<61::AID-CNE4>3.0.CO;2-I
Gutierrez C, Cusick CG (1997) Area V1 in macaque monkeys projects to multiple histochemically
defined subdivisions of the inferior pulvinar complex. Brain Res 765:349–356.
https://doi.org/10.1016/S0006-8993(97)00696-3

Gutierrez C, Yaun A, Cusick CG (1995) Neurochemical subdivisions of the inferior pulvinar in
macaque monkeys. J Comp Neurol 363:545–562. https://doi.org/10.1002/cne.903630404
Guye M (2006) The role of corticothalamic coupling in human temporal lobe epilepsy. Brain
129:1917–1928. https://doi.org/10.1093/brain/awl151
Haak K V., Marquand AF, Beckmann CF (2018) Connectopic mapping with resting-state fMRI.
Neuroimage 170:83–94. https://doi.org/10.1016/j.neuroimage.2017.06.075
Hansen JY, Markello RD, Tuominen L, et al (2022a) Correspondence between gene expression and
neurotransmitter receptor and transporter density in the human brain. Neuroimage
264:119671. https://doi.org/10.1016/j.neuroimage.2022.119671
Hansen JY, Shafiei G, Markello RD, et al (2022b) Mapping neurotransmitter systems to the
structural and functional organization of the human neocortex. Nat Neurosci 25:1569–1581.
https://doi.org/10.1038/s41593-022-01186-3
Homman-Ludiye J, Bourne JA (2019) The medial pulvinar: function, origin and association with
neurodevelopmental disorders. J Anat 235:507–520. https://doi.org/10.1111/joa.12932
Hong S-J, Xu T, Nikolaidis A, et al (2020) Toward a connectivity gradient-based framework for
reproducible biomarker discovery. Neuroimage 223:117322.
https://doi.org/10.1016/j.neuroimage.2020.117322
Iglesias JE, Insausti R, Lerma-Usabiaga G, et al (2018) A probabilistic atlas of the human thalamic
nuclei combining ex vivo MRI and histology. Neuroimage 183:314–326.
https://doi.org/10.1016/j.neuroimage.2018.08.012
Jaramillo J, Mejias JF, Wang X-J (2019) Engagement of Pulvino-cortical Feedforward and Feedback
Pathways in Cognitive Computations. Neuron 101:321-336.e9.
https://doi.org/10.1016/j.neuron.2018.11.023
Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved optimization for the robust and
accurate linear registration and motion correction of brain images. Neuroimage
Jenkinson M, Smith S (2001) A global optimisation method for robust affine registration of brain
images. Med Image Anal. https://doi.org/10.1016/S1361-8415(01)00036-6
Jeurissen B, Tournier JD, Dhollander T, et al (2014) Multi-tissue constrained spherical
deconvolution for improved analysis of multi-shell diffusion MRI data. Neuroimage.
https://doi.org/10.1016/j.neuroimage.2014.07.061
Ji B, Li Z, Li K, et al (2016) Dynamic thalamus parcellation from resting-state fMRI data. Hum Brain
Mapp 37:954–967. https://doi.org/10.1002/hbm.23079
Kalamatianos T, Mavrovounis G, Skouras P, et al (2023) Medial Pulvinar Stimulation in Temporal
Lobe Epilepsy: A Literature Review and a Hypothesis Based on Neuroanatomical Findings.
Cureus. https://doi.org/10.7759/cureus.35772
Kanel P, Carli G, Vangel R, et al (2023) Challenges and innovations in brain PET analysis of
neurodegenerative disorders: a mini-review on partial volume effects, small brain region
studies, and reference region selection. Front Neurosci 17:.
https://doi.org/10.3389/fnins.2023.1293847
Katsumi Y, Zhang J, Chen D, et al (2023) Correspondence of functional connectivity gradients
across human isocortex, cerebellum, and hippocampus. Commun Biol 6:.
https://doi.org/10.1038/s42003-023-04796-0
Kellner E, Dhital B, Kiselev VG, Reisert M (2016) Gibbs-ringing artifact removal based on local
subvoxel-shifts. Magn Reson Med 76:1574–1581. https://doi.org/10.1002/mrm.26054
Kwak Y Bin, Cho KIK, Hwang WJ, et al (2021) Mapping thalamocortical functional connectivity with
large-scale brain networks in patients with first-episode psychosis. Sci Rep 11:19815.
https://doi.org/10.1038/s41598-021-99170-7

Lavoie B, Parent A (1991) Serotoninergic innervation of the thalamus in the primate: An immunohistochemical study. Journal of Comparative Neurology 312:1–18. https://doi.org/10.1002/cne.903120102

Leh SE, Chakravarty MM, Ptito A (2008) The Connectivity of the Human Pulvinar: A Diffusion Tensor Imaging Tractography Study. Int J Biomed Imaging 2008:1–5. https://doi.org/10.1155/2008/789539

Liang C-L, Sinton CM, German DC (1996) Midbrain dopaminergic neurons in the mouse: co-localization with Calbindin-D28k and calretinin. Neuroscience 75:523–533. https://doi.org/10.1016/0306-4522(96)00228-X

Lyon DC, Kaas JH (2002) Evidence from V1 connections for both dorsal and ventral subdivisions of V3 in three species of new world monkeys. J Comp Neurol 449:281–297. https://doi.org/10.1002/cne.10297

Lysakowski A, Standage GP, Benevento LA (1988) An investigation of collateral projections of the dorsal lateral geniculate nucleus and other subcortical structures to cortical areas V1 and V4 in the macaque monkey: A double label retrograde tracer study. Exp Brain Res 69:. https://doi.org/10.1007/BF00247317

Mai JK, Majtanik M (2019) Toward a common terminology for the Thalamus. Front Neuroanat

Margulies DS, Ghosh SS, Goulas A, et al (2016) Situating the default-mode network along a principal gradient of macroscale cortical organization. Proceedings of the National Academy of Sciences 113:12574–12579. https://doi.org/10.1073/pnas.1608282113

Mendes N, Oligschläger S, Lauckner ME, et al (2019) A functional connectome phenotyping dataset including cognitive state and personality measures. Sci Data 6:180307. https://doi.org/10.1038/sdata.2018.307

Mesulam M (1998) From sensation to cognition. Brain 121:1013–1052. https://doi.org/10.1093/brain/121.6.1013

Müller EJ, Munn B, Hearne LJ, et al (2020) Core and matrix thalamic sub-populations relate to spatio-temporal cortical connectivity gradients. Neuroimage 222:. https://doi.org/10.1016/J.NEUROIMAGE.2020.117224

Murayama C, Iwabuchi T, Kato Y, et al (2022) Extrastriatal dopamine D2/3 receptor binding, functional connectivity, and autism socio-communicational deficits: a PET and fMRI study. Mol Psychiatry 27:2106–2113. https://doi.org/10.1038/s41380-022-01464-3

Nørgaard M, Ganz M, Svarer C, et al (2019) Optimization of preprocessing strategies in Positron Emission Tomography (PET) neuroimaging: A [11C]DASB PET study. Neuroimage 199:466–479. https://doi.org/10.1016/j.neuroimage.2019.05.055

Oke AF, Carver LA, Gouvion CM, Adams RN (1997) Three-dimensional mapping of norepinephrine and serotonin in human thalamus. Brain Res 763:69–78. https://doi.org/10.1016/S0006-8993(97)00404-6

Olszewski J, Baxter D (1982) Cytoarchitecture of the human brain stem, 2nd editio. Karger, New York

Onofrj M, Russo M, Delli Pizzi S, et al (2023) The central role of the Thalamus in psychosis, lessons from neurodegenerative diseases and psychedelics. Transl Psychiatry 13:384. https://doi.org/10.1038/s41398-023-02691-0

Paquola C, Benkarim O, Dekraker J, et al (2020) Convergence of cortical types and functional motifs in the human mesiotemporal lobe. Elife 9:1–26. https://doi.org/10.7554/ELIFE.60673

Paquola C, Bethlehem RA, Seidlitz J, et al (2019a) Shifts in myeloarchitecture characterise adolescent development of cortical gradients. Elife 8:. https://doi.org/10.7554/ELIFE.50482

Paquola C, Vos De Wael R, Wagstyl K, et al (2019b) Microstructural and functional gradients are
increasingly dissociated in transmodal cortices. PLoS Biol 17:.
https://doi.org/10.1371/JOURNAL.PBIO.3000284
Patenaude B, Smith SM, Kennedy DN, Jenkinson M (2011) A Bayesian model of shape and
appearance for subcortical brain segmentation. Neuroimage 56:907–922.
https://doi.org/10.1016/j.neuroimage.2011.02.046
Perez-Rando M, Elvira UKA, García-Martí G, et al (2022) Alterations in the volume of thalamic
nuclei in patients with schizophrenia and persistent auditory hallucinations. Neuroimage Clin
35:103070. https://doi.org/10.1016/j.nicl.2022.103070
Pérez-Santos I, Palomero-Gallagher N, Zilles K, Cavada C (2021) Distribution of the Noradrenaline
Innervation and Adrenoceptors in the Macaque Monkey Thalamus. Cerebral Cortex 31:4115–
4139. https://doi.org/10.1093/cercor/bhab073
Pizzi SD, Sestieri C, Chiacchiaretta P, et al (2023) Modafinil-induced changes in the functional
thalamic connectivity. Neuroscience Applied 2:103876.
https://doi.org/10.1016/j.nsa.2023.103876
Plachti A, Eickhoff SB, Hoffstaedter F, et al (2019) Multimodal Parcellations and Extensive
Behavioral Profiling Tackling the Hippocampus Gradient. Cerebral Cortex 29:4595–4612.
https://doi.org/10.1093/cercor/bhy336
Pons TP, Kaas JH (1985) Connections of area 2 of somatosensory cortex with the anterior pulvinar
and subdivisions of the ventroposterior complex in macaque monkeys. Journal of
Comparative Neurology 240:16–36. https://doi.org/10.1002/cne.902400103
Preller KH, Burt JB, Ji JL, et al (2018) Changes in global and thalamic brain connectivity in LSD-
induced altered states of consciousness are attributable to the 5-HT2A receptor. Elife 7:.
https://doi.org/10.7554/eLife.35082
Przeździk I, Faber M, Fernández G, et al (2019) The functional organisation of the hippocampus
along its long axis is gradual and predicts recollection. Cortex 119:324–335.
https://doi.org/10.1016/J.CORTEX.2019.04.015
Rieck RW, Ansari MS, Whetsell WO, et al (2004) Distribution of Dopamine D2-Like Receptors in the
Human Thalamus: Autoradiographic and PET Studies. Neuropsychopharmacology 29:362–
372. https://doi.org/10.1038/sj.npp.1300336
Rockland KS, Andresen J, Cowie RJ, Robinson DL (1999) Single axon analysis of pulvinocortical
connections to several visual areas in the Macaque. J Comp Neurol 406:221–250.
https://doi.org/10.1002/(SICI)1096-9861(19990405)406:2<221::AID-CNE7>3.0.CO;2-K
Rolls ET, Huang C-C, Lin C-P, et al (2020) Automated anatomical labelling atlas 3. Neuroimage
206:116189. https://doi.org/10.1016/j.neuroimage.2019.116189
Romanski LM, Giguere M, Bates JF, Goldman-Rakic PS (1997a) Topographic organization of medial
pulvinar connections with the prefrontal cortex in the rhesus monkey. J Comp Neurol
379:313–32
Romanski LM, Giguere M, Bates JF, Goldman-Rakic PS (1997b) Topographic organization of medial
pulvinar connections with the prefrontal cortex in the rhesus monkey. J Comp Neurol
379:313–332. https://doi.org/10.1002/(SICI)1096-9861(19970317)379:3<313::AID-
CNE1>3.0.CO;2-6
Rousseeuw PJ (1987) Silhouettes: A graphical aid to the interpretation and validation of cluster
analysis. J Comput Appl Math 20:53–65. https://doi.org/10.1016/0377-0427(87)90125-7
Saalmann YB, Pinsk MA, Wang L, et al (2012) The pulvinar regulates information transmission
between cortical areas based on attention demands. Science (1979) 337:753–756.
https://doi.org/10.1126/science.1223082

Salimi-Khorshidi G, Douaud G, Beckmann CF, et al (2014) Automatic denoising of functional MRI
data: Combining independent component analysis and hierarchical fusion of classifiers.
Neuroimage 90:449–468. https://doi.org/10.1016/j.neuroimage.2013.11.046
Samara A, Eilbott J, Margulies DS, et al (2023) Cortical gradients during naturalistic processing are
hierarchical and modality-specific. Neuroimage 271:120023.
https://doi.org/10.1016/j.neuroimage.2023.120023
Sánchez-González MÁ, García-Cabezas MÁ, Rico B, Cavada C (2005) The Primate Thalamus Is a Key
Target for Brain Dopamine. The Journal of Neuroscience 25:6076–6083.
https://doi.org/10.1523/JNEUROSCI.0968-05.2005
Schaefer A, Kong R, Gordon EM, et al (2018) Local-Global Parcellation of the Human Cerebral
Cortex from Intrinsic Functional Connectivity MRI. Cerebral Cortex 28:3095–3114.
https://doi.org/10.1093/cercor/bhx179
Shipp S (2001) Corticopulvinar connections of areas V5, V4, and V3 in the macaque monkey: A dual
model of retinal and cortical topographies. Journal of Comparative Neurology.
https://doi.org/10.1002/cne.1363
Shipp S (2003) The functional logic of cortico-pulvinar connections. Philosophical Transactions of
the Royal Society B: Biological Sciences 358:1605–1624.
https://doi.org/10.1098/rstb.2002.1213
Smith RE, Tournier J-D, Calamante F, Connelly A (2012) Anatomically-constrained tractography:
Improved diffusion MRI streamlines tractography through effective use of anatomical
information. Neuroimage 62:1924–1938. https://doi.org/10.1016/j.neuroimage.2012.06.005
Smith SM, Beckmann CF, Andersson J, et al (2013a) Resting-state fMRI in the Human Connectome
Project. Neuroimage 80:144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039
Smith SM, Beckmann CF, Andersson J, et al (2013b) Resting-state fMRI in the Human Connectome
Project. Neuroimage 80:144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039
Smith SM, Jenkinson M, Woolrich MW, et al (2004) Advances in functional and structural MR
image analysis and implementation as FSL. In: NeuroImage
Sotiropoulos SN, Jbabdi S, Xu J, et al (2013) Advances in diffusion MRI acquisition and processing in
the Human Connectome Project. Neuroimage 80:125–43.
https://doi.org/10.1016/j.neuroimage.2013.05.057
Stepniewska I, Kaas JH (1997) Architectonic subdivisions of the inferior pulvinar in New World and
Old World monkeys. Vis Neurosci 14:1043–1060.
https://doi.org/10.1017/S0952523800011767
Tamietto M, Pullens P, de Gelder B, et al (2012) Subcortical Connections to Human Amygdala and
Changes following Destruction of the Visual Cortex. Current Biology 22:1449–1455.
https://doi.org/10.1016/j.cub.2012.06.006
Thomas Yeo BT, Krienen FM, Sepulcre J, et al (2011) The organization of the human cerebral cortex
estimated by intrinsic functional connectivity. J Neurophysiol 106:1125–1165.
https://doi.org/10.1152/jn.00338.2011
Tian Y, Margulies DS, Breakspear M, Zalesky A (2020) Topographic organization of the human
subcortex unveiled with functional connectivity gradients. Nat Neurosci.
https://doi.org/10.1038/s41593-020-00711-6
Tournier J-D, Calamante F, Connelly A (2010) Improved probabilistic streamlines tractography by
2nd order integration over fibre orientation distributions. Proceedings of the International
Society for Magnetic Resonance in Medicine
Tournier JD, Smith R, Raffelt D, et al (2019) MRtrix3: A fast, flexible and open software framework
for medical image processing and visualisation. Neuroimage

Turnbull A, Karapanagiotidis T, Wang H-T, et al (2020) Reductions in task positive neural systems occur with the passage of time and are associated with changes in ongoing thought. Sci Rep 10:9912. https://doi.org/10.1038/s41598-020-66698-z

Tustison NJ, Avants BB, Cook PA, et al (2010) N4ITK: Improved N3 Bias Correction. IEEE Trans Med Imaging 29:1310–1320. https://doi.org/10.1109/TMI.2010.2046908

Uğurbil K, Xu J, Auerbach EJ, et al (2013) Pushing spatial and temporal resolution for functional and diffusion MRI in the Human Connectome Project. Neuroimage 80:80–104. https://doi.org/10.1016/j.neuroimage.2013.05.012

Vakilna YS, Chaitanya G, Hafeez MU, et al (2023) Pulvinar neuromodulation for seizure monitoring and network modulation in temporal plus epilepsy. Ann Clin Transl Neurol 10:1254–1259. https://doi.org/10.1002/acn3.51815

Valk SL, Xu T, Margulies DS, et al (2020) Shaping brain structure: Genetic and phylogenetic axes of macroscale organization of cortical thickness. Sci Adv 6:. https://doi.org/10.1126/sciadv.abb3417

Van Essen DC, Smith SM, Barch DM, et al (2013) The WU-Minn Human Connectome Project: An overview. Neuroimage. https://doi.org/10.1016/j.neuroimage.2013.05.041

Van Essen DC, Ugurbil K, Auerbach E, et al (2012) The Human Connectome Project: A data acquisition perspective. Neuroimage 62:2222–2231. https://doi.org/10.1016/j.neuroimage.2012.02.018

Velioglu HA, Ayyildiz B, Ayyildiz S, et al (2023) A structural and resting-state functional connectivity investigation of the pulvinar in elderly individuals and Alzheimer's disease patients. Alzheimer's & Dementia 19:2774–2789. https://doi.org/10.1002/alz.12850

Veraart J, Novikov DS, Christiaens D, et al (2016) Denoising of diffusion MRI using random matrix theory. Neuroimage. https://doi.org/10.1016/j.neuroimage.2016.08.016

Vos De Wael R, Royer J, Tavakol S, et al (2021) Structural Connectivity Gradients of the Temporal Lobe Serve as Multiscale Axes of Brain Organization and Cortical Evolution. Cerebral Cortex (New York, NY) 31:5151. https://doi.org/10.1093/CERCOR/BHAB149

Webster MJ, Bachevalier J, Ungerleider LG (1993) Subcortical connections of inferior temporal areas TE and TEO in macaque monkeys. Journal of Comparative Neurology 335:73–91. https://doi.org/10.1002/cne.903350106

Wong GM, Hofmann K, Shlobin NA, et al (2023) Stimulation of the pulvinar nucleus of the thalamus in epilepsy: A systematic review and individual patient data (IPD) analysis. Clin Neurol Neurosurg 235:108041. https://doi.org/10.1016/j.clineuro.2023.108041

Yang S, Meng Y, Li J, et al (2020) The thalamic functional gradient and its relationship to structural basis and cognitive relevance. Neuroimage 218:116960. https://doi.org/10.1016/j.neuroimage.2020.116960

Zhang Y, Lu Z, Sun Y, et al (2023) Predictive role of pulvinar in social functional outcome of schizophrenia. Psychiatry Res 327:115419. https://doi.org/10.1016/j.psychres.2023.115419

1211

1212

1213

1214

1215

1216