Neural specialisation for concrete and abstract concepts revealed through meta-analysis

Paul HOFFMAN*

Matthew BAIR

School of Philosophy, Psychology & Language Sciences, University of Edinburgh, UK

* Correspondence to:
Dr. Paul Hoffman
School of Philosophy, Psychology & Language Sciences,
University of Edinburgh,
7 George Square, Edinburgh, EH8 9JZ, UK
Tel: +44 (0) 131 650 4654
Email: p.hoffman@ed.ac.uk

Acknowledgements
The research was supported by a BBSRC grant (BB/T004444/1). For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Abstract

Identifying the brain regions that process concrete and abstract concepts is key to understanding the neural architecture of thought, memory and language. We review current theories of concreteness effects and test their neural predictions in a meta-analysis of 72 neuroimaging studies (1400 participants). Concrete concepts preferentially activated visual and action processing regions, particularly when presented in sentences. Abstract concepts preferentially activated networks for social cognition (particularly for sentences), and for language and semantic control (particularly when presented as single words). These results suggest that concrete and abstract concepts vary both in the information-processing modalities they engage and in the demands they place on cognitive control processes. Specialisation for both concept types was present in different parts of the default mode network (DMN), with effects dissociating along a social-spatial axis. Concrete concepts generated greater activation in a medial temporal DMN component, implicated in constructing mental models of spatial contexts and scenes. In contrast, abstract concepts showed greater activation in frontotemporal DMN regions involved in theory-of-mind and language processing. These results support prior claims that generating models of situations and events is a core DMN function and further indicate specialisation within DMN for different aspects of these models.

Public significance statement

The distinction between concrete and abstract concepts is fundamental to language and cognition. Here, we present the largest meta-analysis of neuroimaging studies experiments investigating this distinction, including data from 1400 participants. Our results indicate that concrete and abstract language differentially engages a range of neural systems involved in perception, action, language and social processing. These results provide new insights into how the brain constructs representations of the world from linguistic information, and into the neural basis of imagination.

Introduction

Human language and experience encompasses a vast array of concepts along a spectrum of concreteness. At one end of the spectrum lie highly concrete words that label objects and acts we directly perceive in our environment (e.g., *a parrot, to jump*). At the other, we find highly abstract words whose meanings are intangible and divorced from direct sensorimotor experience (e.g., *a theory, to believe*). Accounting for this conceptual diversity is a central challenge for theories of language comprehension and semantic cognition (Reilly et al., 2023). A range of distinct theories have been developed to explain processing differences between concrete and abstract concepts, motivated by a large body of behavioural and neuroscientific data. In this article, we briefly review these theories and identify their predictions about how concrete and abstract concepts differentially activate the cortex. We then test these predictions in meta-analysis of 72 neuroimaging studies (1400 participants) that contrasted activation to concrete and abstract language. Our analysis includes twice as many studies as previous meta-analyses of these effects (Binder et al., 2009; Bucur & Papagno, 2021; J. Wang et al., 2010), allowing for a more complete mapping of these effects across the brain and allowing us to test for the first time how neural concreteness effects vary systemically along theoretically-relevant stimulus and task dimensions. By cleaving the neural response to meaning along the fundamental dimension of concreteness, we gain new insights into the ways in which the brain constructs representations of the different forms of experience conveyed through language.

Many of the major accounts of concreteness effects stem from an embodied (or grounded) approach to semantics (Barsalou, 1999; Gallese & Lakoff, 2005; Glenberg & Robertson, 2000; Pulvermüller, 2013). Developed in opposition to classical symbolic views of language (Johnson-Laird, 1988; Newell, 1980), the embodied view holds that conceptual representations are closely tied to the perceptual experiences they reference and that understanding a concept involves reactivating those experiences in some way. The embodied semantics umbrella shelters a broad range of specific theories, from strong accounts that emphasise the primacy of embodied reactivation of sensorimotor cortices (Gallese & Lakoff, 2005; Glenberg & Robertson, 2000) to those that propose a central role for multimodal hub regions that bind and integrate the various types of experience associated with each concept (Binder, 2016; Binder & Desai, 2011; Lambon Ralph et al., 2017). These theories share the general assumption that understanding words frequently involves activating neural networks that process the experiences the words refer to.

Embodied semantics accounts have traditionally focused on the involvement of sensorimotor modalities in semantic representation. An early forerunner of these, Dual-Coding theory, proposed that concrete words are represented in a visual semantic system in addition to a verbal system shared by abstract words (Paivio, 1986). More recently, evidence has accumulated for activation of a range of sensorimotor networks during concrete word processing. Motor and motion-processing regions are activated when we process language that describes actions (Giacobbe et al., 2022; Hauk et al., 2004; Saygin et al., 2010; Watson et al., 2013) and parts of the ventral visual stream are activated when people make judgements about the shapes and colours of words' referents (Fernandino et al., 2016; Martin et al., 1995; Thompson-Schill et al., 1999; van Dam et al., 2012). This embodied perspective makes a straightforward prediction about concreteness effects in the brain. Since concrete words are more strongly associated with sensorimotor experiences, they should activate sensorimotor cortices to a greater extent. As vision is the dominant modality influencing a word's concreteness (Connell & Lynott, 2012), we would expect strong concrete > abstract (C>A) activation effects in the visual system. Action processing cortex should show similar effects, particularly for sentences that contain action verbs.

The representation of abstract concepts has frequently been presented as a challenge to the embodied semantics approach, because they do not directly reference sensorimotor experiences. This challenge has been addressed in a number of ways. Conceptual Metaphor Theory (Gibbs, 1994; Lakoff & Johnson, 1999) asserts that abstract concepts activate sensorimotor representations indirectly through metaphor and analogy (e.g., the association of power with vertical height) (Pecher, 2018). Few studies have investigated whether abstract concepts activate sensorimotor cortices indirectly via metaphoric association (Desai, 2022; Joue et al., 2020) but it seems likely that such indirect activation would be weaker than that elicited by concrete concepts. Thus, this view is also compatible with C>A effects in sensorimotor cortex.

Other researchers have considered other forms of experience that abstract words draw upon. The Affective Embodiment view (Kousta et al., 2011; Lenci et al., 2018; Newcombe et al., 2012; Vigliocco et al., 2014) holds that abstract concepts are more closely associated with emotional content than concrete words (but see (Winter, 2023)) and should therefore preferentially engage brain regions involved in experiencing emotions. This perspective is part of a wider move to ground semantics in activation of internal bodily states and experiences as well as in the external world (Barsalou, 2008; Connell et al., 2018; Martin,

2016). A related view emphasises the importance of social experiences in the acquisition and understanding of abstract concepts (Borghi et al., 2019; Pexman et al., 2022; Troche et al., 2014; Wiemer-Hastings & Xu, 2005). Many abstract words refer to social interactions at an interpersonal or societal level (e.g., loan, democracy) and might therefore be supported by simulations of our own or others' mental states. These related views predict abstract > concrete (A>C) activation effects in brain regions that process emotion and in parts of the default mode network involved in social cognition and theory of mind (Andrews-Hanna et al., 2014; Yeshurun et al., 2021).

In addition to direct experience, we gain knowledge about concepts through their expression in language. The recent successes of large computational language models have demonstrated that sophisticated semantic information can be derived solely from the statistical patterns of natural language use (Devlin et al., 2019; Mikolov et al., 2013; Pereira et al., 2016). In line with this position, a number of hybrid theories suggest that semantic processing is supported by language-based representations as well as by simulations in modal brain systems (Barsalou et al., 2008; Binder & Desai, 2011; Connell, 2019; Dove et al., 2022; Louwerse, 2011; Vigliocco et al., 2009). Verbal experience is more critical to the acquisition of abstract words and is thought to play a more central role in determining their meanings (Borghi et al., 2019; Vigliocco et al., 2009; Villani et al., 2019). These hybrid views therefore predict A>C effects in temporal lobe regions involved in language comprehension. Some researchers see linguistic representations of meaning as most suited to supporting superficial processing (Barsalou et al., 2008; Connell, 2019). A>C effects in language cortex might therefore be most prominent for simple tasks, while deeper conceptual processing instead requires more complex representations grounded in other brain systems.

Thus far, we have focused on the types of experience that contribute to our understanding of concrete and abstract words. Other accounts have investigated the role of context in shaping our understanding of each concept type. It has long been known that when people are presented with isolated abstract words, they find it hard to generate contexts in which the word could be used (Schwanenflugel et al., 1988; Schwanenflugel & Shoben, 1983). This is probably because abstract words are used in a wider variety of contexts than concrete words, suggesting that their meanings are highly context-dependent (Davis et al., 2020; Hoffman et al., 2013). Abstract word meanings also vary greatly between individuals (X. Wang & Bi, 2021) and are difficult to process in the absence of context (Bechtold et al., 2021; Hoffman et al., 2010; Schwanenflugel et al., 1988; Schwanenflugel & Shoben, 1983). Because

of this inherent variability and context dependence, we and others have proposed understanding abstract words relies more on control processes that regulate how semantic knowledge is activated (Bechtold et al., 2021; Cousins et al., 2016; Davis et al., 2020; Hoffman, 2016; Hoffman et al., 2010, 2015). These accounts predict A>C activation in brain regions that support controlled semantic processing (Jackson, 2021) and predict that these effects should be largest (a) when no context is available to support processing (i.e., for single words rather than sentences) and (b) when participants are required to make an specific semantic judgement (cf. more passive processing).

Finally, a related account proposes that abstract concepts vary greatly in the real-world situations or events that they describe, as well as in their verbal contexts (Barsalou et al., 2018; Davis et al., 2020). For example, an abstract concept like *justice* could sometimes bring to mind a courtroom situation and at other times a prison (Davis et al., 2020). It has also been suggested that we rely heavily on situational features – i.e., the spatiotemporal context of an experience, the entities present and their interactions – to bring meaning to abstract concepts (Barsalou et al., 2018; Barsalou & Wiemer-Hastings, 2005; Binder, 2016; Davis et al., 2020; McRae et al., 2018). This proposal is related to the claim that social interactions are highly relevant to abstract words (Borghi et al., 2019; Pexman et al., 2022; Troche et al., 2014) and that abstract concepts are structured by their thematic relationships with one another (Crutch & Warrington, 2005). It is hard to derive clear predictions about neural concreteness effects from this situational account. If processing abstract words involves simulating the situations they could refer to, we might expect A>C activation in brain regions that represent situation models (Ranganath & Ritchey, 2012; Yeshurun et al., 2021). However, given that abstract concepts are weakly associated with a wide range of possible situations, such effects might only occur when abstract words are placed in a specific constraining context (e.g., a sentence). During processing of single words, we might expect concrete words to activate situational cortex more, since these words bring specific situations to mind more readily (Binder, 2016; Schwanenflugel et al., 1988).

As this brief review has shown, identifying neural processing differences between concrete and abstract concepts is critical to unlocking a deeper understanding of how we bring meaning to experience and language. The various theoretical claims and their neural predictions are summarised in Table 1. To test these, we performed quantitative co-ordinate-based meta-analyses of 72 neuroimaging studies that manipulated the concreteness of language stimuli. We formally assessed how concreteness effects intersect with the brain regions that

support 7 key processing domains (see Figure 2). By examining the differential contributions of these systems to concrete and abstract processing, we were able to assess the degree of support for each theoretical position. These theories also make specific predictions about concreteness effects depend on the stimuli and tasks used to elicit them. Thus, we systematically investigated how the expression of C>A and A>C effects is influenced by two theoretically-relevant factors: (a) the distinction between single-word and sentence stimuli and (b) the depth of semantic processing participants were required to engage in.

We also assessed where concreteness effects occur within the default-mode network (DMN). DMN is often deactivated for tasks that require sole focus on the external environment, leading to its traditional characterisation as a task-negative network(Buckner et al., 2008; Raichle et al., 2001). However, tasks that require use of internally-generated representations, including spontaneous thoughts, episodic memories and semantic knowledge, positively engage this network (Andrews-Hanna, 2012; Binder et al., 2009; Ranganath & Ritchey, 2012). These findings, along with the observation that DMN regions are convergence points for multiple modal processing streams (Binder & Desai, 2011; Margulies et al., 2016), have led to proposals that DMN integrates incoming information with past experience to mentally simulate complex multimodal events and situations (Baldassano et al., 2018; Fernandino & Binder, 2024; Heidlmayr et al., 2020; Morales et al., 2022; Ranganath & Ritchey, 2012; Yeshurun et al., 2021). DMN regions are therefore prime candidates for supporting the different mental simulations required to understand concrete and abstract concepts.

The DMN can be fractionated into three sub-systems with distinct functional profiles (Andrews-Hanna et al., 2010, 2014; Lee et al., 2021; Shao et al., 2024; Yeo et al., 2011), shown in Figure 2. First, a medial temporal sub-network (MT-DMN) encompasses the hippocampus and adjacent medial temporal cortex, as well as parts of retrospenial, posterior inferior parietal and ventromedial prefrontal cortices. This subnetwork is particularly involved in episodic memory and the mental representation of scenes and spatial contexts. Second, a frontotemporal system (FT-DMN; sometimes termed dorsal medial DMN) consists of lateral temporal, dorsomedial prefrontal and anterior inferior parietal cortices, as well as lateral prefrontal regions that frequently co-activate with them. This subnetwork has been implicated in semantic and social cognition. Finally, a core system (core-DMN) is spatially interposed between the other two systems and shows some functional characteristics of both, but with particular engagement for self-relevant processing. It has recently been suggested that these systems might differentially support concrete and abstract aspects of thought and imagination

(Andrews-Hanna & Grilli, 2021). Specifically, the complex semantic representations of abstract words and their reliance on social cognition might lead these regions to preferentially activate FT-DMN regions, while the grounding of concrete concepts in external spatial contexts might associate these with the MT-DMN system. Our meta-analysis assesses this proposal within the domain of language processing for the first time.

Method

Study selection

The process for identifying studies is summarised in Figure 1. We searched for peer-reviewed studies published between January 1990 and August 2022, which contrasted processing of concrete and abstract language. An initial search was conducted using the Scopus database for articles containing the following terms in their title, abstract or keywords: (concrete* or abstract* or imageab*) and (fMRI or PET or functional MRI or functional magnetic or positron emission) and (word* or language or semantic* or sentence* or memory). This yielded 903 studies, which were screened for inclusion.

The inclusion criteria were:

1. The article reported the results of a functional neuroimaging experiment in healthy adult participants.

2. Participants were required to process meaningful language stimuli in their native language, which could be presented in written form or as audio or video recordings. We placed no restriction on the type of processing required.

3. The language stimuli varied in concreteness, either in a factorial design (concrete vs. abstract) or through a parametric manipulation of concreteness values. We included all studies where the researchers referred to their conditions as concrete and/or abstract. In most cases, researchers reported that their conditions differed in concreteness or imageability ratings obtained from a published database, such as (Brysbaert et al., 2014).

4. The article reported activation peaks for a univariate contrast between concrete and abstract conditions (or effect of parametric variation in concreteness) in a standard stereotactic space (MNI or Talairach).

A total of 64 studies were identified through this process. We added to this 8 further studies that met the inclusion criteria but were not returned in the Scopus search. These were either cited by studies that passed the initial screening process and were present in previous

neuroimaging meta-analyses of concreteness effects (Binder et al., 2009; Bucur & Papagno, 2021; J. Wang et al., 2010). Thus, a total of 72 studies were included in analyses (see Supplementary Materials for a full list).

To investigate how concreteness effects were influenced by relevant study characteristics, we classified studies in two dimensions.

1. <u>Stimulus.</u> *Word* studies used single-word stimuli (including those that presented pairs or triads of single words for semantic decisions). *Sentence* studies used sentence-level stimuli (including three studies that used noun-verb phrases).

2. <u>Task.</u> *Semantic* studies used tasks that explicitly required participants to make a judgement based on the meaning of the stimuli (e.g., semantic relatedness judgements, pleasantness judgements, sensibility judgements). *Non-semantic* studies did not require explicit meaning-based decisions (e.g., passive reading/listening, lexical decision, memory tasks).

For both of these dimensions, where a study could not be classified unambiguously it was assigned to neither category.

<u>ALE analyses</u>

Activation Likelihood Estimation (ALE) analyses were carried out using GingerALE 3.0.2 (Eickhoff et al., 2012). The software takes peaks from neuroimaging contrasts of interest across a set of studies, models the spatial distribution of these peaks and computes activation likelihood maps for the whole brain. These maps are then subjected to voxel-wise statistical tests to identify regions that consistently show effects across studies.

We first used the ALE method to identify regions showing greater activation for concrete relative to abstract words (C>A) and the reverse (A>C) in the full set of 72 studies. To investigate the factors influencing these effects, we then conducted subtraction analyses that contrasted studies on the two dimensions. For example, the Stimulus C>A subtraction analyses identified regions that were more likely to show C>A effects for Word studies than in Sentence studies and vice-versa. There were four subtraction analyses in total (C>A and A>C for Stimulus and Task contrasts).

We used Turkeltaub et al.'s (2012) non-additive ALE algorithm, which limits the influence of individual studies reporting multiple peaks close to one another. Peaks reported in Talairach space were converted to MNI space using the tal2icbm_spm transform (Lancaster et al., 2007). The main C>A and A>C results were thresholded using the permutation-based

method for cluster-level inference (Eickhoff et al., 2012). A family-wise error cluster-corrected threshold of $p < 0.05$ was used (with a cluster-forming threshold of $p < 0.005$ and 10,000 permutations). Cluster-level inference is not available for subtraction analyses so for these we used a voxelwise threshold of $p < 0.05$ combined with a minimum cluster extent of 1600mm^3 and 100,000 permutations.

Comparison with functional brain networks

We used three methods to systematically interpret the functional correlates of regions exhibiting concreteness effects. First, we submitted unthresholded ALE maps to the Neurosynth website (Neurosynth.org) to obtain correlations with its meta-analytic probability maps. Neurosynth is an automated meta-analysis tool that identifies terms commonly used in a large corpus of neuroimaging papers (14,371 studies for the analyses reported here) and relates these to reported peak activation co-ordinates in the same papers. A Neurosynth map for a given term indicates the likelihood that activation in each voxel is preferentially reported by studies that use the term. For example, in the map for "language", each voxel's value indicates the likelihood that activation is reported there in papers that discuss language, relative to studies that don't. This large-scale unsupervised approach to meta-analysis can successfully reproduce known functional networks as well as allowing exploration of new terms (Rubin et al., 2017; Yarkoni et al., 2011). It is important to note, however, that the automated nature of Neurosynth makes it unsuitable for investigating concreteness effects. Papers associated with the term "concreteness" usually report both C>A and A>C peaks and Neurosynth's algorithms do not discriminate between these.

By correlating the Neurosynth term maps with our unthresholded ALE maps, we were able to determine which terms are most consistently associated with regions that show concreteness effects. For the main analysis, we subtracted A>C ALE values from C>A ALE values and correlated the result with Neurosynth maps. Here, positively-correlated terms were more associated with regions showing C>A effects and negatively-correlated terms with A>C effects. To visualise the results, terms relating to anatomical structures were removed and the 30 most strongly correlated terms in each direction were extracted. These were plotted as word clouds in which the size of each term is proportional to the strength of its correlation with concreteness effects. A similar process was followed to visualise associations with the effects of each study dimension.

Second, we compared the spatial distribution of concreteness effects with a series of functional brain networks. We selected networks that support the various processes that concrete and abstract words have been hypothesised to differentially rely on, again mostly based on Neurosynth meta-analytic maps. These were:

1. Action. Regions associated with the "action" term.
2. Visual. Regions associated with the "visual" term.
3. Emotions. Regions associated with the "emotional" term.
4. Social. Regions associated with the "social" term.
5. Language. Regions associated with the "language comprehension" term.
6. Semantic control. No term similar to semantic control was present in the Neurosynth database so we instead used the network identified in a recent meta-analysis of studies manipulating controlled processing demands in semantic tasks (Jackson, 2021).
7. Situations. "Situations" was not present in the Neurosynth database as an individual term. Instead, we searched Neurosynth's topic-based maps (Rubin et al., 2017) for a topics that related to processing of situations and events. We selected a topic from the v5-topics-200 set for which the highest-loading terms were: events, future, personal, past, cultural, thinking, construction, episodic, experiences.

Neurosynth maps (z-values for test of the association between voxel activation and term use, thresholded at FDR-corrected $p < 0.01$) were downloaded from neurosynth.org, smoothed at 4mm FWHM and binarized at a threshold of $z>3$. They were then compared with binarised images of significant concreteness effects to determine the degree to which they overlap. We quantified this by measuring the proportion of voxels showing C>A and A>C effects that fell within each network.

Finally, we compared the spatial distribution of concreteness effects across sub-systems of the default mode network (DMN). Following other researchers (Andrews-Hanna et al., 2014; Poerio et al., 2017; Shao et al., 2024), we used Yeo et al.'s (2011) resting-state parcellation of the brain into 17 networks to define frontotemporal, core and medial temporal DMN sub-networks. We used the same proportional measure to quantify overlap between these sub-networks and concreteness effects.

Transparency and Openness

The raw data for these analyses, including all peak co-ordinates and the studies and contrasts they were obtained from, are available on the OSF website: https://osf.io/78ra9/.

Untrhresholded ALE maps for all analyses are available at the same address, as is the code used to perfmed the network comparisons. The analyses were not pre-registered.

Results

Literature Search

We identified 72 studies that reported peak activation co-ordinates for contrasts of concrete vs. abstract language stimuli. Studies were published between 1995 and 2022 and together comprised 10 languages, 1400 participants, 459 C>A peaks and 435 A>C peaks. The number of included studies is considerably higher than older meta-analyses on this topic (Binder et al., 2009; J. Wang et al., 2010). It is also higher than the 32 studies analysed more recently by Bucur & Papagno (2021), principally because these authors only included studies that used single-word stimuli (whereas we also included phrases and sentence stimuli). As reported in Supplementary Table 1, studies used a range of tasks to contrast concrete and abstract words, including passive reading/listening, lexical decision, semantic decisions of various types and memory encoding and retrieval tasks. A range of stimuli were also employed, with some studies presenting individual words and others presenting phrases or complete sentences. To investigate how these study characteristics influenced the expression of concreteness effects, we classified studies along two theoretically-relevant dimensions: *Stimuli* (23 Sentence studies; 48 Word studies; 1 unclassifiable) and *Task* (30 Semantic; 40 Non-semantic; 2 unclassifiable). There was no statistical association between these two dimensions ($\chi^2(1) = 0.18$, $p = 0.67$).

Elucidation of relevant cortical networks

To aid interpretation of our results, we first defined 7 relevant functional networks (obtained from existing meta-analytic resources; see Method for details) and 3 sub-divisions of the DMN, obtained from parcellation of resting-state fMRI data (Yeo et al., 2011). These are shown in Figure 2. There are two important points to take away from Figure 2 that colour the results reported below. First, the functional networks frequently overlap one another. The social network overlaps considerably with the emotion network, particularly in the amygdala and ventromedial prefrontal cortex. It also shows convergence with situation-processing regions in the lateral anterior temporal lobes and inferior parietal cortices. Parts of left inferior prefrontal cortex are implicated in semantic control but also more generally in

language comprehension, presumably as control is frequently needed to resolve verbal semantic ambiguities (Vitello & Rodd, 2015). Even the sensorimotor networks for vision and action overlap in the lateral occipitotemporal cortex, in motion-sensitive area V5/MT+ (Zeki, 2004). This overlap in functional networks reflects the fact that higher-order cognition involves the co-activation of a broad range of representations and processes. For example, understanding situations involving people may require us to simulate their mental states, leading to overlap between the situation and social networks. This cements the idea that different perspectives on concreteness effects are not mutually exclusive. It is possible, for example, that abstract concepts rely heavily on language representations and, as a consequence, recruit semantic control regions that resolve ambiguity in language. By understanding how different networks work in concert to support concrete and abstract comprehension, we gain insights into how meaning is constructed among multiple interacting systems.

The second critical point is that many of the functions under investigation align with sub-divisions of the DMN. As other researchers have noted, regions involved in social cognition, language processing and semantic control tend to fall within FT-DMN (particularly in the lateral temporal and lateral prefrontal cortices) but not MT-DMN (Andrews-Hanna et al., 2014; Smallwood et al., 2021), which may bias these FT-DMN regions towards preferring abstract concepts. In contrast, the situation-processing network includes parts of all DMN sub-divisions, including lateral temporal regions of FT-DMN and medial temporal and retrospenial parts of MT-DMN. This may reflect the diverse types of processing that are relevant to different types of events and situations. Inter-personal interactions are important in many real-world situations, and these may engage social cognition regions in FT-DMN. In other situations, the physical environment may be more salient and these could engage scene processing regions in MT-DMN. The general alignment of the situation network with DMN fits with the view that a major function of the DMN is to construct situation models (Baldassano et al., 2018; Heidlmayr et al., 2020; Morales et al., 2022; Ranganath & Ritchey, 2012; Yeshurun et al., 2021). The association of the situation network with multiple DMN sub-divisions hints at the diverse types of processing called upon across situations and also raises the possibility that concrete and abstract specialisation might occur in different parts of this network. In contrast, there is no intersection between action and visual networks and the DMN. MT-DMN regions, particularly in the medial temporal lobe, are linked with the

mental representation of visual scenes (Epstein & Baker, 2019). However, temporal lobe MT-DMN regions fall slightly anterior to those implicated in direct visual perception.

Overall Effects of Concreteness

Activation Likelihood Estimation (ALE) analyses (Eickhoff et al., 2009, 2012) were used to identify regions that consistently showed positive (C>A) and negative (A>C) concreteness effects across the full set of 72 studies. Results are shown in Figure 3 (for peak co-ordinates, see Table 2). As in earlier meta-analyses (Binder et al., 2009; Bucur & Papagno, 2021; J. Wang et al., 2010), the most prominent A>C effects were observed in left inferior frontal gyrus (IFG) and anterior parts of the superior and middle temporal gyri (aSTG/MTG). C>A effects were most pronounced in the angular gyri (AG) and posterior parts of the left inferior temporal, fusiform and parahippocampal gyri (ITG/FG/PHG). Neighbouring regions of cortex sometimes showed opposing effects, particularly in posterior MTG/ITG and in the posterior cingulate and precuneus (PCC/PCu). In addition, while previous analyses identified strongly left-lateralised effects, our higher-powered analysis reveals that these effects are often mirrored in the right hemisphere, particularly in right aSTG/MTG (A>C), right PHG (C>A) and right AG (C>A). Thus, although language processing is traditionally thought of as a left-lateralised function, concrete and abstract words differentially engage bilateral brain networks, in line with the bilateral nature of many of the functional networks shown in Figure 2.

The bar plots in Figure 3 show the proportion of voxels showing concreteness effects that fell within each of our functional networks of interest. These results support many of the predictions in Table 1. Abstract words preferentially activated parts of the social network (particularly aSTG/MTG), language network (IFG and lateral temporal cortex) and semantic control network (left IFG and pMTG). Concrete word effects showed overlap with action and visual networks, though this was more limited. In particular, C>A effects were observed around the motion-sensitive cortex of left V5/MT+ and in the left supramarginal gyrus. C>A effects were also found in the ventral temporal cortex but these were in MT-DMN cortex, largely anterior to regions directly implicated in visual perception.

The situation network showed considerable overlap with both C>A and A>C effects, suggesting that both concrete and abstract language can elicit mental representations of events and situations. Importantly, concrete and abstract effects occurred in different parts of this network, a trend that can be seen more clearly when we turn to the DMN sub-networks. Here, we saw a striking dissociation between concrete and abstract concepts. Greater

activation for abstract words was strongly associated with FT-DMN regions. Almost 50% of A>C voxels fell within this network while less than 1% of C>A voxels did. In contrast, more than 20% of C>A voxels fell within MT-DMN while no A>C voxels did. This dissociation is clearest in the anterior temporal cortices, where lateral FT-DMN regions were more active for abstract language while ventromedial MT-DMN regions preferred concrete language (for a similar result, see (Hoffman et al., 2015)). These results have two important implications. First, although DMN is most associated with tracking semantic content in complex narratives (Baldassano et al., 2018; Heidlmayr et al., 2020; Morales et al., 2022), our results show that DMN representations can also be elicited by more simple language stimuli (single words and sentences). Second, they indicate that the type of content that engages this system varies systematically along a social-spatial axis.

Finally, the wordclouds in Figure 3 visualise the terms in the neuroimaging literature that are most strongly associated with regions showing C>A and A>C effects. These were generated by submitting C>A and A>C ALE maps to the decoding function in the Neurosynth meta-analytic database (Rubin et al., 2017; Yarkoni et al., 2011) (see Method for details). This automated decoding approach supports the picture already seen. It confirms that regions preferentially activated by abstract language are associated with theory of mind, mental states and language processing, particularly at the sentence level. Conversely, the pattern for concrete language is associated with aspects of visual and action processing (e.g., tools, pictures, mental imagery). However, C>A regions are most strongly associated with episodic and autobiographical memory, which are core MT-DMN functions. There were 7 studies in the meta-analysis where participants were asked to either encode or retrieve items from episodic memory. We re-ran the analysis excluding these studies and obtained very similar results (see Supplementary Figure 1), indicating that these effects do not stem from task instructions. It could be that concrete words automatically trigger retrieval of specific autobiographical or episodic memories, because they bring to mind specific situations (Binder, 2016). Alternatively, it could be that tasks used to probe episodic/autobiographical memory rely on similar processes to those involved in comprehending concrete language (e.g., the mental representation of visuospatial contexts).

Influence of study characteristics

We next tested how concreteness effects varied as a function of stimulus and task. Figure 4 shows contrasts of Word vs. Sentence studies, conducted separately for A>C and

C>A effects. Red regions in the left-hand panel are those where A>C effects were more likely to be observed for studies that presented sentences (or multi-word phrases). These regions primarily overlapped with the social and situation networks and included the lateral anterior temporal lobes bilaterally and left temporoparietal junction. These findings support the idea that abstract sentences are more likely to bring to mind rich social situations than individual abstract words. Yellow regions were more likely to show A>C effects in studies that presented single words. These effects were concentrated in left IFG and pMTG, which are major constituents of the semantic control network (Jackson, 2021). These results are consistent with the view that the cognitive control demands of processing abstract words are maximal when no context is available to constrain their interpretation.

The right-hand panel of Figure 4 shows stimulus influences on concrete word activation. Green regions show where C>A effects were more likely for Sentence studies. These effects were observed mainly in parts of the action network, specifically left lateral occipitotemporal cortex and supramarginal gyrus. This engagement suggests simulation of sentence content, since concrete sentences invariably describe physical actions. In contrast, Word studies typically present concrete nouns rather than verbs, which would not be expected to engage simulations of actions. However, the visual network was also more likely to show C>A effects when sentences were used, suggesting that concrete sentences might also elicit richer sensory simulations more generally. A region of the left inferior frontal sulcus was also more likely to show C>A activation in Sentence studies. Greater likelihood of C>A activation in Word studies was limited to a small number of voxels in ventral precuneus.

Figure 5 presents analysis of studies split by task type. For A>C effects, a number of regions were more likely to show abstract word preferences when participants made explicit semantic judgements about the stimuli (or opposed to non-semantic tasks or passive reading/listening). These effects were distributed across several functional networks, suggesting that deeper semantic processing accentuates general activation for abstract language but does not make demands on any specific process or network. No regions were more likely to show A>C effects in Non-semantic studies.

The picture for concrete concept preferences was more complex. Semantic studies were more likely to find C>A activation in a distributed set of regions that primarily coincided with the situation network. These included parts of AG, PCu and PHG. Greater activation in Non-semantic studies was confined to a single cluster in the left pre-central gyrus, a region primarily associated with action and motor processing. These effects might reflect a tendency

to focus on motoric aspects of concrete concepts during shallower processing, with broader situational contexts only instated when participants evaluate concepts more deeply. Finally, when comparing across panels in Figure 5, it is clear that the dissociation in concreteness effects across the DMN was more likely to occur when participants made explicit semantic judgements. This suggests the association of the two classes with different DMN sub-divisions is related to deep processing of the conceptual significance of these concepts.

Discussion

The distinction between concrete and abstract concepts is fundamental to language and semantic cognition, with a range of theories predicting specific differences in how these classes of concept are processed in the brain. We tested these predictions in the largest meta-analysis of the literature to date. When aggregating across 72 neuroimaging studies, we found widespread activation differences between concrete and abstract language, falling within a range of functional networks that support different aspects of conceptual processing. In particular, there was a systematic association of concrete and abstract activation with different subdivisions of the DMN. Effects were most widespread in the left hemisphere though in some cases these were mirrored in the right, in line with other findings that conceptual processing in language tasks engages bilateral neural networks (Patel et al., 2023; Yu et al., 2024).

Many of the predictions in Table 1 were supported by our analyses. In line with embodied semantics theories (Barsalou, 1999, 2008; Binder, 2016; Binder & Desai, 2011; Connell et al., 2018; Gallese & Lakoff, 2005; Glenberg & Robertson, 2000; Lambon Ralph et al., 2017; Martin, 2016; Pulvermüller, 2013), concrete language produced greater activation in visual and action networks, while abstract language preferentially activated the social network (though not the emotion network). These effects were more likely when participants processed sentences, which elicit richer representations of actions, events and social interactions. Abstract concepts engaged the language network to a greater degree than concrete though, contrary to some predictions (Barsalou et al., 2008; Connell, 2019), this effect was not more prominent for shallower, less semantic tasks. Abstract concepts also relied more heavily on semantic control regions, particularly when presented as single words, supporting the claim that the complex, variable meanings of these words require more executive regulation (Bechtold et al., 2021; Cousins et al., 2016; Davis et al., 2020; Hoffman, 2016; Hoffman et al., 2010, 2015). In general then, our findings are well accounted for by two

broad tenets of semantic cognition: 1. that semantic representations are partly embodied in modality-specific neural systems that process different aspects of our experiences (Barsalou, 1999, 2008; Binder, 2016; Binder & Desai, 2011; Connell et al., 2018; Gallese & Lakoff, 2005; Glenberg & Robertson, 2000; Lambon Ralph et al., 2017; Martin, 2016; Pulvermüller, 2013) and 2. that semantic activation is shaped by control processes and the need for these varies with concept type and level of contextual support (Giallanza et al., 2023; Hoffman et al., 2010, 2015; Jackson, 2021; Lambon Ralph et al., 2017).

In addition to providing robust empirical assessment of long-standing claims about concreteness effects, our study provides a rich source of data to assess recent theories about the role of situation models in processing these concept classes (Barsalou et al., 2018; Barsalou & Wiemer-Hastings, 2005; Binder, 2016; Davis et al., 2020; McRae et al., 2018). Rather than being limited to one concept class, different parts of the situation processing network were activated by concrete and abstract language. This suggests that concrete and abstract language describe different types of situation, and that, for abstract words in particular, situation models are more likely to be elicited by sentences than individual words. This apparent divergence in situational processing was further supported by a clear dissociation within DMN sub-divisions, along a social-spatial axis. Abstract stimuli preferentially activated FT-DMN regions involved in social, language and semantic cognition while concrete stimuli engaged MT-DMN regions attuned to processing scenes and spatial contexts.

There is an emerging consensus that a general function of the DMN is to generate simulations of real or imagined events and contexts, variously termed situation models (Fernandino & Binder, 2024; Ranganath & Ritchey, 2012; Yeshurun et al., 2021), schemas (Gilboa & Marlatte, 2017), perceptually-decoupled cognition (Smallwood et al., 2013), mental time travel (Addis, 2020) or simply imagination (Andrews-Hanna & Grilli, 2021). In particular, Andrews-Hanna and Grilli (2021) propose that this general DMN function can be decomposed into two sub-systems they term the "mind's eye" (MT-DMN), which represents visuospatial and external aspects of situations, and the "mind's mind" (FT-DMN), which represents conceptual and internally-focused aspects. Previous studies have found that these systems are preferentially attuned to perceptual/spatial vs. thematic/conceptual aspects of autobiographical memories (D'Argembeau et al., 2014; Gurguryan & Sheldon, 2019), to memories of places vs. people (Silson et al., 2019) and to the vividness vs. emotional salience of imagined events (Lee et al., 2021). Our data provide a powerful new source of evidence for this hypothesis by showing that concrete and abstract words differentially engage the

mind's eye and mind's mind. While a previous study has shown a similar pattern during story reading (Tamir et al., 2016), our meta-analysis shows that single sentences and even individual words are sufficient to drive selective responses in these DMN sub-systems, presumably because they spontaneously bring to mind models of relevant situations. Real-life situations involve a complex interplay of external and internally-focused elements, explaining why the DMN frequently emerges as a single coherent network. However, by isolating activity relating to concrete and abstract aspects of thought, we have begun to identify the underlying structure of the DMN simulation engine.

We must also consider alternative explanations for concreteness effects in the DMN. First, in line with its traditional characterisation as a task-negative network, DMN activation is known to negatively correlate with task difficulty (Mckiernan et al., 2003). Abstract language is usually more difficult to process than concrete (Schwanenflugel et al., 1988), which might lead to generic difficulty-related C>A effects in these regions. However, as we observed both C>A *and* A>C effects in parts of the DMN that show difficulty-related effects (i.e., PCC/PCu), this does not provide a straightforward explanation of our results. Second, MT-DMN regions are frequently implicated in episodic memory recall, raising the possibility that concrete concepts trigger associations with specific past experiences in a way that abstract concepts do not. While this is possible, it is also possible that similarities in activation arise from episodic memory tasks drawing on the mind's eye simulation system. MT-DMN is activated by imagining future events as well as remembering past ones (Addis, 2020; Mullally & Maguire, 2014), suggesting a more general role in simulating both experienced and imagined events.

In this meta-analysis, we aimed to test the major established neurocognitive theories of concreteness. There are, however, some recent perspectives which we did not test explicitly. For example, we did not test recent proposals that comprehending abstract words entails greater activation of mouth motor cortex (Borghi et al., 2019) or that abstract concepts are particularly associated with internal bodily sensations (Connell et al., 2018). In addition, the meta-analytic approach identifies broad trends in activation that are consistent across a range of studies, tasks and stimuli. It is less sensitive to effects that may occur for specific subsets of concept within the broader concrete and abstract domains. For example, we did not find abstract word activation in emotion processing regions as has been proposed by others (Vigliocco et al., 2014) and this may be because emotions are only salient for a subset of abstract concepts that are not strongly sampled in most studies.

Unpicking the contributions of neural systems to specific concepts will require a more nuanced approach than the uni-dimensional approach to concreteness that has dominated the field thus far. Some studies have moved in this direction, quantifying the contributions of specific sensorimotor domains (e.g., shape, colour, sound) as well as more intangible social, emotional, and cognitive aspects of experience to specific concepts (Binder et al., 2016; Connell et al., 2018; Hoffman & Lambon Ralph, 2013; Lynott et al., 2020; Troche et al., 2014; Villani et al., 2019). These studies have confirmed that concreteness is a fundamental organising principle, with concrete and abstract concepts strongly differentiated across many domains. However, characterizing concepts at this finer-grained level allows for a more precise dissection of the neural systems underlying semantic representation. This can include identifying activation that correlates with specific domains of experience (Fernandino et al., 2016) and using multivariate approaches to identify regions whose representational structure is determined by experiential attributes (Tong et al., 2022; X. Wang et al., 2018). These developments notwithstanding, the present study indicates that the broader classes of concrete and abstract concept vary reliably and systematically in their engagement of different neural networks. We propose that much of this variation can be attributed to the different forms of mental simulation involved in understanding these concepts.

References

Addis, D. R. (2020). Mental Time Travel? A Neurocognitive Model of Event Simulation. *Review of Philosophy and Psychology*, *11*(2), 233–259. https://doi.org/10.1007/s13164-020-00470-0

Andrews-Hanna, J. R. (2012). The Brain's Default Network and Its Adaptive Role in Internal Mentation. *The Neuroscientist*, *18*(3), 251–270. https://doi.org/10.1177/1073858411403316

Andrews-Hanna, J. R., & Grilli, M. D. (2021). Mapping the Imaginative Mind: Charting New Paths Forward. *Current Directions in Psychological Science*, *30*(1), 82–89. https://doi.org/10.1177/0963721420980753

Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-anatomic fractionation of the brain's default network. *Neuron*, *65*(4), 550–562. https://doi.org/10.1016/j.neuron.2010.02.005

Andrews-Hanna, J. R., Smallwood, J., & Spreng, R. N. (2014). The default network and self-generated thought: Component processes, dynamic control, and clinical relevance. *Annals of the New York Academy of Sciences*, *1316*(1), 29. https://doi.org/10.1111/nyas.12360

Baldassano, C., Hasson, U., & Norman, K. A. (2018). Representation of real-world event schemas during narrative perception. *Journal of Neuroscience*, *38*(45), 9689–9699. https://doi.org/10.1523/jneurosci.0251-18.2018

Barsalou, L. W. (1999). Perceptual symbol systems. *Behavioral and Brain Sciences*, *22*(04), 577–660.

Barsalou, L. W. (2008). Grounded Cognition. *Annual Review of Psychology*, *59*, 617–645. https://doi.org/10.1146/annurev.psych.59.103006.093639

Barsalou, L. W., Dutriaux, L., & Scheepers, C. (2018). Moving beyond the distinction between concrete and abstract concepts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *373*(1752), 20170144. https://doi.org/10.1098/rstb.2017.0144

Barsalou, L. W., Santos, A., Simmons, W. K., & Wilson, C. D. (2008). Language and simulation in conceptual processing. In A. M. De Vega, A. C. Glenberg, & A. Graesser (Eds.), *Symbols, Embodiment and Meaning* (pp. 245–283). Oxford University Press.

Barsalou, L. W., & Wiemer-Hastings, K. (2005). Situating Abstract Concepts. In D. Pecher & R. A. Zwaan (Eds.), *Grounding Cognition* (1st ed., pp. 129–163). Cambridge University Press. https://doi.org/10.1017/CBO9780511499968.007

Bechtold, L., Bellebaum, C., Hoffman, P., & Ghio, M. (2021). Corroborating behavioral evidence for the interplay of representational richness and semantic control in semantic word processing. *Scientific Reports*, *11*(1), Article 1. https://doi.org/10.1038/s41598-021-85711-7

Binder, J. R. (2016). In defense of abstract conceptual representations. *Psychonomic Bulletin & Review*, *23*(4), 1096–1108. https://doi.org/10.3758/s13423-015-0909-1

Binder, J. R., Conant, L. L., Humphries, C. J., Fernandino, L., Simons, S. B., Aguilar, M., & Desai, R. H. (2016). Toward a brain-based componential semantic representation. *Cognitive Neuropsychology*, *33*(3–4), 130–174. https://doi.org/10.1080/02643294.2016.1147426

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends in Cognitive Sciences*, *15*(11), 527–536. https://doi.org/10/djtc3r

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex*, *19*, 2767–2796. https://doi.org/10.1093/cercor/bhp055

Borghi, A. M., Barca, L., Binkofski, F., Castelfranchi, C., Pezzulo, G., & Tummolini, L. (2019). Words as social tools: Language, sociality and inner grounding in abstract concepts. *Physics of Life Reviews*, *29*, 120–153.

Brysbaert, M., Warriner, A. B., & Kuperman, V. (2014). Concreteness ratings for 40 thousand generally known English word lemmas. *Behavior Research Methods*, *46*(3), 904–911. https://doi.org/10/gd6hzk

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy, function and relevance to disease. *Annals of the New York Academy of Sciences*, *1124*, 1–38. https://doi.org/10.1196/annals.1440.011

Bucur, M., & Papagno, C. (2021). An ALE meta-analytical review of the neural correlates of abstract and concrete words. *Scientific Reports*, *11*, 15727. https://doi.org/10/gs4tfk

Connell, L. (2019). What have labels ever done for us? The linguistic shortcut in conceptual processing. *Language, Cognition and Neuroscience*, *34*(10), 1308–1318. https://doi.org/10.1080/23273798.2018.1471512

Connell, L., & Lynott, D. (2012). Strength of perceptual experience predicts word processing performance better than concreteness or imageability. *Cognition*, *125*(3), 452–465. https://doi.org/10/f4dvqc

Connell, L., Lynott, D., & Banks, B. (2018). Interoception: The forgotten modality in perceptual grounding of abstract and concrete concepts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *373*(1752), 20170143. https://doi.org/10.1098/rstb.2017.0143

Cousins, K. A., York, C., Bauer, L., & Grossman, M. (2016). Cognitive and anatomic double dissociation in the representation of concrete and abstract words in semantic variant and behavioral variant frontotemporal degeneration. *Neuropsychologia*, *84*, 244–251. https://doi.org/10/f8m24b

Crutch, S. J., & Warrington, E. K. (2005). Abstract and concrete concepts have structurally different representational frameworks. *Brain*, *128*, 615–627. https://doi.org/10/cqfm9m

D'Argembeau, A., Cassol, H., Phillips, C., Balteau, E., Salmon, E., & Van der Linden, M. (2014). Brains creating stories of selves: The neural basis of autobiographical reasoning. *Social Cognitive and Affective Neuroscience*, *9*(5), 646–652. https://doi.org/10.1093/scan/nst028

Davis, C. P., Altmann, G. T. M., & Yee, E. (2020). Situational systematicity: A role for schema in understanding the differences between abstract and concrete concepts. *Cognitive Neuropsychology*, *37*(1–2), 142–153. https://doi.org/10.1080/02643294.2019.1710124

Desai, R. H. (2022). Are metaphors embodied? The neural evidence. *Psychological Research*, *86*(8), 2417–2433. https://doi.org/10/gnw5s5

Devlin, J., Chang, M.-W., Lee, K., & Toutanova, K. (2019). *BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding* (arXiv:1810.04805). arXiv. https://doi.org/10.48550/arXiv.1810.04805

Dove, G., Barca, L., Tummolini, L., & Borghi, A. M. (2022). Words have a weight: Language as a source of inner grounding and flexibility in abstract concepts. *Psychological Research*, *86*(8), 2451–2467. https://doi.org/10.1007/s00426-020-01438-6

Eickhoff, S. B., Bzdok, D., Laird, A. R., Kurth, F., & Fox, P. T. (2012). Activation likelihood estimation meta-analysis revisited. *Neuroimage*, *59*(3), 2349–2361. https://doi.org/10.1016/j.neuroimage.2011.09.017

Eickhoff, S. B., Laird, A. R., Grefkes, C., Wang, L. E., Zilles, K., & Fox, P. T. (2009). Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical estimates of spatial uncertainty. *Human Brain Mapping*, *30*(9), 2907–2926. https://doi.org/10.1002/hbm.20718

Epstein, R. A., & Baker, C. I. (2019). Scene Perception in the Human Brain. *Annual Review of Vision Science*, *5*(Volume 5, 2019), 373–397. https://doi.org/10.1146/annurev-vision-091718-014809

Fernandino, L., & Binder, J. R. (2024). How does the "default mode" network contribute to semantic cognition? *Brain and Language*, *252*, 105405. https://doi.org/10.1016/j.bandl.2024.105405

Fernandino, L., Binder, J. R., Desai, R. H., Pendl, S. L., Humphries, C. J., Gross, W. L., Conant, L. L., & Seidenberg, M. S. (2016). Concept representation reflects multimodal abstraction: A framework for embodied semantics. *Cerebral Cortex*, *26*(5), 2018–2034. https://doi.org/10.1093/cercor/bhv020

Gallese, V., & Lakoff, G. (2005). The brain's concepts: The role of the sensory-motor system in conceptual knowledge. *Cognitive Neuropsychology*, *22*(3–4), 455–479. https://doi.org/10.1080/02643290442000310

Giacobbe, C., Raimo, S., Cropano, M., & Santangelo, G. (2022). Neural correlates of embodied action language processing: A systematic review and meta-analytic study. *Brain Imaging and Behavior*, *16*(5), 2353–2374. https://doi.org/10.1007/s11682-022-00680-3

Giallanza, T., Campbell, D., Cohen, J. D., & Rogers, T. (2023). *An Integrated Model of Semantics and Control*. PsyArXiv. https://doi.org/10.31234/osf.io/jq7ta

Gibbs, R. W. (1994). *The poetics of mind: Figurative thought, language, and understanding*. Cambridge University Press.

Gilboa, A., & Marlatte, H. (2017). Neurobiology of Schemas and Schema-Mediated Memory. *Trends in Cognitive Sciences*, *21*(8), 618–631. https://doi.org/10.1016/j.tics.2017.04.013

Glenberg, A. M., & Robertson, D. A. (2000). Symbol grounding and meaning: A comparison of high-dimensional and embodied theories of meaning. *Journal of Memory and Language*, *43*(3), 379–401. https://doi.org/10/dsr5gr

Gurguryan, L., & Sheldon, S. (2019). Retrieval orientation alters neural activity during autobiographical memory recollection. *NeuroImage*, *199*, 534–544. https://doi.org/10.1016/j.neuroimage.2019.05.077

Hauk, O., Johnsrude, I., & Pulvermuller, F. (2004). Somatotopic representation of action words in human motor and premotor cortex. *Neuron*, *41*(2), 301–307. https://doi.org/10.1016/s0896-6273(03)00838-9

Heidlmayr, K., Weber, K., Takashima, A., & Hagoort, P. (2020). No title, no theme: The joined neural space between speakers and listeners during production and comprehension of multi-sentence discourse. *Cortex*, *130*, 111–126. https://doi.org/10.1016/j.cortex.2020.04.035

Hoffman, P. (2016). The meaning of 'life' and other abstract words: Insights from neuropsychology. *J Neuropsychol*, *10*, 317–343. https://doi.org/10.1111/jnp.12065

Hoffman, P., Binney, R. J., & Lambon Ralph, M. A. (2015). Differing contributions of inferior prefrontal and anterior temporal cortex to concrete and abstract conceptual knowledge. *Cortex*, *63*, 250–265. https://doi.org/10.1016/j.cortex.2014.09.001

Hoffman, P., Jefferies, E., & Lambon Ralph, M. A. (2010). Ventrolateral prefrontal cortex plays an executive regulation role in comprehension of abstract words: Convergent neuropsychological and repetitive TMS evidence. *Journal of Neuroscience*, *30*(46), 15450–15456. https://doi.org/10.1523/jneurosci.3783-10.2010

Hoffman, P., & Lambon Ralph, M. A. (2013). Shapes, scents and sounds: Quantifying the full multi-sensory basis of conceptual knowledge. *Neuropsychologia*, *51*(1), 14–25. https://doi.org/10/f4mns4

Hoffman, P., Lambon Ralph, M. A., & Rogers, T. T. (2013). Semantic diversity: A measure of semantic ambiguity based on variability in the contextual usage of words. *Behavior Research Methods*, *45*(3), 718–730. https://doi.org/10.3758/s13428-012-0278-x

Jackson, R. L. (2021). The neural correlates of semantic control revisited. *Neuroimage*, *224*, 117444. https://doi.org/10.1016/j.neuroimage.2020.117444

Johnson-Laird, P. N. (1988). *The computer and the mind: An introduction to cognitive science*. Harvard University Press.

Joue, G., Boven, L., Willmes, K., Evola, V., Demenescu, L. R., Hassemer, J., Mittelberg, I., Mathiak, K., Schneider, F., & Habel, U. (2020). Metaphor processing is supramodal semantic processing: The role of the bilateral lateral temporal regions in multimodal communication. *Brain and Language*, *205*, 104772. https://doi.org/10.1016/j.bandl.2020.104772

Kousta, S. T., Vigliocco, G., Vinson, D. P., Andrews, M., & Del Campo, E. (2011). The Representation of Abstract Words: Why Emotion Matters. *J Exp Psychol Gen*, *140*(1), 14–34. https://doi.org/10/brjczk

Lakoff, G., & Johnson, M. (1999). *Philosophy in the flesh: The embodied mind and its challenge to western thought*. Basic Books.

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and computational bases of semantic cognition. *Nature Reviews Neuroscience*, *18*, 42–55. https://doi.org/10/gmhhtx

Lee, S., Parthasarathi, T., & Kable, J. W. (2021). The Ventral and Dorsal Default Mode Networks Are Dissociably Modulated by the Vividness and Valence of Imagined Events. *Journal of Neuroscience*, *41*(24), 5243–5250. https://doi.org/10.1523/JNEUROSCI.1273-20.2021

Lenci, A., Lebani, G. E., & Passaro, L. C. (2018). The emotions of abstract words: A distributional semantic analysis. *Topics in Cognitive Science*, *10*(3), 550–572. https://doi.org/10.1111/tops.12335

Louwerse, M. M. (2011). Symbol interdependency in symbolic and embodied cognition. *Topics in Cognitive Science*, *3*(2), 273–302. https://doi.org/10.1111/j.1756-8765.2010.01106.x

Lynott, D., Connell, L., Brysbaert, M., Brand, J., & Carney, J. (2020). The Lancaster Sensorimotor Norms: Multidimensional measures of perceptual and action strength for 40,000 English words. *Behavior Research Methods*, *52*(3), 1271–1291. https://doi.org/10.3758/s13428-019-01316-z

Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G., Eickhoff, S. B., Castellanos, F. X., & Petrides, M. (2016). Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*, *113*(44), 12574–12579. https://doi.org/10.1073/pnas.1608282113

Martin, A. (2016). GRAPES—Grounding representations in action, perception, and emotion systems: How object properties and categories are represented in the human brain. *Psychonomic Bulletin & Review*, *23*, 979–990. https://doi.org/10.3758/s13423-015-0842-3

Martin, A., Haxby, J. V., Lalonde, F. M., Wiggs, C. L., & Ungerleider, L. G. (1995). Discrete cortical regions associated with knowledge of color and knowledge of action. *Science*, *270*, 102–105. https://doi.org/10/fv2rf4

Mckiernan, K. A., Kaufman, J. N., Kucera-Thompson, J., & Binder, J. R. (2003). A parametric manipulation of factors affecting task-induced deactivation in functional neuroimaging. *Journal of Cognitive Neuroscience*, *15*(3), 394–408. https://doi.org/10/b6j6ht

McRae, K., Nedjadrasul, D., Pau, R., Lo, B. P.-H., & King, L. (2018). Abstract Concepts and Pictures of Real-World Situations Activate One Another. *Topics in Cognitive Science*, *10*(3), 518–532. https://doi.org/10.1111/tops.12328

Mikolov, T., Chen, K., Corrado, G., & Dean, J. (2013). Efficient estimation of word representations in vector space. *arXiv Preprint arXiv:1301.3781*.

Morales, M., Patel, T., Tamm, A., Pickering, M., & Hoffman, P. (2022). Similar neural networks respond to coherence during comprehension and production of discourse. *Cerebral Cortex*, *32*(19), 4317–4330. https://doi.org/10.1093/cercor/bhab485

Mullally, S. L., & Maguire, E. A. (2014). Memory, Imagination, and Predicting the Future: A Common Brain Mechanism? *The Neuroscientist*, *20*(3), 220–234. https://doi.org/10.1177/1073858413495091

Newcombe, P. I., Campbell, C., Siakaluk, P. D., & Pexman, P. M. (2012). Effects of emotional and sensorimotor knowledge in semantic processing of concrete and abstract nouns. *Frontiers in Human Neuroscience*, *6*, 275. https://doi.org/10.3389/fnhum.2012.00275

Newell, A. (1980). Physical symbol systems. *Cognitive Science*, *4*(2), 135–183.

Paivio, A. (1986). *Mental representations: A dual-coding approach*. Oxford University Press.

Patel, T., Morales, M., Pickering, M. J., & Hoffman, P. (2023). A common neural code for meaning in discourse production and comprehension. *NeuroImage*, *279*, 120295. https://doi.org/10.1016/j.neuroimage.2023.120295

Pecher, D. (2018). Curb your embodiment. *Topics in Cognitive Science*, *10*(3), 501–517. https://doi.org/10/gd5pgp

Pereira, F., Gershman, S., Ritter, S., & Botvinick, M. (2016). A comparative evaluation of off-the-shelf distributed semantic representations for modelling behavioural data. *Cognitive Neuropsychology*, *33*(3–4), 175–190. https://doi.org/10/ghcv8k

Pexman, P. M., Diveica, V., & Binney, R. J. (2022). Social semantics: The organization and grounding of abstract concepts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *378*(1870), 20210363. https://doi.org/10.1098/rstb.2021.0363

Poerio, G. L., Sormaz, M., Wang, H.-T., Margulies, D., Jefferies, E., & Smallwood, J. (2017). The role of the default mode network in component processes underlying the wandering mind. *Social Cognitive and Affective Neuroscience*, *12*(7), 1047–1062. https://doi.org/10.1093/scan/nsx041

Pulvermüller, F. (2013). How neurons make meaning: Brain mechanisms for embodied and abstract-symbolic semantics. *Trends in Cognitive Sciences*, *17*(9), 458–470. https://doi.org/10.1016/j.tics.2013.06.004

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences*, *98*(2), 676–682. https://doi.org/10.1073/pnas.98.2.676

Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour. *Nature Reviews Neuroscience*, *13*(10), 713–726. https://doi.org/10.1038/nrn3338

Reilly, J., Shain, C., Borghesani, V., Kuhnke, P., Vigliocco, G., Peelle, J., Mahon, B., Buxbaum, L., Majid, A., Brysbaert, M., Borghi, A., De Deyne, S., Dove, G., Papeo, L., Pexman, P., Poeppel, D., Lupyan, G., Boggio, P., Hickock, G., … Vinson, D. (2023). *What We Mean When We Say Semantic: A Consensus Statement on the Nomenclature of Semantic Memory*. PsyArXiv. https://doi.org/10.31234/osf.io/xrnb2

Rubin, T. N., Koyejo, O., Gorgolewski, K. J., Jones, M. N., Poldrack, R. A., & Yarkoni, T. (2017). Decoding brain activity using a large-scale probabilistic functional-anatomical atlas of human cognition. *PLoS Computational Biology*, *13*(10), e1005649. https://doi.org/10/gcgdtm

Saygin, A. P., McCullough, S., Alac, M., & Emmorey, K. (2010). Modulation of BOLD response in motion-sensitive lateral temporal cortex by real and fictive motion sentences. *Journal of Cognitive Neuroscience*, *22*(11), 2480–2490. https://doi.org/10.1162/jocn.2009.21388

Schwanenflugel, P. J., Harnishfeger, K. K., & Stowe, R. W. (1988). Context Availability and Lexical Decisions for Abstract and Concrete Words. *Journal of Memory and Language*, *27*(5), 499–520. https://doi.org/10/dt3z52

Schwanenflugel, P. J., & Shoben, E. J. (1983). Differential Context Effects in the Comprehension of Abstract and Concrete Verbal Materials. *Journal of Experimental Psychology: Learning, Memory and Cognition*, *9*(1), 82–102. https://doi.org/10/d4jwbt

Shao, X., Krieger-Redwood, K., Zhang, M., Hoffman, P., Lanzoni, L., Leech, R., Smallwood, J., & Jefferies, E. (2024). Distinctive and complementary roles of default mode network subsystems in semantic cognition. *Journal of Neuroscience*, *44*(20), e1907232024. https://doi.org/10.1523/JNEUROSCI.1907-23.2024

Silson, E. H., Steel, A., Kidder, A., Gilmore, A. W., & Baker, C. I. (2019). Distinct subdivisions of human medial parietal cortex support recollection of people and places. *eLife*, *8*, e47391. https://doi.org/10.7554/eLife.47391

Smallwood, J., Bernhardt, B. C., Leech, R., Bzdok, D., Jefferies, E., & Margulies, D. S. (2021). The default mode network in cognition: A topographical perspective. *Nature Reviews Neuroscience*, *22*(8), 503–513. https://doi.org/10.1038/s41583-021-00474-4

Smallwood, J., Tipper, C., Brown, K., Baird, B., Engen, H., Michaels, J. R., Grafton, S., & Schooler, J. W. (2013). Escaping the here and now: Evidence for a role of the default mode network in perceptually decoupled thought. *NeuroImage*, *69*, 120–125. https://doi.org/10.1016/j.neuroimage.2012.12.012

Tamir, D. I., Bricker, A. B., Dodell-Feder, D., & Mitchell, J. P. (2016). Reading fiction and reading minds: The role of simulation in the default network. *Social Cognitive and Affective Neuroscience*, *11*(2), 215–224. https://doi.org/10.1093/scan/nsv114

Thompson-Schill, S. L., Aguirre, G. K., D'Esposito, M., & Farah, M. J. (1999). A neural basis for category and modality specificity of semantic knowledge. *Neuropsychologia*, *37*, 671–676. https://doi.org/10/fts7nx

Tong, J., Binder, J. R., Humphries, C., Mazurchuk, S., Conant, L. L., & Fernandino, L. (2022). A Distributed Network for Multimodal Experiential Representation of Concepts. *The Journal of Neuroscience*, *42*(37), 7121–7130. https://doi.org/10.1523/JNEUROSCI.1243-21.2022

Troche, J., Crutch, S., & Reilly, J. (2014). Clustering, hierarchical organization, and the topography of abstract and concrete nouns. *Frontiers in Psychology*, *5*, 360. https://doi.org/10/gbf4r5

Turkeltaub, P. E., Eickhoff, S. B., Laird, A. R., Fox, M., Wiener, M., & Fox, P. (2012). Minimizing within-experiment and within-group effects in activation likelihood estimation meta-analyses. *Human Brain Mapping*, *33*(1), 1–13. https://doi.org/10.1002/hbm.21186

van Dam, W. O., van Dijk, M., Bekkering, H., & Rueschemeyer, S. A. (2012). Flexibility in embodied lexical-semantic representations. *Human Brain Mapping*, *33*(10), 2322–2333. https://doi.org/10/bw8sdp

Vigliocco, G., Kousta, S. T., Della Rosa, P. A., Vinson, D. P., Tettamanti, M., Devlin, J. T., & Cappa, S. F. (2014). The neural representation of abstract words: The role of emotion. *Cerebral Cortex*, *24*, 1767–1777. https://doi.org/10.1093/cercor/bht025

Vigliocco, G., Meteyard, L., Andrews, M., & Kousta, S. (2009). Toward a theory of semantic representation. *Language and Cognition*, *1*(02), 219–247. https://doi.org/10/fvwk3v

Villani, C., Lugli, L., Liuzza, M. T., & Borghi, A. M. (2019). Varieties of abstract concepts and their multiple dimensions. *Language and Cognition*, *11*(3), 403–430. https://doi.org/10.1017/langcog.2019.23

Vitello, S., & Rodd, J. M. (2015). Resolving semantic ambiguities in sentences: Cognitive processes and brain mechanisms. *Language and Linguistics Compass*, *9*(10), 391–405. https://doi.org/10.1111/lnc3.12160

Wang, J., Conder, J. A., Blitzer, D. N., & Shinkareva, S. V. (2010). Neural Representation of Abstract and Concrete Concepts: A Meta-Analysis of Neuroimaging Studies. *Hum Brain Mapp*, *31*(10), 1459–1468. https://doi.org/10/cw55dk

Wang, X., & Bi, Y. (2021). Idiosyncratic tower of Babel: Individual differences in word-meaning representation increase as word abstractness increases. *Psychological Science*, *32*(10), 1617–1635. https://doi.org/10.1177/09567976211003877

Wang, X., Wu, W., Ling, Z., Xu, Y., Fang, Y., Wang, X., Binder, J. R., Men, W., Gao, J.-H., & Bi, Y. (2018). Organizational principles of abstract words in the human brain. *Cerebral Cortex*, *28*(12), 4305–4318. https://doi.org/10.1093/cercor/bhx283

Watson, C. E., Cardillo, E. R., Ianni, G. R., & Chatterjee, A. (2013). Action concepts in the brain: An activation likelihood estimation meta-analysis. *Journal of Cognitive Neuroscience*, *25*(8), 1191–1205. https://doi.org/10/ghms3p

Wiemer-Hastings, K., & Xu, X. (2005). Content differences for abstract and concrete concepts. *Cognitive Science*, *29*(5), 719–736. https://doi.org/10/cd4zxg

Winter, B. (2023). Abstract concepts and emotion: Cross-linguistic evidence and arguments against affective embodiment. *Philosophical Transactions of the Royal Society B*, *378*(1870), 20210368. https://doi.org/10/gs4td3

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature Methods*, *8*(8), 665–670. https://doi.org/10.1038/nmeth.1635

Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165. https://doi.org/10.1152/jn.00338.2011

Yeshurun, Y., Nguyen, M., & Hasson, U. (2021). The default mode network: Where the idiosyncratic self meets the shared social world. *Nature Reviews Neuroscience*, *22*(3), 181–192. https://doi.org/10.1038/s41583-020-00420-w

Yu, S., Gu, C., Huang, K., & Li, P. (2024). Predicting the next sentence (not word) in large language models: What model-brain alignment tells us about discourse comprehension. *Science Advances*, *10*(21), eadn7744. https://doi.org/10.1126/sciadv.adn7744

Zeki, S. (2004). Thirty years of a very special visual area, Area V5. *The Journal of Physiology*, *557*(Pt 1), 1–2. https://doi.org/10.1113/jphysiol.2004.063040

Figure 1: Process used to identify studies

Figure 2: Functional networks used to interpret concreteness effects *DMN = default mode network; FT = fronto-temporal; MT = medial temporal.*

Figure 3: Meta-analysis of 72 studies of concreteness effects *Greater activation for abstract concepts (A>C) shown in hot colours and for concrete concepts (C>A) shown in cool colours. Results are shown at a voxel-level threshold of p<0.005, corrected for multiple comparisons at the cluster level. Bar plots indicate the proportion of voxels showing effects that fall within each of the networks shown in Figure 2. Wordclouds show Neurosynth terms most associated with A>C and C>A effects.*

Figure 4: Effects of stimulus type *Figure shows regions where A>C (left side) and C>A (right side) effects were more likely for Sentence vs. Word studies. Bar plots indicate the proportion of voxels showing effects that fall within each of the networks shown in Figure 2. Wordclouds show Neurosynth terms most associated with A>C and C>A effects.*

Figure 5: Effects of task type *Figure shows regions where A>C (left side) and C>A (right side) effects were more likely for Semantics vs. Non-semantic studies. Bar plots indicate the proportion of voxels showing effects that fall within each of the networks shown in Figure 2. Wordclouds show Neurosynth terms most associated with A>C and C>A effects. N.S. = nothing significant.*

Table 1: Summary of predictions about concreteness effects and degree to which they were supported in the meta-analysis
✓ *indicates strong support;* (✓) *indicates weak or partial support;* ✗*indicates no support.*

Claim	C>A predictions	Supported?	A>C predictions	Supported?
Understanding concrete language is more reliant on activating its sensorimotor correlates	Action network especially sentences Visual network	(✓) ✓ ✓		
Understanding abstract language is more reliant on activating its social and emotional associations			Social network especially sentences Emotional network especially sentences	✓ ✓ ✗ ✗
Understanding abstract language is more reliant on activating linguistic knowledge			Language network especially non-semantic tasks	✓ ✗
The meaning of abstract language is more variable than that of concrete language			Semantic Control especially words especially semantic tasks	✓ ✓ (✓)
Situation models can be more engaged when processing concrete or abstract language, depending on the level of contextual support	Situation network? especially words?	✓ (✓)	Situation network? especially sentences?	✓ ✓

Table 2: Peak effect co-ordinates for the main analysis

Cluster	Regions	Volume (mm³)	x	y	z	ALE value
	Abstract > Concrete					
1	Left inferior frontal gyrus & lateral anterior temporal lobe	18448	-54	22	6	0.053
			-50	10	-22	0.051
			-48	24	-8	0.042
			-50	8	-36	0.023
			-58	-8	-18	0.020
			-58	-10	-10	0.018
2	Left posterior middle and superior temporal gyri	7408	-58	-40	0	0.050
			-60	-26	8	0.014
			-56	-26	22	0.012
			-60	-22	16	0.012
3	Right lateral anterior temporal lobe	5200	56	0	-18	0.032
			52	8	-24	0.022
			52	-12	-14	0.019
			54	-12	-4	0.018
			62	-2	-4	0.016
4	Posterior cingulate & precuneus	3328	2	-54	30	0.025
			-10	-52	36	0.022
			-4	-40	32	0.014
5	Left parietal operculum	2216	-48	-38	22	0.022
	Concrete > Abstract					
1	Left posterior middle and inferior temporal gyri, fusiform & parahippocampal gyrus	14520	-30	-36	-18	0.049
			-24	-32	-22	0.045
			-54	-62	-2	0.033
			-52	-60	-12	0.029
2	Left angular gyrus & lateral occipital cortex	9000	-38	-74	34	0.036
			-44	-76	24	0.035
			-34	-82	28	0.023
			-38	-74	48	0.015
3	Posterior cingulate & precuneus	4800	-8	-54	10	0.040
			8	-54	10	0.021
			-14	-50	26	0.014
4	Right angular gyrus	4424	46	-66	24	0.030
			44	-60	40	0.021
			48	-68	46	0.017
5	Left inferior frontal sulcus	3368	-42	32	12	0.034
6	Left superior frontal gyrus	3296	-20	16	52	0.029

			-24	28	52	0.018
			-26	8	50	0.018
			-28	34	46	0.017
			-24	36	48	0.017
7	Right parahippocampal gyrus	2848	32	-30	-18	0.032
			30	-40	-24	0.018
8	Left supramarginal gyrus	2416	-60	-30	38	0.044

Supplementary Materials

Supplementary Table 1: Details of all studies included in the meta-analysis

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Fletcher	1996	PET	6	Paired associate recall	Word pairs varying in semantic relatedness (English)	1	6	Word	Non-semantic	Yes
D'Esposito	1997	fMRI	7	Imagine (C); Passive listening (A)	Single words (English)	2	3	Word	Non-semantic	
Mellet	1998	PET	8	Understand (A); Understand and imagine (C only)	Words and their definitions (English)	4	9	Sentence	Non-semantic	
Kiehl	1999	fMRI	6	Lexical decision	Single words (English)	1		Word	Non-semantic	
Perani	1999	PET	14	Lexical decision	Single nouns and verbs (Italian)	7		Word	Non-semantic	
Jessen	2000	fMRI	14	Memory encoding (view to remember)	Single words (German)	2	4	Word	Non-semantic	Yes
Wise	2000	PET	18	Passive listening; Semantic similarity judgement; Passive listening/reading	Single words; triads; single words (English)		2	Word		
Tyler	2001	PET	9	Lexical decision	Single nouns and verbs (English)	1	1	Word	Non-semantic	
Grossman	2002	fMRI	16	Pleasantness judgement	Single words (English); C = animals, implements	5		Word	Semantic	

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Noppeney	2002	PET	12	Semantic property judgement	Single words (English); C = visual or auditory associates; A = non-emotional	2	1	Word	Semantic	
Fiebach	2003	fMRI	12	Lexical decision	Single words (German); C = inanimate	1	1	Word	Non-semantic	
Kounis	2003	fMRI	16	Pleasantness judgement	Single words (English); C = animals, implements	1	4	Word	Semantic	
Giesbrecht	2004	fMRI	8	Semantic relatedness judgement	Word pairs (English), related and unrelated		4	Word	Semantic	
Noppeney	2004	fMRI	15	Semantic similarity judgement	Triads (English); C = visual attributes, sounds, hand movements	4		Word	Semantic	
Whatmough	2004	PET	15	Semantic relatedness judgement	Word pairs (English), related and unrelated	1	1	Word	Semantic	
Binder1	2005	fMRI	24	Lexical decision	Single words (English)	9	15	Word	Non-semantic	
Binder2	2005	fMRI	24	Reading aloud	Single words (English), regular and irregular spellings	11	36	Word	Non-semantic	
Sabsevitz	2005	fMRI	28	Semantic similarity judgement	Triads (English)	10	26	Word	Semantic	
Tettamanti	2005	fMRI	17	Passive listening	Single sentences (Italian); C = hand, mouth and foot actions	1	1	Sentence	Non-semantic	
Wallentin	2005	fMRI	18	Sensibility judgement (understood/not)	Single sentences (Danish) describing real or fictive motion	23	10	Sentence	Semantic	
Bedny	2006	fMRI	13	Semantic similarity judgement	Triads (English), nouns and verbs	2	3	Word	Semantic	

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Fliessbach	2006	fMRI	21	Memory encoding (view to remember) & recognition	Single words (German)	2	3	Word	Non-semantic	Yes
Harris	2006	fMRI	22	Pleasantness judgement	Single words (English); A = mental states and metaphysical states	2	7	Word	Semantic	
Goldberg	2007	fMRI	14	Property verification	Animal names (English) paired with sensory (C) or verbal (A) properties	12	4	Word	Semantic	
Pexman	2007	fMRI	20	Consumability judgement	Ambiguous and unambiguous single words (English)	25		Word	Semantic	
Rüschemeyer	2007	fMRI	20	Lexical decision	Simple verbs (German)		9	Word	Non-semantic	
Hauk	2008	fMRI	21	Passive reading	Single words (English)		5	Word	Non-semantic	
Kuperberg	2008	fMRI	16	Sensibility judgement	Single sentences with congruous or incongruous endings (English)	7	18	Sentence	Semantic	
Tettamanti	2008	fMRI	18	Passive listening	Affirmative or negative single sentences (Italian)	6	10	Sentence	Non-semantic	
Desai	2009	fMRI	30	Sensibility judgement	Single sentences (English); C = motor-related and visual-related	29	4	Sentence	Semantic	
Mestres-Misse	2009	fMRI	15	Think about meaning	Sentence pairs ending in novel or known word (Spanish)	1	2	Sentence	Non-semantic	Yes

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Olivetti Belardinelli	2009	fMRI	9	Mental imagery (A and C)	Phrases describing experiences and concepts (Italian)		23	Sentence	Non-semantic	
Graves	2010	fMRI	20	Reading aloud	Single words (English)	2	3	Word	Non-semantic	
van Dam	2010	fMRI	14	Semantic decision (mouth action or not)	Single verbs (Dutch)		5	Word	Semantic	
Desai	2011	fMRI	22	Sensibility judgement	Single sentences (literal)	4	9	Sentence	Semantic	
Rodriguez-Ferriero	2011	fMRI	14	Passive reading	Single verbs (Spanish)	9	5	Word	Non-semantic	
Straube	2011	fMRI	17	Passive listening	Single sentences (German)	7	14	Sentence	Non-semantic	
Zhuang	2011	fMRI	14	Lexical decision	Single words (English)		1	Word	Non-semantic	
Kana	2012	fMRI	24	Reading with comprehension questions	Single sentences; C = actions; A = mental states	2	8	Sentence	Non-semantic	
Papeo	2012	fMRI	18	Memory encoding (view to remember)	Single verbs (Italian); C = hand actions; A = mental states	1	23	Word	Non-semantic	
Shafto	2012	fMRI	16	Lexical decision	Single words (English)	1		Word	Non-semantic	
van Dam	2012	fMRI	19	Action or colour decision	Single nouns (Dutch)		18	Word	Semantic	
Desai	2013	fMRI	27	Sensibility judgement	Single sentences (literal)	3	12	Sentence	Semantic	
Gilead	2013	fMRI	20	Passive reading	Single sentences varying in tense (Hebrew)	15	5	Sentence	Non-semantic	

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Nagels	2013	fMRI	17	Passive listening (unrelated perceptual task)	Single sentences (German)	9	8	Sentence	Non-semantic	
Sakreida	2013	fMRI	25	Semantic judgement (detect oddball foot-related stimuli)	Noun-verb phrases (German; concrete-concrete vs. abstract-abstract)	1	3	Sentence	Semantic	
Straube	2013	fMRI	20	Semantic judgement (person vs. object related)	Single sentences (German); C = object-related; A = social-related	6	1	Sentence	Semantic	
Hayashi	2014	fMRI	16	Generate visual imagery (A and C)	Single words (Japanese)	2	7	Word	Non-semantic	
Roxbury	2014	fMRI	17	Lexical decision	Single nouns (English)		5	Word	Non-semantic	
Tomasino	2014	fMRI	24	Mental imagery & Pleasantness judgement; Letter search	Single verbs (Italian)	18	9	Word		
Vigliocco	2014	fMRI	17	Lexical decision	Single nouns (English)	1		Word	Non-semantic	
Hoffman	2015	fMRI	19	Semantic similarity judgement	Triads with single sentence primes (English)	21	13		Semantic	
Binney	2016	fMRI	19	Semantic relatedness judgement	Triads (English); C = animal functions, A = social concepts	12		Word	Semantic	
Caplan	2016	fMRI	20	Paired associates learning	Noun pairs (English)	14	6	Word	Non-semantic	Yes

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Ghio	2016	fMRI	36	Sentence verification (1-back task)	Single sentences (Italian); C = hand, mouth and foot actions, A = emotions, mental states, mathematical concepts	9		Sentence	Non-semantic	
Kumar	2016	fMRI	20	Orthographic judgement	Single nouns (Hindi)	4		Word	Non-semantic	
van Dam	2016	fMRI	14	Passive reading	Single sentences (English)	16	11	Sentence	Non-semantic	
Westbury	2016	fMRI	14	Lexical decision	Single words (English)	10	1	Word	Non-semantic	
Cesare	2017	fMRI	22	Passive listening	Single verbs (Italian)		16	Word	Non-semantic	
Mayer	2017	fMRI	42	Translation to native language	Single nouns (German)	2		Word	Semantic	
Auger	2018	fMRI	32	Sensibility judgement	Single sentences (English); C = landmark-related and action-related	3	3	Sentence	Semantic	
Clark	2018	fMRI	45	Memory encoding (view to remember)	Single words (English); C = object words and scene words	4	21	Word	Non-semantic	Yes
Della Rosa	2018	fMRI	26	Lexical decision	Single nouns (Italian)	12	1	Word	Non-semantic	
Lin	2018	fMRI	19	Semantic relatedness judgement	Verb pairs (Chinese; high and low social)		12	Word	Semantic	
Pauligk	2019	fMRI	21	Lexical decision	Single words (German)	2		Word	Non-semantic	

1st author	Year	Imaging	N	Task	Stimuli	A>C peaks	C>A peaks	Stimuli	Task	Episodic Memory
Wang	2019	fMRI	22	Semantic relatedness judgement	Triads (Chinese); A = non-social neutral words	23		Word	Semantic	
Garbarini	2020	fMRI	17	Naming to definition	Definitions of words (Italian)	1	3	Sentence	Semantic	
Joue	2020	fMRI	32	Concreteness decision	Single sentences (German; literal only)	25	1	Sentence	Semantic	
Meersmans	2020	fMRI	26	Repetition; reading	Single nouns (Dutch)	5		Word	Non-semantic	
Tian	2020	fMRI	29	Passive reading	Noun-verb phrases (Chinese; literal only)	1	4	Sentence	Non-semantic	
Choudhury	2021	fMRI	36	Gesture-sentence relatedness decision	Single sentences (German)	3		Sentence	Semantic	
Meersmans	2022	fMRI	22	Concreteness decision; valence decision	Single nouns (Dutch)	15	19	Word	Semantic	
TOTAL			**1400**			**435**	**459**			

Supplementary Figure 1: Results of meta-analysis excluding 7 studies that used episodic memory tasks

List of studies included in meta-analysis

Auger, S. D., & Maguire, E. A. (2018). Retrosplenial Cortex Indexes Stability beyond the Spatial Domain. *The Journal of Neuroscience*, *38*(6), 1472–1481. https://doi.org/10.1523/JNEUROSCI.2602-17.2017

Bedny, M., & Thompson-Schill, S. L. (2006). Neuroanatomically separable effects of imageability and grammatical class during single-word comprehension. *Brain and Language*, *98*(2), 127–139. https://doi.org/10.1016/j.bandl.2006.04.008

Binder, J. R., Medler, D. A., Desai, R., Conant, L. L., & Liebenthal, E. (2005). Some neurophysiological constraints on models of word naming. *NeuroImage*, *27*(3), 677–693. https://doi.org/10.1016/j.neuroimage.2005.04.029

Binder, J. R., Westbury, C. F., McKiernan, K. A., Possing, E. T., & Medler, D. A. (2005). Distinct Brain Systems for Processing Concrete and Abstract Concepts. *Journal of Cognitive Neuroscience*, *17*(6), 905–917. https://doi.org/10.1162/0898929054021102

Binney, R. J., Hoffman, P., & Lambon Ralph, M. A. (2016). Mapping the Multiple Graded Contributions of the Anterior Temporal Lobe Representational Hub to Abstract and Social Concepts: Evidence from Distortion-corrected fMRI. *Cerebral Cortex*, *26*(11), 4227–4241. https://doi.org/10.1093/cercor/bhw260

Caplan, J. B., & Madan, C. R. (2016). Word Imageability Enhances Association-memory by Increasing Hippocampal Engagement. *Journal of Cognitive Neuroscience*, *28*(10), 1522–1538. https://doi.org/10.1162/jocn_a_00992

Choudhury, M., Steines, M., Nagels, A., Riedl, L., Kircher, T., & Straube, B. (2021). Neural Basis of Speech-Gesture Mismatch Detection in Schizophrenia Spectrum Disorders. *Schizophrenia Bulletin*, *47*(6), 1761–1771. https://doi.org/10.1093/schbul/sbab059

Clark, I. A., Kim, M., & Maguire, E. A. (2018). Verbal Paired Associates and the Hippocampus: The Role of Scenes. *Journal of Cognitive Neuroscience*, *30*(12), 1821–1845. https://doi.org/10.1162/jocn_a_01315

Della Rosa, P. A., Catricalà, E., Canini, M., Vigliocco, G., & Cappa, S. F. (2018). The left inferior frontal gyrus: A neural crossroads between abstract and concrete knowledge. *NeuroImage*, *175*, 449–459. https://doi.org/10.1016/j.neuroimage.2018.04.021

Desai, R. H., Binder, J. R., Conant, L. L., Mano, Q. R., & Seidenberg, M. S. (2011). The Neural Career of Sensory-motor Metaphors. *Journal of Cognitive Neuroscience*, *23*(9), 2376–2386. https://doi.org/10.1162/jocn.2010.21596

Desai, R. H., Binder, J. R., Conant, L. L., & Seidenberg, M. S. (2010). Activation of Sensory-Motor Areas in Sentence Comprehension. *Cerebral Cortex*, *20*(2), 468–478. https://doi.org/10.1093/cercor/bhp115

Desai, R. H., Conant, L. L., Binder, J. R., Park, H., & Seidenberg, M. S. (2013). A piece of the action: Modulation of sensory-motor regions by action idioms and metaphors. *NeuroImage*, *83*, 862–869. https://doi.org/10.1016/j.neuroimage.2013.07.044

D'Esposito, M., Detre, J. A., Aguirre, G. K., Stallcup, M., Alsop, D. C., Tippet, L. J., & Farah, M. J. (1997). A functional MRI study of mental image generation. *Neuropsychologia*, *35*(5), 725–730. https://doi.org/10.1016/S0028-3932(96)00121-2

Di Cesare, G., Errante, A., Marchi, M., & Cuccio, V. (2017). Language for action: Motor resonance during the processing of human and robotic voices. *Brain and Cognition*, *118*, 118–127. https://doi.org/10.1016/j.bandc.2017.08.001

Fiebach, C. J., & Friederici, A. D. (2004). Processing concrete words: FMRI evidence against a specific right-hemisphere involvement. *Neuropsychologia*, *42*(1), 62–70. https://doi.org/10.1016/S0028-3932(03)00145-3

Fletcher, P. C., Shallice, T., Frith, C. D., Frackowiak, R. S. J., & Dolan, R. J. (1996). Brain activity during memory retrieval: The influence of imagery and semantic cueing. *Brain*, *119*(5), 1587–1596. https://doi.org/10.1093/brain/119.5.1587

Fliessbach, K., Weis, S., Klaver, P., Elger, C. E., & Weber, B. (2006). The effect of word concreteness on recognition memory. *NeuroImage*, *32*(3), 1413–1421. https://doi.org/10.1016/j.neuroimage.2006.06.007

Garbarini, F., Calzavarini, F., Diano, M., Biggio, M., Barbero, C., Radicioni, D. P., Geminiani, G., Sacco, K., & Marconi, D. (2020). Imageability effect on the functional brain activity during a naming to definition task. *Neuropsychologia*, *137*, 107275. https://doi.org/10.1016/j.neuropsychologia.2019.107275

Ghio, M., Vaghi, M. M. S., Perani, D., & Tettamanti, M. (2016). Decoding the neural representation of fine-grained conceptual categories. *NeuroImage*, *132*, 93–103. https://doi.org/10.1016/j.neuroimage.2016.02.009

Giesbrecht, B. (2004). Separable Effects of Semantic Priming and Imageability on Word Processing in Human Cortex. *Cerebral Cortex*, *14*(5), 521–529. https://doi.org/10.1093/cercor/bhh014

Gilead, M., Liberman, N., & Maril, A. (2013). The language of future-thought: An fMRI study of embodiment and tense processing. *NeuroImage*, *65*, 267–279. https://doi.org/10.1016/j.neuroimage.2012.09.073

Goldberg, R. F., Perfetti, C. A., Fiez, J. A., & Schneider, W. (2007). Selective Retrieval of Abstract Semantic Knowledge in Left Prefrontal Cortex. *The Journal of Neuroscience*, *27*(14), 3790–3798. https://doi.org/10.1523/JNEUROSCI.2381-06.2007

Graves, W. W., Desai, R., Humphries, C., Seidenberg, M. S., & Binder, J. R. (2010). Neural Systems for Reading Aloud: A Multiparametric Approach. *Cerebral Cortex*, *20*(8), 1799–1815. https://doi.org/10.1093/cercor/bhp245

Grossman, M., Koenig, P., DeVita, C., Glosser, G., Alsop, D., Detre, J., & Gee, J. (2002). The Neural Basis for Category-Specific Knowledge: An fMRI Study. *NeuroImage*, *15*(4), 936–948. https://doi.org/10.1006/nimg.2001.1028

Harris, G. J., Chabris, C. F., Clark, J., Urban, T., Aharon, I., Steele, S., McGrath, L., Condouris, K., & Tager-Flusberg, H. (2006). Brain activation during semantic processing in autism spectrum disorders via functional magnetic resonance imaging. *Brain and Cognition*, *61*(1), 54–68. https://doi.org/10.1016/j.bandc.2005.12.015

Supplementary Materials

Hauk, O., Davis, M. H., Kherif, F., & Pulvermüller, F. (2008). Imagery or meaning? Evidence for a semantic origin of category-specific brain activity in metabolic imaging. *European Journal of Neuroscience*, *27*(7), 1856–1866. https://doi.org/10.1111/j.1460-9568.2008.06143.x

Hayashi, A., Okamoto, Y., Yoshimura, S., Yoshino, A., Toki, S., Yamashita, H., Matsuda, F., & Yamawaki, S. (2014). Visual imagery while reading concrete and abstract Japanese kanji words: An fMRI study. *Neuroscience Research*, *79*, 61–66. https://doi.org/10.1016/j.neures.2013.10.007

Hoffman, P., Binney, R. J., & Lambon Ralph, M. A. (2015). Differing contributions of inferior prefrontal and anterior temporal cortex to concrete and abstract conceptual knowledge. *Cortex*, *63*, 250–266. https://doi.org/10.1016/j.cortex.2014.09.001

Jessen, F., Heun, R., Erb, M., Granath, D.-O., Klose, U., Papassotiropoulos, A., & Grodd, W. (2000). The Concreteness Effect: Evidence for Dual Coding and Context Availability. *Brain and Language*, *74*(1), 103–112. https://doi.org/10.1006/brln.2000.2340

Joue, G., Boven, L., Willmes, K., Evola, V., Demenescu, L. R., Hassemer, J., Mittelberg, I., Mathiak, K., Schneider, F., & Habel, U. (2020). Metaphor processing is supramodal semantic processing: The role of the bilateral lateral temporal regions in multimodal communication. *Brain and Language*, *205*, 104772. https://doi.org/10.1016/j.bandl.2020.104772

Kana, R. K., Blum, E. R., Ladden, S. L., & Ver Hoef, L. W. (2012). "How to do things with Words": Role of motor cortex in semantic representation of action words. *Neuropsychologia*, *50*(14), 3403–3409. https://doi.org/10.1016/j.neuropsychologia.2012.09.006

Kiehl, K. A., Liddle, P. F., Smith, A. M., Mendrek, A., Forster And, B. B., & Hare, R. D. (1999). Neural pathways involved in the processing of concrete and abstract words. *Human Brain Mapping*, *7*(4), 225–233. https://doi.org/10.1002/(SICI)1097-0193(1999)7:4<225::AID-HBM1>3.0.CO;2-P

Kounios, J., Koenig, P., Glosser, G., DeVita, C., Dennis, K., Moore, P., & Grossman, M. (2003). Category-specific medial temporal lobe activation and the consolidation of semantic memory: Evidence from fMRI. *Cognitive Brain Research*, *17*(2), 484–494. https://doi.org/10.1016/S0926-6410(03)00164-2

Kumar, U. (2016). Neural dichotomy of word concreteness: A view from functional neuroimaging. *Cognitive Processing*, *17*(1), 39–48. https://doi.org/10.1007/s10339-015-0738-1

Kuperberg, G. R., West, W. C., Lakshmanan, B. M., & Goff, D. (2008). Functional Magnetic Resonance Imaging Reveals Neuroanatomical Dissociations During Semantic Integration in Schizophrenia. *Biological Psychiatry*, *64*(5), 407–418. https://doi.org/10.1016/j.biopsych.2008.03.018

Lin, N., Wang, X., Xu, Y., Wang, X., Hua, H., Zhao, Y., & Li, X. (2018). Fine Subdivisions of the Semantic Network Supporting Social and Sensory–Motor Semantic Processing. *Cerebral Cortex*, *28*(8), 2699–2710. https://doi.org/10.1093/cercor/bhx148

Mayer, K. M., Macedonia, M., & Von Kriegstein, K. (2017). Recently learned foreign abstract and concrete nouns are represented in distinct cortical networks similar to the native language: Abstract and Concrete Nouns in the Brain. *Human Brain Mapping*, *38*(9), 4398–4412. https://doi.org/10.1002/hbm.23668

Supplementary Materials

Meersmans, K., Bruffaerts, R., Jamoulle, T., Liuzzi, A. G., De Deyne, S., Storms, G., Dupont, P., & Vandenberghe, R. (2020). Representation of associative and affective semantic similarity of abstract words in the lateral temporal perisylvian language regions. *NeuroImage*, *217*, 116892. https://doi.org/10.1016/j.neuroimage.2020.116892

Meersmans, K., Storms, G., De Deyne, S., Bruffaerts, R., Dupont, P., & Vandenberghe, R. (2022). Orienting to different dimensions of word meaning alters the representation of word meaning in early processing regions. *Cerebral Cortex*, *32*(15), 3302–3317. https://doi.org/10.1093/cercor/bhab416

Mellet, E., Tzourio, N., Denis, M., & Mazoyer, B. (1998). Cortical anatomy of mental imagery of concrete nouns based on their dictionary definition: *NeuroReport*, *9*(5), 803–808. https://doi.org/10.1097/00001756-199803300-00007

Mestres-Missé, A., Münte, T. F., & Rodriguez-Fornells, A. (2009). Functional Neuroanatomy of Contextual Acquisition of Concrete and Abstract Words. *Journal of Cognitive Neuroscience*, *21*(11), 2154–2171. https://doi.org/10.1162/jocn.2008.21171

Nagels, A., Chatterjee, A., Kircher, T., & Straube, B. (2013). The role of semantic abstractness and perceptual category in processing speech accompanied by gestures. *Frontiers in Behavioral Neuroscience*, *7*. https://doi.org/10.3389/fnbeh.2013.00181

Noppeney, U., & Price, C. J. (2002). Retrieval of Visual, Auditory, and Abstract Semantics. *NeuroImage*, *15*(4), 917–926. https://doi.org/10.1006/nimg.2001.1016

Noppeney, U., & Price, C. J. (2004). Retrieval of abstract semantics. *NeuroImage*, *22*(1), 164–170. https://doi.org/10.1016/j.neuroimage.2003.12.010

Olivetti Belardinelli, M., Palmiero, M., Sestieri, C., Nardo, D., Di Matteo, R., Londei, A., D'Ausilio, A., Ferretti, A., Del Gratta, C., & Romani, G. L. (2009). An fMRI investigation on image generation in different sensory modalities: The influence of vividness. *Acta Psychologica*, *132*(2), 190–200. https://doi.org/10.1016/j.actpsy.2009.06.009

Papeo, L., Rumiati, R. I., Cecchetto, C., & Tomasino, B. (2012). On-line Changing of Thinking about Words: The Effect of Cognitive Context on Neural Responses to Verb Reading. *Journal of Cognitive Neuroscience*, *24*(12), 2348–2362. https://doi.org/10.1162/jocn_a_00291

Pauligk, S., Kotz, S. A., & Kanske, P. (2019). Differential Impact of Emotion on Semantic Processing of Abstract and Concrete Words: ERP and fMRI Evidence. *Scientific Reports*, *9*(1), 14439. https://doi.org/10.1038/s41598-019-50755-3

Perani, D., Cappa, S. F., Schnur, T., Tettamanti, M., Collina, S., Rosa, M. M., & Fazio1, F. (1999). The neural correlates of verb and noun processing. *Brain*, *122*(12), 2337–2344. https://doi.org/10.1093/brain/122.12.2337

Pexman, P. M., Hargreaves, I. S., Edwards, J. D., Henry, L. C., & Goodyear, B. G. (2007). Neural Correlates of Concreteness in Semantic Categorization. *Journal of Cognitive Neuroscience*, *19*(8), 1407–1419. https://doi.org/10.1162/jocn.2007.19.8.1407

Rodríguez-Ferreiro, J., Gennari, S. P., Davies, R., & Cuetos, F. (2011). Neural Correlates of Abstract Verb Processing. *Journal of Cognitive Neuroscience*, *23*(1), 106–118. https://doi.org/10.1162/jocn.2010.21414

Roxbury, T., McMahon, K., & Copland, D. A. (2014). An fMRI study of concreteness effects in spoken word recognition. *Behavioral and Brain Functions*, *10*(1), 34. https://doi.org/10.1186/1744-9081-10-34

Rüschemeyer, S.-A., Brass, M., & Friederici, A. D. (2007). Comprehending Prehending: Neural Correlates of Processing Verbs with Motor Stems. *Journal of Cognitive Neuroscience*, *19*(5), 855–865. https://doi.org/10.1162/jocn.2007.19.5.855

Sabsevitz, D. S., Medler, D. A., Seidenberg, M., & Binder, J. R. (2005). Modulation of the semantic system by word imageability. *NeuroImage*, *27*(1), 188–200. https://doi.org/10.1016/j.neuroimage.2005.04.012

Sakreida, K., Scorolli, C., Menz, M. M., Heim, S., Borghi, A. M., & Binkofski, F. (2013). Are abstract action words embodied? An fMRI investigation at the interface between language and motor cognition. *Frontiers in Human Neuroscience*, *7*. https://doi.org/10.3389/fnhum.2013.00125

Shafto, M., Randall, B., Stamatakis, E. A., Wright, P., & Tyler, L. K. (2012). Age-related Neural Reorganization during Spoken Word Recognition: The Interaction of Form and Meaning. *Journal of Cognitive Neuroscience*, *24*(6), 1434–1446. https://doi.org/10.1162/jocn_a_00218

Straube, B., Green, A., Bromberger, B., & Kircher, T. (2011). The differentiation of iconic and metaphoric gestures: Common and unique integration processes. *Human Brain Mapping*, *32*(4), 520–533. https://doi.org/10.1002/hbm.21041

Straube, B., He, Y., Steines, M., Gebhardt, H., Kircher, T., Sammer, G., & Nagels, A. (2013). Supramodal neural processing of abstract information conveyed by speech and gesture. *Frontiers in Behavioral Neuroscience*, *7*. https://doi.org/10.3389/fnbeh.2013.00120

Tettamanti, M., Buccino, G., Saccuman, M. C., Gallese, V., Danna, M., Scifo, P., Fazio, F., Rizzolatti, G., Cappa, S. F., & Perani, D. (2005). Listening to Action-related Sentences Activates Fronto-parietal Motor Circuits. *Journal of Cognitive Neuroscience*, *17*(2), 273–281. https://doi.org/10.1162/0898929053124965

Tettamanti, M., Manenti, R., Della Rosa, P. A., Falini, A., Perani, D., Cappa, S. F., & Moro, A. (2008). Negation in the brain: Modulating action representations. *NeuroImage*, *43*(2), 358–367. https://doi.org/10.1016/j.neuroimage.2008.08.004

Tian, L., Chen, H., Zhao, W., Wu, J., Zhang, Q., De, A., Leppänen, P., Cong, F., & Parviainen, T. (2020). The role of motor system in action-related language comprehension in L1 and L2: An fMRI study. *Brain and Language*, *201*, 104714. https://doi.org/10.1016/j.bandl.2019.104714

Tomasino, B., Fabbro, F., & Brambilla, P. (2014). How do conceptual representations interact with processing demands: An fMRI study on action- and abstract-related words. *Brain Research*, *1591*, 38–52. https://doi.org/10.1016/j.brainres.2014.10.008

Tyler, L. K. (2001). The neural representation of nouns and verbs: PET studies. *Brain*, *124*(8), 1619–1634. https://doi.org/10.1093/brain/124.8.1619

Van Dam, W. O., & Desai, R. H. (2016). The Semantics of Syntax: The Grounding of Transitive and Intransitive Constructions. *Journal of Cognitive Neuroscience*, *28*(5), 693–709. https://doi.org/10.1162/jocn_a_00926

Van Dam, W. O., Rueschemeyer, S.-A., & Bekkering, H. (2010). How specifically are action verbs represented in the neural motor system: An fMRI study. *NeuroImage*, *53*(4), 1318–1325. https://doi.org/10.1016/j.neuroimage.2010.06.071

Van Dam, W. O., Van Dijk, M., Bekkering, H., & Rueschemeyer, S.-A. (2012). Flexibility in embodied lexical-semantic representations. *Human Brain Mapping*, *33*(10), 2322–2333. https://doi.org/10.1002/hbm.21365

Vigliocco, G., Kousta, S.-T., Della Rosa, P. A., Vinson, D. P., Tettamanti, M., Devlin, J. T., & Cappa, S. F. (2014). The Neural Representation of Abstract Words: The Role of Emotion. *Cerebral Cortex*, *24*(7), 1767–1777. https://doi.org/10.1093/cercor/bht025

Wallentin, M., Ostergaard, S., Lund, T., Ostergaard, L., & Roepstorff, A. (2005). Concrete spatial language: See what I mean? *Brain and Language*, *92*(3), 221–233. https://doi.org/10.1016/j.bandl.2004.06.106

Wang, X., Wang, B., & Bi, Y. (2019). Close yet independent: Dissociation of social from valence and abstract semantic dimensions in the left anterior temporal lobe. *Human Brain Mapping*, *40*(16), 4759–4776. https://doi.org/10.1002/hbm.24735

Westbury, C. F., Cribben, I., & Cummine, J. (2016). Imaging Imageability: Behavioral Effects and Neural Correlates of Its Interaction with Affect and Context. *Frontiers in Human Neuroscience*, *10*. https://doi.org/10.3389/fnhum.2016.00346

Whatmough, C., Verret, L., Fung, D., & Chertkow, H. (2004). Common and Contrasting Areas of Activation for Abstract and Concrete Concepts: An H215O PET Study. *Journal of Cognitive Neuroscience*, *16*(7), 1211–1226. https://doi.org/10.1162/0898929041920540

Wise, R. J. S., Howard, D., Mummery, C. J., Fletcher, P., Le, A., & Scott, S. K. (2000). Noun imageability and the temporal lobes. *Neuropsychologia*, *38*(7), 985–994. https://doi.org/10.1016/S0028-3932(99)00152-9

Zhuang, J., Randall, B., Stamatakis, E. A., Marslen-Wilson, W. D., & Tyler, L. K. (2011). The Interaction of Lexical Semantics and Cohort Competition in Spoken Word Recognition: An fMRI Study. *Journal of Cognitive Neuroscience*, *23*(12), 3778–3790. https://doi.org/10.1162/jocn_a_00046