SSR42 is a Novel Regulator of Cytolytic Activity in *Staphylococcus aureus*

Mary-Elizabeth Jobson[1,2], Brooke R. Tomlinson[1], Emilee M. Mustor[1,2], Emily A. Felton[1,2], Andy

Weiss[1], Clayton C. Caswell[3], Lindsey N. Shaw[1,2]

[1]Department of Molecular Biosciences, University of South Florida, Tampa, Florida, United

States

[2]Center for Antimicrobial Resistance, University of South Florida, Tampa, Florida, United States

[3]Department of Biomedical Sciences and Pathobiology, Center for One Health Research, VA-

MD College of Veterinary Medicine, Virginia Tech, Blacksburg, Virginia, United States

Abstract

SSR42 is the longest noncoding RNA in the *S. aureus* cell and the second-most abundant transcript in the stationary phase transcriptome, second only to RNAIII. It is highly conserved across strains and exhibits pronounced stability in stationary phase, however the mechanism behind its regulatory role has yet to be fully elucidated. Herein, we used transcriptomic and proteomic approaches to probe the role of SSR42, revealing that it is a powerful, novel activator of the primary leukocidin LukAB. SSR42 is required for cytotoxicity towards, and escape from within, human neutrophils, and also mediates survival within human blood. We show that SSR42 wields this role via derepression by the peroxide repressor PerR in response to the presence of human neutrophils and governs *lukAB* induction in this niche. Importantly, this regulation is driven by direct RNA-RNA interaction, as we show binding of the 5' UTR of the *lukAB* transcript with the 3' end of SSR42, which ultimately modulates transcript stability as well as translational activity. Finally, we demonstrate that this behavior is absolutely required for full virulence of *S. aureus* in murine models of both pneumonia and sepsis. Collectively, we present SSR42 as a pleiotropic regulatory RNA that acts as a nexus between environmental sensing and the regulation of pathogenesis, responding to environmental stimuli and host immune factors to bolster cytotoxic behavior and facilitate infection in *S. aureus.*

Abstract word count: 225

Importance

S. aureus is a master pathogen due to its formidable collection of virulence factors. These are tightly controlled by a diverse group of regulators that titrate their abundance to adapt to unique infectious niches. The role of regulatory RNAs in stress adaptation and pathogenesis is becoming increasingly more relevant in *S. aureus*. In this study, we provide the most comprehensive global analysis to date of just such a factor, SSR42. Specifically, we uncover

that SSR42 is required for mediating cytotoxicity - one of the pillars of infection - in response to

phagocytosis by human neutrophils. We find that SSR42 is induced by components of the host

immune system and facilitates downstream activation of cytotoxic factors via RNA-RNA

interactions. This illustrates that SSR42 forms a pivotal link between sensing the external

environment and mediating resistance to oxidative stress while promoting virulence, solidifying it

as a major global regulator in *S. aureus.*

Introduction

Over the last few decades, more than 300 regulatory RNAs have been identified in

Staphylococcus aureus strains(1–6). While progress has been made exploring these elements

mechanistically, there is still a dearth of knowledge on the role of regulatory RNAs in complex

regulatory systems, as compared to classical transcription factors. Commonly referred to as

small RNAs (sRNAs), these factors range from *trans*-encoded- to antisense- transcripts,

typically acting in a posttranscriptional manner to allow for rapid adaptation to environmental

cues (6–13). The most well-studied example of an sRNA in *S. aureus* is RNAIII, which is the

effector for the Agr system. It is known to modulate mRNA abundance by relieving inhibitory

secondary structures at the Shine-Dalgarno sequence (e.g. *hla*), or by occluding this site to

block access of the ribosome (e.g. *spa, rot, sbi*) ultimately promoting transcript degradation (14–

17).

Many other virulence-related regulatory RNAs have also been documented in *S. aureus*, such

as the Teg family of transcripts, which include Teg41 that impacts hemolytic ability through

engagement with the PSMs, and Teg58 which monitors arginine dysbiosis and modulates

biofilm production (18–20). The Spr-group of regulators form another family of regulatory RNAs,

and includes SprC, which plays a role in controlling metabolism and immune evasion, SprY,

which works to titrate RNAIII activity through direct 'sponge' interactions, and SprD, which works

in consort with RNAIII to repress *sbi* (21–24) . RsaC is a ~1,100-nt transcript that is expressed

during manganese starvation and enhances the oxidative stress response, leading to increased

survival during confrontation with the host immune system; thus forming a link between metal

homeostasis and pathogenesis (25). From a broader perspective, a recent global study utilizing

CLASH technology to uncover RNA-RNA interactions in host-like conditions identified hundreds

of binding events, occurring not only between regulatory RNAs and mRNAs, but between

regulatory RNAs themselves as well. Some of these interactions linked the metabolic status of

the cell to the expression of membrane-permeabilizing toxins, while furthermore uncovering an

RNA-sponge relationship between RsaE and RsaI (13)

The largest regulatory RNA in *S. aureus* is SSR42, which is 1,233 nt long, contains no functional

ORFs, and bears 98% sequence conservation across strains, with only two single nucleotide

variants found in UAMS-1 (26). It was first characterized by Morrison et al. from a larger group

of Small Stable RNAs (SSRs), where it was shown that the SSR42 transcript exhibits enhanced

stability during stationary phase growth, with a half-life of over 30 minutes, compared to a ~5

minutes in exponential phase. Using microarray analysis, 80 mRNA species were found to be

altered in an SSR42 mutant strain, with key virulence factors such as *hla, lukF, hlgC, aur,* and

sraP found to be differentially expressed. Phenotypically, an SSR42 mutant demonstrated an

abrogated ability to lyse human erythrocytes and impaired virulence in a murine model of skin

abscess formation (26). Others have shown that SSR42 is tightly controlled via the Repressor of

Surface Proteins (Rsp), and in turn may control expression of the two-component system

SaeRS, potentially explaining the ahemolytic nature of SSR42 mutant strains (27). Work by our

group identified that SSR42 is unique to *S. aureus* and does not exist in other Staphylococcal

species (5). We also demonstrate that expression of SSR42 is profoundly increased during the

growth of *S. aureus* in human serum (1).

Despite SSR42 being well linked to *S. aureus* pathogenesis, no direct target or mechanism of

action has been demonstrated for this molecule. Herein, we use a combination of large-scale

omics studies coupled with genetic and biochemical analyses to demonstrate that SSR42 is a

novel regulator of *lukAB* expression and cytolytic activity through direct RNA-RNA interaction.

We show that this regulation is controlled by the peroxide sensor PerR and is stimulated in

response to human neutrophils, mediating escape from the intracellular niche. Finally, the

impact of this interaction is illustrated by impaired virulence in murine models of both sepsis and

pneumonia.

Methods

(Additional methods can be found in the supplemental methods file.)

Bacterial strains and growth conditions: All bacterial strains and plasmids used for this study

are listed in **Table 1**. Overnight cultures were routinely grown at 37°C with shaking at 250 rpm in

5mL of tryptic soy broth (TSB) for *S. aureus* or lysogeny broth (LB) for *E. coli.* When required,

media was supplemented with the following antibiotics: for *E. coli*: 100μg/mL ampicillin; for *S.*

aureus: 10μg/mL chloramphenicol, 5μg/mL erythromycin, 25μg/mL lincomycin, and 5μg/mL

tetracycline. To obtain synchronous growth of *S. aureus,* overnight cultures were diluted 1:100

into fresh TSB and grown for 3h before being standardized to a uniform OD_{600} of 0.05 in fresh

TSB. Unless otherwise described, cell pellets were harvested via centrifugation at 4000 x *g* for

10 minutes.

Toxicity towards neutrophils: Cytotoxicity assays were performed as previously described

(28), with some variations. The immortalized HL-60 cell line was used as a model of human

neutrophil engagement. Cells were grown to confluence in RPMI supplemented with 10% FBS

and 100U Pen/Strep in 37°C and 5% CO_2. Upon reaching sufficient cell density, cells were

differentiated into neutrophil-like cells by the addition of 1.25% DMSO for 4 days; which was

confirmed via observation of cell morphology. For use, $1x10^5$ cells were seeded in 100µl RPMI

with 10% FBS into a 96-well plate. Bacterial cultures were grown in biological triplicate overnight

and standardized to 0.05 before being grown for 15h. Following incubation, cultures were

pelleted via centrifugation and supernatant was diluted two-fold in TSB. Seeded neutrophils

were intoxicated in technical duplicate with 5µl of diluted bacterial supernatant (5% total volume)

followed by incubation at 37°C, 5% CO_2 for 1h. Cell viability was measured using the CellTiter96

Aqueous One Solution Cell proliferation reagent (Promega). Following 1h incubation, 20µl of the

CellTiter reagent was added to each well, and the plate was returned to the incubator for 2h.

Color development was assessed via OD_{490} using a Biotek Cytation 5 plate reader. Data is

reported as percent death of supernatant-treated cells with respect to neutrophils treated with

5µl TSB only.

Intracellular survival within neutrophils: To assess intracellular survival within human

neutrophils, HL-60 cells were grown and differentiated as described above, and $2x10^6$ cells

were seeded into 500µl in a 24-well plate. Bacterial cultures were also grown and processed as

above, with the following modifications. After centrifugation, pelleted cells were washed 2X with

PBS, following which all strains were standardized to 10x the desired final MOI of 30

(OD_{600}=0.135 for *S. aureus*), thus all strains were standardized to an OD_{600}=13.5 in PBS. From

these standardized cultures, 5µl was added to each well of neutrophils and incubated for 1h at

37°C and 5% CO_2. Following incubation, cells were removed from wells and pelleted at 2.5 x *g*

for 1 minute before being resuspended in pre-warmed PBS. This PBS wash was repeated, and

the cells were finally resuspended in pre-warmed RPMI supplemented with 30µg/mL gentamicin

and returned to incubate for 24h. At this point in time the cells were removed and washed twice

with pre-warmed PBS once again, however the final resuspension was in 0.5% Triton X-100 to

lyse neutrophils. Lysates were serially diluted and plated to enumerate CFU of internalized

bacteria.

Blood survival assays: Blood survival assays were performed as described previously by our

group (29). Briefly, bacterial cultures were grown in biological triplicate overnight before being

diluted 1:100 and grown for a further 3h. These synchronized cultures were then used to

inoculate 1mL of human, gender-pooled whole blood (BioIVT) at a final OD_{600} of 0.05. Samples

were taken immediately after inoculation (T=0) as well as after 4h of rotation at 37°C. Samples

were serially diluted and plated onto TSA to calculate CFU/mL. Data is represented as percent

survival in comparison to T=0.

Luminescent reporter assays with neutrophil exposure: HL60 cells were differentiated as

described above before being seeded into black-walled, clear bottomed 96-well plates at a

concentration of $2x10^5$ in 100µl of phenol red-free RPMI. Bacterial strains were grown overnight

as previously described and washed 2x with PBS and standardized to a final OD_{600} of 1.0.

Following this, standardized bacterial cells were added to seeded HL-60 cells to reach a final

OD_{600} of 0.1. Luminescence and OD_{600} T=0 readings were taken immediately after inoculation

(T=0) on a Biotek Cytation 5 plate reader, with additional readings taken hourly, between which

plates were incubated at 37°C and $5\%CO_2$. Luminescence was normalized to OD_{600} for each

well.

Measuring translational control by SSR42 via GFP reporters: SSR42 mutant strains

containing both the *lukAB* mRNA-GFP fusion plasmid (pCN33) alongside the SSR42 expression

or empty vector plasmid (pICS) were grown overnight in biological triplicate. Cultures were

synchronized for 3h and standardized to an OD_{600} of 0.05 in a black-walled, clear bottom 96-well

plate. Strains were then monitored for OD_{600} and fluorescence (excitation:485/20,

emission:528/20, Gain:75) measured every 15min for 18h. Fluorescence levels of the mRNA

reporter in the presence of SSR42 *in trans* were compared to fluorescence levels of the mRNA

reporter in the presence of an empty vector. Strains were normalized by growth via OD_{600}.

Electrophoretic mobility shift assays (EMSAs):　All RNA was generated via *in vitro*

transcription using a T7 RNA polymerase promoter and the MAXIscript T7 kit (ThermoFisher).

SSR42 was amplified using OL7499/OL7500 while fragments of the *lukA* UTR and *splE* UTR

were amplified using primers 7501/7502 and 7523/7524, respectively. These T7-controlled

fragments were cloned into pGEM-T Easy and liberated from the plasmid using restriction

enzymes NcoI and PstI (Thermo Fisher Scientific). *In vitro* transcription was performed using the

MAXIscript T7 kit, ^{32}P-labeled UTP for the mRNA, and unlabeled UTP for SSR42. TurboDNase

was added to degrade any remaining DNA. Reactions were cleaned using a Nucleotide Clean-

up Kit (Qiagen). RNA-RNA binding reactions were performed with labeled mRNA and SSR42 at

varying concentrations, which were heated to 90°C for 2 minutes and then incubated at RT for

30 min. Native acrylamide gels were equilibrated on ice for 30 minutes in 0.5x TBE before

loading binding reactions, which were mixed with RNA loading dye and then run at 100V until

the dye front has moved ¾ of the way through the gel. Gels were dried and affixed to filter paper

using a gel dryer before being visualized via autoradiography.

Murine models: All animal studies in this work were performed in accordance with and

approved by the Institutional Animal Care and Use Committee of the University of South Florida.

To prepare inocula for a model of bacterial sepsis, bacteria was washed 2x with PBS before

being adjusted to $5x10^8$ CFU/mL in PBS. Female, six-week-old CD-1 mice (Charles River

Laboratories) were inoculated with 100µl of bacteria, resulting in a final dose of $5x10^7$. Infection

was allowed to progress for 7 days, or until mice reach a premoribund state, at which point they

were euthanized. After the infection period, all mice were euthanized and the brain, heart, liver,

spleen, lungs, and kidneys were harvested. For murine models of pneumonia, inocula were

prepared in a similar manner, however the bacterial density was instead adjusted to $1x10^9$.

Female and male 6-week old C57BL/6 mice were intranasally inoculated with 100µl of bacteria,

resulting in a final dose of $1x10^8$. Infections were allowed to proceed for 24h, after which mice

were euthanized and lungs harvested. Organs for both models were homogenized in 2mL PBS

and serially diluted and plated for CFU/mL. All infection experiments were performed with 8

mice per strain and a Mann-Whitney test was used to measure statistical significance.

Data Availability: RNAseq data from this study is available under the GEO accession

#GSE237701. Proteomics data is available via proteomeXchange under the ID PXD043724.

Results

SSR42 is global regulator of virulence factor expression: Although previous studies exist

exploring SSR42, the only large-scale omics examination of this factor is a microarray

performed over a decade ago (26). Thus, to bring new light to the role of SSR42, we herein

performed RNA-sequencing on an SSR42 mutant in the USA300 background. Strikingly, in the

wild-type strain, SSR42 was the second most abundant transcript in the cell, second only to

RNAIII **(Supplemental figure 1).** Moreover, 20% of all RNA in stationary phases cells (absent

rRNA) belongs to SSR42, and 45% to SSR42 and RNAIII combined. Beyond this, we found a

total of 220 genes were significantly differentially expressed between wild-type and mutant (**Fig.**

1A), greatly surpassing the 80 genes previously shown to be controlled by SSR42 via

microarray. These included those involved in metabolism, transport, signal transduction and, by

far the largest functional category (33 such factors), human disease. Regarding this latter group,

SSR42 primarily serves as an activator, with 83% of transcripts decreased in abundance in the

mutant strain **(Fig. 1B,C).**

Key virulence factors regulated by SSR42 included secreted toxins, such as alpha hemolysin

(*hla,* -3.44), gamma hemolysin (*hlgA* -5.39, *hlgB* -4.77), the Panton-Valentine leukocidin (*lukS* -

3.62, *lukF* -3.86), and the *lukAB* leukocidin (-5.49, -4.29). Secreted proteases were also

impacted, with all serine protease (*spl*) operon genes downregulated in the mutant (~-3-fold),

whilst aureolysin (*aur*) and the V8 protease (*sspA)* were both increased in abundance (+4.5-fold

and +2.81-fold, respectively). Moreover, SSR42 appears to activate immune evasion factors,

including igG-binding protein *sbi* (-5.36), extracellular fibrinogen-binding protein *efb* (-2.85), and

staphylokinase *sak* (-2.48). In the same vein, we identified a profound change in the secreted

extracellular matrix and plasma binding protein *emp* (-34.6). Other virulence factors repressed

by SSR42 include the serine-rich adhesin for platelets *sraP* (+4.12) and the Immunodominant

Antigen B *isaB* (+2.92).

We also uncovered alterations in known transcriptional regulators, the most striking being the

SarA-family proteins *sarZ* and *sarR*, which were downregulated 9.6-fold and 62.3-fold,

respectively, in the mutant. Interestingly, the Repressor of Surface Proteins (*rsp*), which is

divergently transcribed from SSR42 and has been shown to be a primary activator of its

expression (27), was found to be upregulated 4.31-fold upon SSR42 deletion. Finally, we

identified changes in expression of the metalloregulators *fur* (+3.14) and *mntR* (-2.13), as well

as the alternative sigma factor *sigS* (-2.75), in the mutant strain. A random subset of these fold

changes were explored and confirmed by qPCR analysis (**Supplemental Figure 2**).

We next used proteomic analysis to validate these changes, revealing strong correlation to our

RNAseq data **(Fig 1 A,B,C, and Table S1).** Notably, while leukocidins LukSF and LukAB exhibit

decreased protein abundance in the mutant that matches RNAseq trends, LukAB in particular

exhibits a profound decrease in fold-changes at the protein level (RNAseq – *lukA* = -5.51, *lukB* =

-4.3; Proteomics - LukA = -9.87, LukB = -7.83).

SSR42 is a novel regulator of cytolytic activity: Given that LukAB had such profound

changes in our omic studies we next set out to explore if these had an impact of leukocidin

activity. As LukAB is the primary effector for the lysis of human neutrophils (30), human

promyeloblast (HL-60) cells were differentiated into neutrophils before being treated with

bacterial supernatant from the wild type, mutant, and complemented strains. Strikingly, after a

1h incubation, supernatant from the SSR42 mutant elicited a 5-fold reduction in killing compared

to the wild-type and complement **(Fig. 2A).** To see how this translated into bacterial viability, we

next explored how deletion of SSR42 impacted *S. aureus* survival during exposure to whole

human blood. Here we noted the mutant exhibited a marked 7-fold increase in survival

compared to the wild type and complementing strains **(Fig. 2B).** Of note, *lukAB* mutants of *S.*

aureus have been shown to have increased intracellular survival within neutrophils (a key

component of whole blood), due to an inability to use the toxin to escape the phagosome (31).

To determine if this was the case for the SSR42 mutant, we next measured intracellular survival

of our strains within HL-60 cells. Here we determined that, after 24h incubation, the mutant

displayed a dramatic increase in survival in comparison to the wild type and complement within

this niche **(Fig. 2C).** As such, it appears that SSR42 is a major new regulator of LukAB activity

in *S. aureus*.

SSR42 is upregulated upon exposure to neutrophils via PerR derepression: It has

previously been shown that *lukAB* expression is induced following exposure of *S. aureus* to

neutrophils, and that this is mediated, indirectly, via the peroxide regulator PerR, which is known

to derepress its targets in the presence of reactive oxygen species (32–34). Given that SSR42

plays an intimate role in intracellular survival and cytotoxicity, we next set out to determine if

SSR42 was the missing link in this chain of regulation. To this end, we created a luciferase

transcriptional reporter for the SSR42 promoter and exposed it to differentiated neutrophil-like

HL-60 cells. In so doing, we determined that P_{SSR42} exhibited a >3-fold increase in activity after

just 1h in the presence of neutrophils **(Fig 3A)**. Next, to investigate if this regulation was

governed by PerR, we re-assessed SSR42 promoter activity in a *perR* mutant. This time we

determined that P_{SSR42} expression is significantly higher in the mutant strain when exposed to

neutrophils as compared to wild-type **(Fig 3B)**, which is consistent with PerR derepression.

SSR42 controls *lukAB* expression in response to neutrophils: We next set out to

investigate whether SSR42 might mediate the impact of PerR on *lukAB* expression during

engagement with neutrophils. As such, we created a P$_{lukAB}$-*lux* fusion and assessed its response

to neutrophils in the context of SSR42 and PerR. In so doing, we noted that upon addition of

neutrophils, there was a significant increase in *lukAB* expression in the wild-type strain that

mirrored that of P$_{SSR42}$ **(Fig. 4A).** However, this response is strikingly abrogated in an SSR42

mutant, where the luciferase signal is 5-fold lower in comparison to the wild-type after 2h of

neutrophil exposure **(Fig. 4B).** To uncouple the effect of SSR42 from PerR, we next assessed

lukAB expression in a *perR* mutant as well as an SSR42/*perR* double mutant. This revealed

that, once again in a similar manner to SSR42, *lukAB* expression was higher in a *perR* mutant,

however, *lukAB* expression in the SSR42/*perR* mutant mirrors that of an SSR42 mutant **(Fig. 4**

A-C). This suggests that the response of *lukAB* to the presence of neutrophils is likely a result of

direct regulation by SSR42, which is in turn controlled by PerR. This is further proven upon

overexpression of PerR in an SSR42 mutant, where there is no significant difference in P$_{lukAB}$

activity **(Fig S3.)**.

SSR42 regulates *lukAB* post-transcriptionally: To determine if the changes in *lukAB*

expression are mediated by SSR42 binding to, and affecting stability of, the *lukAB* mRNA, we

fused a fragment of the *lukA* 5' UTR and a portion of its coding sequence in frame to GFP and

expressed it under the control of a constitutive promoter in *S. aureus.* We then quantified GFP

fluorescence in an SSR42 mutant vs. wild-type, revealing a significant decrease in the stability

of the translated LukA-GFP product in the mutant strain **(Fig 5A).** Next, we used a two-plasmid

system in which we expressed SSR42 under its native promoter alongside the LukA-GFP

reporter in the SSR42 mutant strain. Here we observed a striking increase in LukA-GFP

translation in the presence of SSR42 when compared to empty vector controls **(Fig. 5B).** Taken

together, this suggests that SSR42 putatively binds post-transcriptionally to the *lukAB* mRNA to

regulate its stability and thus translation.

The 3' end of SSR42 is required for interaction with *lukAB*: To explore interaction between

SSR42 and the *lukAB* mRNA further, we first used IntaRNA to predict possible base pairing

interactions between the two RNA molecules (35). This predicted interaction with the 3' end of

SSR42, ~380bp from the end of the transcript, and the 5' UTR of *lukAB*, ending 7bp upstream of

the ribosome binding site (RBS) **(Fig. 6A)**. To investigate the relevance of this interaction *in*

vivo, we generated a series of truncated SSR42 variants and introduced these into the SSR42

mutant, before reassessing cytotoxicity. In so doing we noted that a 648-nt SSR42 fragment

(from the +1 onwards) was unable to restore wild-type levels of cytotoxicity, while a longer 793-

nt transcript partially restored toxicity. **(Fig 6B).** However, upon complementation with a 940-nt

SSR42 transcript that contains the region identified by IntaRNA, wild-type cytotoxicity was

observed. This suggests that the far 3' end of SSR42 is primarily responsible for mediating

interaction with *lukAB*, and thus governing cytotoxicity.

SSR42 directly binds to the *lukAB* mRNA and enhances its stability: To explore further the

direct interaction between SSR42 and the *lukAB* mRNA, electrophoretic mobility shift assays

(EMSAs) were performed. A portion of the *lukA* UTR that included the region identified in

IntaRNA was radiolabeled using *in vitro* transcription and incubated with increasing

concentrations of *in vitro* transcribed SSR42. When these samples were analyzed, we observed

a concentration-dependent interaction between the two RNA molecules (**Fig. 7A).** By

comparison, no binding was seen with radiolabeled *splE* mRNA under identical conditions,

supporting the specificity of the SSR42-*lukAB* interaction. We next set out to investigate whether

the observed binding had any impact on transcript stability. To that end, transcriptional arrest

studies were performed on the wild type, mutant, and complemented strains grown for 15h.

Samples were taken at t=0 before the addition of Rifampin, and then again at 5, 10, 30, and 45

minutes posttreatment. RNA abundance was then quantified via qPCR, revealing that the *lukA*

transcript degraded far more quickly, and to a greater overall degree, in the mutant strain when

compared to the wild type and complement **(Fig 7B).** When the mRNA half-life was calculated

using a one phase decay curve, we found that the *lukAB* mRNA $t_{1/2}$ was 2.9 minutes in the

SSR42 mutant, but 4.5 minutes in the wild type. This suggests SSR42 is required for

maintaining stability of the *lukAB* mRNA and facilitating translation of its protein products.

SSR42 is required to cause invasive *S. aureus* infections: Previous work has shown that

SSR42 is important for localized infections using a murine model of abscess formation (26). To

assess the role SSR42 plays in invasive disease we assessed the pathogenic capacity of the

mutant in a murine model of septic infection. Mice were infected with the SSR42 mutant, or wild-

type strain and the bacterial burden was determined across multiple organ systems after 7 days.

Here we found that the SSR42 mutant exhibited an abrogated ability to disseminate to the liver,

heart, and spleen **(Fig. 8A).** Specifically, in the heart, ΔSSR42 exhibits a 4-log decrease in

bacterial burden compared to WT, with 1-2-log decreases found in the liver and spleen. To

explore this virulence defect more fully, we next used an acute model of murine pneumonia,

intranasally inoculating mice with the wild type and SSR42 mutant strains. After 24h of infection

we assessed bacterial recovery from the lungs and once again found a major defect in the

mutant's ability to survive, with the ΔSSR42 strain exhibiting a 3-log decrease in CFU/mL in

comparison to the wild type **(Fig. 8B)**. These findings, alongside those from others demonstrate

that SSR42 is required for *S. aureus* infection in three different animal models, representing

both localized and invasive forms of disease.

Discussion

Recent years have seen an abundance of new discoveries regarding the complex nature of

RNA-based regulation, revealing that many of these elements exhibit pleiotropic behavior and

nuanced roles in pathogenesis. Herein, we explore just such a factor, broadening the known

regulome of the long noncoding RNA SSR42 from *S. aureus*. Using both transcriptomic and

proteomic approaches, with high sensitivity technology, we expand the SSR42 network of

control almost three-fold from previous studies.

Unraveling the SSR42 regulome reveals a far more complex network than was previously

ascertained. Of note, we identified no significant changes in any component relating to the

SaeRS TCS, which previous studies suggested was responsible for SSR42-related regulation of

at least some virulence factors (27). Instead, we note the strong downregulation of two SarA-

homologs, *sarZ* and *sarR*, at -9.6-fold and -62.6-fold, respectively, in our SSR42 mutant. Both

transcription factors have been characterized to play a role in promoting virulence factor

expression in various *S. aureus* strains (36–39). Specifically, SarZ upregulates the expression of

aur, sspAB, and *hla* while SarR activates *sspAB* and the *spl* proteases. Upon assessment of our

own transcriptomics, the regulation of the proteases *aur* (via SarZ) and *sspAB* (via SarR) are

consistent with the trends seen in our data, however the changes in *hla* (via SarZ) and *spl* (via

SarR) expression are not. It is intriguing that these transcripts are so strikingly decreased in

abundance, suggesting that SSR42 plays a strong role in controlling their expression or stability.

Indeed, bioinformatic searches for RNA-RNA binding do predict very strong interactions

between SSR42 and the mRNAs of both transcription factors **(Fig. S4)**. Thus, it is possible that

changes in *sarR* and *sarZ* expression may be causative for some, but not all, of the alterations

in virulence factor transcription seen in the absence of SSR42.

It is known that the Repressor of Surface Proteins, Rsp, is absolutely required for SSR42

expression (27). Both are transcribed divergently from each other, sharing a putative promoter

region, and yet in an SSR42 mutant *rsp* transcript levels are increased ~4-fold. This suggests

some element of negative feedback by which SSR42 effectively represses its own expression

during stationary phase by repressing *rsp*. It is well-established that SSR42 exhibits profound

transcript stability during stationary phase (>30min half-life) (26), thus this auto-repression via

rsp could be a method to titrate SSR42 abundance in the cell when it is not necessary to

produce new transcript. More importantly, however, is the fact that *rsp* has previously been

shown to be induced in the presence of hydrogen peroxide and is required for full cytotoxicity in

an Agr-independent manner (40). We posit that these phenotypes are at least partially SSR42-

mediated. To explore this, we assessed expression from the *rsp* promoter in the presence of

neutrophils and found that it too is induced by their presence **(Fig S5)**. Importantly, however, P$_{rsp}$

is not derepressed by PerR in this context, suggesting that the PerR-mediated control of *lukAB*

by SSR42 is independent of Rsp. This demonstrates that, depending on the environment, Rsp

is necessary, but not sufficient, for SSR42 expression.

When assessing impact on LukAB production, we show that SSR42 binding to the *lukAB* mRNA

affects its stability, producing a concomitant impact on protein production. The latter effect, we

posit, is reflective of this interaction altering the secondary structure of the *lukAB* mRNA,

perhaps making it more favorable for the ribosome to access and initiate translation. Indeed, a

very similar mechanism is seen for the well-studied interaction between RNAIII and the *hla*

mRNA, where there is binding directly upstream of the RBS for this same purpose(17).

Interestingly, in addition to the impact of SSR42 on *lukAB* mRNA stability and/or translational,

we also observe alterations in P$_{lukAB}$ promoter activity. This, at first glance, is contradictory to

typical sRNA behavior, which usually only manifest post-transcriptionally. However, the P$_{lukAB}$

fragment used in our luciferase studies also contains the 5'UTR region of the transcript,

including the SSR42 binding site. Thus, as the absence of SSR42 is proven to destabilize the

lukAB mRNA, the promoter fusion transcript will inevitably be destabilized in our luciferase

assays as well.

That said, it is a cornerstone of regulatory RNA behavior to exert multiple methods of regulation

on a single target. This is exemplified by RNAIII, which binds the hla mRNA and opens its

secondary structure near the SD sequence to allow translation to initiate, while also binding and

inhibiting translation of *rot* to prevent transcriptional inhibition of *hla* (16, 17). Exerting diverse

modes of control over mRNA targets can ensure a robust regulatory response, which we readily

see in the case of SSR42 and LukAB. At the transcript level, we identified a ~5-fold decrease in

the lukAB mRNA, which likely reflects mRNA instability in the absence of SSR42; yet we see a

nearly 10-fold decrease in abundance of both LukAB components in secretomes, indicating how

lost translation can have an amplified effect. However, LukAB is somewhat unique in that it is

not only secreted but also stored within the cell envelope, where it can exert proximity-based

toxicity when in confrontation with host immune cells (41, 42). In our own proteomic analysis,

the cell-associated fraction has a striking -27-fold decrease in protein abundance for LukA and a

-6.6-fold decrease for LukB in an SSR42 mutant. This suggests that SSR42 influences not only

the canonically secreted variant of this toxin, but also the sorted and stored, yet active, form of

LukAB. Thus, in an SSR42 mutant, this weapon is markedly decreased in availability, and thus it

follows that the SSR42 mutant is unable to escape the phagocytic niche.

Recent work by Savin et al. identified the peroxide sensor PerR as a regulator of LukAB activity

that responds to the presence of hPMNs and leads to enhanced leukocidin expression (34).

Their study suggested no direct regulation between PerR and lukAB as there are no PerR

binding boxes detected in the leukocidin promoter. In our study, we observed that the SSR42

promoter is also induced in the presence of human neutrophils and, importantly, is required for

the lukAB promoter to respond under these conditions. Investigating P_{SSR42} architecture

revealed that there are 3 putative PerR binding boxes (sequence identified by Horsburgh et. al

(33)) upstream and/or within the SSR42 transcript **(Fig S6)**. It has previously been shown that

known PerR targets such as katA, ahpC, and ftnA can possess multiple binding boxes within

their promoters and are de-repressed during oxidative stress (32–34). When we mutated each

of these binding sequences individually, we identified no changes in promoter induction (data

not shown), however, we were unable to combinatorially mutate more than one box within the

promoter sequence at the same time. It is possible that profound derepression in the absence of

these PerR binding sites results in toxic over production of SSR42.

Importantly, the SSR42 promoter has previously been shown to be induced in response to the

presence of hydrogen peroxide (40). Thus, we propose that SSR42 is the missing link between

PerR-mediated ROS sensing and subsequent activation of LukAB activity to mediate survival

and escape from phagocytic cells. Indeed, there is precedence for regulatory RNA involvement

in oxidative stress resistance across bacterial species. First, in *E. coli,* the PerR homolog OxyR

derepresses the pleiotropic sRNA regulator OxyS in response to hydrogen peroxide (43–45). In

Pseudomonas stutzeri, the sRNA NfiS regulates catalase production and interacts with OxyR in

response to H_2O_2-induced conditions (46). Finally, in *S. aureus,* RsaC is an sRNA that inhibits

the translation of *sodA and* promotes oxidative stress resistance through SodM activity (25).

Evidence for PerR-mediated regulation of SSR42 also arises from a recent study by our group.

Here, upon assessing transcriptomic changes in response to treatment with Calprotectin (i.e.

metal starved), SSR42 was found to be downregulated 2-fold. When samples were treated with

a mutated form of Calprotectin that only sequesters Zn and Cu, SSR42 was upregulated >3-fold

(47). In *S. aureus,* PerR requires Fe and Mn for oxidation in response to endogenous H_2O_2 (32);

thus, it follows that derepression of SSR42 would be detected only in conditions including these

metals.

Metal homeostasis and the PerR-mediated oxidative stress response are intimately linked

through a network of metalloregulators: Fur, Zur, and MntR (33, 48–51). Interestingly, the *fur*

transcript is upregulated 3.14-fold in our SSR42 mutant while *mntR* expression is decreased -2-

fold. It is important to note that in addition to regulating SSR42, PerR plays a role in regulating

Fur and MntR expression, while all three players act in a coordinated manner to induce

overlapping regulons contributing to oxidative stress resistance, including factors such as

catalase (KatA*)*, thioredoxin reductase (TrxB), and alkylhydroperoxide reductase (AhpCF) (48,

51). While these elements were not altered in the SSR42 mutant transcriptome, we did uncover

three alternative antioxidant-like factors that were decreased in expression in the SSR42

mutant. These include a nitroreductase linked to iron acquisition (*ntrA,* -3.41-fold), a

nitroreductase with glutathione antioxidant behavior (SAUSA300_0790, -2.26-fold), and a

thioredoxin-like protein (SAUSA300_0795, -2.27-fold) (52–54). Thus, we posit that SSR42

weaves a common thread connecting these metallo-oxidative regulators and subsequently

impacts virulence factor abundance in response to ROS-generating niches. Key examples of

this behavior in *S. aureus* have been shown in the case of IsrR and RsaC, two sRNAs that are

expressed in low iron or manganese environments, respectively, and function to enhance the

oxidative stress response and pathogenesis upon activation (25, 55–57).

Another regulator, the alternative sigma factor *sigS*, was downregulated -2.75-fold in the SSR42

mutant. σ^S has previously been shown by our group to play a role in stress adaptation, immune

evasion, and pathogenesis, and has recently been shown to have a regulon that includes

various noncoding RNAs (58, 59). Although SSR42 was not found to be differentially expressed

in a *sigS* mutant, our findings suggest that SSR42 is instead responsible for full σ^S abundance.

A similar phenomenon is observed with the *E. coli* sRNA OxyS, which senses hydrogen

peroxide and in turn activates expression of the alternative sigma factor, RpoS (44). Moreover,

we found that *prli42*, which encodes a membrane protein that is required for the formation of the

σ^B controlling stressosome in *Listeria monocytogenes*, but has not yet received characterization

in *S. aureus,* was decreased in expression ~ 3-fold in the SSR42 mutant (60–62). In addition to

our data showing H_2O_2/neutrophil-mediated SSR42 induction and subsequent stimulation of

LukAB activity, these findings demonstrate that SSR42 may also play a role in activating

components of other stress-responsive systems. Taken together, these findings expand the role

of SSR42 beyond a direct virulence-related pathway and illustrates a much more complex role

for this sRNA as it pertains to sensing oxidative stress, nutritional immunity, and other

exogenous signals. A proposed mechanism of SSR42's involvement in the regulation of these

processes is summarized in **Fig 9.**

To our knowledge, this is the first direct RNA-based regulation of *lukAB* characterized to date in

S. aureus. Not only do we uncover a novel regulator or leukocidin activity, but we begin to

unravel the labyrinthine network that controls the response to changing environments,

particularly within the host. Indeed, past work on SSR42 demonstrated its role in murine

abscess formation, while our own study broadens this to include complex and disseminated

infections including sepsis and pneumonia. Thus, we present SSR42 as a pleiotropic regulatory

RNA that responds to pathophysiologically relevant conditions to bolster virulence factor

abundance and facilitate pathogenesis in *S. aureus.*

Acknowledgements: This study was supported by grants AI124458 and AI157506 (both to

L.N.S.) from the National Institute of Allergy and Infectious Diseases.

References

1.   Carroll RK, Weiss A, Broach WH, Wiemels RE, Mogen AB, Rice KC, Shaw LN. 2016. Genome-wide
annotation, identification, and global transcriptomic analysis of regulatory or small RNA gene
expression in *Staphylococcus aureus*. mBio 7.

2.   Liu W, Rochat T, Toffano-Nioche C, Le Lam TN, Bouloc P, Morvan C. 2018. Assessment of bona fide
sRNAs in *Staphylococcus aureus*. Front Microbiol 9:1–13.

3.   Pichon C, Felden B. 2005. Small RNA genes expressed from *Staphylococcus aureus* genomic and
pathogenicity islands with specific expression among pathogenic strains. Proc Natl Acad Sci U S A
102:14249–14254.

4.   Sorensen HM, Keogh RA, Wittekind MA, Caillet AR, Wiemels RE, Laner EA, Carroll RK.
2020.Reading between the Lines: Utilizing RNA-Seq Data for Global Analysis of sRNAs in
Staphylococcus aureus. mSphere5:10.1128/msphere.00439-20.

5.   Broach WH, Weiss A, Shaw LN. 2016. Transcriptomic analysis of staphylococcal sRNAs: Insights
into species-specific adaption and the evolution of pathogenesis. Microb Genom 2.

6.   Bronsard J, Pascreau G, Sassi M, Mauro T, Augagneur Y, Felden B. 2017. SRNA and cis-Antisense
sRNA identification in *Staphylococcus aureus* highlights an unusual sRNA gene cluster with one
encoding a secreted peptide. Sci Rep 7:1–17.

7. Menard G, Silard C, Suriray M, Rouillon A, Augagneur Y. 2022. Thirty Years of sRNA-Mediated
Regulation in *Staphylococcus aureus*: From Initial Discoveries to In Vivo Biological
ImplicationsInternational Journal of Molecular Sciences.

8. Barrientos L, Mercier N, Lalaouna D, Caldelari I. 2021. Assembling the Current Pieces: The Puzzle
of RNA-Mediated Regulation in *Staphylococcus aureus*. Front Microbiol 12:1–7.

9. Bastock RA, Marino EC, Wiemels RE, Holzschu DL, Keogh RA, Zapf RL, Murphy ER, Carroll RK.
2021. *Staphylococcus aureus* Responds to Physiologically Relevant Temperature Changes by
Altering Its Global Transcript and Protein Profile. mSphere 6.

10. Bordeau V, Cady A, Revest M, Rostan O, Sassi M, Tattevin P, Donnio PY, Felden B. 2016.
Staphylococcus aureus regulatory RNAs as potential biomarkers for bloodstream infections.
Emerg Infect Dis 22:1570–1578.

11. Kim S, Reyes D, Beaume M, Francois P, Cheung A. 2014. Contribution of teg49 Small RNA in the 5'
Upstream Transcriptional Region of sarA to Virulence in *Staphylococcus aureus*. Infect Immun
82:4369–4379.

12. Le Huyen KB, Gonzalez CD, Pascreau G, Bordeau V, Cattoir V, Liu W, Bouloc P, Felden B,
Chabelskaya S. 2021. A small regulatory RNA alters *Staphylococcus aureus* virulence by titrating
RNAIII activity. Nucleic Acids Res 49:10644–10656.

13. McKellar SW, Ivanova I, Arede P, Zapf RL, Mercier N, Chu LC, Mediati DG, Pickering AC, Briaud P,
Foster RG, Kudla G, Ross Fitzgerald J, Caldelari I, Carroll RK, Tree JJ, Granneman S. 2022. RNase III
CLASH in MRSA uncovers sRNA regulatory networks coupling metabolism to toxin expression. Nat
Commun 13.

14. Gupta RK, Luong TT, Lee CY. 2015. RNAIII of the *Staphylococcus aureus* agr system activates global
regulator MgrA by stabilizing mRNA. Proc Natl Acad Sci U S A 112:14036–14041.

15. Huntzinger E, Boisset S, Saveanu C, Benito Y, Geissmann T, Namane A, Lina G, Etienne J,
Ehresmann B, Ehresmann C, Jacquier A, Vandenesch F, Romby P. 2005. *Staphylococcus aureus*
RNAIII and the endoribonuclease III coordinately regulate spa gene expression. EMBO Journal
24:824–835.

16. Boisset S, Geissmann T, Huntzinger E, Fechter P, Bendridi N, Possedko M, Chevalier C, Helfer AC,
Benito Y, Jacquier A, Gaspin C, Vandenesch F, Romby P. 2007. *Staphylococcus aureus* RNAIII
coordinately represses the synthesis of virulence factors and the transcription regulator Rot by an
antisense mechanism. Genes Dev 21:1353–1366.

17. Morfeldt E, Taylor D, Von Gabain A, Arvidson S. 1995. Activation of alpha-toxin translation in
Staphylococcus aureus by the trans-encoded antisense RNA, RNAIII. EMBO Journal 14:4569–4577.

18. Briaud P, Zapf RL, Mayher AD, McReynolds AKG, Frey A, Sudnick EG, Wiemels RE, Keogh RA, Shaw
LN, Bose JL, Carroll RK. 2022. The Small RNA Teg41 Is a Pleiotropic Regulator of Virulence in
Staphylococcus aureus. Infect Immun 19.

19. Zapf RL, Wiemels RE, Keogh RA, Holzschu DL, Howell KM, Trzeciak E, Caillet AR, King KA, Selhorst
SA, Naldrett MJ, Bose JL, Carroll RK. 2019. The small RNA Teg41 regulates expression of the alpha
phenol-soluble modulins and is required for virulence in *Staphylococcus aureus*. mBio 10:1–19.

20. Manna AC, Leo S, Girel S, González-Ruiz V, Rudaz S, Francois P, Cheung AL. 2022. Teg58, a small
regulatory RNA, is involved in regulating arginine biosynthesis and biofilm formation in
Staphylococcus aureus. Sci Rep 12:1–18.

21. Zhou J, Zhao H, Yang H, He C, Shu W, Cui Z, Liu Q. 2022. Insights Into the Impact of Small RNA SprC
on the Metabolism and Virulence of *Staphylococcus aureus*. Front Cell Infect Microbiol 12:1–13.

22. Le Huyen KB, Gonzalez CD, Pascreau G, Bordeau V, Cattoir V, Liu W, Bouloc P, Felden B,
Chabelskaya S. 2021. A small regulatory RNA alters *Staphylococcus aureus* virulence by titrating
RNAIII activity. Nucleic Acids Res 49:10644–10656.

23. Chabelskaya S, Gaillot O, Felden B. 2010. A *Staphylococcus aureus* small RNA is required for
bacterial virulence and regulates the expression of an immune-evasion molecule. PLoS Pathog
6:1–11.

24. Chabelskaya S, Bordeau V, Felden B. 2014. Dual RNA regulatory control of a *Staphylococcus
aureus* virulence factor. Nucleic Acids Res 42:4847–4858.

25. Lalaouna D, Baude J, Wu Z, Tomasini A, Chicher J, Marzi S, Vandenesch F, Romby P, Caldelari I,
Moreau K. 2019. RsaC sRNA modulates the oxidative stress response of *Staphylococcus aureus*
during manganese starvation. Nucleic Acids Res 47:9871–9887.

26. Morrison JM, Miller EW, Benson MA, Alonzo F, Yoong P, Torres VJ, Hinrichs SH, Dunmana PM.
2012. Characterization of SSR42, a Novel virulence factor regulatory RNA that contributes to the
pathogenesis of a *Staphylococcus aureus* USA300 representative. J Bacteriol 194:2924–2938.

27. Horn J, Klepsch M, Manger M, Wolz C, Rudel T, Fraunholz M.2018.Long Noncoding RNA SSR42
Controls *Staphylococcus aureus* Alpha-Toxin Transcription in Response to Environmental Stimuli. J
Bacteriol200:10.1128/jb.00252-18.

28. DuMont AL, Nygaard TK, Watkins RL, Smith A, Kozhaya L, Kreiswirth BN, Shopsin B, Unutmaz D,
Voyich JM, Torres VJ. 2011. Characterization of a new cytotoxin that contributes to *Staphylococcus
aureus* pathogenesis. Mol Microbiol 79:814–825.

29. Brittney D. Gimza SMM& LNS. 2021. An Ex Vivo Model for Assessing Growth and Survivability of
Staphylococcus aureus in Whole Human Blood. Methods in Molecular Biology 2341:127–131.

30. DuMont AL, Yoong P, Day CJ, Alonzo F, McDonald WH, Jennings MP, Torres VJ. 2013.
Staphylococcus aureus LukAB cytotoxin kills human neutrophils by targeting the CD11b subunit of
the integrin Mac-1. Proc Natl Acad Sci U S A 110:10794–10799.

31. DuMont AL, Yoong P, Surewaard BGJ, Benson MA, Nijland R, Strijp JAG van, Torres VJ. 2013.
Staphylococcus aureus elaborates leukocidin AB to mediate escape from within human
neutrophils. Infect Immun 81:1830–1841.

32.   Ji CJ, Kim JH, Won Y Bin, Lee YE, Choi TW, Ju SY, Youn H, Helmann JD, Lee JW. 2015. *Staphylococcus*
564   *aureus* PerR is a hypersensitive hydrogen peroxide sensor using iron-mediated histidine oxidation.
565   Journal of Biological Chemistry 290:20374–20386.

566   33.   Horsburgh MJ, Clements MO, Crossley H, Ingham E, Foster SJ. 2001. PerR controls oxidative stress
567   resistance and iron storage proteins and is required for virulence in *Staphylococcus aureus*. Infect
568   Immun 69:3744–3754.

569   34.   Savin A, Anderson EE, Dyzenhaus S, Podkowik M, Shopsin B, Pironti A, Torres VJ. 2024.
570   *Staphylococcus aureus* senses human neutrophils via PerR to coordinate the expression of the
571   toxin LukAB. Infect Immun92:e00526-23.

572   35.   Mann M, Wright PR, Backofen R. 2017. IntaRNA 2.0: Enhanced and customizable prediction of
573   RNA-RNA interactions. Nucleic Acids Res 45:W435–W439.

574   36.   Manna A, Cheung AL. 2001. Characterization of *sarR*, a modulator of *sar* expression in
575   *Staphylococcus aureus*. Infect Immun 69:885–896.

576   37.   Reyes D, Andrey DO, Monod A, Kelley WL, Zhang G, Cheung AL. 2011. Coordinated regulation by
577   AgrA, SarA, and SarR to control *agr* expression in *Staphylococcus aureus*. J Bacteriol 193:6020–
578   6031.

579   38.   Tamber S, Cheung AL. 2009. SarZ promotes the expression of virulence factors and represses
580   biofilm formation by modulating SarA and *agr* in *Staphylococcus aureus*. Infect Immun 77:419–
581   428.

582   39.   Chen PR, Nishida S, Poor CB, Cheng A, Bae T, Kuechenmeister L, Dunman PM, Missiakas D, He C.
583   2009. A new oxidative sensing and regulation pathway mediated by the MgrA homologue SarZ in
584   *Staphylococcus aureus*. Mol Microbiol 71:198–211.

585   40.   Das S, Lindemann C, Young BC, Muller J, Österreich B, Ternette N, Winkler AC, Paprotka K,
586   Reinhardt R, Förstner KU, Allen E, Flaxman A, Yamaguchi Y, Rollier CS, Diemen P Van, Blättner S,
587   Remmele CW, Selle M, Dittrich M, Müller T, Vogel J, Ohlsen K, Crook DW, Massey R, Wilson DJ,
588   Rudel T, Wyllie DH, Fraunholz MJ. 2016. Natural mutations in a *Staphylococcus aureus* virulence
589   regulator attenuate cytotoxicity but permit bacteremia and abscess formation. Proc Natl Acad Sci
590   U S A 113:E3101–E3110.

591   41.   Zheng X, Marsman G, Lacey KA, Chapman JR, Goosmann C, Ueberheide BM, Torres VJ. 2021. The
592   cell envelope of *Staphylococcus aureus* selectively controls the sorting of virulence factors. Nat
593   Commun 12:1–16.

594   42.   Zheng X, Ma SX, John AS, Torres VJ. 2022. The Major Autolysin Atl Regulates the Virulence of
595   *Staphylococcus aureus* by Controlling the Sorting of LukAB. Infect Immun 90:1–13.

596   43.   Altuvia S, Weinstein-Fischer D, Zhang A, Postow L, Storz G. 1997. A small, stable RNA induced by
597   oxidative stress: Role as a pleiotropic regulator and antimutator. Cell 90:43–53.

598   44.   Altuvia S, Zhang A, Argaman L, Tiwari A, Storz G. 1998. The Escherichia coli OxyS regulatory RNA
599   represses fhlA translation by blocking ribosome binding. EMBO Journal 17:6069–6075.

45. González-Flecha B, Demple B. 1999. Role for the oxyS gene in regulation of intracellular hydrogen
peroxide in Escherichia coli. J Bacteriol 181:3833–3836.

46. Zhang H, Zhan Y, Yan Y, Liu Y, Hu G, Wang S, Yang H, Qiu X, Liu Y, Li J, Lu W, Elmerich C, Lin
M.2019.The *Pseudomonas stutzeri*-Specific Regulatory Noncoding RNA NfiS Targets *katB* mRNA
Encoding a Catalase Essential for Optimal Oxidative Resistance and Nitrogenase Activity. J
Bacteriol201:10.1128/jb.00334-19.

47. Reyes Ruiz VM, Freiberg JA, Weiss A, Green ER, Jobson M, Felton E, Shaw LN, Chazin WJ, Skaar EP.
2024.Coordinated adaptation of *Staphylococcus aureus* to calprotectin-dependent metal
sequestration. mBio0:e01389-24.

48. Horsburgh MJ, Ingham E, Foster SJ. 2001. In *Staphylococcus aureus*, Fur is an interactive regulator
with PerR, contributes to virulence, and is necessary for oxidative stress resistance through
positive regulation of catalase and iron homeostasis. J Bacteriol 183:468–475.

49. Torres VJ, Attia AS, Mason WJ, Hood MI, Corbin BD, Beasley FC, Anderson KL, Stauff DL, McDonald
WH, Zimmerman LJ, Friedman DB, Heinrichs DE, Dunman PM, Skaar EP. 2010. *Staphylococcus
aureus* fur regulates the expression of virulence factors that contribute to the pathogenesis of
pneumonia. Infect Immun 78:1618–1628.

50. Lindsay JA, Foster SJ. 2001. Zur: A Zn^{2+}-responsive regulatory element of *Staphylococcus aureus*.
Microbiology (N Y) 147:1259–1266.

51. Horsburgh MJ, Wharton SJ, Cox AG, Ingham E, Peacock S, Foster SJ. 2002. MntR modulates
expression of the PerR regulon and superoxide resistance in *Staphylococcus aureus* through
control of manganese uptake. Mol Microbiol 44:1269–1286.

52. Tavares AFN, Nobre LS, Melo AMP, Saraiva LM. 2009. A novel nitroreductase of *Staphylococcus
aureus* with S-nitrosoglutathione reductase activity. J Bacteriol 191:3403–3406.

53. Liochev SI, Hausladen A, Fridovich I. 1999. Nitroreductase A is regulated as a member of the
soxRS regulon of *Escherichia coli.* Proc Natl Acad Sci U S A 96:3537–3539.

54. Nishinaka Y, Masutani H, Nakamura H, Yodoi J. 2001. Regulatory roles of thioredoxin in oxidative
stress-induced cellular responses. Redox Report 6:289–295.

55. Ganske A, Busch LM, Hentschker C, Reder A, Michalik S, Surmann K, Völker U, Mäder U. 2024.
Exploring the targetome of IsrR , an iron-regulated sRNA controlling the synthesis of iron-
containing proteins in *Staphylococcus aureus* 1–19.

56. Coronel-Tellez RH, Pospiech M, Barrault M, Liu W, Bordeau V, Vasnier C, Felden B, Sargueil B,
Bouloc P. 2022. sRNA-controlled iron sparing response in Staphylococci. Nucleic Acids Res
50:8529–8546.

57. Rios-Delgado G, McReynolds AKG, Pagella EA, Norambuena J, Briaud P, Zheng V, Munneke MJ,
Kim J, Racine H, Carroll R, Zelzion E, Skaar E, Bose JL, Parker D, Lalaouna D, Boyd JM. 2024. The
Staphylococcus aureus small non-coding RNA IsrR regulates TCA cycle activity and virulence.
bioRxiv 2024.07.03.601953.

58.   Shaw LN, Lindholm C, Prajsnar TK, Miller HK, Brown MC, Golonka E, Stewart GC, Tarkowski A,
Potempa J. 2008. Identification and characterization of σS, a novel component of the
Staphylococcus aureus stress and virulence responses. PLoS One 3.

59.   Al Ali A, Alsulami J, Aubee JI, Idowu A, Tomlinson BR, Felton EA, Jackson JK, Kennedy SJ, Torres NJ,
Shaw LN, Thompson KM. 2023. *Staphylococcus aureus* SigS Induces Expression of a Regulatory
Protein Pair That Modulates Its mRNA Stability. J Bacteriol 205.

60.   Impens F, Rolhion N, Radoshevich L, Bécavin C, Duval M, Mellin J, Garciá Del Portillo F, Pucciarelli
MG, Williams AH, Cossart P. 2017. N-terminomics identifies Prli42 as a membrane miniprotein
conserved in Firmicutes and critical for stressosome activation in *Listeria monocytogenes*. Nat
Microbiol 2.

61.   Quin MB, Berrisford JM, Newman JA, Baslé A, Lewis RJ, Marles-Wright J. 2012. The bacterial
stressosome: A modular system that has been adapted to control secondary messenger signaling.
Structure 20:350–363.

62.   Ranganathan N, Johnson R, Edwards AM. 2020. The general stress response of *Staphylococcus
aureus* promotes tolerance of antibiotics and survival in whole human blood. Microbiology
(United Kingdom) 166:1088–1094.

Figure 1. SSR42 is a global regulator of virulence factor production. (A) CIRCOS file illustrating transcriptional changes between wild-type and ΔSSR42 at 15h of growth. The inner circle exhibits transcript expression represented as a histogram of TPM expression values for the wild type (black bars) and SSR42 mutant (blue bars). Outermost circle is a heatmap describing fold change in expression where green indicates a positive fold change and red indicates a negative fold change in comparison to the wild type. **(B)** The number of genes upregulated (green) or downregulated (pink) $\geq$2-fold in the SSR42 null strain at 15h, sorted by function based on KEGG ontology. **(C)** Heat map illustrating changes in transcript and protein abundance of key virulence factors of interest.

Figure 2. SSR42 plays a central role during engagement with the innate immune system. (A) HL-60 cells were differentiated into neutrophil-like cells with 1.25% DMSO for 4 days before being treated with 5% v/v bacterial supernatant from 15 h cultures of the strains shown. Intoxicated cells were incubated for 1 h before cell viability was measured using the CellTiter reagent. Percent cell death in comparison to TSB-treated cells is shown. **(B)** Bacterial strains were synchronized and then standardized to an OD_{600} of 0.05 in 1 mL pooled donor whole human blood. Bacterial recovery after 4 h was enumerated as CFU/mL. **(C)** Human neutrophil-like cells were incubated with the strains shown at an MOI of 30, and intracellular bacterial survival was determined after 24h. All experiments were performed in biological triplicate. Error bars represent ± SEM. Student's t test with Welch's correction was used to determine statistical significance relative to wildtype. ** $P \leq 0.01$, ,***, $P \leq 0.001$.

Figure 3. SSR42 is upregulated in response to neutrophils via PerR Derepression. (A) Differentiated HL60 cells were seeded at a concentration of 2×10^5 before the *S. aureus* WT harboring a P_{SSR42}-lux reporter fusion was added at a final OD_{600} of 0.1. Luminescence and OD_{600} reads were taken at T=0 and then hourly for 3 hours. Luminescence was normalized to OD_{600}. Data is from biological triplicate samples. **(B)** As in A but comparing P_{SSR42} expression between the wild-type and a *perR* mutant strain. A student's T-test with Welch's correction was used to determine significance. Error bars are ±SEM. * $P\leq0.05$ ** $P\leq0.01$, ,***, $P\leq0.001$.

Figure 4. SSR42 controls LukAB expression in response to neutrophils. (A) Differentiated HL60 cells were seeded at a concentration of 2×10^5 before the *S. aureus* WT harboring a P_{lukAB}-lux reporter fusion was added at a final OD_{600} of 0.1. Luminescence and OD_{600} reads were taken at T=0 and then hourly for 3 hours. Luminescence was normalized to OD_{600}. Data is from biological triplicate samples. **(B)** As in A but after 2h of exposure to neutrophils. **(C)** Data from B represented as a percentage of wild-type expression. Experiments were performed with 12 biological replicates. A student's T-test with Welch's correction was used to determine significance. Error bars are ±SEM. ****, P<0.0001

Figure 5. SSR42 post-transcriptionally controls LukAB abundance. (A) A portion of the *lukA* 5' UTR and coding sequence was translationally fused to GFP and was introduced into the wild-type or ΔSSR42 strains **(B)** The *lukA-GFP* reporter fusion plasmid was co-expressed with either an empty vector (EV) or a plasmid containing SSR42 in the ΔSSR42 strain. Fluorescence was measured every 15 minutes. Error bars are ± SEM. All experiments were performed in biological triplicate.

Figure 6. The 3' end of SSR42 mediates binding to the *lukAB 5'* UTR. (A) IntaRNA was used to identify thermodynamically likely regions of base paring between SSR42 and the *lukAB* mRNA, identifying an interaction region between the 3' end of SSR42 and the 5' UTR of *lukAB.*. Created using Biorender.com. **(B)** As in 2A but using the strains noted. Error bars are ±SEM. Student's t test with Welch's correction was used to determine statistical significance relative to wildtype. Experiments performed in biological triplicate. * P≤0.05, ,***, P≤0.001.

Figure 7. SSR42 directly binds and stabilizes the *lukAB mRNA*. (A) *In vitro* transcribed SSR42 was added in increasing amounts (1,000-fold to 1,500-fold excess) to radiolabeled portions of the *lukAB* mRNA. Samples were heated to 90° C for 2 min and then incubated at room temperature for 30 min before being loaded on to a non-denaturing gel. At identical concentrations, no binding is seen for radiolabeled *splE*. Images shown are representatives of 5 experimental replicates. **(B)** Bacterial cultures were grown in biological triplicate for 15 h before transcription was halted by the addition of 250 µg/mL Rifampin. Samples were collected 0, 5, 10, 30, and 45 minutes post-transcriptional arrest and *lukA* mRNA abundance was assessed via qPCR analysis. Data was compared to initial RNA abundance at t=0 and normalized to 16S rRNA. Shown is a one phase decay curve. Error bars are ± SEM.

Figure 8. SSR42 is required for full virulence in murine models of invasive disease. (A) CD-1 mice were inoculated with the wild-type or SSR42 mutant at 5×10^8 CFU via tail vein injection. Following 7 days of infection, organs were harvested and CFU determined. **(B)** C57BL/6J mice were intranasally inoculated with the wild-type or SSR42 mutant at 1×10^8 CFU. After 24h, lungs were harvested and CFU determined. Whiskers represent minimum and maximum values, Solid lines represent median values. Mann Whitney tests were used to determine statistical significance relative to wildtype. * $P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$. n=8 for each strain, error bars are ±SEM.

Figure 9. Proposed role of SSR42 in *S. aureus*. The presence of neutrophils and/or reactive oxygen species leads to PerR-mediated derepression of SSR42, which then binds and stabilizes *lukAB* mRNA in order to defend against phagocytosis and/or facilitate escape. The regulatory links between the metalloregulators, stress response, and oxidative stress resistance is represented. Created with BioRender.com

Table 1. Bacterial strains and plasmids used in this study.

Strain Name	Description	Source
E. coli		
DH5α	Cloning strain	(1)
JM109	Cloning strain	Promega pGEM-T Easy Vector Kit
MEJ3834	JM109 pGEM-T::*lukA* UTR	This study
MEJ3845	JM109 pGEM-T::SSR42	This study
S. aureus		
RN4220	Cloning strain, restriction-deficient	Lab stock
TCH1516	USA300 HOU CA-MRSA Erm^s, Wildtype parent strain	(2)
BRT2544	TCH1516 ΔSSR42	This study
NE665	JE2 *perR*::tn	(3)
MEJ4002	TCH1516 *perR*::tn	This study
MEJ4003	TCH1516 ΔSSR42, *perR*::tn	This study
LNS1927	TCH1516 pMK4 EV	This study
BRT2545	TCH1516 ΔSSR42 pMK4 EV	This study
BRT2546	TCH1516 ΔSSR42 pMK4::SSR42	This study
MEJ4000	TCH1516 pXen-1::P$_{SSR42}$	This study
MEJ4001	TCH1516 *perR*::tn, pXen-1::P$_{SSR42}$	This study
MEJ3546	TCH1516 pXen-1::P$_{lukAB}$	This study
MEJ3550	TCH1516 ΔSSR42 pXen-1::P$_{lukAB}$	This study
MEJ3826	TCH1516 *perR*::tn, pXen-1::P$_{lukAB}$	This study
MEJ3827	TCH1516 ΔSSR42, *perR*::tn, pXen-1::P$_{lukAB}$	This study
MEJ3828	TCH1516 ΔSSR42, *perR*::tn, pXen-1::P$_{lukAB}$, pJB67::PerR	This study
MEJ3831	TCH1516 pCN33::*lukA* UTR	This study
MEJ3832	TCH1516 ΔSSR42 pCN33::*lukA* UTR	This study
MEJ3833	TCH1516 ΔSSR42 pICS3::*SSR42*	This study
MEJ3823	TCH1516 ΔSSR42 pICS3::SSR42-648-nt	This study
MEJ3824	TCH1516 ΔSSR42 pICS3::SSR42-793-nt	This study
MEJ3825	TCH1516 ΔSSR42 pICS3::SSR42-940-nt	This study

Plasmids	Description	Source
pJB38	Allelic replacement vector, *E. coli* Amp^R, *S. aureus* CM^R	(4)
pMK4	Shuttle vector, *E. coli* Amp^R, *S. aureus* CM^R	(5)
pXen-1	Luciferase reporter plasmid, *E. coli* Amp^R, *S. aureus* CM^R	(6)
pJB67	pCN51 with optimized RBS and Cd-inducible promoter, *E. coli* Amp^R, *S. aureus* Ery^R	(7)
pICS3	RNA expression vector, *E. coli* Amp^R, *S. aureus* CM^R	(8)
pCN33	mRNA target GFP reporter, *E. coli* Amp^R, *S. aureus* Ery^R	(8)
pGEM-T	pGEM-T Easy Vector cloning plasmid	Promega pGEM-T Easy Vector Kit

Table 2. Primers used in this study.

Primer	Sequence*	Description[†]
OL2939	ATG<u>GGTACC</u>CGTTATCTTGTTGGAAGT	F1 SSR42 deletion
OL3592	TCAGCAATTCAT<u>ACGCGT</u>GCACCAAATAATTTAATTAGACTC	R1 SSR42 deletion
OL2389	ATG<u>GAATTC</u>AATCGCTGTAACGGATTCAT	F2 SSR42 deletion
OL3593	ATTATTTGGTGC<u>ACGCGT</u>ATGAATTGCTGAACTCCAATG	R2 SSR42 deletion
OL2380	GCCACCTGACGTCTAAGAAACC	F MCS pJB38ΔSSR42
OL2381	GCGATTTACATATGAGTTATGC	R MCS pJB38ΔSSR42
OL2888	ACT<u>GGATCC</u>GTGTCTATAATACTTTGACCT	F SSR42 complementation
OL3734	ATG<u>GTCGAC</u>GTTCAATATTTACTACAAAGTCGTG	R SSR42 complementation
OL2393	TCGTATGTTGTGTGGAATTG	F MCS pMK4
OL2394	GTGCTGCAAGGCGATTAAG	R MCS pMK4
OL398	TCCTACGGGAGGCAGCAG T	F 16S
OL399	GGACTACCAGGGTATCTA ATCCTGTT	R 16S
OL7625	CAAATAGCTACTCCAAAACGATTAGTTATAA	F RT-qPCR *lukA*
OL7506	CAGGGTTTTCTACAGTAGCAATTC	R RT-qPCR *lukA*
OL2123	GAG TTG TTA TCA ATG GTC AC	F RT-qPCR *sarA*
OL2124	ACT GCT TTA ACA ACT TGT GG	R RT-qPCR *sarA*
OL2487	TAGTGGTCCATCAACAG	F RT-qPCR *lukS*-PV
OL2488	ACGTTCTACTTCACTGATA	R RT-qPCR *lukS*-PV
OL4036	GGCTCTATGAAAGCAGCAGATA	F RT-qPCR *hla*
OL4037	CTGTAGCGAAGTCTGGTGAAA	R RT-qPCR *hla*
OL5048	GGCAGTGGCTCATTCAACTAC	F RT-qPCR *hlgA*
OL5049	CACCTTTAGAGTTCTGACTTTCTAC	R RT-qPCR *hlgA*
OL2170	GGC TAA AGC TGA ACA TAA TG	F RT-qPCR *splE*
OL2171	ATA TTC ACC ATT GGG ATG C	R RT-qPCR *splE*
OL6159	GTCGCACATTCACAAGTTTATC	F RT-qPCR *aur*
OL6160	GCCTGACTGGTCCTTATATTC	R RT-qPCR *aur*
OL4526	ATG<u>GAATTC</u>GTTGGCATGTCATATTTCTCTCTCCTG	F SSR42 promoter for pXen-1
OL7767	ATG<u>GGATCC</u>CGAATATTTTGTAACAGGGCTACTAAG	R SSR42 promoter for pXen-1
OL6607	AATT<u>GAATTC</u>ATTAACTGTCTAGTATCAACGATC	F *lukAB* promoter for pXen-1
OL6608	AATT<u>GGATCC</u>AAACACGTTTTTTATTTTTCAT	R *lukAB* promoter for pXen-1
OL7693	CACGCGCGCAATACTACAG	*perR* screening
OL7694	CGCTTACTTATATGAAGTAAGCTAACG	*perR* screening
OL7501	AATT**TAATACGACTCACTATAGGG**TCGGATATGCTTAATTTTATAGTAAATTGTATG	F *lukA* 5' UTR fragment + T7

		promoter
OL7502	AAATTGCACATGATAATGATGACG	R *lukA* 5' UTR fragment
OL7244	ACT<u>CTGCAG</u>GTGTCTATAATACTTTGACCT	F SSR42 for pICS3
OL7245	ATG<u>GAATTC</u>GTTCAATATTTACTACAAAGTCGTG	R SSR42 for pICS3
OL7337	AATTGTCGACTCTCTTGATGTAAAATCTAAGATGTTTATGC	F SSR42 648-nt truncation
OL7338	AATTGTCGACATGTGGATGCCTGTCTTGATG	F SSR42 793-nt truncation
OL7339	AATTGTCGACATGTAGAAATTGAGTGTGAAAGTTAATAATAGAT	F SSR42 940-nt truncation
OL7499	AATT**TAATACGACTCACTATAGGG**GATTTCAAACCTATGTATTTC	F SSR42 +T7 promoter for pGEM-T
OL7500	AATATGATGTTCAATATTTACTACAAAGTCG	R SSR42 for pGEM-T
OL7501	AATTT**AATACGACTCACTATAGGG**TCGGATATGCTTAATTTTATAGTAAATTGTATG	F *lukA* UTR +T7 promoter for pGEM-T
OL7502	AAATTGCACATGATAATGATGACG	R *lukA* UTR for pGEM-T
OL7523	AATTT**AATACGACTCACTATAGGG**GCAGATAATTTAGATAAATAAATCATCCATCCATA	F *splE* UTR +T7 promoter for pGEM-T
OL7524	TACCCTCAACCACTGTTGT	R *splE* UTR for pGEM-T
OL6163	ATG<u>GAATTC</u>ATGCACCAAATAATTTAATTAGACTCTATCG	F *rsp* promoter for pXen-1
OL6164	ATG<u>GGATCC</u>CATGTCATATTTCTCTCTCCTGCC	R *rsp* promoter for pXen-1

*Underline indicates restriction enzyme site, bold indicate T7 promoter.
†F = Forward; R = Reverse; MCS = Multiple Cloning Site

1. Salisbury V, Hedges RW, Datta N. Two modes of "curing" transmissible bacterial plasmids. J Gen Microbiol. 1972;70(3):443-52.
2. Kolar SL, Nagarajan V, Oszmiana A, Rivera FE, Miller HK, Davenport JE, Riordan JT, Potempa J, Barber DS, Koziel J, Elasri MO, Shaw LN. NsaRS is a cell-envelope-stress-sensing two-component system of Staphylococcus aureus. Microbiology (Reading). 2011 Aug;157(Pt 8):2206-2219. doi: 10.1099/mic.0.049692-0. Epub 2011 May 12. PMID: 21565927; PMCID: PMC3167884.
3. Fey P. D., Endres J. L., Yajjala V. K., Widhelm T. J., Boissy R. J., Bose J. L., et al.. (2013). A genetic resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. mBio 4, e00537–e00512. doi: 10.1128/mBio.00537-12
4. Bose JL, Fey PD, Bayles KW. Genetic tools to enhance the study of gene function and regulation in *Staphylococcus aureus.* Appl Environ Microbiol. 2013;79(7):2218-24.
5. Sullivan MA, Yasbin RE, Young FE. New shuttle vectors for *Bacillus subtilis* and *Escherichia coli* which allow rapid detection of inserted fragments. Gene. 1984;29(1-2):21-6.
6. Francis K. P., Joh D., Bellinger-Kawahara C., Hawkinson M. J., Purchio T. F., Contag P. R. (2000). Monitoring bioluminescent *Staphylococcus aureus* infections in living mice using a novel luxABCDE construct. Infect. Immun. 68, 3594–3600. doi: 10.1128/IAI.68.6.3594-3600.2000.
7. Windham IH, Chaudhari SS, Bose JL, Thomas VC, Bayles KW. 2016. SrrAB modulates *Staphylococcus aureus* cell death through regulation of cidABC transcription. J Bacteriol 198:1114–1122. doi: 10.1128/JB.00954-15.
8. Ivain L, Bordeau V, Eyraud A, Hallier M, Dreano S, Tattevin P, Felden B, Chabelskaya S. An in vivo reporter assay for sRNA-directed gene control in Gram-positive bacteria: identifying a novel sRNA target in *Staphylococcus aureus.* Nucleic Acids Res. 2017 May 5;45(8):4994-5007. doi: 10.1093/nar/gkx190. PMID: 28369640; PMCID: PMC5416835.