Commensal-pathogen dynamics structure disease outcomes during *Clostridioides difficile* colonization

Skye R. S. Fishbein[1,2*], Anna L. DeVeaux[1*], Sakshi Khanna[1,2*], Aura L. Ferreiro[1,2], James Liao[1,2], Wesley Agee[1], Jie Ning[1,2], Bejan Mahmud[1], Miranda J. Wallace[1,2], Tiffany Hink[3], Kimberly A. Reske[3], Janaki Guruge[1,2], Sidh Leekha[1], Erik R. Dubberke[3], Gautam Dantas[1,2,4-6**]

[1]The Edison Family Center for Genome Sciences and Systems Biology, Washington University School of Medicine, St. Louis, Missouri, USA.

[2]Department of Pathology and Immunology, Division of Laboratory and Genomic Medicine, Washington University School of Medicine, St. Louis, Missouri, USA.

[3]Division of Infectious Diseases; Washington University School of Medicine, St. Louis, Missouri, USA

[4]Department of Molecular Microbiology; Washington University School of Medicine, St. Louis, Missouri, USA

[5]Department of Biomedical Engineering; Washington University in St Louis, St. Louis, Missouri, USA

[6]Department of Pediatrics; Washington University School of Medicine, St. Louis, Missouri, USA

* these authors contributed equally

** corresponding author: GD (dantas@wustl.edu)

Author contact during review: Skye R. S. Fishbein (skye.f@wustl.edu)

Abstract

Gastrointestinal colonization by *Clostridioides difficile* is common in healthcare settings and ranges in clinical presentation from asymptomatic carriage to lethal *C. difficile* infection (CDI). We used a systems biology approach to investigate why patients colonized with *C. difficile* have a range of outcomes. Microbiota-humanization of germ-free mice with fecal samples from toxigenic C. difficile carriers revealed a spectrum of virulence among clade 1 lineages and identified commensal *Blautia* associated with markers of non-pathogenic colonization. Using gnotobiotic mice engrafted with defined human microbiota, we observed strain-specific CDI severity across clade 1 strains. Yet, mice engrafted with a higher diversity community were protected from severe disease across all strains without suppression of *C. difficile* colonization. These results indicate that when colonization resistance has been breached without overt infection, commensals can attenuate a diversity of virulent strains without inhibiting pathogen colonization, providing insight into determinants of stable *C. difficile* carriage.

Introduction

Healthcare-associated diarrhea caused by *Clostridioides difficile* remains difficult to predict, prevent, and treat, resulting in a significant and global public health burden[1]. While *C. difficile* infection (CDI) can be fatal, especially in the case of recurrence, *C. difficile* colonization in the absence of CDI (hereafter, *carriage*) is common. Recent studies in U.S. hospitals, by our group and others, have shown that toxigenic *C. difficile* carriage is more prevalent than symptomatic CDI, affecting approximately 10% of admissions in acute care and cancer-related cohorts[2,3]. Concerningly, these asymptomatic carriers are at substantially increased risk of developing future CDI, despite the possibility that these patients can also remain carriers without CDI for months at a time[2,3]. Despite the initial success of live biotherapeutics for recurrent CDI[4], there is an unmet need to prophylactically intervene with such vulnerable patient groups, especially in a healthcare setting. Understanding what biological factors influence where patients lie along the spectrum from asymptomatic *C. difficile* carriage to CDI and identifying intervention targets is thus a public health priority.

The epidemiology of CDI has changed over the past two decades, with hypervirulent *C. difficile* lineages from clade 2 that were responsible for past epidemics (e.g., PCR ribotype 027/ST1[5]) no longer causing most healthcare-associated CDI. Genomic surveillance studies have instead found that clade 1 *C. difficile* strains (e.g., ST2, ST8, ST3, ST42)[2,3,6] are predominant among currently circulating lineages isolated from both patients with CDI and asymptomatic carriers in multiple regionally distinct healthcare settings in the U.S.[2,3,6,7]. *C. difficile* is genetically diverse[16], with strains showing marked differences in toxin production and metabolic preferences[8]. Quantifying the relative virulence of clinically prevalent *C. difficile* lineages, and their modulation by commensal microbes, is required to identify strategies aimed at preventing the transition from carriage to CDI in patients.

CDI is a paradigmatic disease of the gut microbiome, as loss of colonization resistance through microbiome perturbation often precedes CDI development[9]. Indeed, the remarkable efficacy of fecal microbiota transfers in preventing recurrent CDI depends upon restoration of colonization resistance[10]. Microbiota-conferred colonization resistance prevents *C. difficile* germination and vegetative growth through commensal bile acid, carbohydrate, amino acid, and micronutrient metabolism[11-14]. Contrastingly, certain commensal species such as *Bacteroides thetaiotaomicron* or *Enterococcus faecalis* have been shown to cross-feed *C. difficile*, supplying amino acids and carbohydrates to enable pathogenic growth and in some instances fuel disease onset[15-18]. The high prevalence of *C. difficile* carriage, where colonization resistance has been lost yet destructive toxigenesis is blunted, suggests that understanding these complex commensal-pathogen interactions may be key to preventing and treating CDI in vulnerable patients.

Validation of predicted commensal-pathogen interactions necessitates *in vivo* systems that successfully model patient microbiome structures while controlling for pathogen and commensal variation. Innovative gnotobiotic animal models using diverse defined communities have allowed us to understand the strain-specific nature of carbohydrate, amino acid, and bile acid metabolism in the gut microbiome, and such experimental systems would be ideal for gaining a deeper quantitative understanding of colonization resistance or resilience against *C. difficile*[19,20]. Further, it remains unclear how virulent circulating lineages of *C. difficile* are relative to earlier epidemic variants, complicating our understanding of how to pre-empt disease diagnostically or therapeutically. The identification

of microbiome communities with desired resilience against diverse strains *in vivo*, and further modeling community dynamics using defined consortia, holds promise for examining how specific members of the microbiome can modulate the pathogenicity of *C. difficile*.

In this study, we use a systems biology approach to investigate how commensal-pathogen interactions influence the outcome of *C. difficile* colonization. We demonstrate how human gut microbiome structures impact *C. difficile* virulence by associating microbiome features with CDI status in colonized patients and identifying stable commensal-pathogen dynamics from patient microbiomes in microbiota-humanized gnotobiotic animal models. We present evidence for antagonistic roles of *Blautia* spp. in modulating *C. difficile* growth *in vivo* across a diverse collection of clinically relevant pathogen lineages. Importantly, our construction of defined gut microbial communities from patient stools allowed us to control for microbiome variation and identify that community structure can prevent toxin-mediated disease across these strains without inhibiting its colonization. These observations identify a dual role for the microbiome in supporting *C. difficile* growth while suppressing pathogenesis, which may have implications for the prevention of CDI in colonized patients and the design of novel microbiome-targeting therapies.

Clinical and microbial features are discriminatory of CDI status in colonized patients

Identifying commensal, pathogen, and clinical features that render *C. difficile*-colonized patients vulnerable to CDI is essential to improving CDI treatment and prevention. Our previous examination of *C. difficile* carriers and patients with CDI revealed interactions between the microbiome, antibiotic exposure, and pathogen correlates that could drive patient outcomes[21]. We sought to investigate how these multilayered interactions underlie CDI risk in a broader group of hospitalized patients and controls. We leveraged stool samples and microbiome data from adult hospitalized cohorts between 2014 and 2019 of patients with CDI (CDI+, *C. difficile*+; $n = 53$) and *C. difficile* carriers (CDI-, *C. difficile*+; $n = 96$), with corresponding high-quality toxigenic *C. difficile* genome assemblies from 141/149 samples. We define CDI- as patients whose stool was either toxin enzyme immunoassay negative (EIA-) or not tested (based on a lack of clinical suspicion of CDI); toxigenic *C. difficile* carriage (*C. difficile*+) was confirmed by culture and/or stool nucleic acid amplification test (NAAT, see Methods for more detail). To contextualize the hospitalized patient microbiome, we additionally selected patient samples from these cohorts who did not have CDI symptoms and were *C. difficile* culture negative [hospitalized controls (HC); CDI-, *C. difficile*-; $n = 39$]. When available, we included longitudinal samples in our analysis and controlled for patient ID. Lastly, we extended our investigation to a control set of healthy volunteers without recent antibiotic exposure within the same geographic area (St. Louis, MO, USA)[22] [healthy volunteers (HV); CDI-, *C. difficile*-; $n = 20$] (Fig. 1A). In total, we analyzed 208 stool metagenomes from 187 adults representing a spectrum of *C. difficile* colonization and disease status (Fig. 1A).

HVs had significantly increased taxonomic richness relative to carriers and patients with CDI, whereas HCs represented an intermediate group (Fig. 1B). This trend was also reflected by differences in species Shannon diversity (Fig. S1A) and microbiome structure as measured by beta diversity for HVs compared to all hospitalized patients ($p = 1e-04$, PERMANOVA; Fig. S1B). As many of these differences may be mediated by antibiotic exposure, we compared quantitative 30-day antibiotic exposures and identified significant differences in cephalosporin, fluoroquinolone, and intravenous (IV) vancomycin exposure across hospitalized patient groups (Table S1). Using ShortBRED[23] to profile antibiotic-resistance genes (ARGs) in the microbiome, we observed an increase in both ARG burden and richness concomitant with *C. difficile* colonization and CDI, highlighting the substantial influence of antimicrobial exposure on differences in microbiome structure within this population (Fig. S1C-D). Finally, we integrated pathogen genomic information from the corresponding *C. difficile* strains of colonized patient samples[2] and noted diverse representation of clades 1, 2, 4, 5, and cryptic clade 1 among our sample set, with a predominance of clade 1 among both carriers and patients with CDI (33/53 CDI cases and 68/78 carriers with cognate isolate assemblies; Fig. 1C)[6], which matches other reports in the United States indicating the abundance of clade 1 in health-care settings[3,7]. Additionally, we observed that pangenomic differences in gene content among these selected strains clustered by clade (Fig. 1D). As previously observed[21], patients colonized with strains encoding the accessory binary toxin, *cdtB*, were more likely to have CDI ($p = 0.0028$, Fisher's exact test).

Because of these significant microbiome, antibiotic exposure, and pathogen correlates of CDI, we computed the relative

importance of these variables for distinguishing between patients with CDI and asymptomatic carriers. *C. difficile* relative abundance is

highly correlated with CDI status using linear mixed effect models (coef = 2.26, *qval* = 1.4e-07; Fig. S1E) and may overshadow the

importance of other features; thus, we excluded it from downstream modeling tasks. We trained a random forest classifier on the relative

abundance of feature-selected commensal taxa (*n* = 6 species), recent antibiotic exposures, participant demographic data, and the

presence of *cdtB* for a randomized subset of colonized patients (60%) and assessed its accuracy in predicting CDI in the remaining data

(40%) (see Methods). The highest performing model (of 100 iterations) had an accuracy of 84.3% with a positive predictive value of

80% and negative predictive value of 87.1% (Fig. S1F), demonstrating its effectiveness in discriminating between carriers and patients

with CDI. Interestingly, in the absence of *C. difficile* relative abundance data, we observed that cephalosporins, IV vancomycin, *Blautia*

wexlerae, and *Veillonella parvula* were all significantly important features in predicting disease status, while surprisingly, the mean

feature importance of *cdtB* did not outperform those of models trained with randomly shuffled class labels (Fig. 1E). Exposure to

cephalosporins or IV vancomycin and the abundance of *V. parvula* were all enriched in patients with CDI relative to carriers (Fig. S1E

and Table 1), which corroborates the known influence of these antibiotic classes on CDI risk and identifying *V. parvula* as a commensal

signature of CDI (Fig. S1E). In contrast, *B. wexlerae* was depleted in patients with CDI and negatively associated with recent

cephalosporin exposure in colonized patients (Fig. S1E, G). In all, we identified specific antibiotic and commensal variables that were

discriminatory of CDI status and found that *cdtB* was not an important feature despite its association with CDI. However, these insights

also underscore the difficulty in identifying meaningful pathogen-microbiome interactions from metagenomic data, which are limited

in ascribing functional importance to associations. Accordingly, we designed an *in vivo* microbiome engraftment screen to better

understand the ecological determinants of *C. difficile* colonization dynamics.

Microbiome engraftment screen reveals strain-specific drivers of commensal-pathogen dynamics

We hypothesized that we could leverage the natural microbiome-pathogen variation in patient stool samples to identify

functional drivers of *C. difficile* proliferation *in vivo*. We devised a screen based on the premise that germ-free mice administered

toxigenic strains of *C. difficile* succumb to lethal infection within 48 hours of gavage even at relatively low titers[12]. Thus, any non-lethal

engraftment would reflect a stable equilibrium between the commensal microbiota and *C. difficile* (Fig. 2A). We selected 15 stools, or

donors, from *C. difficile* carriers without CDI that reflect the ecological variation of colonized patient microbiomes and one stool from

a HV (Donor HV), which lacks *C. difficile*, to engraft into germ-free mice (*n* = 3-8 mice per arm). We followed mice for 21 days post-

engraftment to track disease, measuring clinical health, *C. difficile* burden, toxin production, and microbiome composition. Upon

engraftment, we observed a spectrum of lethality (Fig. 2B), with 33% (5/15) of donor communities resulting in some lethality within

the first week, while the rest (10/15) resulted in non-lethal engraftment. There were also community-specific differences in weight loss

trajectories during the first week, with lethal donors exhibiting significant weight loss relative to Donor HV, which did not contain *C.*

difficile (Fig. S2A). We first investigated how well these stool engraftment experiments modeled human microbiome ecologies in the

context of *C. difficile* colonization. Comparison of human microbiome structures of the selected stools revealed that the *C. difficile*

burden of the original donor stools correlated with distance (as measured by beta diversity) from the HV cluster, where no *C. difficile*

was detected (Fig. 2C, D; Pearson *R* = 0.76, *p* = 0.0011), suggesting that a higher degree of dysbiosis correlates with *C. difficile*

abundance. While there was significant correlation (Spearman *R* = -0.71, *p* = 0.0031) between donor lethality in the first week and *C.*

difficile colony-forming units (CFU)/mL in the inoculum gavage (Fig. S2B), one donor (Donor 3) caused lethal disease despite a

relatively low *C. difficile* inoculum titer. At 21 days, mice from donors surviving initial engraftment (thus excluding donor 5 and 7) had

a community species richness that fell within the range of colonized human microbiome richness (species richness of colonized patient

microbiomes ranged from 12-74; Fig. 2E). To investigate *C. difficile* fitness and virulence in non-lethal donor groups, we compared *C.*

difficile CFUs and TcdB levels in mouse stools at an early point (days 4-5) and a late timepoint (day 21) and calculated ratios of *C.*

difficile growth and toxin production throughout engraftment (where a value above 1 indicates an increase in pathogen abundance or
toxin levels over time), revealing donor-specific trajectories in *C. difficile* proliferation and toxin levels post-engraftment (Fig. S2C, D).
Our human microbiome-genomic data suggested that microbiome structure and not *C. difficile* strain identity drives
symptomatic severity. In this functional microbiome screen, we sought to reexamine the contribution of these two factors to longitudinal
differences in *C. difficile* growth and virulence following stool microbiome engraftment. Further, given our and others' observations that
clade 1 isolates are the major circulating strains in the United States[2,3,6,7], our microbiome engraftment screen contained mostly stools
with *C. difficile* strains from this clade (14/15 stools), including both closely related isolates and unrelated isolates clusters, based on
whole-genome information (Fig. 2F). To understand strain-specific influences on outcomes in our microbiome engraftment screen
(weight-loss, *C. difficile* titers, and toxin levels), we aggregated clade 1 donor strains into groups described by their multi-locus sequence
type (ST), based on a threshold of >99.9% ANI by core-gene alignment of selected clade 1 genomes. Examination of weight-loss
differences revealed that stools containing ST8 and ST43 strains caused significantly more weight-loss relative to other strains and the
ST2 strain group (Fig. 3A). Though some stools containing these two strain groups had high levels of *C. difficile* in the gavage inoculum,
Donor 3 contained a low level of its ST8 strain (Fig. S2B), suggesting that this strain group may be more virulent than other clade 1
isolates examined here. To examine strain influence on growth and toxin production, we restricted our measurements to strain groups
with multiple non-lethal stool microbiome backgrounds. Interestingly, the ST2 strain group had decreased *C. difficile* proliferation and
toxin production over time relative to *C. difficile* strains in all other donors, suggestive of lower virulence independent of the microbiome
(Fig. 3B, 3C). Juxtaposed with the increased lethality associated with ST8 strains from Donor 3 and Donor 8, our microbiome
engraftment screen revealed significant variation in clade 1 strain fitness and toxin production *in vivo*.
We leveraged the above pathogen genomic information to more precisely define the commensal contribution to *C. difficile*
proliferation and toxin production in our screen. Using beta diversity across all sequenced murine microbiomes to generate a principal
coordinate analysis, we confirmed human donor-specific clustering of gnotobiotic mouse microbiome structures (Fig. 3D) and identified
a significant correlation between gnotobiotic mouse microbiome structure at day 21 and corresponding mouse stool levels of *C. difficile*
(Fig. 3E, $R = 0.65$, $p < 0.001$) and, to a lesser extent, its toxin (Fig. 3E, $R = 0.38$, $p = 0.0036$). We then used linear mixed effect modeling
to investigate taxonomic features that were associated with quantitative differences in *C. difficile* growth in stool while controlling for
C. difficile strain identity and other microbiome-relevant factors (see Methods). Importantly, our screen revealed that *Blautia* was
strongly associated with decreased *C. difficile* growth (Fig. 3F), validating correlative data from our human microbiome analyses
indicating that *Blautia wexlerae* depletion (perhaps related to antibiotic exposure) was a signature of CDI in patient microbiomes (Fig.
1E, S1E). Based on these metagenomic findings from both patient samples and gnotobiotic colonized communities *in vivo*, we predict
that *Blautia* spp. interact with *C. difficile* during stable colonization and modulate its growth. We further find that several carbohydrate
catabolism pathways, including glycogen degradation and three proteoglycan-related sugar acid degradation pathways, were associated
with increased *C. difficile* growth (Fig. S2E). These taxonomic and metabolic pathway associations indicate that *Blautia* and complex
glycan metabolism may be involved in stabilizing *C. difficile* during non-pathogenic colonization *in vivo*.
178
Defined microbiome enables *C. difficile* colonization and suppresses strain-specific virulence
While we identified multiple correlates of *C. difficile* proliferation during colonization *in vivo*, including both strain identity
and microbiome composition, we sought to further understand strain differences in the globally prevalent clade 1 and identify strategies
by which the microbiome alters *C. difficile* virulence in a controlled manner. To achieve this, we built two defined bacterial culture
collections representative of patients carrying *C. difficile* in our cohort as a background for subsequent *C. difficile* challenge *in vivo*.
These defined communities were designed to either 1) recapitulate the phenotype of a candidate non-lethal donor in our microbiome
engraftment screen or 2) broadly represent the most prevalent taxa seen in *C. difficile* colonized patients based on metagenomic analysis.
Thus, for the first defined community (DefCom1), we sought to reassemble the community (except for *C. difficile*) from donor 4, for
which we observed intermediate weight loss *in vivo* (Fig. S2A); DefCom1 comprised 24 species representing 98% of the relative

abundance of species observed in the original donor stool (Supplementary Table 2). For the second community (DefCom2), we

supplemented DefCom1 with additional taxa to represent at least 70% of the 30 most prevalent genera among colonized patients, totaling

46 commensal species (Fig. 4A). We pooled each set of isolates and engrafted germ-free mice with two consecutive daily gavages of

either the DefCom1 or DefCom2 defined culture collection. After 28 days, both communities were representative of human microbiome

states found in colonized patients (Fig. S3A).

To investigate differences in virulence among the diverse clade 1 lineage, we chose three clade 1 strains from our engraftment

experiments that reflected differences in mouse weight loss, lethality, and *C. difficile* growth. Specifically, we selected strains natively

colonizing donor 4 (the source community for DefCom1) from the least 'virulent' strain group (ST2, Fig. 2D, F); donor 6 (ST53), which

had the highest TcdB load in non-lethal donors; and donor 3 (ST8), for which we observed intermediate lethality at a relatively low *C.*

difficile load and thus predicted higher virulence (Fig. 2D). Following engraftment with DefCom1 or DefCom2, mice were challenged

with 10^4 CFU of ST2, ST53, ST8, or a ribotype 027 reference strain (ST1, UK1)[24]. Relative to the hypervirulent ST1 infection, we

observed strain-specific virulence of clade 1 isolates (Fig. 4C) across both communities, with ST8 resulting in intermediate disease and

ST2 and ST53 causing mild disease. Notably, DefCom1 colonization resulted in more severe disease, as measured by lethality and

weight loss, upon *C. difficile* challenge for all strains relative to DefCom2. DefCom2 suppressed strain-specific differences in weight

loss relative to microbiome structures in both DefCom1 and human fecal engraftment backgrounds (Fig. 4D, data from Fig. S2);

interestingly, this effect was most pronounced for ST8, the most virulent clade 1 isolate.

We sought to validate the observed difference in microbiome protection between DefCom1 and DefCom2 using ST2 and ST8

strains in an independent experiment; a comparison of microbiome engraftment at Day 28 indicated that we reproduced community

structure between experimental replicates (Fig. S3B, C). In addition to recapitulating weight loss severity differences (Fig. 4E), we found

that DefCom2 resulted in decreased toxin production and decreased intestinal inflammation (as measured by calprotectin, which is

induced during CDI [25,26]) relative to DefCom1 for both strains (Fig. 4F). Surprisingly, for the moderately virulent ST8 strain, DefCom2

supported a higher level of *C. difficile* as measured by CFUs (Fig. S3D) relative to DefCom1. To rule out the possibility that absolute

bacterial loads explained the difference in ST8 *C. difficile* CFU load between communities (i.e., carrying a similar abundance of *C.*

difficile relative to community absolute abundance), we plated pre-challenge (day 28) stools from each repeat experiment on LYBHI

with sodium taurocholate to recover the majority of spore and vegetative cells in the community. Interestingly, we observed equivalent

bacterial load between DefCom1 and DefCom2 after two days of growth (Fig. S3E), suggesting preferential growth of ST8 in DefCom

2 background.

Given the results from our microbiome engraftment screens, we postulated that the community-specific difference in disease

severity and the promotion of ST8's growth by DefCom2 reflects the microbiome's capacity to cross-feed *C. difficile*, thus encouraging

its proliferation while attenuating virulence. Specifically, based on the well-studied repression of toxin production in the presence of

metabolizable nutrients[27], we predict that the increased flux of host- and dietary-derived nutrient degradation products released by

commensals represses *C. difficile* toxin production. Investigation of metagenomically-inferred metabolic pathway differences between

DefCom1 and DefCom2 predominantly revealed an enrichment in glycan degradation pathways, including that of host-derived

glycosaminoglycans and plant polysaccharides (Fig. S3F). These associations support a potential role for glycan metabolism in

promoting *C. difficile*-commensal interactions during stable colonization, which may contribute to its attenuated virulence across Clade

1 strains in DefCom2.

Discussion:

C. difficile 630 (a clade 1 isolate of ST54[28,29]) and a number of well-studied clade 2 strains[29-31] are genetically tractable, and

these strains have enabled a foundational understanding of how this bacterial pathogen grows and causes disease[29]. Importantly, previous

work in a conventional antibiotic-induced CDI murine model has demonstrated differential virulence between *C. difficile* 630 and these

epidemic strains, recapitulating both CDI and non-pathogenic colonization[32]. However, ribotype 012, which includes *C. difficile* 630, is

an uncommon lineage identified in patients today and may not be sufficiently representative of other clade 1 strains in terms of studying
C. difficile colonization outcomes. Given our growing understanding of the vast diversity of *C. difficile* lineages (owing to its open
pangenome[6]), we sought to expand our understanding of clade 1 virulence in the context of gnotobiotic models of patient-derived
microbiomes. Our data show that, across two microbiome backgrounds, only one of the clade 1 isolates tested (a strain from ST8) causes
levels of initial weight loss in mice equivalent to those of the hypervirulent ST1 from clade 2 (Fig. 4). These data are consistent with
previous reports that ribotype 002, to which our ST8 strain belongs, may have significant virulence relative to other common circulating
clade 1 lineages[33,34]. In contrast, the ST2 strain produced minimal weight loss, supporting the trend of overt differences in virulence that
others have observed between these clades of *C. difficile* and underlining the variation in clade 1 virulence[32]. Notably, ST8 and ST2 are
the most globally prevalent clade 1 lineages of *C. difficile*[6]. Although ribotype 027 (clade 2) was still the single most common ribotype
identified in patients with healthcare-associated CDI in 2018, from an epidemiologic perspective, it appears that historically less virulent
strains are the predominant source of *C. difficile* carriage and CDI. Future investigation of how clade 1 relative to other clade 1 and
clade 2 isolates cause severe disease in equivalent microbiome backgrounds is warranted to both adapt surveillance strategies in
vulnerable patient populations and to improve our fundamental understanding of *C. difficile* pathogenesis across a greater representation
of clinically and epidemiologically significant pathogen genomic diversity.

We identified an inverse correlation between *Blautia* spp. and CDI in colonized patients and with *C. difficile* growth in our
microbiome engraftment screen. *Blautia* has previously been implicated in human and animal microbiome-association studies as
significantly altered in *C. difficile*-related disease outcomes[35-37], primarily associated with protection from disease. Consistent with such
findings, these taxa appear to blunt *V. cholera* proliferation through bile acid metabolism [38-41]. More recent genome-scale metabolic
modeling indicates that *Blautia* may compete with *C. difficile* for amino acid pools[42]. These studies suggest that in our microbiome
engraftment screen, *Blautia* spp. either alter germination rates of *C. difficile* or slow down vegetative growth by consuming overlapping
nutrient pools. Additionally, data from our clinical cohort indicates several Lachnospiraceae to be depleted in patients with recent
cephalosporin exposure, which was most indicative of CDI in colonized patients, underlining the clinical significance of these ecological
relationships.

Gnotobiotic animal models of *C. difficile*-commensal interactions have made critical progress in illuminating the mechanisms
by which commensals support *C. difficile* proliferation through carbon-source crossfeeding[12,18] or inhibit *C. difficile* disease through
competitive Stickland fermentation[12] and secondary bile acid generation[25]. These approaches relied on simple communities (n = 1-13
commensal species) to profile the transcriptional and metabolomic landscape of *C. difficile* colonization *in vivo*[12,18,25]. More recently, a
gnotobiotic model using three gut commensals revealed that *C. difficile* consumption of host nutrients decreased virulence[43]. Based on
these approaches, we hypothesized that we could model diverse human microbiome variation that would help explain differences in
CDI susceptibility among colonized patients and provide a controlled background for assessing the virulence of clinically prevalent *C.*
difficile. The defined culture collection approach allowed us to recapitulate *C. difficile* carriage, where gnotobiotic animals colonized
with DefCom2 attenuated the strain-specific differences in weight loss (among clade 1 and 2 isolates) seen with DefCom1. DefCom2's
distinct suppression of toxin production and toxin-mediated inflammation across diverse strains (from clade 1 and 2) indicates that, in a
given patient, certain microbiome structures that enable *C. difficile* colonization may overcome strain-specific virulence attributes.
Further prediction of pathway utilization in these different microbiomes reveals that DefCom2 is enriched in polysaccharide/sugar
degradation and glycosaminoglycan degradation. Given that the ST8 strain grew to higher abundance in the DefCom2 background, we
speculate that DefCom2 feeds *C. difficile* host- and/or diet-derived glycan degradation products and thus represses cellular stress and
subsequent toxin production. This hypothesis fits with our longstanding understanding of nutrient-mediated transcriptional repression
of toxin production[8,44,45] and is supported by recent data indicating that commensal competition may increase toxin production[42]. These
defined microbiome data in complex with quantitative measurements of strain virulence indicate that *C. difficile* is not exclusively an
acutely destructive gastrointestinal pathogen but can instead operate as a pathobiont capable of cooperating in a complex microbial
community.

We acknowledge the limitations of our study and highlight opportunities for future work to further examine how commensals

can support stable *C. difficile* colonization and suppress virulence. Our human cohorts were derived from a single healthcare setting in

the U.S., and while our observed toxigenic *C. difficile* colonization rates and the prevalence of clade 1 isolates[2] mirror other recent

reports in Michigan[3] and Massachusetts[7], it will be important to extend such analyses outside the U.S. and beyond the healthcare setting.

It is notable that the patients who served as donors for engraftment experiment did not have clinical CDI, yet a fraction of those donor

stools produced severe disease in mice, suggesting that not all species present in the donor engrafted into mice. While our defined

communities were rationally designed to be highly representative of microbiome states observed in stably colonized patients, they are

not inclusive of all taxa found in these communities, and the construction of additional consortia or augmentation of our existing

communities may reveal differences in resilience against CDI, such as the taxa that are most affected by antibiotic exposure. Although

we identify DefCom2 to attenuate disease from virulent clade 1 and clade 2 *C. difficile* strains compared to DefCom1, strain dropout

experiments[19] will be crucial for systematically assigning which taxa are necessary and sufficient for such resilience and in which

combinations, and, lastly, whether this protection extends to other clinical *C. difficile* lineages (such as the globally prevalent ST11 from

clade 5)[6].

Our work highlights the provocative potential for future anti-CDI therapies which could suppress toxin production while still

supporting asymptomatic *C. difficile* growth and colonization in the gut. If surveillance approaches continue to report stable colonization

as a defining feature of circulating *C. difficile* lineages, it will be important to better predict disease based on microbiome structure and

relevant antibiotic exposure. Furthermore, the development of novel microbiome therapeutics that specifically focus on suppressing

toxin production are warranted, with concomitant consideration of strain-specific features of the commensal microbiome that cooperate

with or antagonize pathogenic proliferation and epithelial destruction.

<u>Methods:</u>

Integration of human microbiome samples

Stool samples were collected from inpatients at Barnes-Jewish Hospital (BJH) in St. Louis, MO, USA in two cohorts (retrospective and

prospective)[2,46]. Stool specimens that were submitted for toxin enzyme immunoassay (EIA) testing were collected from patients in the

retrospective cohort who had clinically significant diarrhea and no alternate identifiable cause of diarrhea from 2014-2016 as previously

described[46]. Samples that were metagenomically sequenced to sufficient depth and had corresponding toxigenic *C. difficile* genome

assemblies were included here for analysis for a total of 48 primary CDI episode (EIA+) and 54 carriage (EIA-) samples[21]. Microbiome

and isolate assembly data were downloaded from BioProject accession numbers PRJNA748262 and PRJNA980715, respectively. The

prospective cohort was drawn from a surveillance study of patients admitted to the BJH hematopoietic stem cell transplant and leukemia

units[2] during a four and six month period, respectively, in 2019. Stool and/or rectal swab samples were collected at the time of admission

and weekly thereafter and were cultured as previously described[2]. As with the retrospective cohort, specimens submitted to the BJH

microbiology laboratory for EIA testing due to clinical suspicion of CDI were cultured for *C. difficile*. In the prospective cohort, carriers

were classified as being culture positive for toxigenic *C. difficile*, did not develop CDI within 30 days of stool collection date, had no

documented CDI episodes in the previous 8 weeks, and were EIA- in the case of suspected CDI. One patient-timepoint was flanked by

two positive toxigenic cultures and both *C. difficile* and the *tcdA/B* genes were detected in the stool metagenome, so the patient was also

classified as a carrier. Hospitalized control samples were within 7 days of a *C. difficile* culture-negative sample and did not become

culture-positive during hospitalization; furthermore, samples for which *C. difficile* was detected metagenomically were excluded.

Samples of patients with CDI (EIA+) were selected if they were stool specimens submitted for EIA testing (i.e., before CDI treatment

initiation) and represented a first episode of CDI (i.e., no documented CDI within the previous 8 weeks). Longitudinal samples, which

applied to a subset of carriers and hospitalized controls from the prospective cohort, were included as available for linear mixed effect

modeling (which controlled for patient ID, see below) and one sample per patient was randomly selected as the "primary" sample for

all other analyses (for one patient, two samples were included in the PCoA visualizations as they were both used as donor stools). For this cohort, *C. difficile* genome assemblies were downloaded from BioProject accession number PRJNA980715*;* a core-genome alignment was constructed using panaroo and fastree was used to construct a phylogenetic tree, as previously[47]. Antibiotic exposures in the 30 days prior to stool collection were encoded as binary variables and compared between patient groups using a Fisher's exact test for subclasses with at least 10% prevalence across both hospitalized cohorts. Healthy volunteer microbiome data was drawn from a previously published cohort of adult individuals in the same geographic area without recent prior antibiotic exposure[22]; sequencing data was downloaded from BioProject PRJNA664754.

Metagenomic sequencing and analysis

Metagenomic DNA was extracted from patient stool from the prospective cohort and mouse fecal matter (for all animal experiments) using the Qiagen PowerSoil Pro kit. Libraries were prepared using a modified Nextera protocol as previously[21] and sequenced as 2x150 paired-end reads on an Illumina NovaSeq 6000 or NextSeq 550 instrument. Reads were demultiplexed by their index pair and quality-trimmed using Trimmomatic[48] (v0.39, SLIDINGWINDOW: 4:20, LEADING: 10, TRAILING: 10, MINLEN:60) followed by decontamination of human or mouse genomic reads using DeconSeq[49](v4.3; -dbs hsref38 or mm_ref_s1 & mm_ref_s2, respectively). MetaPhlAn4 (mpa_vJan21_CHOCOPhlAnSGB_202103) database was used to profile taxonomic composition and lowly abundant taxa were removed by filtering out species calls below 0.1% relative abundance[50]. HUMAnN3 was used to profile community metabolic function and normalization of pathways was performed to relative abundance[50]. Antimicrobial resistance gene (ARG) profiling of human metagenomes was performed using shortBRED against a marker gene set built from the NCBI and CARD databases[23]. ARG burden was determined using reads mapping per kilobase per million. Rarefaction was performed to identify sequencing thresholds for human microbiomes or mouse microbiomes based on subsampling a random set of ten samples from nine million paired-end reads to 1000 reads using seqtk (https://github.com/lh3/seqtk). Subsampled metagenomic reads were profiled using MetaPhlAn4 to assess taxonomic diversity, and differences between species richness were assessed for significance using a Wilcoxon rank sum test. Correspondingly, samples for human analyses that accrued less than 2.5 million paired-end reads per metagenome were removed from further analyses. Samples for mouse analyses that accrued less that 2 million paired-end reads per metagenome were removed from further analyses. Pathways below 0.01% relative abundance or assigned as 'UNINTEGRATED' were removed. For comparison of DefCom1 and DefCom2 functional pathways, aggregation of pathways was performed using a MetaCyc "parent orthology" classification[51].

Random forest classifier

Random forest classifier analyses were carried out as described previously[52] in R with modifications using caret v6.0.94. The dataset consisted of all colonized patient samples with corresponding high-quality *C. difficile* genome assemblies and available race information for the participant; the "primary" sample was selected for a given patient (n = 128 [50 CDI, 78 carriers]). This dataset was randomly split into training (60%) or testing (40%) subsets (seed = 42). Taxonomic features were selected with Boruta (v8.0.0) using species relative abundances from the training data (after excluding *C. difficile*) and identifying taxa selected in at least 25 of 100 iterations (seeds 1:100). The six selected taxa were subjected to preprocessing using the caret "center" and "scale" transformations, while 30-day antibiotic exposures (>10% prevalence), demographic data, and *C. difficile* binary toxin (*cdtB*) presence or absence were encoded as categorical variables. Random forest classifiers were trained on 100 random subsets of 80% of the training cohort using a different seed in each instance (seeds 1:100) with 10-fold cross-validation in each case using the "Accuracy" metric. Each model was tested against the retained validation cohort and the most accurate model was used to summarize model performance. To determine feature importance, null models were trained on datasets with randomly shuffled class labels and the resulting feature importances were determined using the caret varImp function with scale = FALSE. The mean feature importance per variable was determined for each model tested and all empirical and null distributions were compared using Wilcoxon rank sum test and Bonferroni correction for multiple hypotheses.

Microbiome engraftment screen

Stools were selected from carriers in this study and used for fecal microbiota engraftment of germ-free mice under Washington University in St. Louis IRB #202203020. Gnotobiotic mouse experiments were performed using protocols approved by the Washington

University School of Medicine Institutional Animal Care and Use Committee (21-0160). Adult male and female germ-free mice were housed in Allentown SPP Sentry racks and fed Lab Diet 5K67 autoclavable chow with a 12 hour on, 12 hour off 6am – 6pm light cycle. Stools from donors HV, 1, 2, 3, 4, 5, 6, 7, 8, and HV were engrafted into equal numbers of male and female mice (N=8) over the course of two experiments. Stools from donors 10, 13, 14, 15, 16, 22, and 24 were engrafted into equal numbers of male and female mice (N=4) over the course of one experiment; one mouse in the donor 22 arm suffered a lethal cecal torsion before the experiment was initiated, thus decreasing the number of mice (N=3). To prepare stools for gavage or culturing, fecal matter was resuspended in PBS-30% glycerol-0.5% cysteine and filtered through a 100uM filter (Falcon) to remove debris. Filtered gavage material was stored at -80 degrees C until gavage. Mice were gavaged with 100uL of gavage material on two consecutive days. Following gavage, mice were monitored for weight-loss and clinical signs of disease (marked by hunched posture and slowed movement)[53,54]. Upon 15% weight-loss (relative to weight on the first day of gavage) and or severe clinical disease (moribund), mice were euthanized. Stool was collected at least once per week following engraftment, and each week following through 21 days. Across three separate experiments, a stool from the first week and a stool or cecal content sample from the third week was used to quantify *C. difficile* CFU per miligram (CFU/mg), via serial dilution onto *C. difficile* selective plates (Biomerieux), and TcdB levels per milligram of stool (TcdB/mg) using an ELISA kit (TCGbiomics). For the TcdB ELISAs, fecal samples in the microbiome engraftment screen were diluted 1:5 in dilution buffer, and manufacturer instructions were followed. To calculate ratios of CFU/mg or TcdB/mg for each mouse, the average level was determined across mice within each donor arm from the same experiment at an early timepoint (day 4 or 5) and the day 21 value was divided by this value. This was because some mice early after engraftment did not produce stool.

For the phylogenetic analyses of donor stool *C. difficile* isolates, a core-gene alignment between all 15 isolate genomes was generated using panaroo as above, and a tree was constructed using RAxML[55]. Average nucleotide identity (ANI) between closely related clade 1 isolate genomes was determined by running snp-dist(https://github.com/tseemann/snp-dists) with the panaroo core-gene alignment as input. ANI=(total number of basepairs in alignment- total number of SNPs)/ total number of basepairs in alignment.

Statistical analyses of microbiome data

Species alpha diversity (richness and Shannon diversity) and the ARG richness and burden (RPKM) were compared across human cohort groups using the Wilcoxon rank sum test with Bonferroni correction for multiple hypotheses. Species beta diversity (Bray-Curtis distance matrix computed using vegdist) was compared between groups by PERMANOVA using adonis2 with 9999 permutations. In the human microbiome data, MetaPhlAn4 features were used in MaAsLin2[52] to investigate differences between disease status or cephalosporin exposure using the formulas: Taxa~ Group+(1 | Participant_ID) for all colonized patients or Taxa~ ceph_30d+(1 | Participant_ID) for all hospitalized patients, respectively, with normalization=NONE and all other parameters set to default. Similarly, MetaPhlAn4 features (from all stool/cecal microbiomes from surviving mice, as filtered above) and HUMAnN3 features were used to understand microbiome correlates of *C. difficile* growth (CFU/mg) and toxin (TcdB/mg). The formula for the analyses in the microbiome engraftment screen was: Taxa/Pathways ~ Growth + Timepoint +Toxin+(1 | DonorID) + (1 | Cdiffstrain) + (1 | CageAZ) + (1 | Sex) + (1 | MouseID), with min_prevalence=0.2, normalization=NONE, transform=LOG, and all other options used in default settings. The formula for the analyese in the defined community experiments to investigate community differences in metabolic pathway composition at day 28 was: Aggregated Pathways ~ Community+ (1 | MouseID) + (1 | Cage) + (1 | Sex) + (1 | Strain). The q-value and coefficient were depicted in plots. Sparse estimates of correlations among microbiomes (SECOM) was used to investigate correlational relationships between species features[56]. Linear correlations of Metaphlan4 features was performed using secom_linear() with specifications: assay_name = "counts", tax_level = "Species", pseudo = 0,lib_cut=0, method = "pearson", soft = FALSE, thresh_len = 100, n_cv = 10, thresh_hard = 0, max_p = 0.005. P-value filtered correlations of interest were visualized using Cytoscape[57].

Defined community culturing

Stool gavage material used for animal experiments (from donors 4, 13, 15, and 16) was used to culture taxa representative of microbiomes of this patient population. For each donor, ethanol-treated stool[58] was plated on LYBHI[59] containing 0.1% sodium taurocholate, and non-ethanol-treated stool was plated on LYBHI in 10-fold dilutions. For ethanol-treated stool, stool was resuspended

in an equal volume of 100% ethanol and left to sit for 4 hours. Following this incubation, serial dilutions were plated onto agar plates.

Plates were incubated for 2-3 days; distinct colonies were subcultured into LYBHI in deep 96 well plates and allowed to grow for two

days. Well contents were identified using V3-V4 16S rRNA sequencing[60]. Candidate taxa were subcultured to a clonal morphology and

confirmed by 16S full-length Sanger sequencing via Genewiz. Purified clones were stored in LYBHI+25% glycerol at -80 degrees C.

Defined community mouse experiments

All isolates were inoculated onto LYBHI agar plates for 3 days to allow growth. A subset of isolates were inoculated onto LYBHI+0.5%

gastric mucin agar and grown in LYBHI+0.5% gastric mucin broth: both *Akkermansia muciniphila* strains, *Blautia hansenii, Alistipes*

putredinis, Anaerofustis stericorihominis, *Anaerotruncus colihominis.* A single colony was inoculated into a deep-well plate with 1mL

of LYBHI and diluted back 1:10 into fresh media every 24 hours for two days. Isolates were pooled at equivalent loads based on OD_{600},

to achieve ~10^9 CFU/mL in PBS. Germ-free adult (10-12 weeks of age) male and female C57BL-6J mice were gavaged once daily for

two days with each community. Mice were followed over time for health (weight and clinical signs of distress). At 28 days, mice were

gavaged with 10^4 CFU/mL of each *C. difficile* strain. Stool was also collected at Day 28 before infection and enumerated for total

CFU/mg on LYBHI with 0.5% sodium taurocholate after 2 days of growth. Mice were assessed for health daily following gavage for

seven days and euthanized as above. At days 2, 4, and 6, stool was collected and analyzed for *C. difficile* CFU/mg. Cecal content samples

were collected following euthanasia. Toxin concentration (TcdB) in cecal samples was detected using ELISA kit for 'Separate detection

of *Clostridium difficile* TcdA or TcdB' (tgcBIOMICS, Germany). Cecal samples were diluted 1:10 in dilution buffer and loaded onto

ELISA plates for detection of toxin, as per the manufacturer's instructions. TcdB levels were normalized to total protein in the

corresponding cecal sample estimated by Bradford assay and expressed as ng/mg protein. For calprotectin estimation, cecal samples

were diluted 1:5 in reagent diluent and quantified by ELISA using the DuoSet Mouse S100A8/S100A9 Heterodimer kit (R&D Systems,

Minneapolis, MN, USA) according to the manufacturer's recommendations. Calprotectin levels were normalized to total protein and

expressed as pg/mg protein.

Data analyses and visualization

All data distributions were depicted using RStudio or Prism and indicate the median with a line (whether in a box-and-whiskers plot or

alone). Boxplots display the 25th and 75th percentile in the boxed region and depict the maximum and minimum value with the whiskers.

Ellipses displayed on PCoA plots depict the 95% confidence interval for the specified group. All statistical tests were two-sided unless

otherwise mentioned in the figure legends. All distributions were assessed for normality using the Shapiro-Wilkes test and then

correspondingly tested for statistically significant differences using an experiment-specific parametric or non-parametric test.

Data availability:

Metagenomic data and its associated metadata is available at BioProject PRJNA1067813.

Code availability:

The random forest classifier code and accompanying antibiotic exposure information and metadata can be found at

https://github.com/dantaslab/2024_Fishbein_DeVeaux_Cdiff_DefCom_Microbiome.

Acknowledgements:

The authors are grateful for members of the Dantas lab for their helpful feedback on the data analysis and preparation of the manuscript,

with special thanks to Drew Schwartz and Mark Gorelik for their conceptual insights and Philip Frasse for assisting in sample

preparation. The authors would also like to thank the Edison Family Center for Genome Sciences and Systems Biology staff, Eric Martin,

Brian Koebbe, MariaLynn Crosby, and Jessica Hoisington-López for their expertise and support in sequencing/data analysis, Jacky

Theodore, Bonnie Dee, James Anderson, and Keith Page for administrative support, and Sondra Seiler for her contributions to the

prospective clinical study. Lastly, the authors express their gratitude towards Summer Rucknagel, J. Michael White, and Sian Madden from the Washington University School of Medicine Gnotobiotic Research, Education, and Transgenic Center for their valuable support in conducting the gnotobiotic mouse experiments described in this study. G.D. and ERD were supported in part by the National Institute of Allergy and Infectious Diseases (NIAID: https://www.niaid.nih.gov/) of the National Institutes of Health (NIH) under award number R01AI155893 (PI: G. Dantas) and an award from the Foundation for Barnes-Jewish Hospital and Institute of Clinical and Translational Sciences. This publication was also supported by the NIH/National Center for Advancing Translational Sciences (NCATS), grant UL1 TR002345 (PI: B. Evanoff). This work was also supported by funding through the CDC BAA #200-2018-02926 (PI: E. Dubberke). SRSF is supported by the National Institute of Child Health and Human Development (NICHD: https://www.nicdhd.nih/gov) of the NIH under award number T32 HD004010 (PI: P. Tarr) and through the NIAID of the NIH under award number K99AI17674 (PI: S. Fishbein). ALD is supported by the NIH under award number T32GM139774 (PI: H. True-Krob). The conclusions from this study represent those of the authors and do not represent positions of the funding agencies.

Author contributions:

SRSF, ALD, ALF, ERD, and GD conceived the study. SRSF, ALD, SK, ALF, JL, JN, MJW, and BM performed the gnotobiotic mouse work. SRSF, ALD, SK, JL, BM, and SL performed *in vitro* experiments and prepared sequencing data. SRSF, ALD, JL, WA, and SL prepared the defined communities with expert guidance from JG. SRSF, ALD, and WA performed computational analyses. SRSF and ALD wrote the manuscript with feedback from all authors. TH and KAR retrieved patient-associated metadata, provided patient stool samples, and cultured *C. difficile* isolates from primary human stools. All authors participated in reviewing and editing the manuscript. ERD and GD supervised the research.

Declaration of interests statement:

ERD has served as a consultant for Seres Therapeutics, Ferring Pharmaceuticals, AstraZeneca, Summit, Merck, Recursion, and Abbott and has received research support from Theriva Biologics, Ferring Pharmaceuticals, and Pfizer, all unrelated to this study. The authors declare no other competing interests.

References:

Balsells, E. *et al.* Global burden of Clostridium difficile infections: a systematic review and meta-analysis. *J Glob Health* **9**, 010407, doi:10.7189/jogh.09.010407 (2019).

Newcomer, E. P. *et al.* Genomic surveillance of Clostridioides difficile transmission and virulence in a healthcare setting. *medRxiv*, doi:10.1101/2023.09.26.23295023 (2023).

Miles-Jay, A. *et al.* Longitudinal genomic surveillance of carriage and transmission of Clostridioides difficile in an intensive care unit. *Nat Med*, doi:10.1038/s41591-023-02549-4 (2023).

Feuerstadt, P. *et al.* SER-109, an Oral Microbiome Therapy for Recurrent Clostridioides difficile Infection. *N Engl J Med* **386**, 220-229, doi:10.1056/NEJMoa2106516 (2022).

Best, E. L., Freeman, J. & Wilcox, M. H. Models for the study of Clostridium difficile infection. *Gut Microbes* **3**, 145-167, doi:10.4161/gmic.19526 (2012).

Knight, D. R. *et al.* Major genetic discontinuity and novel toxigenic species in Clostridioides difficile taxonomy. *Elife* **10**, doi:10.7554/eLife.64325 (2021).

Worley, J. *et al.* Genomic Determination of Relative Risks for Clostridioides difficile Infection From Asymptomatic Carriage in Intensive Care Unit Patients. *Clin Infect Dis* **73**, e1727-e1736, doi:10.1093/cid/ciaa894 (2021).

Powers, D. A., Jenior, M. L., Kolling, G. L. & Papin, J. A. Network analysis of toxin production in Clostridioides difficile identifies key metabolic dependencies. *PLoS Comput Biol* **19**, e1011076, doi:10.1371/journal.pcbi.1011076 (2023).

9 Schubert, A. M., Sinani, H. & Schloss, P. D. Antibiotic-Induced Alterations of the Murine Gut Microbiota
and Subsequent Effects on Colonization Resistance against Clostridium difficile. *mBio* **6**, e00974,
doi:10.1128/mBio.00974-15 (2015).
10 Benech, N. *et al.* Update on Microbiota-derived therapies for recurrent Clostridioides difficile infections,
2023. *Clin Microbiol Infect*, doi:10.1016/j.cmi.2023.12.007 (2023).
11 Theriot, C. M., Bowman, A. A. & Young, V. B. Antibiotic-Induced Alterations of the Gut Microbiota Alter
Secondary Bile Acid Production and Allow for Clostridium difficile Spore Germination and Outgrowth in
the Large Intestine. *mSphere* **1**, doi:10.1128/mSphere.00045-15 (2016).
12 Girinathan, B. P. *et al.* In vivo commensal control of Clostridioides difficile virulence. *Cell Host Microbe*
29, 1693-1708 e1697, doi:10.1016/j.chom.2021.09.007 (2021).
13 Aguirre, A. M. *et al.* Bile acid-independent protection against Clostridioides difficile infection. *PLoS*
Pathog **17**, e1010015, doi:10.1371/journal.ppat.1010015 (2021).
14 Theriot, C. M. *et al.* Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase
susceptibility to Clostridium difficile infection. *Nat Commun* **5**, 3114, doi:10.1038/ncomms4114 (2014).
15 Smith, A. B. *et al.* Liberation of host heme by Clostridioides difficile-mediated damage enhances
Enterococcus faecalis fitness during infection. *mBio*, e0165623, doi:10.1128/mbio.01656-23 (2023).
16 Smith, A. B. *et al.* Enterococci enhance Clostridioides difficile pathogenesis. *Nature* **611**, 780-786,
doi:10.1038/s41586-022-05438-x (2022).
17 Engevik, M. A. *et al.* Mucin-Degrading Microbes Release Monosaccharides That Chemoattract
Clostridioides difficile and Facilitate Colonization of the Human Intestinal Mucus Layer. *ACS Infect Dis* **7**,
1126-1142, doi:10.1021/acsinfecdis.0c00634 (2021).
18 Ng, K. M. *et al.* Microbiota-liberated host sugars facilitate post-antibiotic expansion of enteric pathogens.
Nature **502**, 96-99, doi:10.1038/nature12503 (2013).
19 Wang, M. *et al.* Strain dropouts reveal interactions that govern the metabolic output of the gut
microbiome. *Cell* **186**, 2839-2852 e2821, doi:10.1016/j.cell.2023.05.037 (2023).
20 Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut microbiome. *Cell* **185**,
3617-3636 e3619, doi:10.1016/j.cell.2022.08.003 (2022).
21 Fishbein, S. R. *et al.* Multi-omics investigation of Clostridioides difficile-colonized patients reveals
pathogen and commensal correlates of C. difficile pathogenesis. *Elife* **11**, doi:10.7554/eLife.72801 (2022).
22 Anthony, W. E. *et al.* Acute and persistent effects of commonly used antibiotics on the gut microbiome
and resistome in healthy adults. *Cell Rep* **39**, 110649, doi:10.1016/j.celrep.2022.110649 (2022).
23 Kaminski, J. *et al.* High-Specificity Targeted Functional Profiling in Microbial Communities with
ShortBRED. *PLoS Comput Biol* **11**, e1004557, doi:10.1371/journal.pcbi.1004557 (2015).
24 Chen, X. *et al.* A mouse model of Clostridium difficile-associated disease. *Gastroenterology* **135**, 1984-
1992, doi:10.1053/j.gastro.2008.09.002 (2008).
25 Studer, N. *et al.* Functional Intestinal Bile Acid 7alpha-Dehydroxylation by Clostridium scindens
Associated with Protection from Clostridium difficile Infection in a Gnotobiotic Mouse Model. *Front Cell*
Infect Microbiol **6**, 191, doi:10.3389/fcimb.2016.00191 (2016).
26 Zackular, J. P. *et al.* Dietary zinc alters the microbiota and decreases resistance to Clostridium difficile
infection. *Nat Med* **22**, 1330-1334, doi:10.1038/nm.4174 (2016).
27 Sorbara, M. T. *et al.* Functional and Genomic Variation between Human-Derived Isolates of
Lachnospiraceae Reveals Inter- and Intra-Species Diversity. *Cell Host Microbe* **28**, 134-146 e134,
doi:10.1016/j.chom.2020.05.005 (2020).
28 Janezic, S. & Rupnik, M. Genomic diversity of Clostridium difficile strains. *Res Microbiol* **166**, 353-360,
doi:10.1016/j.resmic.2015.02.002 (2015).
29 Sebaihia, M. *et al.* The multidrug-resistant human pathogen Clostridium difficile has a highly mobile,
mosaic genome. *Nat Genet* **38**, 779-786, doi:10.1038/ng1830 (2006).
30 Edwards, A. N., Krall, E. G. & McBride, S. M. Strain-Dependent RstA Regulation of Clostridioides difficile
Toxin Production and Sporulation. *J Bacteriol* **202**, doi:10.1128/JB.00586-19 (2020).

31 Sun, X. *et al.* Mouse relapse model of Clostridium difficile infection. *Infect Immun* **79**, 2856-2864,
doi:10.1128/IAI.01336-10 (2011).
32 Theriot, C. M. *et al.* Cefoperazone-treated mice as an experimental platform to assess differential
virulence of Clostridium difficile strains. *Gut Microbes* **2**, 326-334, doi:10.4161/gmic.19142 (2011).
33 Kong, K. Y. *et al.* Biological characteristics associated with virulence in Clostridioides difficile ribotype 002
in Hong Kong. *Emerg Microbes Infect* **9**, 631-638, doi:10.1080/22221751.2020.1739564 (2020).
34 Griffiths, D. *et al.* Multilocus sequence typing of Clostridium difficile. *J Clin Microbiol* **48**, 770-778,
doi:10.1128/JCM.01796-09 (2010).
35 Barron, M. R. *et al.* Intestinal Inflammation Reversibly Alters the Microbiota to Drive Susceptibility to
Clostridioides difficile Colonization in a Mouse Model of Colitis. *mBio* **13**, e0190422,
doi:10.1128/mbio.01904-22 (2022).
36 Berkell, M. *et al.* Microbiota-based markers predictive of development of Clostridioides difficile infection.
Nat Commun **12**, 2241, doi:10.1038/s41467-021-22302-0 (2021).
37 Lesniak, N. A. *et al.* The Gut Bacterial Community Potentiates Clostridioides difficile Infection Severity.
mBio **13**, e0118322, doi:10.1128/mbio.01183-22 (2022).
38 Reeves, A. E., Koenigsknecht, M. J., Bergin, I. L. & Young, V. B. Suppression of Clostridium difficile in the
gastrointestinal tracts of germfree mice inoculated with a murine isolate from the family
Lachnospiraceae. *Infect Immun* **80**, 3786-3794, doi:10.1128/IAI.00647-12 (2012).
39 Reed, A. D., Nethery, M. A., Stewart, A., Barrangou, R. & Theriot, C. M. Strain-Dependent Inhibition of
Clostridioides difficile by Commensal Clostridia Carrying the Bile Acid-Inducible (bai) Operon. *J Bacteriol*
202, doi:10.1128/JB.00039-20 (2020).
40 Buffie, C. G. *et al.* Precision microbiome reconstitution restores bile acid mediated resistance to
Clostridium difficile. *Nature* **517**, 205-208, doi:10.1038/nature13828 (2015).
41 Reed, A. D. & Theriot, C. M. Contribution of Inhibitory Metabolites and Competition for Nutrients to
Colonization Resistance against Clostridioides difficile by Commensal Clostridium. *Microorganisms* **9**,
doi:10.3390/microorganisms9020371 (2021).
42 Jenior, M. L. *et al.* Systems-ecology designed bacterial consortium protects from severe Clostridioides
difficile infection. *bioRxiv*, doi:10.1101/2023.08.08.552483 (2023).
43 Pruss, K. M. *et al.* Oxidative ornithine metabolism supports non-inflammatory C. difficile colonization.
Nat Metab **4**, 19-28, doi:10.1038/s42255-021-00506-4 (2022).
44 Bouillaut, L., Dubois, T., Sonenshein, A. L. & Dupuy, B. Integration of metabolism and virulence in
Clostridium difficile. *Res Microbiol* **166**, 375-383, doi:10.1016/j.resmic.2014.10.002 (2015).
45 Ransom, E. M., Kaus, G. M., Tran, P. M., Ellermeier, C. D. & Weiss, D. S. Multiple factors contribute to
bimodal toxin gene expression in Clostridioides (Clostridium) difficile. *Mol Microbiol* **110**, 533-549,
doi:10.1111/mmi.14107 (2018).
46 Dubberke, E. R. *et al.* Clostridium difficile colonization among patients with clinically significant diarrhea
and no identifiable cause of diarrhea. *Infect Control Hosp Epidemiol* **39**, 1330-1333,
doi:10.1017/ice.2018.225 (2018).
47 Newcomer, E. P., Fishbein, S. R. S. ,Zhang, K.,Hink, T.,Reske, K.T.,Cass, C.,Iqbal, Z.H., Struttmann, E.L.,
Dubberke, E.R.,Dantas, G. Genomic surveillance of Clostridioides difficile transmission and virulence in a
healthcare setting. *medRxiv*, doi:https://doi.org/10.1101/2023.09.26.23295023 (2023).
48 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).
49 Schmieder, R. & Edwards, R. Fast identification and removal of sequence contamination from genomic
and metagenomic datasets. *PLoS One* **6**, e17288, doi:10.1371/journal.pone.0017288 (2011).
50 Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse microbial
communities with bioBakery 3. *Elife* **10**, doi:10.7554/eLife.65088 (2021).
51 Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes - a 2019 update. *Nucleic Acids*
Res **48**, D445-D453, doi:10.1093/nar/gkz862 (2020).

52 Ferreiro, A. L. *et al.* Gut microbiome composition may be an indicator of preclinical Alzheimer's disease.
Sci Transl Med **15**, eabo2984, doi:10.1126/scitranslmed.abo2984 (2023).
53 Aston, W. J. *et al.* A systematic investigation of the maximum tolerated dose of cytotoxic chemotherapy
with and without supportive care in mice. *BMC Cancer* **17**, 684, doi:10.1186/s12885-017-3677-7 (2017).
54 Fletcher, J. R. *et al.* Clostridioides difficile exploits toxin-mediated inflammation to alter the host
nutritional landscape and exclude competitors from the gut microbiota. *Nat Commun* **12**, 462,
doi:10.1038/s41467-020-20746-4 (2021).
55 Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).
56 Lin, H., Eggesbo, M. & Peddada, S. D. Linear and nonlinear correlation estimators unveil undescribed taxa
interactions in microbiome data. *Nat Commun* **13**, 4946, doi:10.1038/s41467-022-32243-x (2022).
57 Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction
networks. *Genome Res* **13**, 2498-2504, doi:10.1101/gr.1239303 (2003).
58 Browne, H. P. *et al.* Culturing of 'unculturable' human microbiota reveals novel taxa and extensive
sporulation. *Nature* **533**, 543-546, doi:10.1038/nature17645 (2016).
59 Beller, Z. W. *et al.* Inducible CRISPR-targeted "knockdown" of human gut Bacteroides in gnotobiotic mice
discloses glycan utilization strategies. *Proc Natl Acad Sci U S A* **120**, e2311422120,
doi:10.1073/pnas.2311422120 (2023).
60 Vernier, C. L. *et al.* The gut microbiome defines social group membership in honey bee colonies. *Sci Adv*
6, doi:10.1126/sciadv.abd3431 (2020).
601
602

Figures and Tables

604
Figure 1- Microbiological variations underlie *Clostridioides difficile* colonization outcomes in a tertiary care hospital. A) Sampling
scheme for human microbiome analysis including multiple hospitalized cohorts and a group of healthy volunteers within the same
geographic area. B) Taxonomic richness of stool metagenomes (N = 53 CDI, 82 carrier, 32 HC, 20 HV) provided at the species level
using MetaPhlAn4. Statistical tests performed using the Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons.
C) Phylogenetic tree (mid-point rooted) built from core gene alignment of our complete collection of *C. difficile* clinical isolates collected
at Barnes-Jewish hospital. Inner ring corresponds to *C. difficile* clade (C1 = cryptic clade 1). Outer ring denotes isolates selected for this
study and whether they derived from *C. difficile* carriers or patients with CDI. Isolates are labeled by donor ID for samples selected as
microbiome engraftment donors in the following section and their lethality phenotype *in vivo*. D) Principal coordinate analysis of pan-
genome alignment of selected *C. difficile* isolates plotted by Jaccard distance based on their pairwise average nucleotide identity (ANI).

Each point corresponds to an individual genome and is colored by clade and filled based on CDI status. Percent variance explained is
displayed parenthetically for the two axes shown. (C-D) depict data from 248 bacterial isolates. E) Mean feature importance plot based
on a random forest classifier built to predict CDI status among colonized (carrier or CDI) samples using all features shown. The black
and maroon points and standard deviation represent predictive importance of the empirical and null model, respectively. Features in bold
are significantly more predictive than the corresponding null distribution ($p < 0.05$, Wilcoxon-rank sum test with Bonferroni correction
for multiple comparisons). Color strip denotes feature type as referenced in the legend.

630   with recent cephalosporin exposure in colonized patients as carried out in E. HC, hospitalized control; HV, healthy volunteer. For A-D,
631   N = 53 CDI, 82 carrier, 32 HC, 20 HV.
632

Antibiotic exposures in the previous 30 days by class				
	HC	Carrier	CDI	*p*-value
n	32	82	53	
Any antibiotics (%)	16 (50.0)	18 (22.0)	21 (39.6)	**0.006**
Intravenous vancomycin (%)	8 (25.0)	10 (12.2)	21 (39.6)	**0.001**
Cephalosporin (%)	3 (9.4)	6 (7.3)	29 (54.7)	**<0.001**
Carbapenem (%)	4 (12.5)	6 (7.3)	8 (15.1)	0.363
Fluoroquinolone (%)	6 (18.8)	5 (6.1)	11 (20.8)	**0.023**

Specific categories refer to classes with ≥10% prevalence. *p*-values determined by Fisher's exact test.

634
635 **Supplementary Table 1 – Antibiotic exposures in the previous 30 days for hospitalized patient groups.** *P*-values determined across
636 all three groups with Fisher's exact test ($p \leq 0.05$ bolded). HC, hospital control.
637

Figure 2- **Microbiome engraftment screen recapitulates the spectrum of human *C. difficile*-associated disease**. A) Humanization of germ-free mice with carrier patient microbiomes allows observation of pathogen-microbiome dynamics. B) Kaplan-Meyer plot of microbiome humanization, where each color reflects a stool (Donor) that caused >0% lethality in mice (*n* = 3-8 mice per group, see Methods). C) Principal coordinate analysis of Bray-Curtis dissimilarity across human microbiomes (●), where donors used in the screen are indicated as lethal (▲) if any mice died or nonlethal (■). (■) (*n* = 53 CDI, 83 carrier, 32 HC, 20 HV). D) Relationship between microbiome community structure and donor stool *C. difficile* abundance (CFU/mL), as measured by distance from HV communities in C. *R*-squared value and p-value from a Pearson correlation test. E) Richness of microbiomes in surviving mice at day 21 (N indicates number of mouse stool or fecal microbiome quantified for each donor), as measured by MetaPhlAn4 species. F) Midpoint-rooted phylogenetic tree built from core gene alignment of 15 *C. difficile* isolates associated with donor stools. Boxes indicate a relatedness of >99.9% ANI; grey boxes indicate isolates associated with non-lethal donors while red boxes indicate >0% lethality associated with donor stools.

650
Supplementary Figure 2 - Donor-specific features of microbiome engraftment A) Minimum weight loss across 21 days for each
stool microbiome. Significant statistical differences for each donor relative to the control microbiome (Donor HV) as measured by
multiple t-tests against Donor HV as a reference, with p-value adjustment performed by Bonferroni correction. B) Relationship between
percentage of mice surviving over 21 days and *C. difficile* concentration in the inoculum. Boxplots of growth (C) and toxin (D) ratio of
day 21 *C. difficile* levels relative to day 4-5 levels for each stool microbiome engrafted. E) Linear coefficients of MetaCyc pathways

(HUMAnN3-produced) associated with increased *C. difficile* growth, q-value <0.1, as described in the Methods via MaAsLin2 (analyzed
88 stool metagenomes over time across 13 donor backgrounds)
658

Figure 3- Strain variation and microbiome structure drive differences in colonization outcome *in vivo*. *C. difficile* strain groups display significant differences in A) weight-loss, B) titer ratios, and C) toxin/mg ratios. P-values indicate Benjamini-Hochberg corrected results of a Dunn's test comparison between mouse measurements across different strain groups, and *n* indicates the number of mice from which ratios was calculated. D) Principal coordinate analyses of Bray-Curtis dissimilarity between all fecal microbiomes of surviving mice; each point represents a stool microbiome from one mouse timepoint at day 21. E) Relationship between microbiome structure (PCoA from [D]) and stool *C. difficile* (top) and toxin levels (bottom). *R* value and p-value indicate the results of a Pearson correlation test. F) Linear coefficients of taxa related to growth or toxin levels in mice with a q-value <0.1 using MaAsLin2 as described in the Methods (analyzed 88 stool metagenomes over time across 13 donor backgrounds).

Figure 4-Construction of defined human gut microbiota in gnotobiotic models recapitulates spectrum of *C. difficile*-associated disease. A) Prevalence and abundance of genera in patients colonized with *C. difficile* from Figure 1, where color indicates use in

DefCom1 or DefCom2. B) Metagenomic abundance of species at >0.1% relative abundance in DefCom1 and DefCom2 (representative
of average abundance across DefCom1 [N=43 mouse stool metagenomes] and DefCom2 [N=52 mouse stool metagenomes] at day 28
in all experiments). C) Weight-loss after infection with *C. difficile* isolates following engraftment in DefCom1 (top, N=8 mice in each
arm) and DefCom2 (bottom, N=8 mice in each arm). D) Minimum weight-loss in the first week after engraftment (from Figure 2, N= 8
mice per donor arm) or infection from (C). P-values indicate Kruskal-Wallis test comparison of weight-loss within a single microbiome
background. Weight-loss percentage following infection with least virulent (ST2) and most virulent (ST8) clade 1 isolates, repeat of (C);
N=6 mice per arm. F) Cecal levels of *C. difficile* toxin and calprotectin levels across *C. difficile* strains and microbiome backgrounds at
day 7 post-infection from (E). P-values indicate Wilcoxon rank-sum test performed across different microbiome backgrounds for the
same *C. difficile* strain. For (C) and (E), one-way ANOVAs were performed with Tukey's multiple comparisons test with a single pooled
adjustment.
684
685

Supplementary Figure 3- DefCom2 supports less virulent *C. difficile* proliferation
A) Principal coordinate analysis of Bray-Curtis dissimilarity between taxonomic features (aggregated to genus level) from human samples (small points) with representative DefCom1 and DefCom2 microbiomes (large points, averaged across all mice at day 28 [N=43 mice from DefCom1, N=52 mice from DefCom2]). B) Principal coordinate analysis of Bray-Curtis dissimilarity between mouse stool microbiomes from (A) at day 28 during each defined community experiment. C) Quantification of beta diversity (Bray Curtis dissimilarity) differences between different communities (Between DefCom), between experimental replicates within the same DefCom (Between Replicate), and between microbiomes from the same defined community across experimental replicates (Within DefCom). P-values indicated a Wilcoxon rank sum test performed comparing Between DefCom group to other groups. D) Stool levels of *C. difficile* post-infection (left, from Figure 4E) with area under the curve (AUC, right panel), measured across all three time points for ST8 infection in DefCom1 (N=5 mice) and DefCom2 (N=6). **, p<0.01, as obtained by Mann-Whitney test. E) Stool levels of commensal bacteria in each defined community at day 28 of second experiment (N= before *C. difficile* infections. F) Linear coefficients of MetaCyc-aggregated pathway differences between DefCom1 and DefCom2 microbiomes, q-value<1e-15 are depicted.

700
701 **Supplementary Table 2- Species used in each defined community.**

DefCom1		DefCom2			
Species	Source	Species	Source	Species	Source
Akkermansia muciniphila	ATCC (BAA-835)	*Akkermansia muciniphila*	Donor 15	*Bifidobacterium pseudocatenulatum*	Donor 10
Alistipes putredinis	Donor 4	***Alistipes putredinis***	Donor 4	*Blautia hansenii*	Donor 10
Bacteroides caccae	Donor 4	***Bacteroides caccae***	Donor 4	*Blautia obeum*	Donor 15
Bacteroides fragilis	Donor 4	***Bacteroides fragilis***	Donor 4	*Desulfovibrio desulfuricans*	Donor 15
Bacteroides ovatus	Donor 4	***Bacteroides ovatus***	Donor 4	*Emergencia timonensis*	Donor 15
Bacteroides uniformis	Donor 4	***Bacteroides uniformis***	Donor 4	*Enterocloster clostridioformis*	Donor 13
Bacteroides xylanisolvens	Donor 4	***Bacteroides xylanisolvens***	Donor 4	*Enterococcus faecalis*	Donor 13
Bifidobacterium longum	Donor 4	***Bifidobacterium longum***	Donor 4	*Enterococcus faecium*	Donor 13
Blautia coccoides	Donor 4	***Blautia coccoides***	Donor 4	*Faecalimonas umbilicata*	Donor 13
Clostridium innocuum	Donor 4	***Clostridium innocuum***	Donor 4	*Hungatella effluvii*	Donor 10
Clostridium perfringens	Donor 4	***Clostridium perfringens***	Donor 4	*Klebsiella pneumoniae*	Donor 13
Clostridium symbiosum	Donor 4	***Clostridium symbiosum***	Donor 4	*Lacticaseibacillus paracasei*	Donor 15
Eggerthella lenta	Donor 4	***Eggerthella lenta***	Donor 4	*Limosilactobacillus fermentum*	Donor 15
Eisenbergiella tayi	Donor 4	***Eisenbergiella tayi***	Donor 4	*Parabacteroides distasonis*	Donor 15
Enterocloster lavalensis	Donor 4	***Enterocloster lavalensis***	Donor 4	*Ruminococcus gnavus*	Donor 13
Erysipelatoclostridium ramosum	Donor 4	***Erysipelatoclostridium ramosum***	Donor 4	*Sellimonas intestinalis*	Donor 10
Escherichia coli	Donor 4	***Escherichia coli***	Donor 4	*Streptococcus anginosus*	Donor 10
Eubacterium callanderi	Donor 4	***Eubacterium callanderi***	Donor 4	*Streptococcus caecimuris*	Donor 15
Flavonifractor plautii	Donor 4	*Flavonifractor plautii*	ATCC (265)	*Streptococcus mutans*	Donor 10
Phocaeicola vulgatus	Donor 4	***Phocaeicola vulgatus***	Donor 4	*Sutterella wadsworthensis*	Donor 10
Ruminococcus gauvreauii	Donor 4	***Ruminococcus gauvreauii***	Donor 4		
Ruthenibacterium lactatiformans	Donor 4	***Ruthenibacterium lactatiformans***	Donor 4		
Clostridium butyricum	Donor 4	***Anaerofustis stercorihominis***	Donor 4		
Collinsella aerofaciens	Donor 4	*Anaerotruncus colihominis*	Donor 4		
Anaerofustis stercorihominis	Donor 4	*Bacteroides thetaiotaomicron*	Donor 10		
Enterocloster aldenesis	Donor 4	*Barnesiella intestinihominis*	Donor 10		
Note: Bolded species represent isolates used in both communities					

702