1   **Appearance of green tea compounds in plasma following green tea consumption**
2   **is modulated by the gut microbiome in mice**
3

John D. Sterrett,[1,2,#] Kevin D. Quinn,[3] Katrina A. Doenges,[3] Nichole M. Nusbacher,[4] Cassandra
L. Levens,[5] Mike L. Armstrong,[3] Richard M. Reisdorph,[3] Harry Smith,[3] Laura M. Saba,[3] Kristine
A. Kuhn,[5] Catherine A. Lozupone,[4] Nichole A. Reisdorph[3]
1. Department of Integrative Physiology, University of Colorado, Boulder, CO
2. Interdisciplinary Quantitative Biology, University of Colorado, Boulder, CO
3. Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado,
10   Aurora, CO
11   4. Department of Biomedical Informatics, Anschutz Medical Campus, University of
12   Colorado, Aurora, CO
13   5. Division of Rheumatology, Department of Medicine, University of Colorado, Aurora, CO
14
15   #: Address correspondence to John Sterrett, john.sterrett@colorado.edu

16
17   Running Head: Gut microbiome affects green tea compounds in plasma
18

Abstract

Studies have suggested that phytochemicals in green tea have systemic anti-inflammatory and
neuroprotective effects. However, the mechanisms behind these effects are poorly understood,
possibly due to differential metabolism of phytochemicals due to varying gut microbiome
composition. To unravel this complex relationship, our team utilized a novel combined
microbiome analysis and metabolomics approach applied to low diversity, pseudo-germ-free
(GF) and human colonized (HU) gnotobiotic mice treated with an acute dose of powdered
matcha green tea. A total of 20 GF mice received 10 distinct human fecal slurries, for an n=2
mice per human gut microbiome; 9 mice remained GF throughout the experiment. We used
untargeted metabolomics on green tea and plasma to identify green tea compounds that were
found in GF and HU mouse plasma following acute dosing of green tea. 16s ribosomal RNA
gene sequencing was performed on feces of all mice at study end to assess microbiome
composition. We found multiple green tea compounds in plasma associated with microbiome
presence and diversity; we also detected strong associations between green tea compounds in
plasma and specific taxa in the gut. Many of these compounds are bioactive in the host,
including spiramycin and lactiflorin. Additionally, some of the physiologically relevant green tea
compounds are likely derived from plant-associated microbes, highlighting the importance of
considering foods and food products as meta-organisms. Overall, we describe a novel workflow
for discovering relationships between individual food compounds and composition of the gut
microbiome.

Importance
Foods contain thousands of unique and biologically important compounds beyond the main
nutrients often considered, such as those listed on nutrition facts labels. In mammals, many of
these compounds are altered or broken down by the community of microbes in the colon. This
means that these microbes impact the thousands of biologically important compounds we
consume, and that understanding microbial breakdown of food-derived compounds will be
important for understanding how foods impact health. We used metabolomics to track green tea

compounds in plasma of mice with and without complex microbiomes. From this, we can start to
recognize certain groups of green tea-derived compounds that are impacted by mammalian
microbiomes. This research presents a novel technique for understanding microbial metabolism
of food-derived compounds in the gut, which can be applied to other health-relevant foods.

Introduction
Assessing the impact of food on health is challenging.[1] This is exacerbated by the unique co-
metabolism of foods by the host and the gut microbiome across individuals with distinct
microbiota. Owing to recent advances in 'omics technologies, determining the identity of
microbial and microbial:host metabolites following consumption of specific foods is now
possible.[1–3] For example, Wang, *et. al.* found that a product of microbiome:host co-metabolism
of phosphatidylcholine, trimethylamine N-oxide (TMAO), predicted risk for cardiovascular
disease (CVD).[4] In addition, integrated metabolomics and microbiome approaches are being
applied with promising results.[5–12] However, challenges remain, including identifying metabolites
that are specifically produced by the microbiome or by host:microbiome interactions. Our
approach addresses this challenge through use of human colonized (HU) mice fed a food with
known health effects, namely green tea (GT). It aims to determine how the gut microbiome
affects which GT compounds are found in the plasma of HU and germ free (GF) mice.

Alteration of the gut microbiome has been associated with the development of multiple
disorders,[13] including depression and anxiety,[14] metabolic syndrome,[15] and inflammation.[16] A
broad understanding of how the microbiome affects the host metabolome has come from
comparing GF mice to those colonized with the microbiota of various humans[17] and/or treated
with antibiotics compared to non-antibiotic treated controls.[17–19] These studies found that
microbial composition has a profound influence on the presence and relative abundances of
many metabolites in various sites including the blood, urine, feces, and the gastrointestinal tract
(GIT). Additionally, dietary differences clearly influence microbial community structure, as
certain substrates favor specific taxa. For example, Wu *et al.*[20] reported greater relative
abundance of *Bacteroidetes* with high animal protein consumption and greater relative
abundance of *Prevotella* with plant consumption.[21] Together, these findings suggest that unique
microbiome profiles may be bidirectionally linked to specific dietary components.

Several studies have suggested that various components of tea, including flavonoids and
phenolic acids, have both anti-inflammatory and neuroprotective effects.[22–25] Polyphenol
metabolism, which is conducted through a combination of the host and microbiome, is
exquisitely complex.[26] However, the exact mechanism by which GT can affect specific
molecular networks is not yet understood. To our knowledge, only a few studies have focused
on the metabolism of tea in the context of the microbiome. For example, Axling *et. al.* found that
supplementing mice with GT powder along with *Lactobacillus plantarum* promoted growth of
Lactobacillus and attenuated high fat diet-induced inflammation.[27] Studies utilizing humanized
mice is a logical next step for determining the metabolites responsible for these effects and
which microbes might be involved in their metabolism.

These scientific premises resulted in the formation of our **hypotheses** *that specific GT*
compounds will associate with specific bacterial genera or be altered in concentration by the

presence of a gut microbiome. We tested these hypotheses by colonizing gnotobiotic mice with
microbiomes from 10 healthy humans and then treating both colonized and GF mice with an
acute dose of GT extract by oral gavage (**Figure 1**). The chemical composition of GT and
plasma from mice fed GT was determined using mass spectrometry-based metabolomics.
Mouse plasma metabolomics data was mined for GT compounds and associations between GT
compounds and microbiome composition and diversity were determined.

Because diet is among the most significant modifiers of human health, detailed understanding of
the microbiome in the context of food metabolism is critical for disease modification and
prevention.

99
Methods
Chemicals, standards, and reagents: Chemicals, standards and reagents used for sample
preparation and liquid chromatography-mass spectrometry (LCMS) analysis were of high-
performance liquid chromatography (HPLC) or LCMS grade. These included water from
Honeywell Burdick & Jackson (Muskegon, MI, USA), methanol, and methyl tert-butyl ether
(MTBE) from VWR (Radnor, PA, USA), formic acid from ThermoFisher Scientific (Waltham, MA,
USA), acetonitrile from Fisher Scientific (Fair Lawn, NJ, USA), and 2-Propanol from Millipore
Sigma (Burlington, MA, USA). Authentic standards for sample preparation were from Avanti
Polar Lipids Inc. (Alabaster, AL, USA), Cambridge Isotope Laboratories (Tewksbury, MA, USA),
Sigma-Aldrich (St. Louis, MO, USA) and CDN Isotopes (Pointe-Claire, Quebec, Canada).
110
111 *Human fecal sample collection:* Human fecal samples were collected from 10 healthy
112 participants using a commode specimen collector. Fecal samples were then shipped cold within
113 48 hours, aliquoted and stored at -80 °C prior to use in gnotobiotic mouse experiments.
114 Collection of the human fecal samples used in these experiments was approved by the
115 Colorado Multiple Institutional Review Board (COMIRB) COMIRB #14-1595.
116
Mouse Colonization: All mouse studies were approved by the University of Colorado Institutional
Animal Care and Use Committee (IACUC). C57BL/6 GF mice from the colony maintained at the
University of Colorado Gnotobiotics Core were placed on an irradiated, low polyphenol diet
(TD.97184, Envigo, Indianapolis, IN) at the time of weaning (3 weeks of age) through 6 weeks
of age. At 4 weeks of age, mice were orally gavaged with 200 µL of either human fecal slurry
(100 mg stool homogenized in 1 mL reduced PBS in an anaerobic chamber) or PBS for GF
controls (n=9). A total of 20 mice received 10 distinct human fecal slurries, for an n=2 mice per
human gut microbiome. Due to limited availability of GF mice, the experiment was conducted in
three cohorts as described in the Supplemental Methods. Mice were housed individually in a
sterile caging system to maintain their GF or individualized colonization status for the remainder
of the experiment. Mice were weighed three times weekly and on the day of GT dosing.
128
Mouse GT Gavage: Gavage solutions of GT were prepared under sterile conditions in a Nuair
Biological Safety Cabinet (Plymouth, MN). LCMS water was heated to 70 °C and sterile filtered
through a Steriflip disposable vacuum filtration tube (Merck KgaA, Darmstadt, Germany). Jade
Leaf matcha green tea (Seattle, WA) was irradiated by Envigo at the same time as the low

polyphenol diet and stored in a sterile 50 mL Falcon Tube (Corning, Glendale, AZ) at 4 °C. GT
slurries were prepared at 10 mg/mL with the warm sterile water, vortexed for 15 seconds to mix
and then stored at 4 °C until gavage. A total of 29 mice (20 HU, 9 GF) were gavaged with 100
µL of GT slurry (50 mg/kg equivalent). Leftover slurries were stored at -80 °C for GT
metabolomics analysis.
138
Plasma and Tissue Collection: Mice were sacrificed approximately 2 hours after the GT gavage.
Blood was collected in 1.3 mL K3 EDTA micro sample tubes (Sarstedt Inc., Nuembrecht,
Germany) via submandibular bleeding, inverted 5 times and immediately placed on ice. Blood
was centrifuged at 3,000 xg for 30 mins at 4 °C, within 30 minutes of collection. Plasma was
aliquoted into 1.5 mL microcentrifuge tubes (Fisher Scientific) and stored at -80 °C until
analysis. Fecal microbiome samples were collected in 1.5 mL microcentrifuge tubes, flash
frozen in liquid nitrogen and stored at -80 °C until analysis. The 2 hour time point was selected
following a small pilot utilizing wild type (WT) mice whereby GT compounds were detected in
higher numbers and abundance compared to 4 and 24 hour timepoints (Supplemental
Methods).
149
Metabolomics: Frozen plasma samples and four GT extracts were thawed on ice and prepared
for analyses as previously described.[28–30] Briefly, proteins were precipitated and small
molecules were extracted from supernatants using an organic liquid-liquid extraction technique
with methyl tert-butyl ether (MTBE) and water. This resulted in hydrophilic (aqueous) and
hydrophobic (lipid) fractions. Nine hydrophobic and 6 hydrophilic labeled spike-in standards
were used to monitor instrument performance and sample preparation batch variability.[31] Spiked
and un-spiked methanol preparatory blanks were prepared alongside study samples. Quality
control (QC) included analysis of purchased plasma (BioIVT, Westbury, NY), spiked with
authentic standards and prepared alongside study plasma samples (Supplemental Methods).
159
Aqueous and lipid fractions were analyzed by LCMS using published methods.[29,31,32] Briefly, the
hydrophobic fraction was analyzed using reverse phase C18 chromatography and a quadrupole
time-of-flight mass spectrometer (QTOF 6545, Agilent Technologies, Santa Clara, CA) in
positive ionization mode. The hydrophilic fraction was analyzed using an SB-Aq column (Agilent
Technologies) on the same instrument. LCMS method details are provided in the Supplemental
Methods.
166
Metabolomics Data Processing: Metabolomics spectral data for both plasma and GT extracts
were processed using a recursive workflow and area-to-height conversion with Agilent's
MassHunter Profinder ver. 10.0 service pack 1 (Profinder) and Mass Profiler Professional Ver.
15.1 (MPP, Agilent Technologies).[10] Data from lipid and aqueous fractions were extracted
separately in Profinder using Batch Molecular Feature Extraction followed by Batch Targeted
Feature Extraction. Compounds found in blanks were removed. Plasma samples were limited to
compounds eluting before 10.4 min and 13 min for lipid and aqueous fractions, respectively. GT
samples were limited to compounds eluting before 10.4 min and 11 min for lipid and aqueous
fractions, respectively. Compounds eluting past these times had poor signal to noise ratios and
were not used.

Plasma compounds were annotated using MassHunter ID Browser ver. 10.0 (Agilent
Technologies) by searching custom in-house databases and public databases consisting of
compounds from METLIN, Lipid Maps, Kyoto Encyclopedia of Genes and Genomes (KEGG),
and Human Metabolome Database (HMDB) using accurate mass and isotope ratios. An initial
database search was conducted using H+ as the primary charge carrier. Unannotated
compounds were re-searched using the same databases with loss of water included. These
compounds were designated Metabolomics Standards Initiative (MSI) level three and are listed
by software-assigned compound number (e.g. C1287 is compound number 1287).[11] Plasma
compound lists were exported for statistical analysis and to determine which GT compounds
were detected in plasma.

Green Tea compounds in Plasma: Data from the four GT extracts were processed as above
with the following adjustments: GT compounds had to be present in 100% of GT samples and
the compounds were restricted to those eluting before 10 min for both the aqueous and lipid
fractions. In addition to the databases used to annotate plasma, the GT samples were also
searched against Phenol Explorer, FooDB and a Traditional Chinese Medicine natural products
database (Agilent Technologies). The GT compound lists were exported as .cef files and
imported into Quantitative Analysis (Agilent Technologies).

LCMS data for plasma samples was processed as described above and imported into
Quantitative Analysis. Extracted ion chromatograms were generated for each GT compound
from plasma LCMS data using a "targeted data extraction" strategy. The method parameters
included: left and right extraction window of 10 ppm, retention time extraction window $\pm$ 0.2
minutes, and retention time outlier >0.3 minutes. For each GT compound, the monoisotopic
peak (M1) and C13 peak (M2) were used for qualification. The M1 peak was used as the
quantifier ion, while M1 and M2 peaks (or only M1 when no M2 peak was detected) were used
as qualifier ions. When a second charge carrier such as sodium was detected, its monoisotopic
peak was used as a qualifier. Qualifier relative uncertainty was set to 20%. Analysis results
were imported into MPP and filtered to retain only targets that were present in at least 3 plasma
samples with an area ≥20,000 counts for the aqueous fraction and ≥ 55,000 counts for the lipid
fraction. These compound lists, including area under the curve (AOC) for relative quantitative
comparison, were exported for statistical and informatic analysis.

Whole metabolomics datasets were visualized using principal components analysis (PCA) for
the purpose of quality control (Supplemental Methods).

Microbiome - DNA extraction and sequencing: DNA was extracted from the fecal samples using
the Power Soil Pro Kit protocol (Qiagen, Germantown, MD). Barcoded primers targeting the V4
region of the16S rRNA gene were used to PCR amplify the extracted bacterial DNA using the
Earth Microbiome Project (EMP) standard protocols (http://www.earthmicrobiome.org).[33] PCR
product quantification was completed using PicoGreen (Invitrogen, Carlsbad, CA) and used to
pool equal amounts of amplified DNA from each sample. The QIAquick PCR Purification Kit

(Qiagen) was used to clean the pooled libraries. Three runs were used to generate sequences
using the Illumina MiSeq platform (San Diego, CA).

Microbiome - Sequence data processing: Microbiome sequencing data were processed in
QIIME2-2021.8.[34] Sequences were demultiplexed using the q2-demux plugin, then denoised
using q2-dada2[35] with a truncation length of 230bp. A phylogenetic tree was created using Saté-
enabled phylogenetic placement (SEPP) via the q2-fragment-insertion plugin.[36] Taxonomy was
assigned to reads using a naïve-Bayes classifier trained on the latest Greengenes 1 database
as of January 2022 (Greengenes 13.8).[37] Reads with no taxonomy below the kingdom level and
reads classified as mitochondria and chloroplast were removed from further consideration. For
diversity analyses, samples were rarefied to 40,625 reads per sample, which was the highest
rarefaction depth that did not exclude any samples. A taxa bar plot was created using
Microshades.[38]

Differential abundance testing: Analysis of Compositions of Microbiomes with Bias Correction
(ANCOM-BC)[39] was used to assess differentially abundant taxa between GF and HU mice.
ANCOM-BC was performed at the family and genus level, and a Benjamini-Hochberg FDR
correction was applied to p values.

Diversity metrics: Alpha diversity was calculated using Faith's phylogenetic diversity,[40] and
microbiome beta diversity was calculated using unweighted UniFrac distances.[41] Metabolome
beta diversity was calculated using the Bray-Curtis distance metric.

Ordination: Microbiome principal coordinate analysis (PCoA) plots were generated using the
unweighted UniFrac distance matrix via the Python package scikit-bio (v0.5.6),[42] and
metabolomics PCoA plots were generated using Bray-Curtis distances.

Calculation of centroids for distance-based statistics: To avoid pseudoreplication[43] affecting
precision of estimates in permutation-based tests where random effects or cluster-robust
standard errors cannot be used, we collapsed microbiome UniFrac distances to centroids within
humanized microbiome donors, or germ-free centroids within each experimental cohort, using
the dist_multi_centroids function from the usedist R package (v0.4.0).[44] This reduced sample
size from n=29 (20 mice with microbiomes humanized in pairs from 10 unique donors + 9 GF
across 3 experimental cohorts) to n=13 (10 averaged HU centroids corresponding to each
donor microbiome + 3 GF centroids corresponding to each cohort). Corresponding non-distance
data (such as compound abundance) were averaged (arithmetic mean) within groups.

Procrustes randomization test: Procrustes randomization tests were performed on the
metabolomics and microbiome PCoA coordinates from centroid distances using the protocol
described by Peres-Neto and Jackson.[45] Specifically, we first performed a Procrustes
transformation on the observed datasets, and then permuted the GT lipid or GT aqueous
compound Bray-Curtis distance PCoA coordinates 10^4 times and performed a Procrustes
transformation on the permuted datasets. The *p* value was calculated based on the portion of

permuted Procrustes-transformed datasets with resulting m^2 (Gower's statistic, also referred to
as disparity) scores lower than the Procrustes m^2 score of the observed datasets.

Compounds associated with microbiome composition: To identify compounds and
neurochemicals associated with microbiome community composition, PERMANOVA tests were
performed using Adonis2 from the R package vegan (v2.6-4)[46] to assess correlations between
plasma abundances of each individual compound and the microbiome unweighted UniFrac
distance matrix (using 10^4 permutations). _p_ values were adjusted with a Benjamini-Hochberg
correction to reduce false discovery rates. Due to reduced power from using centroids, we
considered uncorrected _p_ < 0.01 to be associated with microbiome composition. For any
compounds associated with microbiome composition, we used the R package Selbal (v0.1.0)[47]
with 10-fold cross validation to identify genus-level balances that explained compound
abundance.

Compounds associated with microbiome diversity: We performed linear regression with cluster
robust standard errors on each individual compound and Faith's phylogenetic diversity to
assess which compounds associated with microbiome diversity. Standard errors were clustered
by humanized microbiome donor, or by experiment for GF mice, and _p_ values were calculated
based on clustered standard errors using the R package fixest (v0.11.2).[48] _p_ values were
adjusted with a Benjamini-Hochberg correction to reduce false discovery rates (FDR).
Regressions with any data points that had a dffit (difference in model fit if the point was
removed) absolute value greater than 1 were removed from further consideration.

Mixed-effects multi-omics modeling: Associations between metabolites and microbes were
assessed by pairwise linear mixed effects models between each microbe and each metabolite,
with the formula:

$z\,score(metabolite) \sim \beta * arcsinh(microbe * 100) + (1|humanized\,id) + \epsilon$　　(_eqn._1)

Metabolite concentrations were z-score transformed so that β coefficients could be compared
across compounds. Microbe relative abundances were multiplied by 100 to transform to percent
relative abundance and then arcsinh transformed ($arcsinh(x) = ln\sqrt{x^2 + 1}$) to alter the
distributions of relative abundances to minimize violations of linear modeling assumptions
across features. (1|_humanized id_) indicates a random effect for the humanized microbiome ID or
GF status. The maximum influence of any datapoint was calculated for each regression using
the mdffits function from the R package HLMdiag (v0.5.0).[49] Any regressions with any data
points that had an influence (difference in model fit if the point was removed) absolute value
greater than or equal to 4 were removed from further consideration. Models were run using the
R package lme4 (v1.1-29),[50] and _p_ values were calculated based on Satterthwaite's degrees of
freedom using the R package LmerTest (v3.1-3).[51] _p_ values were adjusted with a Benjamini-
Hochberg correction to reduce FDR.

Results

Metabolomics of mouse plasma and GT: Metabolomics of the mouse plasma resulted in 4,282
lipid and aqueous compounds while metabolomics of GT extract resulted in 4,415 compounds
(**Figure 1**). Subsequent analyses focused on only the 624 GT compounds that were detected in
at least 3 plasma samples following GT gavage. Of these, 432 were found in the lipid-rich
extract (GT lipids) while 192 were found in the aqueous extract (GT aqueous). All 624 GT
plasma compounds were next evaluated for their relationships with microbiome composition and
diversity.

Contamination of germ-free controls: Following 16S rRNA gene amplicon sequencing of GF and
HU fecal microbiomes, it was discovered that the GF mice had microbiomes (**Figure 2**). The GF
controls also had small cecums, which are uncharacteristic of GF mice (data not shown).
Specifically, the GF controls were dominated by 6 Amplicon Sequence Variants (ASVs) in the
Firmicutes phylum, which made up over 90% of the detected reads. **Supplemental Table 1**
shows the taxonomic assignment and relative abundances of these ASVs. Following extensive
investigation, it was determined that the contamination resulted from the irradiated low
polyphenol food that was fed to mice. For example, bacterial colonies grew when irradiated food
was plated and PCR analysis showed a range of 3.5 ng/μL-4.6 ng/μL in the 3 irradiated samples
tested. No growth was seen when irradiated GT was plated. Discussions with the vendor were
largely inconclusive, but it was surmised that the source of bacteria may have been casein used
in the diet.[52] While not ideal, because all mice in the study received the same food, it was
determined that all mice were exposed to this contamination, and therefore we continued with
data analysis, focusing on statistical methods that consider microbiome community composition
or diversity rather than solely group status (GF vs HU).

Taxonomic composition of fecal microbiomes: GF and HU mouse microbiomes differed in
community composition (pseudo-F = 13.8, $p < 0.001$). GF mice were colonized almost entirely
by Firmicutes, with six of the nine mice having high levels of unassigned ASVs in the family
Peptostreptococcaceae **(Figure 2**, p = 0.005**)**. Additionally, GF mice had high relative
abundances of *Epulopiscium* (ANCOM-BC; p = 0.015), and *Turicibacter* (ANCOM-BC; p =
0.002) in their microbiomes compared to HU mice **(Figure 2)**.

Microbiome composition was very similar within HU mouse replicates, and variable between
mice colonized with different human donors (**Figure 2**). The microbiome of all but two pairs of
HU mice had *Akkermansia* present at high relative abundances. All HU pairs had representation
of Bacteroidetes, primarily including *Bacteroides*, *Parabacteroides,* and *Alistipes*. Proteobacteria
were present in the microbiome of all HU but not GF mice, and this phylum predominantly
consisted of ASVs belonging to *Bilophila* and *Sutterella*. Some HU pairs also had low relative
abundances (<5%) of *Fusobacteria* in their fecal microbiome.

Diversity and composition of the fecal microbiome, plasma metabolome, and plasma lipidome:
Both alpha and beta diversity of the fecal microbiome differed between HU and GF mice.
Specifically, microbiome phylogenetic composition (UniFrac distance) strongly differed between
GF and HU mice (pseudo-F = 5.8, p < 0.005), and clear separation between GF and HU can be
seen in the unweighted UniFrac PCoA (**Figure 3A**). Additionally, Faith's phylogenetic diversity
was significantly higher in HU mice (mean [95% confidence interval (CI)] = 12.0 [12.3 – 13.3])
than GF mice (mean [95% CI] = 4.1 [2.3 – 5.9]; regression β = 8.7, p < 0.001; **Figure 3B**). The
composition of GT-specific lipids in the plasma showed a stronger difference between GF and
HU mice (Bray-Curtis distance PERMANOVA on centroids R^2 = 0.12, pseudo-F = 1.5, p = 0.13;
Figure 3C) than the composition of aqueous GT-specific did (Bray-Curtis distance
PERMANOVA on centroids R^2 = 0.12, pseudo-F = 0.37, p = 0.92; **Figure 3D**), though neither of
these were significant. This can be seen in PCoA plots, where the GT lipid compounds in
plasma weakly separate, and the GT aqueous compounds in plasma do not cluster according to
GF or HU group (**Figure 3C and 3D**). The portion of shared GT compounds present in plasma
was high in both the lipid and aqueous datasets (Jaccard similarity lipid mean [min – max] = 1.0
[0.99 – 1.0]; aqueous mean [min – max] = 0.95 [0.88 – 0.98].
365
366
Multivariate associations between microbiome and plasma metabolome composition: A
Procrustes randomization test revealed a stronger relationship between microbiome
composition (UniFrac) and GT lipids in plasma (**Figure 4A**, m^2 = 0.55, p = 0.053) than between
microbiome composition and aqueous GT compounds in plasma (using Procrustes-transformed
Bray-Curtis PCoA, **Figure 4B**, m^2 = 0.37, p = 0.87). This is supported by the results of mantel
tests on centroids, which found a nearly significant relationship between UniFrac distance and
GT lipids in plasma (r = 0.36, p = 0.06) and no association between UniFrac distance and
aqueous GT compounds (r = -0.18, p = 0.90).
375
Green tea compounds associated with microbiome composition: One goal of the study was to
determine if GT compounds in plasma associate with microbiome diversity and composition.
Understanding these associations are a first step towards determining which bacterial species
may be responsible for the metabolism of specific GT compounds. PERMANOVA tests on
centroids assessing associations between the concentration of each GT compound and
microbiome composition (unweighted UniFrac) found that out of 624 individual aqueous
or lipid GT compounds in plasma, 59 GT compounds were significantly associated with
microbiome composition (p < 0.05), though none of these relationships remained significant
after FDR correction (**Figure 5 and Supplemental Table 2**)**.** For example, montecristin,
PG(20:0), shikonofuran D, acetylagmatine, PC(38:4), clitidine, ethyl-p-methoxycinnamate,
gamma-glutamyl-alanine, crotonic acid, and lactiflorin were the 10 annotated GT compounds
most associated with microbiome composition and with p < 0.01.
388
389 Selbal-identified balances were able to explain 45% to 70% of the variation in the abundance of
390 these compounds, with gamma-glutamyl-alanine having the most variation explained by a
391 positive correlation with a balance of the genera *Alistipes* to *Butyricimonas* **(Figure 6)**. Balances

392   identified by Selbal are ratios of taxa, meaning that as gamma-glutamyl-alanine increased,
393   *Alistipes* increased relative to *Butyricimonas,* or *Butyricimonas* decreased relative to *Alistipes.*
394
Green tea compounds associated with microbiome phylogenetic diversity: When testing GT
compound associations with microbiome alpha diversity (Faith's phylogenetic diversity), 11
compounds had FDR-corrected *p* < 0.05 **(Figure 7 and Supplemental Table 3)**, including
lactiflorin, acetylagmatine, aspartic acid, 7,8- dihydroparasiloxanthin, montecristin, CL(82:16),
PG(20:0), and PS(41:7) **(Figure 8).** Most of these compounds had negative associations with
microbiome diversity, except for CL(82:16) and an unannotated compound.
401
Relationships between individual microbes and GT compounds: When controlling for the
humanized microbiome with which mice were colonized with a random effect and applying a
Benjamini-Hochberg FDR correction, across all mice (including GF mice) we identified 286
significant relationships between microbes and GT compounds in plasma out of the potential
39,936 relationships tested. Of these, the strongest associations were between an ASV
assigned to the class Clostridia and ganoderic acid, an ASV in the genus *Catenibacterium* and
ganoderic acid, an ASV in the genus *Bifidobacterium* and an unannotated compound, an ASV in
the genus *Allobacterium* and montecristin, and an ASV in the family Peptostreptococcaceae and
acetylagmatine (data not shown).
411
Despite including GF status as a covariate in each compound-taxon regression, there was still a
large signal related to the high relative abundance taxa in the GF mice, such as
Peptostreptococcaceae. Thus, we also performed this analysis in only HU mice, as differences
in overall architecture of the GF vs HU mice fecal microbiomes may have biased these
individual metabolite-taxon relationships. When mixed effects regressions were performed for
only HU mice, several GT compounds were significantly associated with specific bacterial
ASVs. Of 39,963 potential relationships tested, 24 were significant after Benjamini-Hochberg
FDR correction (**Figure 9A**). For example, the GT compound picrotoxinin was positively
associated with ASVs in the genera *Chistensenella*. Spiramycin was positively associated with
taxa in the genera *Lactobacillus* and *Gemmiger* (**Figure 9B**). While multiple unannotated
compounds had multiple significantly associated microbes, spiramycin was the annotated
compound with more than one significant relationship with individual taxa (after FDR correction).
The genus *Gemmiger* had the highest number of significant relationships with GT compounds,
though 3 of these 5 relationships were primarily with unannotated compounds, and the
relationships between compounds are not known.
427

Discussion:
We show that GT compounds significantly associated with specific gut bacterial genera
following the acute feeding of GT extract to mice. Our analysis strategy included an important
data reduction step, whereby analysis focused on only GT compounds that were found in the
plasma of mice who had consumed GT. This greatly increased power (via decreasing the need
for as stringent FDR correction) by limiting analysis from 4,282 plasma compounds to 624
compounds that were specific to the intervention. Importantly, these are compounds detected in
dietary formulations of green tea, not solely compounds produced by the plant from which green

tea is derived, *Camellia sinensis.* Much like the mice in this study with associated microbiomes,
C. sinensis plants are meta-organisms that coexist with microbial communities.[53,54] Microbial
metabolites and even excreted metabolites from other nearby plants or pollinators may be
present in the leaves used to create the tea used in this study, so the GT compounds discussed
may come from a variety of sources across multiple kingdoms of life.

Sequencing of GF and HU microbiomes from fecal pellets allowed us to perform analyses
considering microbiome composition and diversity, rather than solely GF status. This was
particularly useful because the GF mice in this study were populated with a low diversity
microbiome due to contamination of the low polyphenol food used. On a large scale, we found a
stronger multivariate association between phylogenetic composition of the gut microbiome and
the lipid GT compounds, relative to the aqueous GT compounds detected in plasma. One
reason for this could be that aqueous compounds could have been more thoroughly absorbed in
the small intestine (prior to interactions with the microbiome), whereas lipids may not have been
as thoroughly absorbed due to the lack of a solubilization by bile acids. Bile acid secretion,
which is important for lipid absorption, is stimulated *via* cholecystokinin by moderate to high lipid
concentration in the small intestine;[55] however, because the green tea gavage used in this study
would not have resulted in high concentrations of lipids in the small intestine, low secretion of
bile acids could have slowed absorption of GT lipids, resulting in more interactions with the
colonic microbiome prior to absorption. Notably, there are other potential explanations, such as
altered lipid absorption-relevant enterocyte physiology or altered host metabolism of GT lipids,
so future studies should consider measuring bile acid concentrations before and after GT
gavage, assessing intestinal histology, and assessing relevant host processes (such as through
transcriptomics) to construct a systems-level understanding of the effects of the gut microbiome
on lipid *versus* aqueous food compound absorption and metabolism. Additionally, expanding
this research to mice consuming or not consuming dietary lipids simultaneously with the GT
gavage could be used assess the generality of these results across fed and fasted states.

Multiple compounds found in green tea extracts were individually associated with microbiome
composition and diversity. Importantly, these compounds are known to be present in plants and
are relevant to mammalian physiology. For example, the annotated GT compound with the
strongest association with microbiome composition, montecristin, is an annonaceous
acetogenin (a group of compounds that are waxy derivatives of fatty acids and are reported to
have an array of health effects, including antimalarial, antiparasitic, and anti-cancer activities).[56]
Other plant compounds from GT that differ as a function of the microbiome include the following:
shikonofuran,[57] a bioactive component of the herbs *Lithospermum erythrorhizon*, *Arnebia*
euchroma, and *A. guttata*; acetylagmatine,[58,59] which can be synthesized by plants, bacteria,
and animals but with low synthesis rates in mammals; and lactiflorin, a plant compound
originally characterized in the plant genus *Paeonia*, that activates the antioxidant-controlling
transcription factor nuclear factor erythroid 2-related factor (Nrf2) in rats.[60] Lactiflorin, a
monoterpene glycoside, has been used in the treatment of rheumatoid arthritis via inhibiting
leukocyte recruitment and angiogenesis via, potentially, the VEGFR and PI3K-Akt signaling
pathways, interleukin signaling, and platelet activation.[61]

Previous studies have focused on the health effects of green tea catechins, a subgroup of
flavonoids with known antioxidant properties.[62]   While several catechins, including
epigallocatechin 3-p-coumarate, catechin 3-gallate, gallocatechin 3'-gallate, and
epigallocatechin 3-cinnamate were found in green tea extract, these were not found at
detectable levels in at least 3 mouse plasma samples and hence were not included in
microbiome analyses. This could be due to the concentration of extract used, the sensitivity of
the assay, or the timing of blood draw.
487
Most of the compounds associated with microbiome presence, diversity, or composition were
found in lower abundance in the plasma of HU mice, relative to GF mice. For example, lactiflorin
was nearly 7 times less abundant in HU mice. Our data suggest that the presence of a gut
microbiome either (1) inhibits the absorption of lactiflorin, (2) increases metabolism of lactiflorin
by the host, or (3) allows for bacterial degradation of lactiflorin. In fact, most compounds
associated with microbiome presence, diversity, or composition were less abundant in HU
relative to GF mice, suggesting potential co-metabolism of these compounds by the
microbiome. Shikonofuran D was the one annotated compound that was negatively associated
with microbiome diversity or presence.
497
Another compound identified to be associated with microbiome composition was gamma-
glutamyl-alanine. Notably, many plants and mammals possess gamma-glutamyl transpeptidase
(GGT),[63] which is required for synthesis of gamma-glutamyl amino acids and often involved in
glutathione metabolism; however, most bacteria also encode GGT,[64] making the microbiome a
candidate to alter concentrations of gamma-glutamyl-alanine either *via* direct metabolism or *via*
an unknown effect on host metabolism. 70% of the variation in plasma gamma-glutamyl-alanine
was explainable by the ratio of *Alistipes* to *Butyricimonas*. Notably, human and murine gut
colonizers within the genus *Butyricimonas* contain the gene gamma-D-glutamyl-L-lysine
dipeptidyl-peptidase (K20742),[65] which releases gamma-glutamyl-alanine from cell walls, and
Alistipes is negatively correlated with plasma GGT in humans,[66] supporting the directionality of
the relationship of this ratio with plasma gamma-glutamyl alanine abundance. However, this
balance may just be a proxy for overall microbiome functional composition, as other members of
the microbial community that interact with *Alistipes* and *Butyricimonas* may be more directly
responsible for differences in gamma-glutamyl alanine.
512
We also found many significant relationships (after FDR correction) between GT compounds
and individual taxa with only HU mice included in analysis **(Figure 9)**. In contrast to the
multivariate associations where most compounds were lower in HU mice, we found primarily
positive associations between individual taxa and GT compounds in plasma. The annotated GT
compound with the highest number of significant relationships with individual taxa was
spiramycin, an antimicrobial produced by *Streptomyces ambofaciens*, a member of the soil
microbiome/rhizosphere.[67] However, the mechanism of the positive association between
spiramycin and the genera *Gemmiger* and *Lactobacillus* is unclear. Dieporeticenin, another
member of the annonaceous acetogenin class, which has demonstrated effects on bacterial
biofilm formation in a variety of contexts,[68] was positively associated with the family RF32 of the
class Alphaproteobacteria. We found positive relationships with multiple other bioactive plant

compounds such as wilforlide B (a norditerpenoid) and *Anaerohabdus*, picrotoxinin (neurotoxin
originally found in *Anamirta cocculus* that inhibits the action of gamma-aminobutyric acid
(GABA))[69] and *Christensenella*, and petasitenine (a pyrrolizidine alkaloid isolated from *Petasites*
japonicus) and the family Coriobacteriales. While this data is compelling in that the majority of
compounds have been found in plants, the mechanistic relationships between these compounds
and individual taxa are unclear, especially given that metabolism pathways are poorly
understood. Future work could track closely related compounds or apply machine learning-
based approaches to identify potential metabolic pathways for these natural products and
assess microbial genomes for genes responsible for these reactions.
533
One limitation of the study is the small number of replicate mice that were colonized with a
specific human microbiome (n=2), and that mice were colonized with only 10 different human
microbiomes. While significant relationships between specific bacteria and GT compounds were
found, increasing the sample size would increase power to identify relationships, especially in
the permutation-based tests, where we needed to reduce our effective sample size to control for
pseudoreplication. More samples would potentially allow for using within-estimator (within each
humanized microbiome source) to estimate the effects of small changes to individual taxa within
a similar microbiome community structure subtype; this is akin to estimating the effects of
increasing one species of bird within different terrestrial biomes like grassland and savannah.
Another weakness is the lack of plasma and microbiome sampling before green tea gavage.
Having samples from mice in each group that had not received a green tea gavage, either
before/after or untreated controls, would also increase our ability to identify metabolism of GT
compounds, as some of the compounds identified were likely already present in the plasma of
our mice. However, our focus on only GT compounds found in plasma does reduce this
possibility. Moreover, we chose the two-hour timepoint to maximize the GT compound signal in
plasma, though multiple plasma samples after feeding would allow for more detailed resolution
of GT compound abundance trajectories based on microbiome composition, including time lags
to assess causal directionality. An additional limitation of the study is the presence of gut
microbiomes, albeit of low diversity, in the GF mouse group. Although it was assumed that all
mice were exposed to the same contaminated low polyphenol mouse food, the contaminating
taxa were able to dominate the fecal microbiome of the GF mice, such that there were no mice
free of microbial metabolism of GT compounds. Currently underway studies utilize mice
unexposed to GT as a control in a similarly designed acute feeding study, as well as a two-week
feeding of GT to compare the effects of the microbiome on sub-chronic versus acute
metabolism of GT.
559
One strength of our experimental design lies in the use of 10 different human microbiomes to
enable the evaluation of relationships between individual microbial species and GT compounds.
While many humanized microbiome studies are pseudoreplicated,[43] our use of multiple donor
microbiomes while controlling for replication covers a broad variety of possible human-like
microbiomes. Together, these results suggest that specific bacterial species may affect the
metabolism of these bioactive compounds, thereby influencing their health effects.
566

Overall, this study represents a successful workflow for discovering relationships between food
compounds and composition of the gut microbiome, as well as individual gut bacterial genera. In
addition, this methodology has allowed us to track GT compounds from the tea to plasma.
Though our study was limited in size and timepoints, we believe this methodology can and
should be applied to study the effects on the gut microbiome on food metabolism. We identified
multiple relationships between microbiome composition and GT compounds in plasma after GT
consumption, supporting that bacterial taxa affect the absorption and metabolism of GT
compounds, thereby possibly influencing their positive or negative health effects.
575

576 **Acknowledgements**

577 This project was funded by the National Science Foundation-sponsored Interdisciplinary
578 Quantitative Biology PhD program, the Integrated Data Science (Int dS) Graduate Training
579 Fellowship, and the William J. Freytag Fellowship.

Figure 1. Compounds from green tea were detected in plasma after green tea gavage in mice. Flowchart shows study design. A total of 29 mice (20 humanized, 9 GF) were gavaged with 100 μL of green tea then sacrificed 2 hours later, and plasma, whole brain, and fecal microbiome samples were collected. Liquid chromatography mass spectrometry-based metabolomics was performed on green tea samples and plasma. Plasma metabolomics data was mined to determine which green tea compounds could be found in plasma. Relationships between the fecal microbiome and green tea compounds found in plasma were assessed. Abbreviations: GF, germ free; LCMS, liquid chromatography-mass spectrometry; MeOH, methanol; rRNA, ribosomal ribonucleic acid.

591
**Figure 2. Taxa bar plot shows differences in phylum and genus level composition
between GF and humanized mice microbiomes.** Bars are colored by phylum and
microshaded by genus, and height of each colored bar represents the relative abundance of the
corresponding taxon. One stacked bar represents the composition of one mouse's microbiome,
and humanized microbiome IDs are shown along the x-axis. Germ-free mice are labeled as
"Control" and have vastly different composition compared to humanized mice.
598

599
Figure 3. Microbiome PCoA and alpha diversity show clear separation between germ free
and humanized mice, whereas metabolome PCoA plots do not separate as clearly. A)
Microbiome unweighted UniFrac PCoA is colored by humanized vs germ free status. B) Boxplot
shows higher Faith's phylogenetic diversity in humanized vs germ free mice. C) PCoA of lipid
fraction of green tea compounds in plasma is colored by humanized vs germ free status. D)
PCoA of aqueous fraction of green tea compounds in plasma is colored by humanized vs germ
free status. Abbreviation: PCoA, principal coordinates analysis; PCo, principal coordinates axis.
607

608
Figure 4. Procrustes analysis indicates strong association between green tea lipids in
plasma and the microbiome but not between green tea aqueous compounds in plasma
and the microbiome. Panel A shows the connection between microbiome composition and the

composition of aqueous GT compounds detected in plasma, where each triangle represents the
composition of one microbiome sample, and a line connects to that sample's corresponding
aqueous GT compounds in plasma. Panel B shows the connection between microbiome
composition and the composition of lipid GT compounds detected in plasma, where each
triangle represents the composition of one microbiome sample, and a line connects to that
sample's corresponding lipid GT compounds in plasma. A Procrustes randomization test
revealed a nearly-significant relationship between the lipid GT compounds in plasma and gut
microbiome composition ($p = 0.053$), whereas there was not a significant relationship between
the aqueous GT compounds and microbiome composition ($p = 0.87$). In both panels,
unweighted UniFrac microbiome PCoA coordinates were left untransformed, and metabolomics
sample Bray-Curtis PCoA coordinates were transformed with a Procrustes function. Samples of
the same type positioned close together were compositionally similar, whereas samples far
apart were dissimilar. Shorter lines between samples indicate better overlap between the
microbiome and metabolomics datasets. For statistical testing, samples belonging to mice from
the same donor microbiome or germ-free mice from the same experimental cohort were
collapsed into one centroid per treatment unit. Abbreviation: PCoA, principal coordinates
analysis; PCo, principal coordinates axis.
629

total = 624 variables

Figure 5. Several green tea compounds are associated with microbiome composition and are different between germ free and humanized mice. Each point represents one green tea compound. The y-axis represents p-values from a PERMANOVA assessing the correlation between that compound and microbiome unweighted UniFrac composition, and the x-axis represents fold change in the compound's mean abundance from germ free to humanized mice, where positive values indicate higher abundance in humanized mice. The horizontal line represents $p = 0.01$, and vertical lines represent a 2-fold change in mean compound abundance. Compounds with $p < 0.01$ and fold change absolute value > 2 are labeled with their annotation.

Figure 6. Compositional balances in microbial relative abundances explain green tea compound abundance in plasma. Selbal with cross validation was used to identify balances (log-ratios) of taxa that predict compounds that were associated with microbiome composition. The x-axis of each plot shows the microbiome balance, with the numerator and denominator labeled on the plot as <family>_<genus>. The y-axis shows the abundance of each green tea compound, which is labeled at the top of each plot, along with the regression R^2. The line on each plot shows the regression line of best fit and the shaded region indicates the 95% confidence interval for that regression line. Each point represents the average balance for each donor microbiome or experimental control.

total = 624 variables

Figure 7. Several green tea metabolites are associated with phylogenetic diversity of the gut microbiome. Each point represents one green tea compound. The y-axis represents p-values from a regression between that compound and microbiome phylogenetic diversity, and the x-axis represents the effect size of that compound's association with microbiome diversity. Compounds abundances were z-score transformed to allow for comparison of effect sizes, and p-values were calculated using cluster robust standard errors, where cluster ID was the donor microbiome source, or experimental cohort for germ-free mice. The horizontal line represents p = 0.01, and vertical lines represent an effect size of 1, indicating that a 1 standard deviation increase in compound abundance was associated with a 1 unit increase in Faith's phylogenetic diversity. Compounds with p < 0.01 and fold change absolute value > 2 are labeled with their annotation.

Figure 8. Phylogenetic diversity of the gut microbiome is associated with green tea compound abundance in plasma. Fixed effects regression with cluster robust standard errors (clustered by humanized microbiome donor ID or germ-free experimental cohort) was used to identify green tea compounds that were associated with Faith's phylogenetic diversity of the gut microbiome, and compounds that had significant (Benjamini-Hochberg $p < 0.05$) relationship with Faith's phylogenetic diversity are shown. The x-axis of each plot shows Faith's phylogenetic diversity, and the y-axis shows the abundance of each green tea compound, which is labeled at the top of each plot. The line on each plot shows the regression line of best fit and the shaded region indicates the 95% confidence interval for that regression line. Each point represents one sample.

Figure 9. Individual green tea compounds are associated with specific genera when controlling for overall microbiome composition. A) A stacked bar plot shows a summary of relationships between green tea compounds and individual genus-level relative abundances, indicating the significance and direction of compound-microbe relationships. B) A shaped heatmap shows the direction and significance of any microbes and compounds that had significant relationships after Benjamini-Hochberg p-value correction. To generate these results, 39,860 pairwise linear mixed effects regressions were run on z-score(metabolite) ~ arcsinh(microbe) + (1|humanized_id), where (1|humanized_id) indicates the microbiome mice were humanized with, or if mice were germ free. Metabolites were transformed to have a mean of 0 and standard deviation of 1 to allow for comparison of effect sizes across metabolites. Each point represents the regression run between the metabolite on the x-axis and the microbe shown on the y-axis. The color of each point represents the beta extracted from the model (coefficient of the microbe), and the size of points represents the p-value of that coefficient. Circles represent p > 0.05, whereas triangles represent p < 0.05.

References

1. Jones DP, Park Y, Ziegler TR. Nutritional metabolomics: Progress in addressing complexity
in diet and health. *Annu Rev Nutr*. 2012;32:183-202. doi:10.1146/annurev-nutr-072610-
145159

2. Bouchard-Mercier A, Rudkowska I, Lemieux S, Couture P, Vohl MC. The metabolic
signature associated with the Western dietary pattern: a cross-sectional study. *Nutr J*.
2013;12:158. doi:10.1186/1475-2891-12-158

3. Rangel-Huerta OD, Gil A. Nutrimetabolomics: An Update on Analytical Approaches to
Investigate the Role of Plant-Based Foods and Their Bioactive Compounds in Non-
Communicable Chronic Diseases. *Int J Mol Sci*. 2016;17(12):2072.
doi:10.3390/ijms17122072

4. Wang Z, Klipfell E, Bennett BJ, et al. Gut flora metabolism of phosphatidylcholine promotes
cardiovascular disease. *Nature*. 2011;472(7341):57-63. doi:10.1038/nature09922

5. Dopkins N, Nagarkatti PS, Nagarkatti M. The role of gut microbiome and associated
metabolome in the regulation of neuroinflammation in multiple sclerosis and its implications
in attenuating chronic inflammation in other inflammatory and autoimmune disorders.
Immunology. 2018;154(2):178-185. doi:10.1111/imm.12903

6. Shaffer M, Armstrong AJS, Phelan VV, Reisdorph N, Lozupone CA. Microbiome and
metabolome data integration provides insight into health and disease. *Transl Res J Lab Clin
Med*. 2017;189:51-64. doi:10.1016/j.trsl.2017.07.001

7. Markle JGM, Frank DN, Mortin-Toth S, et al. Sex Differences in the Gut Microbiome Drive
Hormone-Dependent Regulation of Autoimmunity. *Science*. 2013;339(6123):1084-1088.
doi:10.1126/science.1233521

8. Markle JGM, Frank DN, Adeli K, von Bergen M, Danska JS. Microbiome manipulation
modifies sex-specific risk for autoimmunity. *Gut Microbes*. 2014;5(4):485-493.
doi:10.4161/gmic.29795

9. Bowler RP, Jacobson S, Cruickshank C, et al. Plasma sphingolipids associated with chronic
obstructive pulmonary disease phenotypes. *Am J Respir Crit Care Med*. 2015;191(3):275-
284. doi:10.1164/rccm.201410-1771OC

10. Hughes G, Cruickshank-Quinn C, Reisdorph R, et al. MSPrep--summarization,
normalization and diagnostics for processing of mass spectrometry-based metabolomic
data. *Bioinforma Oxf Engl*. 2014;30(1):133-134. doi:10.1093/bioinformatics/btt589

11. Quinn KD, Schedel M, Nkrumah-Elie Y, et al. Dysregulation of metabolic pathways in a
mouse model of allergic asthma. *Allergy*. 2017;72(9):1327-1337. doi:10.1111/all.13144

12. Reisdorph N, Wechsler ME. Utilizing metabolomics to distinguish asthma phenotypes:
strategies and clinical implications. *Allergy*. 2013;68(8):959-962. doi:10.1111/all.12238

13. Frank DN, Zhu W, Sartor RB, Li E. Investigating the biological and clinical significance of
human dysbioses. *Trends Microbiol*. 2011;19(9):427-434. doi:10.1016/j.tim.2011.06.005

14. Vlainić JV, Šuran J, Vlainić T, Vukorep AL. Probiotics as an Adjuvant Therapy in Major
Depressive Disorder. *Curr Neuropharmacol*. 2016;14(8):952-958.
doi:10.2174/1570159x14666160526120928

15. Weiss R, Dziura J, Burgert TS, et al. Obesity and the metabolic syndrome in children and
adolescents. *N Engl J Med*. 2004;350(23):2362-2374. doi:10.1056/NEJMoa031049

16. Strober W. Impact of the gut microbiome on mucosal inflammation. *Trends Immunol*.
2013;34(9):423-430. doi:10.1016/j.it.2013.07.001

17. Marcobal A, Kashyap PC, Nelson TA, et al. A metabolomic view of how the human gut
microbiota impacts the host metabolome using humanized and gnotobiotic mice. *ISME J*.
2013;7(10):1933-1943. doi:10.1038/ismej.2013.89

18. Wikoff WR, Anfora AT, Liu J, et al. Metabolomics analysis reveals large effects of gut
microflora on mammalian blood metabolites. *Proc Natl Acad Sci*. 2009;106(10):3698-3703.
doi:10.1073/pnas.0812874106

19. Theriot CM, Bowman AA, Young VB. Antibiotic-Induced Alterations of the Gut Microbiota
Alter Secondary Bile Acid Production and Allow for Clostridium difficile Spore Germination
and Outgrowth in the Large Intestine. *mSphere*. 2016;1(1):10.1128/msphere.00045-15.
doi:10.1128/msphere.00045-15

20. Wu GD, Chen J, Hoffmann C, et al. Linking long-term dietary patterns with gut microbial
enterotypes. *Science*. 2011;334(6052):105-108. doi:10.1126/science.1208344

21. Filippis FD, Pellegrini N, Vannini L, et al. High-level adherence to a Mediterranean diet
beneficially impacts the gut microbiota and associated metabolome. *Gut*. 2016;65(11):1812-
1821. doi:10.1136/gutjnl-2015-309957

22. Naveed M, BiBi J, Kamboh AA, et al. Pharmacological values and therapeutic properties of
black tea (Camellia sinensis): A comprehensive overview. *Biomed Pharmacother
Biomedecine Pharmacother*. 2018;100:521-531. doi:10.1016/j.biopha.2018.02.048

23. Cao J, Han J, Xiao H, Qiao J, Han M. Effect of Tea Polyphenol Compounds on Anticancer
Drugs in Terms of Anti-Tumor Activity, Toxicology, and Pharmacokinetics. *Nutrients*.
2016;8(12):762. doi:10.3390/nu8120762

24. de Mejia EG, Ramirez-Mares MV, Puangpraphant S. Bioactive components of tea: Cancer,
inflammation and behavior. *Brain Behav Immun*. 2009;23(6):721-731.
doi:10.1016/j.bbi.2009.02.013

25. Jochmann N, Baumann G, Stangl V. Green tea and cardiovascular disease: from molecular
targets towards human health. *Curr Opin Clin Nutr Metab Care*. 2008;11(6):758-765.
doi:10.1097/MCO.0b013e328314b68b

26. van Duynhoven J, Vaughan EE, Jacobs DM, et al. Metabolic fate of polyphenols in the
human superorganism. *Proc Natl Acad Sci*. 2011;108(supplement_1):4531-4538.
doi:10.1073/pnas.1000098107

27. Axling U, Olsson C, Xu J, et al. Green tea powder and Lactobacillus plantarum affect gut
microbiota, lipid metabolism and inflammation in high-fat fed C57BL/6J mice. *Nutr Metab*.
2012;9(1):105. doi:10.1186/1743-7075-9-105

28. Uffelman CN, Doenges KA, Armstrong ML, et al. Metabolomics Profiling of White Button,
Crimini, Portabella, Lion's Mane, Maitake, Oyster, and Shiitake Mushrooms Using
Untargeted Metabolomics and Targeted Amino Acid Analysis. *Foods*. 2023;12(16):2985.
doi:10.3390/foods12162985

29. Turi KN, Michel CR, Manke J, Doenges KA, Reisdorph N, Bauer AK. Multi-Omics Analysis
of Lung Tissue Demonstrates Changes to Lipid Metabolism during Allergic Sensitization in
Mice. *Metabolites*. 2023;13(3):406. doi:10.3390/metabo13030406

30. Yang Y, Cruickshank C, Armstrong M, Mahaffey S, Reisdorph R, Reisdorph N. New sample
preparation approach for mass spectrometry-based profiling of plasma results in improved
coverage of metabolome. *J Chromatogr A*. 2013;1300:217-226.
doi:10.1016/j.chroma.2013.04.030

31. Reisdorph NA, Hendricks AE, Tang M, et al. Nutrimetabolomics reveals food-specific
compounds in urine of adults consuming a DASH-style diet. *Sci Rep*. 2020;10(1):1157.
doi:10.1038/s41598-020-57979-8

32. Sutliff AK, Saint-Cyr M, Hendricks AE, et al. Lipidomics-Based Comparison of Molecular
Compositions of Green, Yellow, and Red Bell Peppers. *Metabolites*. 2021;11(4):241.
doi:10.3390/metabo11040241

33. Gilbert JA, Meyer F, Jansson J, et al. The Earth Microbiome Project: Meeting report of the
"1st EMP meeting on sample selection and acquisition" at Argonne National Laboratory
October 6th 2010. *Stand Genomic Sci*. 2010;3(3):249-253. doi:10.4056/aigs.1443528

34. Bolyen E, Rideout JR, Dillon MR, et al. Reproducible, interactive, scalable and extensible
microbiome data science using QIIME 2. *Nat Biotechnol*. 2019;37(8):852-857.
doi:10.1038/s41587-019-0209-9

35. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016;13(7):581-583.
doi:10.1038/nmeth.3869

36. Janssen S, McDonald D, Gonzalez A, et al. Phylogenetic Placement of Exact Amplicon
Sequences Improves Associations with Clinical Information. *mSystems*. 2018;3(3):e00021-
18. doi:10.1128/mSystems.00021-18

37. McDonald D, Price MN, Goodrich J, et al. An improved Greengenes taxonomy with explicit
ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J*.
2012;6(3):610-618. doi:10.1038/ismej.2011.139

38. Dahl EM, Neer E, Bowie KR, Leung ET, Karstens L. microshades: An R Package for
Improving Color Accessibility and Organization of Microbiome Data. *Microbiol Resour*
Announc. 2022;11(11):e00795-22. doi:10.1128/mra.00795-22

39. Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction. *Nat*
Commun. 2020;11(1):3514. doi:10.1038/s41467-020-17041-7

40. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv.* 1992;61(1):1-
10. doi:10.1016/0006-3207(92)91201-3

41. Lozupone C, Knight R. UniFrac: a New Phylogenetic Method for Comparing Microbial
Communities. *Appl Environ Microbiol.* 2005;71(12):8228-8235.
doi:10.1128/AEM.71.12.8228-8235.2005

42. Rideout JR, Caporaso G, Bolyen E, et al. scikit-bio. Published online August 2, 2023.
doi:10.5281/ZENODO.8209901

43. Walter J, Armet AM, Finlay BB, Shanahan F. Establishing or Exaggerating Causality for the
Gut Microbiome: Lessons from Human Microbiota-Associated Rodents. *Cell.*
2020;180(2):221-232. doi:10.1016/j.cell.2019.12.025

44. Apostol TM, Mnatsakanian MA. Sums of Squares of Distances in m-Space. *Am Math Mon.*
2003;110(6):516-526. doi:10.2307/3647907

45. Peres-Neto PR, Jackson DA. How well do multivariate data sets match? The advantages of
a Procrustean superimposition approach over the Mantel test. *Oecologia.* 2001;129(2):169-
178. doi:10.1007/s004420100720

46. Oksanen J, Simpson GL, Blanchet FG, et al. *Vegan: Community Ecology Package.*; 2024.
https://vegandevs.github.io/vegan/

47. Rivera-Pinto J, Egozcue JJ, Pawlowsky-Glahn V, Paredes R, Noguera-Julian M, Calle ML.
Balances: a New Perspective for Microbiome Analysis. *mSystems.*
2018;3(4):10.1128/msystems.00053-18. doi:10.1128/msystems.00053-18

48. Bergé L. Efficient estimation of maximum likelihood models with multiple fixed-effects: the R
package FENmlm. *CREA Discuss Pap.* 2018;(13).

49. Loy A, Hofmann H. HLMdiag: A Suite of Diagnostics for Hierarchical Linear Models in R. *J*
Stat Softw. 2014;56(5):1-28.

50. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J*
Stat Softw. 2015;67(1):1-48. doi:10.18637/jss.v067.i01

51. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed
Effects Models. *J Stat Softw.* 2017;82(13):1-26. doi:10.18637/jss.v082.i13

52. Bisanz JE, Upadhyay V, Turnbaugh JA, Ly K, Turnbaugh PJ. Meta-Analysis Reveals
Reproducible Gut Microbiome Alterations in Response to a High-Fat Diet. *Cell Host*
Microbe. 2019;26(2):265-272.e4. doi:10.1016/j.chom.2019.06.013

53. Bag S, Mondal A, Banik A. Exploring tea (*Camellia sinensis*) microbiome: Insights into the
functional characteristics and their impact on tea growth promotion. *Microbiol Res.*
2022;254:126890. doi:10.1016/j.micres.2021.126890

54. Hassani MA, Özkurt E, Seybold H, Dagan T, Stukenbrock EH. Interactions and
Coadaptation in Plant Metaorganisms. *Annu Rev Phytopathol*. 2019;57:483-503.
doi:10.1146/annurev-phyto-082718-100008

55. McLaughlin JT, Lomax RB, Hall L, Dockray GJ, Thompson DG, Warhurst G. Fatty acids
stimulate cholecystokinin secretion via an acyl chain length-specific, Ca2+-dependent
mechanism in the enteroendocrine cell line STC-1. *J Physiol*. 1998;513(Pt 1):11-18.
doi:10.1111/j.1469-7793.1998.011by.x

56. Moghadamtousi SZ, Fadaeinasab M, Nikzad S, Mohan G, Ali HM, Kadir HA. *Annona*
muricata (Annonaceae): A Review of Its Traditional Uses, Isolated Acetogenins and
Biological Activities. *Int J Mol Sci*. 2015;16(7):15625-15658. doi:10.3390/ijms160715625

57. Liao M, Li A, Chen C, et al. Systematic identification of shikonins and shikonofurans in
medicinal Zicao species using ultra-high performance liquid chromatography quadrupole
time of flight tandem mass spectrometry combined with a data mining strategy. *J*
Chromatogr A. 2015;1425:158-172. doi:10.1016/j.chroma.2015.11.028

58. Haenisch B, von Kügelgen I, Bönisch H, et al. Regulatory mechanisms underlying agmatine
homeostasis in humans. *Am J Physiol Gastrointest Liver Physiol*. 2008;295(5):G1104-1110.
doi:10.1152/ajpgi.90374.2008

59. Galgano F, Caruso M, Condelli N, Favati F. Focused review: agmatine in fermented foods.
Front Microbiol. 2012;3:199. doi:10.3389/fmicb.2012.00199

60. Yu J, Elix JA, Iskander MN. Lactiflorin, a monoterpene glycoside from paeony root.
Phytochemistry. 1990;29(12):3859-3863. doi:10.1016/0031-9422(90)85347-I

61. Hao J, Qi F, Wang H, et al. Network pharmacology-based prediction of inhibiting leukocyte
recruitment and angiogenesis of total glucosides of peony against rheumatoid arthritis. *Ann*
Palliat Med. 2022;11(10):3085-3101. doi:10.21037/apm-21-2203

62. Musial C, Kuban-Jankowska A, Gorska-Ponikowska M. Beneficial Properties of Green Tea
Catechins. *Int J Mol Sci*. 2020;21(5):1744. doi:10.3390/ijms21051744

63. Masi A, Trentin AR, Agrawal GK, Rakwal R. Gamma-glutamyl cycle in plants: a bridge
connecting the environment to the plant cell? *Front Plant Sci*. 2015;6.
doi:10.3389/fpls.2015.00252

64. Saini M, Kashyap A, Bindal S, Saini K, Gupta R. Bacterial Gamma-Glutamyl
Transpeptidase, an Emerging Biocatalyst: Insights Into Structure–Function Relationship and
Its Biotechnological Applications. *Front Microbiol*. 2021;12. doi:10.3389/fmicb.2021.641251

65. Le Roy T, Van der Smissen P, Paquot A, et al. *Butyricimonas faecalis* sp. nov., isolated
from human faeces and emended description of the genus *Butyricimonas. Int J Syst Evol*
Microbiol. 2019;69(3):833-838. doi:10.1099/ijsem.0.003249

66. Yang C, Xu J, Xu X, et al. Characteristics of gut microbiota in patients with metabolic
associated fatty liver disease. *Sci Rep*. 2023;13(1):9988. doi:10.1038/s41598-023-37163-4

67. Vurukonda SSKP, Giovanardi D, Stefani E. Plant Growth Promoting and Biocontrol Activity
of *Streptomyces* spp. as Endophytes. *Int J Mol Sci*. 2018;19(4):952.
doi:10.3390/ijms19040952

68. Neske A, Ruiz Hidalgo J, Cabedo N, Cortes D. Acetogenins from Annonaceae family. Their
potential biological applications. *Phytochemistry*. 2020;174:112332.
doi:10.1016/j.phytochem.2020.112332

69. Crossley SWM, Tong G, Lambrecht MJ, Burdge H, Shenvi RA. Synthesis of (–)-Picrotoxinin
by Late-Stage Strong Bond Activation. *J Am Chem Soc*. 2020;142(26):11376-11381.
doi:10.1021/jacs.0c05042

889