HIV broadly neutralizing antibody escape dynamics drive the outcome of AAV vectored immunotherapy in humanized mice

Nicolas M.S. Galvez[1][*], Adam D. Nitido[1][*], Seo Bin Yoo[1], Yi Cao[1], Cailin E. Deal[1], Christine L. Boutros[1], Scott W. MacDonald[1], Yentli E. Soto Albrecht[1], Evan C. Lam[1], Maegan L. Sheehan[1], Dylan Parsons[1], Allen Z. Lin[1], Martin J. Deymier[1], Jacqueline M. Brady[1], Benjamin Moon[1], Christopher B. Bullock[1], Serah Tanno[1], Amarendra Pegu[2], Xuejun Chen[2], Cuiping Liu[2], Richard A. Koup[2], John R. Mascola[2], Vladimir D. Vrbanac[1], Daniel Lingwood[1] and Alejandro B. Balazs[1][#]

[1] Ragon Institute of Massachusetts General Hospital, Massachusetts Institute of Technology and Harvard University, Cambridge, MA 02139, USA

[*] These authors contributed equally.

[2] Vaccine Research Center, National Institute of Allergy and Infectious Diseases and National Institutes of Health, Bethesda, MD 20892, USA

[#]Direct all correspondence to: Dr. Alejandro Balazs, Ragon Institute of MGH, MIT & Harvard, 600 Main St, Cambridge, MA 02139; phone: (857) 268-7000; e-mail: abalazs@mgh.harvard.edu

Summary

Broadly neutralizing antibodies (bNAbs) have shown promise for prevention and treatment of HIV. Potency and breadth measured *in vitro* are often used as predictors of clinical potential; however, human studies demonstrate that clinical efficacy of bNAbs can be undermined by both pre-existing and *de novo* resistance. Here we find that HIV-infected humanized mice receiving bNAbs delivered via AAV as Vectored ImmunoTherapy (VIT) can be used to identify antibody escape paths, which are largely conserved for each bNAb. Path selection, and consequent therapeutic success, is driven by the fitness cost and resistance benefit of emerging mutations. Applying this framework, we independently modulated bNAb resistance or the fitness cost of escape mutants, resulting in enhanced efficacy of VIT. This escape path analysis successfully explains the therapeutic efficacy of bNAbs, and enables a tractable means of quantifying and comparing the potential for viral escape from therapeutics *in vivo*.

Introduction

Despite substantial efforts to develop therapies and vaccines, HIV-1 remains a global pandemic. Current combination antiretroviral therapy (cART) regimens can effectively prevent and treat HIV infections, and emerging long-acting agents such as Lenacapavir show remarkable efficacy in preventing viral acquisition[1,2]. However, their therapeutic success still requires strict adherence to the regimen, as viral rebound is swiftly observed upon treatment interruption and may lead to the development of cART resistance[1]. This is of particular relevance when access to healthcare is interrupted[3]. Over the past decade, broadly neutralizing antibodies (bNAbs) have been extensively described and characterized[4,5]. These antibodies target different sites of vulnerability on the HIV envelope (Env) protein and can prevent infection across HIV clades[4]. Their clinical success is often predicted by using metrics determined *in vitro* across large panels of diverse viral isolates[6]. These measurements consist of potency, the average inhibitory concentration needed to block 50% infection (IC_{50}) across all tested isolates, as well as breadth, the fraction of isolates neutralized at a given antibody concentration[7]. Recent advances in antibody isolation have identified new bNAbs with median potencies as low as 0.003 µg/mL[8,9] and breadth capable of neutralizing up to 96% of a panel of 208 viruses at an IC_{50} of <1 µg/mL[10].

Given their potential, significant efforts have been made to advance bNAb-based therapies, with multiple studies demonstrating their success in preventing HIV infection in mouse[11–16] and non-human primate (NHP) models[17–21]. These findings paved the way for the Antibody Mediated Protection (AMP) studies (NCT02716675 and NCT02568215), two international harmonized phase 2b randomized controlled trials

assessing the capacity of the CD4 binding site (CD4bs)-targeting bNAb VRC01 to

prevent HIV-1 acquisition[22]. While VRC01 prevented transmission of neutralization-

sensitive strains, overall efficacy was limited by a large fraction of circulating strains with

mutations conferring resistance to neutralization[23,24]. Numerous additional studies have

also demonstrated the potential for bNAb administration to transiently suppress HIV

viremia in animal models[25–32] as well as in people living with HIV (PLWH) who harbor

sensitive viruses[33–40]. However, these studies also revealed that even combinations of

two or three bNAbs targeting different sites of vulnerability can fail to control viremia due

to the emergence of escape mutations[38,41,42].

The clinical translation of bNAbs is also hindered by the need for repeated

infusions to maintain serum concentrations above a therapeutic threshold[43]. To

overcome this, we and others have described Vectored Immunoprophylaxis (VIP)[12],

which utilizes a single intramuscular (IM) injection of recombinant adeno-associated

viral (AAV) vectors engineered to encode a bNAb transgene [44–47]. VIP induces the

production of durable and protective concentrations of antibodies *in vivo*[48]. Studies of

VIP with VRC07, a CD4bs-targeting bNAb clonally related to VRC01 but with greater

potency and breadth[8,17], have demonstrated that delivery via AAV8 prevents

intravenous and mucosal transmission of HIV in humanized mice[11,12,16] and of SIV/SHIV

in NHP models[49,50]. Recently, the VRC603 trial administered AAV8-VRC07[11,51] to cART-

suppressed patients (NCT03374202). In this study, volunteers reported up to 3 µg/mL of

VRC07 in serum for at least three years following a single IM administration[51]. Despite

these promising results, anti-drug antibodies (ADAs) emerged in a subset of

participants, highlighting one of the challenges of this approach. However, recent

studies suggest that ADAs may be less likely to emerge in settings of increased immune

tolerance, such as transient immunosuppression of adult NHPs[52] or in pediatric

settings[53].

The therapeutic potential of bNAbs delivered via AAV as Vectored

ImmunoTherapy (VIT) to cure an established HIV infection is an area of active

investigation, given its potential use in resource-limited settings where ART adherence

is suboptimal[53]. In particular, the success of VIT can be hindered by the selection of

escape variants, as seen during previous trials of passively transferred bNAbs[54]. Here,

we present the use of VIT in HIV-infected humanized mice as an experimental model to

evaluate viral suppression and profile the emergence of Env mutations leading to viral

escape from individual vectored antibodies. We chose two widely used Clade B, Tier 2

isolates for this: $HIV_{JR\text{-}CSF}$, a chronic virus cloned from the CSF of a person who died of

severe AIDS encephalopathy[55]; and $HIV_{REJO.c}$, a transmitter/founder isolate that was

computationally derived as the likely strain that initiated a heterosexual transmission[56].

Both of these isolates represent clinically relevant viruses, as they were cloned soon

after isolation and have not been extensively cultured *in vitro* unlike laboratory adapted

strains (e.g., $HIV_{NL4\text{-}3}$)[57]. To our surprise, AAV8-VRC07 treatment drove viral evolution

towards a high fitness cost escape path that often failed to emerge, enabling long-term

suppression of $HIV_{REJO.c}$. Interestingly, while necessary for activity, traditional breadth

and potency metrics were insufficient to predict therapeutic outcome; instead, the

specific fitness cost and resistance benefit tradeoffs of the primary escape paths best

explained the success of VIT. To understand this dynamic, we adapted the fitness

landscape[58] framework to create an 'escapability' map for each antibody-strain

combination, effectively explaining VIT outcomes by identifying the most traversed

escape paths during vectored antibody therapy. Our data show that escape pathway

dynamics are a key component to understanding the *in vivo* efficacy of antibody-

mediated therapies against HIV, and that future therapeutic strategies that increase the

fitness cost of the primary escape paths are essential for clinical success.

Results

VRC07, but neither N6 nor PGDM1400, suppresses HIV$_{REJO.c}$ replication in humanized mice

To explore whether bNAbs delivered via AAV8 as VIT could be used as a long-lived therapeutic strategy for suppressing ongoing HIV replication, we employed the Bone marrow-Liver-Thymus (BLT) humanized mouse model[59]. This model exhibits robust engraftment of human lymphocytes across tissues, enabling sustained infection with HIV[60,61]. BLT mice were infected with HIV$_{REJO.c}$, a clade B, Tier 2, CCR5-tropic, transmitted/founder infectious molecular clone (IMC) of HIV[56]. Prior to treatment, the virus was allowed to propagate and diversify in the host over a period of four weeks (Figure 1A). Groups of BLT mice harboring established HIV$_{REJO.c}$ infections were given a single IM injection of $5x10^{11}$ genome copies (GC) AAV8 expressing either VRC07, N6, or PGDM1400 bNAbs or Luciferase as a negative control. Importantly, all the tested bNAbs potently neutralized HIV$_{REJO.c}$ *in vitro* (Figure S1A, Table S1). After vector administration, antibody expression levels in all mice increased over a period of six weeks (Figure 1B).

For AAV8-VRC07-treated mice, the bNAb expression level plateaued at a geometric mean of ~55 µg/mL in plasma, which was sustained for the remainder of the study; a value approximately 2,000 times over its *in vitro* IC$_{50}$ value (Figure S1B). Infected mice exhibited mean plasma viral loads averaging $5.34x10^{5}$ copies per mL of plasma by week four and sustained viremia following AAV8-Luciferase treatment for the remaining 22 weeks of the study (Figure 1C). In contrast, following AAV8-VRC07 treatment, viral loads decreased precipitously over a three-week period, which diverged

into two outcomes: viral rebound in 56% (30/54) or sustained control in 44% (24/54) of

mice throughout the study (Figure 1D, Figure S1C). Notably, this trend was observed

across five independent experiments, each using a unique donor for each batch of BLT

mice (Table S2).

To explore whether another CD4bs-directed bNAb could achieve long-lived viral

suppression we tested N6[62], an exceptionally broad bNAb with an *in vitro* IC_{50} against

$HIV_{REJO.c}$ similar to that of VRC07 (Figure S1A). In AAV8-N6 treated mice, the plasma

concentrations of N6 achieved a geometric mean of ~15 µg/mL, over 300 times above

the *in vitro* IC_{50} against $HIV_{REJO.c}$ (Figure 1B, Figure S1B). However, AAV8-N6 treatment

only resulted in a transient decline in viral load over the first two weeks, which returned

to pre-treatment levels within four weeks of vector injection (Figure 1E).

Finally, we also tested PGDM1400[63], a somatic variant of the PGT145 antibody

family that recognizes the HIV envelope trimer V1/V2 apex with exceptional potency.

This bNAb neutralized $HIV_{REJO.c}$ with approximately five-fold greater potency than those

measured for VRC07 and N6 (Figure S1A, Table S1). Following AAV8-PGDM1400

administration, expression of PGDM1400 rapidly increased, reaching a geometric mean

plasma concentration of ~28 µg/mL, or nearly 3,700 times above the *in vitro* IC_{50} against

$HIV_{REJO.c}$ (Figure 1B, Figure S1B). However, all mice treated with AAV8-PGDM1400

displayed no significant change in viremia over the 26-week period of observation

(Figure 1F). Comparison of the geometric mean viral loads of all evaluated mice

demonstrated that only those animals receiving AAV8-VRC07 exhibited a significant

decline in viral load compared to the AAV8-Luciferase control. (Figure S1D).

To determine whether differences in viral population structure prior to antibody

treatment might explain the observed outcome, we compared the average diversity

(Shannon entropy) of the HIV *env* gene three weeks after infection, but prior to vector

administration, and noted no significant differences between any of the different groups

(Figure S1E). When comparing rebounding AAV8-VRC07 treated mice to suppressed

mice, we found that mice in each group had indistinguishable expression levels of

VRC07 antibody (Figure S1F), but rebounding mice had a 1.6 times higher viral load

compared to suppressed mice at the time of AAV administration (Figure S1F).

Together, these studies show that only AAV8-VRC07 could suppress actively

replicating $HIV_{REJO.c}$ for an extended duration. In contrast, AAV8-N6 treatment resulted

in a small but significant delay in viral escape as compared to the AAV8-Luciferase

control; while AAV8-PGDM1400 was unable to significantly delay viral escape, despite

comparable viral heterogeneity and higher bNAb potency (Figure 1G).

Antibody escape paths are conserved for each broadly neutralizing antibody

To understand the basis for the observed variation in bNAbs effectiveness

against $HIV_{REJO.c}$, we mapped the escape paths traversed by the virus to acquire

resistance to each individual bNAb. Plasma viral RNA was obtained from each mouse

endpoint and used to amplify viral Env for deep sequencing to analyze non-synonymous

mutations that arose during viral escape. Individual HIV-infected mice receiving AAV8-

Luciferase exhibited mutations distributed throughout the $HIV_{REJO.c}$ envelope (Figure

S2A, Control); however, these mutations followed no discernable pattern on the HIV_{REJO}

env gene (Figure S2B) and did not significantly impact the distribution of potential N-

linked glycosylation sites (PNGs) on this protein (Figure S2C).

Most Env sequences obtained from $HIV_{REJO.c}$-infected animals that escaped from

VRC07 exhibited two pairs of potential escape mutations: either two D-loop mutations

(N276D and D279A) or a D-loop and a V5-loop mutation (D279A and N460D,

respectively) (Figure S2A, VRC07). When assessed collectively across all escaped

mice, these three mutations encompassed the majority of amino acid divergence from

the parental (WT) sequence (Figure 2A). Notably, the N460D mutation also resulted in a

PNGs loss in the V5-loop of $HIV_{REJO.c}$ (Figure S2D).

To determine the impact of these mutations on VRC07 resistance, each

individual mutation and combination of mutations observed in haplotypes were

engineered into $HIV_{REJO.c}$ IMC plasmids to produce replication-competent Env mutant

viruses. Each mutant virus was tested for neutralization sensitivity against VRC07 *in*

vitro (Figure 2B). When assessed individually, neither N276D nor N460D conferred any

resistance to VRC07 neutralization relative to the WT $HIV_{REJO.c}$. However, D279A led to

partial escape by mediating a 51-fold increase in the IC_{50} to 1.4 µg/mL and a reduction

of the dose-response curve slope (Table S1) [64]. Of note, this new IC_{50} still remained

well below the geometric mean serum concentration of VRC07 achieved *in vivo*.

However, complete escape from VRC07 was only observed for envelopes containing

both D279A and either N460D or N276D (Figure 2B).

To measure the fitness cost associated with VRC07 escape mutations emerging

in $HIV_{REJO.c}$, we determined the growth rate of each IMC envelope mutant using

QuickFit, an RT-qPCR-based assay that quantifies viral growth in activated primary

CD4$^+$ T cells across large numbers of replicates (Figure S2E)[65]. The doubling time of

each IMC harboring escape mutations was then normalized to the WT virus to

determine specific fitness costs (or benefits) (Figure 2C). The D279A mutation incurred

a significant fitness cost, whereas no changes were observed for the N276D or N460D

mutations alone. Strains containing paired mutations also exhibited a higher fitness cost

compared to the WT strain. Altogether, these results show that HIV$_{REJO.c}$ follows

conserved evolutionary paths to achieve complete escape from VRC07 with differential

impact on fitness cost and antibody resistance benefit.

The escape paths traversed by HIV$_{REJO.c}$ to acquire resistance to N6 were also

evaluated (Figure S2A, N6). Three mutations in or around the D-loop were the most

frequent (Figure S2F), with one of them resulting in loss of a PNGs (Figure S2G). Unlike

the HIV$_{REJO.c}$ escape paths from VRC07, one mutation on the D-loop, namely A281D,

was sufficient to escape from levels of N6 achieved *in vivo* (Figure S2H, Table S1).

Moreover, when A281D was paired with T278K, resistance increased, whereas pairing

with N276D modestly decreased resistance. Additionally, A281K led to complete

escape from N6, although it required three nucleotide mutations from the parental

sequence (Figure S2H). Using the aforementioned QuickFit assay, the fitness cost of all

identified individual and haplotype combinations of mutations were evaluated (Figure

S2I,J). Similar to the HIV$_{REJO.c}$ VRC07 escape paths, the initial A281D and the

combinations with N276D or T278K, resulted in a higher fitness cost relative to the WT

strain (Figure S2I,J).

Lastly, the escape paths traversed by HIV$_{REJO.c}$ to acquire resistance to

PGDM1400 were evaluated (Figure S2A, PGDM1400). Mutations were observed

surrounding the V1/V2 trimer apex region, with N160K and D167G being the most

prevalent (Figure S2K), with the former resulting in the loss of a PNGs (Figure S2L). As

expected, both of these mutations were sufficient to confer complete escape from

PGDM1400, reaching IC_{50} values above the average level of expression achieved *in*

vivo (Figure S2M, Table S1). N160K and D167G resulted in a fitness benefit compared

to WT strain (Figure S2N,O).

Together, these results show that *in vivo* escape from bNAbs in humanized mice

follows reproducible paths, each with distinct fitness costs and resistance benefits. In

light of the abundance of escape paths, we sought to understand why a given pathway

was traversed as opposed to another.

The Escape Barrier of each bNAb is determined by the fitness landscape of escape

paths

To create an escapability map of the fitness landscape in the context of VIT, we

projected individual escape mutations onto a two-dimensional axis of replicative fitness

cost and resistance benefit (Figure S2P). This was replicated for each evaluated bNAb,

with each point representing an identified escape mutation and arrows representing the

potential escape paths (Figure 2D-F). We categorized each mouse sample into an

escape path by using the viral haplotype frequencies of the dominant escape mutations

(Figure S3A-F).

To escape VRC07 *in vivo*, 81% of the $HIV_{REJO.c}$ sequences harbored the D279A

mutation, conferring partial escape from VRC07, albeit with a high fitness cost (Figure

2D). These escaped viruses also acquired a secondary mutation, typically N460D (Path

A) or N276D (Path B), leading to complete escape from VRC07. Path A escape variants

(D279A + N460D) incurred an even higher fitness cost than the D279A-only escape

variants; whereas path B escape variants (N276D + D279A) improved their fitness

relative to the D279A-only variants. Notably, 19% of the escapes did not harbor the

D279A amino acid change as a partial escape (Path D), but rather found an alternative

escape pathway. These non-canonical escape paths exhibited a significant delay in

time to escape (Figure 2G), suggesting that these viral populations incurred a higher

fitness cost than D279A-based escape paths. To identify the potential forces driving a

given population into a specific path, we compared the VRC07 expression levels (Figure

S3G), the post-escape viral load (Figure S3J), and the total number of accumulated

mutations (Figure S3M) for each path, but failed to find any significant differences

across the paths.

The same analysis was performed for N6 (Figure 2E), where 86% of the escape

variants harbored an A281D mutation, conferring complete escape from N6, with an

IC_{50} 4.5 times greater than the average *in vivo* expression. These A281D variants

typically had a second mutation, either N276D (Path A) or T278K (Path B). Compared

to A281D-only variants, path A escapes were marginally more fit and less resistant to

N6; while path B escapes were both more fit and more resistant to N6 (Figure 2E). A

mix of both path A and path B haplotypes was found within most samples (Figure S3E).

Notably, samples with higher serum concentrations of N6 were associated with lower

post-escape viral loads (Figure 2H), suggesting that N6 may impair viral replication.

This was particularly evident for samples with higher frequencies of N276D (Path A)

(Figure 2I). A small number of N6 escape sequences had three nucleotide mutations

resulting in A281K, which completely escaped N6 with a minor fitness cost (Figure 2E).

There were also some escaped samples harboring only the A281D variant, with no

other prevalent mutations; however, the N6 serum levels of these samples were 6.4-fold

lower compared to those of path A samples (Figure S2H). Of note, given that these

were only two samples, statistical differences could not be assessed. The post-escape

viral load (Figure S2K) and the total number of accumulated mutations (Figure S2N)

were similar across the different paths.

Finally, the PGDM1400 escape paths were clustered based on the haplotypes of

N160K and D167G (Figure 2F, Figure S3C and S3F). N160K (Path A) was found in

56% of the escape variants and the remaining 44% contained D167G (Path B). Both

paths led to complete escape from PGDM1400, with an IC_{50} 2.0- or 2.9-times greater

than the average *in vivo* expression (Figure 2F). We did not observe any significant

difference between the PGDM1400 expression levels and post-escape viral loads for

the two escape paths (Figure S3I and 3L, respectively). However, Path B escapes

accumulated significantly more mutations compared to Path A escapes (Figure S3O).

In order to understand the cumulative fitness cost of $HIV_{REJO.c}$ escape from

VRC07, N6, and PGDM1400, we used the escapability maps to determine Escape

Barrier scores. Briefly, for each escape path on the map, we calculated the area under

the curve (AUC) summing the fitness cost across the IC_{50} values from the WT to the

escaped strain (Figure S4A-C). We then combined the AUC of the top two escape

paths, which accounted for the majority of the observed haplotypes, into a final Escape

Barrier score (Figures S4D-F). Interestingly, the ranking of the Escape Barrier scores for

297 VRC07, N6, and PGDM1400 corresponded with *in vivo* experimental outcomes (Figure

298 2J).

300 *HIV$_{JR-CSF}$ rapidly escapes from bNAbs due to low fitness cost escape mutations*

301   Given the ability of AAV8-VRC07 to reproducibly suppress HIV$_{REJO.c}$ in a subset

302 of mice, we sought to determine whether this could be replicated with another HIV

303 isolate. HIV$_{JR-CSF}$ is a Tier 2, Clade B, CCR5-tropic primary isolate originally obtained

304 from the cerebrospinal fluid of a person living with HIV[55]. Groups of humanized mice

305 were infected with an HIV$_{JR-CSF}$ IMC viral stock four weeks prior to receiving AAV8-

306 Luciferase or AAV8-bNAbs (Table S2). Antibody expression stabilized within four weeks

307 of AAV administration, achieving geometric mean plasma concentrations tens to

308 hundreds of times above the *in vitro* IC$_{50}$ (Figure 3A, Figure S5A,B). In contrast to

309 HIV$_{REJO.c}$, HIV$_{JR-CSF}$-infected mice exhibited no changes in viral load following

310 administration of any of the AAV8-bNAbs evaluated and were indistinguishable from the

311 AAV8-Luciferase controls (Figure 3B, Figure S5C-F).

312   We deep sequenced HIV$_{JR-CSF}$ Env from each mouse to identify mutations

313 associated with escape from VRC07, N6, and PGDM1400 (Figure S6A). Interestingly,

314 mice receiving AAV8-Luciferase exhibited some consistent mutations distributed

315 throughout the HIV$_{JR-CSF}$ envelope despite the absence of bNAb selection pressure,

316 suggesting adaptation to the host (Figure S6B). This was most apparent at position

317 N339 that translated into loss of a PNGs near the V3-loop, which was not observed for

318 HIV$_{REJO.c}$ (Figure S6C). Other mutations around the V2- and V4-loops were also found

319 at higher rates than for HIV$_{REJO.c}$.

To escape VRC07, HIV_{JR-CSF} acquired a single D-loop mutation at position D279,

which was observed across all samples (Figure 3C), and a minority of samples

exhibited a loss of a PNGs at N276 as also seen for $HIV_{REJO.c}$ (Figure S6D). The IC_{50} of

HIV_{JR-CSF} harboring D279 escape mutations were evaluated, with D279A and D279H

achieving complete escape with an IC_{50} greater than 200 µg/mL, while D279N exhibited

a modest increase in IC_{50} relative to the WT strain (Figure S6E, Table S2). D279A

exhibited an increased fitness cost relative to the parental strain, while D279H and

D279N had no statistically significant fitness cost (Figure S6F,G). Unlike $HIV_{REJO.c}$, a

single mutation at site D279 was sufficient to mediate complete escape from VRC07

with a modest fitness cost (Figure 3D), in line with the inability of AAV8-VRC07 to

suppress HIV_{JR-CSF}.

Next, we evaluated the escape paths following AAV8-N6 treatment, finding

selection for a single nucleotide change to yield a A281D mutation (Figure S6H), along

with loss of a PNGs at site N276 (Figure S6I). The A281D mutation was sufficient to

achieve escape (Figure S6J) albeit at a high fitness cost (Figure S6K,L). We also

observed that the combination of A281D with N276D resulted in complete escape at a

lower fitness cost (Figure S6M).

Finally, escape from AAV8-PGDM1400 treatment resulted in a variety of V2-loop

mutations at sites N160 or T162 (Figure S6N), resulting in loss of a PNGs (Figure S6O).

All of the mutations evaluated at sites N160 and T162 resulted in escape from

PGDM1400 (Figure S6P), with no fitness cost (Figure S6Q,R). The resulting

escapability map recapitulates that of $HIV_{REJO.c}$ (Figure S6S).

342   Considering the lack of viral suppression observed for some VIT-treated HIV-

343 infected mice, we evaluated the *in vitro* neutralizing activity of sera from a subset of

344 these animals (Figure S7A-H). As expected, sera from all mice expressing bNAbs were

345 able to neutralize both $HIV_{REJO.c}$ and HIV_{JR-CSF} with an IC_{50} value similar to purified

346 proteins. No neutralization activity was seen in sera from control mice. Given prior

347 reports of antiviral activity of bNAbs following passive transfer in HIV-infected

348 humanized mice[31], we explored whether the kinetics of vectored antibody expression

349 played a role in the lack of antiviral activity seen for N6 and PGDM1400 in our studies.

350 We performed passive transfer of either a control antibody (2A10, a malaria-specific

351 antibody), N6, or PGDM1400 in HIV_{JR-CSF} infected humanized mice (Figure S7I-L and

352 Table S3). As expected, high antibody concentrations were detected starting as early as

353 one week post-infusion, and remained elevated throughout the course of weekly

354 administrations (Figure S7I). Interestingly, we did not observe any change in viral load

355 in any of the mice during the length of the experiment (Figure S7K-L). Despite this, deep

356 sequencing of viral envelopes revealed the selection of escape mutations that largely

357 mirrored those seen for AAV-delivered bNAbs (Figure 7M,N). These results suggest

358 that differences in antibody expression kinetics do not explain the lack of activity seen in

359 our studies. Instead, it is possible that HIV infection in the BLT humanized mouse model

360 may be particularly challenging to suppress relative to other models[31].

362 *Increasing $HIV_{REJO.c}$ mutation resistance benefit enables complete escape from VRC07*

363    Our experiments demonstrate that $HIV_{REJO.c}$ and HIV_{JR-CSF} exhibit

364 differences in escape paths under VRC07 selection. However, these studies could not

discern whether the differences in Env sequence or the inherent fitness differences

between these isolates contributed to escapability. Notably, in the context of the

$HIV_{REJO.c}$ Env, the D279A mutation only resulted in a partial escape, whereas for $HIV_{JR\text{-}CSF}$,

the same mutation resulted in complete escape. To evaluate whether this could be

explained by differences in the diversification of these viruses in humanized mice,

relative to those seen in humans, we compared the baseline frequency of each amino

acid at each position in env for both isolates to the frequencies observed in the

corresponding position in the LANL database. We found that amino acids that were at a

high frequency in $HIV_{REJO.c}$ and $HIV_{JR\text{-}CSF}$ also tended to have a higher frequency in the

LANL database (Figure S8A), suggesting that both isolates generate diverse

quasispecies that are consistent with clinical infections. To determine whether the

fitness cost of escape mutations observed in both viruses were consistent with

conservation in the LANL database, we directly compared the LANL amino acid

frequency of each escape mutation to their observed fitness cost in $HIV_{REJO.c}$ and $HIV_{JR\text{-}CSF}$

backbones, and found a strong and significant negative correlation (Figure S8B),

confirming that high fitness cost mutations in both isolates are also rarely seen in the

LANL database.

To dissect the contribution of the resistance benefit to escape VRC07, we

engineered chimeric IMCs of $HIV_{REJO.c}$ containing the swaps of amino acid sequences

comprising the D-loop (HIV_{RD}), the V5-loop (HIV_{RV}) or both loops (HIV_{RDV}) from $HIV_{JR\text{-}CSF}$

(Figure 3E, Table S4). Each of the three chimeric viruses were neutralized by

VRC07 with a similar IC_{50} to those of the original isolates (Figure 3F). Importantly, both

D-loop chimeras, HIV_{RD} and HIV_{RDV}, exhibited a modest increase in fitness cost,

whereas the V5-loop-only chimera, HIV_{RV}, showed no difference in fitness relative to the

$HIV_{REJO.c}$ strain (Figure S9A,B).

Following infection of BLT humanized mice with each chimeric strain, AAV8-

VRC07 administration resulted in a geometric mean steady-state plasma antibody

concentration of approximately 50 µg/mL for all the groups (Figure S9C). This led to

suppression in a subset of $HIV_{REJO.c}$-infected mice, whereas all HIV_{JR-CSF}-infected mice

escaped (Figure S9D,E). A similar proportion of mice infected with HIV_{RD} and HIV_{RDV}

were suppressed by VRC07, as seen for $HIV_{REJO.c}$ (Figure S9F,G). In contrast, all mice

infected with HIV_{RV} exhibited rapid escape from VRC07, as seen for HIV_{JR-CSF} (Figure

3G, Figure S9H,I).

We focused on the HIV_{RV} chimera to understand why this swap resulted in a

suppression outcome similar to that of HIV_{JR-CSF}. In contrast to $HIV_{REJO.c}$, mutations

arising in HIV_{RV}-infected animals receiving AAV8-VRC07 were confined to the D-loop

(Figure S9J-N). HIV_{RV} mutants harboring single amino acid changes at D279 were

completely escaped from VRC07, albeit with substantial fitness costs (Figure S9O-Q).

Taken together, these data suggest that the sequence context of the HIV

envelope directly influences the path taken to escape from VRC07 (Figure 3H). The

escapability map reveals that D279 mutations result in complete escape for HIV_{RV}, in

contrast to $HIV_{REJO.c}$. Finally, the Escape Barrier score for HIV_{JR-CSF} escaping from all

bNAbs and HIV_{RV} escape from VRC07 was calculated as before (Figure S10A-D). For

HIV_{JR-CSF}, VRC07 exhibited the lowest score; whereas HIV_{RV} score was between

$HIV_{REJO.c}$ and HIV_{JR-CSF} (Figure 3I, Figure S10E,F).

Increasing HIV$_{REJO.c}$ mutation fitness cost results in complete suppression by VRC07

Given the importance of sequence context in defining the resistance benefit of

mutations along the evolutionary escape path, we sought to evaluate the impact of

mutation fitness cost on escape. To this end, we sought to alter the fitness of VRC07-

escape mutants without affecting their resistance benefits. Given the previously

described reduction in viral fitness following escape from antiretroviral therapy (ART)

drug regimens[66], we tested the ability of the Polymerase (Pol) mutations to impact the

fitness of VRC07 escape mutations. We engineered Pol M184I and M184V mutations

into the original HIV$_{REJO.c}$ IMC vector and detected no changes in neutralization

sensitivity to VRC07 or viral fitness (Figure S11A,B). We then evaluated both the fitness

cost and resistance benefit of the previously identified VRC07-HIV$_{REJO.c}$ escape

mutations (i.e., N276D, D279A, N460D, and their combinations) in the context of the Pol

M184V mutation. While we observed no changes in neutralization sensitivity (Figure 2B,

Figure S11C, Table S1), we observed significant increases in the fitness cost for escape

mutants containing Pol M184V relative to WT Pol (Fig 4A, Figure S11D).

The net effect of this was a shift in the escapability map for HIV$_{REJO.c}$ with the Pol

M184V mutation, such that complete escape through a second envelope mutation

incurred substantially higher fitness costs than for the WT Pol strain (Figure 4B). The

Escape Barrier score was calculated for Pol M184V mutants assuming the same

relative escape paths frequencies as seen for the WT Pol, resulting in a 43% increase

(Figure 4C), suggesting that selection of the Pol M184 mutations could improve VRC07

suppression of HIV$_{REJO.c}$.

To evaluate the predicted reduction in $HIV_{REJO.c}$ escape from VRC07, we intentionally drove the emergence of Pol mutations through sub-optimal dosing of ART in humanized mice. BLT mice were infected with $HIV_{REJO.c}$ and over the course of 6 weeks, we titrated the dose of ART drug in their food, from 1% to 100% of the standard human equivalent dose. Eight weeks after infection, either AAV8-VRC07 or AAV8-Luciferase were administered. Two weeks after AAV administration, the ART treatment was interrupted (Figure 4D, Table S5). Importantly, at the time of AAV administration, no meaningful differences in viral loads were seen between the mice that had received escalating antiretroviral drugs relative to control mice (Figure S11E). As expected, HIV *pol* genes sequenced from mice receiving the suboptimal ART dosing displayed a near-complete prevalence of M184V/I at the time of VRC07 administration (Figure 4E).

As anticipated, ART drug treatment had no impact on AAV8-driven expression of VRC07 (Figure S11F). ART-treated mice receiving AAV8-Luciferase maintained steady viral loads, with minor dips at each escalation of drug concentration (Figure 4F). Administration of AAV8-VRC07 without ART treatment resulted in only partial suppression of HIV, as observed in our prior experiments (Figure S11G). Additionally, the post-escape viral load for the Luciferase-treated mice showed no difference irrespective of the presence or absence of the selective pressure (Figure S11H). However, administration of AAV8-VRC07 to ART-treated mice resulted in complete suppression (Figure 4F,G), suggesting that the additional fitness costs were specific to the VRC07 escape mutations, as predicted in our theoretical escape map and Escape Barrier scores. Taken together, these results suggest that by decreasing the overall

455   fitness of $HIV_{REJO.c}$, and therefore modulating the fitness cost axis of the escapability

456   maps, resistance to VRC07 becomes significantly more difficult.

Discussion

Antibody immunotherapy to prevent or treat HIV infection is currently being evaluated in numerous clinical trials. While bNAb combinations have shown promise[42,67–69], individual bNAbs have demonstrated limited activity in viremic PLWH due to their half-life and the emergence of escape mutations[34,35,37,39,70,71]. Here we show that vectored delivery of a single bNAb, well-matched to the infecting strain, is capable of suppressing viremia in approximately half of HIV-infected humanized mice. Notably, this efficacy was not correlated with neutralization potency determined *in vitro* against the infecting viral stock or breadth metrics derived from global pseudovirus panels[17,62,63]. Rather, our results suggest that the *in vivo* efficacy of this treatment is best explained by the bNAb-specific dynamics of viral escape. By using the BLT humanized mouse platform and viral stocks derived from IMCs, we performed dozens of independent infections with a specific virus and AAV8-bNAb treatment. This led to the reproducible characterization of the most commonly traversed escape paths by HIV, demonstrating surprising conservation of viral escape from specific bNAbs.

Of the two CD4bs-targeting antibodies evaluated in this study, N6 was broader, neutralizing 96% of a 181-strain global panel with an IC_{50} of less than 1 μg/mL[62]. Despite its outstanding potency and breadth, we found that a single base pair change in the D-loop of $HIV_{REJO.c}$ or HIV_{JR-CSF} yielded resistance to neutralization that was above the steady-state concentrations achieved in our experiments. Remarkably, this A281 mutation has also been previously reported for antibodies from the VRC01-class family, suggesting conserved escape paths for these antibodies[72,73]. Despite a lack of viral suppression by AAV8-N6, we observed that mice with higher antibody concentrations

exhibited lower post-escape viral loads. As such, a higher steady-state concentration of

N6 would likely result in improved viral suppression; however, expression was lower

relative to other bNAbs despite identical vector dosage, consistent with previous reports

of short N6 half-life following passive transfer[74,75].

PGDM1400 recognizes the HIV envelope V1/V2 trimer apex and neutralizes with

a median IC_{50} of 0.003 µg/mL against a 77-virus panel[63]. A single-point mutation in the

V2-loop (N160 or D167) was enough for both HIV isolates to acquire complete

resistance to PGDM1400, which has been previously observed for trimer apex-targeting

antibodies[76]. The low fitness cost of these mutations is reflected in their high pre-

existing frequency in control mice and PLWH (Table S6) (https://www.hiv.lanl.gov/). In a

clinical trial using PGDM1400, multiple sequences isolated from rebound participants

exhibited a loss of the N160 glycan, suggesting that glycan-based escape is a major

driver of bNAb resistance in humans[68].

VRC07 neutralizes 83% of a panel of 179 strains at less than 1 µg/mL, with a

geometric mean IC_{50} of 0.11 µg/mL[17]. In contrast to N6 and PGDM1400, complete

escape of $HIV_{REJO.c}$ from VRC07 required a two-step evolutionary path. The mutations

we report here include an essential D-loop mutation at D279, resulting in partial

resistance with moderate fitness cost, and a second mutation in either the D- or V5-

loops which confers complete resistance and further modulates the fitness of the virus.

Notably, the tolerability of D- and V5-loop mutations identified through deep mutational

scanning varied significantly across strains, in line with our isolate-specific fitness cost

data[77]. Although it required two steps, $HIV_{REJO.c}$ escape from VRC07 occurred in 44% of

mice across our study. Interestingly, the N276D D279A escape path was observed less

often than the D279A N460D escape path, despite having a lower fitness cost. This may

be a result of the high frequency of pre-existing N460D (Table S6) and the large

distance between these two mutations, which is known to increase the likelihood of

recombination[78]. Of note, D279A, along with the paired mutations at N276 and N460,

have been previously reported as escape mutations from VRC01-class antibodies in

humans[36,73,79]. Finally, $HIV_{REJO.c}$ escapes that did not acquire D279A had a significant

delay in time-to-escape, suggesting that blocking a primary escape path can result in

the emergence of rarer and more costly alternatives. The differences in escape paths

seen for VRC07 suggest that the timing of escape, and particularly how early these

mutations emerge, plays a significant role in the therapeutic success of VIT, and future

analyses should try to dissect this more in depth.

The AMP trials have identified the predicted serum neutralization 80% inhibitory

dilution titer (PT_{80}) as a robust predictor of protection in humans and NHPs[24,80]. This

biomarker is defined as the bNAb serum concentration over the *in vitro* IC_{80} value

against a given HIV strain. A steady-state PT_{80} of 200 or higher correlated with a

prevention efficacy of 90%[24]. Despite reaching average PT_{80} values against $HIV_{REJO.c}$ of

over 500 for VRC07 and over 400 for PGDM1400 over the four week period of

increasing expression, we did not see efficacy against established infections (Figure

S12). Interestingly, there were no differences in PT_{80} values between $HIV_{REJO.c}$

suppressed or escaped mice, demonstrating that in our model, potency- and breadth-

derived parameters were insufficient to predict the *in vivo* efficacy of bNAbs.

Our results show that escape from CD4bs-directed bNAbs is more likely to

impact viral fitness as compared to bNAbs targeting a variable loop, likely due to the

CD4-binding requirement during viral infection[36,81]. In a clinical trial evaluating 3BNC117,

another CD4bs-targeting bNAb, sequences from three out of eight participants

remained sensitive to the bNAb following viral rebound after antibody serum decay[34].

Additionally, patients undergoing ART interruption following 3BNC117 administration

remained suppressed until antibody levels declined, suggesting a lack of pre-existing

escape variants in the latent viral reservoir[72]. Recent characterization of escape from

eCD4-Ig further demonstrates the difficulty of viral escape from CD4bs-targeted

therapeutics[81]. Collectively, these findings highlight the advantage of targeting high

fitness-cost epitopes.

In this study we project the fitness landscape of each bNAb-strain onto two-

dimensional escapability maps by comparing both the fitness cost and resistance

benefit of mutations for each major escape path. Using chimeric strains, we modulated

the neutralization benefit of escape mutations as represented by the X-axis of the

escapability maps. Despite the same neutralization sensitivity to VRC07 and no

difference in fitness compared to $HIV_{REJO.c}$, we found that the HIV_{RV} chimera exhibited

complete and rapid escape from vectored VRC07, like HIV_{JR-CSF}, highlighting the

importance of sequence context on escape mutations[77]. In this study, we utilized two

different HIV isolates and found that the diversity of the quasispecies in infected

humanized mice was similar to the sequence diversity seen in the LANL database, as

reported by other studies that evaluated the conservation of mutations cross-

sectionally[82]. Importantly, this conservation also correlated with the fitness cost of the

mutations, showing that more costly mutations were less represented in the LANL

database. The relevance of the sequence context and diversification could have

implications for studies performed in NHPs, given the chimeric nature of SHIV[83].

Using M184 Pol mutations, we modulated the fitness cost of escape mutations,

as represented by the Y-axis of the escapability maps. We found that M184 Pol viruses

harboring VRC07 escape mutations maintained their susceptibility to VRC07 but grew

more slowly than WT Pol strains *in vitro*. We predicted that this shift in fitness cost

would result in a more challenging escape path from the antibody *in vivo*. Consistent

with this hypothesis, AAV8-VRC07 administration to $HIV_{REJO.c}$-infected animals

harboring these Pol mutations resulted in sustained viral suppression. Together, these

experiments demonstrate that the two escapability map dimensions of resistance

benefit and fitness cost drive the efficacy of AAV-bNAb therapy *in vivo*.

The BLT humanized mouse model enables the systematic comparison of bNAb

therapeutic efficacy and identification of escape paths taken by HIV isolates. Despite

their lack of endogenous humoral immune responses, here we report the same VRC07,

N6, and PGDM1400 escape mutations seen in NHPs and humans, with the benefit of a

shorter time frame and lower cost[84]. Given their diverse and continuously evolving viral

population, as previously reported[31] and seen here (Figure S7), HIV-infected humanized

mice represent an excellent model to evaluate bNAb escape *in vivo*.

The concept of fitness landscapes has been frequently used to understand HIV

evolution, including the dynamics of intrapatient adaptation[85] and antiretroviral escape[86].

Within this framework, it is well known that a fitness cost is typically accrued during

escape from cytotoxic T lymphocytes[87–89], antiretroviral drugs[90–93], or antibodies[94].

Indeed, modern cART leverages this understanding to maximize their clinical efficacy[95].

Our findings suggest that bNAb escape paths are predictable and that maximizing the cost of escape *in vivo* may be the key to improving their clinical efficacy. Whether analogous improvements in the clinical efficacy of bNAbs could be achieved by reducing the fitness of viruses replicating in patients who have failed ART regimens remains to be determined. Recent efforts have focused on antibody combinations that target independent sites of vulnerability, however, future bNAb combinations with orthogonal, high cost, escape paths should be tested, as combinations whose escape mutations result in additive fitness costs may improve the therapeutic efficacy of antibody-based interventions.

Limitations of the study

Despite our best effort to be comprehensive, our study has a number of limitations. First, as previously stated, differences in bNAb expression levels influence the resulting escape paths. Future studies could focus on dissecting the influence of bNAb expression level on escape path selection. In addition, AAV-mediated antibody delivery starts with lower serum levels of bNAb before achieving a high steady-state concentration, whereas passive transfer results in high initial concentration, which then declines over time. This difference could result in divergent outcomes, as passive transfer applies a stronger initial selection pressure, which may alter viral escape dynamics. However, subclinical viral replication has been reported as passively transferred antibody decays, even when still above the currently accepted protective levels[96]. Moreover, at steady-state, the mutations we identified were the same as those which ultimately emerged during passive transfer studies. Second, we report the results of only three antibodies targeting two distinct sites of vulnerability against two clade B isolates of HIV. However, a plethora of bNAbs have been described targeting five sites of vulnerability and many diverse HIV isolates exist [97]. Whether our findings would extend to other bNAbs and HIV clades remains to be determined. However, several reports show that mutations and evolutionary paths are conserved and convergent across different isolates and clades[98–100]. Third, our study focuses on escape mutations present at the endpoint of each independent experiment, therefore, the dynamic events occurring earlier and over the course of escape still need to be elucidated. Fourth, HIV infection in BLT humanized mice could not be assessed beyond six months due to declining health of animals from this model, which may limit the opportunities for escape

mutations to arise. Fifth, humanized mice have a relatively small blood volume and lack

secondary lymphoid organs that may not completely recapitulate HIV infection in

humans[101]. Moreover, BLT mice elicit a poor humoral immune response, and therefore

lack this selective pressure[102], which may influence the selection of escape paths. In the

case of PGDM1400, it has been shown that the polyclonal humoral response elicited in

humans is reactive against the various glycans on the Env protein[103]. Viremic patients

receiving individual bNAbs have exhibited more substantial drops in viral load than what

was seen in our model[34,35,41,70]. Additionally, there have been reports of synergy

between passively transferred bNAbs and endogenous antibody responses in

patients[104], which are unlikely to occur in humanized mice. As such, bNAb-based

suppression of HIV in humanized mice may represent a high bar for therapeutic efficacy

given that suppression is completely dependent on the administered bNAb. Finally,

measurements of viral growth rates were performed *in vitro* with PBMCs from a single

donor. Whether the absolute growth rate of these mutants across genetically distinct

patients varies remains untested, previous studies suggest that their relative fitness

would remain unchanged[65].

Resources availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dr. Alejandro Balazs (abalazs@mgh.harvard.edu).

Materials availability

All genetic constructs and unique biological materials generated in this study are available from the lead contact without restriction upon request.

Data and code availability

A summary of the sequencing analyzed data is available online at https://github.com/Balazs-Lab/Escapability. Raw sequencing files reported in this paper will be shared by the lead contact upon request. All other data are available in the main text or as part of the supplementary data and tables. Source data referencing each figure are also provided for this study as supplementary material. Codes used for the sequencing analyses in this study, along with instructions to run them, can be found in the following link: https://github.com/Balazs-Lab/Escapability

Acknowledgements

We wish to thank Dennis Burton (Scripps) for providing antibody proteins, Dan Barouch for providing anti-PGDM1400 idiotype antibody protein. We also thank K. L. Clayton and D. T. Claiborne for advice on experimental design and the manuscript. We thank the Ragon Institute Protein Core for providing antibody proteins. The following reagents were obtained through the NIH HIV Reagent Program, Division of AIDS, NIAID, NIH: HIV-1, Strain JR-CSF Infectious Molecular Clone (pYK-JRCSF), ARP-2708,

contributed by Dr. Irvin S. Y. Chen and Dr. Yoshio Koyanagi; ARP-11746

(pREJO.c/2864), contributed by Dr. John Kappes and Dr. Christina Ochsenbauer; TZM-

bl Cells, ARP-8129, contributed by Dr. John C. Kappes, Dr. Xiaoyun Wu and Tranzyme

Inc. N.M.S.G. was supported by the Executive Committee on Research (ECOR) of the

Mass General Hospital (MGH) Fund for Medical Discovery (FMD) Fundamental

Research Fellowship Award. A.D.N. was supported by the Ruth L. Kirschstein National

Research Service Award (NRSA) Postdoctoral Training Program Award T32AI007245.

C.E.D. was supported by the NRSA Individual Postdoctoral Fellowship 1F32AI125096-

01A1. J.M.B. was supported by the NSRA Predoctoral Award 1F31AI131747-01A1.

A.B.B. was supported by the National Institute for Allergy and Infectious Disease Career

Transition Award K22AI102769, Research Grants R01AI174875, R01AI174276, the

National Institutes for Drug Abuse (NIDA) Avenir New Innovator Award DP2DA040254,

the NIDA Avant-Garde Award 1DP1DA060607, CDC subcontract 200-2016-91773-

T.O.2, the MGH Transformative Scholars Program as well as funding from the Charles

H. Hood Foundation. This independent research was supported by the Gilead Sciences

Research Scholars Program in HIV.

Author contributions

C.E.D. and A.B.B. designed the experiments. N.M.S.G., A.D.N., Y.C., C.E.D.,

C.L.B., S.W.M., Y.E.S.A., E.C.L., M.L.S., D.P., A.L., M.J.D., S.B.Y., B.M., and C.B.B.

carried out experiments and analyzed data. J.M.B., A.P., X.C., C.L., R.A.K., J.R.M. and

D.L. offered suggestions for experiments and provided key materials. S.T. and V.D.V.

provided humanized mice. N.M.S.G., A.D.N, Y.C., C.E.D. and A.B.B. wrote the paper

with contributions from all authors. The authors of this study fulfill all the authorship

criteria required by Cell Press. Roles and responsibilities were agreed among the

authors ahead of the research within reason and changes on these roles were

previously   agreed   by   the   authors.

Declaration of interests

A.B.B. is a founder of Cure Systems LLC. All other authors declare no conflicts of

interest.

Supplemental information

Document S1. Figures S1–S12 and Tables S1-S6.

Document S2. Excel file containing all source data referring to all the figures.

Document S3. Excel file presenting all the mice, their treatment, antibody

concentration reported at escape, and weeks of suppression (or to escape).

 Figure legends

Figure 1. Vectored delivery of VRC07, but not PGDM1400 or N6, can suppress
established $HIV_{REJO.c}$ infection.
(A) BLT humanized mice were infected with 300 $TCID_{50}$ of $HIV_{REJO.c}$ and the viral
population was allowed to replicate for 4 weeks. Mice were then IM injected with $5x10^{11}$
genome copies (GC) of AAVs encoding for either VRC07, N6, PGDM1400, or
Luciferase as control. Mice were followed for 6 months, and blood samples were
collected weekly.
(B) ELISA-based quantitation of gp120-binding antibodies in the serum of $HIV_{REJO.c}$-
infected humanized mice following administration of AAV8-Luciferase, AAV8-VRC07,
AAV8-N6, or AAV8-PGDM1400 vectors. Black arrow denotes vector administration.
Data are plotted as geometric mean ± geometric SD.
(C-F) HIV viral load in plasma of $HIV_{REJO.c}$-infected mice injected with AAV8-Luciferase
as control **(C)**, AAV8-VRC07 **(D)**, AAV8-N6 **(E)**, or AAV8-PGDM1400 **(F)**. Black arrows
denote vector administration. Each colored line depicts an individual mouse tracked
over time. The sensitivity of qPCR was 1 genome copy per µL of plasma, and 5 µL were
used in the reaction, resulting in a 1000 copy per mL limit of detection (solid line). Data
are presented as mean ± S.E.M.
(G) Kaplan-Meier plot of viral suppression in BLT humanized mice infected with
$HIV_{REJO.c}$ given the indicated bNAb-expressing vector. The total model significance
(p<0.0001) and pairwise comparison against the Luciferase control (****: p<0.0001)
were assessed independently using Log-rank (Mantel-Cox) tests. The percentage of
HIV suppressed was defined as the fraction of mice that were not escaped as described
in the methods.
Figure 2. Fitness cost and resistance benefit accrued along escape paths
determine the difficulty of $HIV_{REJO.}$ escape from each bNAb.
(A) Amino acid divergence from the envelope gene of the $HIV_{REJO.c}$ parental strain
across all AAV8-VRC07 treated mice. Sequences were determined by Illumina Deep
Sequencing of the viral envelope isolated from plasma at the final experimental
timepoint. The X-axis represents the envelope protein amino acid position relative to
HIV_{HXB2} numbering. The Y-axis represents the percentage of average amino acid
divergence from the parental strain, corrected for divergence observed in control mice.
Pie charts represent the most common amino acid mutations for sites with the highest
divergence. See also Figure S2.
(B) *In vitro* neutralization assays of each $HIV_{REJO.c}$ mutant identified in **(A)** against
VRC07. Data are plotted as mean ± S.E.M. Each data point was evaluated in
quadruplicate.
(C) Relative viral growth of each $HIV_{REJO.c}$ mutant identified in **(A)**. Growth rates were
determined in activated $CD4^+$ T cells performing *QuickFit* assays and normalized to the
parental strain. Data are plotted as mean ± S.E.M. and statistical differences were

assessed by a Kruskal-Wallis non-parametric ANOVA with Dunn's *post hoc* test to
correct for multiple comparisons (****: p<0.0001). See also Figure S2.
(D-F) Escapability maps denoting fitness cost (Y-axis, relative doubling time shown in
(**C**)) and resistance benefits (X-axis, neutralization resistance shown in (**B**)) for each
HIV$_{REJO.c}$ mutant observed during escape from VRC07 (**D**), N6 (**E**), or PGDM1400 (**F**).
Dashed vertical line denotes the geometric mean antibody serum concentration after
vectored bNAb administration. Shaded areas represent the neutralization assay limit of
detection. Solid arrows represent the likely initial path taken with dashed arrows
representing the likely second step paths to escape. In the map legend, each escape
path is sorted based on the relative frequency observed in the sequencing data. See
also Figures S2 and S3.
(G) Survival analysis of HIV$_{REJO.c}$-infected mice treated with AAV8-VRC07 by escape
path haplotype. The total model significance (p=0.0126) and pairwise comparisons of
the non-D279A escape paths to each other escape path (*: p<0.05) were assessed
independently using Log-rank (Mantel-Cox) tests. See also Figure S3.
(H) Correlation plot of N6 geometric mean-expression vs post-escape HIV$_{REJO.c}$ viral
load. The line represents the semi-log least squares regression, ρ represents the
Spearman correlation value determined for the data, with the associated p-value below.
See also Figure S3.
(I) Correlation plot of the post-N6 escape viral load of HIV$_{REJO.c}$ vs the frequency of
N276D determined by viral envelope sequencing. The line represents the semi-log least
squares regression, ρ represents the Spearman correlation value determined for the
data, with the associated p-value below. See also Figure S3.
(J) Escape Barrier (AUC) Score denoting the aggregate fitness cost for each escape
path as determined in the escapability maps. This score was calculated by adding the
area under the escapability plot for escape paths A and path B viral escapes from
VRC07, N6**,** and PGDM1400. See also Figure S4.
Figure 3. HIV$_{REJO.c}$ V5-loop restricts escape from VRC07, by decreasing the
resistance benefit of D279 mutations.
(A) ELISA-based quantitation of gp120-binding antibodies in the serum of HIV$_{JR-CSF}$-
infected humanized mice following administration of 5x10^{11} genome copies (GC) of
AAV8-Luciferase, AAV8-VRC07, AAV8-N6, or AAV8-PGDM1400 vectors. Black arrow
denotes vector administration. Data are plotted as geometric mean ± geometric SD.
(B) Kaplan-Meier plot of viral suppression in humanized mice infected with HIV$_{JR-CSF}$
given the indicated bNAb-expressing vector. The total model significance was assessed
using a Long-rank (Mantel-Cox) test (p=0.0714). See also Figure S5.
(C) Amino acid divergence from the envelope gene of the HIV$_{JR-CSF}$ parental strain
across all AAV8-VRC07 treated mice. Sequences were determined by Illumina Deep
Sequencing of the viral envelope isolated from plasma at the final experimental
timepoint. The X-axis represents the envelope protein amino acid position relative to

HIV$_{HXB2}$ numbering. The Y-axis represents the percentage of average amino acid
divergence from the parental strain, corrected for divergence observed in control mice.
Pie chart represents the most common amino acid mutations for the site with the
highest divergence. See also Figure S6.
(D) Escapability map of HIV$_{JR-CSF}$ escape from VRC07. Dashed vertical line denotes the
geometric mean antibody serum concentration after AAV8-VRC07 administration.
Shaded area represents the neutralization assay limit of detection. Arrows represent the
likely path taken to escape. See also Figures S6 and S8.
(E) Alignment of the D-loop and the V5-loop amino acid sequences for HIV$_{REJO.c}$, HIV$_{JR-CSF}$, and the HIV$_{RD}$, HIV$_{RV}$, and HIV$_{RDV}$ chimeras.
765
(F) *In vitro* neutralization of HIV$_{REJO.c}$, HIV$_{RD}$, HIV$_{RV}$, HIV$_{RDV}$, and HIV$_{JR-CSF}$ by VRC07
bNAb. Data are plotted as mean ± S.E.M. Each datapoint was evaluated in
quadruplicate.
(G) Kaplan-Meier plot of viral suppression in humanized mice infected with HIV$_{REJO.c}$,
HIV$_{RV}$, or HIV$_{JR-CSF}$ following vectored VRC07 administration. The total model
significance (p<0.0001) and pairwise comparisons against HIV$_{REJO.c}$ (***: p<0.001) were
assessed independently using Log-rank (Mantel-Cox) tests. See also Figure S7.
(H) Escapability map of HIV$_{RV}$ during escape from VRC07. Lighter square symbols
represent the original HIV$_{REJO.c}$ escape path taken against VRC07. Dashed vertical line
denotes geometric mean bNAb concentrations after vectored VRC07 administration.
Shaded areas represent the limits of detection. Arrows represent the likely path to
escape. See also Figures S7 and S8.
(I) Escape Barrier (AUC) Score quantifying the difficulty of escape from VRC07 for
HIV$_{REJO.c}$, HIV$_{RV}$, and HIV$_{JR-CSF}$. See also Figure S8.
Figure 4. Increasing the fitness cost of HIV$_{REJO.c}$ escape mutations enhances the
efficacy of vectored VRC07.
(A) The fitness of HIV$_{REJO.c}$-VRC07 escape mutations with or without the Pol M184V
mutation were evaluated using *QuickFit*. Data are presented as mean ± SD. Statistical
differences were assessed by a Two-Way ANOVA, with a Šidák *post hoc* test to correct
for multiple comparisons (*: p<0.05; ****: p<0.0001). See also Figure S9.
(B) Escapability map of HIV$_{REJO.c}$-VRC07 escape mutants with or without the Pol M184V
mutation. Solid symbols (diamonds) represent envelope mutations on the Pol M184V
mutant background, while lighter symbols (circles) represent the data for envelope
mutations on the WT Pol background. Shaded areas represent the limits of detection.
Solid arrows represent the likely path taken with dashed arrows representing secondary
steps to escape. See also Figure S9.
(C) Empirical Escape Barrier score for HIV$_{REJO.c}$ escape from VRC07 as compared to
the theoretical Escape Barrier score for HIV$_{REJO.c-PolM184V}$ escape from VRC07.
(D) Experimental setup to determine the impact of Pol M184 mutations on HIV$_{REJO.c}$
escape from VRC07. Humanized mice were infected with HIV$_{REJO.c}$ and then treated

with a sub-optimal ART starting 2 weeks after infection, to select and maintain Pol M184
mutants. At week 8, 5×10^{11} GC of AAV8-VRC07 was injected IM, and the ART regimen
was stopped at week 10. Mice were followed for 6 months, and blood samples were
collected weekly.
(E) Percentage of Pol M184V/I mutations determined from viral sequences isolated from
the plasma of $HIV_{REJO.c}$-infected mice at week 8, prior to AAV8-VRC07 administration.
Statistical differences were assessed by an unpaired two-tailed Student's t-test (****:
p<0.0001). Data are presented as mean ± S.E.M.
(F) $HIV_{REJO.c}$ viral load in plasma of AAV8-VRC07 treated mice. Black arrows denote
vector administration. Each colored line depicts an individual mouse. The qPCR lower
limit of detection was 1 genome copy per µL of plasma and 5µL were used in the
reaction (solid line). Data are presented as mean ± S.E.M.
(G) Kaplan-Meier plot of viral suppression in humanized mice infected with $HIV_{REJO.c}$
with or without ART selection and with or without AAV8-VRC07 administration. The total
model significance (p<0.0001) and pairwise comparisons against the Luciferase +
Selection control (***: p<0.001; ****: p<0.0001) were assessed independently with Log-
rank (Mantel-Cox) tests. See also Figure S9.
813
814
Supplementary figure titles

Figure S1. AAV-VRC07 can suppress $HIV_{REJO.c}$ infection in BLT humanized mice
(Related to Figure 1).
Figure S2. Detailed escape data for $HIV_{REJO.c}$-infected mice treated with VRC07,
N6, and PGDM1400 (Related to Figure 2).
Figure S3. Escape path analysis for $HIV_{REJO.c}$-infected mice treated with various
bNAbs (Related to Figure 2).
Figure S4. Escape Barrier score calculation for the primary $HIV_{REJO.c}$ escape paths
from various bNAbs (Related to Figure 2).
Figure S5. Vectored delivery of VRC07, N6, or PGDM1400 bNAbs fails to suppress
HIV_{JR-CSF} in BLT humanized mice (Related to Figure 3).
Figure S6. Detailed escape data for HIV_{JR-CSF}-infected mice treated with VRC07,
N6, and PGDM1400 (Related to Figure 3).
Figure S7. Vectored antibodies retain neutralizing activity in vivo and passively
transferred bNAbs fail to suppress HIVJR-CSF in the BLT model due to escape.
Figure S8. Comparative analysis of amino acid frequencies in control mice and
LANL database (Related to Figures 1 and 3).
Figure S9. Neutralization, growth rate, and infectivity of chimeric $HIV_{REJO.c}$ viruses
(Related to Figure 3).
Figure S10. Escape Barrier scores for HIV_{JR-CSF} escape paths from various bNAbs
and HIV_{RV} escape paths from VRC07 (Related to Figure 3).

Figure S11. Characterization of HIV$_{REJO.c}$ strains harboring ART-escape mutations
during VRC07 escape (Related to Figure 4).
Figure S12. Fold change over IC50 and calculated PT80 values for HIV$_{REJO.c}$- and
HIV$_{JR-CSF}$-infected humanized mice expressing vectored bNAbs (Related to
Figures 1 and 3).

Methods

BLT Humanized Mice. BLT humanized mice were generated by the Human Immune System Mouse Program at the Ragon Institute of MGH, MIT, and Harvard. Briefly, 6- to 8-week-old female NSG mice were transplanted with human liver and thymus tissue under the kidney capsule and injected intravenously with 100,000 CD34$^+$ cells isolated from liver tissue by AutoMACS (Miltenyi Biotec, Cat#130-100-453). Mice were rested for 10 weeks after surgery to allow for recovery and engraftment. All experiments were done with approval from the Institutional Animal Care and Use Committee of the MGH and conducted in accordance with the guidelines and regulations of the American Association for the Accreditation of Laboratory Animal Care.

HIV production. HIV was produced by transient transfection of HEK 293T/17 cells (ATCC, Cat#CRL-11268 - RRID:CVCL_1926) using 25K MW Linear Polyethyleneimine (PEI, Polysciences Inc., Cat#23966) maintained in DMEM medium (Corning, Cat# 10-013-CV) supplemented with 10% fetal bovine serum (FBS - VWR, Cat#89510-186), and 1% penicillin–streptomycin mix (Corning, Cat#30-002-CI) with infectious molecular clone (IMC) plasmids encoding for HIV$_{REJO.c}$ or HIV$_{JR-CSF}$ (AIDS Reagent Program NIH - BEI Resources Cat#ARP-11746 and #ARP-2708, respectively) or IMC plasmids containing the indicated mutations. After 48 hours, culture supernatants were collected, filtered through a 0.45-m filter, and titered using either an HIV-1 p24 antigen capture assay (AIDS and Cancer Virus Program, Leidos Biomedical Research, Inc., Frederick National Laboratory for Cancer Research) or a 50% tissue culture infective dose

($TCID_{50}$) assay on TZM-bl cells (BEI Resources, Cat#ARP-8129 - RRID:CVCL_B478).

$TCID_{50}$ was calculated using the Spearman-Karber formula[105].

Humanized mouse HIV infection. Prior to HIV infection, blood samples were obtained

from mice and subjected to flow cytometry to determine the baseline $CD3^+$ (BioLegend,

Cat#300434 - RRID:AB_10962690), $CD4^+$ (BioLegend, Cat#300534 - RRID:

AB_2563791), and $CD8^+$ (BioLegend, Cat#300554 - RRID: AB_2564382) T cells

engraftment levels. The next day, mice were intravenously infected with either 10ng of

p24 or 300 $TCID_{50}$ of $HIV_{REJO.c}$ or HIV_{JR-CSF} diluted in PBS (Corning, Cat#21-031-CV) to

a volume of 50 ⌈L. Blood was collected weekly to determine viral loads and bNAb levels

in serum.

AAVs vector production, quantification, validation, and administration. AAV8

vectors encoding either Luciferase, 2A10, or bNAbs were produced and validated as

previously described[12]. Briefly, HEK293T/17 cells were co-transfected with the AAV

backbone vector and helper vectors pHELP and pAAV 2/8 SEED using PEI. AAVs were

collected over a five day period following transfection, filtered through a 0.22 µm filter

(Corning, Cat#431097), and fresh media was gently added back to the cells each time.

After collection, the virus was Polyethylene glycol 8,000 (PEG, VWR, Cat#JTU222-9)

precipitated on ice O.N., and then pelleted at 8,000 x g for 30 min. Pellets were re-

suspended in cesium chloride, split evenly into two Quick-Seal tubes (Beckman,

Cat#342413) and centrifuged at 330,000 x g at 20 °C for 24 h (Beckman Coulter, Optima

LE-80K, 70Ti rotor). AAV-containing fractions were determined with a refractometer,

with refractive indexes between 1.3755 and 1.3655 considered positive. These were

then diluted into 15l mL of Test Formulation Buffer 2 (TFB2, 100 mM sodium citrate

(VWR, Cat#Cat#6132-04-3), 10 mM Tris, pH 8.00 (Fisher Scientific, Cat#S1519-

500GM)), loaded in a 100 kDa MWCO centrifugal filter (Millipore, Cat#UFC910024) and

centrifuged at 500 x g at 4 °C until 1 mL remained in the filter. This wash was repeated

twice. Final retentate was aliquoted and stored at −80 °C. Purified AAVs were quantified

by qPCR using the PerfeCTa SYBR Green SuperMix, Low ROX (Quanta Biosciences,

Cat#95056-500) and primers designed against the CMV enhancer

(AACGCCAATAGGGACTTTCC and GGGCGTACTTGGCATATGAT). Samples were

run in duplicate on an QuantStudio 12K Flex (Applied Biosystems). To validate the

functional activity of each lot, *in vitro* transduction assays were performed in

HEK293T/17 cells. Six days after transduction, supernatants were recovered and

quantified for total IgG production by ELISA. AAV8 IM injections were performed as

previously described[12]. Briefly, aliquots of previously titered viruses were thawed on ice

and diluted in PBS to achieve the predetermined dose in a 40 ⌠L volume. A single 40 ⌠L

injection was administered into the gastrocnemius muscle of BLT humanized mice with

a 28 G insulin syringe.

Antibody quantification by ELISA. Plasma was used to determine bNAb

concentrations. For detection of gp120-binding IgG, ELISA plates were coated with 2

⌠g/mL of HIV gp120 protein (Novus, Cat#NBP1-76371 or Immune Technology, Cat#IT-

001-0025p) per well for 1h at room temperature. For detection of PGDM1400, ELISA

plates were coated with BG505 SOSIP (provided by the Vaccine Research Center) at 5

⌠g/mL or a PGDM1400-specific idiotype (provided by Dan Barouch) at 1 ⌠g/mL for 2hr

at room temperature. Plates were blocked with 1% BSA (LGC Clinical Diagnostics,

Cat#50-61-00) in Tris-buffered saline (TBS - Thermo Scientific, Cat#AAJ75892AE)

overnight at 4°C. Samples were incubated in TBS plus Tween 20 (Thermo Scientific,

Cat#BP-337-100) containing 1% BSA for 1h at room temperature before incubation with

1:2,500 to 1:10,000-diluted horseradish peroxidase (HRP)-conjugated goat anti-human

IgG-Fc antibody (Bethyl, Cat#A80-104A - RRID: AB_67064) for 30 min at room

temperature. Samples were detected by the TMB Microwell Peroxidase substrate

system (SeraCare, Cat#50-76-00). A standard curve was generated using purified

VRC07, N6, or PGDM1400 (provided by the Vaccine Research Center) as appropriate

for the sample.

Viral load test by quantitative RT-PCR (RT-qPCR). Viral RNA was extracted from

plasma samples using the QIAamp viral RNA mini kit (Qiagen, Cat#52906)). Each RNA

sample was treated with 2 U of Turbo DNase (Fischer Scientific, Cat#AM2239) at 37°C

for 30 min followed by heat inactivation at 75°C for 15 min. 10 ⌈L of the treated RNA

were used in a 20 ⌈L RT-qPCR reaction with the qScript XLT one-step RT-qPCR Tough

Mix, low ROX mix (Quanta Biosciences, Cat#95134-500), a TaqMan probe (5'-/56-

FAM/CCCACCAAC/ZEN/AGGCGGCCTTAACTG/3IABkFQ/-3') (IDT) and primers

designed targeting the Pol gene of $HIV_{REJO.c}$ (CAATGGCCCCAATTTCATCA and

GAATGCCGAATTCCTGCTTGA) or HIV_{JR-CSF} (CAATGGCAGCAATTTCACCA and

GAATGCCAAATTCCTGCTTGA). Samples were run in triplicate on a QuantStudio 12K

Flex (Applied Biosystems). The following cycling conditions were used: 50°C for 10 min,

95°C for 3 min followed by 55 cycles of 95°C for 3s and 60°C for 30s. Virus titer was

determined by comparison with a standard curve generated using RNA extracted from

serially diluted mixture of commercially titered viral stock and pure mouse serum. The

limits of detection were 1,000 copies per mL for all viral strains. For the purpose of

generating Kaplan-Meier curves, viral escape was defined as the first week after AAV

administration in which viral load did not decrease at least 75% relative to the prior

week and was above 10^4 copies per mL, provided that a subsequent week was also

above that value. Curves were analyzed for statistical significance using the Mantel-Cox

Log-rank test in Graphpad Prism v10.2.2 (RRID:SCR_002798).

Illumina deep sequencing and identification of HIV envelope mutants. Viral RNA

extracted from blood samples at the conclusion of each study was used to synthesize

cDNA using the SuperScript IV Reverse Transcriptase enzyme with a strain-specific 3'

primer for $HIV_{REJO.c}$ (TTGGTACTTGTGATTGCTCCATGTCTCTCC) or HIV_{JR-CSF}

(CCCTATCTGTTGCTGGCTCAGCTCGTC). cDNA was subjected to nested PCR

amplification with HIV-specific envelope primers that yielded a 2.5-kb fragment. The

first-round primers used for amplification from $HIV_{REJO.c}$ were

GCAATAGTAGCATTAGTAATAGCAGGAATAATAGCAATAGTTGTGTGG and

CTGCTCCACCCCCTCTG; whereas the primers used for envelope amplification from

HIV_{JR-CSF} were GCAATAATTGTGTGGTCCATAGTACTCATAGAATATAGGA and

CCCTATCTGTTGCTGGCTCAGCTCGTC. First-round PCR was performed with 1 to

2.5 μL from cDNA reaction using 1x Q5 reaction buffer, 5 mM dNTPs, 0.5 ⌈M of strain

specific primers and 0.02U/⌈L of Q5 Hot Start High-Fidelity DNA Polymerase (NEB,

Cat#M0493L) in a total reaction volume of 25 ⌈L. PCR conditions for the first round

were: 98°C for 30s, followed by 30 cycles of 98°C for 10s, 70°C for 30s, 72°C for 3 min,

with a final extension of 72°C for 2 min. 1⌈L of first round PCR product was then used

as the template for the second round PCR with identical cycling conditions and PCR mix

except for the primers. The second-round primers used for envelope amplification from

$HIV_{REJO.c}$ were AAAATAGACAGGTTAATTGATAGAATAAGAGATAGAGCAGAAGACA-

GTG and TCATTCTTTCCCTTACAGCAGGCCATC; whereas the primers used for

envelope amplification from HIV_{JR-CSF} were AAAATAGATAGGTTAATTGATAAAATAAG-

AGAGAGAGCAGAAGACAG and TCATTCTTTCCCTTACAGTAGACCATCCAGGC.

PCR products were gel extracted, diluted to 0.15 ng/⌈L in UV-irradiated water and

subjected to Nextera XT Illumina library preparation (Illumina, Cat#FC-131-1096). PCR

products were quantified using Qubit and D5000 ScreenTape System (Agilent,

Cat#5067-5589). The library pool was denatured with 0.2 N NaOH, diluted to 4 nM,

spiked with 10% PhiX to improve sequence heterogeneity and quality and subjected to

2 x 250 or 2 x 300 paired-end sequencing on the Illumina MiSeq (RRID:SCR_016379).

Envelope Escape Mutant Analysis. Sequencing reads were filtered for quality using

fastp (RRID: SCR_016962)[106] and aligned to a reference sequence specific to the viral

strain under analysis using Bowtie2 (RRID: SCR_016368)[107]. The alignments were

sorted and indexed using samtools (RRID: SCR_002105)[108]. Amino acid changes were

called using a custom codon aware variant caller, written in python (RRID:

SCR_008394). For each sample, the total divergence at each amino acid site was

determined by summing the frequency of every non-WT amino acid (relative to the IMC

strain sequence, meaning that synonymous mutations are not considered divergent).

The group amino acid divergence (plotted in the figures) was determined by averaging

the amino acid divergence across each sample at every site. Each site was numbered

using the HIV_{HXB2} nomenclature[109]. The analysis pipeline was run using snakemake

(RRID: SCR_003475)[110]. All the bioinformatics tools used in these analyses are

available online at https://github.com/Balazs-Lab/Escapability.

Viral Diversity and Variant Conservation Analysis. Similar to the envelope escape

mutation analysis, sequencing reads were filtered for quality using fastp[106] and aligned

to a reference sequence specific to the viral strain under analysis using Bowtie2[107]. The

alignments were sorted and indexed using samtools[108]. SNVs were called using

LoFreq[111]. The heterogeneity of each sample was determined using the SNV data to

calculate the average Shannon entropy across all sites in the viral envelope, as

performed previously[112,113]. The Shannon entropy at each site in the genome was

calculated by taking the sum of the frequency of each SNV times the natural log of the

SNV frequency (frequency * ln(frequency)), multiplied by negative one. The analysis

pipeline was run using snakemake[110] and is available online at

https://github.com/Balazs-Lab/Escapability. For the comparison of viral diversification to

the LANL database, the HIV1 FLT 2022 Env Protein Alignment was downloaded from

LANL and filtered for subtype B sequences. All positions were numbered based on their

HIV_{HXB2} alignment and only positions shared across $HIV_{REJO.c}$, HIV_{JR-CSF} , and the LANL

database were analyzed. The average frequency of each $HIV_{REJO.c}$ and HIV_{JR-CSF} amino

acid was binned by rounding the log_{10} transformed frequency using the MROUND()

function in excel. These grouped bins were then used to evaluate the LANL amino acid

frequency data, of which the average and standard deviation were calculated using the

Excel Pivot Table functions.

1004

Construction of HIV mutants. Individual mutations were introduced into the parental

IMC vectors expressing the molecular clone using overlapping PCR with primers

incorporating the desired mutations. After amplification of the *env* or *pol* gene with the

mutagenesis primers using KOD Hot Start Master mix (EMD Millipore, Cat#71975-3) or

Q5 Hot Start Master Mix (NEB, Cat#M0493L), the PCR product was purified by gel

extraction (Promega, Cat#A9282) and cloned by homologous recombination into the

appropriate recipient parental backbone vector using the In-Fusion HD cloning kit

(Clontech, Cat#639650). The ligation product was transformed into DH5⟨ (Zymo,

Cat#T3009) or SURE2 (Agilent, Cat#200152) competent cells, and positive clones were

full plasmid sequenced.

1015

In vitro **neutralization assay.** To compare the sensitivity of point-mutant viruses to

bNAb antibody neutralization, each mutant was produced by transient transfection of

HEK 293T cells as described above and viral supernatants were titered by $TCID_{50}$ on

TZM-bl cells. Then, neutralization assays were performed using a TECAN Fluent 780

liquid handler by mixing 20 µL of virus with 20µL of 2.5-fold serial dilutions of each

antibody and incubating this mixture at 25°C for 1 h. After the incubation, antibody-virus

mixtures were added to previously plated 6,000 TZM-bl cells with 75 ⌠g per mL of DEAE

dextran (Sigma, Cat#D9885-10G) and incubated at 37°C for 48 h. Cells were then lysed

using luciferin-containing buffer[114] and Luciferase signal was quantified using a

PHERAstar FSX plate reader (BMG LabTech). Percentage of infection was determined

by calculating the difference in luminescence between test wells (cells with virus and

antibody) and cell control wells (cells only) and dividing this value by the difference

between the virus control wells (cells with virus) and the cell control wells. These values

were plotted against antibody concentrations and fitted into a four-parameter nonlinear

regression to calculate IC_{50} and Hill Slope using GraphPad Prism v10.2.2.

1031

1032 **Determination of viral fitness.** *In silico* viral growth curves were generated with growth

1033 rates derived from *in vitro* QuickFit assays as described previously[65]. Briefly,

1034 commercially acquired human PBMCs (AllCells, Cat#PB004F; Lot #A2857) were

1035 thawed and $CD4^+$ T cells were isolated using the EasySep™ Human $CD4^+$ T Cell

Isolation Kit (STEMCELL Technologies, Cat#17952). Naive CD4$^+$ T cells were resuspended in complete RPMI 1640 (Corning, Cat#36750) (cRPMI; 10% FBS and 1% penicillin/streptomycin) supplemented with 10 ng/mL recombinant IL-2 (R&D Systems, Cat#202-IL-050) and 4 µg/mL of anti-CD28 antibody (Biolegend, Cat#302934 - RRID: AB_11148949), plated in 24-well plates coated with 2 µg/mL of anti-CD3 antibody (Biolegend, Cat#317326 - RRID: AB_11150592), and incubated at 37ºC and 5% CO_2 for 4 days. Cells were then pooled and incubated for another 4 days before use. Purification and activation efficiency were evaluated by flow cytometry. Previously titered viruses were three-fold serially diluted and then added to 50 ⌈L of activated CD4$^+$ (1×10^5 cells per well) plated in a 96 round-well plate. Viruses and cells were spinoculated at 1,200 RPM for 1 hour at 20°C, and then incubated for 24 hours at 37ºC with 5% CO_2. Cells were washed five times with 200⌈L of cRPMI, resuspended in 200 ⌈L of fresh cRPMI plus 10 ng/mL of IL-2 and finally transferred to a 96 flat-well plate. Plates were incubated at 37°C with 5% CO_2 for 6 days. 32 ⌈L of supernatant were collected daily and fresh media was added to replace the volume. Collected supernatants were immediately RNA extracted using QuickExtract DNA Extraction Solution (Biosearch Technologies, Cat#QE09050)[65]. Extracted RNA was used to determine viral loads by RT-qPCR as stated above. Viral loads were used to determine growth rates and generate *in silico* growth curves using a half-maximal equation in MonolixSuite 2023R1 (Lixoft).

Escape Barrier Analysis. Sample specific sequencing data from the Envelope Escape Mutant Analysis was used to generate sample specific escape haplotypes. Sequences

from each escaped sample were filtered for a minimum mutation frequency of at least

10%, and then manually classified into an escape path using the decision tree

algorithms described in Figure S3 and Figure S8. The area under the curve (AUC) for

each escape path was calculated from the IC_{50} values and relative fitness cost data

using the trapz function in the pracma[115] package in R (RRID:SCR_001905)[116]. The

calculation began at the WT coordinate for each virus and summed the AUC up to the

position of the escaped haplotype. If the escape IC_{50} was greater than 200 µg/mL (the

limit of detection), then the X-coordinate for IC_{50} was assigned a value of 200 µg/mL,

otherwise the IC_{50} of the escape haplotype was used. For all samples (except the HIV_{RV}

escape) only escape paths A and B were used for the Escape Barrier score analysis

because they represented at least 50% of the escape paths. The raw AUC value for

each path was scaled by the relative proportions of the paths (Relative Fraction value in

Figure S4D and S8E). The path scaled AUC scores were then summed to create the

Escape Barrier Score. The scripts and coordinate files are available at

https://github.com/Balazs-Lab/Escapability.

1074

1075   **Passive transfer experiments.** Humanized BLT mice were infected with HIV_{JR-CSF} as

1076   described above, and 4 weeks later were passively infused with 25µg per gram of body

1077   weight of a control antibody (2A10), N6 or PGDM1400, weekly for 12 weeks. Blood was

1078   collected weekly to determine viral loads and bNAb levels in serum.

1079

In vivo **antiretroviral selection-pressure of HIV.** To select for ART-escaping HIV

mutations, individual tablets of emtricitabine (Emtriva[TM] (FTC), Gilead Sciences) or

Tenvir (tenofovir disoproxil fumarate (TDF); Cipla LTD) were crushed into a fine powder

and manufactured with TestDiet 5B1Q feed (Modified LabDiet 5058 with 0.12%

amoxicillin) into powder. The final concentration of these drugs in the stock food was

2.3% (4500 mg/kg TDF, 3000 mg/kg FTC)[28]. To achieve a comparable human dose

(i.e., 200 mg FTC, 300 mg TDF), the Reagan-Shaw formula[117] was used to translate the

dose from human to mouse with the assumption that an average mouse weighs 20 g

and a human weighs 60 kg. On average mice ate 2 g of food per day and the powdered

ART-food was diluted in normal TestDiet 5B1Q food to achieve ingestion of the

corresponding target human dose per day. BLT mice were infected with $HIV_{REJO.c}$ as

previously stated and over the course of 6 weeks, we titrated the dose of ART drug in

their food, from 1% to 100% of the standard human equivalent dose (1% for weeks 2

and 3, 10% for weeks 4 through 6, 100% until week 10). At week 8, mice were injected

with AAV expressing VRC07 or Luciferase as stated above. At week 10, ART treatment

was interrupted. Blood samples were collected weekly to evaluate viral loads and

antibody expression.

References

References

1. Landovitz, R.J., Scott, H., and Deeks, S.G. (2023). Prevention, treatment and cure of HIV
infection. Nat Rev Microbiol *21*, 657–670. https://doi.org/10.1038/s41579-023-00914-1.

2. Bekker, L.-G., Das, M., Abdool Karim, Q., Ahmed, K., Batting, J., Brumskine, W., Gill, K.,
Harkoo, I., Jaggernath, M., Kigozi, G., et al. (2024). Twice-Yearly Lenacapavir or Daily F/TAF
for HIV Prevention in Cisgender Women. N Engl J Med *391*, 1179–1192.
https://doi.org/10.1056/NEJMoa2407001.

3. Jewell, B.L., Mudimu, E., Stover, J., Ten Brink, D., Phillips, A.N., Smith, J.A., Martin-Hughes,
R., Teng, Y., Glaubius, R., Mahiane, S.G., et al. (2020). Potential effects of disruption to HIV
programmes in sub-Saharan Africa caused by COVID-19: results from multiple
mathematical models. The Lancet HIV *7*, e629–e640. https://doi.org/10.1016/S2352-
3018(20)30211-3.

4. McCoy, L.E., and Burton, D.R. (2017). Identification and specificity of broadly neutralizing
antibodies against HIV. Immunological reviews *275*, 11–20.
https://doi.org/10.1111/imr.12484.

5. Sajadi, M.M., Dashti, A., Rikhtegaran Tehrani, Z., Tolbert, W.D., Seaman, M.S., Ouyang, X.,
Gohain, N., Pazgier, M., Kim, D., Cavet, G., et al. (2018). Identification of Near-Pan-
neutralizing Antibodies against HIV-1 by Deconvolution of Plasma Humoral Responses.
Cell. https://doi.org/10.1016/j.cell.2018.03.061.

6. deCamp, A., Hraber, P., Bailer, R.T., Seaman, M.S., Ochsenbauer, C., Kappes, J., Gottardo,
R., Edlefsen, P., Self, S., Tang, H., et al. (2014). Global Panel of HIV-1 Env Reference Strains
for Standardized Assessments of Vaccine-Elicited Neutralizing Antibodies. J Virol *88*, 2489–
2507. https://doi.org/10.1128/JVI.02853-13.

7. Sok, D., and Burton, D.R. (2018). Recent progress in broadly neutralizing antibodies to HIV.
Nat Immunol *19*, 1179–1188. https://doi.org/10.1038/s41590-018-0235-7.

8. Wu, X., Zhang, Z., Schramm, C.A., Joyce, M.G., Kwon, Y.D., Zhou, T., Sheng, Z., Zhang, B.,
O'Dell, S., McKee, K., et al. (2015). Maturation and Diversity of the VRC01-Antibody
Lineage over 15 Years of Chronic HIV-1 Infection. Cell *161*, 470–485.
https://doi.org/10.1016/j.cell.2015.03.004.

9. Diskin, R., Scheid, J.F., Marcovecchio, P.M., West, A.P., Klein, F., Gao, H., Gnanapragasam,
P.N.P., Abadir, A., Seaman, M.S., Nussenzweig, M.C., et al. (2011). Increasing the Potency
and Breadth of an HIV Antibody by Using Structure-Based Rational Design. Science.
https://doi.org/10.1126/science.1213782.

10. Doria-Rose, N.A., Bhiman, J.N., Roark, R.S., Schramm, C.A., Gorman, J., Chuang, G.-Y.,
Pancera, M., Cale, E.M., Ernandes, M.J., Louder, M.K., et al. (2016). New Member of the
V1V2-Directed CAP256-VRC26 Lineage That Shows Increased Breadth and Exceptional

Potency. J Virol *90*, 76–91. https://doi.org/10.1128/JVI.01791-15.

11. Balazs, A.B., Ouyang, Y., Hong, C.M., Chen, J., Nguyen, S.M., Rao, D.S., An, D.S., and
Baltimore, D. (2014). Vectored immunoprophylaxis protects humanized mice from
mucosal HIV transmission. Nature Medicine *20*, 296–300.
https://doi.org/10.1038/nm.3471.

12. Balazs, A.B., Chen, J., Hong, C.M., Rao, D.S., Yang, L., and Baltimore, D. (2011). Antibody-
based protection against HIV infection by vectored immunoprophylaxis. Nature *481*, 81–
84. https://doi.org/10.1038/nature10660.

13. Deruaz, M., Moldt, B., Le, K.M., Power, K.A., Vrbanac, V.D., Tanno, S., Ghebremichael,
M.S., Allen, T.M., Tager, A.M., Burton, D.R., et al. (2016). Protection of Humanized Mice
From Repeated Intravaginal HIV Challenge by Passive Immunization: A Model for Studying
the Efficacy of Neutralizing Antibodies In Vivo. The Journal of Infectious Diseases *214*,
612–616. https://doi.org/10.1093/infdis/jiw203.

14. Stoddart, C.A., Galkina, S.A., Joshi, P., Kosikova, G., Long, B.R., Maidji, E., Moreno, M.E.,
Rivera, J.M., Sanford, U.R., Sloan, B., et al. (2014). Efficacy of broadly neutralizing
monoclonal antibody PG16 in HIV-infected humanized mice. Virology *462-463C*, 115–125.
https://doi.org/10.1016/j.virol.2014.05.036.

15. Veselinovic, M., Neff, C.P., Mulder, L.R., and Akkina, R. (2012). Topical gel formulation of
broadly neutralizing anti-HIV-1 monoclonal antibody VRC01 confers protection against
HIV-1 vaginal challenge in a humanized mouse model. Virology *432*, 505–510.
https://doi.org/10.1016/j.virol.2012.06.025.

16. Brady, J.M., Phelps, M., MacDonald, S.W., Lam, E.C., Nitido, A., Parsons, D., Boutros, C.L.,
Deal, C.E., Garcia-Beltran, W.F., Tanno, S., et al. (2022). Antibody-mediated prevention of
vaginal HIV transmission is dictated by IgG subclass in humanized mice. Sci. Transl. Med.
14, eabn9662. https://doi.org/10.1126/scitranslmed.abn9662.

17. Rudicell, R.S., Kwon, Y.D., Ko, S.-Y., Pegu, A., Louder, M.K., Georgiev, I.S., Wu, X., Zhu, J.,
Boyington, J.C., Chen, X., et al. (2014). Enhanced potency of a broadly neutralizing HIV-1
antibody in vitro improves protection against lentiviral infection in vivo. Journal of Virology
88, 12669–12682. https://doi.org/10.1128/JVI.02213-14.

18. Julg, B., Liu, P.-T., Wagh, K., Fischer, W.M., Abbink, P., Mercado, N.B., Whitney, J.B.,
Nkolola, J.P., McMahan, K., Tartaglia, L.J., et al. (2017). Protection against a mixed SHIV
challenge by a broadly neutralizing antibody cocktail. Science translational medicine *9*,
eaao4235. https://doi.org/10.1126/scitranslmed.aao4235.

19. Julg, B., Tartaglia, L.J., Keele, B.F., Wagh, K., Pegu, A., Sok, D., Abbink, P., Schmidt, S.D.,
Wang, K., Chen, X., et al. (2017). Broadly neutralizing antibodies targeting the HIV-1
envelope V2 apex confer protection against a clade C SHIV challenge. Science translational

medicine *9*, eaal1321. https://doi.org/10.1126/scitranslmed.aal1321.

20. Julg, B., Sok, D., Schmidt, S.D., Abbink, P., Newman, R.M., Broge, T., Linde, C., Nkolola, J.,
Le, K., Su, D., et al. (2017). Protective Efficacy of Broadly Neutralizing Antibodies with
Incomplete Neutralization Activity against Simian-Human Immunodeficiency Virus in
Rhesus Monkeys. Journal of Virology *91*, e01187-17. https://doi.org/10.1128/JVI.01187-
17.

21. Moldt, B., Rakasz, E.G., Schultz, N., Chan-Hui, P.-Y., Swiderek, K., Weisgrau, K.L.,
Piaskowski, S.M., Bergman, Z., Watkins, D.I., Poignard, P., et al. (2012). Highly potent HIV-
specific antibody neutralization in vitro translates into effective protection against
mucosal SHIV challenge in vivo. Proceedings of the National Academy of Sciences of the
United States of America *109*, 18921–18925. https://doi.org/10.1073/pnas.1214785109.

22. Corey, L., Gilbert, P.B., Juraska, M., Montefiori, D.C., Morris, L., Karuna, S.T., Edupuganti,
S., Mgodi, N.M., deCamp, A.C., Rudnicki, E., et al. (2021). Two Randomized Trials of
Neutralizing Antibodies to Prevent HIV-1 Acquisition. N Engl J Med *384*, 1003–1014.
https://doi.org/10.1056/NEJMoa2031738.

23. Mayer, K.H., Seaton, K.E., Huang, Y., Grunenberg, N., Isaacs, A., Allen, M., Ledgerwood,
J.E., Frank, I., Sobieszczyk, M.E., Baden, L.R., et al. (2017). Safety, pharmacokinetics, and
immunological activities of multiple intravenous or subcutaneous doses of an anti-HIV
monoclonal antibody, VRC01, administered to HIV-uninfected adults: Results of a phase 1
randomized trial. PLoS medicine *14*, e1002435.
https://doi.org/10.1371/journal.pmed.1002435.

24. Gilbert, P.B., Huang, Y., deCamp, A.C., Karuna, S., Zhang, Y., Magaret, C.A., Giorgi, E.E.,
Korber, B., Edlefsen, P.T., Rossenkhan, R., et al. (2022). Neutralization titer biomarker for
antibody-mediated prevention of HIV-1 acquisition. Nat Med *28*, 1924–1932.
https://doi.org/10.1038/s41591-022-01953-6.

25. Badamchi-Zadeh, A., Tartaglia, L.J., Abbink, P., Bricault, C.A., Liu, P.-T., Boyd, M., Kirilova,
M., Mercado, N.B., Nanayakkara, O.S., Vrbanac, V.D., et al. (2018). Therapeutic Efficacy of
Vectored PGT121 Gene Delivery in HIV-1-Infected Humanized Mice. Journal of Virology *92*,
118. https://doi.org/10.1128/JVI.01925-17.

26. Barouch, D.H., Whitney, J.B., Moldt, B., Klein, F., Oliveira, T.Y., Liu, J., Stephenson, K.E.,
Chang, H.-W., Shekhar, K., Gupta, S., et al. (2013). Therapeutic efficacy of potent
neutralizing HIV-1-specific monoclonal antibodies in SHIV-infected rhesus monkeys.
Nature, 1–16. https://doi.org/10.1038/nature12744.

27. Borducchi, E.N., Liu, J., Nkolola, J.P., Cadena, A.M., Yu, W.-H., Fischinger, S., Broge, T.,
Abbink, P., Mercado, N.B., Chandrashekar, A., et al. (2018). Antibody and TLR7 agonist
delay viral rebound in SHIV-infected monkeys. Nature *278*, 1295.
https://doi.org/10.1038/s41586-018-0600-6.

28. Halper-Stromberg, A., Lu, C.-L., Klein, F., Horwitz, J.A., Bournazos, S., Nogueira, L.,
Eisenreich, T.R., Liu, C., Gazumyan, A., Schaefer, U., et al. (2014). Broadly Neutralizing
Antibodies and Viral Inducers Decrease Rebound from HIV-1 Latent Reservoirs in
Humanized Mice. Cell, 1–11. https://doi.org/10.1016/j.cell.2014.07.043.

29. Horwitz, J.A., Halper-Stromberg, A., Mouquet, H., Gitlin, A.D., Tretiakova, A., Eisenreich,
T.R., Malbec, M., Gravemann, S., Billerbeck, E., Dorner, M., et al. (2013). HIV-1 suppression
and durable control by combining single broadly neutralizing antibodies and antiretroviral
drugs in humanized mice. Proceedings of the National Academy of Sciences of the United
States of America. https://doi.org/10.1073/pnas.1315295110.

30. Julg, B., Pegu, A., Abbink, P., Liu, J., Brinkman, A., Molloy, K., Mojta, S., Chandrashekar, A.,
Callow, K., Wang, K., et al. (2017). Virological Control by the CD4-Binding Site Antibody N6
in Simian-Human Immunodeficiency Virus-Infected Rhesus Monkeys. Journal of Virology
91, e00498-17. https://doi.org/10.1128/JVI.00498-17.

31. Klein, F., Halper-Stromberg, A., Horwitz, J.A., Gruell, H., Scheid, J.F., Bournazos, S.,
Mouquet, H., Spatz, L.A., Diskin, R., Abadir, A., et al. (2012). HIV therapy by a combination
of broadly neutralizing antibodies in humanized mice. Nature *492*, 118–122.
https://doi.org/10.1038/nature11604.

32. Shingai, M., Nishimura, Y., Klein, F., Mouquet, H., Donau, O.K., Plishka, R., Buckler-White,
A., Seaman, M., Piatak, M., Lifson, J.D., et al. (2013). Antibody-mediated immunotherapy
of macaques chronically infected with SHIV suppresses viraemia. Nature, 1–15.
https://doi.org/10.1038/nature12746.

33. Bar, K.J., Sneller, M.C., Harrison, L.J., Justement, J.S., Overton, E.T., Petrone, M.E., Salantes,
D.B., Seamon, C.A., Scheinfeld, B., Kwan, R.W., et al. (2016). Effect of HIV Antibody VRC01
on Viral Rebound after Treatment Interruption. The New England journal of medicine *375*,
2037–2050. https://doi.org/10.1056/NEJMoa1608243.

34. Caskey, M., Klein, F., Lorenzi, J.C.C., Seaman, M.S., West, A.P., Buckley, N., Kremer, G.,
Nogueira, L., Braunschweig, M., Scheid, J.F., et al. (2015). Viraemia suppressed in HIV-1-
infected humans by broadly neutralizing antibody 3BNC117. Nature.
https://doi.org/10.1038/nature14411.

35. Caskey, M., Schoofs, T., Gruell, H., Settler, A., Karagounis, T., Kreider, E.F., Murrell, B.,
Pfeifer, N., Nogueira, L., Oliveira, T.Y., et al. (2017). Antibody 10-1074 suppresses viremia
in HIV-1-infected individuals. Nature Medicine *23*, 185–191.
https://doi.org/10.1038/nm.4268.

36. Lynch, R.M., Wong, P., Tran, L., O'Dell, S., Nason, M.C., Li, Y., Wu, X., and Mascola, J.R.
(2015). HIV-1 fitness cost associated with escape from the VRC01 class of CD4 binding site
neutralizing antibodies. Journal of Virology *89*, 4201–4213.
https://doi.org/10.1128/JVI.03608-14.

37. Mehandru, S., Vcelar, B., Wrin, T., Stiegler, G., Joos, B., Mohri, H., Boden, D., Galovich, J.,
Tenner-Racz, K., Racz, P., et al. (2007). Adjunctive passive immunotherapy in human
immunodeficiency virus type 1-infected individuals treated with antiviral therapy during
acute and early infection. Journal of Virology *81*, 11016–11031.
https://doi.org/10.1128/JVI.01340-07.

38. Mendoza, P., Gruell, H., Nogueira, L., Pai, J.A., Butler, A.L., Millard, K., Lehmann, C., Suárez,
I., Oliveira, T.Y., Lorenzi, J.C.C., et al. (2018). Combination therapy with anti-HIV-1
antibodies maintains viral suppression. Nature, 1–21. https://doi.org/10.1038/s41586-
018-0531-2.

39. Trkola, A., Kuster, H., Rusert, P., Joos, B., Fischer, M., Leemann, C., Manrique, A., Huber,
M., Rehr, M., Oxenius, A., et al. (2005). Delay of HIV-1 rebound after cessation of
antiretroviral therapy through passive transfer of human neutralizing antibodies. Nature
Medicine *11*, 615–622. https://doi.org/10.1038/nm1244.

40. Happe, M., Lynch, R.M., Fichtenbaum, C.J., Heath, S.L., Koletar, S.L., Landovitz, R.J., Presti,
R.M., Santana-Bagur, J.L., Tressler, R.L., Holman, L.A., et al. (2025). Virologic effects of
broadly neutralizing antibodies VRC01LS and VRC07-523LS on chronic HIV-1 infection. JCI
Insight *10*, e181496. https://doi.org/10.1172/jci.insight.181496.

41. Bar-On, Y., Gruell, H., Schoofs, T., Pai, J.A., Nogueira, L., Butler, A.L., Millard, K., Lehmann,
C., rez, I.S. x000E1, Oliveira, T.Y., et al. (2018). Safety and antiviral activity of combination
HIV-1 broadly neutralizing antibodies in viremic individuals. Nature Medicine, 1–13.
https://doi.org/10.1038/s41591-018-0186-4.

42. Julg, B., Walker-Sperling, V.E.K., Wagh, K., Aid, M., Stephenson, K.E., Zash, R., Liu, J.,
Nkolola, J.P., Hoyt, A., Castro, M., et al. (2024). Safety and antiviral effect of a triple
combination of HIV-1 broadly neutralizing antibodies: a phase 1/2a trial. Nat Med.
https://doi.org/10.1038/s41591-024-03247-5.

43. Ledgerwood, J.E., Coates, E.E., Yamshchikov, G., Saunders, J.G., Holman, L., Enama, M.E.,
DeZure, A., Lynch, R., Gordon, I., Plummer, S., et al. (2015). Safety, Pharmacokinetics, and
Neutralization of the Broadly Neutralizing HIV-1 Human Monoclonal Antibody VRC01 in
Healthy Adults. Clinical and experimental immunology. https://doi.org/10.1111/cei.12692.

44. Lewis, A.D., Chen, R., Montefiori, D.C., Johnson, P.R., and Clark, K.R. (2002). Generation of
Neutralizing Activity against Human Immunodeficiency Virus Type 1 in Serum by Antibody
Gene Transfer. Journal of Virology *76*, 8769–8775.
https://doi.org/10.1128/JVI.76.17.8769-8775.2002.

45. Fang, J., Qian, J.-J., Yi, S., Harding, T.C., Tu, G.H., VanRoey, M., and Jooss, K. (2005). Stable
antibody expression at therapeutic levels using the 2A peptide. Nature Biotechnology *23*,
584–590. https://doi.org/10.1038/nbt1087.

46. Fuchs, S.P., Martinez-Navio, J.M., Piatak, M., Lifson, J.D., Gao, G., and Desrosiers, R.C.
(2015). AAV-Delivered Antibody Mediates Significant Protective Effects against SIVmac239
Challenge in the Absence of Neutralizing Activity. PLoS Pathogens *11*, e1005090.
https://doi.org/10.1371/journal.ppat.1005090.

47. Gardner, M.R., Kattenhorn, L.M., Kondur, H.R., von Schaewen, M., Dorfman, T., Chiang,
J.J., Haworth, K.G., Decker, J.M., Alpert, M.D., Bailey, C.C., et al. (2015). AAV-expressed
eCD4-Ig provides durable protection from multiple SHIV challenges. Nature, 1–16.
https://doi.org/10.1038/nature14264.

48. Brady, J.M., Baltimore, D., and Balazs, A.B. (2017). Antibody gene transfer with adeno-
associated viral vectors as a method for HIV prevention. Immunological reviews *275*, 324–
333. https://doi.org/10.1111/imr.12478.

49. Saunders, K.O., Wang, L., Joyce, M.G., Yang, Z.-Y., Balazs, A.B., Cheng, C., Ko, S.-Y., Kong,
W.-P., Rudicell, R.S., Georgiev, I.S., et al. (2015). Broadly Neutralizing Human
Immunodeficiency Virus Type 1 Antibody Gene Transfer Protects Nonhuman Primates
from Mucosal Simian-Human Immunodeficiency Virus Infection. Journal of Virology *89*,
8334–8345. https://doi.org/10.1128/JVI.00908-15.

50. Welles, H.C., Jennewein, M.F., Mason, R.D., Narpala, S., Wang, L., Cheng, C., Zhang, Y.,
Todd, J.-P., Lifson, J.D., Balazs, A.B., et al. (2018). Vectored delivery of anti-SIV envelope
targeting mAb via AAV8 protects rhesus macaques from repeated limiting dose intrarectal
swarm SIVsmE660 challenge. PLoS Pathogens *14*, e1007395.
https://doi.org/10.1371/journal.ppat.1007395.

51. Casazza, J.P., Cale, E.M., Narpala, S., Yamshchikov, G.V., Coates, E.E., Hendel, C.S., Novik,
L., Holman, L.A., Widge, A.T., Apte, P., et al. (2022). Safety and tolerability of AAV8 delivery
of a broadly neutralizing antibody in adults living with HIV: a phase 1, dose-escalation trial.
Nat Med. https://doi.org/10.1038/s41591-022-01762-x.

52. Fuchs, S.P., Mondragon, P.G., Zabizhin, R., Tomer, S., Wang, L., Cook, E., Dudley, D.M.,
Weisgrau, K.L., Furlott, J., Coonen, J., et al. (2025). Transient rapamycin treatment avoids
unwanted host immune responses toward AAV-delivered anti-HIV antibodies. Nat
Commun *16*, 8906. https://doi.org/10.1038/s41467-025-63970-6.

53. Ardeshir, A., O'Hagan, D., Mehta, I., Shandilya, S., Hopkins, L.L.J., Adamson, L., Kuroda,
M.J., Hahn, P.A., da Costa, L.A.B., Fuchs, S.P., et al. (2025). Determinants of successful
AAV-vectored delivery of HIV-1 bNAbs in early life. Nature *645*, 1020–1028.
https://doi.org/10.1038/s41586-025-09330-2.

54. Thavarajah, J.J., Hønge, B.L., and Wejse, C.M. (2024). The Use of Broadly Neutralizing
Antibodies (bNAbs) in HIV-1 Treatment and Prevention. Viruses *16*, 911.
https://doi.org/10.3390/v16060911.

55.   Koyanagi, Y., Miles, S., Mitsuyasu, R.T., Merrill, J.E., Vinters, H.V., and Chen, I.S. (1987).
1316   Dual infection of the central nervous system by AIDS viruses with distinct cellular tropisms.
1317   Science *236*, 819–822.

1318   56.   Ochsenbauer, C., Edmonds, T.G., Ding, H., Keele, B.F., Decker, J., Salazar, M.G., Salazar-
1319   Gonzalez, J.F., Shattock, R., Haynes, B.F., Shaw, G.M., et al. (2012). Generation of
1320   Transmitted/Founder HIV-1 Infectious Molecular Clones and Characterization of Their
1321   Replication Capacity in CD4 T Lymphocytes and Monocyte-Derived Macrophages. J Virol
1322   *86*, 2715–2728. https://doi.org/10.1128/JVI.06157-11.

1323   57.   McCarthy, M., He, J., and Wood, C. (1998). HIV-1 Strain-Associated Variability in Infection
1324   of Primary Neuroglia. J Neurovirol *4*, 80–89.
1325   https://doi.org/10.3109/13550289809113484.

1326   58.   Wright, S. (1932). The roles of mutation, inbreeding, crossbreeding, and selection in
1327   evolution. Proc. In Sixth Int. Cong. Genet *1*, 356–366.

1328   59.   Melkus, M.W., Estes, J.D., Padgett-Thomas, A., Gatlin, J., Denton, P.W., Othieno, F.A.,
1329   Wege, A.K., Haase, A.T., and Garcia, J.V. (2006). Humanized mice mount specific adaptive
1330   and innate immune responses to EBV and TSST-1. Nature Medicine *12*, 1316–1322.
1331   https://doi.org/10.1038/nm1431.

1332   60.   Baenziger, S., Tussiwand, R., Schlaepfer, E., Mazzucchelli, L., Heikenwalder, M., Kurrer,
1333   M.O., Behnke, S., Frey, J., Oxenius, A., Joller, H., et al. (2006). Disseminated and sustained
1334   HIV infection in CD34+ cord blood cell-transplanted Rag2-/-gamma c-/- mice. Proceedings
1335   of the National Academy of Sciences of the United States of America *103*, 15951–15956.
1336   https://doi.org/10.1073/pnas.0604493103.

1337   61.   Sun, Z., Denton, P.W., Estes, J.D., Othieno, F.A., Wei, B.L., Wege, A.K., Melkus, M.W.,
1338   Padgett-Thomas, A., Zupancic, M., Haase, A.T., et al. (2007). Intrarectal transmission,
1339   systemic infection, and CD4+ T cell depletion in humanized mice infected with HIV-1. The
1340   Journal of experimental medicine *204*, 705–714. https://doi.org/10.1084/jem.20062411.

1341   62.   Huang, J., Kang, B.H., Ishida, E., Zhou, T., Griesman, T., Sheng, Z., Wu, F., Doria-Rose, N.A.,
1342   Zhang, B., McKee, K., et al. (2016). Identification of a CD4-Binding-Site Antibody to HIV
1343   that Evolved Near-Pan Neutralization Breadth. Immunity *45*, 1108–1121.
1344   https://doi.org/10.1016/j.immuni.2016.10.027.

1345   63.   Sok, D., van Gils, M.J., Pauthner, M., Julien, J.-P., Saye-Francisco, K.L., Hsueh, J., Briney, B.,
1346   Lee, J.H., Le, K.M., Lee, P.S., et al. (2014). Recombinant HIV envelope trimer selects for
1347   quaternary-dependent antibodies targeting the trimer apex. Proceedings of the National
1348   Academy of Sciences of the United States of America *111*, 17624–17629.
1349   https://doi.org/10.1073/pnas.1415789111.

64.   Webb, N.E., Montefiori, D.C., and Lee, B. (2015). Dose–response curve slope helps predict

therapeutic potency and breadth of HIV broadly neutralizing antibodies. Nat Commun *6*,
8443. https://doi.org/10.1038/ncomms9443.

65. Galvez, N.M.S., Sheehan, M.L., Lin, A.Z., Cao, Y., Lam, E.C., Jackson, A.M., and Balazs, A.B.
(2024). QuickFit: A High-Throughput RT-qPCR-Based Assay to Quantify Viral Growth and
Fitness In Vitro. Viruses *16*, 1320. https://doi.org/10.3390/v16081320.

66. Wainburg, M.A. (2004). The impact of the M184V substitution on drug resistance and viral
fitness. Expert Review of Anti-infective Therapy *2*, 147–151.
https://doi.org/10.1586/14787210.2.1.147.

67. Mendoza, P., Gruell, H., Nogueira, L., Pai, J.A., Butler, A.L., Millard, K., Lehmann, C., Suárez,
I., Oliveira, T.Y., Lorenzi, J.C.C., et al. (2018). Combination therapy with anti-HIV-1
antibodies maintains viral suppression. Nature, 1–21. https://doi.org/10.1038/s41586-
018-0531-2.

68. Julg, B., Stephenson, K.E., Wagh, K., Tan, S.C., Zash, R., Walsh, S., Ansel, J., Kanjilal, D.,
Nkolola, J., Walker-Sperling, V.E.K., et al. (2022). Safety and antiviral activity of triple
combination broadly neutralizing monoclonal antibody therapy against HIV-1: a phase 1
clinical trial. Nat Med *28*, 1288–1296. https://doi.org/10.1038/s41591-022-01815-1.

69. Pegu, A., Xu, L., DeMouth, M.E., Fabozzi, G., March, K., Almasri, C.G., Cully, M.D., Wang, K.,
Yang, E.S., Dias, J., et al. (2022). Potent anti-viral activity of a trispecific HIV neutralizing
antibody in SHIV-infected monkeys. Cell Reports *38*, 110199.
https://doi.org/10.1016/j.celrep.2021.110199.

70. Lynch, R.M., Boritz, E., Coates, E.E., DeZure, A., Madden, P., Costner, P., Enama, M.E.,
Plummer, S., Holman, L., Hendel, C.S., et al. (2015). Virologic effects of broadly neutralizing
antibody VRC01 administration during chronic HIV-1 infection. Science translational
medicine *7*, 319ra206-319ra206. https://doi.org/10.1126/scitranslmed.aad5752.

71. Schoofs, T., Klein, F., Braunschweig, M., Kreider, E.F., Feldmann, A., Nogueira, L., Oliveira,
T., Lorenzi, J.C.C., Parrish, E.H., Learn, G.H., et al. (2016). HIV-1 therapy with monoclonal
antibody 3BNC117 elicits host immune responses against HIV-1. Science.
https://doi.org/10.1126/science.aaf0972.

72. Scheid, J.F., Horwitz, J.A., Bar-On, Y., Kreider, E.F., Lu, C.-L., Lorenzi, J.C.C., Feldmann, A.,
Braunschweig, M., Nogueira, L., Oliveira, T., et al. (2016). HIV-1 antibody 3BNC117
suppresses viral rebound in humans during treatment interruption. Nature.
https://doi.org/10.1038/nature18929.

73. Otsuka, Y., Schmitt, K., Quinlan, B.D., Gardner, M.R., Alfant, B., Reich, A., Farzan, M., and
Choe, H. (2018). Diverse pathways of escape from all well-characterized VRC01-class
broadly neutralizing HIV-1 antibodies. PLoS Pathog *14*, e1007238.
https://doi.org/10.1371/journal.ppat.1007238.

74. Wise, M.C., Xu, Z., Tello-Ruiz, E., Beck, C., Trautz, A., Patel, A., Elliott, S.T.C., Chokkalingam,
N., Kim, S., Kerkau, M.G., et al. (2020). In vivo delivery of synthetic DNA–encoded
antibodies induces broad HIV-1–neutralizing activity. Journal of Clinical Investigation *130*,
827–837. https://doi.org/10.1172/JCI132779.

75. Wu, R.L., Houser, K.V., Gaudinski, M.R., Widge, A.T., Awan, S.F., Carter, C.A., Holman, L.A.,
Saunders, J., Hendel, C.S., Eshun, A., et al. (2025). Safety and pharmacokinetics of N6LS, a
broadly neutralising monoclonal antibody for HIV: a phase 1, open-label, dose-escalation
study in healthy adults. The Lancet HIV. https://doi.org/10.1016/s2352-3018(25)00041-4.

76. Moore, P.L., Gray, E.S., Sheward, D., Madiga, M., Ranchobe, N., Lai, Z., Honnen, W.J.,
Nonyane, M., Tumba, N., Hermanus, T., et al. (2011). Potent and broad neutralization of
HIV-1 subtype C by plasma antibodies targeting a quaternary epitope including residues in
the V2 loop. J Virol *85*, 3128–3141. https://doi.org/10.1128/JVI.02658-10.

77. Radford, C.E., and Bloom, J.D. (2025). Comprehensive maps of escape mutations from
antibodies 10-1074 and 3BNC117 for Envs from two divergent HIV strains. bioRxiv,
2025.01.30.635715. https://doi.org/10.1101/2025.01.30.635715.

78. Neher, R.A., and Leitner, T. (2010). Recombination Rate and Selection Strength in HIV
Intra-patient Evolution. PLoS Comput Biol *6*, e1000660.
https://doi.org/10.1371/journal.pcbi.1000660.

79. Li, Y., O'Dell, S., Walker, L.M., Wu, X., Guenaga, J., Feng, Y., Schmidt, S.D., McKee, K.,
Louder, M.K., Ledgerwood, J.E., et al. (2011). Mechanism of neutralization by the broadly
neutralizing HIV-1 monoclonal antibody VRC01. J Virol *85*, 8954–8967.
https://doi.org/10.1128/JVI.00754-11.

80. Pegu, A., Borate, B., Huang, Y., Pauthner, M.G., Hessell, A.J., Julg, B., Doria-Rose, N.A.,
Schmidt, S.D., Carpp, L.N., Cully, M.D., et al. (2019). A Meta-analysis of Passive
Immunization Studies Shows that Serum-Neutralizing Antibody Titer Associates with
Protection against SHIV Challenge. Cell Host & Microbe *26*, 336-346.e3.
https://doi.org/10.1016/j.chom.2019.08.014.

81. Fellinger, C.H., Gardner, M.R., Weber, J.A., Alfant, B., Zhou, A.S., and Farzan, M. (2019).
eCD4-Ig limits HIV-1 escape more effectively than CD4-Ig or a broadly neutralizing
antibody. Journal of Virology. https://doi.org/10.1128/JVI.00443-19.

82. Bricault, C.A., Yusim, K., Seaman, M.S., Yoon, H., Theiler, J., Giorgi, E.E., Wagh, K., Theiler,
M., Hraber, P., Macke, J.P., et al. (2019). HIV-1 Neutralizing Antibody Signatures and
Application to Epitope-Targeted Vaccine Design. Cell Host & Microbe *25*, 59-72.e8.
https://doi.org/10.1016/j.chom.2018.12.001.

83. Asmal, M., Luedemann, C., Lavine, C.L., Mach, L.V., Balachandran, H., Brinkley, C., Denny,
T.N., Lewis, M.G., Anderson, H., Pal, R., et al. (2015). Infection of monkeys by simian-

human immunodeficiency viruses with transmitted/founder clade C HIV-1 envelopes.
Virology *475*, 37–45. https://doi.org/10.1016/j.virol.2014.10.032.

84.   Zhang, C., Zaman, L.A., Poluektova, L.Y., Gorantla, S., Gendelman, H.E., and Dash, P.K.
1426   (2023). Humanized Mice for Studies of HIV-1 Persistence and Elimination. Pathogens *12*,
1427   879. https://doi.org/10.3390/pathogens12070879.

1428   85.   Wodarz, D., and Nowak, M.A. (1998). The effect of different immune responses on the
1429   evolution of virulent CXCR4–tropic HIV. Proc. R. Soc. Lond. B *265*, 2149–2158.
1430   https://doi.org/10.1098/rspb.1998.0552.

1431   86.   Theys, K., Deforche, K., Beheydt, G., Moreau, Y., Van Laethem, K., Lemey, P., Camacho,
1432   R.J., Rhee, S.-Y., Shafer, R.W., Van Wijngaerden, E., et al. (2010). Estimating the
1433   individualized HIV-1 genetic barrier to resistance using a nelfinavir fitness landscape. BMC
1434   Bioinformatics *11*, 409. https://doi.org/10.1186/1471-2105-11-409.

1435   87.   Ganusov, V.V., and De Boer, R.J. (2006). Estimating Costs and Benefits of CTL Escape
1436   Mutations in SIV/HIV Infection. PLoS Comput Biol *2*, e24.
1437   https://doi.org/10.1371/journal.pcbi.0020024.

1438   88.   Read, E.L., Tovo-Dwyer, A.A., and Chakraborty, A.K. (2012). Stochastic effects are
1439   important in intrahost HIV evolution even when viral loads are high. Proc. Natl. Acad. Sci.
1440   U.S.A. *109*, 19727–19732. https://doi.org/10.1073/pnas.1206940109.

1441   89.   Batorsky, R., Sergeev, R.A., and Rouzine, I.M. (2014). The Route of HIV Escape from
1442   Immune Response Targeting Multiple Sites Is Determined by the Cost-Benefit Tradeoff of
1443   Escape Mutations. PLoS Comput Biol *10*, e1003878.
1444   https://doi.org/10.1371/journal.pcbi.1003878.

1445   90.   Gardner, E.M., Burman, W.J., Steiner, J.F., Anderson, P.L., and Bangsberg, D.R. (2009).
1446   Antiretroviral medication adherence and the development of class-specific antiretroviral
1447   resistance. AIDS *23*, 1035–1046. https://doi.org/10.1097/QAD.0b013e32832ba8ec.

1448   91.   Armstrong, K.L., Lee, T.-H., and Essex, M. (2011). Replicative Fitness Costs of
1449   Nonnucleoside Reverse Transcriptase Inhibitor Drug Resistance Mutations on HIV Subtype
1450   C. Antimicrob Agents Chemother *55*, 2146–2153. https://doi.org/10.1128/AAC.01505-10.

1451   92.   Hill, A.L., Rosenbloom, D.I.S., and Nowak, M.A. (2012). Evolutionary dynamics of HIV at
1452   multiple spatial and temporal scales. J Mol Med *90*, 543–561.
1453   https://doi.org/10.1007/s00109-012-0892-1.

1454   93.   Zanini, F., Puller, V., Brodin, J., Albert, J., and Neher, R.A. (2017). *In vivo* mutation rates and
1455   the landscape of fitness costs of HIV-1. Virus Evolution *3*, vex003.
1456   https://doi.org/10.1093/ve/vex003.

94.   Meijers, M., Vanshylla, K., Gruell, H., Klein, F., and Lässig, M. (2021). Predicting in vivo

escape dynamics of HIV-1 from a broadly neutralizing antibody. Proc. Natl. Acad. Sci.
U.S.A. *118*, e2104651118. https://doi.org/10.1073/pnas.2104651118.

95. Cong, M., Heneine, W., and García-Lerma, J.G. (2007). The Fitness Cost of Mutations
Associated with Human Immunodeficiency Virus Type 1 Drug Resistance Is Modulated by
Mutational Interactions. J Virol *81*, 3037–3041. https://doi.org/10.1128/jvi.02712-06.

96. Gonelli, C.A., King, H.A.D., Ko, S., Fennessey, C.M., Iwamoto, N., Mason, R.D., Heimann, A.,
Flebbe, D.R., Todd, J.-P., Foulds, K.E., et al. (2025). Antibody prophylaxis may mask
subclinical SIV infections in macaques. Nature *639*, 205–213.
https://doi.org/10.1038/s41586-024-08500-y.

97. Caskey, M. (2020). Broadly neutralizing antibodies for the treatment and prevention of HIV
infection. Curr Opin HIV AIDS *15*, 49–55.
https://doi.org/10.1097/COH.0000000000000600.

98. Han, C., Johnson, J., Dong, R., Kandula, R., Kort, A., Wong, M., Yang, T., Breheny, P.J.,
Brown, G.D., and Haim, H. (2020). Key Positions of HIV-1 Env and Signatures of Vaccine
Efficacy Show Gradual Reduction of Population Founder Effects at the Clade and Regional
Levels. mBio *11*, e00126-20. https://doi.org/10.1128/mBio.00126-20.

99. Haddox, H.K., Dingens, A.S., Hilton, S.K., Overbaugh, J., and Bloom, J.D. (2018). Mapping
mutational effects along the evolutionary landscape of HIV envelope. eLife *7*.
https://doi.org/10.7554/eLife.34420.

100. DeLeon, O., Hodis, H., O'Malley, Y., Johnson, J., Salimi, H., Zhai, Y., Winter, E., Remec, C.,
Eichelberger, N., Van Cleave, B., et al. (2017). Accurate predictions of population-level
changes in sequence and structural properties of HIV-1 Env using a volatility-controlled
diffusion model. PLoS Biol *15*, e2001549. https://doi.org/10.1371/journal.pbio.2001549.

101. Karpel, M.E., Boutwell, C.L., and Allen, T.M. (2015). BLT humanized mice as a small animal
model of HIV infection. Curr Opin Virol *13*, 75–80.
https://doi.org/10.1016/j.coviro.2015.05.002.

102. Seung, E., and Tager, A.M. (2013). Humoral immunity in humanized mice: a work in
progress. J Infect Dis *208 Suppl 2*, S155-159. https://doi.org/10.1093/infdis/jit448.

103. McCoy, L.E., van Gils, M.J., Ozorowski, G., Messmer, T., Briney, B., Voss, J.E., Kulp, D.W.,
Macauley, M.S., Sok, D., Pauthner, M., et al. (2016). Holes in the Glycan Shield of the
Native HIV Envelope Are a Target of Trimer-Elicited Neutralizing Antibodies. Cell Reports
16, 2327–2338. https://doi.org/10.1016/j.celrep.2016.07.074.

104. Freund, N.T., Wang, H., Scharf, L., Nogueira, L., Horwitz, J.A., Bar-On, Y., Golijanin, J.,
Sievers, S.A., Sok, D., Cai, H., et al. (2017). Coexistence of potent HIV-1 broadly neutralizing
antibodies and antibody-sensitive viruses in a viremic controller. Science translational
medicine *9*, eaal2144. https://doi.org/10.1126/scitranslmed.aal2144.

105. Ramakrishnan, M.A. (2016). Determination of 50% endpoint titer using a simple formula.
World journal of virology *5*, 85–86. https://doi.org/10.5501/wjv.v5.i2.85.

106. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics (Oxford, England) *34*, i884–i890.
https://doi.org/10.1093/bioinformatics/bty560.

107. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nature
Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

108. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) *25*, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352.

109. Korber, B.T., Foley, B.T., Kuiken, C.L., Pillai, S.K., and Sodroski, J.G. (2014). Numbering
Positions in HIV Relative to HXB2CG. LANL.
https://www.hiv.lanl.gov/content/sequence/HIV/REVIEWS/HXB2.html.

110. Köster, J., and Rahmann, S. (2012). Snakemake–a scalable bioinformatics workflow engine.
Bioinformatics (Oxford, England) *28*, 2520–2522.
https://doi.org/10.1093/bioinformatics/bts480.

111. Wilm, A., Aw, P.P.K., Bertrand, D., Yeo, G.H.T., Ong, S.H., Wong, C.H., Khor, C.C., Petric, R.,
Hibberd, M.L., and Nagarajan, N. (2012). LoFreq: a sequence-quality aware, ultra-sensitive
variant caller for uncovering cell-population heterogeneity from high-throughput
sequencing datasets. Nucleic Acids Research *40*, 11189–11201.
https://doi.org/10.1093/nar/gks918.

112. Piantadosi, A., Freije, C.A., Gosmann, C., Ye, S., Park, D., Schaffner, S.F., Tully, D.C., Allen,
T.M., Dong, K.L., Sabeti, P.C., et al. (2019). Metagenomic Sequencing of HIV-1 in the Blood
and Female Genital Tract Reveals Little Quasispecies Diversity during Acute Infection.
Journal of Virology *93*. https://doi.org/10.1128/JVI.00804-18.

113. McCrone, J.T., and Lauring, A.S. (2016). Measurements of Intrahost Viral Diversity Are
Extremely Sensitive to Systematic Errors in Variant Calling. Journal of Virology *90*, 6884–
6895. https://doi.org/10.1128/JVI.00667-16.

114. Siebring-van Olst, E., Vermeulen, C., de Menezes, R.X., Howell, M., Smit, E.F., and van
Beusechem, V.W. (2013). Affordable Luciferase Reporter Assay for Cell-Based High-
Throughput Screening. J Biomol Screen *18*, 453–461.
https://doi.org/10.1177/1087057112465184.

115. Borchers, H.W. (2023). pracma: Practical Numerical Math Functions. Version R package
version 2.4.4.

116. R Core Team (2024). R: A Language and Environment for Statistical Computing. (R
1530   Foundation for Statistical Computing).

1531   117. Reagan-Shaw, S., Nihal, M., and Ahmad, N. (2008). Dose translation from animal to human
1532   studies revisited. The FASEB journal : official publication of the Federation of American
1533   Societies for Experimental Biology *22*, 659–661. https://doi.org/10.1096/fj.07-9574LSF.

1534
1535

Graphical Abstract.

Figure 1. Vectored delivery of VRC07, but not PGDM1400 or N6, can suppress established HIV$_{REJO.c}$ infection.

(A) BLT humanized mice were infected with 300 TCID$_{50}$ of HIV$_{REJO.c}$ and the viral population was allowed to replicate for 4 weeks. Mice were then IM injected with 5x10^{11} genome copies (GC) of AAVs encoding for either VRC07, N6, PGDM1400, or Luciferase as control. Mice were followed for 6 months, and blood samples were collected weekly.

(B) ELISA-based quantitation of gp120-binding antibodies in the serum of HIV$_{REJO.c}$-infected humanized mice following administration of AAV8-Luciferase, AAV8-VRC07, AAV8-N6, or AAV8-PGDM1400 vectors. Black arrow denotes vector administration. Data are plotted as geometric mean ± geometric SD.

(C-F) HIV viral load in plasma of HIV$_{REJO.c}$-infected mice injected with AAV8-Luciferase as control **(C)**, AAV8-VRC07 **(D)**, AAV8-N6 **(E)**, or AAV8-PGDM1400 **(F)**. Black arrows denote vector administration. Each colored line depicts an individual mouse tracked over time. The sensitivity of qPCR was 1 genome copy per µL of plasma, and 5 µL were used in the reaction, resulting in a 1000 copy per mL limit of detection (solid line). Data are presented as mean ± S.E.M.

(G) Kaplan-Meier plot of viral suppression in BLT humanized mice infected with HIV$_{REJO.c}$ given the indicated bNAb-expressing vector. The total model significance (p<0.0001) and pairwise comparison against the Luciferase control (****: p<0.0001) were assessed independently using Log-rank (Mantel-Cox) tests. The percentage of HIV suppressed was defined as the fraction of mice that were not escaped as described in the methods.

Figure 2. Fitness cost and resistance benefit accrued along escape paths determine the difficulty of HIV$_{REJO}$. escape from each bNAb.

(A) Amino acid divergence from the envelope gene of the HIV$_{REJO.c}$ parental strain across all AAV8-VRC07 treated mice. Sequences were determined by Illumina Deep Sequencing of the viral envelope isolated from plasma at the final experimental timepoint. The X-axis represents the envelope protein amino acid position relative to HIV$_{HXB2}$ numbering. The Y-axis represents the percentage of average amino acid divergence from the parental strain, corrected for divergence observed in control mice. Pie charts represent the most common amino acid mutations for sites with the highest divergence. See also Figure S2.

(B) *In vitro* neutralization assays of each HIV$_{REJO.c}$ mutant identified in **(A)** against VRC07. Data are plotted as mean ± S.E.M. Each data point was evaluated in quadruplicate.

(C) Relative viral growth of each HIV$_{REJO.c}$ mutant identified in **(A)**. Growth rates were determined in activated CD4$^+$ T cells performing *QuickFit* assays and normalized to the parental strain. Data are plotted as mean ± S.E.M. and statistical differences were assessed by a Kruskal-Wallis non-parametric ANOVA with Dunn's *post hoc* test to correct for multiple comparisons (****: p<0.0001). See also Figure S2.

(D-F) Escapability maps denoting fitness cost (Y-axis, relative doubling time shown in (**C**)) and resistance benefits (X-axis, neutralization resistance shown in (**B**)) for each HIV$_{REJO.c}$ mutant observed during escape from VRC07 (**D**), N6 (**E**), or PGDM1400 (**F**). Dashed vertical line denotes the geometric mean antibody serum concentration after vectored bNAb administration. Shaded areas represent the neutralization assay limit of detection. Solid arrows represent the likely initial path taken with dashed arrows representing the likely second step paths to escape. In the map legend, each escape path is sorted based on the relative frequency observed in the sequencing data. See also Figures S2 and S3.

(G) Survival analysis of HIV$_{REJO.c}$-infected mice treated with AAV8-VRC07 by escape path haplotype. The total model significance (p=0.0126) and pairwise comparisons of the non-D279A escape paths to each other escape path (*: p<0.05) were assessed independently using Log-rank (Mantel-Cox) tests. See also Figure S3.

(H) Correlation plot of N6 geometric mean-expression vs post-escape HIV$_{REJO.c}$ viral load. The line represents the semi-log least squares regression, ρ represents the Spearman correlation value determined for the data, with the associated p-value below. See also Figure S3.

(I) Correlation plot of the post-N6 escape viral load of HIV$_{REJO.c}$ vs the frequency of N276D determined by viral envelope sequencing. The line represents the semi-log least squares regression, ρ represents the Spearman correlation value determined for the data, with the associated p-value below. See also Figure S3.

(J) Escape Barrier (AUC) Score denoting the aggregate fitness cost for each escape path as determined in the escapability maps. This score was calculated by adding the area under the escapability plot for escape paths A and path B viral escapes from VRC07, N6, and PGDM1400. See also Figure S4.

Figure 3. HIV_REJO.c V5-loop restricts escape from VRC07, by decreasing the resistance benefit of D279 mutations.

(A) ELISA-based quantitation of gp120-binding antibodies in the serum of HIV_JR-CSF-infected humanized mice following administration of 5x10[11] genome copies (GC) of AAV8-Luciferase, AAV8-VRC07, AAV8-N6, or AAV8-PGDM1400 vectors. Black arrow denotes vector administration. Data are plotted as geometric mean ± geometric SD.

(B) Kaplan-Meier plot of viral suppression in humanized mice infected with HIV_JR-CSF given the indicated bNAb-expressing vector. The total model significance was assessed using a Long-rank (Mantel-Cox) test (p=0.0714). See also Figure S5.

(C) Amino acid divergence from the envelope gene of the HIV_JR-CSF parental strain across all AAV8-VRC07 treated mice. Sequences were determined by Illumina Deep Sequencing of the viral envelope isolated from plasma at the final experimental timepoint. The X-axis represents the envelope protein amino acid position relative to HIV_HXB2 numbering. The Y-axis represents the percentage of average amino acid divergence from the parental strain, corrected for divergence observed in control mice. Pie chart represents the most common amino acid mutations for the site with the highest divergence. See also Figure S6.

(D) Escapability map of HIV_JR-CSF escape from VRC07. Dashed vertical line denotes the geometric mean antibody serum concentration after AAV8-VRC07 administration. Shaded area represents the neutralization assay limit of detection. Arrows represent the likely path taken to escape. See also Figures S6 and S8.

(E) Alignment of the D-loop and the V5-loop amino acid sequences for HIV_REJO.c, HIV_JR-CSF, and the HIV_RD, HIV_RV, and HIV_RDV chimeras.

(F) *In vitro* neutralization of HIV_REJO.c, HIV_RD, HIV_RV, HIV_RDV, and HIV_JR-CSF by VRC07 bNAb. Data are plotted as mean ± S.E.M. Each datapoint was evaluated in quadruplicate.

(G) Kaplan-Meier plot of viral suppression in humanized mice infected with HIV_REJO.c, HIV_RV, or HIV_JR-CSF following vectored VRC07 administration. The total model significance (p<0.0001) and pairwise comparisons against HIV_REJO.c (***: p<0.001) were assessed independently using Log-rank (Mantel-Cox) tests. See also Figure S7.

(H) Escapability map of HIV_RV during escape from VRC07. Lighter square symbols represent the original HIV_REJO.c escape path taken against VRC07. Dashed vertical line denotes geometric mean bNAb concentrations after vectored VRC07 administration. Shaded areas represent the limits of detection. Arrows represent the likely path to escape. See also Figures S7 and S8.

(I) Escape Barrier (AUC) Score quantifying the difficulty of escape from VRC07 for HIV_REJO.c, HIV_RV, and HIV_JR-CSF. See also Figure S8.

Figure 4. Increasing the fitness cost of HIV$_{REJO.c}$ escape mutations enhances the efficacy of vectored VRC07.

(A) The fitness of HIV$_{REJO.c}$-VRC07 escape mutations with or without the Pol M184V mutation were evaluated using *QuickFit*. Data are presented as mean ± SD. Statistical differences were assessed by a Two-Way ANOVA, with a Šidák *post hoc* test to correct for multiple comparisons (*: p<0.05; ****: p<0.0001). See also Figure S9.

(B) Escapability map of HIV$_{REJO.c}$-VRC07 escape mutants with or without the Pol M184V mutation. Solid symbols (diamonds) represent envelope mutations on the Pol M184V mutant background, while lighter symbols (circles) represent the data for envelope mutations on the WT Pol background. Shaded areas represent the limits of detection. Solid arrows represent the likely path taken with dashed arrows representing secondary steps to escape. See also Figure S9.

(C) Empirical Escape Barrier score for HIV$_{REJO.c}$ escape from VRC07 as compared to the theoretical Escape Barrier score for HIV$_{REJO.c-PolM184V}$ escape from VRC07.

(D) Experimental setup to determine the impact of Pol M184 mutations on HIV$_{REJO.c}$ escape from VRC07. Humanized mice were infected with HIV$_{REJO.c}$ and then treated with a sub-optimal ART starting 2 weeks after infection, to select and maintain Pol M184 mutants. At week 8, 5x10^{11} GC of AAV8-VRC07 was injected IM, and the ART regimen was stopped at week 10. Mice were followed for 6 months, and blood samples were collected weekly.

(E) Percentage of Pol M184V/I mutations determined from viral sequences isolated from the plasma of HIV$_{REJO.c}$-infected mice at week 8, prior to AAV8-VRC07 administration. Statistical differences were assessed by an unpaired two-tailed Student's t-test (****: p<0.0001). Data are presented as mean ± S.E.M.

(F) HIV$_{REJO.c}$ viral load in plasma of AAV8-VRC07 treated mice. Black arrows denote vector administration. Each colored line depicts an individual mouse. The qPCR lower limit of detection was 1 genome copy per µL of plasma and 5µL were used in the reaction (solid line). Data are presented as mean ± S.E.M.

(G) Kaplan-Meier plot of viral suppression in humanized mice infected with HIV$_{REJO.c}$ with or without ART selection and with or without AAV8-VRC07 administration. The total model significance (p<0.0001) and pairwise comparisons against the Luciferase + Selection control (***: p<0.001; ****: p<0.0001) were assessed independently with Log-rank (Mantel-Cox) tests. See also Figure S9.