**Title:**

**Integrating fish biology, population genomics and fishery policies for informed**
**management of interjurisdictional multi-species fisheries**

**Authors:**

Mayuresh Gangal[1237]*, Rohan Arthur[2], Venkatachalam Suri[2], Awadhesh Pandit[3], Abhinav
Tyagi[3], Praveen Prakash[3], Mihir Sule[4], Iravatee Majgaonkar[5], Jagdish Krishnaswamy[6], Uma
Ramakrishnan[3]

**Keywords**

IRIIS

Interjurisdictional fisheries management

Tropical multi-species fisheries

Genetic population structure

Fisheries genomics

**Summery**

Globally coastal fisheries are managed according to political jurisdictions of nations and their
provinces. This practice often results in multiple states/provinces managing a single
genetically distinct fish population autonomously and without coordination. In multi-species,
decentralized and data poor fisheries such management lead to poor assessment and
eventually overharvest of fish. To assess overexploitation risk for such interjurisdictional
populations, we propose a risk indicator IRIIS which incorporates 1) species-specific
vulnerability 2) distribution of genetically-distinct populations with respect to jurisdictional
boundaries and 3) differences in management practices across jurisdictions. As a case study
we evaluated overexploitation risk for individual populations of nine commonly caught fish
species spanning across multiple jurisdictions along India's west coast. Our results indicate
that interjurisdictional differences in management alters innate biological susceptibility. IRIIS
is also sensitive to population-level differences and policy changes. Our analyses emphasize
the need for population-level assessments and interjurisdictional cooperation to achieve
sustainable fisheries.

[1] Manipal Academy of Higher Education, Manipal, India
[2] Oceans and Coasts programme, Nature Conservation Foundation, Mysore, India
[3] National Centre for Biological Sciences, Bengaluru, India
[4] Faculty of Biology, University of Gdansk, Poland
[5] Ashoka Trust for Research in Ecology And the Environment, Bengaluru, India
[6] School of Environment and Sustainability, Indian Institute for Human Settlements, Bengaluru, India

[7] Lead contact and corresponding author – Mayuresh Gangal*Correspondence: mayuresh@ncf-india.org

**Science and society statement**

Coastal fisheries worldwide are managed through politically defined maritime jurisdictions,
while fish populations are structured by biological processes such as dispersal, ocean
currents, and natural barriers. This fundamental mismatch between biological population
boundaries and jurisdictional management boundaries often results in the same population
being exploited under multiple, uncoordinated governance regimes. In the context of the
global overfishing crisis, such governance fragmentation can lead to inaccurate stock
assessments and heightened risk of local population collapse. This challenge is particularly
acute in developing and tropical countries, where fisheries are multispecies, data-limited,
and increasingly decentralized to support community-based governance. While
decentralization promotes social justice, it can inadvertently intensify the misalignment
between biological and administrative boundaries, complicating sustainable management.

In this manuscript, we introduce a robust framework — the Integrative Risk Indicator for
Interjurisdictional Stocks (IRIIS) — which has a high applicability for multispecies, data-poor,
and decentralized fisheries. Building on, and extending beyond, species-level Productivity
Susceptibility Analysis (PSA), IRIIS integrates population genomics to delineate biologically
meaningful population boundaries and combines these with fisheries policies and regulations
across jurisdictions to quantify management-associated risks of overexploitation at the
population level. We apply IRIIS to case studies of nine commercially important fish species
harvested along the west coast of India. Our results demonstrate that the framework can (i)
distinguish differential vulnerability among populations of the same species, (ii) explicitly
capture management disparities across jurisdictions exploiting shared populations, and (iii)
detect sensitivity to temporal changes in fisheries policies. By linking genomics, governance,
and risk assessment, IRIIS provides a practical framework for identifying interjurisdictional
management gaps and actionable entry points for policy coordination. This approach is
particularly relevant for the Global South, where reconciling social equity objectives with
ecological sustainability is both essential and challenging.

## 1. Introduction

Historically described as an inexhaustible natural resource [1], marine fish today face a crisis of overfishing [2]. A recent report by the Food and Agriculture Organization of the United Nations flags 37.7 percent of global fish stocks as fished at unsustainable levels [3]. By some estimates, ~260 million people depend on marine fisheries for livelihood [4], while others suggest that ~600 million people depend on fisheries and aquaculture at least partially or completely [3]. These figures highlight that overfishing is not merely an ecological problem, but a crisis of food security, livelihoods and equity.

Since fish has been always viewed as a natural resource, when it comes to managing capture fisheries, economic and political factors override ecological factors [5,6]. Also, management of marine resources is often done according to terrestrial concepts of management like sovereignty, boundaries, private property etc. Mismatch of these concepts with biological and ecological dynamics of marine ecosystems often impedes effective management of fisheries [7–9]. Sustainably managing fishes at spatio-temporal scales that are biologically meaningful, economically viable and managerially effective is the principal challenge confronting global fisheries [10].

Single species demographic stock assessments provide most effective information for fisheries management [10]. However, in 2012, 80% of global fish catch lacked formal assessment [11] and by 2020, only 25% of world fisheries were numerically assessed [12]. The most reliable RAM legacy database also involves only 20% of world's fisheries mostly coming from developed countries in temperate region [13].

For multi species tropical fisheries with limited capacities to undertake robust stock assessments, Productivity Susceptibility Analysis (PSA) can be a valuable tool. PSA attempts to evaluate overexploitation risk of multiple species simultaneously by calculating natural susceptibility of each species to resource offtake on a single comparable scale [14–19].. They attempt to be biologically comprehensive, incorporating population growth rates, trophic level, breeding strategies and other crucial elements of fish biology in their assessments [15].

To bridge the gap between being an analytical useful tool to one that is directly applicable to fisheries policy, a few limitations of the PSA approach need to be addressed. It has been long understood that, for effective monitoring and management, genetically distinct populations of the same species should be considered as separate monitoring and management units [20–22]. For most PSAs, the unit of analysis is the species and not individual population (but see Patrick et al. 2010 [23], limiting their reliability [14,16–19]. Most PSAs also assume a uniform harvest regime along the entire geographical range of analysis. In most tropical countries with large coastlines like Brazil, Indonesia, Philippines, India and Mexico, coastal fisheries are often decentralized and governed by provinces which have significant autonomy in terms of fisheries management [24–28]. Rarely is there a uniform harvest regime across the geographical area considered within a PSA.

Between the multiple genetically distinct populations of a species and decentralized harvest regimes, there is often a mismatch between the biological boundaries of fish populations and the jurisdictional boundaries of fisheries management. These mismatches lead to an increased risk of local population depletion [29].

Here we propose an Integrative Risk Indicator for Interjurisdictional Stocks (IRIIS) as a
heuristic extension of the PSA by incorporating 1) the inherent resilience and susceptibility
attributes of species characteristic of a typical PSA, 2) information from population genetics
to determine geographical boundaries between populations, to indicate whether the
genetically distinct population is managed by single or multiple management regimes. 3) the
biological relevance of present management regulations on given species.

As a case study, we present a IRIIS assessment for nine commercially important nearshore
fish species harvested along the west coast of India. Marine fisheries in India is a paradigm
case of a multi-species multi-management fishery. In India, individual coastal states
(provinces) have jurisdictional autonomy over adjacent territorial waters. They legislate
fisheries regulations in these waters separately, based on local needs, demands from local
stakeholder groups and with little dialogue with neighboring states [28]. This management
gives little regard to the underlying geographical spread of fish populations [28].

For undertaking IRIIS assessment, we first identified genetically distinct populations of nine
chosen species along 3400 km stretch of the west coast of India. Then for each genetically
distinct population, we estimated biological congruence management policies of single or
multiple management regimes acting on them. This information was then used to construct a
population-wise IRIIS. We then discuss the utility of this matrix in guiding policy and
management decisions.

**2. Results**

**2.1. Representing risk of overexploitation associated with species biology (y
axis)**

IRV values can range between 0 (most vulnerable) to 1 (least vulnerable) [18]. Among the 133
species Mohammad et al. 2021 analyzed, *Solenocera crassicornis* (coastal mud prawn) had
the highest IRV value of 0.512 where *Carcharhinus sorrah* (spot-tail shark) had lowest IRV
value of 0.022. Among the nine target species analysed here, *Nemipterus japonicus* had the
highest IRV index value of 0.31 and thus the lowest vulnerability to overexploitation on the
vertical axis whereas *Epinephelus diacanthus* had the lowest IRV index value of 0.195
making it most vulnerable on the vertical axis. Using the approach outlined in Mohammed et
al. 2021, we calculated an IRV index of 0.237 for *Sillago sihama* (Supplementary Material
Section 1). Our target species occupied a narrower portion of the range of species analyzed
by Mohammad et al. 2021.

**2.2. Establishing geographical boundaries of target fish populations**

We sequenced 311 samples belonging to nine study species across nine locations. After
removing samples with low quality during the filtering process, we used a total of 273
samples for the final analysis. On average, we analyzed 4-5 samples per species per
location (See figure S4 in supplementary material).

Single Nucleotide Polymorphism (SNP) analysis indicates that the degree of gene flow
between individuals across the west coast of India varies considerably with species (Figure
1). The PCA results are presented in Figure 1. For detailed species-wise results which

include results of pairwise $F_{ST}$ and Admixture analysis refer to the section 3 of the
supplementary material.

*Rastrelliger kanagurta* and *Chirocentrus nudus* show no evidence of population
differentiation across the west and east coast of India (Figure 1B & 1C). Due to lack of
samples from the east coast, we restricted analysis of *Epinephelus diacanthus* to the west
coast. *Epinephelus diacanthus* shows evidence of a single population across the entire west
coast (Figure 1D). *Trichiurus lepturus* and *Nemipterus japonicus* also show a single
population across the entire west coast, however they have clear divergences from
conspecific populations from the east coast (Figure 1E & 1F).

The remaining four species (*Pampus argenteus, Sardinella longiceps, Scoliodon laticaudus,*
*Sillago sihama)* show multiple genetically distinct populations along the west coast of India.
Genetic data indicate two potential *Pampus argenteus* populations along the west coast of
India, one between northern Gujarat to central Maharashtra, and another from southern
south Maharashtra to Kerala (Figure 1G). *Sardinella longiceps* also shows evidence of two
populations, with a Gujarat population that separates clearly from the rest of the west coast
population (Figure 1H). Genetic data from *Scoliodon laticaudus* shows evidence of three
distinct west coast populations: a northern Gujarat population, a middle peninsular
(Maharashtra-Goa-Karnataka) population, and a southern Kerala population (Figure 1I).
Finally, *Sillago sihama* shows evidence of an isolated population near the town of Kumta
which is separate from the coastwide Maharashtra - Karnataka population (Figure 1J).

Legend:

Panel A – Map of sampling location along the Indian coastline with the color legend for samples from specific locations.

Porbandar = Dark Red, Surat = Red, Mumbai = Light Red, Harne = Orange, Vengurle = Light yellow, Goa = Yellow, Kumta = Light Cyan, Mangalore = Light Blue, Kozhikode = Medium Blue, Alappuzha = Blue, Odisha = Black

Panel B to J – Principal Component Analysis plot of *individual species* with information about number of SNPs and important filters used for variant calling. See also figures S5 to S13 in supplementary material

Figure 1. Principal Component plots for nine chosen species across the west coast of India

**2.3. Identifying percent disparity between species biology and regulations**

We identified the percent disparity between biology and regulation for each individual
population - state (management regime) combination based on the rationale described in the
methods section 2.1.2. To clearly demonstrate how disparity was calculated, results of the
disparity analysis for the *Sardinella longiceps* are shown in Figure 2 as a model species. In
*Sardinella longiceps* The Gujarat population (which is housed in a single management
regime) has a high disparity (66%), while the peninsular India population, despite being
shared across four states, has a low disparity across the entire coast. Further results of all
species are provided in the Supplementary Material, Section 2.

Legend: Each donut represents a state along the west coast of India. Nine sections are for nine
different regulations and the color of each section represents congruence or disparity of the sardine
biology with the regulation. Legend for all the symbols and colors are embedded in the figure. See
also figure S2 in supplementary material

Figure 2. Disparity analysis for Indian oil sardine, with populations housed in single as well
as multiple management regimes

**2.4. Populating the vulnerabilities of west coast populations using IRIIS**

We used species specific IRV values from Mohammad et al. 2021 [18] and state-wise percent
disparity for each population identified by genetic analysis, to construct a IRIIS for the west
coast (Figure 3). variations in individual populations of different species were evident across
both the axis of IRIIS. Critically, species with multiple populations along the coast could have
varied overharvest risk on the management disparity axis. This was most evident for
*Scoliodon laticaudus*, where based on the disparity scores, the peninsular population had a
much higher vulnerability than the Kerala and Gujarat populations. For populations of
*Pampus argenteus* and *Sardinella longiceps*, differences in management practices between
states still clearly influenced the vulnerability of each population. Widely ranging species
(with single populations across the coast) had very different IRV values and were thus very
different in their biological susceptibility. However, because their populations spanned
several jurisdictions along the coast, they were all much more prone to management
disparities.

Legend: Five colored bars represent five risk associated with management disparity which is increasing from left to right. Vertical axis of each bar represents IRV score. For species presented on the matrix the IRV score is written on the left hand side. The color gradient on each vertical bar represents overall risk of overexploitation. Different population of each species is indicated using maps in legend section. See also figure S2 in the supplementary materials

Figure 3 - Integrative Risk Indicator for Interjurisdictional Stocks (IRIIS). for nine species along the west coast of India

## 3. Discussion

Achieving sustainability in multi-species, poorly monitored and data-poor tropical coastal fisheries is often considered a wicked problem with no easy solutions [30],. Although recent approaches like the PSA comprehensively incorporate complex species biology, they do not account for management complexities arising from the interjurisdictional nature of most coastal fish populations. IRIIS combines genomic techniques and policy analysis for a more realistic assessment of fish stock vulnerability.

Our case study demonstrates that populations of many commonly targeted coastal species across the Indian west coast are interjurisdictional in nature, spanning multiple harvest regimes and fisheries policies. This has critical implications for tropical fisheries management. Our study demonstrates that within a species, overharvest risk for individual populations varies based on which and how many management regimes are harvesting them. By capturing these complexities IRIIS provides a framework which allows fisheries managers to evaluate different fisheries options and negotiate pragmatic interjurisdictional arrangements based on known boundaries of at-risk species.

### 3.1. A robust approach to predict vulnerability to overharvest

Our analysis demonstrates the vital importance of evaluating overexploitation risk for every population individually, something that most PSA assessments do not allow [14,16,17,19]. A case in point is the *Pampus argenteus*. With an IRV score of 0.24 (based on the IRV index, see Methods), it is a species of intermediate risk. Our population genetic analysis shows that there are two populations of *Pampus argenteus* along the west coast of India, a northern population that extends from Gujarat to the central coast of Maharashtra, and a southern population that extends further south to Kerala. Although both of these populations share the same IRV score, they are differentially susceptible to overharvest. Kerala have regulations on minimum mesh size for gill nets, and there are *Pampus argenteus* -specific minimum legal-size regulations in place in Goa, Karnataka and Kerala [31–33]. These regulations are absent in the northern states (refer table S2 in supplementary material). In other words, the southern population (southern Maharashtra to Kerala) is managed by states with harvest rules more congruent to *Pampus argenteus* biology (low disparity), making it much less vulnerable to overfishing than the northern population (Gujarat to central Maharashtra). This example demonstrates how by shifting the assessment focus to spatially explicit populations, our matrix considerably enhances the value of the original IRV assessment. Further, by identifying the specific management attributes responsible for differences in risk, the matrix provides actionable recommendations to policy makers on how to reduce overharvest risk

with changes in policy and interjurisdictional cooperation. While the matrix is unable to
assess how these management efforts perform on the ground, but provides a useful
framework to assess their effectiveness.

Another aspect of this IRIIS is that it is dynamic and sensitive to periodic revisions in fishery
policy across management regimes. The case of *Sardinella longiceps* demonstrates this
well. The west coast has a largely panmictic population except for Gujarat, which has a
distinct population. Until 11 years ago, no west coast Indian states had regulations on fishing
methods like bull trawling, light fishing etc [31,34–36]. In addition, there were no mesh size
regulations for purse nets or gill nets and none of the states had defined minimum legal
sizes for the *Sardinella longiceps*. As a result, based on IRIIS, until 2016, the *Sardinella*
*longiceps* was in the highest category of interjurisdictional management associated
overharvest risk. The subsequent adoption of these regulations by all west coast states has
significantly reduced this risk and moved *Sardinella longiceps* on the management axis from
"High disparity in majority of the management regimes" to "Low disparity in all management
regimes" category (Figure 4). Along with the sensitivity of the matrix, this example shows
how the matrix can be used as a tool to formulate biologically meaningful regulations /
policies for target species. In addition, the multi-species nature of the matrix can help policy
makers to track what impact the new regulations will have on other non-targeted species.

Legend: Five colored bars represent five risk associated with management disparity which

is increasing from left to right. Vertical axis of each bar represents IRV score. For species presented on the matrix the IRV score is written on the left hand side. The color gradient on each vertical bar represents overall risk of overexploitation.. See also table S2,S3 and S5 in supplementary materials

253 Figure 4.  Shift in overexploitation risk of the peninsular population of *Sardinella longiceps*
254 along the management disparity axis as a direct result of changing management practices
255
256

We assume that IRV values which are based on critical life history traits remain relatively
constant for a species. However, it is possible that these traits could vary between genetic
stocks.  For instance, Patrik et al., 2010 show that the PSA vulnerability of different
populations of the same species could differ considerably [23]. In evaluating the disparity
between biology and management regime information for our nine target species, we
identified considerable differences in basic life history characters like length at first maturity
(L50) between *Pampus argenteus* [37,38] and *Scoliodon laticaudus* [39,40] caught at different
locations. These traits could potentially also influence their catchability in gear operating
along the coast. Such differences could potentially lead to varying inherent population-level
susceptibilities and slightly different IRV values for each species. One benefit of using matrix
like IRIIS is that as it incorporates genetically distinct population in risk assessment actually
makes these regional differences in life history more apparent.

While IRV values may vary slightly as we learn more about each population, they represent
a largely unchanging inherent property of each taxon. However, a central feature of our
matrix is that real-world vulnerability is as much a characteristic of the management
jurisdictions in question as it is of the biological character of the species. As an example, the
IRV value for *Chirocentrus nudus* is high, making it apparently less vulnerable to
overexploitation. However, its population is spread all along the west coast which have five
different management regimes of five states. In addition, the regulation of four of the five
states has very low congruence with *Chirocentrus nudus* biology (higher disparity). This lack
of congruence makes it far more vulnerable to overexploitation compared to its species-
associated vulnerability indicated by IRV.  This explicitly recognizes the fact that fisheries
regimes vary widely between locations, and that transboundary agreements (or the lack of
them), can strongly mediate the inherent vulnerability of fish stocks. It is critical to keep this
in mind when applying the matrix to a specific fishery. For instance, we classified the Gujarat
population of the *Sardinella longiceps* as belonging to a single jurisdiction, reducing its
overall vulnerability within the matrix. Broadening the geographical scale of our analysis may
well have shown that this population is contiguous with *Sardinella longiceps* from Pakistan
and beyond, making it potentially subject to trans-national management differences that we
did not evaluate here. More comprehensive analysis would require establishing genetic
boundaries for the entire range of the species in question before determining its overall
overharvest risk. This will have important implications for the kinds of interjurisdictional
arrangements required to reduce the overall susceptibility of the population – between
coastal nations or between sub-national jurisdictions [41,42].

The fact that management differences can so strongly mediate inherent vulnerability
indicates that assessments of this kind are most meaningful when designed for every region,
making them jurisdictionally and biologically appropriate across the range of the species.

Productivity Susceptibility Analyses or PSAs are limited in their scope of use since they
cannot incorporate demographic information and population trends [43,44]. Single species stock
assessments and Bio-economic harvest models based on Maximum Sustainable Yield
(MSY) are limited in applicability because of focus on single species [45,46]. Although the
Kobe/Kobi plot incorporates demographic information of multiple species simultaneously it is
considered as too complex from the point of view of policy [47]. IRIIS approach increases
applicability of all such assessments by providing a usable framework for policy makers. It
incorporates the PSA approach on one axis to incorporate life history-based susceptibility of
species. It incorporates policies of individual management regimes and disparity between
them. If a single species stock assessment shows a decreasing trend in stocks, it helps to
identify potential policy actions and assess how other non-species will react to such policy
changes.
306
The central condition of the IRIIS approach is identifying genetically distinct management
units. Larkin highlighted the need of defining stocks based on genetically distinct populations
45 years ago[49]. Multiple studies since them have also demonstrated various issues fisheries
face in the absence of this information[29,50]. Even so majority of developing countries operate
without this information. We believe our work highlights the importance of identifying
genetically distinct populations on both international and regional scales. The techniques we
describe in this study offer a relatively cost-effective means of establishing the spatial limits
of fish populations. This could inform more nuanced population-level life-history studies and
improved risk assessments.

**3.2.   Trans-jurisdictional management of fish populations in tropical waters**

How could this or exercises such as IRIIS inform sustainable tropical fisheries? Risk
assessments of this sort go a long way in sharply defining the management needs for each
population across its natural range and help define the scale at which fisheries management
would be most biologically meaningful. It offers a clear set of directions to better align
species biology with management, setting the stage for cross-jurisdictional cooperation on
specific aspects of policy and practice.

Although we focused on populations which span across multiple jurisdictions within a
sovereign nation, the matrix can be applied for trans-boundary situations as well. Countries
in western Europe have had a long history of hard-won transboundary fishing arrangements
[48], driven by decades of conflict over shared resources. There is also substantially better
information on geographically identifiable, genetically distinct populations of fish that form the
basis of these arrangements [50]. For some tropical countries transboundary agreements are
important means of earning revenue [51–53]. Engaging in such agreements knowing the
geographical extent of genetically distinct populations of transboundary fish and how these
species-specific agreements impact other species in broader multi species fisheries, can
increase the potential effectiveness of such agreements.

In many countries coastal fisheries are managed in a highly decentralized manner where
regions and provinces have considerable autonomy over local fisheries management [24–27,54].
In addition, the allied problems of equity and social justice associated with coastal fisheries
are seeing an increasing movement towards even further localization of fisheries, with
approaches like TURFs (Territorial User Rights for Fishing) gaining popularity [55]. These are
usually smaller than regional government fisheries management units. Most TURFs in Chile,
Mexico and Japan are smaller than the dispersal ability of fish larvae and adults [56]. While
these efforts are a welcome direction to secure the livelihood of marginalized coastal
communities, they are also subject to boundary paradoxes, where increasing fisheries
boundaries come at the cost of increasingly complex interjurisdictional negotiations [57]. Our
analysis raises serious questions of how to ensure the success of these pockets of
community managed fishing in a wider ocean of a rapidly expanding fishery sector [58].
Without adequate coordination across biological bounds, these efforts may also result in
worsening of current rates of overharvest, and it is important to create forums of dialogue
between these community managed and larger regional fisheries, keeping in mind the
genetic boundaries of fish stocks.

As far as our case study from India is concerned, recent Indian fisheries policies heavily
emphasize ecosystem-based management of fisheries. The vision statement of the 2020
National Fisheries Policy emphasizes "developing an ecologically healthy, economically
viable and socially inclusive fisheries sector" with "facilitating coordination among
States/UTs, National Agencies and other stakeholders" as its major policy intervention [59,60].
This policy can create a perfect platform to incorporate spatial information of genetically
distinct populations of individual fish species in the fisheries policy framework. So far, no
forum has been created to facilitate coordination among states and union territories. We
hope our results can provide a strong reason to start this coordination process.

### 3.3. Conclusion

For multi-species tropical fisheries, transitioning to population-specific management of fish
stocks may seem like a distant horizon. Eventually though, if coastal tropical nations are
serious about stemming the collapse of coastal fish stocks, population-wise management is
likely to be critical, at least for the most vulnerable of coastal species. Coastal
interjurisdictional arrangements of the kind that are already in place for open-water migratory
species like tuna [61] offer a possible template that can be built on. Our results demonstrate
the need for such agreements for nearshore species of fish and for individual provinces and
community managed areas within a single nation state, which was largely considered less
important in tropical context. For this, IRIIS can be a useful prioritization tool. Because the
management axis allows for dynamic mobility based on changes in policy, it can be used to
strategically evaluate which set of policy directions could reduce the susceptibility of the
widest number of fish populations across a shared coastline. As our analysis shows, the
inherent vulnerability of fish can be strongly mediated by what managers do at local levels,
and how closely they coordinate their actions with neighbors that share the same genetic
resource pool. This offers the principal challenge but also the best hope for rational
management of dwindling fish stocks.

### 4. Methods

We first describe – the Integrative Risk Indicator for Interjurisdictional Stocks (IRIIS) framework to assess the relative Risk of genetically distinct, interjurisdictional fish stocks based on an extension of a standard PSA. We then demonstrate the utility of this framework based on a case study of nine nearshore species along the west coast of India.

#### 4.1.1. Constructing the Integrative Risk Indicator for Interjurisdictional Stocks (IRIIS)

We propose that the risk of overharvest for the population of a target species is a function of i). the inherent life history and catchability of each species (PSA) and ii). the biological sustainability of management practices across all the jurisdictional regimes the population spans. Figure 5 outlines the rationale IRIIS assessment. The detailed description can be found in 4.1.1.a and 4.1.1.b

Legend: The matrix uses indicators like PSA analysis as a starting point to denote biological susceptibility (represented on the y axis). The management disparity axis (x axis) is based on an ordered set of decision rules that evaluate the jurisdictional overlap of each population, and the congruence of management practices in each jurisdiction with known species biology. Five colored bars represent five risks associated with management disparity which is increasing from left to right. The color gradient on each vertical bar represents overall risk of overexploitation.

Figure 5 – Rationale of development of Integrative Risk Indicator for Interjurisdictional Stocks
(IRIIS).

**Rationale for constructing the IRIIS**

**4.1.1.a. Vertical axis of biological susceptibility:** The vertical axis of the Integrative Risk
Indicator for Interjurisdictional Stocks (IRIIS) is the score assigned by classical Productivity
Susceptibility Analyses which typically assign a vulnerability score to their sampling units
(usually species) [16,18,23].

**4.1.1.b. Horizontal axis of management disparity:** The horizontal axis of the matrix is
premised on the observation that in addition to species-based vulnerability indicated by PSA
analysis, the vulnerability of a particular population is mediated by the degree to which the
spatial extent of the population matches the spatial extent of the management regime.
Therefore, populations can be grouped in two broad vulnerability categories i) Populations
located entirely within a single management jurisdiction ii) Population ranging across multiple
management jurisdictions. Note that the species-specific vulnerability (based on PSA) does
not change in these assessments between populations.

If the assessed population is entirely located within a single jurisdiction, we classify its
management-based vulnerability by calculating the percent disparity: Low disparity <33%;
High disparity >=33%. Management-based vulnerability increases as we move from bar 1 to
bar 2.

If a population spans multiple management regimes, we assess its interjurisdictional
management vulnerability by calculating the percent disparity of the population within each
jurisdiction separately. Based on the proportion of jurisdictions with more than 33% disparity
with species biology we assign the population in three assessment categories. 1) All
jurisdictions with <33 % disparity (vertical bar no. 3 in Figure 5), 2) majority of management
regimes with <33% disparity (vertical bar no. 4 in Figure 5), 3) Majority of management
regimes with >33% disparity (vertical bar no. 5 in Figure 5). Management-based vulnerability
increases from left to right.

We assume that populations entirely located in a single management jurisdiction overall
have an inherent lower vulnerability than populations spread across jurisdiction of multiple
management regimes due to inherent differences in implementation between jurisdictions
and the added complications of interjurisdictional co-ordination, if any.

### 4.1.2. Evaluating the biological relevance of present management regimes

For every management regime within the geographical area of the assessment, we first
catalogue all available policy instruments, regulations or subsidies related to the fishery. We
then evaluate each of these practices and policies based on their congruence (or disparity)
with sustainability goals based on what is known of species' biology. E.g. A policy to restrict
mechanized fishing during June-August, is biologically congruent with species that breed
during this period. Conversely, the policy would show biological disparity with species that do
not breed during this period. The proportion of all policy/practice attributes in that
management regime that do not align with the biology of the given species is used to
calculate a percent disparity score.

Table 1 demonstrates how we calculated the percent disparity for a combination of a
hypothetical province with only two policies related to harvest and a hypothetical species
which breeds between June and August and has Length at Maturity (L50) of 20 cm. For
simplicity, in the current assessment, we assume that all attributes have an equal weightage
on sustainable harvest. However, it is quite possible to assign different weights to each
attribute if there is a sufficient justification to do so. Weightage can also be assigned based
on reliable knowledge of how effectively each policy attribute is implemented on the ground.

Table 1 - Evaluating the biological efficacy of management practices for a hypothetical province with only two policies related to harvest and a hypothetical species which breeds between June and September, and has Length at Maturity (L50) of 20 cm

| Policy Number | Attribute and its intended consequence | Prescribed policy | Species biology | Assessment |
|---|---|---|---|---|
| 1 | Temporal/seasonal ban on mechanized fishing to protect breeding individuals | Ban period: June – August | Breeding period of target species: June- September | Congruence between species biology and policy attributes |
| 2 | Prescribed Minimum Legal Size (MLS) to protect juveniles from overfishing | No prescription (all size individuals are legal to catch) | Length at Maturity (L50) – 20 cm | Disparity between species biology and policy attribute |
| | Total disparity percentage | | | 50% policy attributes show disparity **– High disparity combination** |

445
446
447
448

### 4.2. Case study: West coast of India

We apply IRIIS to evaluate the vulnerability of populations of nine commercially important fish species along the west coast of India. We use SNP-based genomic data to identify species population boundaries for these target species and an analysis of management practices across all regimes to populate the matrix.

### 4.2.1. Study area

The study region extends from the western shores of the Kutch peninsula to Kanyakumari at the southern tip of India - a stretch of approximately 3400 km of coastline (Figure.6) The region represents a single eco-region based on its faunal community composition [62]. The governance of the coast is distributed across five states (Gujarat, Maharashtra, Goa, Karnataka and Kerala). Each state regulates its own fisheries with policy instruments in the form of state specific Marine Fisheries Regulation Acts (MFRAs). Such disjunct governance in a single ecoregion makes the west coast an ideal case study to understand how differences in fisheries management can mediate overharvest risk of individual fish populations.

### 4.2.2. Choice of species

Indian fisheries are typically multi species, with several species constituting the target catch.
For such fisheries, the comparison between biological boundaries of fish and the
jurisdictional boundaries of fisheries needs to happen for multiple species simultaneously.
Our attempt was to include as many fisheries target species as was logistically possible.

We identified nine fish species commonly caught in trawl fisheries along the west coast of
India. The choice of species was dependent on their ubiquity, the frequency with which they
occur in trawl catches across the west coast, and ease of identification in the field. They
were also chosen to represent different trophic groups, habitats and taxonomic groups.

The selected fish species were silver pomfret (*Pampus argenteus*), ribbonfish (*Trichiurus*
*lepturus*), Japanese threadfin bream (*Nemipterus japonicus*), spiny cheek grouper
(*Epinephelus diacanthus*), spade nose shark (*Scoliodon laticaudus)*, wolf herring
(*Chirocentrus nudus*), Indian oil sardine (*Sardinella longiceps),* ladyfish (*Sillago sihama*), and
Indian mackerel (*Rastrelliger kanagurta*).

**4.2.3. Calculating risk of overexploitation associated with species biology.**

Mohamed et al. in 2021 [18] formulated an Index of Resilience and Vulnerability (IRV) based
on the growth coefficient (K), longevity: asymptotic length (L∞), reproductive load, number of
spawning months (NSM), fecundity and mean trophic level (MTL) as vulnerability attributes
and species geographic distribution, body length depth ratio (BLD), exploitation ratio,
recruitment, susceptibility to gear and price as resilience attributes. Although not explicitly
framed as PSA, the IRV uses same basic principles. This is the most rigorous assessment of
species susceptibility of marine fish from the region. we used it to formulate our y axis which
represents risk of overexploitation associated with species biology.

We used IRV values for all nine species (Figure 3). Mohammad et al. 2021 did not assess
the IRV index of *Silago sihama*, so we calculated its IRV based on the formula given by
Mohammad et al 2021 (Supplementary Material, Section 1).

**4.2.4. Identifying the geographical extent of individual fish populations.**
491
**4.2.4.1. Data collection**

We sampled individuals of each species from ten sites located at roughly 200 km intervals
along the entire west coast from Gujarat to Kerala. We have listed the sites with the map in
Figure 6. At each site we located the places where fishing boats land their catch (landing
sites) and we sampled freshly landed fish from these landing sites. Since the study uses
genetic data to identify geographical boundaries between populations, we also sampled
individuals from Balasore - a single location on India's east coast. Balasore is located at
least 1700 km away from southernmost sampling locations on the west coast and was
expected to act as a separate population in the/our analysis.

Legend: Map of sampling location along the Indian coastline with the color legend for samples from specific locations.

Figure 6 – Location of sampling sites along the west coast of India

Tissue samples were collected using fin clips extracted from the fish directly at the landing
site. Fin clips of 10-18 individuals from each species were collected at each sampling
location using sterilized instruments to avoid cross-contamination, and stored in absolute
ethanol in a 2 ml tube. We ensured that the fish sampled were sourced from local waters by
sampling from boats where the sailors confirm that they were fishing in waters no further
than 50 km from sampling location. As an additional safeguard we sampled only boats with
lower engine capacity and cannot travel beyond 50-60 km from the landing site due to fuel
constraints. Sampling fish without considering their life stages posed the risk of mixing
individuals from two distinct populations, leading to potential misidentification. To avoid this,
we collected samples from only the adult population at each location.

**4.2.4.2.    Data generation**

We used genome-wide SNP (Single Nucleotide Polymorphism) markers to identify
genetically distinct populations and their geographical extent for all nine chosen species. We
chose SNP-based methods to get a high number of regions of differentiation across the DNA
to overcome the constraints of sample size [64]. We used the Double Digest Restriction
Associated DNA (ddRAD) approach to generate genomic data [65]. The sample collection
protocol along with detailed protocols of DNA extraction, library preparation is given in the
Supplementary Material, Section 3.

**4.2.4.3.    Data analysis**

We used three approaches for SNP calling: 1) For silver pomfret (*Pampus argenteus*) and
Indian mackerel (*Rastrelliger kanagurta*), we used available reference genomes from the
NCBI database for SNP calling (*Rastrelliger kanagurta* - GCA_024605255.1, *Pampus*

*argenteus* - GCA_000697985.1). 2) For spiny cheek grouper (*Epinephelus diacanthus*) and
spade nose shark (*Scoliodon laticaudus)* we used the reference genomes of the nearest
taxonomically related species. We used the *Epinephelus coioides* (GCA_900536245.1)
genome for *Epinephelus diacanthus* and *Scyliorhinus canicula* (GCA_902713615.1) genome
for *Scoliodon laticaudus* from the NCBI database. 3) There were no available reference
genomes that we could use for any of the other species, so we used a *de novo* variant
calling pipeline by STACKS. The detailed bioinformatic protocol (quality control of reads and
SNP discovery) are provided in Supplementary Material, Section 3 and 4.

After SNP discovery, we used three indices to identify evidence of genetic similarity and
dissimilarity between samples and ultimately identify genetically distinct populations for all
nine species. These indices were:
535
1. Visualization based on a principal component analysis (PCA) using PLINK [66].
2. Visualization based on admixture probabilities using the program ADMIXURE [67]. We ran
an admixture for K ranging from 1 to n, where n = number of sites in which we have
information of that particular species, for 10 replicates each, the best K was determined by
estimating the cross-validation error. The results were analyzed and plotted using a web-
based server, CLUMPAK [68].
3. A pairwise mean $F_{ST}$ calculation conducted using STACKS [69].
543

### 4.2.5.  Identifying disparity between species biology and management attributes

Across the west coast of India, we examined fisheries regulations and policy instruments
relevant to our study species. Following Gangal et. al [28], we identified nine different
regulations as the key management attributes of the regimes prevalent in the five states of
the Indian west coast [28]. They were 1) partial or complete fishing ban during monsoon
months (June-August) 2) maximum allowable engine capacity of a fishing vessel 3) minimum
allowable mesh size for cod end of trawl operation 4) minimum allowable mesh size for gill
nets 5) minimum allowable mesh size for purse nets / ring seine nets 6) minimum allowable
legal size for a species 7) ban on bull / paired trawling 8) ban on fishing using lights 9)
restriction on mechanized fishing in shallow waters. These nine regulations were used to
assess disparity between regulations and fish biology for each state. The details and
justifications for the percent disparity calculations are provided in detail in Supplementary
Material, section 2.

### 4.2.6.  Populating the IRIIS matrix

We first mapped all nine species based on their IRV values along the vertical axis. Species
with higher IRV values were positioned higher on the vertical axis. The horizontal
(management) axis was populated based on the geographic spread of each population in
relation to jurisdictional boundaries. Based on the geographic extent of each population, as
per rationale explained in figure 5, we classified populations into two categories - populations
ranging within a single management regime and populations ranging across multiple
management regimes.

Within the first category (populations ranging within a single management regime), we
further classified the populations into two subcategories (from left to right): 1. populations
with low disparity (disparity index less than 33%) and 2. populations with high disparity
(disparity index more than 33%).

In the second category, along the management axis (population ranging across multiple
management regimes), we classified populations along the horizontal axis into 3
subcategories (from left to right): 1. Low disparity in all the management regimes, 2. Low
disparity in majority of the management regimes, 3. High disparity in majority of the
management regimes.

**Resource availability statement**

The raw demultiplexed data generated in this study will be deposited in the Sequence Read
Archive (SRA) database of the online repository of NCBI (National Center for Biotechnology
Information) and the accession numbers will be provided upon acceptance. The references
of government regulations used for disparity analysis are provided as citations in
Supplementary material, section 2. All analytical scripts are included in the Supplementary
Materials

**Acknowledgements**

We thank all the fisherfolk, fish venders and retailors who allowed us to collect samples from
their fish. We thank Isha Bopardikar, Alisa Barns, Al Badush Gafoor, Rajeshwari BT and
Amit Nayak for help at the time of sample collection. We thank Meghana Natesh, Vinay
Sagar, Aparna Krishnan, Tejali Naik, Laxminarayan CP, Vinay Kumar, Gauri Gharpure for
their help in Lab work. We thank Shri Amm Murugappa Chettiar Research Centre,
Cholamandalam Investment and Finance Company Limited (Chola) and Mr.Arvind Dattar for
providing financial assistance for the project. We also thank Central Marine Fisheries
Research Institute, Department of Fisheries – Government of India, Department of Fisheries
of Gujarat, Maharashtra, Goa, Karnataka and Kerala for publicly available information on
policies, schemes and subsidies and rules and regulations.

**Authors contributions**

MG - Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation,
Methodology, Project administration, Validation, Visualization, Writing – original draft, Writing
– review & editing

RA - Conceptualization, Funding acquisition, Methodology, Project administration,
Supervision, Validation, Visualization, Writing – review & editing

VS - Methodology, Supervision, Validation, Writing – review & editing

AP - Formal analysis, Methodology, Resources, Software, Validation

AT - Formal analysis, Methodology, Software, Validation

PP - Formal analysis, Software, Validation

MS - Data curation, Writing – review & editing

IM - Data curation, Writing – review & editing

JK - Methodology, Supervision, Validation, Writing – review & editing

UR - Conceptualization, Methodology, Project administration, Resources, Supervision,
Validation, Writing – review & editing

**Declaration of Interest**

The authors declare no competing interests

**Supplemental information**

Document S1. Figures S1–S13, Tables S1 to S8, and supplemental references

**Figures**

Figure 1. Principal Component plots for nine chosen species across the west coast of India

Legend:

Panel A – Map of sampling location along the Indian coastline with the colour legend for
samples from specific locations. Porbandar = Dark Red, Surat = Red, Mumbai = Light Red,
Harne = Orange, Vengurle = Light yellow, Goa = Yellow, Kumta = Light Cyan, Mangalore =
Light Blue, Kozhikode = Medium Blue, Alappuzha = Blue, Odisha = Black. Panel B to J –
Principal Component Analysis plot of individual species with information about number of
SNPs and important filters used for variant calling. See also figures S5 to S13 in
supplementary material

Figure 2. Disparity analysis for Indian oil sardine, with populations housed in single as well
as multiple management regimes

Legend: Each donut represents a state along the west coast of India. Nine sections are for
nine different regulations and the color of each section represents congruence or disparity of
the sardine biology with the regulation. Legend for all the symbols and colors are embedded
in the figure. See also figure S2 in supplementary material

Figure 3 - Integrative Risk Indicator for Interjurisdictional Stocks (IRIIS). for nine species
along the west coast of India

Legend: Legend: Five colored bars represent five risks associated with management
disparity which is increasing from left to right. vertical axis of each bar represents IRV score.
For species presented on the matrix the IRV score is written on the left-hand side. The color
gradient on each vertical bar represents overall risk of overexploitation. Different population
of each species is indicated using maps in legend section. See also figure S2 in the
supplementary materials

Figure 4.  Shift in overexploitation risk of the peninsular population of *Sardinella longiceps*
along the management disparity axis as a direct result of changing management practices

Legend: Legend: Five colored bars represent five risks associated with management
disparity which is increasing from left to right. Vertical axis of each bar represents IRV score.
For species presented on the matrix the IRV score is written on the left-hand side. The color
gradient on each vertical bar represents overall risk of overexploitation. See also table S2,
S3 and S5 in supplementary materials

Figure 5 – Rationale of development of Integrative Risk Indicator for Interjurisdictional Stocks
(IRIIS).

Legend: Legend: The matrix uses indicators like PSA analysis as a starting point to denote
biological susceptibility (represented on the y axis). The management disparity axis (x axis)
is based on an ordered set of decision rules that evaluate the jurisdictional overlap of each
population, and the congruence of management practices in each jurisdiction with known
species biology. Five colored bars represent five risks associated with management disparity
which is increasing from left to right. The color gradient on each vertical bar represents
overall risk of overexploitation.

Figure 6 – Location of sampling sites along the west coast of India

Legend: Legend: Map of sampling location along the Indian coastline with the color legend
for samples from specific locations.

**Tables**

Table 1 - Evaluating the biological efficacy of management practices for a hypothetical
province with only two policies related to harvest and a hypothetical species which breeds
between June and September, and has Length at Maturity (L50) of 20 cm

**References**

1.      Huxley, F.R.S. (1883). Inaugural meeting of the fishery congress. Address. Gt. Int.
Fish. Exhib., 1–22.

2.      FAO (2023). SDG INDICATOR 14.4.1 - Proportion of fish stocks within biologically
sustainable levels Target. In Tracking progress on food and agriculture-related SDG
indicators 2023, pp. 107–110. https://doi.org/https://doi.org/10.4060/cc7088en-fig55.

3.    FAO (2024). The State of World Fisheries and Aquaculture 2024 – Blue
Transformation in action https://doi.org/https://doi.org/10.4060/cd0683en.

4.    Teh, L.C.L., and Sumaila, U.R. (2011). Contribution of marine fisheries to worldwide
employment. Fish Fish. *14*, 77–88. https://doi.org/10.1111/j.1467-2979.2011.00450.x.

5.    Campling, L., Havice, E., and Mccall Howard, P. (2012). The Political Economy and
Ecology of Capture Fisheries: Market Dynamics, Resource Access and Relations of
Exploitation and Resistance. J. Agrar. Chang. *12*, 177–203.
https://doi.org/10.1111/j.1471-0366.2011.00356.x.

6.    OECD (2025). OECD Review of Fisheries 2025.

7.    Westholm, A., and Argüello, G. (2023). Dynamic Ocean Management in Areas
Beyond National Jurisdiction. Ocean Dev. Int. Law *54*, 448–468.
https://doi.org/10.1080/00908320.2023.2296392.

8.    Warren, L.M. (2023). Property rights and marine protected areas. Environ. Law Rev.
*25*, 234–245. https://doi.org/10.1177/14614529231187745.

9.    Hung, P.Y. (2025). Bordered-In, Bordered-Out, and Overlapping Territorialities in
Ocean Space: The Case of Fisheries. In Ocean Governance ( Beyond ) Borders, K.
Peters and J. Turner, eds. (Palgrave macmillan), pp. 75–98.

10.    Hilborn, R., Oscar, R., Anderson, C.M., Baum, J.K., and Branch, T.A. (2020). Effective
fisheries management instrumental in improving fish stock status. PNAS *117*, 2218–
2224. https://doi.org/10.1073/pnas.1909726116.

11.    Costello, C., Ovando, D., Hilborn, R., Gaines, S.D., Deschenes, O., and Lester, S.E.
(2012). Status and Solutions for the World's Unassessed Fisheries. Science (80-. ).
*338*, 517–520. https://doi.org/10.7551/mitpress/3629.003.0022.

12.    FAO (2020). The State of World Fisheries and Aquaculture 2020. Sustainability in
action.

13.    Hilborn, R., Amoroso, R.O., Anderson, C.M., Baum, J.K., Branch, T.A., Costello, C.,
De Moor, C.L., Faraj, A., Hively, D., Jensen, O.P., et al. (2020). Effective fisheries
management instrumental in improving fish stock status. PNAS *117*, 2218–2224.
https://doi.org/10.1073/pnas.1909726116.

14.    Rosenberg, A.A., Acosta, A., Babcock, E., Harrington, J., Hobday, A., Mogensen,
C.B., O'Boyle, R., Rader, D., Swasey, J.H., Trumble, R.J., et al. (2009). Use of
Productivity-Susceptibility Analysis ( PSA ) in Setting Annual Catch Limits for U. S.
Fisheries : A Workshop Report.

15.    Patrick, W.S., Spencer, P., Ormseth, O., Cope, J., Field, J., Kobayashi, D., Gedamke,
T., Cortes, E., Bigelow, K., Overholtz, W., et al. (2009). Use of Productivity and
Susceptibility Indices to Determine Stock Vulnerability , with Example Applications to
Six U . S . Fisheries.

16.    Duffy, L.M., and Griffiths, S.P. (2019). Assessing attribute redundancy in the
application of productivity- susceptibility analysis to data-limited fisherie. Aquat. Living
Resour. *32*, 1–11. https://doi.org/https://doi.org/10.1051/alr/2019018.

17.    Faruque, H., and Matsuda, H. (2021). Assessing the vulnerability of bycatch species

from Hilsa gillnet fishing using productivity susceptibility analysis: Insights from
Bangladesh. Fish. Res. *234*, 105808. https://doi.org/10.1016/j.fishres.2020.105808.

18.    Mohamed, K.S., Sathianandan, T.V., Vivekanandan, E., Kuriakose, S., Ganga, U.,
714           Pillai, S.L., and Nair, R.J. (2021). Application of biological and fisheries attributes to
715           assess the vulnerability and resilience of tropical marine fish species
716           https://doi.org/10.1371/journal.pone.0255879.

717    19.    Leach, A., Kell, L.T., and Mumford, J. (2023). Productivity-Susceptibility Analysis tool.
718           ICCAT *80*, 98–116.

719    20.    Laikre, L., Palm, S., and Ryman, N. (2005). Genetic Population Structure of Fishes :
720           Implications for Coastal Zone Management. AMBIO A J. Hum. Environ. *34*, 111–119.
721           https://doi.org/http://dx.doi.org/10.1579/0044-7447-34.2.111.

722    21.    Palsbøll, P.J., Berube, M., and Allendorf, F.W. (2006). Identification of management
723           units using population genetic data. TRENDS Ecol. Evol. *22*, 11–16.
724           https://doi.org/10.1016/j.tree.2006.09.003.

725    22.    Waples, R.S., Punt, A.E., and Cope, J.M. (2008). Integrating genetic data into
726           management of marine resources: how can we do it better? Fish Fish. *9*, 423–449.

727    23.    Patrick, W.S., Spencer, P., Link, J., Cope, J., Field, J., Kobayashi, D., Lawson, P.,
728           Gedamke, T., Cortés, E., Ormseth, O., et al. (2010). Using productivity and
729           susceptibility indices to assess the vulnerability of United States fish stocks to
730           overfishing. Fish. Bull. *108*, 305–322.

731    24.    Arceo, H.O., Cazalet, B., Alin, P.M., Mangialajo, L., and Francour, P. (2013). Moving
732           beyond a top-down fisheries management approach in the northwestern
733           Mediterranean: Some lessons from the Philippines. Mar. Policy *39*, 29–42.
734           https://doi.org/10.1016/j.marpol.2012.10.006.

735    25.    Jaya, I., Satria, F., Wudianto, Nugroho, D., Sadiyah, L., Buchary, E.A., White, A.T.,
736           Franklin, E.C., Courtney, C.A., Green, G., et al. (2022). "Are the working principles of
737           fisheries management at work in Indonesia?" Mar. Policy *140*, 105047.
738           https://doi.org/https://doi.org/10.1016/j.marpol.2022.105047.

739    26.    Monteferri, B., and Muller, M.R. (2020). Marine, coastal and fisheries issues in Chile,
740           Mexico and Peru: An initial institutional and policy review. In Marine and Fisheries
741           Policies in Latin America : A Comparison of Selected Countries, M. R. Muller, R.
742           Oyanedel, and B. Monteferri, eds. (Routledge. Taylor and Francis Group), pp. 6–14.

743    27.    Nakamura, J., and Hazin, F. (2020). Assessing the Brazilian federal fisheries law and
744           policy in light of the Voluntary Guidelines for Securing Sustainable Small-scale
745           fisheries. Mar. Policy *113*, 103798. https://doi.org/10.1016/j.marpol.2019.103798.

746    28.    Gangal, M., Suri, V., and Arthur, R. (2023). How well does Indian fisheries policy
747           engage with fisheries biology? Exploring the science-policy interface of coastal
748           capture fisheries along the west coast of India. Mar. Policy *156*, 105796.
749           https://doi.org/10.1016/j.marpol.2023.105796.

750    29.    Berger, A.M., Deroba, J.J., Bosley, K.M., Goethel, D.R., Langseth, B.J., Schueller,
751           A.M., and Hanselman, D.H. (2021). Incoherent dimensionality in fisheries
752           management: consequences of misaligned stock assessment and population
753           boundaries. ICES J. Mar. Sci. *78*, 155–171. https://doi.org/10.1093/icesjms/fsaa203.

30. Torre-castro, M. De (2012). Governance for Sustainability☐: Insights from Marine
Resource Use in a Tropical Setting in the Western Indian Ocean. Coast. Manag. *40*,
612–633. https://doi.org/10.1080/08920753.2012.727738.

31. Fisheries-and-ports(B)-Department-Kerala (2018). Kerala Marine Fisheries Regulation
Act Rule 2018 (Government of Kerala).

32. Department-of-Fisheries-karnataka (2018). Minimum legal size (Department of
Fisheries, Government of Karnataka).

33. Department-of-Fisheries-Goa (2021). No. DF/ENF/NOTI-IMPL-MLS/2020-21/5042
(Directorate of Fisheries, Government of Goa).

34. Law-and-Judiciary-Department-Maharashtra (2021). Maharashtra Marine Fishing
Regulation (Amendment) Act, 2021 (Law and Judiciary Department. Government of
Maharashtra).

35. No. DF/ENF/ORDER/2016-17 (2016). (Department of Fisheries, Directorate of
Fisheries, Goa).

36. Department-of-Fisheries-karnataka (2014). Ban on bull trawling (Department of
Fisheries, Government of Karnataka).

37. Ghosh, S., Mohanraj, G., Asokan, P.K., Dhokia, H.K., Zala, M.S., and Bhint, H.M.
(2009). Fishery and stock estimates of the silver pomfret , Pampus argenteus (
Euphrasen ), landed by gill netters at Veraval. Indian J. Fish. *56*, 177–182.

38. Gladston, Y., Devi, M.S., and Xavier, K.A.M. (2016). Design and gillnet selectivity of
Pampus argenteus along the Satpati coast, Maharashtra, India. Reg. Stud. Mar. Sci.
https://doi.org/10.1016/j.rsma.2016.12.006.

39. Barnes, A., Sutaria, D., Harry, A. V., and Jabado, R.W. (2018). Demographics and
length and weight relationships of commercially important sharks along the north-
western coast of India. Aquat. Conserv. Mar. Freshw. Ecosyst. *28*, 1374–1383.
https://doi.org/10.1002/aqc.2940.

40. Sen, S., Chakraborty, S.K., Vivekanandan, E., Zacharia, P.U., Kizhakudan, S.J.,
Jaiswar, A.K., Dash, G., G, J., and Bharadiya, S.A. (2019). Population dynamics and
stock assessment of spadenose shark Scoliodon laticaudus Muller and Henle 1839
along Gujarat coast of India. Indian J. Geo Mar. Sci. *48*, 423–433.

41. Ramesh, N., Rising, J.A., and Oremus, K.L. (2019). The small world of global marine
fisheries: The cross-boundary consequences of larval dispersal. Science (80-. ). *364*,
1192–1196.

42. Palacios-Abrantes, J., Reygondeau, G., Wabnitz, C.C.C., and Cheung, W.W.L.
(2020). The transboundary nature of the world's exploited marine species. Sci. Rep.
*10*, 17668. https://doi.org/10.1038/s41598-020-74644-2.

43. Corte´s, E., Brooks, E.N., and Shertzer, K.W. (2015). Risk assessment of
cartilaginous fish populations. ICES J. ofMarine Sci. *72*, 1057–1068.
https://doi.org/doi:10.1093/icesjms/fsu157.

44. Baillargeon, G.A., Tlusty, M.F., Dougherty, E.T., and Rhyne, A.L. (2020). Improving
the productivity – susceptibility analysis to assess data-limited fisheries. Mar. Ecol.

Prog. Ser. *644*, 143–156. https://doi.org/10.3354/meps13362.

45.  Hoshino, E., Pascoe, S., Hutton, T., and Kompas, T. (2018). Estimating maximum
economic yield in multispecies fisheries: a review. Rev. Fish Biol. Fish. *28*, 261–276.
https://doi.org/10.1007/s11160-017-9508-8.

46.  Legović, T., Klanjšček, J., and Geček, S. (2010). Maximum sustainable yield and
species extinction in ecosystems. Ecol. Modell. *221*, 1569–1574.
https://doi.org/10.1016/j.ecolmodel.2010.03.024.

47.  Hilborn, R. (2019). Food for Thought Measuring fisheries performance using the "
Goldilocks plot ." ICES J. Mar. Sci. *76*, 45–49. https://doi.org/10.1093/icesjms/fsy138.

48.  Lado, E.P. (2016). Why the common fisheries policy is important. In The Common
Fisheries Policy: The Quest for Sustainability (John Wiley & Sons, Ltd.), pp. 1–17.
https://doi.org/10.1002/9781119085676.

49.  Larkin, P.A. (1981). A Perspective on Population Genetics and Salmon Management.
Can. J. Fish. Aquat. Sci. *38*, 1469–1475. https://doi.org/10.1139/f81-198.

50.  Reiss, H., Hoarau, G., Dickey-Collas, M., and Wolff, W.J. (2009). Genetic population
structure of marine fish: Mismatch between biological and fisheries management
units. Fish Fish. *10*, 361–395. https://doi.org/10.1111/j.1467-2979.2008.00324.x.

51.  Chávez, C., Dresdner, J., González, N., Leiva, M., and Quiroga, M. The performance
of shared fish stock fisheries under varying institutional and socioeconomic
conditions: Evidence from the South Eastern Pacific Anchoveta Fishery. Mar. Policy.
https://doi.org/10.1016/j.marpol.2020.104262.

52.  Yeeting, A.D., Weikard, H.P., Bailey, M., Ram-Bidesi, V., and Bush, S.R. (2018).
Stabilising cooperation through pragmatic tolerance: the case of the Parties to the
Nauru Agreement (PNA) tuna fishery. Reg. Environ. Chang. *18*, 885–897.
https://doi.org/10.1007/s10113-017-1219-0.

53.  Gagern, A., and Bergh, J. Van Den (2013). A critical review of fishing agreements with
tropical developing countries. Mar. Policy *38*, 375–386.

54.  McBride, R.S. (2014). Managing a Marine Stock Portfolio: Stock Identification,
Structure, and Management of 25 Fishery Species along the Atlantic Coast of the
United States. North Am. J. Fish. Manag. *34*, 710–734.
https://doi.org/10.1080/02755947.2014.902408.

55.  Arnason, R., and Runolfsson, B.T. (2023). Delineating the impacts of strong user
rights in fisheries. Mar. Policy *155*, 105666.
https://doi.org/https://doi.org/10.1016/j.marpol.2023.105666.

56.  Aceves-Buenoa, E., Cornejo-Donoso, J., Miller, S.J., and Gaines, S.D. (2017). Are
Territorial Use Rights in Fisheries (TURFs) sufficiently large? Mar. Policy *78*, 189–
195. https://doi.org/10.1016/j.marpol.2017.01.024.

57.  Song, A.M., Scholtens, J., Stephen, J., Bavinck, M., and Chuenpagdee, R. (2017).
Transboundary research in fisheries. Mar. Policy *76*, 8–18.
https://doi.org/10.1016/j.marpol.2016.10.023.

58.  Swartz, W., Sala, E., Tracey, S., Watson, R., and Pauly, D. (2010). The spatial

expansion and ecological footprint of fisheries (1950 to present). PLoS One *5*,
e15143. https://doi.org/10.1371/journal.pone.0015143.

59.　Department-of-Animal-Husbandry-Dairying-and-Fisheries (2017). National policy on
Marine Fisheries 2017 (Ministry of agriculture and Farmers Welfare, Government of
India).

60.　Department-of-Fisheries (2020). National Fisheries Policy, 2020 (Ministry of Fisheries,
Animal Husbandry and Dairying, Government of India).

61.　Cullis-Suzuki, S., and Pauly, D. (2010). Failing the high seas: A global evaluation of
regional fisheries management organizations. Mar. Policy *34*, 1036–1042.
https://doi.org/10.1016/j.marpol.2010.03.002.

62.　Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M.,
Halpern, B.S., Jorge, M.A., Lombana, A., Lourie, S.A., et al. (2007). Marine
ecoregions of the world: A bioregionalization of coastal and shelf areas. Bioscience
*57*, 573–583. https://doi.org/10.1641/B570707.

63.　Froese, R., and D. Pauly (2024). Sillago sihama summary page. Fishbase.
https://fishbase.se/popdyn/KeyfactsSummary_1.php?ID=4544&GenusName=Sillago&
SpeciesName=sihama&vStockCode=4729&fc=307.

64.　Willing, E., Dreyer, C., and Oosterhout, C. Van (2012). Estimates of Genetic
Differentiation Measured by F ST Do Not Necessarily Require Large Sample Sizes
When Using Many SNP Markers. PLoS One *7*, e42649.
https://doi.org/10.1371/journal.pone.0042649.

65.　Tyagi, A., Yadav, N., Pandit, A., and Ramakrishnan, U. (2024). On the road to losing
connectivity: Fecal samples provide genome-wide insights into anthropogenic impacts
on two large herbivore species in central India. Mol. Ecol., 1–17.
https://doi.org/10.1111/mec.17461.

66.　Purcell, S., Neale, B., Todd-brown, K., Thomas, L., Ferreira, M.A.R., Bender, D.,
Maller, J., Sklar, P., Bakker, P.I.W. De, Daly, M.J., et al. (2007). PLINK□: A Tool Set
for Whole-Genome Association and Population-Based Linkage Analyses. Am. J.
Hum. Genet. Vol. *81*, 559–575. https://doi.org/10.1086/519795.

67.　Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of
ancestry in unrelated individuals. Genome Res., 1655–1664.
https://doi.org/10.1101/gr.094052.109.vidual.

68.　Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., and Mayrose, I. (2015).
CLUMPAK: a program for identifying clustering modes and packaging population
structure inferences across K. Mol. Ecol. Resour. *15*, 1179–1191.
https://doi.org/10.1111/1755-0998.12387.

69.　Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., and Cresko, W.A. (2013).
Stacks: An analysis tool set for population genomics. Mol. Ecol. *22*, 3124–3140.
https://doi.org/10.1111/mec.12354.

# *Sardinella longiceps (Indian oil sardine)*

| Population housed in single management unit – Gujarat population | Population housed in multiple management units – Peninsular India population | | | |
|---|---|---|---|---|
|

 |

 |

 |

 |

 |
| High Disparity | Low disparity | Low disparity | Low disparity | Low disparity |

# Legend

 Mesh size regulation for purse nets

 Monsoon fishing ban

 Maximum engine capacity

 Mesh size regulation for cod-end net of trawler

 Mesh size regulation for gillnets

 Minimum Legal size

 Ban on Bull trawling

 Restriction on fishing in shallow waters

 Ban on light fishing

 Disparity between species biology and regulation

 Congruence between species biology and regulation

 Low disparity with percent disparity value written

 High disparity with percent disparity value written

# lowest vulnerability

Biological susceptibility (IRV index)

0.31
0.28
0.25
0.24
0.23
0.22
0.19

disparity low | disparity high | Low disparity in all management regimes | Low disparity in majority of management regimes | High disparity in majority of management regimes

**Single management regime** — **Multiple management regimes**

# highest vulnerability

Management disparity

---

# Legend

## Genetically distinct populations

| *Rastrelliger kanagurta* | *Chirocentrus nudus* | *Epinephelus diacanthus* |
|---|---|---|
| Single west coast population | Single west coast population | Single west coast population |
| *Trichiurus lepturus* | *Nemipterus japonicus* | *Pampus argenteus* |
| Single west coast population | Single west coast population | Gujarat - Maharashtra population / South Maharashtra to Kerala population |
| *Sardinella longiceps* | *Scoliodon laticaudus* | *Sillago sihama* |
| Gujarat Population / Peninsular population | Gujarat population / Maharashtra- Goa – Karnataka Population / Kerala population | Peninsular population / Kumta population |

## Overexploitation risk

Low risk

High risk

lowest
vulnerability

0.31

Biological susceptibility
(IRV index)

| 2023 | | | |
|------|----|----|----|
| MH | GO | KR | KL |

| 2016 | | | |
|------|----|----|----|
| MH | GO | KR | KL |

0.28

0.25
0.24
0.23

0.22

0.19

Disparity low

Disparity High

Disparity in all
management regimes

Low Disparity in
majority of
management regimes

High Disparity in
majority of
management regimes

highest
vulnerability

Single management regime    Multiple management regimes

Management disparity

Legend

MH – Maharashtra      GO – Goa      KR – Karnataka      KL – Kerala

XX – high disparity percent in that state      XX – Low disparity percent in that state