Title - Unbounding fish: Assessing vulnerability to overexploitation for transboundary
fish populations

Authors: Mayuresh Gangal[1,2,3], Rohan Arthur[2], Venkatachalam Suri[2], Awadhesh Pandit[3]
Abhinav Tyagi[3], Praveen Prakash[3], Mihir Sule[4], Iravatee Majgaonkar[5], Jagadish
Krishnaswamy[6], Uma Ramakrishnan[3]

Corresponding author – Mayuresh Gangal, Email: mayuresh@ncf-india.org

Affiliations:

1. Manipal Academy of Higher Education, Manipal, India.
2. Nature Conservation Foundation, Mysore, India
3. National Centre for Biological Sciences, Bengaluru, India
4. Independent Researcher, Mumbai, India
5. Ashoka Trust For Research In Ecology And The Environment, Bengaluru, India
6. Indian Institute for Human Settlements, Bengaluru, India

Abstract

Tropical fish populations often traverse management boundaries, making it difficult to evaluate
species vulnerabilities and optimise sustainable fishing goals. Most vulnerability assessments
rely on life histories and gear susceptibility, but ignore transboundary differences in
management, which can strongly affect vulnerability. We propose a novel vulnerability matrix,
the Biology – Transboundary Management (BTM) index for individual fish populations based
on 1) Species specific vulnerability 2) distribution of genetically-distinct populations with
respect to jurisdictional boundaries and 3) sustainability of management practices across
jurisdictions. Combining single-nucleotide polymorphism (SNP) analyses and policy
assessments, we evaluate relative vulnerabilities of nine commercially relevant fish
populations over a 3400km stretch of India's west coast, across multiple jurisdictions. The
BTM matrix indicates that transboundary differences in management strongly influences
biological vulnerability. The matrix is sensitive to population-level differences and policy
changes. Our analyses emphasise the need for population-level assessments and
transboundary cooperation in achieving sustainable multi species fisheries.

Introduction

The global defaunation of biodiversity once characteristic of terrestrial ecosystems, now
pervades the seas and oceans as well. Historically described as an inexhaustible natural
resource[1], marine fish today face a crisis of overfishing[2]. Recent global analyses indicate that
of 4,713 global fish stocks (77%) show worrying declines in catch despite increasing fishing
efforts[3]. Some groups like elasmobranchs are more susceptible than others. More than 390
species of elasmobranchs are threatened with extinction[4]. With the livelihoods of ~260 million
people involved in marine fisheries worldwide, this is not merely an ecological problem, but a
crisis of food security, livelihoods and equity[5]. This crisis is particularly acute in the global
south, where dependent coastal communities target a diverse community of fish, and where

the resources available to adequately monitor and regulate the fishery are limited[6]. Finding
ways to effectively manage fish resources at spatio-temporal scales that are biologically
meaningful, economically viable and managerially effective is the principal challenge
confronting global fisheries[7].

Bio-economic harvest models based on Maximum Sustainable Yield (MSY) calculations are
the most widely used tool for fisheries management[8,9]. However, this approach has been
widely criticised because it is simplistic and does not take into account differences in life history
traits, ecological factors, and variations between regions[8–10]. Also, while this approach may
work for managing single species typical of many temperate fisheries, its applicability to
complex multi-species tropical fisheries is questionable[8,11]. Identifying biologically and
ecologically relevant approaches that work for the tropics is crucial to inform fisheries policies
and help managers protect tropical fish stocks.

In evaluating the sustainability of fisheries, most recent work has focused on matching
fisheries effort with the productivity of each fish stock based on its natural susceptibility to
resource offtake[12–17]. These approaches fall under the broad umbrella of Productivity
Susceptibility Analyses or PSA. They attempt to be biologically comprehensive, incorporating
population growth rates, trophic level, breeding strategies and other crucial elements of fish
biology in their assessments[13]. A typical PSA approach incorporates multiple species
harvested in the area of concern, making them much more appropriate than MSY approaches
for multi species fisheries. They are also much more realistic about fisheries practices and the
susceptibility of fish species to different gear.

To bridge the gap between being an analytical useful tool to one that is directly applicable to
fisheries policy, a few limitations of the PSA approach need to be addressed. For most
available PSA's, the unit of analysis is the geographic range of the species (but see Patric et
al. 2010[18]) within the area of concern[12,14–17]. While this is a convenient simplification, it has
been long understood that, to be effective, genetically distinct populations of the same species
should be managed as separate management units[19–21]. Not incorporating this population
level information in PSA analyses significantly limits their applicability to real-world fisheries
management.

Along with species-centric analyses, most PSAs also assume a uniform harvest regime along
the entire geographical range of analysis. However, for coastal and nearshore fisheries this
assumption is seldom true. Apart from the EU, most countries manage the coastal waters
under their jurisdiction with little transboundary coordination. This transboundary problem
repeats within national borders as well. In most tropical countries with large coastlines like
Brazil, Indonesia, Philippines, India and Mexico, coastal fisheries are often decentralised and
governed by provinces which have significant autonomy in terms of fisheries management[22–
25,26]. Rarely is there a uniform harvest regime across the geographical area considered within
a PSA.

Between the multiple genetically distinct populations of a species and decentralised harvest
regimes there is often a mismatch between the biological boundaries of fish populations and
the management boundaries of fisheries. These mismatches lead to an increased risk of local
depletion of populations[27]. The consequences of these mismatches have been well
documented in the North Eastern Atlantic, where species like the Atlantic cod (*Gadus*

morhua)[28] and the North Atlantic mackerel (*Scomber scombrus*)[29] have experienced serious
overharvest and overfishing as a result of a misalignment between management and biological
boundaries.

Here we propose population-wise vulnerability matrix as a heuristic extension of the PSA by
incorporating 1) the inherent resilience and susceptibility attributes of species characteristic of
a typical PSA analysis, 2) the biological relevance of present management regulations[26] and
3) information on whether the population is managed through single or multiple management
regimes. As a case study, we present a vulnerability assessment for nine commercially
important fish species harvested along the west coast of India. Marine Fisheries in India is a
paradigm case of a multi species multi-management fishery where individual states
(provinces) manage fisheries in their waters semi-autonomously with little dialogue amongst
them. The current fishery is managed within provincial waters (Indian states) with little regard
to the underlying geographical spread of fish populations[26]. This makes Indian fisheries an
ideal system in which to test the application and relevance of the combined vulnerability matrix.
We used single nucleotide polymorphism (SNP) analysis of the target species to identify their
population boundaries along the 3400 km stretch of the west coast of India. We combined this
with a detailed analysis of management policies across all west-coast states to determine; i)
how closely these policies and management regimes matched the biological susceptibility of
the species and ii) whether the genetically contiguous population straddled a single or multiple
management regime. This information was used to construct a population-wise vulnerability
matrix. We then discuss the utility of this matrix in guiding policy and management decisions.

Methods

We first describe a heuristic framework to assess the relative vulnerability of genetically
distinct, transboundary fish stocks based on an extension of a standard PSA analysis. We
then demonstrate the utility of this framework based on a case study of species from an
extensive coastline along the west coast of India. We use Single Nucleotide Polymorphism
(SNP) analysis to identify the genetic boundaries of nine commercially targeted species and
determine their population-level vulnerability after evaluating management practices across
the range of each population.

1.   Constructing the Biology- Transboundary management vulnerability matrix (BTM)

We propose that the vulnerability of fish populations to overharvest is a function of both, the
inherent life history and catchability of each species (PSA), as well as number of management
regimes and their impacts that the population is exposed to across its range. This
management associated vulnerability is determined largely by (i) how prevalent management
practices align with sustainability based on known species biology and (ii) different
management and harvest regimes that population is subject to across its geographical
distribution. Fig. 5 outlines a schematic representation of this matrix and the Rationale section
details the steps we employed in its construction.

Evaluating the biological relevance of present management regimes

We first evaluate the congruence of current management practices with known species
biology. For every management regime within the geographical area of the assessment we

first catalogue all available policy instruments, regulations or subsidies related to the fishery.
These can also encompass catch and effort attributes of a particular fishing regime. We then
evaluate each of these practices and policies based on their congruence (or disparity) with
sustainability goals based on what is known of species' biology. We then assess the overall
biological veracity of the management regime in question to the assessed population by
calculating a percent disparity. The percent disparity is the proportion of all policy/practice
attributes that do not align with the biology of the assessed population of a given species
presented in percentage form. For simplicity, in the current assessment, we assume that all
attributes have an equal weightage on sustainable harvest. However, it is quite possible to
assign different weights to each attribute if there is a sufficient justification to do so.
Weightages can also be assigned based on reliable knowledge of how effectively each policy
attribute is implemented on the ground. Table 1 provides a tabular explanation of this
procedure with the help of a hypothetical example.

Table 1 - Evaluating the biological veracity of present management regime

Sr. No	Attribute and its intended consequence	Prescribed policy	Species biology	Assessment
1	Temporal/seasonal ban on mechanised fishing to protect breeding individuals	Ban period: June – August	Breeding period of target species: June to September	Congruence between species biology and policy attributes
2	Prescribed Minimum Legal Size (MLS) to protect juveniles from overfishing	No prescription (all size individuals are legal to catch)	Length at Maturity (L50) – 20 cm	Disparity between species biology and policy attribute
	Total disparity percentage			50% policy attributes show disparity **– High disparity combination**

Rationale for constructing the BTM vulnerability matrix

Figure 1 – Steps in the construction of the Biology - Transboundary Management (BTM) vulnerability matrix.

Legend: Silhouettes represent different fish species while different colours represent genetically distinct populations within the study area

Rationale for construction of proposed vulnerability matrix

Figure 1A: The vertical axis of the Biology - Transboundary vulnerability matrix is the score assigned by classical Productivity Susceptibility Analyses which typically assign a vulnerability score to their sampling units (usually species)[14,16,18] (figure 1A).

Figure 1B With the help of genetic tools, we identify and separately depict distinct populations in the study area. In species-based PSAs these populations would occupy the same positions on the vulnerability axis as shown in figure 1B.

Figure 1C The horizontal axis of the matrix is premised on the observation that in addition to species-based vulnerability indicated by PSA analysis, the vulnerability of a particular population is mediated by the degree to which the spatial extent of the population matches the spatial extent of the management regime. Therefore, populations can be grouped in two broad vulnerability categories i) Populations located entirely within a single management jurisdiction ii). Note that the species-specific vulnerability (based on PSA) does not change in these assessments between populations.

Figure 1D If the assessed population is entirely located within a single jurisdiction, we classify
its management-based vulnerability by calculating the percent disparity: Low disparity <33%;
High disparity >=33%. Management-based vulnerability increases as we move from bar 1 to
bar 2.

Figure 1E If a population spans multiple management regimes, we assess its transboundary
management vulnerability by calculating the percent disparity of the population within each
jurisdiction separately. Based on the proportion of jurisdictions with more than 33% disparity
with species biology we assign the population in three assessment categories. 1) All
jurisdictions with <33 % disparity (vertical bar no. 3 in figure 1E), 2) majority of management
regimes with <33% disparity (vertical bar no. 4 in figure 1E), 3) Majority of management
regimes with >33% disparity (vertical bar no. 5 in figure 1E). Management-based vulnerability
increases from left to right.

Figure 1F The Biology - Transboundary Management (BTM) vulnerability matrix combines
management-based vulnerability assessments of i) Populations that are entirely located under
jurisdiction of single management regime (Figure 1D) ii) Populations that are spread across
justification of multiple management regimes (Figure 1E), keeping the common axis of
species-specific vulnerability (as shown in Figure 1C). We assume that populations entirely
located in a single management jurisdiction have an inherently overall lower vulnerability than
populations spread across jurisdiction of multiple management regimes due to inherent
differences in implementation between jurisdictions and the added complications of
transboundary co-ordination, if there is any.

2. Case study: West coast of India

We apply the BTM matrix to evaluate the vulnerability of populations of nine commercially
important fish species along the west coast of India. We use SNP based genomic data to
identify species population boundaries for these target species and an analysis of
management practices across all regimes to populate the matrix.

Study area

The study region extends from the western shores of the Kutch peninsula to Kanyakumari at
the southern tip of India - a stretch of approximately 3400 km of coastline (Refer Figure.2A)
The region represents a single eco-region based on its faunal community composition[30]. While
fish composition and bathymetry are relatively consistent, the governance of the coast is
distributed across five states (Gujarat, Maharashtra, Goa, Karnataka and Kerala). Each state
manages its own fisheries with policy instruments in the form of Marine Fisheries Regulation
Acts (MFRAs), as well as schemes and subsidies to manage the coastal waters under its
jurisdiction. The uniformity of the habitat across the coastline together with its disjunct
governance regimes makes the west coast an ideal case study to understand how differences
in fisheries management can mediate the vulnerabilities of individual fish populations. As
potential outliers, we also sampled individuals from a single location on India's east coast, in
Balasore, Odisha, approximately 1700 km away from southernmost sampling locations on the
west coast. These east coast samples are expected to act as a separate population in our
analysis.

Choice of species

We identified nine fish species commonly caught in trawls on the west coast of India. The
choice of species was dependent on their ubiquity, the frequency with which they occur in
trawl catches across the west coast, and ease of identification in the field. They were also
chosen to represent different trophic groups, habitats, taxonomic groups and commercial
value.

The selected fish species were silver pomfret (*Pampus argenteus*), ribbonfish (*Trichiurus*
lepturus), Japanese threadfin bream (*Nemipterus japonicus*), spiny cheek grouper
(*Epinephelus diacanthus*), spade nose shark (*Scoliodon laticaudus)*, wolf herring
(*Chirocentrus nudus*), Indian oil sardine (*Sardinella longiceps)*, ladyfish (*Sillago sihama*), and
Indian mackerel (*Rastrelliger kanagurta*),

Table 2 – Target species with their life history characters and comparative commercial
values

	Common name	Scientific name	Habitat	Food	Value
	Silver Pomfret	*Pampus argenteus*	Muddy bottoms	Zooplankton	Very high
	Ribbonfish	*Trichiurus lepturus*	Muddy bottoms	Fish, squid etc	High
	Japanese threadfin bream	*Nemipterus japonicus*	Mud and Sand bottoms	Small fish, Crustaceans etc	Low
	Spiny cheek grouper	*Epinephelus diacanthus*	Muddy substrate	Fish	Low
	Spade nose shark	*Scoliodon laticaudus*	Rocky substrate	Small fish, shrimp	Moderate
	wolf herring	*Chirocentrus nudus*	Inshore waters	Small fish, crustaceans etc	High
	Indian oil sardines	*Sardinella longiceps*	Coastal pelagic	Phytoplankton, crustaceans etc	Low
	ladyfish	*Sillago sihama*	Beach, sandbank, estuaries	Polychaete, shrimps,	low
	Indian Mackerel	*Rastrelliger kanagurta*	Coastal pelagic	Small Zooplankton	Moderate

Identifying the geographical extent of individual fish populations.

i. Data collection

We collected tissues samples using fin-clips taken from the fish directly at the landing site. Sample collection along the west coast of India was conducted in April 2018, February-March 2020 and November 2020. The samples from one east coast site (Balasore, Odisha) were collected in January 2023

Fin clips of ~ 10-18 individuals were collected from each species at each sampling location with the help of sterilised instruments and stored in 100% ethanol. We limited our sampling to fish landed from small boats with low engine capacity to ensure (as much as possible) that the fish sampled were sourced from local waters. We sampled only adult fish from each location.

ii. Data generation

We used genome-wide SNP (Single Nucleotide Polymorphism) markers to identify genetically distinct populations and their geographical extent for all nine chosen species. We used the Double Digest Restriction Associated DNA (ddRAD) approach to generate genomic data[31]. The sample collection protocol along with detailed protocols of DNA extraction, library preparation is given in the supplementary material, section 3.

iii. Data analysis

We used two approaches for SNP calling: with reference genomes and deNOVO SNP calling. For silver pomfret (*Pampus argenteus*) and Indian mackerel (*Rastrelliger kanagurta,*) we used available reference genomes from the NCBI database for SNP calling (*Rastrelliger kanagurta - GCA_024605255.1, Pampus argenteus* - GCA_000697985.1). For spiny cheek grouper (*Epinephelus diacanthus*) and spade nose shark (*Scoliodon laticaudus)* we used the reference genomes of the nearest taxonomically related species. We used the *Epinephelus coioides* (GCA_900536245.1) genome for *Epinephelus diacanthus* and *Scyliorhinus canicular* (GCA_902713615.1) genome for *Scoliodon laticaudus* from the NCBI database. There were no available reference genomes that we could use for any of the other species, so we used a de novo variant calling pipeline by STACKS. The detailed bioinformatic protocol (quality control of reads and SNP discovery) are provided in supplementary materials, section 3 and section 4.

After SNP discovery, we used three indices to identify evidence of genetic similarity and dissimilarity between samples and ultimately identify genetically distinct populations for all nine species. These indices were:

1. Visualisation based on a principal component analysis (PCA) using PLINK[32].
2. Visualisation based on admixture probabilities using the programme ADMIXURE[33]. We ran admixture for K ranging from 1 to n, where n = the number of sites in which we have information of that particular species, for 10 replicates each, the best K was determined by estimating the cross-validation error. The results were analysed and plotted using a web-based server, CLUMPAK[34].
3. A pairwise mean F_{ST} calculation conducted using STACKS[35].

Calculating "Productivity - Susceptibility (PSA) vulnerability axis

We used the Productivity Susceptibility Analysis conducted by[16] Mohammad et al. (2021) for 133 species found in Indian waters as the base vulnerability for our target species. They formulated an index of Resilience and Vulnerability (IRV) based on the Growth coefficient (K), Longevity: Asymptotic length (L∞), Reproductive load, Number of spawning months (NSM), Fecundity and Mean trophic level (MTL) as vulnerability attributes and species geographic distribution, body length depth ratio (BLD), exploitation ratio, recruitment, susceptibility to gear and price as resilience attributes. This study is the most rigorous assessment of species susceptibility of marine fish from the region.

We used IRV index values along the species-specific vulnerability axis (the vertical axis of our matrix) for all nine species (Refer Figure 3A). Mohammad et al. did not assess the IRV index of *Silago sihama*, so we calculated its IRV index based on the formula given by Mohammad et al. (See supplementary material, section 1).

Identifying disparity between species biology and management attributes

Across the west coast of India, we surveyed fisheries regulations and policy instruments relevant to our study species. In a related study, we reviewed fisheries policies across the west coast of India and identified a total of nine different regulations as the key management attributes of the regimes prevalent in the five states of the Indian west coast[26]. They were 1) partial or complete fishing ban during monsoon months (June to August) 2) maximum allowable engine capacity of a fishing vessel 3) minimum allowable mesh size for cod end of trawl operation 4) minimum allowable mesh size for gill nets 5) minimum allowable mesh size for purse nets / ring seine nets 6) minimum allowable legal size for a species 7) ban on bull / paired trawling 8) ban on fishing using lights 9) restriction on mechanised fishing in shallow waters. These nine regulations were used to assess state-wide disparity with fish biology. Although we would ideally have included catch and effort details with respect to MSY as independent attributes, no adequate data exists for these nine species at the individual population level. See discussion for more details. The details and justifications for the percent disparity calculations are provided in detail in Supplementary material, section 2.

Populating the Biology - Transboundary Management (BTM) vulnerability matrix

We used information of the geographical extent of individual populations from genomic data, the IRV index and percent disparity (as described in above sections), to map the population-level vulnerability of the 9 target species using the Biology - Transboundary Management (BTM) vulnerability matrix.

Results

2. Establishing geographical boundaries of target fish populations

Single Nucleotide Polymorphism (SNP) analysis indicates that the degree of gene flow between individuals across the west coast of India varies considerably with species, as is evident from the Principal Component Analysis (PCA) as shown in Fig. 2. We substantiated our inference of population structure using pairwise F_{ST} and Admixture analysis[33] in addition to PCA. For more details about these analyses refer to the section 3 of the supplementary material.

Rastrelliger kanagurta and *Chirocentrus nudus* show a completely panmictic population across the west and east coast (Figure 2B & 2C). *Epinephelus diacanthus* shows evidence of a single population across the entire west coast (Figure 2D). *Trichiurus lepturus* and *Nemipterus japonicus* also show a single population across the entire west coast however they have clear divergences from conspecific populations from the east coast (Figure 2E & 2F).

The remaining four species show multiple genetically distinct population along the west coast of India. Genetic data indicate two potential *Pampus argenteus* populations along the west coast of India, one between northern Gujarat to central Maharashtra, and another from southern south Maharashtra to Kerala (Figure 2G). *Sardinella longiceps* also shows evidence of two populations, with a Gujarat population that separates clearly from the rest of the west coast population (Figure 2H). *Sillago sihama* shows evidence of an isolated population near the town of Kumta which is separate from the coastwide Maharashtra - Karnataka population (Figure 2J). Finally, genetic data from *Scoliodon laticaudus* shows evidence of three distinct west coast populations: a northern Gujarat population, a middle Maharashtra-Goa-Karnataka population, and a southern Kerala population (Figure 2I).

334   Figure 2. Principal Component plots for nine chosen species across the west coast of India

3. Formulating the vertical "Productivity - Susceptibility (PSA) vulnerability" axis

IRV values can range between 0 (most vulnerable) to 1 (least vulnerable)[16]. Among the 133
species Mohammad et al. analysed, *Solenocera crassicornis* (coastal mud prawn) had the
highest IRV value of 0.512 where *Carcharhinus sorrah* (spot-tail shark) had lowest IRV value
of 0.022. the nine target species we chose, *Nemipterus japonicus* had the highest IRV index
value of 0.31 and thus the lowest vulnerability to overexploitation on the vertical axis whereas
Epinephelus diacanthus had the lowest IRV index value of 0.195 making it most vulnerable
on the vertical axis. We calculated an IRV index of 0.237 for *Sillago sihama* (See
supplementary material, section 1). Our target species occupied a narrower portion of the
range of species analysed by Mohammad et al. 2021. Figure 3A illustrates the spread of target
species along the IRV axis.

Figure 3. Biology – Transboundary Management (BTM) vulnerability matrix for nine species along the west coast of India

4.   Identifying percent disparity between species biology and regulations

We identified the percent disparity between biology and regulation for each individual
population - state (management regime) combination based on the rationale described in the
methods section. Results of the disparity analysis for the Indian oil sardine are shown in Figure
4. The Gujarat population (which is housed in a single management regime) has a high
disparity (66%), while the peninsular India population, despite being shared across four states,
has a low disparity across the entire coast. Further results of all species are provided in the
supplementary materials, section 2.

Figure 4. Disparity analysis for Indian oil sardine, with populations housed in single as well
as multiple management regimes.

364
365
366

5.   Mapping the vulnerability of west coast populations using the BTM matrix

We used species specific IRV values from Mohammad et al. and state wise percent disparity
for each population identified by genetic analysis, to construct a Biology - Transboundary
Management (BTM) vulnerability matrix for the west coast[16] (Figure 3B). There was a
considerable spread on both axes of the matrix. Critically, species with multiple populations
along the coast could have vastly different vulnerabilities based on the management disparity
axis. This was most clear for the spade nose shark, where the transboundary southern
population had a much higher vulnerability than the Kerala and Gujarat populations. For
populations of pomfret and sardines the differences were not as stark, although differences in
management practices between states still clearly influenced the vulnerability of each
population. While panmictic species (with a single population across the coast), occupied a
wide range of PSA vulnerabilities, the inherent transboundary nature of these species strongly
mediated their inherent vulnerability.

Discussion

Achieving sustainability in multi species, poorly monitored and data-poor tropical coastal
fisheries is often considered a wicked problem with no easy solutions[36],[6]. Although recent
approaches like the Productivity Susceptibility Analysis (PSA) comprehensively incorporate
complex species biology, they do not account for management complexities arising from the
transboundary nature of most coastal fish populations. The BTM matrix we propose combines
genomic techniques and policy analysis for a more realistic assessment of fish stock
vulnerability. Our case study demonstrates that many target coastal species across the Indian
coast are transboundary in nature, spanning multiple harvest regimes with differing
approaches to biological sustainability. This has critical implications for tropical fisheries
management. On the one hand, within a species, each population could face very different
vulnerability threats. On the other, different fisheries harvesting the same population can
potentially influence vulnerability of that population to overexploitation. These complexities are
captured elegantly in the BTM matrix. It provides a framework for sustainable management of
transboundary stocks, allowing fisheries managers to more transparently assess the
vulnerability of populations, evaluate fisheries options and negotiate pragmatic transboundary
arrangements based on known the boundaries of susceptible species.

A novel approach to predict vulnerability to overharvest

Our analysis demonstrates the vital importance of evaluating fish vulnerability for every
population individually, something that most assessments do not allow. A case in point is the
silver pomfret. With an vulnerability score of 0.24 (based on the IRV index, see methods), it is
a species of intermediate vulnerability. However, our population genetic analysis shows that
there are two populations of silver pomfret along the west coast of India, a northern population
that extends from Gujarat to the central coast of Maharashtra, and a southern population that
extends further south to Kerala. Although both of these populations share the same IRV index,
they vary in their vulnerability along the transboundary management axis. Specifically, the
southern population is governed by more stringent harvest rules for pomfret than the northern
population. Karnataka and Kerala both have regulations on minimum mesh size for gill nets,
and there are pomfret-specific minimum legal-size regulations in place in Goa, Karnataka and
Kerala. These regulations are absent in the northern states. As a result, the southern

population experiences considerably lower disparity between pomfret biology and fisheries
regulations across its range than the northern population, making it much less vulnerable to
overexploitation. This example demonstrates how by shifting the assessment focus to spatially
explicit populations, our vulnerability matrix considerably enhances the value of the original
IRV assessment. Further, by identifying the specific management attributes responsible for
differences in vulnerability, the BTM vulnerability matrix approach is highly transparent,
providing actionable recommendations to policy makers on how to reduce species
vulnerability with changes in policy and transboundary cooperation.

Another aspect of this vulnerability matrix is that it is dynamic and sensitive to periodic
revisions in fishery policy across management regimes. The Indian oil sardine demonstrates
this well. The west coast has a largely panmictic population except for Gujarat, which has a
distinct population. Fewer than 10 years ago no west coast Indian states had regulations on
fishing methods like bull trawling, light fishing etc. In addition, there were no mesh size
regulations for purse nets or gill nets and none of the states had defined minimum legal sizes
for the Indian oil sardine. Based on our assessment, until 2016, the Indian oil sardine was in
the highest category of transboundary management vulnerability. The subsequent adoption of
these regulations by all west coast states has significantly reduced its vulnerability to
overharvest. The advantage of this analysis is that it can be used to keep track of the direction
of fisheries policies across boundaries, and can update population vulnerabilities based on
policy decisions (Figure 5).

444   Figure 5. Shift in vulnerability of the peninsular population of Indian oil sardine along the
445   management axis as a direct result of changing management practices

Legend: Matrix legend is similar to figure 3C. The peninsula population of Indian oil
sardine saw a recent improvement in its vulnerability as a direct result of actions by states
within its population range. While before 2016, there was high management disparity
(represented by greyed out boxes) in all states in its range before 2016, there has been a
gradual improvement in management practices, with Karnataka (KR) and Kerala (KL)
reducing its disparity in 2018, and Maharashtra (MH) and Goa (GO) following suit in 2023,
resulting in an overall reduction in the transboundary vulnerability of this population.

We assume that IRV values remain relatively constant for a species. However, it is possible
that critical life-history traits could vary between genetic stocks. For instance, Patrik et al.
show that the PSA vulnerability of different populations of the same species could differ
considerably[18]. In evaluating the disparity between biology and management regime
information for our nine target species, we identified considerable differences in basic life
history characters like length at first maturity (l50) between Pomfret [37,38] and spade nose
shark[39,40]   caught at different locations. These traits could potentially also influence their
catchability in gear operating along the coast. Such differences could potentially lead to
varying inherent population-level susceptibilities and slightly different IRV values for each
species. Unfortunately, comprehensive life-history data of this granularity is unavailable for
most tropical species. Before it is possible to biologically characterise each individual
population more accurately, a more fundamental problem is identifying the population
boundaries. Additional life-history studies could help update IRV values for a more nuanced
vulnerability assessment.

While IRV values may vary slightly as we learn more about each population, they represent a
largely unchanging inherent property of each taxon. However, a central feature of our matrix
is that real-world vulnerability is as much a characteristic of the management jurisdiction in
question as it is of the biological character of the species. This explicitly recognises the fact
that fisheries regimes vary widely between locations, and that transboundary agreements (or
the lack of them), can strongly mediate the inherent vulnerability of fish stocks. It is critical to
keep this in mind when applying the matrix to a specific fishery. For instance, we classified the
Gujarat population of the Indian oil sardine as belonging to a single jurisdiction, reducing its
overall vulnerability within the matrix. Broadening the geographical scale of our analysis may
well have shown that this population is contiguous with sardines from Pakistan and beyond,
making it potentially subject to trans-national management differences that we did not evaluate
here. This could equally apply to the spade nose shark of Gujarat and Kerala or the isolated
ladyfish / silver sillago population from Kumta. A fuller analysis would require establishing
genetic boundaries for the entire range of the species in question before determining its overall
vulnerability. This will have important implications for the kinds of transboundary arrangements
required to reduce the overall susceptibility of the population – between coastal nations or
between sub-national jurisdictions. The fact that management differences can so strongly
mediate inherent vulnerability indicates that assessments of this kind are most meaningful
when designed for every region, making them jurisdictionally and biologically appropriate
across the range of the species.

Transboundary management of fish populations in tropical waters

How could this vulnerability matrix (or exercises such as these) inform sustainable tropical
fisheries? Vulnerability assessments of this sort go a long way in sharply defining the
management needs for each population across its natural range and help define the scale at
which fisheries management would be most biologically meaningful. It offers a clear set of
directions to better align species biology with management, setting the stage for cross-
boundary cooperation on specific aspects of policy and practice. How tropical regions
negotiate these arrangements however continues to remain tricky.

Countries in Western Europe have had a long history of hard-won transboundary fishing
arrangements[41], driven by decades of conflict over shared resources. There is also
significantly better information on geographically identifiable, genetically distinct populations
of fish that form the basis of these arrangements[28]. Tropical fisheries clearly have a
considerable distance to travel before they arrive at this state of knowledge or policy
engagement. For one, there exists no precedent of international transboundary arrangements
for coastal tropical fisheries although there are non-binding international arrangements for
pelagic species, like tuna[42].

More importantly, most tropical fisheries are organised and managed in very different ways
from their temperate counterparts. In many countries coastal fisheries are managed in a highly
decentralised manner where regions and provinces have considerable autonomy over local
fisheries management[22–25]. In addition, the allied problems of equity and social justice
associated with coastal fisheries are seeing an increasing movement towards even further
localisation of fisheries, with approaches like TURFs (Territorial User Rights for Fishing)
gaining popularity[43]. These are usually smaller than regional government fisheries
management units. Most TURFs in Chile, Mexico and Japan are smaller than the dispersal

ability of fish larvae and adults[44]. While these efforts are a welcome direction to secure the
livelihood of marginalised coastal communities, they are also subject to boundary paradoxes,
where increasing fisheries boundaries come at the cost of increasingly complex transboundary
negotiations[45]. Our analysis raises serious questions of how to ensure the success of these
pockets of community managed fishing in a wider ocean of a rapidly expanding fishery sector.
Without adequate coordination across biological bounds, these efforts may also result in
inadvertent overharvest, and it is important to create forums of dialogue between these
community managed and larger regional fisheries, keeping in mind the genetic boundaries of
fish stocks.

Conclusion

For multi-specific tropical fisheries, transitioning to population-specific management of fish
stocks may seem like a distant horizon. Eventually though, if coastal tropical nations are
serious about stemming the collapse of coastal fish stocks, population-wise management is
likely to be critical, at least for the most vulnerable of coastal species. Coastal transboundary
arrangements of the kind that are already in place for open-water migratory species like tuna[42]
offer a possible template that can be built on. Our results demonstrate the need for such
agreements for nearshore species of fish and for individual provinces and community
managed areas within a single nation state, which was largely considered less important in
tropical context. For this, the BTM vulnerability matrix can be a useful prioritisation tool.
Because the management axis allows for dynamic mobility based on changes in policy, it can
be used to strategically evaluate which set of policy directions could reduce the susceptibility
of the widest number of fish populations across a shared coastline. As our analyses shows,
the inherent vulnerability of fish can be strongly mediated by what managers do at local levels,
and how closely they coordinate their actions with neighbours that share the same genetic
resource pool. This offers the principal challenge but also the best hope for rational
management of dwindling fish stocks.

Bibliography

1.   Huxley, F. R. S. Inaugural meeting of the fishery congress. Address. *The Great International Fisheries Exhibition,* 1–22 (1883).

2.   FAO. SDG INDICATOR 14.4.1 - Proportion of fish stocks within biologically sustainable levels Target. in *Tracking progress on food and agriculture-related SDG indicators 2023* 107–110 (2023). doi:https://doi.org/10.4060/cc7088en-fig55

3.   Costello, C. *et al.* Global fishery prospects under contrasting management regimes. *PNAS* **113**, 5125–5129 (2016).

4.   Dulvy, N. K. *et al.* Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Curr. Biol.* **31**, 4773-4787.e8 (2021).

5.   Teh, L. C. L. & Sumaila, U. R. Contribution of marine fisheries to worldwide employment. *Fish Fish.* **14**, 77–88 (2011).

6.   Mora, C. *et al.* Management Effectiveness of the World ' s Marine Fisheries. *PloS Biol.* **7**, e10001311 (2009).

7.   Hilborn, R., Oscar, R., Anderson, C. M., Baum, J. K. & Branch, T. A. Effective fisheries management instrumental in improving fish stock status. *PNAS* **117**, 2218–2224 (2020).

8.   Legovic, T., Klanjšček, J. & Geček, S. Maximum sustainable yield and species extinction in ecosystems. *Ecol. Modell.* **221**, 1569–1574 (2010).

9.   Ramesh, M. & Namboothri, N. Maximum sustainable yield: A myth and its manifold effects. *Econ. Polit. Wkly.* **53**, 58–63 (2018).

10. Finley, C. & Oreskes, N. Maximum sustained yield : a policy disguised as science. *ICES J. Mar. Sci.* **70**, 245–250 (2013).

11. Hoshino, E., Pascoe, S., Hutton, T. & Kompas, T. Estimating maximum economic yield in multispecies fisheries : a review. *Rev. Fish Biol. Fish.* **28**, 261–276 (2018).

12. Rosenberg, A. A. *et al. Use of Productivity-Susceptibility Analysis (PSA) in Setting Annual Catch Limits for U. S. Fisheries : A Workshop Report.* (2009).

13. Patrick, W. S. *et al. Use of Productivity and Susceptibility Indices to Determine Stock Vulnerability , with Example Applications to Six U . S . Fisheries.* (2009).

14. Duffy, L. M. & Griffiths, S. P. Assessing attribute redundancy in the application of productivity- susceptibility analysis to data-limited fisherie. *Aquat. Living Resour.* **32**, 1–11 (2019).

15. Faruque, H. & Matsuda, H. Assessing the vulnerability of bycatch species from Hilsa gillnet fishing using productivity susceptibility analysis: Insights from Bangladesh. *Fish. Res.* **234**, 105808 (2021).

16. Mohamed, K. S. *et al. Application of biological and fisheries attributes to assess the vulnerability and resilience of tropical marine fish species. PLoS ONE* **16**, (2021).

17. Leach, A., Kell, L. T. & Mumford, J. Productivity-Susceptibility Analysis tool. *ICCAT* **80**, 98–116 (2023).

18. Patrick, W. S. *et al.* Using productivity and susceptibility indices to assess the vulnerability of United States fish stocks to overfishing. *Fish. Bull.* **108**, 305–322 (2010).

19. Laikre, L., Palm, S. & Ryman, N. Genetic Population Structure of Fishes : Implications for Coastal Zone Management. *AMBIO A J. Hum. Environ.* **34**, 111–119 (2005).

20. Waples, R. S., Punt, A. E. & Cope, J. M. Integrating genetic data into management of marine resources: how can we do it better? *Fish Fish.* **9**, 423–449 (2008).

21. Palsbøll, P. J., Bérubé, M. & Allendorf, F. W. Identification of management units using population genetic data. *TRENDS Ecol. Evol.* **22**, 11–16 (2006).

22. Arceo, H. O., Cazalet, B., Alin, P. M., Mangialajo, L. & Francour, P. Moving beyond a top-down fisheries management approach in the northwestern Mediterranean: Some lessons from the Philippines. *Mar. Policy* **39**, 29–42 (2013).

23. Jaya, I. *et al.* 'Are the working principles of fisheries management at work in Indonesia?' *Mar. Policy* **140**, 105047 (2022).

24. Monteferri, B. & Muller, M. R. Marine, coastal and fisheries issues in Chile, Mexico

and Peru: An initial institutional and policy review. in *Marine and Fisheries Policies in*
Latin America : A Comparison of Selected Countries (eds. Muller, M. R., Oyanedel, R.
& Monteferri, B.) 6–14 (Routledge. Taylor and Francis Group, 2020).

25. Nakamura, J. & Hazin, F. Assessing the Brazilian federal fisheries law and policy in
light of the Voluntary Guidelines for Securing Sustainable Small-scale fisheries. *Mar.*
Policy **113**, 103798 (2020).

26. Gangal, M., Suri, V. & Arthur, R. How well does Indian fisheries policy engage with
fisheries biology? Exploring the science-policy interface of coastal capture fisheries
along the west coast of India. *Mar. Policy* **156**, 105796 (2023).

27. Berger, A. M. *et al.* Incoherent dimensionality in fisheries management:
consequences of misaligned stock assessment and population boundaries. *ICES J.*
Mar. Sci. **78**, 155–171 (2021).

28. Reiss, H., Hoarau, G., Dickey-Collas, M. & Wolff, W. J. Genetic population structure of
marine fish: Mismatch between biological and fisheries management units. *Fish Fish.*
10, 361–395 (2009).

29. Østhagen, A., Spijkers, J. & Totland, O. A. Collapse of cooperation ? The North-
Atlantic mackerel dispute and lessons for international cooperation on transboundary
fish stocks. *Marit. Stud.* **19**, 155–165 (2020).

30. Spalding, M. D. *et al.* Marine ecoregions of the world: A bioregionalization of coastal
and shelf areas. *Bioscience* **57**, 573–583 (2007).

31. Tyagi, A., Yadav, N., Pandit, A. & Ramakrishnan, U. On the road to losing
connectivity: Fecal samples provide genome-wide insights into anthropogenic impacts
on two large herbivore species in central India. *Mol. Ecol.* 1–17 (2024).
doi:10.1111/mec.17461

32. Purcell, S. *et al.* PLINK : A Tool Set for Whole-Genome Association and Population-
Based Linkage Analyses. *Am. J. Hum. Genet. Vol.* **81**, 559–575 (2007).

33. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry
in unrelated individuals. *Genome Res.* 1655–1664 (2009).
doi:10.1101/gr.094052.109.vidual

34. Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A. & Mayrose, I.
CLUMPAK: a program for identifying clustering modes and packaging population
structure inferences across K. *Mol. Ecol. Resour.* **15**, 1179–1191 (2015).

35. Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A. & Cresko, W. A. Stacks: An
analysis tool set for population genomics. *Mol. Ecol.* **22**, 3124–3140 (2013).

36. Torre-castro, M. De. Governance for Sustainability : Insights from Marine Resource
Use in a Tropical Setting in the Western Indian Ocean. *Coast. Manag.* **40**, 612–633
(2012).

37. Ghosh, S. *et al.* Fishery and stock estimates of the silver pomfret , Pampus argenteus
(Euphrasen), landed by gill netters at Veraval. *Indian J. Fish.* **56**, 177–182 (2009).

38. Gladston, Y., Devi, M. S. & Xavier, K. A. M. Design and gillnet selectivity of Pampus
argenteus along the Satpati coast, Maharashtra, India. *Reg. Stud. Mar. Sci.* (2016).
doi:10.1016/j.rsma.2016.12.006

39. Barnes, A., Sutaria, D., Harry, A. V. & Jabado, R. W. Demographics and length and
weight relationships of commercially important sharks along the north-western coast
of India. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **28**, 1374–1383 (2018).

40. Sen, S. *et al.* Population dynamics and stock assessment of spadenose shark
Scoliodon laticaudus Muller and Henle 1839 along Gujarat coast of India. *Indian J.*
Geo Mar. Sci. **48**, 423–433 (2019).

41. Lado, E. P. Why the common fisheries policy is important. in *The Common Fisheries*
Policy: The Quest for Sustainability 1–17 (John Wiley & Sons, Ltd., 2016).
doi:10.1002/9781119085676

42. Cullis-Suzuki, S. & Pauly, D. Failing the high seas: A global evaluation of regional
fisheries management organizations. *Mar. Policy* **34**, 1036–1042 (2010).

43. Arnason, R. & Runolfsson, B. T. Delineating the impacts of strong user rights in
fisheries. *Mar. Policy* **155**, 105666 (2023).

44.   Aceves-Buenoa, E., Cornejo-Donoso, J., Miller, S. J. & Gaines, S. D. Are Territorial
Use Rights in Fisheries (TURFs) sufficiently large? *Mar. Policy* **78**, 189–195 (2017).
45.   Song, A. M., Scholtens, J., Stephen, J., Bavinck, M. & Chuenpagdee, R.
Transboundary research in fisheries. *Mar. Policy* **76**, 8–18 (2017).

655
656
657

A

Sampling locations
Map and Legend

- Porbandar
- Surat
- Mumbai
- Harne
- Vengurle
- Goa
- Kumta
- Mangalore
- Kozhikode
- Alappuzha
- Odisha

B

Indian Mackerel
(*Rastrelliger kanagurta*)

Ref. Genome –
Rastrelliger kanagurta
No. of SNPs – 13702

Important filters
Missing data – Nil
Minor Allele Count – 3

C

Wolf herring
(*Chirocentrus nudus*)

Ref. Genome –
None (DeNOVO)
No. of SNPs – 6747

Important filters
Missing data – 10%
Minor Allele Count – 3

D

Spinycheek grouper
(*Epinephelus diacanthus*)

Ref. Genome –
Epinephelus coioides
No. of SNPs – 5252

Important filters
Missing data – Nil
Minor Allele Count – 3

E

Ribbonfish
(*Trichiurus lepturus*)

Ref. Genome –
None (DeNOVO)
No. of SNPs – 5076

Important filters
Missing data – 10%
Minor Allele Count – 3

F

Japanese threadfin bream
(*Nemipterus japonicus*)

Ref. Genome –
None (DeNOVO)
No. of SNPs – 9824

Important filters
Missing data – 10%
Minor Allele Count – 3

G

Silver pomfret
(*Pampus argenteus*)

Ref. Genome –
Pampus argenteus
No. of SNPs – 4945

Important filters
Missing data – Nil
Minor Allele Count – 3

H

Indian oil sardine
(*Sardinella longiceps*)

Ref. Genome –
None (DeNOVO)
No. of SNPs – 4470

Important filters
Missing data – 25%
Minor Allele Count – 3

I

Spadenose Shark
(*Scoliodon laticaudus*)

Ref. Genome –
Scyliorhinus canicular
No. of SNPs – 1166

Important filters
Missing data – 25%
Minor Allele Count – 3

J

Silver Silago
(*Sillago sihama*)

Ref. Genome –
None (DeNOVO)
No. of SNPs – 2755

Important filters
Missing data – 10%
Minor Allele Count – 3

Sardinella longiceps (Indian oil sardine)

Population housed in single management regime – Gujarat population	Population housed in multiple management regimes – Peninsular India population			

 |

 |

 |

 | |
| High Disparity | Low disparity | Low disparity | Low disparity | Low disparity |
| | High disparity in majority of management regimes – 0/4 | | | |

Legend

 Congruence between species biology and regulation

 Disparity between species biology and regulation

 Monsoon fishing ban

 Maximum engine capacity

 Mesh size regulation for cod-end net of trawler

 Mesh size regulation for purse nets

 Mesh size regulation for gillnets

 Minimum Legal size

 Ban on Bull trawling

 Ban on light fishing

 Restriction on fishing in shallow waters

 High disparity with percent disparity value written

 Low disparity with percent disparity value written

Management Disparity Axis: Illustrates the gap between
on ground species management and biologically sustainable practices,
highlighting the divergence between operational strategies and ecological sustainability.