CUTS RNA Biosensor for the Real-Time Detection of TDP-43 Loss-of-Function

Longxin Xie*[1,2,3], Jessica Merjane*[1,3], Cristian A Bergmann*[1,3], Jiazhen Xu[1,3,4,]: Bryan Hurtle[1,3,5];
Christopher J Donnelly[1,3-7]

[1]Department of Neurobiology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

[2]School of Medicine, Tsinghua University, China

[3]LiveLikeLou Center for ALS Research, University of Pittsburgh School of Medicine, Pittsburgh,
PA, USA

[4]Interdisciplinary Biomedical Graduate Program Cellular and Molecular Pathology, University of
Pittsburgh, Pittsburgh, PA, USA

[5]Center for Neuroscience at the University of Pittsburgh, Pittsburgh, PA, USA

[6]Pittsburgh Institute for Neurodegeneration, University of Pittsburgh, Pittsburgh, PA, USA

[7]Center for Protein Conformational Diseases, University of Pittsburgh, Pittsburgh, PA, USA

* Authors share equal contributions

Highlights
• CUTS is a cryptic exon RNA biosensor enabling real-time detection of TDP-43 loss of splicing
function.
• CUTS exhibits a significant linear relationship with TDP-43 loss.
• CUTS can deliver an autoregulated gene payload in response to TDP-43 loss-of-function.
• TDP-43 homotypic phase transitions induce loss of splicing function via CUTS detection.

Abstract

Given the mounting evidence implicating TDP-43 dysfunction in several neurodegenerative
diseases, there is a pressing need to establish accessible tools to sense and quantify TDP-43
loss-of-function (LOF). These tools are crucial for assessing potential disease contributors and
exploring therapeutic candidates in TDP-43 proteinopathies. Here, we develop a sensitive and
accurate real-time sensor for TDP-43 LOF: the CUTS (CFTR UNC13A TDP-43 Loss-of-Function)
system. This system combines previously reported cryptic exons regulated by TDP-43 with a
reporter, enabling the tracking of TDP-43 LOF through live microscopy and RNA/protein-based

assays. We demonstrate CUTS' effectiveness in detecting LOF caused by TDP-43
mislocalization and RNA binding dysfunction, and pathological aggregation. Our results highlight
the sensitivity and accuracy of the CUTS system in detecting and quantifying TDP-43 LOF,
opening avenues to explore unknown TDP-43 interactions that regulate its function. In addition,
by replacing the fluorescent tag in the CUTS system with the coding sequence for TDP-43, we
show significant recovery of its function under TDP-43 LOF conditions, highlighting CUTS'
potential for self-regulating gene therapy applications. In summary, CUTS represents a versatile
platform for evaluating TDP-43 LOF in real-time and advancing gene-replacement therapies in
neurodegenerative diseases associated with TDP-43 dysfunction.

<u>Introduction</u>

Amyotrophic lateral sclerosis (ALS) is a progressive and fatal neurodegenerative disease
(NND) characterized by a persistent degeneration of the neurons of the spinal cord and motor
cortex (Feldman et al., 2022). The dysregulation of the RNA binding protein (RBP), TAR
DNA-binding protein 43 (TDP-43) is a hallmark pathobiology observed in ~97% of all ALS patients
that (Arai et al., 2006; Neumann et al., 2006) ~45% of Frontotemporal lobar degeneration (FTLD)
patients (S.- C. Ling et al., 2013; Neumann et al., 2006), and 40%-60% of Limbic Associated
TDP-43 Encephalopathy (LATE) patients (McKee et al., 2010; Nelson et al., 2019; Tremblay et
al., 2011). Under physiological conditions, TDP-43 orchestrates many cellular processes critical
for neuronal health and homeostasis, including regulating RNA metabolism, splicing, and stress
response pathways. However, in disease, TDP-43 is depleted from the nucleus and mislocalizes
to the cytoplasmic compartment, losing the ability to perform its canonical functions and
transitioning into insoluble aggregates (Hurtle et al., 2023).

Recent efforts increasingly focus on the implications of TDP-43's loss of splicing function
and its consequence on disease onset and progression (Brown et al., 2022; Buratti et al., 2001;
Klim et al., 2019; J. P. Ling et al., 2015; Ma et al., 2022; Melamed et al., 2019).   Physiologically,
TDP-43 selectively binds to specific UG-rich sequences within pre-mRNA transcripts, providing
precise control over the alternative splicing of a subset of conserved targets, thereby modulating
their gene expression and cellular function (Kuo et al., 2009; Lukavsky et al., 2013). This
regulatory function is key in repressing the incorporation of TDP-43-mediated aberrant 'cryptic
exons' (CE), non-conserved intronic regions whose inclusion has been linked to cellular toxicity
and pathological consequences (Brown et al., 2022; Klim et al., 2019; J. P. Ling et al., 2015; Ma
et al., 2022; Mehta et al., 2023; Melamed et al., 2019; Polymenidou et al., 2011).   Noteworthy

examples of transcripts susceptible to aberrant CEs in the absence of functional TDP-43 include

STMN2 (Klim et al., 2019; Melamed et al., 2019) and *UNC13A* (Brown et al., 2022; Ma et al.,

2022), both pivotal in maintaining the integrity and physiological function of neurons, such as axon

regeneration and motor neuron firing (Shin et al., 2012, 2014; Varoqueaux et al., 2002; Willemse

et al., 2023). Therefore, their aberrant splicing patterns and loss-of-function (LOF) in disease

highlights the critical role of TDP-43 in preserving neuronal health and function through the

regulation of key neuronal health.

A well-established and currently available method to accurately monitor and quantify

TDP-43's splicing function has frequently relied on employing cryptic exon 9 inclusion in the

human cystic fibrosis transmembrane conductance regulator (CFTR) transgene (Buratti et al.

2001; Ayala et al. 2006). However, this approach can only be used at the RNA level, limiting its

functionality as a reporting system for high-throughput screens or *in vivo* applications.

Furthermore, endogenous cryptic exons exhibit variable responses to TDP-43 loss, may be cell

type-specific, and are still being defined. To address this, we designed a screening tool, the

'CFTR-UNC13A TDP-43 loss-of-function Sensor' (CUTS) system, engineered to detect TDP-43

LOF and output a proportional and quantifiable GFP signal or other reporters. By combining

previously described TDP-43 binding targets, we significantly improved the sensitivity

measurable in real-time by standard assays. Physiological TDP-43 promotes the splicing of

CUTS CE, resulting in a frameshift and early stop codon that prevents GFP expression. TDP-43

LOF activates the CUTS biosensor by maintaining the CE inclusion and allowing GFP to express.

Here, we highlight the CUTS system sensitivity in response to TDP-43 loss. The CUTS design

allows for GFP expression with direct proportionality to TDP-43 LOF, enabling the detection of

changes in TDP-43 function, even when alterations in TDP-43 itself cannot be directly measured

by traditional means, including qPCR, western blot, and immunofluorescent imaging.

In this study, we highlight the utility and sensitivity of the CUTS biosensor. We also show

that aberrant TDP-43 phase transitions or mislocalization disrupt TDP-43 splicing function via

expression of well-established RNA binding deficient and cytoplasmic TDP-43 constructs. Lastly,

we replaced the GFP cassette with the wildtype *TARDBP* coding sequence (CUTS-TDP43) to

highlight the utility of CUTS biosensor as a regulator of a payload in response to TDP-43 loss of

function implicating its potential use in safeguarded gene replacement therapies associated with

TDP-43 LOF. The CUTS biosensor presented here represents a significant technological

advancement in available TDP-43 biosensors and offers a reliable and highly sensitive means to

dynamically monitor TDP-43 function. Implementing CUTS may facilitate the rapid screening of

potential therapeutics to restore TDP-43 functionality, monitor TDP-43 function to understand

disease-associated dysfunction, or deliver a genetic payload in response to TDP-43 loss.

Results

CUTS TDP-43 LOF sensor design utilizing known Cryptic Exons

To improve the detection of TDP-43 LOF, we designed a novel TDP-43 LOF sensor (TS)

for real-time screening-based detection using previously reported genes known to undergo

TDP-43 regulated splicing (*UNC13A* and *CFTR*). The TS cassettes are constructed with a

constitutively expressed mCherry, followed by a TDP-43 regulated CE and a GFP linked to a 3x

nuclear localization signal (NLS), each separated by a T2A self-cleavage sequence (Figure 1A).

We positioned the GFP reporter outside the mCherry open reading frame (ORF), introducing an

early stop codon upstream to the GFP (Figure 1A). This strategic design achieved three key

outcomes: (1) Under physiological TDP-43 levels, the binding of TDP-43 to the CE and UG-rich

sequence should promote complete intronic splicing, maintaining the in-frame stop codon and

allowing only mCherry expression. (2) TDP-43 loss and/or failure to bind the CE sites, will result

in CE retention and a subsequent frameshift, causing resulting in an out-of-frame stop-codon

GFP codon inclusion. (3) The GFP output of the TS should be proportional to the level of

TDP-43's LOF (Figure 1A).

Figure 1: CUTS is more precise than UNC13A-TS and CFTR-TS as a TDP-43 splicing sensor. *Comparison of stable polyclonal HEK cells expressing UNC13A-TS, CFTR-TS, or CUTS following treatment with siRNA control (siControl) (20nM) or TDP-43 (siTDP-43) (0.6nM - 20nM).*

Cells were reverse transfected with siRNA treatment in complete media supplemented with
doxycycline (1000 ng/mL). After 72 h, cells were analyzed by live imagining and protein lysate
*was harvested for western blot analysis. **(A)** Schematic of the TDP-43 loss of function Sensor*
*(TS) system design. **(B)** Overview of the UNC13A-TS, CFTR-TS, and CUTS cassette design. **(C)***
*Representative live imaging of TS comparison (10X). **(D)** Mean intensity quantification of GFP*
*signal intensity as shown in (C). **(E-G)** Western blot analysis of (E) UNC13A-TS, (F) CFTR-TS,*
*and (G) CUTS developing again GFP and TDP-43 proteins. **(H-J)** Relative pixel quantification of*
GFP and TDP-43 band normalized to total protein (Ponceau S) for the indicated TS from E-G.
Statistical significance was determined by one-way ANOVA and Tukey's multiple comparison
test (* = P < 0.03; ** = P < 0.002; *** = P < 0.0002; **** = P < 0.0001). Green = GFP signal; red =
mCherry signal. Scale bar = 100 µm. N=3 biological replicates.

Supplementary Figure 1: Dose-response of CUTS and TDP-43 knockdown. *Comparison of*
stable polyclonal HEK cells expressing UNC13A-TS, CFTR-TS, or CUTS following treatment
with siRNA control (siControl) (20nM) or TDP-43 (siTDP-43) (0.6nM - 20nM). Cells were reverse
transfected with siRNA treatment in complete media supplemented with doxycycline (1000
ng/mL). After 72 h, cells were analyzed by live imagining and protein lysate was harvested for
*western blot analysis. (**A**) Mean intensity quantification of GFP signal intensity from live imaging,*
*presented as fold change from mock. (**B**) Relative pixel quantification of GFP normalized to total*
protein (Ponceau S), presented as fold change from mock. Statistical significance was
determined by two-way ANOVA and Tukey's multiple comparison test (* = P < 0.03; ** = P <
0.002; *** = P < 0.0002; **** = P < 0.0001). N=3 biological replicates.

CUTS is more precise than CFTR and UNC13A as a TDP-43 LOF biosensor

To assess the functionality of TS, we generated CFTR-TS and UNC13A-TS, which

utilized the TDP-43 regulated CEs from *CFTR* and *UNC13A*, respectively (Figure 1B).

Additionally, we engineered a combined construct termed CUTS (CFTR-UNC13A TS),

integrating both CE sequences (Figure 1B). Additional base modifications were incorporated into

the cassette designs to prevent unexpected stop codons within the CE regions (see Table S1 for

full sequence details). Each construct was coupled with a Tet3g promotor, cloned into a

Piggybac vector, and stably expressed in HEK293 cells. To evaluate the accuracy and reliability

of the three TS constructs, we performed a TDP-43 LOF assay using increasing siRNA

concentrations, followed by live confocal imaging and western blot (WB) analysis (Figure 1 and

S1). Across all cell lines, we observed constitutive mCherry expression and an increased trend in

GFP signals with higher siTDP43 concentrations. The CFTR-TS construct exhibited a notably

high baseline, with detectable GFP leakage in control groups without TDP-43 loss (Figure 1C-1D,

1F, 1I, S1). In contrast, the UNC13A-TS construct demonstrated superior accuracy compared to

CFTR-TS, showing no detectable GFP expression under control conditions in both

imaging-based and WB analyses (Figure 1C-1E, 1H, S1). However, the UNC13A-TS showed

limited sensitivity, with only a modest GFP signal (17% of CUTS) under high quantities of

siTDP43 treatment. Interestingly, cells expressing the CUTS construct exhibited a synergistic

effect from both the CFTR and UNC13A CE sequences, achieving high sensitivity evidenced by

a clear dose-responsive GFP expression with increasing siTDP43 concentrations, while

maintaining high accuracy with minimal leakage via imaging and GFP immunoblotting (Figure

1C-1D, 1G, 1J, S1A-S1B). Given the promising accuracy of the CUTS sensor, we proceeded to

further characterize the CUTS RNA biosensor.

CUTS demonstrates ultra-sensitivity in detecting low-level TDP-43 loss-of-function

To challenge the stability and sensitivity of CUTS, we next conducted an ultra-low dose

TDP-43 siRNA transfection, ranging from 38 – 1200 pM. Immunofluorescence staining (IF)

revealed a consistent increase in both GFP intensity and GFP-positive cell ratios in

CUTS-expressing cells in response to elevated siTDP43, with minimal baseline expression

observed (Figure 2A-2B). We confirmed the ultra-sensitivity and accuracy of CUTS using WB

analysis, demonstrating measurable GFP even at the lowest doses of siTDP43 assessed. While

changes in TDP-43 levels were undetectable by WB at siTDP43 doses of 37.5 – 75 pM

(measured as 2% TDP-43 KD by WB), the CUTS system demonstrated a 7 to 55-fold increase in

GFP expression at these doses compared to baseline (Figure 2C-2D). This increase in GFP

expression continued consistently up to the highest dose of siTDP43 (1,200 pM; 98% TDP-43

KD), showcasing a 118,224-fold increase in GFP compared to baseline. Pearson's correlation

analysis confirmed a highly significant relationship between GFP expression and siTDP43 dose

(P = 0.0011), while the correlation between measurable TDP-43 and siTDP43 concentration was

less significant (P = 0.0429). Linear regression analysis between the logarithmic GFP fold

increase and siRNA doses demonstrated an exceptional linear relationship (R^2 = 0.9998). These

results indicate that GFP expression produced by CUTS is a more sensitive method for detecting

TDP-43 KD (and therefore LOF) than TDP-43 WB detection. The linear relationship between

GFP and siTDP-43 dose also demonstrates the potential of CUTS to be used as a predictive

model for TDP-43 LOF.

Figure 2: CUTS enables detection of TDP-43 loss-of-function with ultra-low siRNA treament. *Low-dose siRNA TDP-43 (siTDP-43) treatment was performed in stable polyclonal HEK cells expressing CUTS. CUTS-expressing cells were reverse transfected with siRNA control (siControl) or siTDP43 in a dose-response curve (38 to 1200pM) in doxycycline*

supplemented media (1000ng/ml) for 72hr. (A) Representative immunofluorescence images of
CUTS-expressing HEK cells under low doses of siRNA TDP-43 treatment. (60X). (B) Mean
intensity quantification of GFP signal from (A) with normalization to the number mCherry positive
cells. (C) Western blot of GFP and TDP-43 proteins from HEK cell lysate expressing CUTS
under low doses of siRNA TDP-43. Ponceau S is shown as a loading control. (D) Pixel intensity
quantification of the GFP and TDP-43 bands shown in (C), presented as fold-change from the
mock-treated sample. (E) Schematic showing the position of qPCR primers, developed to detect
CUTS cryptic exon inclusion (referred to as 'CUTS-CE' and 'CUTS-J'). (F) Representative
agarose gel showing qPCR product from melting curve detecting CUTS cryptic exon inclusion
using the primers shown in (E). (G) qPCR quantification of the siTDP-43 dose curve presented
as fold-change from the mock-treated sample. Purple text indicates the GFP fold change from
the mock-treated sample. Red text indicates the percentage of total detectable TDP-43
knockdown. Linear regression analysis shown in (D) and (G) was performed on Log values.
Fitting method = least squares regression. Green = GFP; red = mCherry. Scale bar = 50 µm. N=3
biological replicates.

We assessed CUTS' sensitivity at the transcript level using the siTDP43 dose curve and
RT-qPCR assessment in Figure 2C-D. To determine the relative amount of CUTS' CE retention,
we designed primers targeting either the entire transcript or the specific junction sites of the CE
(Figure 2E-2F). The RT-qPCR quantification demonstrated increased sensitivity at detecting
changes in TDP-43 levels compared to WB analysis, with the capability of detecting changes in
TDP-43 between each siTDP43 dose (1 – 70% TDP-43 KD) in a linear manner (R^2 = 0.9303).
Using CUTS detection, we observed a clear linear logarithmic relationship between the amount
of CUTS CE-retention and the increasing siTDP43 doses (R^2=0.9994). Even at 1% KD in
TDP-43, the CUTS system detected a 3-fold increase in CE retention compared to baseline,
which increased to a 1,488-fold increase at 70% TDP-43 KD (Figure 2G). As with our WB
analysis (Figure 2D), there was a significant correlation between both GFP expression and
TDP-43 when correlated to siTDP43 dose (P < 0.0001 and P = 0.0279, respectively). Thus,
these data indicate that CUTS is a reliable approach to quantifying TDP-43's loss across an
extensive range, highlighting its ability as a TDP-43 LOF biosensor. Additionally, CUTS
demonstrated ultra-sensitivity under low-level TDP-43 KD, beyond the detection limit of both WB
and RT-qPCR.

Pathological TDP-43 phase transitions or mislocalization activate the CUTS biosensor

In ALS/FTLD, the absolute TDP-43 level remains largely unaffected. Instead, TDP-43

undergoes pathological mislocalization and/or phase transitions likely due to a reduction in RNA

binding, which reduces the functional cellular TDP-43. To evaluate whether these events

contribute to TDP-43 loss-of-function, we tested CUTS's ability to detect TDP-43 LOF caused by

TDP-43 mislocalization or aggregation via aberrant phase transitions. We transfected CUTS

HEK293 cells with four tagged TDP-43 isoforms: (1) TDP-43^{WT}, (2) TDP-43^{cyto}, (3) TDP-43^{5FL},

and (4) TDP-43$^{cyto\ 5FL}$. The TDP-43^{cyto} variants contain point mutations located within the nuclear

localization signal (NLS) of TDP-43, resulting in cytoplasmic mislocalization (Igaz et al., 2011;

Mann et al., 2019). The 5FL form contains five phenylalanine-to-leucine mutations within the two

RNA recognition motif (RRM) domains of TDP-43 that greatly impaired TDP-43's RNA binding

ability and were previously reported to form aggregated "anisomes" inside the nucleus (Cohen et

al., 2015; Elden et al., 2010; Mann et al., 2019; Yu et al., 2021). The TDP-43$^{cyto\ 5FL}$ combines both

modifications, leading to insoluble cytoplasmic inclusions (Elden et al., 2010; Keating et al.,

2023; Lu et al., 2022; Mann et al., 2019; Yu et al., 2021). Excluding TDP-43^{WT}, all three modified

versions have proven to sequestrate endogenous TDP-43 into mislocalized or aggregated

inclusions (Keating et al., 2023). Therefore, our objective was to utilize CUTS to determine

whether the expression of these aggregation-prone TDP-43 variants elicits TDP-43 LOF.

The introduction of TDP-43^{cyto}, TDP-43^{5FL}, and TDP-43$^{cyto\ 5FL}$ induced nuclear GFP signal

when assessed by immunofluorescence analysis, while neither the tagged plasmid backbone

nor TDP-43^{WT} caused any detectable GFP (Figure 3A). We confirmed the expression of

exogenous and endogenous TDP-43 levels by WB and quantified the relative GFP level in each

condition (Figure 3B-3C). As we have previously shown that CUTS demonstrates a proportional

response to TDP-43's LOF (Figure 2), we were able to directly interpret the relative ability of the

different TDP-43 mutants to trigger TDP-43 LOF by comparison of their GFP levels. All three

TDP-43 mutants' expression triggered significant LOF compared to the control conditions, albeit

at varying significance levels. The most modest LOF effect was achieved by TDP-43^{cyto}, followed

by TDP-43$^{cyto\ 5FL}$, and TDP-43^{5FL} (Figure 3C). Interestingly, although both TDP-43^{5FL} and

TDP-43$^{cyto\ 5FL}$ caused significantly elevated LOF, the TDP-43^{5FL} mutant alone mediated greater

LOF than when combined with the NLS mutations highlighting the potential role of nuclear

homotypic TDP-43 interactions potentially contributing to TDP-43 LOF in disease absent its

cytoplasmic mislocalization. To further validate the functionality of exogenous TDP-43, we

transfected the same tagged TDP-43^{WT} into a HeLa TDP-43 knock-out cell line expressing CUTS

(Roczniak- Ferguson & Ferguson, 2019). We detected a significantly decreased GFP signal
compared to the backbone or non-transfected controls, which confirmed the full splicing function
of TDP-43^{WT} (Figure S2A-S2D). Owing to CUTS' performance, we show the functional
consequence caused by TDP-43's mislocalization and/or aberrant phase transitions,
demonstrating that aggregation-prone TDP-43 variants mediated direct LOF toxicity in addition
to any gain-of-function (GOF) toxic events. Furthermore, these results strongly support CUTS's
capability in measuring functional TDP-43 levels under broader contexts beyond TDP-43 KD.

Figure 3: Mislocalization or aberrant phase transitions induce TDP-43 LOF.
Stable HEK cells expressing CUTS were induced with doxycycline (1000 ng/mL) for 24 hours
before transfection with the following plasmids: pCMV backbone, TDP-43^{WT}, TDP-43$^{\Delta NLS}$,

TDP-43^{5FL}, TDP-43$^{\Delta NLS\ 5FL}$, or non-transfected. Following transfection, plasmids were expressed

*for 72 h, followed by live imaging and protein analysis. **(A)** Live-imaging of CUTS HEK cells*

*expressing WT or mutant TDP-43 gene cassettes. **(B)** Representative western blot of exogenous*

*and endogenous GFP and TDP-43. Ponceau S is shown as a loading control. **(C)** Relative GFP*

pixel intensity quantification of the band is shown in (B). Statistical significance was determined

by one-way ANOVA and Tukey's multiple comparison test (* = P < 0.03; ** = P < 0.002; *** = P <

0.0002; **** = P < 0.0001). Green = GFP; red = mCherry. Scale bar = 100 μm. N=3 biological

replicates.

Supplementary Figure 2: TDP-43^{wt} expression rescues loss-of-function in a HeLa TARDBP

Knockout Cell Line. *Transient CUTS expression in Hela TDP-43 KO we induced with*

doxocycline (1000n/ml) for 24 hours before transfection of pCM backbone or TDP-43^{WT}

plasmids. Following transfection, plasmids were expressed for 72 h, followed by live imaging and

*protein analysis. **(A)** Live-imaging of Hela TDP-43 KO expressing CUTS in combination with*

*TDP-43^{WT} or pCMV backbone control. **(B)** Representative WB of exogenous and endogenous*

*GFP and TDP-43. Ponceau S is shown as a loading control. **(C)** Relative GFP pixel intensity*

*quantification of the protein bands shown in (B). **(D)** Agarose gel showing RT-PCR product from*

CUTS cryptic exon inclusion from wildtype or TDP-43 KO Hela cell lines using the CUTS-CE

primers shown in Figure 2E. Statistical significance was determined by one-way ANOVA and

Tukey's multiple comparison test (* = P < 0.03; ** = P < 0.002; *** = P < 0.0002; **** = P <
0.0001). Green = GFP; red = mCherry. Scale bar = 100 μm. N=3 biological replicates.

CUTS-mediated autoregulated restoration of TDP-43 splicing.

Given the growing recognition of the role TDP-43 LOF is believed to play in disease
progression, numerous efforts have been committed to developing rescue methods aimed at
re-delivering TDP-43 or other gene payloads to restore its physiological splicing function
(Baughn et al., 2023; Mehta et al., 2023; Sun et al., 2017). However, a significant challenge in
LOF therapies lies in maintaining precise TDP-43 levels within neurons, as even slight
overexpression can lead to GOF toxicity (Johnson et al., 2009; Park et al., 2017; Yang et al.,
2022). Consequently, a generalized TDP-43 gene-replacement therapy without genome
integration carries a substantial risk of overexpression toxicity. Therefore, a CE biosensor such
as CUTS may be used to control cell- and temporal-specific regulation of a gene payload. To test
this, we generated a CUTS-controlled TDP-43 (CUTS-TDP43) transgene (Figure 4A).
Considering the ultra-sensitivity to TDP-43 LOF and minimal leakage under physiological
TDP-43 levels, CUTS-TDP43 may have the potential to autonomously negatively regulate its
expression, ensuring levels will not surpass physiological levels.

Figure 4: Autoregulatory rescue of TDP-43 loss-of-function by CUTS-TDP43 expression.
(A) Schematic of CUTS as an autoregulatory controller of TDP-43 expression (CUTS-TDP43).
(B-C) TDP-43 siRNA (siTDP43) dose-response curve in stable polyclonal HEK cells expressing
CUTS, CUTS-TDP43, or CUTS-TDP43 (codon optimized). The codon-optimized variation allows
for continued expression during siTDP-43 treatment. HEK cells expressing the CUTS,
CUTS-TDP-43 and CUTS-TDP-43 codon optimize system were reverse transfected with control
siRNA (siControl) or siTDP43 in a dose-response curve (0.6nM-20nM) in a doxycycline
(1000ng/ml) supplement media for 72hr. Cells were then used for live imaging or protein analysis.
(B) Live imaging of the CUTS variants. (C) Immunoblot assay of GFP and TDP-43. Ponceau S is
shown as a loading control. (D-E) CFTR minigene assay in stable CUTS or CUTS-TDP43 (codon
optimized) expressing HEK cells. Cells were induced with doxycycline (1000 ng/mL) for 24 h
before transfection with the CFTR minigene. Following an additional 24h of expression, cells

were transfected with 20nM siControl or siTDP-43. Cells were harvested 48 h following siRNA

transfection for RNA extraction and RT-PCR analysis. (D) PCR agarose gel of CFTR minigene.

(E) PCR analysis of the ratio between the CFTR cryptic exon inclusion and the correctly spliced

product from CFTR as shown in (D). Statistical significance was determined by student t-test (* =

P < 0.03; ** = P < 0.002; *** = P < 0.0002; **** = P < 0.0001). Green = GFP; red = mCherry.

Scale bar = 100 µm. N=3 biological replicates.

To test this, we created a new polyclonal stable line in HEK293 cells (CUTS-TDP43) by

replacing the 3xNLS in the original CUTS cassette with the *TARDBP* ORF fused to a GFP

reporter (Figure 4A). However, as our siTDP43 targets the sequence within the coding region,

CUTS-TDP43 was also knocked down upon siRNA transfection, shown by a generalized

decreased mCherry signal (Figure 4B). Therefore, we designed a codon-optimized

CUTS-TDP43^{CO} that is not targeted by siTDP43 (Figure 4B). Live imaging analyses showed that

CUTS GFP signal demonstrated a steady increase in expression in response to increasing

doses of siTDP43; however, the GFP signal from CUTS-TDP43^{CO} remained undetectable

(Figure 4B). WB analysis further demonstrated successful TDP-43 rescue under endogenous

TDP-43 KD, as shown by the increasing exogenous TDP-43 observed following decreases in

endogenous TDP-43 (Figure 4C). The amount of total TDP-43 appeared to remain consistent

throughout the increasing siTDP43 doses, indicating tight regulation of the rescue parameters.

To further confirm whether CUTS-TDP43^{CO} could rescue TDP-43 splicing functionality, we

performed a *CFTR* minigene assay (Ayala et al., 2006; Buratti & Baralle, 2001). The expression

of CUTS-TDP43^{SO} demonstrated partial, yet significant rescue of cryptic exon 9 splicing in *CFTR*

minigene, supporting its controlled efficacy in rescuing splicing LOF (Figure 4D-4E). Taken

together, these data indicate that CUTS can autoregulate a TDP-43 payload to physiological

levels in response to TDP-43 knockdown.

Discussion:

We developed and characterized the CUTS system, a novel approach to detect TDP-43

LOF. The CUTS system utilizes TDP-43-dependent CE events to correlate the level of TDP-43

LOF directly with the expression of a reporting gene. By combining the CFTR-TS and

UNC13A-TS, our findings demonstrate that the CUTS system provides an optimal balance of

sensitivity and accuracy. This was evidenced by the ability of CUTS to detect modest levels of

TDP-43 LOF, as shown by the dose-responsive increase in the expression of the reporter gene,

GFP, under various siTDP-43 concentrations (Figure 2). Furthermore, our results suggest that
CUTS can effectively discern TDP-43 LOF induced by pathological phase transitions or
mislocalization, a critical aspect in the context of neurodegenerative diseases containing TDP-43
pathology, such as ALS and FTLD. The CUTS system's potential for application in
gene-replacement therapies was also highlighted, offering a promising avenue for autoregulated
rescue of TDP-43 function, which is critical for avoiding the deleterious effects of TDP-43
overexpression.

This tool findings enables for the significant advancement in the capacity to detect
TDP-43 LOF using biosensor assays across diverse experimental settings and through multiple
analytical methods. Previously, the *CFTR* minigene assay has been the predominant approach
for detecting TDP-43 LOF (Ayala et al., 2006; Buratti & Baralle, 2001; Cohen et al., 2015;
Conicella et al., 2020; Jiang et al., 2017; Hermann Broder Schmidt et al., 2019). However, this
approach is associated with several limitations, all of which are effectively addressed by utilizing
the CUTS system. The first advantage of the CUTS system is its ability to detect TDP-43 LOF in
real-time through live-imaging analysis. Unlike *CFTR* minigene assays, which typically
necessitate endpoint experimental analysis, CUTS facilitates continuous monitoring, eliminating
the need for multiple fixed time points. Additionally, CUTS can be seamlessly integrated with
various analytical methods, including RT-qPCR (at the RNA level), WB analysis (at the protein
level), live imaging (for real-time assessment), and immunofluorescence imaging (to correlate
with relevant markers). While not evaluated in this study, it is conceivable that CUTS could be
adapted for use with flow cytometry-based techniques, leveraging GFP-positive cells as an
output for analysis, as previously demonstrated with a *CFTR*-modified sensor (H Broder Schmidt
& Rohatgi, 2020).

In addition to the expanded array of analytical methods offered by CUTS compared to
CFTR minigene assays, we anticipate that the CUTS system will exhibit superior sensitivity and
accuracy. This is supported by the comparison of CUTS with the CFTR-TS or UNC13A-TS
cassette (Figure 1), underscoring the potential of CUTS to outperform single minigene-based
approaches in TDP-43 LOF detection. While recent work in two recent preprints suggests other
CE biosensors are in development, CUTS appears to exhibit enhanced sensitivity (Wilkins et al.,
2023; Zhang et al., 2023). The expression of the GFP reporter in CUTS achieved up to
118,224-fold increase upon TDP-43 knockdown, compared to *ADNP2* (< 5-fold) (Zhang et al.,
2023); TDP-REGv1 (<20-fold); and TDP-REGv2 (<300-fold) (Wilkins et al., 2023). We also show
that CUTS can detect ultra-low levels of TDP-43 knockdown (increasing > 7-fold), below the WB

or RT-qPCR detection limit. Furthermore, CUTS exhibits a robust log-linear relationship to siRNA
doses, making it suitable for quantitative purposes.

Due to the high flexibility of the CUTS system, its application can be expanded *in vitro*
and *in vivo* when coupled with disease models. Integrating the CUTS system with disease
models enables the evaluation of the model's fidelity in recapitulating TDP-43 LOF phenotypes.
Such assessments are crucial for selecting appropriate models that faithfully replicate the
desired study context. Furthermore, coupling the CUTS system with TDP-43 models presents a
valuable approach for diverse screening studies. For example, CUTS can be leveraged for
high-throughput drug screening and CRISPR screening methodologies. Such approaches hold
promise for uncovering critical insights into cell-specific disease mechanisms, identifying pivotal
disease modifiers, and delineating potential therapeutic genetic targets (Aldewachi et al., 2021;
Bock et al., 2022).

A significant advantage of the CUTS system lies in its capacity to deliver precisely
regulated gene therapy for rescuing TDP-43 LOF. This study illustrates this capability by placing
a functional TDP-43 transcript downstream of the CUTS regulatory elements. The system's
self-regulating ability enhances its safety profile as a gene therapy approach, ensuring gene
expression occurs only when necessary and exclusively in cells lacking TDP-43 function.
Furthermore, this system can be expanded by substituting the TDP-43 transcript with other
genetic modifiers of disease, such as antibodies and PROTACs (Pozzi et al., 2019; Tseng et al.,
2023), or genes with established therapeutic potential, including heat shock proteins (HSPs) or
heterogeneous nuclear ribonucleoproteins (hnRNPs) (Koike et al., 2023; Lu et al., 2022; Yu et al.,
2021). This adaptability holds promise for achieving safe therapeutic outcomes without the need
for direct TDP-43 expression.

Methods:

Generation of plasmids

The CUTS sequence, plasmid, and map were originally generated in this study. The CFTR-TS,
UNC13A-TS and CUTS DNA sequences were designed *in silico* and *de novo* synthesized by
Genewiz. These sequences were and assembled into Tet3G vector between EcoRI and NotI
with NEBuilder HiFi DNA Assembly Master Mix (NEB, E2621L) following the manufacturer's

protocol. The full DNA sequence for CUTS, CFTR-TS, and UNC13A-TS can be found in Table
S1.

The CFTR minigene assay plasmid (pTB-CFTR-A455E) was a kind gift from Dr. Yuna Ayala.

The exogenous TDP-43 plasmids were constructed in a pCMV backbone by linking a
3xFlag-APEX2 protein (Addgene #164622) (Bonet- Ponce et al., 2020) to TDP-43 coding
sequences with WT, cyto, 5FL, or cyto 5FL modifications (Mann et al., 2019).

The codon-optimized *TARDBP* coding sequence (Table S1) was synthesized by IDT and
assembled downstream of the GFP sequence of CUTS with NEBuilder HiFi DNA Assembly
Master Mix.

All the primers were synthesized by IDT. All plasmids were verified using whole-plasmid
sequencing via Oxford Nanopore, provided by Plasmidsaurus.

Cell culture and transfection

Human Embryonic Kidney 293 (HEK293) cells (female genotype, acquired from the American
Type Culture Collection (ATCC)) and Hela TDP-43 knock-out (KO) cells (a kind gift from Dr.
Shawn M Ferguson) (Roczniak- Ferguson & Ferguson, 2019) were cultivated in Dulbecco's
Modified Eagle Medium high glucose, pyruvate (DMEM, Thermo Fisher Scientific, 10-313-039)
supplemented with 10% HyClone Bovine Growth Serum (Cytiva HyClon, SH3054103HI) and 1X
GlutaMAX (Thermo Fisher Scientific, 10-313-039). Cells were incubated at 37°C in a 5% CO_2
atmosphere with high humidity. For transfection assays, cells were plated on collagen-coated
coverslips or dishes (50 µg/mL, GIBCO) and transfected with designated DNA quantities using
Lipofectamine 3000 (Thermo Scientific, L3000015) following the provider's protocol.

HEK293 stable cell line generation via Piggybac transposition

For stable cell line creation, HEK293 cells were pre-plated on 6-well plates and transfected at
approximately 70% confluence with 2.5 µg of Piggybac plasmids encoding CUTS, CFTR-TS,
UNC13A-TS, and CUTS-TDP43 alongside 0.5 µg of the Super PiggyBac Transposase
Expressing plasmid (PB200PA-1) using Lipofectamine 3000, according to the manufacturer's
guidelines. A transfection control without transposase was included. Following a 48-hour
post-transfection period, cells were selected with puromycin (Sigma, P8833) at 5 µg/mL, with

media changes every two days. Selection resulted in control cell death within approximately 5

days, while surviving populations were expanded and maintained in reduced puromycin

concentrations (2.5 μg/mL) to establish stable lines. Expression of the transgenes was confirmed

by immunofluorescence staining and Western blot analysis.

SDS-PAGE and Western blot

For protein analysis, cells were lysed directly on the plate using fresh and pre-chilled Urea-RIPA

buffer: 2M fresh urea in 1XRIPA buffer (Boston Bioproducts, BP-115X), supplemented with 1%

protease inhibitor cocktail (Sigma, P8340) and sonicated. Protein concentrations were quantified

using the Pierce BCA Protein Assay Kit (Thermo Scientific, 23227). Proteins were resolved by

SDS-PAGE and transferred to nitrocellulose membranes for WB analysis. Membranes were

blocked and probed with primary antibodies: mouse-anti-GFP (Santa Cruz, sc-9996, 1:200),

mouse-anti-α-tubulin (Sigma, T5168, 1:1000), rabbit-anti-TDP-43 (Proteintech, 10782-2-AP,

1:2500), and rabbit-anti-mCherry (Cell Signaling, 43590, 1:1000), followed by HRP-conjugated

secondary antibodies: donkey-anti-mouse (JacksonImmunoResearch 715035151, 1:5000) or

donkey-anti-rabbit (JacksonImmunoResearch 711035152, 1:5000). Detection was achieved

using Western Lightning ECL Pro (Revvity, NEL1201001EA) or Supersignal West Femto

Maximum Sensitivity Chemiluminescent Substrate (Thermo Scientific, 34095) in an Amersham

ImageQuant 800 GxP biomolecular imager system (Amersham, 29653452).

Confocal microscopy

Confocal imaging was performed on a Nikon A1 laser-scanning microscope using either a 60X

oil immersion or a 10X/20X objective for live-cell observations. A Tokai HIT stage-top incubator

maintained the required environmental conditions. Nikon Elements software facilitated image

acquisition and analysis. Representative images were chosen from at least two independent

experiments with a minimum of three biological replicates each.

siRNA reverse transfection

siRNA reverse transfections were conducted using Lipofectamine RNAiMAX reagent (Thermo

Scientific, 13778150), adhering to the supplier's protocol. To knockdown TDP-43, the following

siRNAs were used: ON-TARGETplus SMARTpool siRNA against *TARDBP* (Dharmaco,
L-012394-00-0005) and siGENOME non-Targeting siRNA for control (Dharmaco,
D-001206-13-05).

RNA extraction, RT-PCR, and qPCR

RNA extraction was performed using the RNeasy Mini Kit (Qiagen, 74106) with concentration
determinations via Nanodrop Spectrophotometer (Nanodrop, ND-1000). Reverse transcription of
extracted RNA (0.2 µg) to cDNA utilized iScript Reverse Transcription Supermix (Bio-Rad,
1708841) in accordance with the manufacturer's guidelines.

The RT-PCR or RT-qPCR was conducted with cDNA diluted 10-fold. For RT-PCR assay, CFTR
cryptic exon region was amplified with the following primer pair: P690-F
(5'-CAACTTCAAGCTCCTAAGCCACTGCCTGC) and P691-R
(5'-TAGGATCCGGTCACCAGGAAGTTGGTTAAATCA). CUTS' cryptic exon region was
amplified with the following primer pair: CUTS-CE-F (5'-ATCCCGGCCCTGGATCCG) and
CUTS-CE-R (5'-GTCAGCTTGCCGTAGGTGGC). PCR products were separated by agarose gel
electrophoresis, and the bands were visualized with Amersham ImageQuant 800 GxP
biomolecular imager system.

For RT-qPCR assay, SsoAdvanced™ Universal SYBR Green Supermix (Biorad, 1725272) was
used following the supplier's protocol on a CFX96 Touch Real-Time PCR Detection System
(Biorad). Three technical replicates were included for each sample with the following program:
95°C for 30 s, 40 cycles of 95°C for 15 s and 60°C for 20 s. CUTS-CE-F
(5'-ATCCCGGCCCTGGATCCG) and CUTS-CE-R (5'-GTCAGCTTGCCGTAGGTGGC) were
used to quantify normal CUTS transcript. CUTS-J-F (5'-TCCGGCGAGGGATTTGGG) and
CUTS-J-R (5'-CCCCACCTAGACCCATCTCTCC) were primers targeting the cryptic exon
junctions to quantify cryptic exon-specific CUTS transcript. Relative quantification of CUTS
cryptic exon was determined by the ΔCt value of CUTS-J normalized to CUTS-CE.

Statistical analysis

Statistical significance was evaluated using GraphPad Prism 9 software, and specific tests used
for each experiment are outlined in the respective figure legends.

Authorship contributions

501   LX, JM, and CAB share equal contributions and authorship. LX and BTH conceptualized the

503   study. LX, JM, and CAB were responsible for data curation, data analysis, conceptualizing and

504   performing the described methodologies, compiling and generating data, and writing the original

505   draft. JX assisted with data analyses and experimental optimization. CJD conceptualized the

506   approach, funded the research, and prepared the final manuscript.

Acknowledgments

We thank Dr. Yuna Ayala, Ph.D. (St. Louis University School of Medicine) for kindly providing the

CFTR minigene plasmids. We thank Dr. Shawn M Ferguson, Ph.D. (Yale University School of

Medicine) for kindly providing the TARDBP$^{-/-}$ HeLa cell lines used in this study. We thank Olivia R.

Shapiro and Jocelyn C. Mauna (Donnelly Lab, University of Pittsburgh School of Medicine) for

their kind assistance and helpful discussions. This work was supported by funds to C.J.D. by the

LiveLikeLou Fund at the Pittsburgh Foundation and grants from NIH (R01NS105756,

R01NS127187).

Conflict of Interest Statement

A provisional patent has been filed for the CUTS1 biosensor with CJD, LX, CM, and CAB are

listed as co-inventors.

Reference

Aldewachi, H., Al-Zidan, R. N., Conner, M. T., & Salman, M. M. (2021). High-Throughput

Screening Platforms in the Discovery of Novel Drugs for Neurodegenerative Diseases.

Bioengineering, *8*(2). https://doi.org/10.3390/bioengineering8020030

Arai, T., Hasegawa, M., Akiyama, H., Ikeda, K., Nonaka, T., Mori, H., Mann, D., Tsuchiya, K.,

Yoshida, M., Hashizume, Y., & Oda, T. (2006). TDP-43 is a component of ubiquitin-positive

tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis.

Biochemical and Biophysical Research Communications, *351*(3), 602–611.

https://doi.org/10.1016/j.bbrc.2006.10.093

Ayala, Y. M., Pagani, F., & Baralle, F. E. (2006). TDP43 depletion rescues aberrant CFTR exon 9

skipping. *FEBS Letters*, *580*(5), 1339–1344. https://doi.org/10.1016/j.febslet.2006.01.052

Baughn, M. W., Melamed, Z., López-Erauskin, J., Beccari, M. S., Ling, K., Zuberi, A., Presa, M.,

Gonzalo-Gil, E., Maimon, R., Vazquez-Sanchez, S., Chaturvedi, S., Bravo-Hernández, M.,

Taupin, V., Moore, S., Artates, J. W., Acks, E., Ndayambaje, I. S., Agra de Almeida Quadros,

A. R., Jafar-Nejad, P., … Cleveland, D. W. (2023). Mechanism of STMN2 cryptic

splice-polyadenylation and its correction for TDP-43 proteinopathies. *Science*, *379*(6637),

1140–1149. https://doi.org/10.1126/science.abq5622

Bock, C., Datlinger, P., Chardon, F., Coelho, M. A., Dong, M. B., Lawson, K. A., Lu, T., Maroc, L.,

Norman, T. M., Song, B., Stanley, G., Chen, S., Garnett, M., Li, W., Moffat, J., Qi, L. S.,

Shapiro, R. S., Shendure, J., Weissman, J. S., & Zhuang, X. (2022). High-content CRISPR

screening. *Nature Reviews Methods Primers*, *2*(1), 8.

https://doi.org/10.1038/s43586-021-00093-4

Bonet-Ponce, L., Beilina, A., Williamson, C. D., Lindberg, E., Kluss, J. H., Saez-Atienzar, S.,

Landeck, N., Kumaran, R., Mamais, A., Bleck, C. K. E., Li, Y., & Cookson, M. R. (2020).

LRRK2 mediates tubulation and vesicle sorting from lysosomes. *Science Advances*, *6*(46). https://doi.org/10.1126/sciadv.abb2454

Brown, A.-L., Wilkins, O. G., Keuss, M. J., Hill, S. E., Zanovello, M., Lee, W. C., Bampton, A., Lee, F. C. Y., Masino, L., Qi, Y. A., Bryce-Smith, S., Gatt, A., Hallegger, M., Fagegaltier, D., Phatnani, H., NYGC ALS Consortium, Newcombe, J., Gustavsson, E. K., Seddighi, S., … Fratta, P. (2022). TDP-43 loss and ALS-risk SNPs drive mis-splicing and depletion of UNC13A. *Nature*, *603*(7899), 131–137. https://doi.org/10.1038/s41586-022-04436-3

Buratti, E., & Baralle, F. E. (2001). Characterization and functional implications of the RNA binding properties of nuclear factor TDP-43, a novel splicing regulator of CFTR exon 9. *The Journal of Biological Chemistry*, *276*(39), 36337–36343. https://doi.org/10.1074/jbc.M104236200

Buratti, E., Dörk, T., Zuccato, E., Pagani, F., Romano, M., & Baralle, F. E. (2001). Nuclear factor TDP-43 and SR proteins promote in vitro and in vivo CFTR exon 9 skipping. *The EMBO Journal*, *20*(7), 1774–1784. https://doi.org/10.1093/emboj/20.7.1774

Cohen, T. J., Hwang, A. W., Restrepo, C. R., Yuan, C.-X., Trojanowski, J. Q., & Lee, V. M. Y. (2015). An acetylation switch controls TDP-43 function and aggregation propensity. *Nature Communications*, *6*, 5845. https://doi.org/10.1038/ncomms6845

Conicella, A. E., Dignon, G. L., Zerze, G. H., Schmidt, H. B., D'Ordine, A. M., Kim, Y. C., Rohatgi, R., Ayala, Y. M., Mittal, J., & Fawzi, N. L. (2020). TDP-43 α-helical structure tunes liquid-liquid phase separation and function. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(11), 5883–5894. https://doi.org/10.1073/pnas.1912055117

Elden, A. C., Kim, H.-J., Hart, M. P., Chen-Plotkin, A. S., Johnson, B. S., Fang, X., Armakola, M., Geser, F., Greene, R., Lu, M. M., Padmanabhan, A., Clay-Falcone, D., McCluskey, L., Elman, L., Juhr, D., Gruber, P. J., Rüb, U., Auburger, G., Trojanowski, J. Q., … Gitler, A. D. (2010).

Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for

ALS. *Nature*, *466*(7310), 1069–1075. https://doi.org/10.1038/nature09320

Feldman, E. L., Goutman, S. A., Petri, S., Mazzini, L., Savelieff, M. G., Shaw, P. J., & Sobue, G.

(2022). Amyotrophic lateral sclerosis. *The Lancet*, *400*(10360), 1363–1380.

https://doi.org/10.1016/S0140-6736(22)01272-7

Hurtle, B. T., Xie, L., & Donnelly, C. J. (2023). Disrupting pathologic phase transitions in

neurodegeneration. *The Journal of Clinical Investigation*.

Igaz, L. M., Kwong, L. K., Lee, E. B., Chen-Plotkin, A., Swanson, E., Unger, T., Malunda, J., Xu,

Y., Winton, M. J., Trojanowski, J. Q., & Lee, V. M.-Y. (2011). Dysregulation of the

ALS-associated gene TDP-43 leads to neuronal death and degeneration in mice. *The Journal*

of Clinical Investigation, *121*(2), 726–738. https://doi.org/10.1172/JCI44867

Jiang, L.-L., Xue, W., Hong, J.-Y., Zhang, J.-T., Li, M.-J., Yu, S.-N., He, J.-H., & Hu, H.-Y. (2017).

The N-terminal dimerization is required for TDP-43 splicing activity. *Scientific Reports*, *7*(1),

6196. https://doi.org/10.1038/s41598-017-06263-3

Johnson, B. S., Snead, D., Lee, J. J., McCaffery, J. M., Shorter, J., & Gitler, A. D. (2009). TDP-43

is intrinsically aggregation-prone, and amyotrophic lateral sclerosis-linked mutations

accelerate aggregation and increase toxicity. *The Journal of Biological Chemistry*, *284*(30),

20329–20339. https://doi.org/10.1074/jbc.M109.010264

Keating, S. S., Bademosi, A. T., San Gil, R., & Walker, A. K. (2023). Aggregation-prone TDP-43

sequesters and drives pathological transitions of free nuclear TDP-43. *Cellular and Molecular*

Life Sciences, *80*(4), 95. https://doi.org/10.1007/s00018-023-04739-2

Klim, J. R., Williams, L. A., Limone, F., Guerra San Juan, I., Davis-Dusenbery, B. N., Mordes, D.

A., Burberry, A., Steinbaugh, M. J., Gamage, K. K., Kirchner, R., Moccia, R., Cassel, S. H.,

Chen, K., Wainger, B. J., Woolf, C. J., & Eggan, K. (2019). ALS-implicated protein TDP-43

sustains levels of STMN2, a mediator of motor neuron growth and repair. *Nature*

Neuroscience, *22*(2), 167–179. https://doi.org/10.1038/s41593-018-0300-4

Koike, Y., Pickles, S., Estades Ayuso, V., Jansen-West, K., Qi, Y. A., Li, Z., Daughrity, L. M., Yue,

M., Zhang, Y.-J., Cook, C. N., Dickson, D. W., Ward, M., Petrucelli, L., & Prudencio, M. (2023).

TDP-43 and other hnRNPs regulate cryptic exon inclusion of a key ALS/FTD risk gene,

UNC13A. *PLoS Biology*, *21*(3), e3002028. https://doi.org/10.1371/journal.pbio.3002028

Kuo, P.-H., Doudeva, L. G., Wang, Y.-T., Shen, C.-K. J., & Yuan, H. S. (2009). Structural insights

into TDP-43 in nucleic-acid binding and domain interactions. *Nucleic Acids Research*, *37*(6),

1799–1808. https://doi.org/10.1093/nar/gkp013

Ling, J. P., Pletnikova, O., Troncoso, J. C., & Wong, P. C. (2015). TDP-43 repression of

nonconserved cryptic exons is compromised in ALS-FTD. *Science*, *349*(6248), 650–655.

https://doi.org/10.1126/science.aab0983

Ling, S.-C., Polymenidou, M., & Cleveland, D. W. (2013). Converging mechanisms in ALS and

FTD: disrupted RNA and protein homeostasis. *Neuron*, *79*(3), 416–438.

https://doi.org/10.1016/j.neuron.2013.07.033

Lukavsky, P. J., Daujotyte, D., Tollervey, J. R., Ule, J., Stuani, C., Buratti, E., Baralle, F. E.,

Damberger, F. F., & Allain, F. H.-T. (2013). Molecular basis of UG-rich RNA recognition by the

human splicing factor TDP-43. *Nature Structural & Molecular Biology*, *20*(12), 1443–1449.

https://doi.org/10.1038/nsmb.2698

Lu, S., Hu, J., Arogundade, O. A., Goginashvili, A., Vazquez-Sanchez, S., Diedrich, J. K., Gu, J.,

Blum, J., Oung, S., Ye, Q., Yu, H., Ravits, J., Liu, C., Yates, J. R., & Cleveland, D. W. (2022).

Heat-shock chaperone HSPB1 regulates cytoplasmic TDP-43 phase separation and

liquid-to-gel transition. *Nature Cell Biology*, *24*(9), 1378–1393.

https://doi.org/10.1038/s41556-022-00988-8

Mann, J. R., Gleixner, A. M., Mauna, J. C., Gomes, E., DeChellis-Marks, M. R., Needham, P. G.,

Copley, K. E., Hurtle, B., Portz, B., Pyles, N. J., Guo, L., Calder, C. B., Wills, Z. P., Pandey, U.

B., Kofler, J. K., Brodsky, J. L., Thathiah, A., Shorter, J., & Donnelly, C. J. (2019). RNA Binding

Antagonizes Neurotoxic Phase Transitions of TDP-43. *Neuron*, *102*(2), 321-338.e8.

https://doi.org/10.1016/j.neuron.2019.01.048

Ma, X. R., Prudencio, M., Koike, Y., Vatsavayai, S. C., Kim, G., Harbinski, F., Briner, A.,

Rodriguez, C. M., Guo, C., Akiyama, T., Schmidt, H. B., Cummings, B. B., Wyatt, D. W., Kurylo,

K., Miller, G., Mekhoubad, S., Sallee, N., Mekonnen, G., Ganser, L., … Gitler, A. D. (2022).

TDP-43 represses cryptic exon inclusion in the FTD-ALS gene UNC13A. *Nature*, *603*(7899),

124–130. https://doi.org/10.1038/s41586-022-04424-7

McKee, A. C., Gavett, B. E., Stern, R. A., Nowinski, C. J., Cantu, R. C., Kowall, N. W., Perl, D. P.,

Hedley-Whyte, E. T., Price, B., Sullivan, C., Morin, P., Lee, H.-S., Kubilus, C. A., Daneshvar, D.

H., Wulff, M., & Budson, A. E. (2010). TDP-43 proteinopathy and motor neuron disease in

chronic traumatic encephalopathy. *Journal of Neuropathology and Experimental Neurology*,

69(9), 918–929. https://doi.org/10.1097/NEN.0b013e3181ee7d85

Mehta, P. R., Brown, A.-L., Ward, M. E., & Fratta, P. (2023). The era of cryptic exons:

implications for ALS-FTD. *Molecular Neurodegeneration*, *18*(1), 16.

https://doi.org/10.1186/s13024-023-00608-5

Melamed, Z., López-Erauskin, J., Baughn, M. W., Zhang, O., Drenner, K., Sun, Y., Freyermuth,

F., McMahon, M. A., Beccari, M. S., Artates, J. W., Ohkubo, T., Rodriguez, M., Lin, N., Wu, D.,

Bennett, C. F., Rigo, F., Da Cruz, S., Ravits, J., Lagier-Tourenne, C., & Cleveland, D. W.

(2019). Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of

TDP-43-dependent neurodegeneration. *Nature Neuroscience*, *22*(2), 180–190.

https://doi.org/10.1038/s41593-018-0293-z

Nelson, P. T., Dickson, D. W., Trojanowski, J. Q., Jack, C. R., Boyle, P. A., Arfanakis, K.,

Rademakers, R., Alafuzoff, I., Attems, J., Brayne, C., Coyle-Gilchrist, I. T. S., Chui, H. C.,

Fardo, D. W., Flanagan, M. E., Halliday, G., Hokkanen, S. R. K., Hunter, S., Jicha, G. A.,

Katsumata, Y., … Schneider, J. A. (2019). Limbic-predominant age-related TDP-43

encephalopathy (LATE): consensus working group report. *Brain: A Journal of Neurology*,

142(6), 1503–1527. https://doi.org/10.1093/brain/awz099

Neumann, M., Sampathu, D. M., Kwong, L. K., Truax, A. C., Micsenyi, M. C., Chou, T. T., Bruce,

J., Schuck, T., Grossman, M., Clark, C. M., McCluskey, L. F., Miller, B. L., Masliah, E.,

Mackenzie, I. R., Feldman, H., Feiden, W., Kretzschmar, H. A., Trojanowski, J. Q., & Lee, V.

M.-Y. (2006). Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic

lateral sclerosis. *Science*, *314*(5796), 130–133. https://doi.org/10.1126/science.1134108

Park, S.-K., Hong, J. Y., Arslan, F., Kanneganti, V., Patel, B., Tietsort, A., Tank, E. M. H., Li, X.,

Barmada, S. J., & Liebman, S. W. (2017). Overexpression of the essential Sis1 chaperone

reduces TDP-43 effects on toxicity and proteolysis. *PLoS Genetics*, *13*(5), e1006805.

https://doi.org/10.1371/journal.pgen.1006805

Polymenidou, M., Lagier-Tourenne, C., Hutt, K. R., Huelga, S. C., Moran, J., Liang, T. Y., Ling,

S.-C., Sun, E., Wancewicz, E., Mazur, C., Kordasiewicz, H., Sedaghat, Y., Donohue, J. P.,

Shiue, L., Bennett, C. F., Yeo, G. W., & Cleveland, D. W. (2011). Long pre-mRNA depletion

and RNA missplicing contribute to neuronal vulnerability from loss of TDP-43. *Nature*

Neuroscience, *14*(4), 459–468. https://doi.org/10.1038/nn.2779

Pozzi, S., Thammisetty, S. S., Codron, P., Rahimian, R., Plourde, K. V., Soucy, G., Bareil, C.,

Phaneuf, D., Kriz, J., Gravel, C., & Julien, J.-P. (2019). Virus-mediated delivery of antibody

targeting TAR DNA-binding protein-43 mitigates associated neuropathology. *The Journal of*

Clinical Investigation, *129*(4), 1581–1595. https://doi.org/10.1172/JCI123931

Roczniak-Ferguson, A., & Ferguson, S. M. (2019). Pleiotropic requirements for human TDP-43 in

the regulation of cell and organelle homeostasis. *Life Science Alliance*, *2*(5).

https://doi.org/10.26508/lsa.201900358

Schmidt, Hermann Broder, Barreau, A., & Rohatgi, R. (2019). Phase separation-deficient TDP43

remains functional in splicing. *Nature Communications, 10*(1), 4890.

https://doi.org/10.1038/s41467-019-12740-2

Schmidt, H Broder, & Rohatgi, R. (2020). High-throughput Flow Cytometry Assay to Investigate

TDP43 Splicing Function. *Bio-Protocol*, *10*(8), e3594. https://doi.org/10.21769/BioProtoc.3594

Shin, J. E., Geisler, S., & DiAntonio, A. (2014). Dynamic regulation of SCG10 in regenerating

axons after injury. *Experimental Neurology*, *252*, 1–11.

https://doi.org/10.1016/j.expneurol.2013.11.007

Shin, J. E., Miller, B. R., Babetto, E., Cho, Y., Sasaki, Y., Qayum, S., Russler, E. V., Cavalli, V.,

Milbrandt, J., & DiAntonio, A. (2012). SCG10 is a JNK target in the axonal degeneration

pathway. *Proceedings of the National Academy of Sciences of the United States of America*,

109(52), E3696-705. https://doi.org/10.1073/pnas.1216204109

Sun, M., Bell, W., LaClair, K. D., Ling, J. P., Han, H., Kageyama, Y., Pletnikova, O., Troncoso, J.

C., Wong, P. C., & Chen, L. L. (2017). Cryptic exon incorporation occurs in Alzheimer's brain

lacking TDP-43 inclusion but exhibiting nuclear clearance of TDP-43. *Acta Neuropathologica*,

133(6), 923–931. https://doi.org/10.1007/s00401-017-1701-2

Tremblay, C., St-Amour, I., Schneider, J., Bennett, D. A., & Calon, F. (2011). Accumulation of

transactive response DNA binding protein 43 in mild cognitive impairment and Alzheimer

disease. *Journal of Neuropathology and Experimental Neurology*, *70*(9), 788–798.

https://doi.org/10.1097/NEN.0b013e31822c62cf

Tseng, Y.-L., Lu, P.-C., Lee, C.-C., He, R.-Y., Huang, Y.-A., Tseng, Y.-C., Cheng, T.-J. R.,

Huang, J. J.-T., & Fang, J.-M. (2023). Degradation of neurodegenerative disease-associated

TDP-43 aggregates and oligomers via a proteolysis-targeting chimera. *Journal of Biomedical*

Science, *30*(1), 27. https://doi.org/10.1186/s12929-023-00921-7

Varoqueaux, F., Sigler, A., Rhee, J.-S., Brose, N., Enk, C., Reim, K., & Rosenmund, C. (2002).

Total arrest of spontaneous and evoked synaptic transmission but normal synaptogenesis in

the absence of Munc13-mediated vesicle priming. *Proceedings of the National Academy of*

Sciences of the United States of America, *99*(13), 9037–9042.

https://doi.org/10.1073/pnas.122623799

Wilkins, O. G., Chien, M. Z. Y. J., Wlaschin, J. J., Pisliakova, M., Thompson, D., Digby, H.,

Simkin, R. L., Diaz, J. A., Mehta, P. R., Keuss, M. J., Zanovello, M., Brown, A.-L., Harley, P.,

Darbey, A., Karda, R., Fisher, E. M. C., Cunningham, T. J., Le Pichon, C. E., Ule, J., & Fratta,

P. (2023). Creation of de novo cryptic splicing for ALS/FTD precision medicine. *BioRxiv*.

https://doi.org/10.1101/2023.11.15.565967

Willemse, S. W., Harley, P., van Eijk, R. P. A., Demaegd, K. C., Zelina, P., Pasterkamp, R. J.,

van Damme, P., Ingre, C., van Rheenen, W., Veldink, J. H., Kiernan, M. C., Al-Chalabi, A., van

den Berg, L. H., Fratta, P., & van Es, M. A. (2023). UNC13A in amyotrophic lateral sclerosis:

from genetic association to therapeutic target. *Journal of Neurology, Neurosurgery, and*

Psychiatry, *94*(8), 649–656. https://doi.org/10.1136/jnnp-2022-330504

Yang, C., Qiao, T., Yu, J., Wang, H., Guo, Y., Salameh, J., Metterville, J., Parsi, S., Yusuf, I.,

Brown, R. H., Cai, H., & Xu, Z. (2022). Low-level overexpression of wild type TDP-43 causes

late-onset, progressive neurodegeneration and paralysis in mice. *Plos One*, *17*(2), e0255710.

https://doi.org/10.1371/journal.pone.0255710

Yu, H., Lu, S., Gasior, K., Singh, D., Vazquez-Sanchez, S., Tapia, O., Toprani, D., Beccari, M. S.,

Yates, J. R., Da Cruz, S., Newby, J. M., Lafarga, M., Gladfelter, A. S., Villa, E., & Cleveland, D.

W. (2021). HSP70 chaperones RNA-free TDP-43 into anisotropic intranuclear liquid spherical

shells. *Science*, *371*(6529). https://doi.org/10.1126/science.abb4309

Zhang, X., Das, T., Kalab, P., & Hayes, L. R. (2023). Multivalent GU-rich oligonucleotides

sequester TDP-43 in the nucleus by inducing high molecular weight RNP complexes. *BioRxiv*.

https://doi.org/10.1101/2023.08.01.551528

Supplementary Table 1. Information of DNA sequence

Domains of the sequence	Sequence
1. UNC13A-TS mCherry Linker-T2A HBA1_exon2 HBA1_intron2 HBA1_exon3 UNC13A_modified_CE UNC13A_intron20 Linker-T2A EGFP 3XNLS	ATGGTGAGCAAGGGCGAGGAGGATAACATGGCCATCATCAAGG AGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGG CCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCCTA CGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGTGG CCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGT ACGGCTCCAAGGCCTACGTGAAGCACCCCGCCGACATCCCCGA CTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGC GTGATGAACTTCGAGGACGGCGGCGTGGTGACCGTGACCCAGG ACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTG CGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGA AGACCATGGGCTGGGAGGCCTCCTCCGAGCGGATGTACCCCGA GGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCT GAAGGACGGCGGCCACTACGACGCTGAGGTCAAGACCACCTAC AAGGCCAAGAAGCCCGTGCAGCTGCCCGGCGCCTACAACGTCA ACATCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATC GTGGAACAGTACGAACGCGCCGAGGGCCGCCACTCCACCGGC GGCATGGACGAGCTGTACAAGGGAGGTGGGGGATCTGGTGAG GGCAGAGGCAGTCTGCTGACATGCGGTGACGTGGAAGAGAATC CCGGCCCTGGATCCGGCCCGGTCAACTTCAAGGTGAGCGGCGG GCCGGGAGCGATCTGGGTCGAGGGGCGAGATGGCGCCTTCCT CGCAGGGCAGAGGATCACGCGGGTTGCGGGAGGTGTAGCGCA GGCGGCGGCTGCGGGCCTGGGCCCTCGGCCCCACTGACCCTC TTCTCTGCACAGCTCCTAAGCCACTGCCTGCTGGTGACCTCTGC CTGGGTTTCCTGGAAAGAACTCTTATCCCCAGGAACTAGTTTGTT GAATAAATGCTGGTGTGAATGTGAATGGATGATTGAACAGATGA ATGAGTGTGATGAGTATGATAAAAGCATGGATGGAGAGATGGGT GAGTACATGGATGGATAGATGGATGAGTTGGTGGGTAGATTCGT GGCTAGATGGATGATGGATGGATGGATGGATGGATGGATGGAT GGACAGATGGATGGATATATGATTGAACTATTGAAAGTATAGATG TATGGATGGGTGAATTTGGGGGTAATTGTTAGATGATGGATGTG GATGGTTGTGAGTGGCTGGTGGACAGACGAAAATGGATGGGT GGATAAATTGATGGGTGGATGGATGGTTGGTTGTATGAAAGAAT GAATGATTGGGTAGATAAAAGAGTAGATGAATGAATTAATGAATA AACAGGCAGATGGATGATGTAAGCTGCCCCAGACCCTGGGACC TCTGACCCCCGGCGACCCCTTGCACTCTCCATGAGCCCTTTCCT

	TTTTTTTCCTCAGGTCTAGGTGGGGGATCTGGTGAGGGCAGAGG CAGTCTGCTGACATGCGGTGACGTGGAAGAGAATCCCGGCCCT TCTAGAAGCAAGGGCGAGGAGCTGTTCACCGGGGTGGTGCCCA TCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAG CGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCT GACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCC TGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTGCT TCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAG TCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTT CAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTC GAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCG ACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTA CAACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGA AGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAG GACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCC CCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCT GAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGC GATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATCA CTCTCGGCATGGACGAGCTGTACAAGCTGCAGAGCAGGGCCGA CCCCAAGAAGAAGAGGAAGGTGGACCCCAAGAAGAAGAGGAAG GTGGACCCCAAGAAGAAGAGGAAGGTGTGA
2. CFTR-TS mCherry Linker-T2A CFTR_intron 8 CFTR_modifi ed_exon9 CFTR_intron 9 Linker-T2A EGFP 3XNLS	ATGGTGAGCAAGGGCGAGGAGGATAACATGGCCATCATCAAGG AGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGG CCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCCTA CGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGGTGG CCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGT ACGGCTCCAAGGCCTACGTGAAGCACCCCGCCGACATCCCCGA CTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGC GTGATGAACTTCGAGGACGGCGGCGTGGTGACCGTGACCCAGG ACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTG CGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGA AGACCATGGGCTGGGAGGCCTCCTCCGAGCGGATGTACCCCGA GGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCT GAAGGACGGCGGCCACTACGACGCTGAGGTCAAGACCACCTAC AAGGCCAAGAAGCCCGTGCAGCTGCCCGGCGCCTACAACGTCA ACATCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATC GTGGAACAGTACGAACGCGCCGAGGGCCGCCACTCCACCGGC GGCATGGACGAGCTGTACAAGGGAGGTGGGGGATCTGGTGAG GGCAGAGGCAGTCTGCTGACATGCGGTGACGTGGAAGAGAATC

	CCGGCCCTGGATCCGGCGAGGTGAGTATGGTACATAAAACAAG CATCTATTGAAAATATCTGACAAACTCATCTTTTATTTTTGATGTG TGTGTGTGTGTGTGTGTGTGTTTTTTTTTTAACAGGGATTTGGG GAATTATTTGAGAAAGCAAAACAAAACAATAACAATAGAAAAACT TCTAATGGTGATGACAGCCAAGATAGAAAGAGGACAGTTGTTGG AGGTTGCTGGATCCACTGGAGCAGGCAAGGTAGTTCTTTTGTTC TTCACTATTAAGAACTTAATTTGGTGCCCATGTCTCTTTTTTTTTC TAGTTTGTAGTGCTGGAAGGTATTTTTGGAGAAATTCTTACATGA GCATTAGGAGAATGTATGGGTGTAGTGTCTTGTATAATAGAAATT GTTCCACTGATAATTTACTCTAGTTTTTTTATTTCCTCATATTATTTT CAGTGGCTTTTTCTTCCACATCTTTATATTTTGCACCACATTCAAC ACTGTATCTTGCACATGGCGAGCATATGGTCGATGATGTAAGCT GCCCCAGACCCTGGGACCTCTGACCCCCGGCGACCCCTTGCAC TCTCCATGAGCCCTTTCCTTTTTTTTTCCTCAGGTCTAGGTGGGGG ATCTGGTGAGGGCAGAGGCAGTCTGCTGACATGCGGTGACGTG GAAGAGAATCCCGGCCCTTCTAGAAGCAAGGGCGAGGAGCTGT TCACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGT AAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGAT GCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCG GCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGAC CTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAG CAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCA GGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCC GCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCAT CGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTG GGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATAT CATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGA TCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCA CTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTG CCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAG ACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGT GACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAG CTGCAGAGCAGGGCCGACCCCAAGAAGAAGAGGAAGGTGGACC CCAAGAAGAAGAGGAAGGTGGACCCCAAGAAGAAGAGGAAGGT GTGA
3. CUTS mCherry Linker-T2A	ATGGTGAGCAAGGGCGAGGAGGATAACATGGCCATCATCAAGG AGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGG CCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCCTA CGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGTGG

CFTR_intron 8 CFTR_modified_exon9 UNC13A_modified_CE UNC13A_intron20 Linker-T2A EGFP 3XNLS	CCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGT ACGGCTCCAAGGCCTACGTGAAGCACCCCGCCGACATCCCCGA CTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGC GTGATGAACTTCGAGGACGGCGGCGTGGTGACCGTGACCCAGG ACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTG CGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGA AGACCATGGGCTGGGAGGCCTCCTCCGAGCGGATGTACCCCGA GGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCT GAAGGACGGCGGCCACTACGACGCTGAGGTCAAGACCACCTAC AAGGCCAAGAAGCCCGTGCAGCTGCCCGGCGCCTACAACGTCA ACATCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATC GTGGAACAGTACGAACGCGCCGAGGGCCGCCACTCCACCGGC GGCATGGACGAGCTGTACAAGGGAGGTGGGGGATCTGGTGAG GGCAGAGGCAGTCTGCTGACATGCGGTGACGTGGAAGAGAATC CCGGCCCTGGATCCGGCGAGGTGAGTATGGTACATAAAACAAG CATCTATTGAAAATATCTGACAAACTCATCTTTTATTTTTGATGTG TGTGTGTGTGTGTGTGTGTGTGTTTTTTTTTTAACAGGGATTTGGG GAATTATTTGAGAAAGCAAAACAAAACAATAACAATAGAAAAACT TCTAATGGTGATGACAGCACTGCCTGGGTTTCCTGGAAAGAACT CTTATCCCCAGGAACTAGTTTGTTGAATAAATGCTGGTGTGAATG TGAATGGATGATTGAACAGATGAATGAGTGTGATGAGTATGATAA AAGCATGGATGGAGAGATGGGTGAGTACATGGATGGATAGATG GATGAGTTGGTGGGTAGATTCGTGGCTAGATGGATGATGGATGG ATGGATGGATGGATGGATGGATGGACAGATGGATGGATATATGA TTGAACTATTGAAAGTATAGATGTATGGATGGGTGAATTTGGGG GTAATTGTTAGATGATGGATGTGGATGGTTGTGAGTGGCTGGTG GACAGACGAAAAATGGATGGGTGGATAAATTGATGGGTGGATGG ATGGTTGGTTGTATGAAAGAATGAATGATTGGGTAGATAAAAGAG TAGATGAATGAATTAATGAATAAACAGGCAGATGGATGATGTAAG CTGCCCCAGACCCTGGGACCTCTGACCCCCGGCGACCCCTTGC ACTCTCCATGAGCCCTTTCCTTTTTTTTCCTCAGGTCTAGGTGGG GGATCTGGTGAGGGCAGAGGCAGTCTGCTGACATGCGGTGACG TGGAAGAGAATCCCGGCCCTTCTAGAAGCAAGGGCGAGGAGCT GTTCACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGAC GTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGC GATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCAC CGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTG ACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGA AGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTC CAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGAC

	CCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGC ATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCC TGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTAT ATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAA GATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGAC CACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGC TGCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAA AGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTC GTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACA AGCTGCAGAGCAGGGCCGACCCCAAGAAGAAGAGGAAGGTGGA CCCCAAGAAGAAGAGGAAGGTGGACCCCAAGAAGAAGAGGAAG GTGTGA
4. Codon-opti mized TDP-43	ATGTCTGAATATATTCGGGTAACCGAAGATGAGAACGATGAACC AATCGAAATTCCAAGTGAAGACGACGGAACAGTCTTGCTCTCCA CAGTAACGGCGCAGTTTCCCGGTGCGTGCGGATTGAGATATCG CAACCCGGTGTCTCAGTGCATGCGAGGTGTAAGGTTGGTCGAA GGGATCCTGCACGCCCCCGATGCAGGCTGGGGTAACCTCGTGT ACGTGGTTAACTACCCGAAGGATAACAAGAGGAAAATGGATGAA ACCGATGCTAGTTCCGCCGTGAAGGTGAAACGAGCGGTTCAAAA AACTTCAGATCTTATAGTTCTGGGCCTCCCTTGGAAGACCACGG AACAGGATTTGAAAGAGTATTTTTCAACCTTTGGCGAGGTTCTCA TGGTGCAGGTTAAAAAAGATCTTAAAACCGGACACTCAAAAGGA TTTGGCTTCGTGAGGTTCACCGAGTACGAAACGCAGGTAAAAGT GATGTCACAGCGCCATATGATAGACGGACGGTGGTGCGATTGTA AGCTGCCCAATTCCAAGCAAAGCCAAGACGAGCCTCTGAGGAGT CGCAAAGTTTTCGTTGGACGCTGTACGGAAGACATGACTGAAGA TGAGCTTAGGGAGTTTTTTAGTCAGTATGGAGACGTTATGGATGT CTTTATCCCAAAACCGTTTCGGGCCTTCGCCTTTGTCACCTTTGC CGACGACCAGATCGCCCAGAGCTTGTGTGGCGAAGACCTGATA ATTAAAGGTATATCTGTCCATATCAGCAATGCGGAACCGAAACAC AACTCTAATCGACAGCTTGAGAGGTCAGGGAGGTTTGGGGGGTAA CCCGGGTGGATTCGGTAACCAAGGGGGGGTTCGGAAACTCCCGG GGCGGAGGAGCCGGATTGGGTAATAACCAGGGATCCAACATGG GCGGCGGGATGAACTTTGGTGCCTTCTCCATAAATCCTGCGATG ATGGCCGCAGCGCAAGCTGCCCTTCAGAGCTCCTGGGGTATGA TGGGGATGCTTGCTTCTCAGCAAAACCAATCCGGGCCCAGCGG CAACAATCAGAATCAAGGTAATATGCAGCGAGAACCGAACCAGG CCTTCGGAAGCGGGAATAATTCCTACTCAGGATCAAATAGTGGT GCAGCCATTGGCTGGGGCTCAGCGTCTAATGCAGGCTCAGGCA

	GTGGGTTTAATGGGGGCTTTGGGAGCTCCATGGACAGCAAGTCATCAGGCTGGGGGTAA