Title: REM sleep quality is associated with balanced tonic activity of the locus coeruleus during

wakefulness

Running head: REM sleep is linked to locus coeruleus activity

Nasrin Mortazavi, MSc,[1] Puneet Talwar, PhD,[1] Ekaterina Koshmanova, PhD, [1] Roya Sharifpour,

MSc, [1] Elise Beckers, MSc, [1,2] Alexandre Berger, PhD, [1,3,4] Islay Campbell, PhD, [1] Ilenia Paparella,

MSc, [1] Fermin Balda, MSc, [1] Ismael Dardour Hamzaoui, MSc, [1] Christian Berthomier, PhD, [5]

Christine Bastin, PhD, [1,6] Christophe Phillips, PhD,[1] Pierre Maquet, MD, PhD,[1,7] Fabienne Collette,

PhD, [1,4] Mikhail Zubkov, PhD, [1] Laurent Lamalle, PhD, [1] and Gilles Vandewalle, PhD [1*]

[1] GIGA-Institute, CRC-Human Imaging, University of Liège, Liège, Belgium.

[2] Faculty of Health, Medicine and Life Sciences, Mental Health and Neuroscience Research

Institute, Alzheimer Centre Limburg, Maastricht University; Maastricht, The Netherlands.

[3] Department of Clinical Neuroscience, Institute of Neuroscience (IoNS), Université Catholique

de Louvain (UCLouvain), Brussels, Belgium

[4] Synergia Medical SA, Mont-Saint-Guibert, Belgium

[5] Physip, Paris, France.

[6] PsyNCog, University of Liège, Liège, Belgium

[7] Neurology department, CHU de Liège, Belgium

*Corresponding author: Gilles Vandewalle, GIGA-Cyclotron Research Centre-Human Imaging,

Bâtiment B30, 8 Allée du Six Août, University of Liège-Sart Tilman, 4000 Liège, Belgium.

27
28

Abstract

Objective

Animal studies established that the locus coeruleus (LC) plays important roles in sleep and wakefulness regulation. Whether it contributes to sleep variability in humans is not yet established. Here, we investigated if the *in vivo* activity of the LC is related to the variability in the quality of Rapid Eye Movement (REM) sleep.

Methods

We assessed the LC activity of 34 healthy younger (~22y) and 18 older (~61y) individuals engaged in bottom-up and top-down cognitive tasks using 7-Tesla functional Magnetic Resonance Imaging (fMRI). We further recorded their sleep electroencephalogram (EEG) to evaluate associations between LC fMRI measures and REM sleep EEG metrics.

Results

Theta oscillation energy during REM sleep was positively associated with LC response in the top-down task. In contrast, REM sleep theta energy was negatively associated with LC activity in older individuals during the bottom-up task. Importantly, sigma oscillations power immediately preceding a REM sleep episode was positively associated with LC activity in the top-down task.

Interpretation

LC activity during wakefulness was related to REM sleep intensity and to a transient EEG change preceding REM sleep, a feature causally related to LC activity in animal studies. The associations depend on the cognitive task, suggesting that a balanced level of LC tonic activity during wakefulness is required for optimal expression of REM sleep. The findings may have implications for the high prevalence of sleep complaints reported in aging and for disorders such as insomnia, Alzheimer's, and Parkinson's disease, for which the LC may play pivotal roles through sleep.

Keywords: Locus coeruleus, sleep, 7 Tesla functional magnetic resonance imaging, aging

Introduction

Up to 35% of the general population report unsatisfying sleep[1] and one out of two adults aged over 50y complains about sleep or daytime sleepiness.[2] Insomnia is the obvious extreme form of these complaints and constitutes the second most prevalent psychiatric disorder.[3] Beyond the behaviors that do not favor appropriate sleep, some people are more vulnerable to poor sleep or more at risk of developing insomnia.[4,5] The biological origin of sleep variability is however not fully established.[6]

The locus coeruleus (LC), a small 1.5cm-long spaghetti-shaped nucleus of the dorsal pons,[7] produces norepinephrine (NE) as a part of the ascending arousal system and projects to the neocortex, hippocampus, amygdala, thalamus, and cerebellum.[8] It plays a crucial role in the control of behavioral states, including alertness and attention,[9] as well as in the regulation of sleep.[6] LC-NE tone must decrease to allow sleep onset, while LC activity dynamically changes during sleep to shape the alternation between Rapid Eye Movement (REM) and non-REM sleep and some of the microstructure elements of sleep.[10,11] A lot of our understanding of the role of the LC in sleep regulation arises from experiments conducted in rodents. Therefore translation to diurnal human beings, with a more developed cortex, is not straightforward[6]. In addition, imaging the LC *in vivo* remains difficult because of its deep location and small size, as well as because of the difficulty to image human sleeping brains. Recent advances in neuroimaging techniques have lifted part of these limitations.[12] For instance, degeneration of the LC was recently suggested to contribute to alteration in rest-activity patterns.[12] We further reported that a higher response of the LC to a salience detection task during wakefulness, which presumably reflects in part LC activity during sleep, was associated with lower intensity of REM

sleep in healthy older individuals.[13] REM sleep intensity does not however solely depend on LC

activity, and no association with sleep features shown to be under the direct control of the LC in

animal were reported in the study.

LC neurons can follow phasic and tonic activity patterns, which lead to distinct release of NE.[14]

Phasic activity typically occurs in response to transient salient stimulation, while the tonic mode

is mostly required for sustained processes. The dynamics of the tonic mode is considered to

follow an inverted U-shape where low LC tonic discharge is associated with low arousal and poor

performance, moderate LC tonic activity allowing pronounced phasic activity leads to optimal

arousal and performance, and high tonic activity hinders phasic activity and leads to anxiety-like

behavior and poorer performance .[15] Likewise, during sleep the LC undergoes periods of higher

or lower tonic activity.[10,11]

How the balance between phasic and tonic modes of activity of the LC during wakefulness may

be related to REM sleep is not known. In the present study, we used Ultra-High-Field (UHF) 7

Tesla (7T) Magnetic Resonance Imaging (MRI) to extract *in vivo* the LC activity in healthy younger

and older adults during two cognitive tasks that rely on different tonic tones of the LC. We

related the LC activity during both tasks to electrophysiology metrics of REM sleep. We

hypothesized that the LC activity in both tasks would be mainly associated with REM sleep

intensity. Moreover, we expected that the associations would differ between two tasks, and

would be more pronounced in older individuals. To test for associations with sleep features

under direct control of the LC, we further tested the association between LC activity and the

power of the sigma band of the EEG that preceded REM sleep episodes, because it is causally

related to NE tone in rodents[16].

Methods

This study was approved by the faculty-hospital ethics committee of ULiège. All participants provided their written informed consent and received financial compensation. The study is part of a larger project that has led to previous publications.[13,17] Part of the results dealing with one cognitive task were reported in one of these publications[13] and are included for comparisons purposes and to conduct additional analyses. Most of the methods were described in details in[13]. More details are also available in the **Supplementary Methods**.

Participants

Fifty-two healthy participants including 34 healthy young (22.2±3.1y, 28 women) and 18 late middle-aged (60.9±5.4y, 14 women) individuals, completed the study **(Table 1)**. The exclusion criteria were as follows: history of major neurologic/psychiatric diseases or stroke; recent history of depression/anxiety; sleep disorders; medication affecting the central nervous system; smoking, excessive alcohol (>14 units/week) or caffeine (>5 cups/day) consumption; night shift work in the past 6 months; BMI ≤18 and ≥29 (for older individuals) and ≥25 (for younger individuals). All older participants had to show normal performance on the Mattis Dementia Rating Scale (score > 130/144).[18]

Protocol

Participants' sleep was recorded in the lab twice. First, participants completed a night of sleep under polysomnography (called habituation night) to screen and exclude for sleep abnormalities (apnea hourly index and periodic leg movement >15; no parasomnia or REM behavioral disorder). All participants further completed a whole-brain structural MRI (sMRI) acquisition and

a specific acquisition centered on the LC. Participants were then requested to sleep regularly for

7 days before the baseline night (±30min from their sleep schedule) based on their preferred

schedule (compliance was verified using sleep diaries and wrist actigraphy - Actiwatch and AX3,

AXIVITY LTD, Newcastle, UK). For the second night (called baseline night), participants first

remained awake for 3h under dim light (<10 lux) then their habitual sleep was recorded in

darkness (N7000 amplifier, EMBLA, Natus, Middleton, WI). Approximately 3h after wake-up time

under dim light (<10 lux), participants completed a functional MRI (fMRI) session that included 3

tasks (**Figure 1A**). This paper is centered on the analyses of the first two tasks, the perceptual

rivalry and auditory salience detection tasks.

Younger participants completed the fMRI session immediately following the baseline night but

older participants were initially part of a different study[19,20] and completed the MRI procedures

in addition to their initial engagement, which included the baseline night recording.

Consequently, the baseline nights of sleep and MRI sessions were completed about 1.25y apart

(mean±SD: 15.5±5.3 months). Prior to the fMRI session, older participants slept regularly for 1

week and were maintained in dim light (<10 lux) for 45 minutes.

Sleep EEG metrics

Sleep was staged in 30s-epochs using an automatic algorithm (ASEEGA, PHYSIP, Paris)[21] to

provide REM sleep percentage and total sleep time (TST). Averaged power was computed per

30-min bin, adjusting for the proportion of rejected data and it was subsequently aggregated in

a sum for REM[22] to provide REM theta energy (overnight cumulated 4.25-8Hz power). Sigma

power (12.25-16Hz) was computed during the 1-min preceding each REM episode, as the

weighted sum of 4s artefact-free window (2s overlap per 30s epoch).

Cognitive tasks

Visual perceptual rivalry task – <u>high-tonic LC</u>. The task consisted of watching a 3D Necker cube,

which can be perceived in two different orientations (**Figure 1B**), for 10 blocks of 1min

separated by 10s of screen-center cross fixation (~12min). Participants were instructed to report

switches between the two percepts through a button press. Given the top-down nature of the

task, we considered that if the task recruited the LC, it would trigger a relatively lower phasic

response over a relatively larger tonic tone (high-tonic LC).

Auditory salience detection task – <u>low-tonic LC</u>. The task consisted of an oddball paradigm

requiring reports on the perception of rare deviant target tones (1,000Hz, 100ms, 20% of tones)

that were pseudo-randomly interleaved within a stream of standard stimuli (500Hz, 100ms)

through a button press (**Figure 1C**). The task included 270 stimuli (54 targets; ~10min). Given the

bottom-up nature of the task, we considered that if the task recruited the LC, it would trigger a

relatively larger phasic response over a relatively lower tonic tone (low-tonic LC).

Pupil size was recorded during both cognitive tasks in a subset of participants (N=27 in the

salience detection task; N=32 in the perceptual rivalry task) (Eyelink-1000, SR Research,

Osgoode, ON, Canada; sampling rate: 1000Hz).

MRI data acquisition and *preprocessing*

MRI data were acquired using a MAGNETOM Terra 7T MRI system (Siemens Healthineers,

Erlangen, Germany). FMRI and sMRI data were preprocessed using SPM12, ANTs and SynthStrip

brain extraction tool[23], as fully described previously[13]. The preprocessed data were resampled to

a 1mm^3 resolution. Individual statistical analyses included one regressor of interest consisting of

a switch in perception (perceptual rivalry) or the occurrence of a target deviant tone (salience

detection) modeled as an event (convolved with the canonical hemodynamic response function

- HRF). Participant movement, respiration, and heart rate were used as covariates of no interest

(physiological data of 4 volunteers were not available).

Individual LC masks were manually delineated by 2 experts based on LC-specific images (as in[13])

to extract LC activity estimates (i.e., mean value over the LC mask) during both tasks and to

compute a LC probabilistic map in the group space for visualization. The T1 structural whole-

brain image was used to extract individual total intracranial volume (TIV) using the CAT12

software.[24]

Statistics

Statistical analyses were performed in SAS 9.4 (SAS Institute, NC, USA) and consisted of

Generalized Linear Mixed Models (GLMM) with 4 main sleep features of interest (REM sleep

latency, REM sleep percentage, REM Theta energy, sigma power prior to each REM sleep

episode) as the dependent variables (with adjusted distribution), LC activity as an independent

variable, as well as age group (younger and older individuals), sex, TST and TIV. Outliers among

all variables lying beyond four times the standard deviation were removed from the analysis

(maximum 2 data points were removed). The final number of individuals included in each

analysis is reported in **table 2** and **table 3**. Partial R^2 *(R^{2*})* values were computed as previously

described.[13] We used Pearson correlations to assess the association between transient pupil

dilation and estimates of LC activity as well as for visualization of the GLMM outputs. The

analyses included 4 main dependent variables of interest: Benjamini and Hochberg False

Discovery Rate (FDR) correction considering 4 independent tests was used to test for significant

associations [p<.0125 (for rank 1/4); p<.025 (for rank 2/4); p<.0375 (for rank 3/4); p<.05 (for

rank 4/4)]. All models included an interaction term between LC activity and age group. If the

Bayesian Index Criterion (BIC) of the model without the interaction was better (i.e. lower) and

the interaction term was not significant it was removed from the model.

Importantly, statistical analyses are slightly modified compared to[13] with the inclusion of TIV as a

covariate (rather than BMI) to take into account non-specific variation in EEG power

computations. Statistical outcomes related to the auditory salience detection task may therefore

slightly vary compared with our previous publication.[13]

We computed a prior sensitivity analysis to get an indication of the minimum detectable effect

size in our main analyses given our sample size. According to G*Power 3 (version 3.1.9.4),[25]

taking into account a power of .8, an error rate α=.05, and a sample size of 52, we could detect

medium effect sizes r>.39 (2-sided; CI:.13–.6; R^2>.15, CI:.02–.36) within a linear multiple-

regression framework including 2 tested predictor (LC activity, age group) and 2/3 covariates

(sex, TST, TIV where relevant).

Figure 1. Overview of the study protocol.

(A) After screening, participants completed an in-lab screening and habituation night under polysomnography to minimize the effect of the novel environment for the subsequent baseline night and to exclude volunteers with sleep disorders. They further completed a structural 7T MRI session including a whole-brain structural MRI and a LC-specific sequence. The latter was used to create individual LC masks in each participant's brain space. After 7 nights of regular sleep-wake time at home, which was confirmed by actigraphy data and/or sleep-wake diary, participants came to the lab three hours before their sleep time and were maintained in dim light (<10 lux) until sleep time. Participants' habitual baseline sleep were recorded overnight in-lab under EEG to extract our main sleep features of interest. All participants underwent an fMRI session approximately 3h after wake-up time (following ≥45min in dim light - < 10lux), during which they completed the visual perceptual rivalry task and the auditory salience detection task.

(B) The Visual perceptual rivalry task consisted of watching a 3D Necker cube, which can be perceived in two different orientations (blue arrow), for 10 blocks of 1min separated by 10s of screen-center cross fixation (total duration ~12min). Participants reported switches in perception through a button press. We considered that task would involve a relatively lower phasic response over a relatively larger tonic tone (high-tonic LC).

(C) The auditory salience detection task consisted of an oddball paradigm requiring button-press reports on the perception of rare deviant target tones (20% occurrence) within a stream of frequent tones (total duration ~10min). We considered that task would involve a relatively larger phasic response over a lower relatively tonic tone (low-tonic LC).

Results

Fifty-two healthy individuals aged 18 to 31y (N=34; 28 women) and 50 to 70y (N=18; 14 women) completed an fMRI protocol consisting of a top-down visual perceptual rivalry task[26], which we considered could trigger a lower phasic response of the LC over a relatively larger tonic tone (high-tonic LC), and a bottom-up auditory salience detection task[27], which we considered could trigger a larger phasic response of the LC over a relatively lower tonic tone (low-tonic LC task). Both tasks successfully recruited the left LC when inspecting the whole-brain statistical analyses over the entire sample (**Figure 2A and B**). The recruitment of the LC was further supported by the fact that, in a subset of participants, we successfully detected significant transient pupil dilation around the event of interest (**Figure 2C, D, E and G**). The LC is indeed considered to drive part of the transient pupil dilation associated with the detection of salient stimuli or with changes in perception in a perceptual rivalry context.[27,28] Transient pupil dilation during the salience detection task was further significantly correlated to the activity estimate of the left LC (Pearson's r=.4, **p=.04**; **Figure 2H**). Since pupil responses are associated with phasic responses of LC,[29] the absence of significant correlation in the perceptual rivalry task (**Figure 2F**, Pearson's r=-.20, p=.29) may support our hypothesis that this task triggers more tonic LC activity. All these results prompted us to extract the activity of the left LC in all individuals and related it to canonical characteristics of their REM sleep estimated based on a night of sleep in the lab under EEG.

We first found that LC responses during both tasks were not correlated (**Figure 3A**) (Pearson's r=.08, p=.54), indicating that the activity of LC in one task did not predict its activity in the other task and supporting that both tasks probe different aspects of LC activity.

Considering the perceptual rivalry task, we found a significant positive association between REM

theta energy and high-tonic LC activity (t=3.38; **p= .001**; **Figure 3B**) on top of the main effect of

TST, while the other covariates were not significant (**Table 2**). As previously reported,[13] we then

found a significant association between REM theta energy and the interaction between low-

tonic LC activity during the salience detection task and age group (t=2.68; **p=.01; Table 3**) that

came on top of a main effect of LC activity (t=-2.78; **p=.008**) (**Figure 3C**) while the other

covariates were not significant (**Table 3**). Post hoc contrasts revealed that higher REM theta

energy was associated with a lower response of the LC in older (t= -2.78; **p=.008**) but not in

younger (t=.33; p=.74) individuals, indicating that the main effect of LC activity was mainly

driven by the older participants.

For both low-tonic and high-tonic LC activity the inclusion of REM sleep duration and/or REM

sleep percentage instead of or on top of TST did not modify the statistical outputs of the models

(data not shown). Furthermore, REM theta energy associations with the activity of the LC during

both tasks remain similar when including them simultaneously in the same GLMM, i.e. a

significant main effect of high-tonic LC activity ($F_{(1,41)}$=9.52, **p=.0036**) and a weak trend for

interaction between low-tonic LC activity and age: $F_{(1,41)}$=2.72, p=.1), supporting that they

explain different parts of the variance in REM theta energy.

No association was found when considering the other macroscopic REM sleep metrics (REM

sleep onset latency, REM sleep percentage) (**Suppl. Figure S1, Table 2 and 3**). Further analyses

indicated that the associations with REM theta energy were specific to this particular band as no

significant association was uncovered between the energy of the other spectral bands of the

EEG during REM sleep (except for a significant main effect of low tonic LC activity [t= -2.33;

p=.025) and of an age-group-by-low tonic LC activity interaction (t= 2.14; p=.037) when using

REM alpha energy as the dependent variable - **Suppl. Figure S2, Suppl. Table S1 and S2**]. The

significant associations we detected were also specific to REM sleep as NREM slow wave energy

(SWE), i.e. the cumulated overnight power over the delta band (.5-4Hz) typical of NREM sleep,

was not significantly associated with either low-tonic and high-tonic LC activity (**Suppl. Figure S2,**

Suppl. Table S1 and S2).

In the next steps, we turned towards sigma power prior to REM sleep as it has been causally

related to LC activity during sleep in animal models.[16] Statistical analyses yielded a significant

positive main effect of high-tonic LC activity during the perceptual rivalry task (t=2.36; **p=0.022**;

Table 2; Figure 3D) and no significant interaction between LC activity and age group. In contrast,

no significant association with low-tonic LC activity during the salience detection task (p=0.17)

and no significant interaction between LC activity and age group were detected (**Table 3; Figure**

3E). In both analyses, we did not detect significant effects of the other covariates (except for

main effect of sex in the salience detection task). Finally, when including the activity of both

tasks in the same GLMM with sigma power prior to REM as the dependent variable, the analysis

yielded a significant main effect of high-tonic LC activity ($F_{(1,42)}$=7.62,**p=.0085**) as well as a

trending interaction between low-tonic LC activity and age ($F_{(1,42)}$=3.77, p=.058). As for REM

theta energy, this result suggests that both types of LC activity estimates explain different parts

of the variance in sigma power prior to REM.

For completeness, the final steps of our analyses consisted of exploratory analyses for

associations between the activity of the LC estimated during both tasks and other REM sleep

metrics, i.e. REM sleep duration, REM bouts duration, and number of arousals during REM sleep.

These led to no significant associations (**Suppl. Figure S3, Suppl. Table S1 and S2**).

Table 1. Characteristics of the study sample

	Young individuals (n = 34)				Late middle-aged individuals (n = 18)				P-value
	Mean	SD	Min	Max	Mean	SD	Min	Max	
Age (years)	22.20	3.17	18.00	29.00	60.88	5.41	53.00	70.00	**<.0001**
BMI (kg/m2)	21.90	2.74	17.20	28.40	24.75	3.40	19.40	30.90	**0.0047**
Education (years)	14.46	2.31	12.00	20.00	14.61	2.70	9.00	19.00	0.85
Depression level (BDI)	6.64	4.17	0	20.00	5.33	4.01	0	14.00	0.27
Anxiety level (BAI)	4.00	2.92	0	11.00	3.05	3.18	0	9.00	0.30
TIV	1402.1	136.6	1203.3	1946.7	1420.5	125.5	1167.6	1600.1	0.627
Daytime sleepiness (ESS)	6.93	3.90	0	14.00	5.77	3.76	0	13.00	0.30
Sex (F–M)	28 F – 6 M				12 F – 6 M				0.20
Habitual Subjective sleep quality (PSQI)	4.79	2.18	1.00	10.00	3.83	2.22	0	8.00	0.14
TST	454.80	30.97	379.50	499.50	394.63	48.16	282.50	495.50	**<.0001**
Theta power in REMS	7281.5	5061.7	1162.4	22346.4	4789.5	3508.2	765.3	13051.8	**0.0449**
REMS percentage	26.63	4.41	19.93	36.45	21.52	5.50	13.11	35.39	**0.002**
REMS latency	103.0	47.834	0	213.5	81.944	56.806	38.000	295.5	0.190
Sigma power prior to REMS	15.7615	8.8957	5.0656	44.356	16.0544	6.795	3.654	27.856	0.896
SWE	175531	171295	14259.2	555453	126388	109290	23293.3	470116	0.2146

The p-values shown in the table correspond to two-sample t-tests except for sex that were compared using a Chi-square test.
BMI: Body Mass Index; BDI: Beck Depression Inventory; BAI: Beck Anxiety Inventory; TIV: Total Intracranial Volume; PSQI:
Pittsburgh Sleep Quality Index; ESS: Epworth Sleepiness Scale; TST: Total Sleep Time; SWE: slow wave energy; REMS: rapid eye
movement sleep. See supplementary method for references of the questionnaires.

Table 2. Associations between REM sleep metrics and LC activity estimated via <u>the visual perceptual</u>
<u>rivalry (high-tonic) task.</u>

Sleep metric (dependent variable)	LC activity	Age group	Sex	TIV	Total sleep time
REM Theta energy **(N=51)**	$F_{(1,45)}=11.39$ **P=0.001*** **R^{2*}=0.201**	$F_{(1,45)}=0.14$ P=0.709	$F_{(1,45)}=0.22$ P=0.645	$F_{(1,45)}=0.08$ P=0.781	$F_{(1,45)}=5.15$ **P=0.028** **R^{2*}=0.102**
Sigma power prior to REM (N=52)	$F_{(1,46)}=5.55$ **P=0.023*** **R^{2*}=0.107**	$F_{(1,46)}=0.60$ P=0.442	$F_{(1,46)}=3.64$ P=0.062	$F_{(1,46)}=2.18$ P=0.147	
REMS percentage (N=52)	$F_{(1,47)}=2.05$ P=0.159	$F_{(1,47)}=13.27$ **P=0.0007** **R^{2*}=0.220**	$F_{(1,47)}=1.66$ P=0.204	$F_{(1,47)}=3.15$ P=0.082	
REMS latency (N=52)	$F_{(1,46)}=0.10$ P= 0.750	$F_{(1,46)}=2.01$ P= 0.163	$F_{(1,46)}=1.27$ P= 0.265	$F_{(1,46)}=0.27$ P= 0.602	$F_{(1,46)}=0.74$ P= 0.392

Prior to the analysis, we removed the outliers among all variables by excluding the samples lying beyond four times
the standard deviation (the final number of individuals included in each analysis is reported below each dependent
variable).

* Significant following Hochberg False Discovery Rate (FDR) correction for 4 independent tests (p<.025)

In all models the interaction between LC activity and age group was not significant. Goodness of fit metric (BIC)
indicated that the interaction term should be removed.

LC: locus coeruleus; TIV: total intracranial volume; REM: rapid eye movement; REMS: rapid eye movement sleep.

Table 3. Associations between REM sleep metrics and LC activity estimated via <u>the auditory salience detection (low-tonic) task</u>.

Sleep metric (dependent variable)	LC activity	Age group	LC activity*age group	Sex	TIV	Total sleep time
REM Theta energy (N=51)	$F_{(1,44)}=6.74$ **P=0.012*** **R^{2*}=0.132**	$F_{(1,44)}=0.54$ P=0.468	$F_{(1,44)}=7.19$ **P=0.010*** **R^{2*}=0.140**	$F_{(1,44)}=0.14$ P=0.707	$F_{(1,44)}=0.10$ P=0.750	$F_{(1,44)}=3.48$ P=0.068
Sigma power prior to REM (N=50)	$F_{(1,44)}=1.90$ P=0.175	$F_{(1,44)}=0.26$ P=0.611	$F_{(1,44)}=2.35$ P=0.132	$F_{(1,44)}=8.01$ **P=0.007** **R^{2*}=0.154**	$F_{(1,44)}=3.11$ P=0.084	
REMS percentage (N=52)	$F_{(1,46)}=1.38$ P=0.245	$F_{(1,46)}=15.60$ **P=0.0003** **R^{2*}=0.253**	$F_{(1,46)}=2.07$ P=0.157	$F_{(1,46)}=0.23$ P=0.635	$F_{(1,46)}=5.56$ **P=0.022** **R^{2*}=0.107**	
REMS latency (N=52)	$F_{(1,45)}=2.88$ P= 0.096	$F_{(1,45)}=0.75$ P= 0.389	$F_{(1,45)}=1.40$ P=0.242	$F_{(1,45)}=2.07$ P= 0.157	$F_{(1,45)}=0.88$ P= 0.351	$F_{(1,45)}=0.96$ P= 0.331

Prior to the analysis, we removed the outliers among all variables by excluding the samples lying beyond four times the standard deviation (the final number of individuals included in each analysis is reported below each dependent variable).

* Significant following Hochberg False Discovery Rate (FDR) correction for 4 independent tests (p<.025)

In all models, the results without the interaction between LC activity and age group were not significant. Goodness of fit metric (BIC) indicated that the interaction term provides a better fitness.

LC: locus coeruleus; TIV: total intracranial volume; REM: rapid eye movement; REMS: rapid eye movement sleep.

Figure 2. (A) Whole-brain and LC responses to the perceptual switches during the visual perceptual rivalry task. Sagittal, coronal, and axial views [MNI coordinates: (−4 −37 -21 mm)]. The images at the top show the whole-brain results using significance for a threshold of p< 0.05 FDR-corrected (t > 2.16) over the group average brain structural image coregistered to the MNI space. Insets at the bottom show the LC probabilistic template (blue) created based on individual LC masks and the significant activation detected within this mask (dark grey). (B) Whole-brain and LC responses to the target sound during the auditory salience detection task. Sagittal, coronal, and axial views [MNI coordinates: (−4 −34 -21 mm)]. The images at the top show the whole-brain results using significance for a threshold of p< 0.05 FDR-corrected (t > 2.33) over the group average brain structural image coregistered to the MNI space. Insets at the bottom show the LC probabilistic template (blue) and the significant activation detected within this mask (dark grey). The legend shows the t-values associated with color maps. (C) A representative example of the pupil dilation in response to the perceptual switches during the visual perceptual rivalry task in one participant. (D) A representative example of the pupil dilation in response to the standard and target sounds during the auditory salience detection task in one subject. (E) Minimum and maximum pupil size before and after the perceptual switches during the visual perceptual rivalry task. Maximum pupil size was significantly higher than the minimum (N=32, p < 0.0001). (F) Association between LC activity during visual perceptual rivalry task and mean ratio of pupil size in response to the perceptual switches. The association was not significant (Pearson's correlation r=-0.201, P=0.29). (G) Mean ratio of pupil size in response to standard sound and target sound during the auditory salience detection task. The mean ratio of pupil size was computed as the change in the pupil diameter from before to after the auditory stimulus presentation. The change in pupil size was significantly higher for target versus standard sound (N=27, p < 0.0001). (H) Association between LC activity during auditory salience detection task and mean ratio of pupil size in response to the target sound. We found a significant positive Pearson's correlation (r=0.403, P=0.040).

Figure 3. (A) LC activity estimates during the visual perceptual rivalry task and auditory salience detection task were not significantly correlated (p=0.54). (B) Association between REM theta energy and the LC activity estimates during the visual perceptual rivalry task. The GLMM yielded a significant main effect of LC activity (p=0.0015). (C) Association between REM theta energy and the LC activity estimates during the auditory salience detection task. The GLMM yielded a significant age group by LC activity interaction (p=0.0103), and post hoc analyses led to a significant association for the older (p=0.008) but not the young group (p=0.75). (D) Association between sigma power prior to REM sleep and the LC activity estimates during the visual perceptual rivalry task. The GLMM yielded a significant main effect of LC activity (p=0.026). (E) Association between sigma power prior to REM sleep and the LC activity estimates during the auditory salience detection task. The GLMM showed neither a significant main effect of LC activity (p=0.859) nor a significant age group by LC activity interaction (p=0.132). Simple regression lines are used for a visual display and do not substitute the GLMM outputs. The black line represents the regression irrespective of age groups (young + old, n = 52). Solid and dashed regression lines represent significant and non-significant outputs of the GLMM, respectively.

Discussion

Whether the variability in the functioning of the human LC may contribute to the interindividual

variability in sleep is not established. In the present cross-sectional study, we show that the

activity of the LC, assessed during wakefulness through 2 distinct cognitive tasks, is related to

the variability of REM sleep intensity, as reflected by the energy of the theta band of the EEG.

We find that the associations depend on the task considered - presumably because they do not

rely on the same balance between phasic and tonic LC activity, and on age - when considering LC

activity during the low-tonic task. In addition, we find that the amount of sigma band oscillations

immediately preceding REM sleep is associated with the activity of the LC assessed during the

high-tonic task. These findings add to the existing view that LC function is essential for sleep

regulation that was mainly brought by animal experiments. We show that the *in vivo* variability

in the activity of the human LC may govern part of the variability in sleep quality assessed using

electrophysiology. The findings may have implications for the neuropathology of several brain

disorders such as insomnia, Alzheimer's disease (AD), and Parkinson's disease (PD) for which the

LC may play a pivotal role through sleep.

LC cells can discharge action potentials in both tonic and phasic modes during wakefulness. The

tonic mode consists of irregular but constant baseline activity (1-6 spikes/s), while phasic activity

is characterized by short (<300ms) bursts of high-frequency activity followed by a long period of

sustained inhibition of spontaneous activity.[30] A balanced tonic activity is required for an optimal

arousal level allowing the expression of phasic activity in response to salient stimuli or

behavioral changes.[15] While the LC was considered for long as a quiet region during sleep, recent

animal studies revealed that the activity of the LC varies importantly during NREM sleep and

may even transiently reach activity levels similar to wakefulness.[10] NE level, driven by LC neuron

activity, fluctuates during NREM sleep between periods of higher and lower NE tone following

an infraslow rhythm of ~50s.[10] This periodic decrease in NE tone must reach levels that are low

enough to open the gate for REM sleep, which is considered to require sustained NE-free

periods to allow, for instance, synaptic pruning.[4] The periodic decrease of the LC activity is

reflected in periodic increases in the expression of sleep spindles and in sigma power such that

each REM episode is preceded by a pronounced transient increase in sigma power[31] that is well

known to somnologists. The overall picture emerging from this recent literature is that LC

activity shapes the macro- and microstructure of sleep and that inappropriate fluctuations in LC

tonic activity can disturb sleep and in particular REM sleep.[4,10]

We find that a higher phasic response of the LC during the top-down perceptual rivalry task,

arguably characterized by a relatively high tonic activity of the LC, is associated with a larger

expression of theta oscillations during REM sleep and of sigma power prior to REM sleep

episodes. We interpret this finding as the reflection of a more appropriate lower tonic activity of

the LC which allows a better expression of its phasic response during the task. In contrast, if

tonic activity during the task is too high, it partly masks the phasic response of the LC. We detect

this during wakefulness but surmise that it constitutes a trait such that the more appropriate

tonic activity of the LC present during wakefulness would also be present during sleep. This

would mean that, in the case of a low tonic LC activity allowing for a large dynamic range of

phasic LC oscillation, NE tone reaches a lower level prior to a REM sleep episode, allowing a

large expression of sigma oscillation and the initiation of a REM sleep episode and complete

silencing during REM. This would in turn enhance the expression (intensity) of REM theta

energy, the most typical oscillatory mode of REM sleep. Theta activity during REM sleep is

considered to reflect hippocampus activity and plays key roles in memory consolidation during

REM sleep.[32] The putative more appropriate tonic activity of the LC we report could therefore

reflect a better expression of REM sleep functions.

In contrast, we find that a larger expression of the phasic response of the LC during the bottom-

up salience detection task, which is arguably underlined by a lower tonic activity of the LC, is

associated with lower expression of theta oscillations during REM sleep, particularly in older

individuals. We link our results to the recent report in rodents that the so-called startle effect,

which is triggered by abrupt sensory changes, is attenuated under higher vs. lower arousal levels

as well as when LC activity is induced prior to the salient stimulation.[33] This was in line with the

established observation that elevated tonic firing of the LC-NE neurons makes the LC sensory-

evoked response less pronounced.[30] This would mean that in individuals with lower tonic activity

of the LC during wakefulness, the dynamic range of the phasic response of the LC is potentially

large. This trait-like phasic LC activity could become maladaptive and during sleep, perturb the

activity of limbic areas, thereby resulting in a reduction in REM theta oscillations. This process

becomes particularly manifest in aging which is characterized by increased sleep fragility and

degradation of sleep quality.[34]

Overall, we interpret our findings as a reflection of the requirement for a balanced tonic LC

activity for optimal REM sleep. This notion is supported, in our view, by the fact that our

statistical outputs remain similar (either significant or as weak trends) when including

concomitantly the activity of the LC as assessed in a low-tonic and high-tonic context. Our

findings may further be related to the diversity in neuron types within the LC that is emerging in

the literature, for instance, if both tasks were recruiting different proportions of LC neurons co-

expressing or not dopamine or projecting to different brain territories[6] and were therefore

characterizing the activity of these different neurons during sleep.

We stress that, although plausible, our interpretations remain putative and that measures of LC

activity during sleep are required to confirm our hypotheses. Future studies could also assess

the connectivity of the LC with other brain structures in humans, as sleep and wakefulness

regulation relies on a network of subcortical nuclei rather than on the LC alone.[35] Furthermore,

while we interpret our finding in "LC centric" manner, i.e. the activity of the LC during

wakefulness reflects its activity during sleep, it may also be that the activity of the LC during

wakefulness shapes the activity of other brain areas during sleep. Sleep oscillations depend

indeed in prior wakefulness in a regional use-dependent manner[36] such that the impact of NE

tone and LC activity on other regions, and particularly over cortical regions, may be reflected in a

larger or lower expression of theta oscillation during REM.

Beyond the healthy population, our findings are in line with the view that inappropriate LC

activity during both wakefulness and sleep can contribute to insomnia disorder.[4] Insomnia is

characterized by a hyperarousal state that is presumably due in part to inappropriately high

tonic activity of the LC during wakefulness, contributing to delayed sleep onset, and during

sleep, contributing to early and/or multiple awakenings. This would also lead to disturbed REM

sleep and unresolved emotional distress.[4] Our participants being healthy and devoid of sleep

disorders, our finding could reflect the continuum between normal sleep and insomnia. They

could also be indicative of the neuropathological mechanisms eventually leading to insomnia,

i.e. a prolonged inappropriate LC tonic activity and REM sleep disturbance are consequences or

contribute to the progressive changes leading to insomnia or maintaining the disorder. The LC is

also central to several neurodegenerative diseases characterized by preclinical alteration in

sleep. The LC is indeed among the first sites showing abnormal tau and alpha-synuclein

aggregates, hallmarks of AD and PD, respectively.[37,38] Abnormal tau inclusions can even be

detected post-mortem as early as during the second decade of life[37] and were suggested to be

associated with larger cortical excitability in late middle-aged healthy individuals.[39] In addition,

post-mortem degeneration of the human LC was recently associated with alteration in the rest-

activity cycle over the ~7y preceding death, particularly in those with higher cortical AD

pathology.[40] Future research could try to link our findings to assessments of tau and alpha-

synuclein burden within the LC, though the level of protein aggregate would go undetected *in*

vivo in young individuals, and would remain challenging to assess over such a deep and small

nuclei in older individuals.

As previously mentioned,[13] although our study provides new insights into the associations

between LC activity and sleep variability in humans, it bears limitations. The baseline night of

sleep was followed by the fMRI acquisition the next day in young individuals, while in the older

group, there was a ~1y gap between the baseline night and the fMRI session. Although sleep

changes over the lifespan,[41] it is stable over a short life period (e.g. a few years).[42] We therefore

consider that this important limitation is unlikely to fully explain our results, particularly given

the fact that no age-related differences were found for the high-tonic perceptual rivalry task.

Moreover, our sample was primarily composed of women. Although it was taken into account in

our statistical analyses, this limits the generalizability of our results. In addition, although it

represents a considerable data collection effort, the size of our sample remains modest,

particularly for the older subsample. Replication in a larger sample is therefore warranted.

Future studies could also apply individually tailored HRF to assess LC response. We used the

canonical HRF to model activity over the entire brain to model average LC response, while

individual LC responses can differ across individuals.[43]

In summary, we provide original evidence that, aside from its integrity, the functioning of the LC

shapes part of the quality of sleep. Our data suggest that an optimal level of LC tonic activity is

required for an optimal expression of REM sleep and of its functions. This reinforces the view

that the LC-NE is promising for interventions aiming at improving sleep quality, including

preventing and/or delaying brain disorders. Our findings suggest that these interventions could

aim at restoring optimal LC functioning during sleep as well as during wakefulness.

Acknowledgments

This work was conducted at the GIGA-In Vivo Imaging platform of ULiège, Belgium. We thank N.

Beliy, E. Lambot, C. Hagelstein, S. Laloux, A. Claes, C. Degueldre, B. Herbillon, G. Hammad, P.

Hawotte, B. Lauricella, T. Pontus, T. Gendron, J. Read, A. Robert and A. Lesoinne for their help in

the different stages of the study. This work was supported by Fonds National de la Recherche

Scientifique (FRS-FNRS, T.0242.19, T.0238.23 and J. 0222.20), Action de Recherche Concertée –

Fédération Wallonie-Bruxelles (ARC SLEEPDEM 17/27-09), Fondation Recherche Alzheimer (SAO-

FRA 2019/0025 & 2022/0014), Fondation Léon Fredericq, ULiège, European Regional

Development Fund (Radiomed, Biomed-Hub, WALBIOIMAGING).

AB is supported by Synergia Medial SA and the Walloon Region (Industrial Doctorate Program,

convention no. 8193). EB is supported by the Maastricht University - ULiège Imaging Valley. RS

and FB were supported by the European Union's Horizon 2020 research and innovation program

under the Marie Skłodowska-Curie grant agreement no. 860613. RS is supported by ULiège. NM,

IP, IC, NM, CP, KK, EK, Cba, MZ, FB and GV are/were supported by the FRS-FNRS. SS was

supported by ULiège-Valeo Innovation Chair and Siemens Healthineers. PT and LL are/were

supported by the EU Joint Programme Neurodegenerative Disease Research (JPND) IRONSLEEP

and SCAIFIELD projects, respectively – FNRS references: PINT-MULTI R.8011.21 & 8006.20. LL is

supported by the European Regional Development Fund (WALBIOIMAGING).

Author contributions

575 Study concept and design by NM, GV and PM. Data acquisition and analysis by NM, PT, EK, RS,

576 EB, AB, IC, IP, FB, ID & SS. Methodological support and/or support in the interpreting the data by

577 CBe, CBa, CP, PM, FC, MZ, LL. Funding was mostly obtained by CBa, FC, CP, PM and GV. NM and

578 GV drafted the first version of the manuscript. All authors revised the manuscript and had final

579 responsibility for the decision to submit for publication.

Potential Conflicts of Interest

581 All authors declare no conflict of interest.

Data availability

The processed data and analysis scripts supporting the results included in this manuscript are

publicly available via the following open repository: https://gitlab.uliege.be/CyclotronResearch-

Centre/Public/fasst/xxx (to be defined upon acceptance of the paper). The raw data could be

identified and linked to a single subject and represent a large amount of data. Researchers

willing to access to the raw should send a request to the corresponding author (GV). Data

sharing will require evaluation of the request by the local Research Ethics Board and the

signature of a data transfer agreement (DTA).

References

1.   Morin CM, Drake CL, Harvey AG, et al. Insomnia disorder. Nat. Rev. Dis. Prim.
2015;1(1):1–18.

2.   Crowley K. Sleep and sleep disorders in older adults. Neuropsychol. Rev. 2011;21:41–53.

3.   American Psychiatric Association D, Association AP. Diagnostic and statistical manual of
mental disorders: DSM-5. American psychiatric association Washington, DC; 2013.

4.   Van Someren EJW. Brain mechanisms of insomnia: new perspectives on causes and
consequences. Physiol. Rev. 2021;101(3):995–1046.

5.   Koshmanova E, Muto V, Chylinski D, et al. Genetic risk for insomnia is associated with
objective sleep measures in young and healthy good sleepers. Neurobiol. Dis.
2022;175:105924.

6.   Poe GR, Foote S, Eschenko O, et al. Locus coeruleus: a new look at the blue spot. Nat.
Rev. Neurosci. 2020;21(11):644–659.

7.   Sharma Y, Xu T, Graf WM, et al. Comparative anatomy of the locus coeruleus in humans
and nonhuman primates. J. Comp. Neurol. 2010;518(7):963–971.

8.   Lindvall O, Björklund A, Divac I. Organization of catecholamine neurons projecting to the
frontal cortex in the rat. Brain Res. 1978;142(1):1–24.

9.   Maness EB, Burk JA, McKenna JT, et al. Role of the locus coeruleus and basal forebrain in
arousal and attention. Brain Res. Bull. 2022;

10. Osorio-Forero A, Cherrad N, Banterle L, et al. When the locus coeruleus speaks up in
sleep: recent insights, emerging perspectives. Int. J. Mol. Sci. 2022;23(9):5028.

11. Gompf HS, Mathai C, Fuller PM, et al. Locus ceruleus and anterior cingulate cortex sustain
wakefulness in a novel environment. J. Neurosci. 2010;30(43):14543–14551.

12. Van Egroo M, Koshmanova E, Vandewalle G, Jacobs HIL. Importance of the locus
coeruleus-norepinephrine system in sleep-wake regulation: Implications for aging and
Alzheimer's disease. Sleep Med. Rev. 2022;62:101592.

13. Koshmanova E, Berger A, Beckers E, et al. Locus coeruleus activity while awake is
associated with REM sleep quality in older individuals. JCI insight 2023;8(20)

14. Aston-Jones G, Bloom F. Activity of norepinephrine-containing locus coeruleus neurons in
behaving rats anticipates fluctuations in the sleep-waking cycle. J. Neurosci.
1981;1(8):876–886.

15. Aston-Jones G, Rajkowski J, Cohen J. Role of locus coeruleus in attention and behavioral
flexibility. Biol. Psychiatry 1999;46(9):1309–1320.

16. Osorio-Forero A, Cardis R, Vantomme G, et al. Noradrenergic circuit control of non-REM
sleep substates. Curr. Biol. 2021;31(22):5009–5023.

17. Berger A, Koshmanova E, Beckers E, et al. Structural and functional characterization of the locus coeruleus in young and late middle-aged individuals. Front. neuroimaging 2023;2:1207844.

18. Mattis S. Mental status examination for organic mental syndrome in the elderly patient. Geriatr. psychiatry 1976;

19. Narbutas J, Van Egroo M, Chylinski D, et al. Cognitive efficiency in late midlife is linked to lifestyle characteristics and allostatic load. Aging (Albany NY) 2019;11(17):7169.

20. Van Egroo M, Narbutas J, Chylinski D, et al. Preserved wake-dependent cortical excitability dynamics predict cognitive fitness beyond age-related brain alterations. Commun. Biol. 2019;2(1)

21. Berthomier C, Drouot X, Herman-Stoïca M, et al. Automatic analysis of single-channel sleep EEG: Validation in healthy individuals. Sleep 2007;30(11):1587–1595.

22. Skorucak J, Arbon EL, Dijk D-J, Achermann P. Response to chronic sleep restriction, extension, and subsequent total sleep deprivation in humans: adaptation or preserved sleep homeostasis? Sleep 2018;41(7):zsy078.

23. Hoopes A, Mora JS, Dalca A V, et al. SynthStrip: Skull-stripping for any brain image. Neuroimage 2022;260:119474.

24. Gaser C, Dahnke R, Thompson PM, et al. CAT–A computational anatomy toolbox for the analysis of structural MRI data. biorxiv 2022;2006–2022.

25. Faul F, Erdfelder E, Buchner A, Lang A-G. Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. Behav. Res. Methods 2009;41(4):1149–1160.

26. Kornmeier J, Bach M. The Necker cube—an ambiguous figure disambiguated in early visual processing. Vision Res. 2005;45(8):955–960.

27. Murphy PR, O'Connell RG, O'Sullivan M, et al. Pupil diameter covaries with BOLD activity in human locus coeruleus. Hum. Brain Mapp. 2014;35(8):4140–4154.

28. Einhauser W, Stout J, Koch C, Carter O. Pupil dilation reflects perceptual selection and predicts subsequent stability in perceptual rivalry. Proc. Natl. Acad. Sci. 2008;105(5):1704–1709.

29. Devilbiss DM, Waterhouse BD. Phasic and tonic patterns of locus coeruleus output differentially modulate sensory network function in the awake rat. J. Neurophysiol. 2011;105(1):69–87.

30. Aston-Jones G, Cohen JD. An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. Annu. Rev. Neurosci. 2005;28:403–450.

31. Kjaerby C, Andersen M, Hauglund N, et al. Memory-enhancing properties of sleep depend on the oscillatory amplitude of norepinephrine. Nat. Neurosci. 2022;25(8):1059–1070.

32.   Boyce R, Glasgow SD, Williams S, Adamantidis A. Causal evidence for the role of REM
664   sleep theta rhythm in contextual memory consolidation. Science (80-.).
665   2016;352(6287):812–816.

666   33.   Yang M, Logothetis NK, Eschenko O. Phasic activation of the locus coeruleus attenuates
667   the acoustic startle response by increasing cortical arousal. Sci. Rep. 2021;11(1):1409.

668   34.   Carrier J, Viens I, Poirier G, et al. Sleep slow wave changes during the middle years of life.
669   Eur. J. Neurosci. 2011;33(4):758–766.

670   35.   Scammell TE, Arrigoni E, Lipton JO. Neural Circuitry of Wakefulness and Sleep. Neuron
671   2017;93(4):747–765.

672   36.   Huber R, Felice Ghilardi M, Massimini M, Tononi G. Local sleep and learning. Nature
673   2004;430(6995):78–81.

674   37.   Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the pathologic process in
675   Alzheimer disease: age categories from 1 to 100 years. J. Neuropathol. Exp. Neurol.
676   2011;70(11):960–969.

677   38.   Hawkes CH, Del Tredici K, Braak H. A timeline for Parkinson's disease. Parkinsonism Relat.
678   Disord. 2010;16(2):79–84.

679   39.   Van Egroo M, Chylinski D, Narbutas J, et al. Early brainstem [18F] THK5351 uptake is
680   linked to cortical hyperexcitability in healthy aging. JCI insight 2021;6(2)

681   40.   Van Egroo M, van Someren EJW, Grinberg LT, et al. Associations of 24-Hour Rest-Activity
682   Rhythm Fragmentation, Cognitive Decline, and Postmortem Locus Coeruleus
683   Hypopigmentation in Alzheimer's Disease. Ann. Neurol. 2024;

684   41.   Zeitzer JM. Control of sleep and wakefulness in health and disease. In: Gillette MU, editor.
685   Progress in Molecular Biology and Translational Science. 2013 p. 137–154.

686   42.   Tucker AM, Dinges DF, Van Dongen HPA. Trait interindividual differences in the sleep
687   physiology of healthy young adults. J. Sleep Res. 2007;16(2):170–180.

688   43.   Prokopiou PC, Engels-Domínguez N, Papp K V, et al. Lower novelty-related locus
689   coeruleus function is associated with Aβ-related cognitive decline in clinically healthy
690   individuals. Nat. Commun. 2022;13(1):1571.