Glycolytic flux controls retinal progenitor cell differentiation via regulating Wnt signaling

Joseph Hanna[1,2,3], Yacine Touahri[1,3,4], Alissa Pak[1,2], Lauren Belfiore[1,2], Edwin van Oosten[1,4], Luke Ajay David[1,2,3], Sisu Han[5], Yaroslav Ilnytskyy[6], Igor Kovalchuk[6], Deborah M Kurrasch[5], Satoshi Okawa[7,8,9,10], Antonio del Sol[10,11,12], Robert A Screaton[1,4], Isabelle Aubert[1,2], Carol Schuurmans[1,2,3,4]*

[1]Biological Sciences, Sunnybrook Research Institute, 2075 Bayview Ave, Toronto, ON, M4N 3M5, Canada
[2]Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, M5S 1A8, Canada
[3]Department of Ophthalmology and Vision Sciences, University of Toronto, Toronto, ON, M5T 3A9, Canada
[4]Department of Biochemistry, University of Toronto, Toronto, ON, M5S 1A8, Canada
[5]Department of Medical Genetics, Hotchkiss Brain Institute, Alberta Childrens' Hospital Research Institute, University of Calgary, Calgary, AB, T2N 1N4, Canada
[6]Department of Biological Sciences, University of Lethbridge, Lethbridge, AB, T1K 3M4, Canada
[7]Pittsburgh Heart, Lung, and Blood Vascular Medicine Institute, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.
[8]Department of Computational and Systems Biology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA
[9]McGowan Institute for Regenerative Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA
[10]Computational Biology Group, Luxembourg Centre for Systems Biomedicine, U of Luxembourg, L-4362 Esch-sur-Alzette, Luxembourg
[11]CIC bioGUNE, Bizkaia Technology Park, 48160 Derio, Spain
[12]IKERBASQUE, Basque Foundation for Science, Bilbao 48013, Spain

*To whom correspondence should be addressed: Carol Schuurmans, Sunnybrook Research Institute, 2075 Bayview Ave, Room A340**,** Toronto, ON, M4N 3M5**.** Phone: (416) 480-6100
E-mail: cschuurm@sri.utoronto.ca

Author Contributions: Conceptualization: JH, CS; Methodology: JH, YT, AP, LB, EvO, LAD, SH, YI, IK, DMK, SO, AdS, RS, CS; Investigation: JH, YT, AP, EvO, LAD, SH, YI, SO; Visualization: JH, YT, AP, LB, SH, LAD; Funding acquisition: IA, RS, CS; Project administration: CS; Supervision: CS, RS, IA, IK, DK, AdS; Writing – original draft: JH, CS; Writing – review & editing: JH, YT, AP, LB, EvO, LAD, SH, YI, IK, DMK, SO, AdS, IA, RS, CS. All the authors contributed to intellectual discussions and the direction of the project.

Competing Interest Statement: All authors declare they have no actual or potential competing interests.

Keywords: *Pten*, retinal progenitor cells, glycolysis, intracellular pH, photoreceptor differentiation, rod outer segments, Wnt signaling

This PDF file includes:
Main text
Figures 1 to 7
Word count: 5160 (excluding materials and methods)

ABSTRACT

Metabolic pathways are remodeled in response to energy and other homeostatic demands and are

dynamically regulated during embryonic development, suggesting a role in guiding cellular

differentiation. Here, we show that glycolytic flux is required and sufficient to bias multipotent

retinal progenitor cells (RPCs) to acquire a rod photoreceptor fate in the murine retina. In RPC-

specific *Phosphatase and tensin homolog* conditional knock-out (*Pten*-cKO) and RPC-specific

conditional gain-of-function of dominant active *PFKB3* (cyto*PFKB3*) mice, glycolytic gene

expression and activity are elevated, correlating with precocious rod photoreceptor

differentiation and outer segment maturation. Conversely, glycolytic inhibition in retinal explants

suppresses RPC proliferation and photoreceptor differentiation, achieved either with 2-deoxy-d-

glucose, a competitive inhibitor of glucose metabolism, by lowering media pH, which disables

PKM2, a rate-limiting enzyme, or by inhibiting lactate/H^+ symporters, which lowers intracellular

pH,. Mechanistically, we show that Wnt signaling, the top-upregulated pathway in *Pten*-cKO

retinas, is a glycolysis-dependent pathway. Pharmacological and genetic perturbation of Wnt

signaling by knocking-out *Ctnnb1,* encoding β-catenin, phenocopies glycolytic inhibition,

suppressing RPC proliferation, photoreceptor differentiation and outer segment maturation.

Thus, developmental rewiring of glycolytic flux modulates Wnt signaling to drive rod

photoreceptor differentiation and maturation, an instructive role that may be exploited

therapeutically for cell replacement strategies.

IMPACT STATEMENT

Transgenic and pharmacological approaches reveal developmental elevations in glycolytic flux

have an instructive role in promoting rod photoreceptor differentiation and maturation via

activation of Wnt signaling.

INTRODUCTION

Multiple neural cell types comprise the mature retina, all of which must be produced in correct numbers with precise identities for effective visual processing. Included are rod and cone photoreceptors, the light-sensing neurons; amacrine, bipolar and horizontal cells, retinal interneurons; ganglion cells, the output neurons of the retina; and supportive glial cells known as Müller glia. Clonal analyses across vertebrate species have revealed that all seven retinal cell types are derived from multipotent retinal progenitor cells (RPCs) that give rise to clones of varying sizes and compositions (Alexiades & Cepko, 1997; Fekete et al., 1994; Gomes et al., 2011; Holt et al., 1988; Jensen & Raff, 1997; Turner & Cepko, 1987; Turner et al., 1990; Wetts & Fraser, 1988). In mice, RPCs begin to differentiate at embryonic day (E) 12, completing their proliferation and differentiation by postnatal day (P) 10 -12 (Cepko, 2014; Young, 1985; X. Zhang et al., 2023). Symmetric proliferative (P/P) divisions predominate early in development, followed by a shift to asymmetric (P/D) divisions that renew the RPC pool and generate differentiated cells, and finally, by symmetric differentiative (D/D) divisions that generate two postmitotic cells (Baye & Link, 2007; Nerli et al., 2020; Paolini et al., 2015). At each stage, individual RPC fate decisions are largely stochastic, albeit with temporal biases, such that the seven retinal cell types are born in overlapping windows (Gomes et al., 2011; He et al., 2012). Ganglion, horizontal and cone cells are born in the earliest embryonic stages, followed closely by amacrine cells, and finally by bipolar cells, rods and Müller glia, which mainly differentiate in the first two postnatal weeks (Rapaport et al., 2004; Young, 1985). How RPCs interpret extrinsic signals and intrinsic contextual cues to ultimately produce the correct numbers of each retinal cell type remains incompletely understood.

Cellular energy and precursor metabolites are produced by the metabolism of glucose, which is converted into pyruvate via glycolysis. Glycolysis is a multi-step enzymatic process that occurs in the cytosol, does not require O_2, and produces 2 molecules of ATP per glucose molecule (Hanna et al., 2022). Pyruvate is either transported to mitochondria for further oxidation to drive oxidative phosphorylation (OXPHOS) by the electron transport chain, or converted to lactate and secreted out of the cell. OXPHOS is an O_2-dependent metabolic pathway that produces 30-38 ATP molecules for each glucose molecule, and hence, produces more cellular energy than glycolysis (Hanna et al., 2022). Yet, the majority of vertebrate retinal cells rely on glycolysis as the main metabolic pathway, even in the presence of O_2 (Hurley et al.,

2015). The preference for aerobic glycolysis has been termed the Warburg effect, an adaptation

commonly observed in cancer cells that is also used by some 'normal' cells (Vander Heiden et

al., 2009; Warburg, 1956a, 1956b). Included are dividing neural progenitor cells, which use

aerobic glycolysis for energy production across tissue types, including in the retina (Hanna et al.,

2022). In Xenopus, RPCs rely on glycogen stores as a fuel source for glycolysis (Agathocleous

et al., 2012), whereas in human retinal organoids, glucose is the essential fuel, and is required for

RPC expansion at the earliest stages of optic vesicle formation (Takata et al., 2023). Finally,

during murine retinal development, mitophagy clears mitochondria to drive a metabolic shift

towards glycolysis, which supports early RPC differentiation into ganglion cells (Esteban-

Martinez et al., 2017). However, whether metabolic reprogramming and levels of glycolytic

activity influence RPC differentiation decisions during later developmental stages is not known.

Metabolic reprogramming, or alterations in metabolic rates and/or pathways, occurs in

stem and progenitor cells during tissue development (Tatapudy et al., 2017), including in the

retina (Esteban-Martinez et al., 2017). In a few lineages, metabolism has been ascribed

instructive roles in driving cellular differentiation, including in skeletal muscle and helper T cell

lineages (Peng et al., 2016; Ryall et al., 2015; Tatapudy et al., 2017). In these lineages, glycolysis

is important for more than energy production, for instance, modifying histone acetylation in

murine skeletal muscle (Ryall et al., 2015), activating Wnt signaling to control chick

neuromesodermal progenitor cell fate choice (Oginuma et al., 2020), and altering protein

subcellular localization in the murine presomitic mesoderm (Miyazawa et al., 2022). To further

investigate how developmental signaling pathways intersect with metabolism to control RPC fate

choice in the retina, we focused on PTEN, an intracellular protein and lipid phosphatase with

pleiotropic actions, including the direct inhibition of glycolysis through dephosphorylation and

consequent inactivation of phosphoglycerate kinase 1 (PGK1), a rate-limiting glycolytic enzyme

(Qian et al., 2019). In *Pten* conditional knockout (KO) retinas, RPCs proliferate excessively

early in embryonic development, accompanied by the accelerated differentiation of

photoreceptors in the outer nuclear layer (ONL) (Cantrup et al., 2012; Jo et al., 2012; Tachibana

et al., 2016; Tachibana et al., 2018). Here, we show that elevated glycolysis accompanies the

accelerated RPC differentiation kinetics in *Pten*-cKO retinas, which ultimately results in the

precocious depletion of the RPC pool. Similarly, in an RPC-specific, conditional gain-of-

function model of dominant active *PFKB3* (cyto*PFKB3*), glycolytic gene expression and activity

are elevated, correlating with precocious rod photoreceptor differentiation and outer segment

(OS) maturation. Conversely, multiple inhibitors of glycolysis reduce RPC proliferation and

photoreceptor differentiation. Finally, we found that glycolysis is required and sufficient to

control Wnt signaling, and show that Wnt activity is required for rod photoreceptor

differentiation and OS maturation. We have thus uncovered a novel glycolytic flux-Wnt

signaling axis as an important regulator of RPC fate choice.

RESULTS

Elevated glycolysis and accelerated rod photoreceptor differentiation correlate in *Pten-*

cKO retinas

Neural progenitor cells in the embryonic brain (Khacho & Slack, 2018) and retina

(Agathocleous et al., 2012) preferentially use glycolysis for energy production. However,

glycolytic metabolites and glycolytic gene expression are not at uniform levels throughout retinal

development, peaking at E18.5-P0 (Esteban-Martinez et al., 2017). Since these measurements

were performed in bulk, and the ratio of RPCs to differentiated cells declines as development

proceeds, it is not clear whether glycolytic activity is temporally regulated within RPCs and/or in

other retinal cell types. To assess glycolytic gene expression within specific cell populations

(Figure 1A), we mined scRNA-seq data collected from developing retinas between E11 and P14

(Clark et al., 2019). High dimensional uniform manifold approximation and projection (UMAP)

plots identified unique cell clusters corresponding to pre-neurogenic (pre-N) RPCs at E11, early-

RPCs from E12-E18 and late-RPCs from P0-P5 (Figure 1B,C). Feature plots were used to

examine glycolytic gene expression in individual retinal cell types (Figure 1C). Several

glycolytic genes were specifically enriched in early compared to late RPC clusters, including

Pgk1, a rate-limiting enzyme that catalyzes the first ATP-producing step in the glycolytic chain

reaction, and *Eno1*, another rate-limiting enzyme (Figure 1C). A less obvious bias towards early

RPC expression was observed for *Hk1*, the initial enzyme in the glycolytic pathway that converts

glucose to glucose-6-phosphate, *Pfk1*, another rate-limiting enzyme, and *Slc16a3,* encoding the

lactate/H^+ symporter that shuttles lactate together with a H^+ ion (Figure 1C). In contrast, *Aldoa,*

Gpi1 and *Ldhb* showed similar expression levels between pre-N, early and late RPCs (Figure

1C). These data confirm that glycolytic gene expression is elevated in RPCs and may peak

during the early-to-late RPC transition.

We noted that the highest levels of *Pgk1* expression at E18/P0 corresponded to the known

pinnacle of RPC differentiation into committed, rod photoreceptor precursors (Kim et al., 2016)

(Figure 1C), which undergo a protracted differentiation period, only expressing mature

photoreceptor markers between P2 and P8 (e.g., *Prph2, Impg2, Pde6g, Gm11744, Gnat2*; Figure

1-figure supplement 1). We thus asked whether levels of glycolytic activity have an instructive

role in rod photoreceptor fate selection. To test this hypothesis, we used two genetic models to

elevate glycolytic activity in RPCs. We first targeted PTEN, a phosphatase that dephosphorylates

and inactivates PGK1, a rate-limiting step (K. Zhang et al., 2023). *Pten* is expressed throughout

retinal development (Figure 1-figure supplement 1), including in late-RPCs (E18.5), which co-

express VSX2 and PAX6 (Figure 1D). To delete *Pten* in RPCs, we generated a *Pten*-cKO model

by crossing a floxed allele of *Pten,* in which exons 4 and 5 were flanked by loxP sites, with a

Pax6-Cre-GFP driver allele (Figure 1E). As a proxy of Cre activity, GFP was expressed in RPCs

and postmitotic newborn cells with a high peripheral to central gradient in *Pax6-Cre-GFP*

transgenics (Figure 1F). Thus, for the remainder of this study, we focused on the peripheral

retina where the *Pax6-Cre* allele was most active, avoiding the central retina, which was readily

distinguishable as it forms a visible hamartoma comprised of remaining wild-type cells in *Pten*-

cKO retinas (Tachibana et al., 2018).

Using cellular birthdating, we previously showed that *Pten*-cKO RPCs are

hyperproliferative and differentiate on an accelerated schedule between E12.5 and E18.5, yet

fewer rod photoreceptors are ultimately present in P7 (Tachibana et al., 2016) and P21 (Hanna et

al., 2025) retinas, suggestive of a developmental defect. Since retinal cell differentiation persists

until ~P12, we conducted similar birthdating experiments, but instead administered BrdU to P4

pups. In wild-type retinas harvested at P14, BrdU cells were mainly distributed between the INL,

where late-born bipolar cells and Müller glia localize, and the ONL, where rod photoreceptors

migrate (Figure 1G,H). Labeled cells in the GCL likely represent displaced amacrine cells, which

are at the end of their differentiation window. In contrast, only a few BrdU-labeled cells were

detected in the GCL in *Pten*-cKO retinas, with no BrdU$^+$ cells present in the INL and ONL

(Figure 1G,H). Thus, the accelerated differentiation observed during embryonic stages is

followed by the precocious depletion of dividing RPCs in *Pten*-cKO retinas by the early

postnatal period, effectively limiting the overall period of rod photoreceptor differentiation

(Figure 1G,H).

Glycolytic gene expression and activity are upregulated in P0 *Pten*-cKO retinas

To investigate how *Pten,* encoding an intracellular phosphatase, controls the timing of

RPC proliferation and differentiation, we began with a transcriptomic approach. To decipher

transcriptomic changes associated with *Pten* loss, we performed bulk RNA sequencing (RNA-

seq) on P0 wild-type (N=4) and *Pten*-cKO (N=5) retinas. A heatmap depiction revealed that

there are distinct blocks of expressed genes in the two genotypes (Figure 2A; Figure 2– Data

supplement 1). Among the differentially expressed genes (DEGs) with a log2-fold change

(log2FC) ≥2 between wild-type and *Pten*-cKO retinas, there were 408 down-regulated and 667

up-regulated genes (Figure 2B). To draw relationships between genes and pathways, we analysed

the Kyoto Encyclopedia of Genes and Genomes (KEGG) database, showing enrichment of terms

such as *axon guidance*, and *regulation of actin cytoskeleton* among the DEGs. In addition,

metabolic pathway terms were enriched, including *glycolysis* (Figure 2C), which involves a

series of enzymatic reactions that break down glucose into pyruvate for energy production

(Figure 2D). Indeed, analysis of Reads Per Kilobase per Million mapped reads (RPKM) revealed

that most of the glycolytic regulatory enzymes were upregulated in *Pten*-cKO retinas, including

Hk1, Gpi1, Pfkl, Aldoa, Pgk1, Pgam1, and *Eno1,* although others, such as *Pkm,* were not

differentially expressed (Figure 2E). Notably, since we employed 75 bp single-end sequencing,

we could not distinguish between alternatively spliced *Pkm1* and *Pkm2* mRNAs, bearing a

common 3' end. Additionally, expression of *Slc16a3*, encoding a monocarboxylate transporter

that symports lactate and one hydrogen ion outside of the cell, was increased in *Pten*-cKO retinas

(Figure 2E). We validated these increases in gene expression for a handful of genes, including

Hk1, Pgk1, Eno1, and *Slc16a3,* using qPCR, confirming that these glycolytic genes were indeed

upregulated in E15.5 *Pten*-cKO retinas (Figure 2F).

To test whether transcriptomic changes in P0 *Pten*-cKO retinas reflected an increase in

glycolytic activity, we performed a Seahorse assay. Extracellular acidification rate (ECAR) was

elevated in P0 *Pten*-cKO RPCs compared to controls, indicative of an increase in lactic acid

levels, which serves as a proxy measure for glycolysis (Figure 2G). We further assessed

mitochondrial O_2-consumption rates in P0 retinal cells, showing that oligomycin, an ATP

synthase inhibitor, suppressed basal respiration rates to the same extent in wild-type and *Pten*-

cKO retinas (Figure 2G). Similarly, maximum or spare respiratory capacity measured upon

stimulation did not differ between P0 wild-type and *Pten*-cKO retinas (Figure 2G). Finally, we

used a colorimetric assay to measure lactate levels in tissue lysates, revealing an elevation of

lactate in *Pten*-cKO retinas compared to wild-type retinas at P0 (Figure 2H). Taken together,

these data indicate that glycolytic gene expression and activity are elevated in *Pten*-cKO retinas,

whereas oxygen-dependent OXPHOS is unaffected.

Pharmacological inhibition of glycolysis decreases RPC proliferation and photoreceptor

differentiation

Glucose is transported across cell membranes via glucose transporters (GLUT), with

Slc2a1, encoding GLUT1, expressed at the highest levels in dividing, *Cdk1*$^+$ RPCs (Figure 3A).

To assess the requirement for glycolysis during retinal development, we developed an *in vitro*

retinal explant assay, dissecting retinas from P0 eyes and removing all non-retinal tissues,

including the retinal pigment epithelium (RPE) (Figure 3B). P0 explants were mounted on

Nucleopore membranes and cultured on top of retinal explant media, providing a source of

nutrients, growth factors and glucose. To block glycolysis, retinal explants were cultured with 2-

deoxy-d-glucose (2DG), a competitive inhibitor that is transported by GLUT1 and converted by

HK1/2 to 2-deoxy-D-glucose-6-phosphate, an inhibitor of the glycolytic pathway (Figure 3C).

2DG is typically used in the range of 5-10 mM in cell culture studies and in general, has anti-

proliferative effects (Zhang et al., 2014). To test whether 5 mM- and 10 mM-2DG impacted RPC

proliferation, explants were exposed to BrdU, which is incorporated into S-phase cells, for 30

minutes prior to harvesting. 2DG treatment resulted in a dose-dependent inhibition of RPC

proliferation as evidenced by a reduction in BrdU$^+$ cells (Figure 3D). To determine whether

glycolytic inhibition reduced the overall percentage of RPCs that continued to proliferate, we

measured the proliferative fraction (p-fraction) by exposing retinal explants to BrdU at the onset

of 2DG treatment, and then immunostaining for BrdU and Ki67, which labels all cycling cells.

The p-fraction was calculated as the ratio of Ki67$^+$ cells that were BrdU$^+$ (*i.e.,* %BrdU$^+$Ki67$^+$/

Ki67$^+$ cells). The total numbers of BrdU$^+$ and Ki67$^+$ cells declined in 2DG-treated explants, as

did the overall p-fraction (Figure 3E), indicating that our 2DG treatment was in the effective
range.

A reduction in the p-fraction could indicate that RPCs had stopped cycling during the 24-
hr window of 2DG treatment, or that these cells had undergone apoptosis. To distinguish
between these possibilities, we first immunostained 2DG-treated retinal explants with VSX2,
which primarily labels RPCs at P0. 2DG treatment for 24 hr did not alter the total number of
VSX2$^+$ RPCs, even at the higher dose of 10 mM, despite the dramatic decline in proliferating
RPCs (Figure 3F). Furthermore, 2DG treatment did not elevate levels of apoptosis, as the
number of retinal cells expressing cleaved-caspase 3 (CC3), an executioner protease, remained
similar in 2DG-treated and control explants (Figure 3G). We then asked whether glycolytic
inhibition impacted cellular differentiation, focusing on CRX-expressing photoreceptor
precursors, which are the main cells generated at P0. At the highest 10 mM 2DG dose, the
number of CRX$^+$ cells declined, indicative of a dose-dependent decrease in the number of
photoreceptor precursors born (Figure 3H). Taken together, these data indicate that glycolytic
inhibition suppresses RPC proliferation, and at higher doses, prevents RPCs from differentiating
into photoreceptors. Stated differently, glycolysis is required to support RPC proliferation and
photoreceptor differentiation.

Previous studies in *Xenopus* similarly found that RPCs rely mainly on glycolysis for
energy production, however, instead of the demand for a constant supply of glucose, Xenopus
RPCs depend on glycogen storage (Agathocleous et al., 2012). To test whether mouse RPCs
similarly require stored glycogen, P0 retinal explants were cultured in the presence of 12.5 µM
and 25 µM glycogen phosphorylase inhibitor (GPI) for 24 hr (Figure 3I), the latter corresponding
to the same concentrations used in Xenopus retina (Agathocleous et al., 2012). In murine retinal
explants, GPI treatment did not impact the number of RPCs that incorporated BrdU (Figure 3I).
These data suggest that rather than depending on glycogen stores, murine RPCs depend on a
constant supply of glucose to maintain proliferation, as shown in human retinal organoids
(Takata et al., 2023).

Precocious rod differentiation and maturation in *cytoPFKB3* gain-of-function mice
phenocopies *Pten*-cKO retinal features

We previously showed that at P7, *Pten*-cKO retinas are comprised of fewer differentiated

rods and amacrine cells, while other cell types are not affected (Tachibana et al., 2016). Given

the increase in glycolysis in *Pten*-cKO retinas observed in the current study, we asked whether

metabolic reprogramming may be sufficient to bias RPC fate choice, thereby altering cellular

composition. As a precedent, glycolytic flux alters neuromesodermal lineage selection in the

chick tail bud (Oginuma et al., 2020) and in mouse mesoderm (Miyazawa et al., 2022). To

determine whether increased glycolysis is a critical downstream effector pathway of *Pten* in

retinal cell fate selection, we used a transgenic approach to elevate glycolytic flux in RPCs.

Glycolytic flux is defined as the conversion of fructose 6-phosphate (F6P) to fructose-1,6-

bisphosphate (FBP), which is catalyzed by phosphofructokinase (PFK1), and is the first rate-

limiting, commitment step in glycolysis (Miyazawa et al., 2022) (Figure 4A). To promote

glycolytic flux, we used a conditional gain-of-function (GOF) transgenic line driving the

expression of a dominant active, cytoplasmic form of PFKB3 (cyto*PFKB3*) (Miyazawa et al.,

2022). PFKB3 converts F6P to fructose-2,6-bisphosphate, which is a strong, allosteric activator

of PFK1 (Shi et al., 2017) (Figure 4A). To overexpress cyto*PFKB3* in RPCs, we removed the

floxed stop cassette using a *Pax6-Cre* driver to create a cyto*PFKB3* GOF model (Figure 4B). Of

note, the same *Pax6-Cre* driver was employed to generate *Pten*-cKOs used in this study. To

confirm that cyto*PFKB3*-GOF retinas undergo increased glycolytic flux, we measured levels of

secreted lactate as a proxy measure using a colorimetric assay. We observed an overall increase

in lactate levels in P0 cyto*PFKB3*-GOF retinas (Figure 4C).

Having validated the GOF model, we next asked whether elevating cyto*PFKB3*

expression in RPCs impacted the timing of retinal cell differentiation using a birthdating assay.

BrdU was administered to pregnant females at E12.5 and pups were sacrificed at P7. In wild-

type retinas, BrdU$^+$ cells were mainly distributed in the GCL, where newborn ganglion cells and

displaced amacrine cells reside, and in the INL, where amacrine cells and horizontal cells are

located (Figure 4D,E). Fewer BrdU$^+$ cells labeled at E12.5 were present in the P7-ONL, with

these early-born ONL cells primarily including cones in wild-type retinas (Figure 4D,E). In

cyto*PFKB3*-GOF retinas, BrdU$^+$ cells were also distributed amongst the three layers, but BrdU$^+$

cells were more abundant in the ONL, where photoreceptors reside, with a concomitant

reduction in the INL (Figure 4D,E). The shift in distribution of E12.5-labeled, BrdU$^+$ cells

phenocopies the altered temporal differentiation observed in *Pten*-cKO retinas (Tachibana et al.,

2016).

The accelerated differentiation of ONL cells in cyto*PFKB3*-GOF retinas prompted us to

ask whether photoreceptor maturation was similarly hastened, focusing on OS formation as a

readout. While most differentiated cells use OXPHOS for energy production, photoreceptors are

an important exception, as these metabolically demanding cells require glycolysis to maintain

OSs, which undergo diurnal turnover (Chinchore et al., 2017). To examine OS formation, we

immunostained P7 wild-type and cyto*PFKB3*-GOF retinas with rhodopsin (Figure 4F). Visually,

photoreceptor OS above the ONL, which are labeled with rhodopsin, were larger in P7

cyto*PFKB3*-GOF retinas, as confirmed by a measurement of the OS area (Figure 4F).

To determine whether rod photoreceptor OSs similarly mature precociously in *Pten*-cKO

retinas, we first examined the P0 transcriptomic data for OS-enriched genes (Datta et al., 2015).

We identified several OS genes that were up-regulated in P0 *Pten*-cKO retinas, although not all,

as several OS genes were also down-regulated (Figure 4G). To assess OS maturation

morphologically, we immunostained P7 retinas with rhodopsin, revealing a similar increase in

the rhodopsin-labeled OS area in *Pten*-cKOs (Figure 4I). Finally, to investigate the role of

glucose metabolism in accelerating the rate of rod photoreceptor maturation, we inhibited

glycolysis during the early postnatal window by *in vivo* administration of 2DG, which was

administered daily from P0 to P7 (Figure 4H,I). Strikingly, 2DG treatment blocked the

precocious maturation of rod photoreceptor OSs in *Pten*-cKO retinas, which returned to wild-

type sizes (Figure 4I). Taken together, these data implicate glucose metabolism and glycolysis as

an important driver of photoreceptor differentiation and OS maturation.

Reducing pHi phenocopies the inhibitory effects of a glycolytic block on RPC proliferation

and differentiation

Glycolysis is not an energy-efficient metabolic pathway as cells produce 2 ATP/glucose

molecule versus 30-38 ATP/glucose when using OXPHOS (Touahri et al., 2024). However, cells

rely on glycolysis for reasons other than energy expenditure. For instance, lactate is a metabolic

byproduct that is shuttled out of the cell along with one H^+ ion by lactate/H^+ symporters (Slc16a

group proteins) when glycolytic activity is high, effectively increasing intracellular pH (pHi)

(Figure 5A). Notably, glycolysis-driven increases in pHi are the central driver of cell fate choice

selection in the chick neuromesodermal lineage (Oginuma et al., 2020). We found *Slc16a* family

lactate/H$^+$ symporters are elevated in *Pten*^{cKO} retinas (Figure 2E), suggesting that more protons

may be extruded to increase pHi. To address whether lactate/H$^+$ symporters control RPC

proliferation, we incubated wild-type P0 explants with 5mM α-cyano-4-hydroxycinnamate

(CNCn), an inhibitor of monocarboxylate transporters, including the Slc16a-family symporters,

for 24 hr (Figure 5B). The impact of CNCn should block the export of lactate into the

extracellular space, effectively lowering pHi and further decreasing glycolysis. To determine

whether RPC proliferation was impacted by CNCn treatment, explants were incubated with

BrdU 30 min before harvesting. We observed a decrease in the number of BrdU$^+$ proliferating

RPCs in CNCn-treated retinal explants compared to controls (Figure 5C). This decline in RPC

proliferation mirrored the effect of inhibiting glycolysis (Figure 3C). Thus, Slc16a-family

symporters play a role in regulating RPC proliferation in the retina.

In keeping with the idea that pH levels influence cell fate selection, a previous study

found that neuromesodermal progenitors in chick tailbud explants cultured in low pH media

undergo cell cycle exit and lineage selection is biased towards neural rather than mesodermal

fates (Oginuma et al., 2020). Given that altered glycolysis influences intracellular pH, which in

turn controls cell fate decisions, we set out to assess the impact of manipulating pH on cell fate

selection in the retina. One of the expected impacts of lowering pH was the aggregation of

PKM2, a rate-limiting enzyme for glycolysis, which aggregates in reversible, inactive amyloids

(Cereghetti et al., 2024). We grew retinal explants in media buffered at pH 6.5 (acidic), pH 7.4

(neutral), or pH 8.0 (basic). Increased glycolysis, leading to more H$^+$ extrusion from the cell and

an increase in pHi was modeled by pH 8.0, while decreased glycolysis, leading to less H$^+$

extrusion and a lower pHi, was modeled by pH 6.5. To assess the impact of pHi changes on RPC

proliferation, P0 retinal explants were cultured in media buffered to different pH levels, with

BrdU added for 30 min and at the end of the 24-hr culture period. More RPCs were BrdU$^+$ in pH

8.0 media compared to neutral pH 7.4 media, consistent with the pro-proliferative effect of

increased glycolysis (Figure 5D). Conversely, in media buffered to pH 6.5, the number of BrdU$^+$

proliferating RPCs declined (Figure 5D).

To determine whether pH changes alter the overall percentage of RPCs that continue to

proliferate, we measured the p-fraction by exposing retinal explants to BrdU at the onset of pH

adjustments, and then calculating the p-fraction as the ratio of Ki67$^+$ cells that were BrdU$^+$. We

found a pH-dependent increase in the number of BrdU$^+$ RPCs and in the p-fraction in retinal

explants cultured in pH 8.0 compared to pH 7.4 (neutral pH control) media (Figure 5E). We also

observed a corresponding increase in VSX2$^+$ RPCs in pH 8.0 media (Figure 5F), and more RPCs

differentiated into CRX$^+$ photoreceptor precursors (Figure 5H). In contrast, in retinal explants

cultured in pH 6.5 media, fewer RPCs entered S-phase as evidenced by the decline in BrdU$^+$

RPCs and lower p-fraction (Figure 5E). However, there were more VSX2$^+$ RPCs in retinal

explants grown in pH 6.5 versus pH 7.4 media (Figure 5F), with no effect on cell death evident

as assessed by cleaved-caspase 3$^+$ cell counts (Figure 5G), likely because fewer of these RPCs

differentiated into CRX$^+$ photoreceptor precursors (Figure 5H). Thus, pH has a dramatic effect

on the decision by RPCs to proliferate versus differentiate, with higher pH more conducive to

rapid, self-renewing, asymmetric divisions (P/D) that generate an RPC and a neuron.

Tissues naturally have pHi and differentiation gradients – for instance, there is an anterior

(low pHi, more differentiated cells) to posterior (high pHi, more neuromesodermal progenitor

cells) gradient in neuromesodermal progenitors in the embryonic chick tailbud (Oginuma et al.,

2020). Given the importance of pH to control RPC proliferation and differentiation dynamics,

and since not all RPCs differentiate at any given time, we asked whether there was a dynamic

representation of pH within RPCs within the developing retina. To test this assumption, we

electroporated P3 retinal explants with pCAGG-pHluorin-IRES2-tdTomato, using tdTomato to

label transfected cells and pHluorin, a ratiometric pH sensor with a 488-nm activation spectrum

in neutral pH. As expected, the 488-nm signal was variable across RPCs that had similar

tdTomato levels (Figure 5I). Taken together, this data is consistent with the idea that pH varies

within RPCs, correlating with the propensity of some RPCs to continue to proliferate while

others differentiate.

Canonical Wnt signaling is active in neurogenic RPCs and controlled by glycolysis

Our finding that glycolysis might be a central driver of RPC fate selection, in part

through the control of pHi, prompted the question of how these biochemical processes might

impact cell fate selection. To glean further insights, we revisited our bulk RNA-seq data set from

P0 *Pten*-cKO retinas and performed a PANTHER (protein analysis through evolutionary

relationships) classification analysis of DEGs (Figure 6-figure supplement 1). As expected,

glycolysis was amongst the up-regulated pathways, but the most up-regulated pathway in P0

Pten-cKO retinas was Wnt signaling (Figure 6-figure supplement 1). Wnts bind to frizzled (Fz)

and low-density lipoprotein receptor-related protein receptors (LRPs), which induces β–catenin

translocation to the nucleus, where it forms an active transcriptional complex with T-cell-specific

transcription factor (TCF)/lymphoid enhancer-binding factor (LEF) transcription factors (Figure

6A). Prior studies had implicated canonical Wnt signaling in maintaining an undifferentiated

RPC state, especially in the peripheral margin at early developmental stages (Ouchi et al., 2011).

However, mining scRNA-seq data from E11 to P14 retinas (Clark et al., 2019) revealed that Wnt

signaling may be active throughout retinal development, with transcripts for *Wnt7b* and *Lef1,*

which is transcribed in response to canonical Wnt signaling (Schmidt et al., 2000), detected in

RPCs at all developmental stages (Figure 6B). Furthermore, we noted that *Dkk*, a Wnt inhibitor,

is only expressed in pre-neurogenic RPCs, and not in RPCs in the early and late neurogenic

periods, suggesting that Wnt signaling may not be active prior to the onset of neurogenesis

(Figure 6B). We thus explored whether Wnt signaling was regulated by glycolysis to play a role

in RPC differentiation.

Components of the Wnt signaling pathway that were upregulated in P0 *Pten*-cKO retinas

included *Dvl3*, an inhibitor of the β-catenin destruction complex, *Apc* and *Gsk3b*, two

components of the destruction complex, and *Tcf4* and *Crebbp*, a downstream effector of Wnt

signaling and its co-activator, respectively (Figure 6C). These data were consistent with our

previous demonstration that canonical Wnt signaling levels are elevated in *Pten*-cKO retinas, as

assessed by increased expression of non-phosphorylated (active) β-catenin (Touahri et al., 2024).

However, unlike in the peripheral retina of *Ctnnb1*-cKOs (encoding for β-catenin), in which the

neurogenic inhibitor *Id3* is activated by Wnt signaling in the periphery to maintain RPCs in an

immature, undifferentiated state (Ouchi et al., 2011), we found that *Id3* is down-regulated in P0

Pten-cKO retinas (Figure 6C), potentially allowing differentiation to proceed unimpeded.

Finally, to establish whether Wnt signaling was indeed regulated by glycolytic activity,

we investigated β-catenin levels in explants treated with 2DG or grown in different pH media. β-

catenin levels declined upon glycolysis inhibition in 2DG treated explants (Figure 6D), and at pH

6.5, a proxy for low glycolysis (Figure 6E). Thus, canonical Wnt signaling is elevated in P0

Pten-cKO retinas (Touahri et al., 2024), in which glycolysis is elevated, and reduced in
conditions in which glycolysis is inhibited.

Wnt signaling controls rod photoreceptor differentiation and maturation

We next asked whether inhibiting Wnt signaling would phenocopy the effects of reducing
glycolysis on RPC proliferation and photoreceptor differentiation. For this purpose, we treated
P0 wild-type CD1 retinal explants with 5 µM or 10 µM FH535, a pharmacological inhibitor of
Wnt/β-catenin signaling (Handeli & Simon, 2008), for 24 hr. To label proliferating RPCs, BrdU
was added 30 min before harvesting similar to previous experiments. FH535 treated retinas
showed a significant decrease in the number of BrdU$^+$ RPCs (Figure 7A) and a concomitant
decrease in the number of CRX$^+$ photoreceptor precursors (Figure 7B) without impacting
apoptosis (Figure 7C). To further study the role of Wnt signaling in photoreceptor maturation,
we crossed a floxed allele of *Ctnnb1*, encoding for β-catenin, with a *Pax6-Cre* driver, which we
previously confirmed leads to β-catenin loss in P14 *Ctnnb1*-cKO retinas (Touahri et al., 2024).
P14 wild-type and *Ctnnb1*-cKO retinal sections were immunolabeled with rhodopsin, showing a
clear reduction in OS area upon the inhibition of Wnt signaling (Figure 7D).

Finally, in chick neuromesodermal progenitors, high glycolytic activity or elevated pHi
promotes the non-enzymatic acetylation of β-catenin, which promotes mesodermal at the
expense of neural cell fate selection (Oginuma et al., 2020). To force protein acetylation,
including of β-catenin, cells can be grown in sodium acetate. We observed that the exposure of
retinal explants to 5-, 10-, or 20 mM sodium acetate increased the percentage of BrdU$^+$
proliferating RPCs (Figure 7E). Thus, the forced activation of β-catenin acetylation is sufficient
to promote RPC proliferation, mimicking the early impact of *Pten* loss/high glycolytic activity in
RPCs (Tachibana et al., 2016). Taken together, our study identifies an unexplored pathway in
which PTEN controls RPC proliferation and rod differentiation and maturation via modulating
glycolysis. Glycolysis in turn controls proliferation and differentiation through modulating pHi
that creates a favorable environment for Wnt signaling activation.

DISCUSSION

The timing of RPC differentiation and neural subtype specification must be precisely

coordinated so that the correct types of retinal neurons and glial cells are born at appropriate

times and in their proper numbers during development. In *Pten*-cKO retinas, RPCs proliferate

and differentiate on an accelerated timeline (Jo et al., 2012; Sakagami et al., 2012; Tachibana et

al., 2016), resulting in the precocious depletion of proliferating RPCs (this study). Ultimately, the

changes in RPC proliferation dynamics culminate in the production of fewer amacrine cells and

rod photoreceptors populating *Pten*-cKO retinas at P7 and P21 (Hanna et al., 2025; Tachibana et

al., 2016; Touahri et al., 2024). To understand how *Pten* controls RPC fate decisions, we first

used a transcriptomic approach, revealing an increase in glycolytic gene expression and activity

in *Pten*-cKO retinas. Consistent with a role for glycolysis in controlling RPC proliferation and

photoreceptor cell differentiation, both were inhibited by 2DG, a pharmacological inhibitor of

glycolysis. Conversely, the promotion of glycolytic flux in RPCs using cyto*PFKB3* GOF

transgenic mice accelerated photoreceptor differentiation, including the enhanced maturation of

rod OSs, an effect phenocopied in *Pten*-cKO retinas. Since elevated glycolytic activity shuttles

more lactate/H^+ out of the cell to increase intracellular pH (pHi), we cultured retinal explants in

media buffered to higher pH levels, which promoted RPC proliferation and accelerated

photoreceptor differentiation. Finally, we linked glycolysis and pHi changes to the activation of

Wnt signaling, which we found is required to promote RPC proliferation, photoreceptor

differentiation, and OS maturation. Taken together, we have uncovered a novel molecular axis

controlling RPC fate decisions in the retina that begins with *Pten*, a signal transduction molecule,

the loss of which elevates glycolytic activity, pHi, and Wnt signaling (Figure 7E).

The role of glycolysis in retinal development across different species

The increased glycolytic gene expression in *Pten*-cKO retinas is likely tied to the

increased expression of hypoxia-induced-factor-1-alpha (*Hif1a*), a known target of mTOR

signaling and transcription factor that induces *Slc1a3* (GLUT1) and glycolytic gene expression

(Hanna et al., 2022). Indeed, mTOR signaling is hyperactive in *Pten*-cKO retinas (Cantrup et al.,

2012; Tachibana et al., 2016; Tachibana et al., 2018; Touahri et al., 2024), and likewise, in *Tsc1*-

cKO retinas, which also increase glycolysis via HIF-1A (Lim et al., 2021). Similar to our

findings in mice, glycolysis is the predominant mode of ATP production in proliferating *Xenopus*

and zebrafish RPCs (Agathocleous et al., 2012). However, in contrast to the results of our study, treatment of Xenopus retinas with 2DG, a glucose inhibitor, does not impact ATP or lactate production, whereas GPI-mediated inhibition of access to glycogen stores reduces energy production (Agathocleous et al., 2012). Thus, Xenopus RPCs rely on glycogen storage to fuel glycolysis, whereas we found that in mouse retina, a constant glucose supply is required to fuel glycolysis and support RPC proliferation. Consistent with our data, in mice carrying mutations in *Glut1*, a glucose transporter, or *Ldha*, a glycolytic enzyme that converts pyruvate to lactate, RPC proliferation and expansion of the optic vesicle are inhibited (Takata et al., 2023). Human RPCs also use glycolysis as revealed by the conversion of ^{13}C-glucose to lactate, a glycolytic intermediate (Takata et al., 2023). Moreover, the inhibition of glucose catabolism by 2DG blocked RPC proliferation and eye morphogenesis in human retinal organoids (Takata et al., 2023). Intriguingly, in this organoid study, RPC identity markers, including RAX, and SIX3, were also suppressed by 2DG, a phenotype attributed to lactate depletion, which indicates that glycolysis is also critical for progenitor cell identity (Takata et al., 2023). One possibility is that the "energy budget" of a cell, dictated by adenosine triphosphate (ATP) and precursor metabolite levels, controls levels of transcription of retinal cell identity genes (Carthew, 2021). Glycolytic metabolites can also impact gene transcription in other ways. For example, lactate, a metabolic byproduct of glycolysis, inhibits histone deacetylase (HDAC) activity to acetylate H3K27ac and activate eye development genes in retinal organoids (Takata et al., 2023). Indeed, the addition of lactate to retinal organoids induced broad transcriptional changes, including an upregulation of genes involved in eye morphogenesis (Takata et al., 2023). Mechanistically, lactate effects on the transcription of eye field markers were attributed to the inhibition of HDACs, which normally inhibit gene expression via histone deacetylation (Takata et al., 2023). In the future, we can further assess the impact of 2DG and other metabolic inhibitors on RPC differentiation, focusing on additional retinal cell types, such as bipolar, horizontal and amacrine cell interneurons, and Müller glia.

The regulation and role of pHi and Wnt signaling in cell fate decisions.

A proxy measure for glycolytic activity are the levels of sentinel metabolites (Kochanowski et al., 2013; Miyazawa et al., 2022; Oginuma et al., 2020; Peeters et al., 2017;

Zhang et al., 2017). In chick embryonic tailbuds (Oginuma et al., 2020) and the murine

presomitic mesoderm (Miyazawa et al., 2022), FBP was identified as a sentinel metabolite since

its levels positively correlate with glycolytic activity. However, while FBP has been ascribed a

major role in controlling mouse embryonic mesoderm development (Miyazawa et al., 2022),

neuromesodermal fate selection in the chick embryonic tailbud is controlled by elevated pHi

induced by high glycolytic activity (Oginuma et al., 2020). In the chick tailbud, high pHi creates

an environment that activates Wnt signaling through non-enzymatic acetylation of β-catenin

(Oginuma et al., 2020). Thus, while phosphorylated and non-phosphorylated β-catenin levels do

not vary, K49 β-catenin acetylation is reduced at low pHi (Oginuma et al., 2020). The onset of

K49 β-catenin acetylation is associated with a switch in cell fate choice from neural to

mesodermal in the chick tailbud (Oginuma et al., 2020). In contrast, in adult cardiomyocytes,

when *Pkm*, a rate-limiting glycolytic enzyme, is mutated, β-catenin can more efficiently shuttle

to the nucleus to activate Wnt-signaling and promote cardiomyocyte proliferation (Hauck et al.,

2021). In cardiomyocytes lacking *Pkm*, transcriptional targets of Wnt signaling, including *Axin2,*

Ccnd1, Myc, Sox2 and *Tnnt3,* are elevated, and β-catenin protein remains in the cytoplasm,

rather than undergoing nuclear translocation (Hauck et al., 2021). Similarly, in a study performed

in cancer cells, high pH leads to increased expression of *Ccnd1*, a β-catenin target gene, and

increases proliferation rates (Koch et al., 2020). These findings are consistent with our

demonstration that β-catenin levels are stabilized at pH 8, and RPC proliferation is enhanced.

Whether pH manipulations also influence cell fate decisions by influencing the stability of other

retinal proteins, such as PKM2, can be further investigated in the future using specific PKM2

inhibitors, such as Shikonin (Chen et al., 2011). Finally, in a separate study performed in cancer

cells, acidification induced by metformin indirectly suppresses Wnt signaling by activating the

DDIT3 transcriptional repressor (Melnik et al., 2018), consistent with our data showing low pH

suppresses β-catenin stability. Melnik et al. (2018) also used Mcl inhibitors, as we did in our

study, and showed that this treatment blocks Wnt signaling. While we did not examine the

impact of CNCn on Wnt signaling, we did observe a decline in proliferation, as expected if Wnt

levels are low.

Our finding that Wnt signaling is required to maintain RPC proliferation and drive

photoreceptor differentiation is contrary to prior studies that ascribed a role to *Ctnnb1* in

maintaining RPCs in an immature, uncommitted state without affecting proliferation (Ouchi et

al., 2011). A major difference is that Ouchi et al. (2011) found that *Id3*, a neurogenic inhibitor, is

upregulated in *Ctnnb1* cKO retinas, whereas we found that *Id3* expression levels decline in *Pten*-

cKO retinas. It may therefore be that the precocious photoreceptor differentiation we observed in

the absence of *Pten* occurs due to the reduction in inhibitory *Id3* levels. There may also be a

biphasic role for Wnt signaling, as has been observed during neocortical development, where

canonical Wnt signaling first induces the symmetric proliferative divisions of neural progenitor

cells and later induces differentiation (Harrison-Uy & Pleasure, 2012). Finally, the ectopic

expression of *Wnt2b* inhibits neuronal differentiation in the chick retina (Kubo et al., 2005),

phenocopying the postnatal *Pten*-cKO retinal phenotype observed in this study, in which Wnt

signaling is also elevated. However, while ectopic *Wnt2b* expression suppresses chick RPC

differentiation by inhibiting the expression of proneural genes, we did not observe a significant

impact on Notch signaling or proneural gene expression in *Pten*-cKO retinas (data not shown).

Thus, the gain of Wnt signaling in *Pten*-cKO retinas impacts RPC proliferation and

differentiation possibly in a proneural gene-independent manner.

The link between glycolysis and Wnt signaling may be species- and lineage-specific. For

instance, in mouse embryos, Miyazawa et al., found that increasing glycolysis in the presomitic

mesoderm inhibits Wnt signaling (Miyazawa et al., 2022). Furthermore, in a study performed in

Madin Darby Canine Kidney (MDCK) cells, higher pH levels were shown to decrease β-catenin

levels in the cytoplasm, nucleus and junctional complexes (Czowski & White, 2024). In this

study, the authors altered pH using inhibitors for a sodium-proton exchanger and a sodium

bicarbonate transporter, in contrast to the Oginuma et al. (2020) study, which instead used the

ionophores nigericin and valinomycin to equilibrate intracellular pHi to media pH. Consistent

with the idea that there is some species specificity, in human retinal organoids, lactate itself was

shown to alter optic vesicle bud gene expression, whereas pH did not appear to be an important

driver of cell fate decisions (Takata et al., 2023). In contrast, in the mouse retina, the GPR81

receptor for lactate is expressed at the highest levels in ganglion cells in which it is required to

promote axon growth, with only low, possibly background expression levels in RPCs and

photoreceptors (Laroche et al., 2021). In the future, to more comprehensively examine the link

between *Pten* loss, glycolytic activity, pHi and Wnt signaling, we could examine levels of

phosphorylated, non-phosphorylated and K49 acetylated β-catenin after each manipulation (*i.e.,*

Pten loss, pH manipulations, CNCn treatment, glycolysis inhibition, acetate treatments). By

collecting CD133⁺ MACS-enriched RPCs from the retinal explants, we can add cell type and

stage specificity to our analysis. Finally, the use of qPCR to examine Wnt signaling genes, such

as *Axin2, Ccnd1, Myc, Sox2* and *Tnnt3*, would provide further support for the impact of

glycolysis and pH on this signaling pathway. Lastly, an important caveat to our study is that

metabolism changes ex vivo versus in vivo, and thus, in the future, in vivo studies can be

performed to assess metabolic changes.

Role of *Pten* in photoreceptor differentiation and maturation

Our finding that *Pten* loss drives photoreceptor differentiation and OS maturation was an

unexpected fate bias given that during development, *Pten* is expressed in multipotent RPCs

(Tachibana et al., 2016) that give rise to all retinal cells in a defined temporal sequence (Javed &

Cayouette, 2017). RPCs that are biased to a generic photoreceptor fate express the transcription

factor (TF) Olig2; if these cells later express the TF Nrl, they become rods, if not, they become

cones (Hafler et al., 2012; Javed & Cayouette, 2017). In *Pten*-cKO retinas, more RPCs become

photoreceptors instead of differentiating into other retinal cell types at early developmental

stages (Tachibana et al., 2016), but this phenotype is not likely to be dependent on Olig2, since

Olig2 is expressed at normal levels from the RNA-seq study performed herein. It is possible that

glycolytic inhibition with 2DG slows down the development and production of most newly

differentiated cells rather than specifically affecting photoreceptor differentiation, which we

could assess in the future. Nevertheless, the finding that *Pten* loss elicits a photoreceptor bias in

early RPCs could have widespread applications in targeting uncurable retinal diseases. Notably,

the pharmacological inhibition of PTEN has been used in pre-clinical studies in lung, muscle,

and nervous systems for regenerative purposes (Borges et al., 2020; Knafo & Esteban, 2017),

including in the retina, to promote ganglion cell axonogenesis (Park et al., 2008).

In summary, we have revealed that elevated glycolysis is a key contributor to the RPC

fate alterations observed in *Pten*-cKO retinas. In particular, we have found that glycolytic flux is

elevated in *Pten*-cKO retinas, which modulates both pHi and Wnt signaling to control the timing

of RPC proliferation and differentiation.

MATERIALS AND METHODS

Mice. All animal procedures were approved by the Sunnybrook Research Institute Animal Care

Committee (16-606) in agreement with the Guidelines of the Canadian Council of Animal Care

(CCAC). The following transgenic animal lines were used from Jackson Laboratory: *Pten*^{fl} allele

(Backman et al., 2001) (B6.129S4-*Pten*^{tm1Hwu}/J. Strain #: 006440, RRID:IMSR_JAX:006440),

Ctnnb1^{fl}: B6(Cg)-Ctnnb1tm1Knw/J. RRID:IMSR_JAX:022775) and *Pax6::Cre* driver

(Marquardt et al., 2001) (STOCK Tg(Pax6-GFP/cre)1Rilm/J. Strain #:024578.

RRID:IMSR_JAX:024578. Common Name: P0-3.9GFPCre) and Ctnnb1^{fl}: B6(Cg)-

Ctnnb1^{tm1Knw}/J. (RRID:IMSR_JAX:022775). For glycolysis gain of function studies,

cyto*PFKB3*- mice (EMBL, Germany) (Miyazawa et al., 2022) were crossed with a *Pax6::Cre*

driver to generate cyto*PFKB3*-GOF transgenic mice. PCR genotyping was performed as

described by Jax for commercial lines and as described for the cyto*PFKB3* allele (Miyazawa et

al., 2022). All lines were maintained on a C57BL/6J background through backcrossing (Strain

#:000664. RRID:IMSR_JAX:000664. Common Name: B6). Animal explants were performed

with CD1 mice (Charles River, IMSR_CRL:022). The day of the vaginal plug was considered to

be embryonic day E0.5 for timed pregnancies.

Bulk RNAseq. P0 *Pten*-cKO and wild-type control eyes were dissected in a clean petri dish

containing Dulbecco's phosphate buffered saline (DPBS). A pair of super-fine tweezers (#5, Fine

Scientific Tools, Canada) was used to remove the cornea, iris, sclera and retinal pigment

epithelium (RPE). Retinas were homogenized using TRIzol (Thermo Fisher Scientific; #15596-

026) according to the manufacturer's protocol. RNA was then purified using an RNAeasy

minikit (74104, Qiagen). Eluted RNA quality and concentration was assessed using a NanoDrop.

Samples with the highest purity (260/280 ratio = 2.0) were used for RNA sequencing (RNAseq)

experiments. Five *Pten* cKO and four wild-type samples were sent to the University of

Lethbridge, Alberta for sequencing. Illumina NextSeq 500 performed sequencing using 75bp

Single-End reads. A NEBNext Ultra II Directional RNA Library Prep Kit was used to construct a

library containing read counts, and was compared to a Mouse GRCM38 reference genome.

Differentially expressed genes (DEGs) were identified using the DESeq1 1.28.1 package. An

adjusted p-value (padj) less than 0.05 for any gene was considered a DEG (Wald test, Benjamini-

Hickberg correction for multiple comparisons). DEGs were visualized by Volcano plot (R

package: EnhancedVolcano 1.6.0) and heatmap (R package: gplots 3.0.3, function heatmap 2).

Gene Ontology (GO) terms for Biological Processes and KEGG (Kyoto Encyclopedia of Genes

and Genomes) pathways were assayed using the DEG up- and down-regulated DEGs (Pathview

1.28.0 package).

In vivo injections. For birthdating and proliferation studies, animals were injected with BrdU at

10mg/kg intraperitoneally at E12.5 and then harvested at P7. For 2DG *in vivo* experiments,

newborn pups were injected with 2-deoxy-d-glucose (2DG) intraperitoneally (30mg/Kg, Sigma

Aldrich, Oakville, Canada) daily starting at P0 until P7.

Tissue processing and immunostaining. Animals were sacrificed using CO_2 euthanasia. Eyes

were dissected and fixed in 4% paraformaldehyde (PFA) overnight at 4ºC. Fixed eyes were

rinsed with PBS and incubated in 20% sucrose overnight. Eyes were then embedded in optical

cutting temperature (O.C.T™) compound (Tissue-Tek®, Sakura Finetek U.S.A. Inc., Torrance,

CA) and frozen on dry ice before being sectioned in a series of 10uM slices using a Leica

CM3050s cryostat (Leica Biosystems, Buffalo Grove, IL, USA). Sections were mounted on

Fisherbrand™ Superfrost™ Plus slides (Thermo Fisher Scientific, Markham, ON). Sections were

blocked for 1 hour in 10% normal horse serum in 1X PBS/0.1% Triton X-100 (PBST). Primary

antibodies were diluted in a blocking solution and incubated at 4ºC overnight. Primary antibodies

included: BrdU (1:400, ab6326, Abcam), VSX2 (1:200, sc-365519, Santa Cruz), Ki67 (1:400,

ab16667, Abcam), cleaved caspase 3 (CC3; 1:400, ab2302, Abcam), Rhodopsin (1:400, MAB

5356, Millipore), PAX6 (1:400, #PRB-278P; Covance Research), PTEN (1:400, 9559S, Cell

Signaling) and CRX (1:400, NBP2-15964. Novus Biologicals). Slides were washed 3 x 10 min

in PBST and then incubated in secondary antibodies diluted in PBST for 1 hour. Secondary

antibodies were conjugated to Alexa568 (1:500, Molecular Probes), Alexa488 (1:500, Molecular

Probes), or Alexa647 (1:500, Molecular Probes). Sections were washed 3 x 10 min washes in

PBST. Sections were counterstained with 4',6-diamidino-2-phenylindole (DAPI, Invitrogen,

D1306) and coverslipped in Aqua-Poly/Mount (Polysciences #18606).

Retinal explant assay. For retinal explants, P0 wild-type CD1 mice pups were sacrificed by
decapitation. Eyes were dissected in ice-cold PBS. Corneas were removed first, followed by
removal of the sclera, iris, lens and RPE. Retinal cups were then mounted on Cytiva Whatman™
90 mm Nucleopore™ Polycarbonate Track-Etched Membranes (Catalog Number 09920022,
Cytiva, Canada). Four cuts were made to flatten cups on the membranes using TMS101 Gills-
Vannas Scissors (SKU: TMS101, Titan Medical, Coral Gables, FL, USA). Retinas mounted on
Nucleopore membranes were then transferred to 24 well plates (Corning® Costar® TC-Treated
Multiple Well Plates, Catalog Number: CLS3524, Millipore Sigma) filled with 1 mL of retinal
explant media: 50% DMEM-high glucose (25 mM), no glutamine, no methionine, no cystine
(Gibco, REF 21013-024), 25% HBSS (Cat 311-512-CL), 25% heat-inactivated horse serum
(Wisent, Cat 065-150), 200 µM l-glutamine (Wisent, Cat 609-065-EL), 0.6 mm HEPES (Wisent,
Cat 330-050-CL), 1% Pen-Strep (Wisent, Cat 450-201-EL). Retinas were incubated at 37ºC, 5%
CO_2. At the experimental endpoint (as indicated), explants were fixed at room temperature with
4% PFA for 15 min, rinsed with DEPC-treated PBS and incubated in 20% DEPC-Sucrose
overnight. Explants were then mounted in OCT on dry ice and cryosectioned at 10 µM as
described above.

Retinal explant drug treatments. For BrdU labeling, 5 µL of 10 mg/mL BrdU was added to the
retinal cell explant media for the time period described. All inhibitors were incubated with retinal
explants for 24 hours. 2DG (D3179-1G, Sigma Aldrich, Canada) was diluted in PBS and added
to retinal explant media to achieve 5 mM and 10 mM final concentrations. 10 mM 2DG was
previously shown to reduce lactate levels in tumor cells (Malm et al., 2015), and we confirmed
the efficacy of our 2DG batch on reducing lactate levels in the retina when administered in vivo
(Hanna et al., 2025). Glycogen phosphorylase inhibitor (GPI) (361515-1MG, Sigma Aldrich,
Canada) was diluted in DMSO and added to the retinal explant media at a final concentration of
12.5 µM or 25 µM for 24 hr. α-cyano-4-hydroxy-cannamic acid (CNCn; C2020-10G; Sigma
Aldrich, Canada) was diluted in explant media and added to explant media at a final
concentration of 5 mM for 24 hr. For pH experiments, pH was adjusted after incubating the
media in cell culture incubator for 1 hour using hydrochloric acid or sodium hydroxide to reach a

pH of 6.5 and 8.0, respectively, and grown for 24 hr. FH535 was diluted in DMSO and added to

the retinal explant media at 5mM and 10mM concentrations for 24 hours (108409, Millipore

Sigma, Canada). For acetate explants, retinas were incubated with sodium acetate at a

concentration of 5 mM, 10 mM or 20 mM diluted in PBS for 48 hours (S2889, Millipore Sigma,

Canada).

Retinal explant electroporation. P2 wild-type retinal explants were electroporated with pCAGG-

pHluorin-IRES2-tdTomato obtained from Olivier Pourquié (Oginuma et al., 2020) as described

(Tachibana et al., 2016). Briefly, retinas were placed in 2 mm gap cuvettes (VWR International)

with 30ul of DNA at 0.5 μg/ μl. Retinas were then electroporated using an ECM830 pulse

generator (BTX Harvard Apparatus) using five square 20 V pulses of 50 ms duration, separated

by 950 ms intervals. After electroporation, retinas were cultured on Nuclepore™ Polycarbonate

Track-Etched Membranes in retinal explant media at 37ºC, 5% CO_2.

qPCR. TRIzol® RNA Isolation Reagent (Thermo Fisher Scientific; #15596-026) was used to

extract RNA according to the manufacturer's instructions. cDNA was synthesized from 0.5 μg

RNA using an RT^2 first-strand kit (Qiagen, #330401) according to the manufacturer's

instructions. qPCR was performed using an RT^2 SYBR® Green PCR Master Mix (Qiagen;

#330500) as described by the manufacturer, with a CFX384 cycler (Biorad Laboratories,

Canada). The following primers were used: *Eno1* (PPM04408B, Qiagen), *Hk1* (PPM05501A,

Qiagen), *Pgk1* (PPM03700A, Qiagen), *Slc16a3* (PPM27328A, Qiagen) and normalized to

housekeeping genes *Hprt* (PPM03559F, Qiagen), *B2m* (PPM03562A, Qiagen), and *Mapk1*

(PPM03571E, Qiagen).

Western blotting. For western blot, retinal cups were dissected on ice and both retinas of the

same animal were pooled together. Retinas were lysed in NP-40 lysis buffer (0.05 M Tris, pH

7.5, 0.15 M NaCl, 1% NP-40, 0.001 M EDTA) with protease (1X protease inhibitor complete, 1

mM PMSF), proteasome (MG132 at 0.05 mM) and phosphatase (50 mM NaF, 1 mM NaOV)

inhibitors. Protein amounts were quantified with a Micro BCA™ Protein Assay Kit

(ThermoFisher Scientific, #23235) following the manufacturer's instructions. 10 μg of lysate

from whole cells or tissue was run on 10% SDS-PAGE gels for western blot analysis. Protein

was transferred to PVDF membrane at 80V for 1 hour. PVDF membranes were then blocked in

5% skim milk powder in tris-buffered saline with 0.1% tween 20 (TBST) for 1 hour at room

temperature. Blots were incubated overnight at 4ºC in a blocking solution with non-

phosphorylated β-Catenin (1:1000, 8814, Cell Signalling) followed by β-actin (1:5000, ab8227,

Abcam). Blots were washed 3 x 10 min in TBST before incubating in secondary antibodies,

including either Goat Anti-Rabbit IgG (H+L)-HRP Conjugate (1/50,000; Bio-Rad #1721019 or

Cell Signaling #7074) or Goat Anti-Mouse IgG (H+L)-HRP Conjugate (1/50,000; Bio-Rad

#1721011). A chemiluminescent signal was obtained from the blots using (ECL Prime Luminor

Enhancer Solution, Cytiva, Canada) and the blot was developed using a ChemiDoc System (Bio-

Rad Laboratories, Canada). Densitometries were calculated using ImageJ (Schneider et al.,

2012), and normalized values (over ß-actin) were plotted.

scRNA-seq analysis. 10x scRNA-seq data of mouse developing retina was obtained from

GSE118614 (Clark et al., 2019). Further processing and analyses were performed with the Seurat

v.3.2.3 R package (Butler et al., 2018). Low-quality cells were excluded by filtering out cells

with fewer than 500 detected genes and cells with mitochondrial RNA more than 5%. The data

was then transformed by the SCTransform function while regressing out the variance due to

mitochondrial RNAs. Clustering was performed by the RunPCA, FindNeighbors, and

FindClusters functions using the first 30 principal components. The 2D projection of the

clustering was carried out by the RunUMAP function. The annotation of cell type to each cluster

was performed by using the same set of markers as in (Clark et al., 2019). Expression of selected

genes was plotted using the FeaturePlot function.

Seahorse assay. P0 *Pten*-cKO and wild-type control pups were sacrificed, and retinal cups were

dissected. Both retinas from the same animal were pooled together in one tube and dissociated

using a Papain kit according to manufacture protocol (Papain Dissociation System, Worthington

Biochemical Corporation, USA). Dissociated cells of each animal were then plated in a 6-well

plate coated with Poly-D-Lysine (10 mg/mL) (Corning, 354210)/Laminin (Corning, 354232)

(2.5 μg/mL) (Sarstedt, Germany). Cells were cultured for 5 days then transferred to Seahorse 96-

well plate at a density of 40,000/well. A Seahorse Mito stress test was run according to the

manufacture protocol. Seahorse basal media was supplemented with Oligomycin, FCCP, R/A at

the indicated time points (Seahorse XF Mito Stress Test Kit, 103015-100, Agilent, USA).

Readings were then normalized to protein concentration/well measured using the Micro BCA™

Protein Assay Kit.

Lactate assay. To measure lactate levels, P0 wild-type, *Pten*-cKO or cyto*PFKFB3*-GOF retinas

were dissected on ice-cold PBS and both retinas of each pup were pooled together in the same

tube. Cell lysates were prepared as described for using NP-40 lysis buffer and protein levels were

quantified using the Micro BCA™ Protein Assay Kit. Measurements of lactate were then

performed using a high sensitivity Lactate Colorimetric/Fluorometric Assay Kit as per the

manufacturer's protocol (K607-100; BioVision; CA, USA) and normalized to protein levels.

Imaging, quantification and figure preparation. Retinal cross sections were imaged using Leica

DM18 inverted fluorescent microscope and Leica Application Suite X (LASX) software for cell

counts (Leica Biosystems, Canada) or a Nikon A1 laser scanning confocal microscope at

Sunnybrook Research Institute Centre for Cytometry and Scanning Microscopy (Nikon,

Canada). OS measurements were performed by imaging three different retinal cross sections

stained with rhodopsin (n=3) of each of three different biological replicates (N=3) for each

animal group. Outer segments were then outlined using ImageJ and the total area was calculated.

Licenses for Adobe Photoshop 2021 and BioRender were used to prepare figures and schematics.

Statistical analysis. Statistical analysis was performed on at least three different animals/explants

(N; biological replicates). Cell counts were performed by counting cells in three different cross

sections (n; technical replicates). For comparisons between two groups, Unpaired two-tailed

Student's t-test was used (GraphPad Prism, USA). For comparisons involving more than two

groups, one-way ANOVA with post-hoc Tukey multiple comparisons of means was used

(GraphPad Prism, USA). Results were plotted on graphs with standard error of the mean (SEM)

error bars (GraphPad Prism, USA). A p-value less than 0.05 was considered statistically
significant.

ACKNOWLEDGEMENTS AND FUNDING SOURCES

We would like to thank Alexander Aulehla for providing the *cytoPFKFB3* transgenic mice. We
thank Olivier Pourquié for providing reagents, Robert Cantrup in the Schuurmans' lab and Petia
Stefanova in the SRI Histology Facility for help with sectioning and immunostaining, and Cathy
Ioria in RS lab for help with the Seahorse assay. This work was supported by the Canadian
Institutes of Health Research (CIHR) PJT 180243 to CS, RS, YT and IK, and PJT – 400973 to
IA and CS. JH was supported by a Canada Graduate Scholarship – Doctoral (CGS-D)/CIHR
award, Vision Science Research Program Scholarship, Ontario Graduate Scholarship and R.O.
Torrance Bursary. CS holds the Dixon Family Chair in Ophthalmology Research at the
Sunnybrook Research Institute.

FIGURE LEGENDS

Figure 1. Glycolysis expression during development and accelerated development in *Pten*-
cKO retinas.

(A) Schematic of the key enzymatic steps in glycolysis.

(B) UMAP plot of scRNA-seq data collected from wild-type retinas between E11 and P14 (Clark
et al., 2019). Stages of data collection are color-coded in the first panel, and cell type annotation
is color coded in the second panel. Pre-neurogenic (N)-RPC, early RPC and late RPC clusters are
labeled.

(C) Feature plots showing the transcript distribution of glycolytic genes, including *Pgk1, Eno1,*
Gpi1, Hk1, Pfkl, Slc16a3, Aldoa, and *Ldhb*, showing enriched *Pgk1, Eno1* and *Hk1* expression in
earlier-staged RPCs. Pre-neurogenic (N)-RPC, early RPC and late RPC clusters are labeled.

(D) E18.5 retinal cross sections immunostained with PAX6, PTEN, and VSX2.

(E) Schematic of the strategy used to generate *Pten*-cKO animals.

(F) Immunostaining of P0 Pax6-Cre-GFP and *Pten*-cKO retinas, showing GFP expression in

RPCs throughout the retina, with higher levels in the periphery in the ciliary marginal zone

(CMZ).

(G) Retinal cross sections of wild-type and *Pten*-cKO animals at P14 injected with BrdU at P4,

showing immunostaining with PTEN (green) and BrdU (red) in wild-type and *Pten*-cKO retinas.

(H) Quantification of the number of BrdU cells in the gcl, inl and onl. Plots show means ± SEM.

N=3 biological replicates/genotype, all with 3 technical replicates. P-value calculated with

unpaired t-test. onl, outer nuclear layer; inl, inner nuclear layer; gcl, ganglion cell layer.

onbl, outer neuroblast layer; inl, inner nuclear layer; gcl, ganglion cell layer; le, lens; cmz, ciliary

marginal zone. Scale bar 50μM in C, 400μM in E and 100μM in F.

Figure 1– Figure supplement 1. Expression of *Pten* and mature photoreceptor markers

during murine retinal development.

UMAP plot of scRNA-seq data collected from wild-type retinas between E11 and P14 (Clark et

al., 2019). Transcript distribution of *Pten*, and mature photoreceptor markers (*Prph2, Impg2,*

Pde6g, Gm11744, Gnat2) at different stages of retinal development.

Figure 2. Glycolytic gene expression and activity are elevated in P0 *Pten*-cKO retinas.

(A) Heat map of bulk RNAseq data comparing gene expression in P0 *Pten*-cKO (N=5) and wild-

type (N=4) retinas.

(B) Volcano plot of DEGs with Log2FC=2, showing both down-regulated (to the left) and up-

regulated (to the right) genes. Venn diagram shows 667 up-regulated genes and 408 down-

regulated genes in P0 *Pten*-cKO retinas.

(C) KEGG pathway enrichment related to DEGs, showing an enrichment of up-regulated

glycolytic pathway genes in P0 *Pten*-cKO retinas (arrow).

(D) Schematic of glycolysis pathway, showing the intracellular enzymes involved in

metabolizing glucose into lactate. Also shown is the Slc16a3 symporter, which extrudes lactate

and H^+ to increase intracellular pHi.

(E) Normalized RPKM (reads per kilobase million) values for glycolytic gene expression in P0 wild-type and *Pten*-cKO retinas. Plots show means ± SEM. N=4 biological replicates for wild-type and N=5 for *Pten*-cKO retinas. P-values calculated with Wald test, with a Benjamini-Hickberg correction for multiple comparisons.

(F) Normalized glycolytic gene expression from qPCR of P0 *Pten*-cKO (N=3) and wild-type control (N=3) retinas. Plots show means ± SEM. N=3 biological replicates/genotype, all with 3 technical replicates. P-values calculated with unpaired t-test.

(G) Seahorse assay on P0 wild-type and *Pten*-cKO RPCs cultured in vitro. Points of treatment with oligomycin (O), FCCP (F), and Rotenone/Antimycin A (R/A) are indicated. Left plot shows elevated extracellular acidification rate (ECAR) in *Pten*-cKO RPCs, while plot to the right shows no significant effect on oxygen consumption rate (OCR). Plots show means ± SEM. N=3 biological replicates for *Pten*-cKO and N=8 biological replicates for wild-type, all with 3 technical replicates. P-values calculated with unpaired t-test (GraphPad Prism, USA).

(H) Lactate assay on P0 wild-type and *Pten*-cKO retinas. Plots show means ± SEM. N=3 biological replicates/genotype, all with 3 technical replicates. P-value calculated with unpaired t-test.

Figure 3. Glycolytic inhibition reduces proliferation and photoreceptor differentiation in P0 retinal explants.

(A) Feature plots showing the transcript distribution of *Cdk1*, a marker of proliferating RPCs, and the genes encoding the major glucose transporters, *Slc2a1* (GLUT1), *Slc2a3* (GLUT3) and *Slc2a4* (GLUT4).

(B) Schematic of retinal explant experiments, showing treatment with pharmacological inhibitors (2DG or GPI), and the labeling of proliferating RPCs (30 min BrdU) or birthdating of newborn retinal cells (24 hr BrdU).

(C) Schematic of glucose uptake replaced by 2DG, which leads to the accumulation of 2-DG-6P that cannot be further metabolized by the glycolytic pathway.

(D) P0 retinal explants treated with 5mM or 10mM 2DG for 1 day in vitro (DIV), showing BrdU immunolabeling and the number of BrdU[+] cells after a 30 min labeling with BrdU.

(E) P0 retinal explants treated with 5mM or 10 mM 2DG for 1 day in vitro (DIV), showing BrdU

immunolabeling and the number of BrdU$^+$ cells after a 24-hr labeling with BrdU, including co-

labeling with Ki67 to calculate the p-fraction (%BrdU$^+$Ki67$^+$cells/total Ki67$^+$ cells).

(F-H) P0 retinal explants treated with 5 mM or 10 mM 2DG for 1 day in vitro (DIV) and

immunostained for Vsx2 (E), CC3 (F) or CRX (G).

(I) P0 retinal explants treated with glycogen phosphorylase inhibitor (GPI) at 12.5 μM and 25

μM for 24 hr and immunostained for BrdU after a 30 min incubation at the end of the treatment

period.

Plots in D-I show means ± SEM. N=3 biological replicates/treatment, all with 3 technical

replicates. P-values calculated with one-way ANOVA with Tukey post hoc test. Scale bars= 25

μM in D,F,H and 50 μM in E,G,I.

Figure 4. Promotion of glycolytic flux with a cyto*PFKB3*-GOF mouse model accelerates

photoreceptor differentiation and outer segment maturation.

(A) Schematic showing function of PFKB3 as an activator of PFK1 through the conversion of

fructose-6-phosphate to fructose-2,6-bisphosphate.

(B) Schematic showing generation of an RPC-specific cyto*PFKB3*-GOF mouse model.

(C) Lactate assay performed on P0 wild-type and cyto*PFKB3*-GOF retinas.

(D,E) Birthdating experiments performed by injecting BrdU into pregnant females at E12.5 and

harvesting retinas at P7. BrdU immunostaining of P7 wild-type and cyto*PFKB3*-GOF retinas (D)

and quantification of the %BrdU$^+$ cells in each of the nuclear layers (E).

(F) Rhodopsin immunostaining of P7 wild-type and cyto*PFKB3*-GOF retinas. An expansion of

outer segment (OS) area is indicated by the arrow and quantified in the graph.

(G) Bulk RNAseq mining of photoreceptor outer segment genes de-regulated in P0 *Pten*-cKO

retinas.

(H) Schematic of mode of action of 2DG.

(**I**) Rhodopsin immunostaining of P7 wild-type and *Pten*-cKO retinas, either without treatment or

following the administration of 2DG between P0-P7. The expanded OS area in *Pten*-cKO retinas

is indicated with an arrow. Quantifications of photoreceptor OS areas are shown in the graph.

Plots show means ± SEM. N=3 biological replicates/treatment group except C, with

cytoPFKFB3-GOF (N=3) and wild-type (N=2), all with 3 technical replicates. P-values

calculated with t-tests in C and F, and one-way ANOVA with Tukey post hoc test in E and G.

Scale bars 100 μM in E and 50 μM in F and G. gcl, ganglion cell layer; inl, inner nuclear layer;

onl, outer nuclear layer; os, outer segments.

Figure 5. Reducing intracellular pH (pHi) inhibits RPC proliferation and differentiation.

(**A**) Schematic showing function of Slc16a1/3 symporters in extruding lactate and H^+ out of the

cell to increase pHi during elevated glycolysis. Also depicted is the impact of acidic pH and

oligomerizing PKM to prevent its activity.

(**B**) Schematic of retinal explant experiments, showing treatment with a

pharmacological inhibitor (CNCn) of Slc16a symporters or growth in different pH media, and

the labeling of proliferating RPCs (30 min BrdU) or birthdating of newborn retinal cells (24 hr

BrdU).

(**C**) P0 retinal explants treated with 5mM CNCn for 1 day in vitro (DIV) followed by a 30 min

BrdU pulse, showing immunolabeling and the quantification of BrdU$^+$ RPCs.

(**D**) P0 retinal explants incubated in media buffered to pH6.5, pH7.4 or pH8.0 for 1 day in vitro

(DIV), showing BrdU immunolabeling and the number of BrdU$^+$ cells after a 30 min labeling

with BrdU.

(**E**) P0 retinal explants incubated in media buffered to pH6.5, pH7.4 or pH8.0 for 1 DIV,

showing BrdU immunolabeling and the number of BrdU$^+$ cells after a 24-hr labeling with BrdU,

including co-labeling with Ki67 to calculate the p-fraction (%BrdU$^+$Ki67$^+$cells/total Ki67$^+$ cells).

(**F-H**) P0 retinal explants incubated in media buffered to pH6.5, pH7.4 or pH8.0 for 1 DIV and

immunostained for Vsx2 (E), CC3 (F) or CRX (G).

(**I**) P3 retinal explant electroporated with pHluorin-tdTomato, imaged after 2DIV to show

electroporated cells (tdTomato$^+$) have varying levels of 488nm emission from pHLuorin (green).

Plots show means ± SEM. N=3 biological replicates/treatment group, all with 3 technical

replicates. P-values calculated with t-tests in C, and one-way ANOVA with Tukey post hoc test

in D,E,F-H. Scale bars= 25 µM in C, D, F and H, and 50 µM in E, G and I.

Figure 6. Wnt signalling is a downstream mediator of Pten and glycolysis.

(A) Schematic illustration of canonical Wnt signaling pathway.

(B) UMAP plot of scRNA-seq data collected from wild-type retinas between E11 and P14 (Clark

et al., 2019). Transcript distribution of *Wnt7b*, *Lef1* and *Dkk1* Wnt genes.

(C) Normalized RPKM (reads per kilobase million) values for Wnt pathway-associated genes in

P0 wild-type (N=4) and *Pten*-cKO (N=5) retinas.

(D) Western blot of β-catenin expression, normalized to β-actin levels, in P0 retinal explants

treated with 5mM and 10mM 2DG or in P0 retinal explants cultured in media buffered to pH8.0,

pH7.4 or pH6.5 **(E)**, all for 24 hrs.

Plots show means ± SEM. N=3 biological replicates/treatment group, all with 3 technical

replicates. P-values calculated with t-tests in C, and one-way ANOVA with Tukey post hoc test

in D,E.

Figure 6 – Figure supplement 1. Upregulation of Wnt-associated genes in *Pten* cKO retinas.
PANTHER (protein analysis through evolutionary relationships) classification analysis of DEGs
of P0 bulk RNAseq showing upregulation of Wnt signaling and glycolysis (arrows).

Figure 7. Wnt signaling modulate retinal proliferation and differentiation.

(A-C) P0 retinal explants treated with 5µM or 10µM FH535 to block Wnt signaling for 1 day in

vitro (DIV), followed by immunostaining for BrdU after a 30 min pulse (A), CRX (B) or CC3

(C).

(D) Immunostaining P14 wild-type and *Ctnnb1*-cKO retinas with rhodopsin. Asterisk points to

ONL rosette in *Ctnnb1*-cKO retina. Graph showing quantification of OS area.

(**E**) Retinal cross sections of P0 wild-type explants treated with 5mM, 10mM and 20mM sodium acetate for 1 day in vitro, along with vehicle controls, and pulsed with BrdU 30 minutes before harvesting. The number of BrdU$^+$ RPCs increased in sodium acetate treated explants.

All experiments included three biological replicates for each group. Statistical analysis was done using one-way ANOVA with Tukey post hoc test used in A-C and unpaired *t*-test in D. Scale bars= 50 μM in A-D. gcl, ganglion cell layer; inl, inner nuclear layer; onl, outer nuclear layer; os, outer segments.

REFERENCES

Agathocleous, M., Love, N. K., Randlett, O., Harris, J. J., Liu, J., Murray, A. J., & Harris, W. A. (2012). Metabolic differentiation in the embryonic retina. *Nat Cell Biol*, *14*(8), 859-864. https://doi.org/10.1038/ncb2531

Alexiades, M. R., & Cepko, C. L. (1997). Subsets of retinal progenitors display temporally regulated and distinct biases in the fates of their progeny. *Development*, *124*(6), 1119-1131. https://doi.org/10.1242/dev.124.6.1119

Backman, S. A., Stambolic, V., Suzuki, A., Haight, J., Elia, A., Pretorius, J., Tsao, M. S., Shannon, P., Bolon, B., Ivy, G. O., & Mak, T. W. (2001). Deletion of Pten in mouse brain causes seizures, ataxia and defects in soma size resembling Lhermitte-Duclos disease. *Nat Genet*, *29*(4), 396-403. https://doi.org/10.1038/ng782

Baye, L. M., & Link, B. A. (2007). Interkinetic nuclear migration and the selection of neurogenic cell divisions during vertebrate retinogenesis. *J Neurosci*, *27*(38), 10143-10152. https://doi.org/10.1523/JNEUROSCI.2754-07.2007

Borges, G. A., Webber, L. P., AE, M. M., Guerra, E. N., Castilho, R. M., & Squarize, C. H. (2020). Pharmacological PTEN inhibition: potential clinical applications and effects in tissue regeneration. *Regen Med*, *15*(2), 1329-1344. https://doi.org/10.2217/rme-2019-0065

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., & Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*, *36*(5), 411-420. https://doi.org/10.1038/nbt.4096

Cantrup, R., Dixit, R., Palmesino, E., Bonfield, S., Shaker, T., Tachibana, N., Zinyk, D., Dalesman, S., Yamakawa, K., Stell, W. K., Wong, R. O., Reese, B. E., Kania, A., Sauve, Y., & Schuurmans, C. (2012). Cell-type specific roles for PTEN in establishing a functional retinal architecture. *PLoS One*, *7*(3), e32795. https://doi.org/10.1371/journal.pone.0032795

Carthew, R. W. (2021). Gene Regulation and Cellular Metabolism: An Essential Partnership. *Trends Genet*, *37*(4), 389-400. https://doi.org/10.1016/j.tig.2020.09.018

Cepko, C. (2014). Intrinsically different retinal progenitor cells produce specific types of progeny. *Nature Reviews Neuroscience*, *15*(9), 615-627. https://doi.org/10.1038/nrn3767

Cereghetti, G., Kissling, V. M., Koch, L. M., Arm, A., Schmidt, C. C., Thüringer, Y., Zamboni, N., Afanasyev, P., Linsenmeier, M., Eichmann, C., Kroschwald, S., Zhou, J., Cao, Y., Pfizenmaier, D. M., Wiegand, T., Cadalbert, R., Gupta, G., Boehringer, D., Knowles, T. P. J., Mezzenga, R., Arosio, P., Riek, R., & Peter, M. (2024). An evolutionarily conserved mechanism controls

1003   reversible amyloids of pyruvate kinase via pH-sensing regions. *Dev Cell*.
1004   https://doi.org/10.1016/j.devcel.2024.04.018

1005 Chen, J., Xie, J., Jiang, Z., Wang, B., Wang, Y., & Hu, X. (2011). Shikonin and its analogs inhibit cancer
1006   cell glycolysis by targeting tumor pyruvate kinase-M2. *Oncogene*, *30*(42), 4297-4306.
1007   https://doi.org/10.1038/onc.2011.137

1008 Chinchore, Y., Begaj, T., Wu, D., Drokhlyansky, E., & Cepko, C. L. (2017). Glycolytic reliance promotes
1009   anabolism in photoreceptors. *Elife*, *6*. https://doi.org/10.7554/eLife.25946

1010 Clark, B. S., Stein-O'Brien, G. L., Shiau, F., Cannon, G. H., Davis-Marcisak, E., Sherman, T., Santiago,
1011   C. P., Hoang, T. V., Rajaii, F., James-Esposito, R. E., Gronostajski, R. M., Fertig, E. J., Goff, L.
1012   A., & Blackshaw, S. (2019). Single-Cell RNA-Seq Analysis of Retinal Development Identifies
1013   NFI Factors as Regulating Mitotic Exit and Late-Born Cell Specification. *Neuron*, *102*(6), 1111-
1014   1126 e1115. https://doi.org/10.1016/j.neuron.2019.04.010

1015 Czowski, B. J., & White, K. A. (2024). Intracellular pH regulates β-catenin with low pHi increasing
1016   adhesion and signaling functions. *bioRxiv*. https://doi.org/10.1101/2024.03.22.586349

1017 Datta, P., Allamargot, C., Hudson, J. S., Andersen, E. K., Bhattarai, S., Drack, A. V., Sheffield, V. C., &
1018   Seo, S. (2015). Accumulation of non-outer segment proteins in the outer segment underlies
1019   photoreceptor degeneration in Bardet-Biedl syndrome. *Proc Natl Acad Sci U S A*, *112*(32),
1020   E4400-4409. https://doi.org/10.1073/pnas.1510111112

1021 Esteban-Martinez, L., Sierra-Filardi, E., McGreal, R. S., Salazar-Roa, M., Marino, G., Seco, E., Durand,
1022   S., Enot, D., Grana, O., Malumbres, M., Cvekl, A., Cuervo, A. M., Kroemer, G., & Boya, P.
1023   (2017). Programmed mitophagy is essential for the glycolytic switch during cell differentiation.
1024   *EMBO J*, *36*(12), 1688-1706. https://doi.org/10.15252/embj.201695916

1025 Fekete, D. M., Perez-Miguelsanz, J., Ryder, E. F., & Cepko, C. L. (1994). Clonal analysis in the chicken
1026   retina reveals tangential dispersion of clonally related cells. *Dev Biol*, *166*(2), 666-682.
1027   https://doi.org/10.1006/dbio.1994.1346

1028 Gomes, F. L., Zhang, G., Carbonell, F., Correa, J. A., Harris, W. A., Simons, B. D., & Cayouette, M.
1029   (2011). Reconstruction of rat retinal progenitor cell lineages in vitro reveals a surprising degree of
1030   stochasticity in cell fate decisions. *Development*, *138*(2), 227-235.
1031   https://doi.org/10.1242/dev.059683

1032 Hafler, B. P., Surzenko, N., Beier, K. T., Punzo, C., Trimarchi, J. M., Kong, J. H., & Cepko, C. L. (2012).
1033   Transcription factor Olig2 defines subpopulations of retinal progenitor cells biased toward
1034   specific cell fates. *Proc Natl Acad Sci U S A*, *109*(20), 7882-7887.
1035   https://doi.org/10.1073/pnas.1203138109

1036 Handeli, S., & Simon, J. A. (2008). A small-molecule inhibitor of Tcf/beta-catenin signaling down-
1037   regulates PPARgamma and PPARdelta activities. *Mol Cancer Ther*, *7*(3), 521-529.
1038   https://doi.org/10.1158/1535-7163.Mct-07-2063

1039 Hanna, J., David, L. A., Touahri, Y., Fleming, T., Screaton, R. A., & Schuurmans, C. (2022). Beyond
1040   Genetics: The Role of Metabolism in Photoreceptor Survival, Development and Repair. *Front
1041   Cell Dev Biol*, *10*, 887764. https://doi.org/10.3389/fcell.2022.887764

1042 Hanna, J., Touahri, Y., Pak, A., David, L. A., van Oosten, E., Dixit, R., Vecchio, L. M., Mehta, D. N.,
1043   Minamisono, R., Aubert, I., & Schuurmans, C. (2025). Pten Loss Triggers Progressive
1044   Photoreceptor Degeneration in an mTORC1-Independent Manner. *Invest Ophthalmol Vis Sci*,
1045   *66*(3), 45. https://doi.org/10.1167/iovs.66.3.45

1046 Harrison-Uy, S. J., & Pleasure, S. J. (2012). Wnt signaling and forebrain development. *Cold Spring Harb
1047   Perspect Biol*, *4*(7), a008094. https://doi.org/10.1101/cshperspect.a008094

1048 Hauck, L., Dadson, K., Chauhan, S., Grothe, D., & Billia, F. (2021). Inhibiting the Pkm2/b-catenin axis
1049   drives in vivo replication of adult cardiomyocytes following experimental MI. *Cell Death Differ*,
1050   *28*(4), 1398-1417. https://doi.org/10.1038/s41418-020-00669-9

1051 He, J., Zhang, G., Almeida, Alexandra D., Cayouette, M., Simons, Benjamin D., & Harris, William A.
1052   (2012). How Variable Clones Build an Invariant Retina. *Neuron*, *75*(5), 786-798.
1053   https://doi.org/https://doi.org/10.1016/j.neuron.2012.06.033

Holt, C. E., Bertsch, T. W., Ellis, H. M., & Harris, W. A. (1988). Cellular determination in the Xenopus
retina is independent of lineage and birth date. *Neuron*, *1*(1), 15-26. https://doi.org/10.1016/0896-
6273(88)90205-x
Hurley, J. B., Lindsay, K. J., & Du, J. (2015). Glucose, lactate, and shuttling of metabolites in vertebrate
retinas. *J Neurosci Res*, *93*(7), 1079-1092. https://doi.org/10.1002/jnr.23583
Javed, A., & Cayouette, M. (2017). Temporal Progression of Retinal Progenitor Cell Identity:
Implications in Cell Replacement Therapies. *Front Neural Circuits*, *11*, 105.
https://doi.org/10.3389/fncir.2017.00105
Jensen, A. M., & Raff, M. C. (1997). Continuous observation of multipotential retinal progenitor cells in
clonal density culture. *Dev Biol*, *188*(2), 267-279. https://doi.org/10.1006/dbio.1997.8645
Jo, H. S., Kang, K. H., Joe, C. O., & Kim, J. W. (2012). Pten coordinates retinal neurogenesis by
regulating Notch signalling. *EMBO J*, *31*(4), 817-828. https://doi.org/10.1038/emboj.2011.443
Khacho, M., & Slack, R. S. (2018). Mitochondrial dynamics in the regulation of neurogenesis: From
development to the adult brain. *Dev Dyn*, *247*(1), 47-53. https://doi.org/10.1002/dvdy.24538
Kim, J. W., Yang, H. J., Brooks, M. J., Zelinger, L., Karakülah, G., Gotoh, N., Boleda, A., Gieser, L.,
Giuste, F., Whitaker, D. T., Walton, A., Villasmil, R., Barb, J. J., Munson, P. J., Kaya, K. D.,
Chaitankar, V., Cogliati, T., & Swaroop, A. (2016). NRL-Regulated Transcriptome Dynamics of
Developing Rod Photoreceptors. *Cell Rep*, *17*(9), 2460-2473.
https://doi.org/10.1016/j.celrep.2016.10.074
Knafo, S., & Esteban, J. A. (2017). PTEN: Local and Global Modulation of Neuronal Function in Health
and Disease. *Trends Neurosci*, *40*(2), 83-91. https://doi.org/10.1016/j.tins.2016.11.008
Koch, L. M., Birkeland, E. S., Battaglioni, S., Helle, X., Meerang, M., Hiltbrunner, S., Ibáñez, A. J.,
Peter, M., Curioni-Fontecedro, A., Opitz, I., & Dechant, R. (2020). Cytosolic pH regulates
proliferation and tumour growth by promoting expression of cyclin D1. *Nat Metab*, *2*(11), 1212-
1222. https://doi.org/10.1038/s42255-020-00297-0
Kochanowski, K., Volkmer, B., Gerosa, L., Haverkorn van Rijsewijk, B. R., Schmidt, A., & Heinemann,
M. (2013). Functioning of a metabolic flux sensor in Escherichia coli. *Proc Natl Acad Sci U S A*,
110(3), 1130-1135. https://doi.org/10.1073/pnas.1202582110
Kubo, F., Takeichi, M., & Nakagawa, S. (2005). Wnt2b inhibits differentiation of retinal progenitor cells
in the absence of Notch activity by downregulating the expression of proneural genes.
Development, *132*(12), 2759-2770. https://doi.org/10.1242/dev.01856
Laroche, S., Stil, A., Germain, P., Cherif, H., Chemtob, S., & Bouchard, J. F. (2021). Participation of L-
Lactate and Its Receptor HCAR1/GPR81 in Neurovisual Development. *Cells*, *10*(7).
https://doi.org/10.3390/cells10071640
Malm, S. W., Hanke, N. T., Gill, A., Carbajal, L., & Baker, A. F. (2015). The anti-tumor efficacy of 2-
deoxyglucose and D-allose are enhanced with p38 inhibition in pancreatic and ovarian cell lines.
Journal of Experimental & Clinical Cancer Research, *34*(1), 31. https://doi.org/10.1186/s13046-
015-0147-4
Marquardt, T., Ashery-Padan, R., Andrejewski, N., Scardigli, R., Guillemot, F., & Gruss, P. (2001). Pax6
is required for the multipotent state of retinal progenitor cells. *Cell*, *105*(1), 43-55.
https://doi.org/10.1016/s0092-8674(01)00295-1
Melnik, S., Dvornikov, D., Müller-Decker, K., Depner, S., Stannek, P., Meister, M., Warth, A., Thomas,
M., Muley, T., Risch, A., Plass, C., Klingmüller, U., Niehrs, C., & Glinka, A. (2018). Cancer cell
specific inhibition of Wnt/β-catenin signaling by forced intracellular acidification. *Cell Discov*, *4*,
37. https://doi.org/10.1038/s41421-018-0033-2
Miyazawa, H., Snaebjornsson, M. T., Prior, N., Kafkia, E., Hammaren, H. M., Tsuchida-Straeten, N.,
Patil, K. R., Beck, M., & Aulehla, A. (2022). Glycolytic flux-signaling controls mouse embryo
mesoderm development. *Elife*, *11*. https://doi.org/10.7554/eLife.83299
Nerli, E., Rocha-Martins, M., & Norden, C. (2020). Asymmetric neurogenic commitment of retinal
progenitors involves Notch through the endocytic pathway. *Elife*, *9*.
https://doi.org/10.7554/eLife.60462

Oginuma, M., Harima, Y., Tarazona, O. A., Diaz-Cuadros, M., Michaut, A., Ishitani, T., Xiong, F., & Pourquie, O. (2020). Intracellular pH controls WNT downstream of glycolysis in amniote embryos. *Nature*, *584*(7819), 98-101. https://doi.org/10.1038/s41586-020-2428-0

Ouchi, Y., Baba, Y., Koso, H., Taketo, M. M., Iwamoto, T., Aburatani, H., & Watanabe, S. (2011). β-Catenin signaling regulates the timing of cell differentiation in mouse retinal progenitor cells. *Molecular and Cellular Neuroscience*, *46*(4), 770-780. https://doi.org/https://doi.org/10.1016/j.mcn.2011.02.010

Paolini, A., Duchemin, A. L., Albadri, S., Patzel, E., Bornhorst, D., Gonzalez Avalos, P., Lemke, S., Machate, A., Brand, M., Sel, S., Di Donato, V., Del Bene, F., Zolessi, F. R., Ramialison, M., & Poggi, L. (2015). Asymmetric inheritance of the apical domain and self-renewal of retinal ganglion cell progenitors depend on Anillin function. *Development*, *142*(5), 832-839. https://doi.org/10.1242/dev.118612

Park, K. K., Liu, K., Hu, Y., Smith, P. D., Wang, C., Cai, B., Xu, B., Connolly, L., Kramvis, I., Sahin, M., & He, Z. (2008). Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. *Science*, *322*(5903), 963-966. https://doi.org/10.1126/science.1161566

Peeters, K., Van Leemputte, F., Fischer, B., Bonini, B. M., Quezada, H., Tsytlonok, M., Haesen, D., Vanthienen, W., Bernardes, N., Gonzalez-Blas, C. B., Janssens, V., Tompa, P., Versees, W., & Thevelein, J. M. (2017). Fructose-1,6-bisphosphate couples glycolytic flux to activation of Ras. *Nat Commun*, *8*(1), 922. https://doi.org/10.1038/s41467-017-01019-z

Peng, M., Yin, N., Chhangawala, S., Xu, K., Leslie, C. S., & Li, M. O. (2016). Aerobic glycolysis promotes T helper 1 cell differentiation through an epigenetic mechanism. *Science*, *354*(6311), 481-484. https://doi.org/doi:10.1126/science.aaf6284

Qian, X., Li, X., Shi, Z., Xia, Y., Cai, Q., Xu, D., Tan, L., Du, L., Zheng, Y., Zhao, D., Zhang, C., Lorenzi, P. L., You, Y., Jiang, B. H., Jiang, T., Li, H., & Lu, Z. (2019). PTEN Suppresses Glycolysis by Dephosphorylating and Inhibiting Autophosphorylated PGK1. *Mol Cell*, *76*(3), 516-527.e517. https://doi.org/10.1016/j.molcel.2019.08.006

Rapaport, D. H., Wong, L. L., Wood, E. D., Yasumura, D., & LaVail, M. M. (2004). Timing and topography of cell genesis in the rat retina. *J Comp Neurol*, *474*(2), 304-324. https://doi.org/10.1002/cne.20134

Ryall, J. G., Dell'Orso, S., Derfoul, A., Juan, A., Zare, H., Feng, X., Clermont, D., Koulnis, M., Gutierrez-Cruz, G., Fulco, M., & Sartorelli, V. (2015). The NAD(+)-dependent SIRT1 deacetylase translates a metabolic switch into regulatory epigenetics in skeletal muscle stem cells. *Cell Stem Cell*, *16*(2), 171-183. https://doi.org/10.1016/j.stem.2014.12.004

Sakagami, K., Chen, B., Nusinowitz, S., Wu, H., & Yang, X. J. (2012). PTEN regulates retinal interneuron morphogenesis and synaptic layer formation. *Mol Cell Neurosci*, *49*(2), 171-183. https://doi.org/10.1016/j.mcn.2011.11.007

Schmidt, M., Tanaka, M., & Münsterberg, A. (2000). Expression of (beta)-catenin in the developing chick myotome is regulated by myogenic signals. *Development*, *127*(19), 4105-4113. https://doi.org/10.1242/dev.127.19.4105

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*, *9*(7), 671-675. https://doi.org/10.1038/nmeth.2089

Shi, L., Pan, H., Liu, Z., Xie, J., & Han, W. (2017). Roles of PFKFB3 in cancer. *Signal Transduction and Targeted Therapy*, *2*(1), 17044. https://doi.org/10.1038/sigtrans.2017.44

Tachibana, N., Cantrup, R., Dixit, R., Touahri, Y., Kaushik, G., Zinyk, D., Daftarian, N., Biernaskie, J., McFarlane, S., & Schuurmans, C. (2016). Pten Regulates Retinal Amacrine Cell Number by Modulating Akt, Tgfbeta, and Erk Signaling. *J Neurosci*, *36*(36), 9454-9471. https://doi.org/10.1523/JNEUROSCI.0936-16.2016

Tachibana, N., Touahri, Y., Dixit, R., David, L. A., Adnani, L., Cantrup, R., Aavani, T., Wong, R. O., Logan, C., Kurek, K. C., & Schuurmans, C. (2018). Hamartoma-like lesions in the mouse retina: an animal model of Pten hamartoma tumour syndrome. *Dis Model Mech*, *11*(5). https://doi.org/10.1242/dmm.031005

Takata, N., Miska, J. M., Morgan, M. A., Patel, P., Billingham, L. K., Joshi, N., Schipma, M. J., Dumar, Z. J., Joshi, N. R., Misharin, A. V., Embry, R. B., Fiore, L., Gao, P., Diebold, L. P., McElroy, G. S., Shilatifard, A., Chandel, N. S., & Oliver, G. (2023). Lactate-dependent transcriptional regulation controls mammalian eye morphogenesis. *Nat Commun*, *14*(1), 4129. https://doi.org/10.1038/s41467-023-39672-2

Tatapudy, S., Aloisio, F., Barber, D., & Nystul, T. (2017). Cell fate decisions: emerging roles for metabolic signals and cell morphology. *EMBO Rep*, *18*(12), 2105-2118. https://doi.org/10.15252/embr.201744816

Touahri, Y., Hanna, J., Tachibana, N., Okawa, S., Liu, H., David, L. A., Olender, T., Vasan, L., Pak, A., Mehta, D. N., Chinchalongporn, V., Balakrishnan, A., Cantrup, R., Dixit, R., Mattar, P., Saleh, F., Ilnytskyy, Y., Murshed, M., Mains, P. E., Kovalchuk, I., Lefebvre, J. L., Leong, H. S., Cayouette, M., Wang, C., Sol, A. D., Brand, M., Reese, B. E., & Schuurmans, C. (2024). Pten regulates endocytic trafficking of cell adhesion and Wnt signaling molecules to pattern the retina. *Cell Rep*, *43*(4), 114005. https://doi.org/10.1016/j.celrep.2024.114005

Turner, D. L., & Cepko, C. L. (1987). A common progenitor for neurons and glia persists in rat retina late in development. *Nature*, *328*(6126), 131-136. https://doi.org/10.1038/328131a0

Turner, D. L., Snyder, E. Y., & Cepko, C. L. (1990). Lineage-independent determination of cell type in the embryonic mouse retina. *Neuron*, *4*(6), 833-845. https://doi.org/10.1016/0896-6273(90)90136-4

Vander Heiden, M. G., Cantley, L. C., & Thompson, C. B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*, *324*(5930), 1029-1033. https://doi.org/10.1126/science.1160809

Warburg, O. (1956a). On respiratory impairment in cancer cells. *Science*, *124*(3215), 269-270. https://www.ncbi.nlm.nih.gov/pubmed/13351639

Warburg, O. (1956b). On the origin of cancer cells. *Science*, *123*(3191), 309-314. https://doi.org/10.1126/science.123.3191.309

Wetts, R., & Fraser, S. E. (1988). Multipotent precursors can give rise to all major cell types of the frog retina. *Science*, *239*(4844), 1142-1145. https://doi.org/10.1126/science.2449732

Young, R. W. (1985). Cell differentiation in the retina of the mouse. *Anat Rec*, *212*(2), 199-205. https://doi.org/10.1002/ar.1092120215

Zhang, C. S., Hawley, S. A., Zong, Y., Li, M., Wang, Z., Gray, A., Ma, T., Cui, J., Feng, J. W., Zhu, M., Wu, Y. Q., Li, T. Y., Ye, Z., Lin, S. Y., Yin, H., Piao, H. L., Hardie, D. G., & Lin, S. C. (2017). Fructose-1,6-bisphosphate and aldolase mediate glucose sensing by AMPK. *Nature*, *548*(7665), 112-116. https://doi.org/10.1038/nature23275

Zhang, D., Li, J., Wang, F., Hu, J., Wang, S., & Sun, Y. (2014). 2-Deoxy-D-glucose targeting of glucose metabolism in cancer cells as a potential therapy. *Cancer Letters*, *355*(2), 176-183. https://doi.org/https://doi.org/10.1016/j.canlet.2014.09.003

Zhang, K., Sun, L., & Kang, Y. (2023). Regulation of phosphoglycerate kinase 1 and its critical role in cancer. *Cell Communication and Signaling*, *21*(1), 240. https://doi.org/10.1186/s12964-023-01256-4

Zhang, X., Leavey, P., Appel, H., Makrides, N., & Blackshaw, S. (2023). Molecular mechanisms controlling vertebrate retinal patterning, neurogenesis, and cell fate specification. *Trends Genet*, *39*(10), 736-757. https://doi.org/10.1016/j.tig.2023.06.002

Hanna et al. Figure 1

A

size factor

wild-type   *Pten*-cKO

B

wild-type vs. *Pten*-cKO Volcano plot

● Log₂ FC   ● p-value and Log₂ FC

Log₂ fold change

total=15,421 genes

408   14346   667

down DEGs   up DEGs

C

Kyoto Encyclopedia of Genes and Genomes (KEGG) Analysis

Axon guidance
Regulation of actin cytoskeleton
HIF-1 signaling pathway
AMPK signaling pathway
Proteoglycans in cancer
Thyroid hormone signaling pathway
Hepatocellular carcinoma
Carbon metabolism
Insulin signaling pathway
Glycolysis/gluconeogenesis
Central carbon metabolism in cancer
ErbB signaling pathway
Biosynthesis of amino acids
EGFR tyrosine kinase inhibitor resistance
Long-term depression
Fructose and mannose metabolism
Type II diabetes mellitus

gene ratio

p-adjust   0.0025 0.0050 0.0075

count   7.5 10.0 12.5 15.0 17.5

D

Glucose — HK → glucose-6-phosphate — GPI → fructose-6-phosphate — PFK1 → fructose-1,6-phosphate — Aldoa → glyceraldehyde 3-phosphate — GAPDH → 1,3-bisphospho-glycerate

dihydroacetone phosphate

2ATP 2ADP   PGK

intracellular pH (pHi) alkaline

Lactate ← Pyruvate ← PKM ← phosphoenol-pyruvate ← Eno ← 2-phospho-glycerate ← PGAM ← 3-phospho-glycerate

LdhA/B

2ATP 2ADP

Slc16a3

extracellular pH (pHe) acidic

Lactate   H⁺

E

Hk1 padj=0.0096
Gpi1 padj=9.84^-10
Pfkl padj=1.06^-05
Aldoa padj=6.98^-12
Pgk1 padj=0.0044
Pgam1 padj=0.0437
Eno1 padj=0.0052
Pkm padj=0.1091
Slc16a3 padj=0.0004

RPKM

wild-type   *Pten*-cKO

F

Hk1 P=0.05
Pgk1 P=0.0002
Eno1 P=0.041
Slc16a3 P=0.043

Normalized expression

wild-type   *Pten*-cKO

G

● Control ● *Pten*-cKO

Oligomycin   FCCP   R/A   p=0.0175

ECAR (mpH/min/Norm. Unit)

Time (minutes)

● Control ● *Pten*-cKO

Oligomycin   FCCP   R/A   p=0.6976

OCR (mpH/min/Norm. Unit)

Time (minutes)

H

Lactate Assay

P=0.024

nmol/ug Protein

wild-type   *Pten* cKO

Hanna et al. Figure 2

Hanna et al. Figure 3

Hanna et al. Figure 4

A

ACIDIC pH 6.5

LdhA/B · Lactate · Pyruvate · phosphoenol pyruvate · PKM · PKM

2ATP 2ADP

H+

Slc16a3

B

P0 eye → retina → retinal flatmount → CNCn or pH6.5 pH7.4 pH8.0 · BrdU 30min or 24 hr → retinal explant culture

C P0 + 30 min

control / 5mM CNCn

BrdU- 30min

p=0.002

D P0 + 30 min

pH6.5 · pH7.4 · pH8.0

BrdU- 30min

p=0.007 p=0.001

E P0 + 24 hr

Ki67/BrdU

p=0.014 p=0.06

BrdU- 24 hr

p=0.002 p=0.015

Ki67

ns ns

F P0 + 24 hr

pH6.5 · pH7.4 · pH8.0

VSX2

p=0.09 p=0.009

G P0 + 24 hr

CC3

ns ns

H P0 + 24 hr

pH6.5 · pH7.4 · pH8.0

CRX

p=0.14 p=0.006

I pCAGG-pHluorin-IRES2-tdTomato (P3 for 2 DIV)

488 nm / tdTomato / merge

Hanna et al. Figure 5

Hanna et al. Figure 7