Improving biodiversity in Central and Eastern European domestic gardens needs regionally scaled strategies

Zsófia Varga-Szilay[a*], Arvīds Barševskis[b], Klára Benedek[c], Danilo Bevk[d], Agata Jojczyk[e], Anton Krištín[f], Jana Růžičková[g,h], Lucija Šerić Jelaska[i], Eve Veromann[j], Silva Vilumets[j], Kinga Gabriela Fetykó[k], Gergely Szövényi[h], Gábor Pozsgai[l*]

[a]Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, 1117, Budapest, Hungary

[b]Coleopterological Research Center, Institute of Life Sciences and Technology, Daugavpils University, Vienības str. 13, 5401, Daugavpils, Latvia

[c]Department of Horticulture, Faculty of Technical and Human Sciences, Sapientia Hungarian University of Transylvania, Calea Sighișoarei nr. 2, 540485, Târgu-Mureș, Romania

[d]Department of Organisms and Ecosystems Research, National Institute of Biology, Vecna pot 121, 1000, Ljubljana, Slovenia

[e]Department of Landscape Art, Institute of Environmental Engineering, Warsaw University of Life Science, Nowoursynowska str. 166., 02-787, Warsaw, Poland

[f]Institute of Forest Ecology Slovak Academy of Sciences, Ľ. Štúra 2, 960 53, Zvolen, Slovakia

[g]HUN-REN-ELTE-MTM Integrative Ecology Research Group, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, 1117, Budapest, Hungary

[h]Department of Systematic Zoology and Ecology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, 1117, Budapest, Hungary

[i]Department of Biology, Faculty of Science University of Zagreb, Rooseveltov trg 6, 10000, Zagreb, Croatia

[j]Chair of Plant Health, Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Fr. R. Kreutzwaldi 1, 51006, Tartu, Estonia

[k]Independent researcher, 6000, Kecskemét, Hungary

[l]Ce3C – Centre for Ecology, Evolution and Environmental Changes, Azorean Biodiversity Group, CHANGE – Global Change and Sustainability Institute, University of the Azores, Faculty of AgricuGltural Sciences and Environment, Rua Capitão João D'Ávila, 9700-042, Angra do Heroísmo, Terceira, Açores, Portugal

*Corresponding authors: Zsófia Varga-Szilay (zsofia@vargaszilay.hu) and Gábor Pozsgai (pozsgaig@coleoptera.hu)

Abstract

Amid ongoing urbanisation and increasing anthropogenic activities, domestic gardens, while they cannot replace undisturbed habitats, play a valuable role in enhancing urban biodiversity by supplementing green areas and improving landscape connectivity. Moreover, biodiversity-friendly gardens also improve human well-being and foster connections between nature and people. To study these benefits, we distributed an online questionnaire between 2022 and 2023, and used a scoring system to evaluate the ecological value of gardens, garden owners' motivations, and pesticide use habits. We used machine learning to explore how these indices interact and what sociodemographic factors drive them across nine Central and Eastern European (CEE) countries. Additionally, we explored the differences and similarities between the ecological values of gardens and gardening practices in building high biodiversity. Our findings reveal significant variability both between and within countries, across all three indices with Romania faring low and Czechia reaching high scores in all three indices. Domestic pesticide use was ubiquitous across all CEE and largely unaffected by sociodemographic factors. However, increased time spent with gardening was associated with reduced pesticide use and a greater potential for fostering high biodiversity. Garden owners over 55 tended to follow and uphold longstanding conventional practices and thus lowered both pesticide and garden index scores. The local-scale differences we observed emphasise the need for regionally tailored guidelines for biodiversity-friendly gardening over standardized regulations across Europe. Optimal strategies for effective environmental educational and community programs can be developed based on local biodiversity and the three indices used in this study. To maximize their impact and meet local needs, these programs should educate garden owners about the environmental and health effects of pesticides and offer comprehensive biodiversity-related information across all regions and social strata. This is particularly crucial in Central and Eastern Europe (CEE), where such initiatives are currently underrepresented.

Highlights

- Domestic gardens can enhance biodiversity and support human well-being.

- Study conducted across nine Central and Eastern European countries.

- Significant variability both between and within countries in gardening practices.

- Improving biodiversity in domestic gardens needs regionally scaled strategies.

- Reducing pesticide use and increasing awareness are key.

Keywords: rural-urban gradient; urban ecosystems; environmental consciousness; sustainable gardening; environmental sensitivity; urbanisation

1. Introduction

Built-up urban areas will cover 7% of the European Union by 2030 (Perpiña Castillo et al., 2019) and this ongoing urbanisation and increasing anthropogenic activities, in turn, contribute to biodiversity loss (Hagen et al., 2012) and the rapid decline of undisturbed ecosystems (Bengtsson et al., 2000). Most parts of Europe are already impoverished, and characterised by small natural areas, broken up by human intervention into a mosaic of agricultural lands and settlements (European Environment Agency, 2012; Eurostat, 2022b; Jongman, 2002; Perpiña Castillo et al., 2019). Thus, it is particularly important to halt, or at least reduce the ongoing degradation and losses of natural habitats in areas heavily modified by human activities, such as agricultural land and cities, and, as much as possible, maximise biodiversity. However, whilst conservation efforts for maintaining and preserving biodiversity have already been a part of many agricultural systems (Concepción et al., 2020; van Elsen, 2000), urban biodiversity rarely gains enough attention (Fischer et al., 2020; Ramalho & Hobbs, 2012). Although, the European Commission adopted a law to stop the loss of green urban spaces by 2030 and achieve a 10% a minimum tree canopy cover by 2050 (Nature Restoration Law, European Commission, 2024), the area of urban green spaces has only increased in some Southern and Western European cities (e.g. Pamplona/Iruña (Spain) and Hague (Netherlands)). From Eastern Europe, Czechia, Poland, and Romania are just about to catch up (European Environment Agency, 2023; Kabisch & Haase, 2013). Yet, in none of the European Union countries, could the amount of re-naturalised areas compensate for what has been lost for urbanisation (European Commission, 2022).

Urban green infrastructure covers an average of 42% of the city area according to the European Environment Agency (EEA) (European Environment Agency, 2022), implying that urban biodiversity can be substantially improved by developing green infrastructure (Baldock, 2020; Baldock et al., 2019; Heidt & Neef, 2008; Liang et al., 2023), such as diversifying city/urban

parks and community gardens, rooftop gardens, establishing pollinator-friendly meadows, creating a network of corridors, and installing green roofs (Aguilera et al., 2019; Beninde et al., 2015; Lin et al., 2015; Seitz et al., 2022). Besides public places, however, private domestic gardens occupy an average of 16-36% of the total urban areas in Europe (Cameron et al., 2012; Goddard et al., 2010), and therefore the role of domestic gardens in preserving and supporting native biodiversity and their contribution to urban sustainability are becoming increasingly important (Fontaine et al., 2016; Hanson et al., 2021). Indeed, enhancing environmental quality in domestic gardens offers a great opportunity to improve human health and well-being (Camps-Calvet et al., 2016; Krols et al., 2022; Marques et al., 2021; Soga et al., 2017) and alleviate biodiversity loss simultaneously (Freeman et al., 2012; Li et al., 2023).

The biodiversity of domestic gardens varies along a wide spectrum (Home et al., 2019; Lindemann-Matthies & Marty, 2013), and their ecological value is determined by multiple environmental factors, such as climate, adjacent landscape mosaic/characteristics (Braschler et al., 2020), natural vegetation (Borysiak et al., 2017; Prendergast et al., 2022), and soil composition (Tresch et al., 2018, 2019). Although these intrinsic attributes are mostly beyond owners' control, the impact of domestic gardens on biodiversity primarily depends on the decisions of garden owners, which dictates design, use, and management (Quistberg et al., 2016). With appropriate management, even small or highly anthropized domestic gardens can support high biodiversity (Donkersley et al., 2023; Griffiths-Lee et al., 2022; Muratet & Fontaine, 2015). On the other hand, numerous anthropogenic stressors (Varga-Szilay et al., 2024), such as the intensity of cultivation (Krištín et al., 2024; van der Veen, 2005), mowing frequency (Lerman et al., 2018) or pesticide use (Tassin de Montaigu & Goulson, 2023), often compromise the ecological value of gardens (Varga-Szilay & Pozsgai, 2022). There is an agreement that the potential ecological value of gardens in conservation correlates positively with the structural diversity of gardens (Majewska & Altizer, 2020), reduced pesticide usage

(Tassin de Montaigu & Goulson, 2023), and reduced mowing frequency (Chollet et al., 2018;

Lerman et al., 2018; Proske et al., 2022). Ultimately, however, garden management is a series

of 'tyranny of small decisions' (Dewaelheyns et al., 2016) with mixed motivations (Gifford &

Nilsson, 2014; Hruška et al., 2021) and external factors such as demographics, household

income, neighbours' expectations, accessibility of information, and education, can influence

owners' behaviour (Goddard et al., 2013; Varga-Szilay et al., 2024). Hence, considering all

factors, evaluating the biodiversity values of gardens necessitates considering the garden

owners' management and attitude toward a diverse garden. Yet, the major drivers in

management may depend on several societal factors.

In Western Europe, leisure opportunities and recreational activities have long been the primary

drivers of gardening, promoting well-being (Beumer, 2018) whereas in Central and Eastern

Europe (CEE) gardening for food self-provisioning (FSP) still may dominate (Jehlička et al.,

2020, 2021; Smith & Jehlička, 2013). The Central and Eastern European countries share a

historical background in FSP, which is thought to be a coping strategy for economic pressures

(Alber & Kohler, 2008; Jehlička et al., 2021). However, the previous differences in gardening

motivation are gradually beginning to blur with the emergence of new gardening trends (e.g.

wildlife-friendly gardening) in both Western and Central- and Eastern Europe (Keshavarz et al.,

2016; Poniży et al., 2021) as the demand for ensuring food security and quality in Western

countries grows (Church et al., 2015; Glavan et al., 2018; Pourias et al., 2016). At the same

time, gardening in CEE countries shifts toward recreational activities (Petzke et al., 2021; Tóth

et al., 2018; Trendov, 2018). Yet, the increasing emphasis on nature conservation across all of

Europe (Galluzzi et al., 2010; Muller et al., 2010; Vávra et al., 2014), seems to be embraced by

CEE countries slower than in their Western counterparts (Hruška et al., 2021; Vávra et al.,

2018).

Thus, the divide in the applied gardening practices between East and West remains significant.

Indeed, due to the limited access to biodiversity-related information (Coisnon et al., 2019;

European Commission, Directorate-General for Communication, 2015), lower willingness to

change to environmentally friendly practices, and the extensive use of pesticides (European

Commission, Directorate-General for Communication, 2015), CEE appears to be lagging

behind in the adoption of biodiversity-friendly practices (for example leave space for wildlife)

(European Commission, Directorate-General for Communication, 2015).

As differences between East and West remain, assessing the ecological value of domestic

gardens, therefore, requires consideration of location-specific anthropogenic and environmental

variables (Varga-Szilay et al., 2024). Whilst there have been numerous studies investigating the

role of gardens in maintaining urban biodiversity in Western Europe since the 1990s (Delahay

et al., 2023), in CEE these are scarce (but see Varga-Szilay et al., 2024; Varga-Szilay & Pozsgai,

2022).

To address this knowledge gap, here, we seek to investigate how gardens' parameters, the

gardening motivation of garden owners, and their pesticide use habits influence each other in

nine Central- and Eastern European countries and explore the differences and similarities

between gardens and gardening practices with a potential for maintaining high biodiversity.

Since these can fundamentally drive local educational strategies, we particularly aim to pinpoint

how geographical differences and sociodemographic parameters best predict domestic gardens'

ecological values, the respondents' attitudes towards supporting insect pollinators in their

gardens, and their pesticide use habits. Hence, insight into the costs and benefits of initiatives

for increasing garden diversities is essential for effective planning and management, our

imperative is to highlight those areas in CEE where these programs can yield the greatest

rewards (e.g. densely populated regions originally hosting high biodiversity).

2. Material and methods

2.1 Questionnaire design and data collection

An online questionnaire was distributed in nine CEE countries (Croatia, Czechia, Estonia, Hungary, Latvia, Poland, Romania, Slovakia, and Slovenia), all of which were formerly part of the Eastern Block and are currently members of the European Union.

The questionnaire was translated from Hungarian to English and then from English to Croatian, Czech, Estonian, Latvian, Polish, Romanian, Slovak, and Slovenian. The participants could choose the language in which they wanted to complete the questionnaire. It was mandatory to respond to 55 of the total 59 questions, organised into nine sections, which took a maximum of 15 minutes to complete. The questionnaire gathered information about the location and the main characteristics of the garden, the sociodemographic parameters, the garden owners' motivations, cultivation habits, pesticide usage, environmental awareness, and pollinator-friendly practices. Although participants had to indicate their gender (male, female, other), their highest level of completed education (elementary, middle, postsecondary, postgraduate), and their residency according to the Nomenclature of Territorial Units for Statistics (henceforth NUTS, Eurostat, 2021), all responses were otherwise anonymously recorded. Because of the large number of NUTS-3 regions in Poland, to simplify the questionnaire, here, residencies were recorded at NUTS-2 levels, while the other eight participating countries were recorded at NUTS-3 levels.

The questionnaire was designed using Google Forms and was actively distributed between 26[th] October 2022 and 18[th] May 2023, for 90 days in each of the nine participating countries (**Supplementary Material Table 1**). The questionnaire was disseminated through channels such as gardening-oriented websites, and various online social media platforms (for instance Facebook and Instagram) where QR codes and hashtags were used to increase the sharing efficiency. Moreover, through targeted email distributions, outreach efforts extended to

professional associations, non-governmental organisations, foundations, and societies

dedicated to domestic gardening and environmental conservation.

The participation was voluntary, and a Data Protection and Privacy Statement was available

alongside the questionnaire.

2.2 Definitions

Not all respondents are, actually, the owners of the gardens they reported on but, for simplicity,

we refer to everyone as a 'garden owner'. For terminological clarity, we used the definitions of

'pesticides' and 'gardening' as given by Varga-Szilay et al. (2024). Pesticides were defined as

'all synthetic and non-synthetic products that are used to control pests', including 'all

commercially available and homemade plant protection products, either those allowed in

organic gardening or used in conventional practices'. Gardening was defined as 'all garden

work and all garden care practices, such as the cultivation of flowers, fruits, vegetables, and

ornamental plants, mowing, and soil management' (Varga-Szilay et al., 2024).

2.3 Data processing

For the analysis, we only used 43 questions relevant to our aims of the original 59 ones. The

original categorical replies of our questionnaire were re-categorised for analytical purposes on

a few occasions (see **Supplementary Methods**). NUTS polygons, number of inhabitants, the

regional gross domestic product (GDP) in purchasing power standards per inhabitant for each

NUTS (PPS, henceforth), and the urban-rural typology for each NUTS-3 region were obtained

from ArcGIS Data and Maps (2022a, 2022b) and Eurostat (2022a, 2024), respectively. PPS was

only available for NUTS-2 categories. When country capitals with separate NUTS categories

were situated within a larger unit, capitals were merged with the largest NUTS surrounding

them (CZ, HR, HU, LV, PL, RO). In these cases, for PSS and number of inhabitants, the mean

and sum of the merged areas were calculated, respectively. In the case of urban-rural typology

data, the originally obtained NUTS-3 categories were merged into NUTS-2 in Poland, and the

NUTS polygons of the above mentioned capitals were merged with the surrounding polygons.

For this, a weighted mean of the numerised typology index was calculated using the areas of

the merged NUTS polygons as weights. The means were rounded to the nearest integer and

converted back to categories. Spatial and polygon calculations were conducted using the 'sf'

(Pebesma, 2018) R package.

2.4 Answer-based scoring system

We rated the potential ecological value of the gardens, the garden owners' motivation for

gardening, and pesticide use habits with an answer-based scoring system. For this, we used a

total of 32 questions from the original 59 (i.e. questions related to sociodemographic variables

were excluded). The garden (GAR) index shows what potential a domestic garden has to build

high biodiversity. This index reflects on the structural diversity of the gardens, including garden

size, the area of undisturbed patches and plants covering the garden, and the habitat types of

adjacent areas, as well as, the disturbance level and the presence of artificial habitats. The

calculation of this index was based on 12 questions and consisted of 10 components, which

contributed with different weights. The weights were determined by the importance of the

components in maintaining high biodiversity (**Supplementary Methods section 1.1**). The

respondent (RES) index assesses the garden owners' knowledge of garden wildlife and their

attitudes to maintaining/creating a biodiversity-friendly garden. The RES index was based on

10 questions and consisted of 10 components, including both theoretical (e.g. 'Do you

think…?'/'Can you imagine…?') and practical (e.g. 'How do you…?') questions/question

groups, which contributed to the index with different weights (**Supplementary Methods**

section 1.2). The pesticide (PES) index shows the degree of pesticide load in gardens by

assessing the amount and diversity of pesticides used and the related knowledge of the garden

owners. It was based on 10 questions and consisted of 11 components, which were contributed

to the index with different weights (**Supplementary Methods section 1.3**). All indices were

scaled between 0 (as the lowest) and 100 (as the highest) points. For the full details of the index

calculation process, the reader should consult with the **Supplementary Methods**.

2.5 Bird index

Since garden diversity depends on the historical biodiversity of a given area (Dobrovodská et

al., 2023) but simplified, human-modified landscapes can achieve moderate levels of ecological

benefits more rapidly (Haenke et al., 2009), the efficiency of biodiversity-friendly activities can

only be measured in the context of local biodiversity indices. Thus, as a proxy for local

biodiversity, we collected a full list of bird species recorded between 2010 and 2023 in each

NUTS-3 area (except in Poland, see section 2.1) from the Global Biodiversity Information

Facility (GBIF.org, 2024). To minimise the observer bias and the effect of rare species, in each

NUTS, we only considered species that had at least five recorded occurrences per year from the

particular area. We standardised the collated number of species with the area of the NUTS (i.e.

divided the species number with the area of NUTS in km squares). Using the same methodology

we also gathered a species list for the whole area of interest (i.e. the nine participating countries)

and standardised for bird species per km^2. We then divided the standardised local bird richness

with the standardised whole-area bird richness. For collecting bird data we used the 'rgbif'

(Chamberlain et al., 2024) and for spatial calculations the 'sf' (Pebesma, 2018) R packages.

2.6 Urban-rural typology

The European urban-rural typology data is a qualitative index which classifies grid cells into

rural and non-rural grids and establishes three categories (predominantly rural, intermediate,

and predominantly urban) of urbanisation based on the percentage of the population living in

rural grid cells (Eurostat, 2024). We employed this index to approximate urbanisation. Since

the qualitative nature of this index, we also converted our four indices (bird, GAR, RES, PES)

to categorical variables by dividing them at their 0.33 and 0.66 quantiles. When calculating the

quantiles, only measured values were included and the theoretical minimum and maximum

were not considered. NUTS areas with less than five respondents were excluded.

2.6 Statistical analysis

We examined the correlation between the three indices (GAR, RES, PES) with the Spearman

correlation test. For calculating the correlation matrix the 'psych' (Revelle, 2021) R package

was used. We used the Kruskal-Wallis method to test if GAR, RES, and PES indices differ

among countries. Pairwise differences were investigated using the pairwise Wilcoxon test with

Holm-corrected p-values.

The three indices were used as response variables in three separate Gradient Boosting Machine

(GBM) learning processes. Seven sociodemographic variables (gender, age, education level,

having children, gardening experience, gardening perception and the average time spent

gardening), the PPS, latitude and longitude of the centroids of the NUTS areas were also

included in the model as numerical explanatory variables. We randomly divided the datasets

into training (70%) and test (30%) sets. After an optimisation and tuning process of the model

parameters, we built the GBM model using a Gaussian distribution with 5 levels of interaction

depth, with 0.3 shrinkage, 0.80 bag fraction, and a 10 fold cross validation on 28 (for GAR

index), 26 (for RES index), and 10 (for PES index) trees. The model fit was evaluated by

calculating the R-squared and Root Mean Standard Error (RMSE) values. We used the SHapley

Additive exPlanations (SHAP) method to interpret our final GBM models with the 'shapviz'

(Mayer & Stando, 2023) and 'kernelshap' (Mayer et al., 2023) R packages. The SHAP assesses

individual variable contributions, accounts for variable interactions, assigns importance values, and facilitates the comparison between all possible variable orders (Lundberg & Lee, 2017). Since GBM predicted the country identity as the most influential variable on all three indices, we examined the relationship between GAR and PES indices (the two non-correlating ones) separately for each country and explored how the other important variables grouped within this data cloud.

For modelling and the visualisation of model results, we used the 'gbm' (Greenwell et al., 2022) and 'caret' (Kuhn et al., 2023) packages in an R environment (R Core Team, 2021).

Preliminary data clean-up was done in a Python (Python Software Foundation, 2019) environment, with the help of 'NumPy' (Harris et al., 2020) and 'Pandas' (The pandas development team, 2022) libraries.

3. Results

3.1 Garden owners' sociodemographic characteristics

Altogether 5255 garden owners completed the questionnaire from the 9 participating countries (**Supplementary Material Fig. 1, Supplementary Material Table 1**), of which 1146 were males (21.80%), 4094 were females (77.91%), and 15 were non-binary gender (0.29%). The majority of respondents were between 36 and 55 years old (n = 2841, 54.06%). The greatest proportion of garden owners had a postsecondary education (n = 3225, 61.37%), followed by the middle education (n = 1466, 27.90%) and postgraduates (n = 512, 9.74%), while only 52 respondents had elementary school as the highest level of their education (0.99%) (**Supplementary Material Table 2**).

In all participating countries, most of the respondents considered gardening as their favourite hobby (between 21.05% and 58.31%) or a pleasant pastime (between 26.46% and 62.54%)

(**Supplementary Material Fig. 2**). In most participating countries, mowing several times a

month was under 40% (**Supplementary Material Fig. 3**) but in Estonia and Latvia, the

majority of garden owners (65.07% and 63.61%, respectively) mowed the lawn several times a

month, and the proportion of those who did so once a month was also high (27.04% and

21.88%). The average pesticide use among the nine countries was 52.60%, ranging from

38.88% in Slovakia to 69.59% in Romania (**Supplementary Material Fig. 4**).

3.2 Indices

The mean of the calculated indices was 56.49 (range: 14.97, 91.32), 63.25 (range: 9.25, 96.75),

76.33 (range: 24.87, 100), for GAR, RES, and PES, respectively (**Supplementary Material**

Table 3). The GAR index with the RES index showed a moderate, positive correlation

(Spearman's rho = 0.55, $p < 0.001$). The PES index showed an almost negligible, negative

correlation with the RES index (Spearman's rho = -0.09, $p < 0.001$). There was no significant

correlation between GAR and PES indices ($p = 0.487$).

There were significant differences between the participating countries in all three indices (GAR,

RES, PES) (Kruskal-Wallis chi-squared = 290.85, 671.51, and 235.13, respectively, $p < 0.001$)

(**Supplementary Material Table 4, 5**), and the index values varied broadly among NUTS areas

(**Fig. 1**). A total of 197 respondents had all three scores below the first quartile of the indices,

with those from Romania and Hungary faring the worst, whilst there were a total of 90

respondents in the fourth quartile of all indices, mostly from Czechia and Slovenia

(**Supplementary Material Table 3, 4**).

Fig. 1: The garden (A), respondents (B) and pesticide (C) indices in the nine participating countries. The colour depth of the maps indicates the mean of the indices calculated for each NUTS-2 area.

3.3 GBM outputs

3.3.1 GAR index

The GBM model for the GAR index explained 12.45% of the variance for new observations (RMSE = 11.29), with SHAP values indicating the country variable being the best and the average time that garden owners spent with gardening being the second-best predictor (SHAP value = 2.15, 1.94, respectively) (**Fig. 2A**).

Countries like Romania, Poland, and Croatia shifted the GAR index toward the lower, while Czechia, Estonia, Latvia, Slovenia, and Slovakia to higher values (**Fig. 2C**). The DIV index increased with increasing time spent with gardening.

The two age categories over 55 shifted the GAR index toward lower values (particularly the age group of over 65), in contrast, the three age categories between 26 and 55 resulted in higher values. Perceiving gardening as a work or a pleasant pastime both lowered the GAR index values while perceiving it as a favourite hobby increased them. Postsecondary and postgraduate education shifted the index to higher values, while middle-level education to lower ones. Albeit

360   it had a low influence in the model, garden owners who had children separated well from those

361   who did not, with those having children shifting the index to higher values. The GAR index

362   values were increased from the west to the east (**Fig. 2B**).

**Fig. 2: Global SHAP (SHapley Additive exPlanations) summary plots for the GBM model

of the GAR index.** Variables are ordered by their (A) importance based on average absolute

SHAP values. Beeswarm plots of SHAP values of (B) numerical and (C) categorical variables.

Each point represents one respondent. For numerical variables, colours indicate variable values,

while for categorical variables, they represent distinct category levels. Countries are

abbreviated as CZ, EE, HR, HU, LV, PL, RO, SI, and SK, for Czechia, Estonia, Croatia,

Hungary, Latvia, Poland, Romania, Slovenia, and Slovakia, respectively. Time spent with

gardening is shown in a self-reported scoring system as indicated by the respondents, with 1

representing the lowest ('one to two hours per day) and 5 representing the highest value ('even

twelve hours per day').

3.3.2 RES index

The GBM model for the RES index explained 24.73% of the variance for new observations

(RMSE = 13.13), with SHAP values indicating the country variable being the best and how

garden owners perceived gardening being the second-best predictors (SHAP value = 3.80, 2.26,

respectively) (**Fig. 3A**). Whilst Hungary and Czechia had the highest SHAP values, and thus

the RES index, Poland and Croatia had the lowest ones, with Romania and Latvia also

indicating lower values. Only perceiving gardening as a favourite hobby shifted the SHAP

values to higher ranges, all other cases lowered them, especially when gardening was perceived

as a duty (**Fig. 3C**). SHAP values increased with the time spent with gardening. Older age

groups predicted lower RES index values, while the 26-35 age group exhibited higher ones.

Gender, education, and having children were less important variables. However, SHAP values

separated well with gender, with women presenting at higher values. SHAP values increased

from west to east, except for a few very eastern locations presenting the highest values on the

right side of the axis (**Fig. 3B**).

Fig. 3: Global SHAP (SHapley Additive exPlanations) summary plots for the GBM model of the RES index. Variables are ordered by their (A) importance based on average absolute SHAP values. Beeswarm plots of SHAP values of numerical (B) and categorical (C) variables. Each point represents one respondent. For numerical variables, colours indicate variable values, while for categorical variables, they represent distinct category levels. Time spent with gardening is shown in a self-reported scoring system as indicated by the respondents, with 1 representing the lowest ('one to two hours per day) and 5 representing the highest value ('even twelve hours per day').

3.3.3 PES index

The GBM model for the PES index explained 0.08% of the variance for new observations (RMSE = 18.19), with SHAP values indicating the country variable being the best and the average time that garden owners spent with gardening being the second best predictors (SHAP value = 3.20, 1.56, respectively) (**Fig. 4A**). Romania, Estonia, and Hungary had the lowest SHAP values, whilst Poland, Slovakia, and Slovenia showed the highest ones (**Fig. 4C**). An average of 1–2 hours of gardening per day predicted a higher PES index, whilst all other times spent with gardening lowered it, with the lowest values associated with those who spent up to 12 hours a day with gardening. More than 10 years of gardening experience shifted the index to lower values. Perceiving gardening as a pleasant pastime shifted SHAP to higher values, whereas in most other cases, the values decreased equally when gardening was perceived as a duty and a favourite hobby. Women shifted SHAP to higher values, however, gender, age, having children, and education level were not deemed to be important variables. Having children separated positive and negative SHAP values, with those respondents who had children associated with higher values. SHAP values increased both from south to north and east to west (**Fig. 4B**).

Fig. 4: Global SHAP (SHapley Additive exPlanations) summary plots for the GBM model of the PES index. Variables are ordered by their (A) importance based on average absolute SHAP values. Beeswarm plots of SHAP values of numerical (B) and categorical (C) variables. Each point represents one respondent. For numerical variables, colours indicate variable values, while for categorical variables, they represent distinct category levels. Time spent with gardening is shown in a self-reported scoring system as indicated by the respondents, with 1 representing the lowest ('one to two hours per day) and 5 representing the highest value ('even twelve hours per day').

3.3.4 The GAR index as a function of the PES index

In our investigation into the influence of country, GAR index, and PES index on the grouping

of gardening experience lengths, we found that respondents who had been gardening for more

than 2 years were characterised by a greater potential for building high biodiversity and more

intensive pesticide use. In Estonia, Latvia and Slovakia, those who had been gardening for less

than a year grouped around low pesticide input (high PES index values). On the other hand, in

Slovenia, among those who had been gardening for a year or less, there was no one who did not

use pesticides at all. In the case of Hungary and Estonia, gardeners who have been gardening

for less than one year scored lower in the GAR index but were also characterised by lower PES

index values. (**Supplementary Material Fig. 5**).

In Hungary, Poland, Romania, and Slovenia for those garden owners who considered gardening

as their favourite hobby the GAR index was greater than in other countries and they also scored

lower in the PES index (**Supplementary Material Fig. 6**).

In all participating countries, except Slovakia, the GAR index was the lowest and the PES index

the highest among garden owners who spent less than 1-2 hours with gardening

(**Supplementary Material Fig. 7**).

In all countries but Croatia older age negatively influenced the GAR index. Indeed in Romania,

garden owners over the age of 65 fared poorly with the GAR index and all of them used

pesticides. In contrast in Croatia, this age group had the highest GAR index values

(**Supplementary Material Fig. 8**).

3.4 Areas with a potential for improvement

When areas within urbanisation levels were cross-referenced with the categorised three indices,

different effects of urbanisation levels of the three indices were revealed (**Fig. 5**). Half of the

predominantly urban NUTS areas (n = 3) were categorised as low GAR values, and half of the

NUTS areas typologised (n = 26) as intermediate reached a medium level on the GAR index.

The RES categories were relatively evenly spread both among intermediate and predominantly

rural areas but most predominantly urban regions (n = 4) were categorised as medium RES

(**Table 1**). The distribution of PES categories was uniform among predominantly urban and

predominantly rural areas, whilst in intermediately urbanised regions intermediate PES

category was the most common (n = 22), followed by the high and low categories (n = 17 and

13, respectively) (**Table 1, Fig. 5**).

Table 1: Contingency tables of urban-rural typology categories and categorised index values of

bird, GAR, RES, and PES indices, as expressed by the number of NUTS areas. Values are

shown as percentages of the columns.

	Urban-rural typology		
Bird	Predominantly urban (n = 6)	Intermediate (n = 56)	Predominantly rural (n = 77)
Low (n = 46)	0	25.00	41.56
Moderate (n = 44)	33.33	35.71	31.17
High (n = 47)	66.67	39.29	27.27
GAR*	Predominantly urban (n = 6)	Intermediate (n = 52)	Predominantly rural (n = 61)
Low (n = 41)	50.00	26.92	39.34
Moderate (n = 42)	16.67	50.00	24.59
High (n = 36)	33.33	23.08	36.07
RES*	Predominantly urban (n = 6)	Intermediate (n = 52)	Predominantly rural (n = 61)
Low (n = 41)	16.67	28.85	39.34
Moderate (n = 42)	66.67	26.92	34.43
High (n = 36)	16.67	44.23	26.23
PES*	Predominantly urban (n = 6)	Intermediate (n = 52)	Predominantly rural (n = 61)
Low (n = 41)	33.33	25.00	32.79
Moderate (n = 42)	33.33	42.31	31.15
High (n = 36)	33.33	32.69	36.07

*Represents a total of 119 areas from which we had at least 5 respondents.

Of the NUTS areas typologised as predominantly urban, Czechia, Hungary, Slovakia and

Slovenia had those with the highest scores of all calculated indices (bird, GAR, RES, and PES)

(e.g. the Plzeň region in Czechia reached the best categories in all variables but the bird index),

while Romania had that with the lowest (the Vrancea county in Romania fall into the worst

categories in all variables). All predominantly urbanised areas were categorised relatively high

in all indices with Slovakia being the best (Bratislava region) and Poland and Romania being

the worst (Silesian Voivodeship, București-Ilfov region).

Fig. 5: The relationship of the bird (A), garden (B), respondent (C), and pesticide (D)

indices with the urban-rural typology index based on NUTS-3 areas (except for Poland, in

476   which NUTS-2 were used). The colours represent the levels of urban-rural typology, as

477   follows: red – predominantly urban, yellow – intermediate, green – predominantly rural. The

478   transparency of the colour indicates the categorised index value (bird/GAR/RES/PES) on a

479   three-level scale (divided into categorised at the 0.33 and 0.66 quantiles). The grey colour

480   indicates NUTS areas from which we had less than five respondents (n = 20).

4. Discussion

Domestic gardens offer significant opportunities for advancing global biodiversity and

sustainability targets. Our work investigated the intertwined relationship between

sociodemographic parameters, sustainable garden management, and biodiversity conservation

from a highly understudied area of the European Union and highlights that there are significant

differences among countries, and even within countries, regardless of the scale, in the examined

parameters.

Indeed, the participating countries differed in all three examined indices, with having the least

differences in GAR and the most in PES, and these differences were supported by both the

pairwise comparisons between countries and the GMB model. There were countries

experiencing either equally low or high values on all indices, such as Romania faring poorly in

all indices, whereas Slovakia showing overly high values. Sometimes indices implied

counteracting effects, such as a high GAR index, suggesting a great prospect for maintaining

high biodiversity in domestic gardens, occurred together with high pesticide use in Hungary.

This is mostly in line with Coisnon et al. (2019) who found similar trade-offs in variables

influencing sustainable gardening practices such as Hungary scoring the worst in leaving spaces

for wildlife and avoiding the use of pesticides, yet having been categorised as the best in

selecting plants that provide food for birds and pollinators and avoiding the introduction of

potentially invasive plants. Other countries such as Romania generally ranked poorly in both

our study and known from the literature (Coisnon et al., 2019). All three indices varied widely,

however, indicating the multidimensional nature of factors driving sustainable gardening.

The extensive application of pesticides beyond the agricultural sector, including domestic

gardens, and the inter-country disparities in pesticide usage within Europe are critical issues for

biodiversity-friendly gardens (Poniży et al., 2021; Varga-Szilay et al., 2024; Varga-Szilay &

Pozsgai, 2022). Our study found substantial differences between countries in this regard, with

20–30% of respondents in Hungary and Romania thinking that the use of pesticides is either

important or crucially important for their gardening, a claim with which only 2% of those from

Slovakia agreed (**Supplementary Material Fig. 9**.).

There is a disparity though between pesticide use and the other two indices we measured.

Although nearly 70% of respondents from Romania considered the widespread use of pesticides

as one of the most threatening factors to pollinators, their average pesticide use was still the

least favourable in our study. Indeed, what gardeners perceive as a biodiversity-friendly garden

may not necessarily align with its real biodiversity potential, suggesting that self-reporting

surveys may be misleading in this term, unless gardens are examined from multiple aspects,

including sociodemographic or cultural factors.

Indeed, in line with Larson et al. (2022), those who perceived gardening as a hobby were more

inclined to change their gardens to be attractive and beneficial to biodiversity, and we also

discovered that garden owners who dedicate more time spent with gardening tend to maintain

gardens with higher biodiversity potential. Our results corroborate with the findings of both

Philpott et al. (2020) and Lin et al. (2017) who indicated that time spent in domestic gardens

positively correlates with plant species richness, higher dissimilarity in crop composition, and

high levels of nature-relatedness. Moreover, our findings agree with those of Geppert et al.

(2024) who reported that the time spent outdoors increased the willingness to pro-pollinator

actions. In our study, however, both when gardening was associated with a hobby and more

time was allocated for gardening were characterised by increased pesticide use. Therefore, there

is a higher possibility for these gardeners to create an ecological trap in these gardens for

visiting organisms (especially insects) (Varga-Szilay & Pozsgai, 2022). There is, however, a

noticeable contrast in behaviour based on gender and parental status (**Fig. 4C**) where women

and respondents with children improve the outcome of the PES index, whilst this separation

was unclear with the other two indices.

Although the country identity proved to be the most important variable in predicting the

potential for maintaining high garden biodiversity, the respondents' attitudes towards supporting

biodiversity (especially pollinators) in their gardens, pesticide use, owners' perception, the

long-term gardening experience and daily management also proved to be important factors.

Furthermore, nature conservation was the prime aspect that influenced gardening habits in four

participating countries, albeit in three of these this did not mean a lower pesticide use, and only

Croatia and Poland were the ones where more than 80% of the respondents felt that nature

conservation does not affect their gardening habits at all (**Supplementary Material Fig. 10**).

Increasing age (especially that over 65) negatively influenced both the potential of gardens for

maintaining high biodiversity and the owners' attitudes toward maintaining biodiversity-

friendly gardens, which suggests the elderly are more inclined to follow and uphold

longstanding conventional practices, regardless of whether or not they are beneficial for

ecological sustainability. Whether these decisions are driven either by the difficulties the elderly

may have with manoeuvring among the continuous flow of complex ecological issues which

were not highlighted during most of their lives, the scarcity of the available information from

the more traditional information channels they use, or some alternative explanations is yet to

be clarified. Nonetheless, the age-independent pesticide use we observed in the GBM model

suggests that even when information is broadly available behavioural change is not granted.

However, gardening practices and their associated sociocultural and economic variables exhibited significant diversity among garden owners and the final outcome for biodiversity-friendliness appeared to be a series of multifactor decisions depending on numerous background drivers (Gifford & Nilsson, 2014). For instance, despite several studies highlighting a substantially positive effect of GDP per capita and household income on sustainable gardening practices and pollinator abundance in domestic gardens (e.g. Baldock et al., 2019; Coisnon et al., 2019), in our GBM model, PPS did not prove to be a significant explanatory variable for any of our indices.

Despite the existing differences among the NUTS regions and countries in all three indices, we did not find strong geographical correlations; neither latitude nor longitude had a high explanatory power in our GBM models. Nonetheless, longitude had a higher influence on the GAR index than latitude, which might not be as expected since the north-south gradient exhibits steeper changes in climate and habitats, than the one from east to west. This greater importance of the east-west axis could be explained by the higher impact of Western European countries (and western culture) on how more westerly regions in CEE perceived biodiversity issues after political and societal changes in the 1990s. Conversely, the lower influence of longitude and latitude on the RES and PES indices could be attributed to the continued strong connection to traditional practices across the studied countries.

Our results show that merely examining the structural diversity of private gardens, sociodemographic parameters, garden owners' motivation for gardening, or biodiversity-negative activities (e.g. pesticide use) individually may be insufficient for obtaining a reliable proxy of whether domestic gardens could potentially maintain/build high diversity in human-altered areas. Moreover, the lack of a clear geographical pattern and within-country diversity in all indices suggests that regionally tailored rather than countrywide strategies are needed to identify pathways toward sustainable green infrastructure and to improve the natural value of

domestic gardens. Since these strategies must be underpinned by interdisciplinary study efforts,

our study can offer fundamental insight into how they can be optimised and where the greatest

rewards can be gained. Indeed, despite the dominance of moderate values of our indices on the

map, local strategies can still be developed based on the preliminary classification.

Areas with generally **low urbanisation** and also **low biodiversity** are likely to be agricultural-

and farmlands in which gardens with high biodiversity can act as habitat islands, ecological

corridors, or stepping stones which facilitate the movement of wildlife through unsuitable

agricultural areas. In these areas, improving the structural complexity of the gardens and

reducing pesticide use could be the most beneficial for biodiversity. Disseminating knowledge

about ecological-friendly practices and sustainable gardening habits would also be key. Indeed,

the predominantly rural areas in northeast Poland, where both bird index and urbanisation are

low, also suffer from low RES. However, information exchange here may need to rely more on

interpersonal relationships or traditional media than online channels (Troumbis, 2021) and

overall biodiversity is likely to be more impacted by agricultural rather than garden practices.

Attempting to improve biodiversity value in areas with generally **low urbanisation** levels and

high biodiversity probably has the least merit, yet reducing pesticide use can prohibit

degradation and may form garden-level biodiversity islands that increase landscape complexity.

Moreover, maintaining and even improving garden diversity in these areas, such as

predominantly rural areas in the northern part of Croatia and Slovenia, can be particularly

important as good examples for environmental education and maintaining source populations

of wildlife for adjacent but less favourable areas.

Thus, at low urbanisation levels improving aspects of the RES would be the priority, along with

a substantial decrease of pesticide use. With increasing biodiversity, gardens will benefit from

improved conservation biological control (Lacey et al., 2015; Landis et al., 2000; Quesada-

Moraga et al., 2022), which may further decrease pesticide use.

Biodiversity-friendly activities can achieve fast and great ecological benefits in areas where

urbanisation is **high** and **biodiversity** is **moderate or low**. Although their biodiversity

potential will remain limited, prioritising increasing gardens' structural and plant diversity will

enable them to serve as ecological corridors or stepping stones for most wildlife to navigate

around highly urbanised areas. Although biodiversity will unlikely increase substantially, the

high visibility of biodiversity in densely populated areas has an outstanding educational value.

Even though few CEE regions fulfil these criteria (e.g. the central part of Czechia), most

Western European areas predominantly classified as urban are likely to fall into this category.

Although **high urbanisation** and **high biodiversity** rarely co-occur, some areas in CEE, such

as Northern Estonia, exhibit such a phenomenon. This offers an opportunity for the

dissemination of activities tied to increasing the RES index (e.g. through campaigns and species

identification trainings), as they can quickly reach many individuals and engage them in

enhancing garden biodiversity. Additionally, these efforts provide a substantial amount of data

for biodiversity monitoring, resulting in additional benefits.

Thus, at high urbanisation levels, increasing gardens' structural diversity and other aspects

beneficial for maintaining high biodiversity are key but improving garden owners' attitudes

towards biodiversity-friendly gardening and reconnecting them to nature are also important.

Since the level of urbanisation across Europe is projected to reach 83.7% by 2025 (United

Nations, 2018), this improvement in attitude is necessary to exploit the potential of domestic

gardens to mitigate large-scale biodiversity loss. However, due to the ubiquitous use of

pesticides in CEE, improving the PES index would be equally necessary in every area, both

urban and rural.

Adequate planning, at least, at the regional scale, with the active involvement of garden owners,

is crucial for increasing the biodiversity value of domestic gardens at a larger scale. For this,

future nature conservation planning strategies and studies should acknowledge the variability

of domestic garden owners' attitudes and gardening practices, as well as the domestic gardens'

potential for biodiversity conservation (Cameron et al., 2012).

4.1 Study limits

Although in our work we collected over 5255 responses, this only represents a small fraction

of the inhabitants and gardeners of the focal area of our study. Moreover, the dissemination

channels we used do not allow random respondent selection and our selection process is most

likely to be biased towards those who had regular access to the Internet, whilst those living in

infrastructurally less developed parts of the countries had a lower chance to learn about our call.

Indeed, parts of Poland and the Southern part of Romania are highly underrepresented. Yet,

several countries like Estonia and Hungary, have a high coverage which likely portrays

faithfully the common gardening behaviours in those regions.

In an effort to engage a broad audience in several languages, we often compromised

completeness for simplicity and brevity. Whilst we believe this did not affect the interpretability

of our questionnaire, it definitely sets limits on the use of scientific terminology and how

detailed information we could gain with some questions.

Although our questionnaire aimed for the response of only one person (the garden owner) for

each domestic garden, these gardens are probably cared for by more than one person (e.g. family

members), and thus, they are not only influenced by one person's motivation but by severals'.

4.2 Future perspectives

646 Involving more participants from the surveyed CEE countries, standardising the questions and

647 expanding the area of interest to all European countries, as well as repeating the same survey

648 within a reasonable time frame would all be highly advantageous. Especially investigating the

649 distinction between the gardening habits in Western and Central-Eastern European regions, as

well as those from the Balkan, such as Serbia and Bulgaria, holds particular potential. While addressing privacy concerns may present challenges, extending our study with measures of biodiversity and landscape configuration from each garden could introduce extra layers of complexity to our exploration.

Furthermore, a direct link between domestic garden diversity, and the potential of thereof, and biodiversity conservation should be established and implemented into urban green space management.

5. Conclusions

While gardens may not replace undisturbed natural habitats, it is imperative to reconsider maintaining high levels of urban and rural biodiversity as valuable complements for increasing the area of semi-natural habitat patches and creating a network of ecological corridors and stepping stones (Beninde et al., 2015). In our study, we, however, show that the potential of domestic gardens for maintaining high biodiversity varies widely among CEE regions, and along with it does their conservation value. Therefore, to maximise this value improving several aspects of CEE gardening practices in concert is necessary. The variability both between and within participating countries, underscores the need for approaches tailored to regional circumstances rather than unified regulations across European countries.

Despite some differences observed between north to south, the idea of promoting biodiversity and nature conservation seems to be spreading from the west towards the east and therefore should continue with additional efforts towards the eastern countries. Besides promoting the importance of biodiversity, some concrete measures in agricultural practice and activities should be promoted as well and offered to gardeners to speed up in situ changes, as gardeners still seem to incline more to traditional approaches. The nature-friendly practice seems to be

supported more by the higher education levels and by the factor of age, or parenting, those facts

can be used to target the promotion of gardening in nature conservation in this part of Europe.

On the other hand, to ensure that domestic gardens truly contribute positively to sustainable

urban environments, the multifaceted issue of pesticide usage that permeates various aspects of

gardening, from food production in allotments to the care of ornamental plants and flea

treatments for pets, must be uniformly addressed across all European Union countries.

Similarly, information for providing a comprehensive understanding among the public

regarding the potential environmental and human health impacts of pesticides should also be

made broadly available.

Biodiversity-friendly gardens also improve human health and well-being (Samus et al., 2022)

and provide urban residents with the opportunity to reconnect to the natural world (Cameron et

al., 2012). Thus, while increasing the diversity of domestic gardens alone may not solve the

biodiversity crisis, by creating ecological corridors it can contribute to conservation, foster a

shift in human attitudes towards preserving natural areas, and enhance the ecological quality of

urban environments. To fully exploit all benefits of domestic gardens and to increase people's

knowledge concerning the environment, population-wide, regular, and high-quality education

and community programs which, shape people's attitudes, willingness to take action, and

interest, are needed (Knapp et al., 2021; Shwartz et al., 2012; Sturm et al., 2021; Tikka et al.,

2000). Similar programs have been running for decades in Western European countries but they

are still scarce in CEE (European Commission, Directorate-General for Communication, 2015;

Jehlička & Jacobsson, 2021). Yet, the need for improving the ecological quality of domestic

gardens is not unique to Western Europe, where it gets the most emphasis, but is also present in

CEE. Hence, local strategies for environmental education emphasising the importance of

biodiversity-friendly gardening should be developed for CEE countries as well. These,

however, should be tailored to local needs and provide access to biodiversity-related

information through a broad variety of channels to reach all strata of society.

Besides serving as a proxy to indicate gardens' environmental quality, our study also can help

in accessing these gardens' potential in designing eco-networks of biodiversity-friendly

gardens. It also can act as a tool for facilitating the decision for optimal strategies to maximise

the environmental benefits of domestic gardens.

Author information

Email address of the authors

- Zsófia Varga-Szilay – zsofia@vargaszilay.hu
- Arvīds Barševskis – arvids.barsevskis@du.lv
- Klára Benedek – benedekklara@ms.sapientia.ro
- Danilo Bevk – danilo.bevk@nib.si
- Agata Jojczyk – agata_jojczyk@sggw.edu.pl
- Anton Krištín – kristin@ife.sk
- Jana Růžičková – jr.tracey@seznam.cz
- Lucija Šerić Jelaska – lucija.seric.jelaska@biol.pmf.hr
- Eve Veromann – eve.veromann@emu.ee
- Silva Vilumets – silva.vilumets@emu.ee
- Kinga Gabriela Fetykó – kingha221@gmail.com
- Gergely Szövényi – szovenyig@gmail.com
- Gábor Pozsgai – pozsgaig@coleoptera.hu

ORCID number of authors

- Zsófia Varga-Szilay: 0000-0001-9712-7654
- Arvīds Barševskis: 0000-0001-9703-0115
- Klára Benedek: 0009-0002-1936-3541
- Danilo Bevk: 0000-0002-5715-3089
- Agata Jojczyk: 0000-0003-4993-3226
- Anton Krištín: 0000-0001-7422-6393
- Jana Růžičková: 0000-0001-9703-4538
- Lucija Šerić Jelaska: 0000-0003-1840-7119
- Eve Veromann: 0000-0002-0977-5270
- Silva Vilumets: 0000-0002-4060-9596
- Kinga Gabriela Fetykó: 0000-0003-4150-6385
- Gergely Szövényi: 0000-0001-9632-4066
- Gábor Pozsgai: 0000-0002-2300-6558

CRediT authorship contribution statement

- **Zsófia Varga-Szilay:** Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Software; Validation; Visualization; Roles/Writing - original draft; and Writing - review & editing

- **Arvīds Barševskis:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Klára Benedek:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Danilo Bevk:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Agata Jojczyk:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Anton Krištín:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Jana Růžičková:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Lucija Šerić Jelaska:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Eve Veromann:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Silva Vilumets:** Data curation; Investigation; Methodology; Project administration; and Writing - review & editing

- **Kinga Gabriela Fetykó:** Conceptualization; Data curation; Investigation; Methodology; Project administration; Validation; and Writing - review & editing

- **Gergely Szövényi:** Conceptualization; Data curation; Investigation; Methodology; Project administration; Supervision; Validation; and Writing - review & editing

- **Gábor Pozsgai:** Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Software; Supervision; Validation; Visualization; Roles/Writing - original draft; and Writing - review & editing

Funding source

- DB was supported by the Slovenian Research Agency (research core funding No.P1-0255) and FRACTAL (Interreg Alpine Space).

–   AK was supported by the Slovak Scientific Grant Agency VEGA 2/0097/23.

–   EV was supported by the Estonian Research Council grant PRG1056.

–   LSJ was supported by The Croatian Science Foundation, MEDITERATRI project (HRZZ
UIP 2017-05-1046).

–   GP was supported by the project Open Access FCT-UIDP/00329/2020-2024 (Thematic
Line 1 – integrated ecological assessment of environmental change on biodiversity,
https://doi.org/10.54499/UIDB/00329/2020) and by the Azores DRCT Pluriannual
Funding (M1.1.A/FUNC.UI&D/010/2021-2024).

Declarations

778 Conflict of interest
779 The authors declare no competing interests.

Data and code availability

782 The data and the underlying computer code is available in the GitHub repository
783 https://github.com/zsvargaszilay/domestic_gardens_in_CEE

Acknowledgements

786 We are grateful to Geoff Gurr and Stephen Venn for their comments on an earlier version of
787 the manuscript.

References

Aguilera, G., et al. (2019). Intensive management reduces butterfly diversity over time in urban green spaces. *Urban Ecosystems*, *22*(2), 335–344. https://doi.org/10.1007/s11252-018-0818-y

Alber, J., & Kohler, U. (2008). Informal food production in the enlarged European Union. *Social Indicators Research*, *89*(1), 113–127. https://doi.org/10.1007/s11205-007-9224-1

ArcGIS Data and Maps. (2022a). *Europe NUTS 2 Demographics and Boundaries—Overview*. https://www.arcgis.com/sharing/rest/content/items/48f0d1a2006c4e89a85712a618fcff89 (accessed on 16 July 2024)

ArcGIS Data and Maps. (2022b). *Europe NUTS 3 Demographics and Boundaries—Overview*. https://www.arcgis.com/home/item.html?id=5eb88decbff54520b49c785241c07bf4 (accessed on 16 July 2024)

Baldock, K. C. R. (2020). Opportunities and threats for pollinator conservation in global towns and cities. *Current Opinion in Insect Science*, *38*, 63–71. https://doi.org/10.1016/j.cois.2020.01.006

Baldock, K. C. R., et al. (2019). A systems approach reveals urban pollinator hotspots and conservation opportunities. *Nature Ecology & Evolution*, *3*(3), Article 3. https://doi.org/10.1038/s41559-018-0769-y

Bengtsson, J., et al. (2000). Biodiversity, disturbances, ecosystem function and management of European forests. *Forest Ecology and Management*, *132*(1), 39–50. https://doi.org/10.1016/S0378-1127(00)00378-9

Beninde, J., Veith, M., & Hochkirch, A. (2015). Biodiversity in cities needs space: A meta-analysis of factors determining intra-urban biodiversity variation. *Ecology Letters*, *18*(6), 581–592. https://doi.org/10.1111/ele.12427

Beumer, C. (2018). Show me your garden and I will tell you how sustainable you are: Dutch citizens' perspectives on conserving biodiversity and promoting a sustainable urban living environment through domestic gardening. *Urban Forestry & Urban Greening*, *30*, 260–279. https://doi.org/10.1016/j.ufug.2017.09.010

Borysiak, J., Mizgajski, A., & Speak, A. (2017). Floral biodiversity of allotment gardens and its

contribution to urban green infrastructure. *Urban Ecosystems*, *20*(2), 323–335.

https://doi.org/10.1007/s11252-016-0595-4

Braschler, B., et al. (2020). Ground-dwelling invertebrate diversity in domestic gardens along a rural-

urban gradient: Landscape characteristics are more important than garden characteristics.

PLOS ONE, *15*(10), e0240061. https://doi.org/10.1371/journal.pone.0240061

Cameron, R. W. F., et al. (2012). The domestic garden – Its contribution to urban green infrastructure.

Urban Forestry & Urban Greening, *11*(2), 129–137.

https://doi.org/10.1016/j.ufug.2012.01.002

Camps-Calvet, M., et al. (2016). Ecosystem services provided by urban gardens in Barcelona, Spain:

Insights for policy and planning. *Environmental Science & Policy*, *62*, 14–23.

https://doi.org/10.1016/j.envsci.2016.01.007

Chamberlain, S., et al. (2024). *rgbif: Interface to the global biodiversity information facility API*.

https://CRAN.R-project.org/package=rgbif

Chollet, S., et al. (2018). From urban lawns to urban meadows: Reduction of mowing frequency

increases plant taxonomic, functional and phylogenetic diversity. *Landscape and Urban*

Planning, *180*, 121–124. https://doi.org/10.1016/j.landurbplan.2018.08.009

Church, A., et al. (2015). 'Growing your own': A multi-level modelling approach to understanding

personal food growing trends and motivations in Europe. *Ecological Economics*, *110*, 71–80.

https://doi.org/10.1016/j.ecolecon.2014.12.002

Coisnon, T., Rousselière, D., & Rousselière, S. (2019). Information on biodiversity and environmental

behaviors: A European study of individual and institutional drivers to adopt sustainable

gardening practices. *Social Science Research*, *84*, 102323.

https://doi.org/10.1016/j.ssresearch.2019.06.014

Concepción, E. D., et al. (2020). Optimizing biodiversity gain of European agriculture through

regional targeting and adaptive management of conservation tools. *Biological Conservation*,

241, 108384. https://doi.org/10.1016/j.biocon.2019.108384

Delahay, R. J., et al. (2023). Biodiversity in residential gardens: A review of the evidence base.

Biodiversity and Conservation, *32*(13), 4155–4179. https://doi.org/10.1007/s10531-023-

02694-9

Dewaelheyns, V., Kerselaers, E., & Rogge, E. (2016). A toolbox for garden governance. *Land Use*

Policy, *51*, 191–205. https://doi.org/10.1016/j.landusepol.2015.11.016

Dobrovodská, M., et al. (2023). Factors affecting the biodiversity of historical landscape elements:

Detailed analyses from three case studies in Slovakia. *Environmental Monitoring and*

Assessment, *195*(6), 674. https://doi.org/10.1007/s10661-023-11035-w

Donkersley, P., et al. (2023). A little does a lot: Can small-scale planting for pollinators make a

difference? *Agriculture, Ecosystems & Environment*, *343*, 108254.

https://doi.org/10.1016/j.agee.2022.108254

European Commission. (2022). *Pioneering proposals to restore Europe's nature by 2050*. European

Commission. https://ec.europa.eu/commission/presscorner/detail/en/ip_22_3746

European Commission. (2024). *Nature Restoration Law*. https://eur-lex.europa.eu/legal-

content/EN/TXT/HTML/?uri=OJ:L_202401991

European Commission, Directorate-General for Communication. (2015). *Special Eurobarometer 436:*

Attitudes of Europeans towards biodiversity (Version 1.00) [Dataset].

http://data.europa.eu/88u/dataset/S2091_83_4_436_ENG (accessed on 16 July 2024)

European Environment Agency. (2012). *Protected areas in Europe – An overview*.

https://www.eea.europa.eu/publications/protected-areas-in-europe-2012 (accessed on 16 July

2024)

European Environment Agency. (2022). *How green are European cities? Green space key to well-*

being – but access varies. https://www.eea.europa.eu/highlights/how-green-are-european-

cities (accessed on 16 July 2024)

European Environment Agency. (2023). *Net land take in cities and commuting zones in Europe*.

https://www.eea.europa.eu/en/analysis/indicators/net-land-take-in-cities (accessed on 16 July

2024)

Eurostat. (2021). *NUTS - Nomenclature of territorial units for statistics – Overview*.

https://ec.europa.eu/eurostat/web/nuts (accessed on 16 July 2024)

Eurostat. (2022a). *Regional gross domestic product (PPS per inhabitant) by NUTS 2 regions* [Dataset].

https://doi.org/10.2908/TGS00005 (accessed on 16 July 2024)

Eurostat. (2022b). *Urban-rural Europe – Introduction*. https://ec.europa.eu/eurostat/statistics-

explained/index.php?title=Urban-rural_Europe_-_introduction (accessed on 16 July 2024)

Eurostat. (2024). *Rural development – Methodology*. https://ec.europa.eu/eurostat/web/rural-

development/methodology (accessed on 16 July 2024)

Fischer, L. K., et al. (2020). Public attitudes toward biodiversity-friendly greenspace management in

Europe. *Conservation Letters*, *13*(4), e12718. https://doi.org/10.1111/conl.12718

Fontaine, B., et al. (2016). Impact of urbanization and gardening practices on common butterfly

communities in France. *Ecology and Evolution*, *6*(22), 8174–8180.

https://doi.org/10.1002/ece3.2526

Freeman, C., et al. (2012). "My garden is an expression of me": Exploring householders' relationships

with their gardens. *Journal of Environmental Psychology*, *32*(2), 135–143.

https://doi.org/10.1016/j.jenvp.2012.01.005

Galluzzi, G., Eyzaguirre, P., & Negri, V. (2010). Home gardens: Neglected hotspots of agro-

biodiversity and cultural diversity. *Biodiversity and Conservation*, *19*(13), 3635–3654.

https://doi.org/10.1007/s10531-010-9919-5

GBIF.org. (2024). *GBIF Home Page*. GBIF.Org. https://www.gbif.org

Geppert, C., et al. (2024). Willingness of rural and urban citizens to undertake pollinator conservation

actions across three contrasting European countries. *People and Nature*, *6*(4), 1502–1511.

https://doi.org/10.1002/pan3.10656

Gifford, R., & Nilsson, A. (2014). Personal and social factors that influence pro-environmental

concern and behaviour: A review. *International Journal of Psychology*, *49*(3), 141–157.

https://doi.org/10.1002/ijop.12034

Glavan, M., et al. (2018). The economic performance of urban gardening in three European cities –

examples from Ljubljana, Milan and London. *Urban Forestry & Urban Greening*, *36*, 100–

122. https://doi.org/10.1016/j.ufug.2018.10.009

Goddard, M. A., Dougill, A. J., & Benton, T. G. (2010). Scaling up from gardens: Biodiversity

conservation in urban environments. *Trends in Ecology & Evolution*, *25*(2), 90–98.

https://doi.org/10.1016/j.tree.2009.07.016

Goddard, M. A., Dougill, A. J., & Benton, T. G. (2013). Why garden for wildlife? Social and

ecological drivers, motivations and barriers for biodiversity management in residential

landscapes. *Ecological Economics*, *86*, 258–273.

https://doi.org/10.1016/j.ecolecon.2012.07.016

Greenwell, B., et al. (https://github.com/gbm-developers), G. B. M. (2022). *gbm: Generalized Boosted*

Regression Models (Version 2.1.8.1) [Software]. https://cran.r-

project.org/web/packages/gbm/index.html

Griffiths-Lee, J., Nicholls, E., & Goulson, D. (2022). Sown mini-meadows increase pollinator

diversity in gardens. *Journal of Insect Conservation*, *26*(2), 299–314.

https://doi.org/10.1007/s10841-022-00387-2

Haenke, S., et al. (2009). Increasing syrphid fly diversity and density in sown flower strips within

simple vs. complex landscapes. *Journal of Applied Ecology*, *46*(5), 1106–1114.

https://doi.org/10.1111/j.1365-2664.2009.01685.x

Hagen, M., et al. (2012). 2—Biodiversity, species interactions and ecological networks in a

fragmented world. In U. Jacob & G. Woodward (Eds.), *Advances in Ecological Research* (Vol.

46, pp. 89–210). Academic Press. https://doi.org/10.1016/B978-0-12-396992-7.00002-2

Hanson, H. I., et al. (2021). Gardens' contribution to people and urban green space. *Urban Forestry &*

Urban Greening, *63*, 127198. https://doi.org/10.1016/j.ufug.2021.127198

Harris, C. R., et al. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362.

https://doi.org/10.1038/s41586-020-2649-2

Heidt, V., & Neef, M. (2008). Benefits of urban green space for improving urban climate. In M. M.

Carreiro, Y.-C. Song, & J. Wu (Eds.), *Ecology, Planning, and Management of Urban Forests:*

International Perspectives (pp. 84–96). Springer. https://doi.org/10.1007/978-0-387-71425-

7_6

Home, R., et al. (2019). Effects of garden management practices, by different types of gardeners, on

human wellbeing and ecological and soil sustainability in Swiss cities. *Urban Ecosystems*,

22(1), 189–199. https://doi.org/10.1007/s11252-018-0806-2

Hruška, V., Smutná, Z., & Vávra, J. (2021). Why I would want to live in the village if I was not

interested in cultivating the plot? A study of home gardening in rural Czechia. *Sustainability*,

13, 1–21. https://doi.org/10.3390/su13020706

Jehlička, P., et al. (2021). Beyond hardship and joy: Framing home gardening on insights from the

European semi-periphery. *Geoforum*, *126*, 150–158.

https://doi.org/10.1016/j.geoforum.2021.05.018

Jehlička, P., et al. (2020). Thinking food like an East European: A critical reflection on the framing of

food systems. *Journal of Rural Studies*, *76*, 286–295.

https://doi.org/10.1016/j.jrurstud.2020.04.015

Jehlička, P., & Jacobsson, K. (2021). The importance of recognizing difference: Rethinking Central

and East European environmentalism. *Political Geography*, *87*, 102379.

https://doi.org/10.1016/j.polgeo.2021.102379

Jongman, R. H. G. (2002). Homogenisation and fragmentation of the European landscape: Ecological

consequences and solutions. *Landscape and Urban Planning*, *58*(2), 211–221.

https://doi.org/10.1016/S0169-2046(01)00222-5

Kabisch, N., & Haase, D. (2013). Green spaces of European cities revisited for 1990–2006. *Landscape*

and Urban Planning, *110*, 113–122. https://doi.org/10.1016/j.landurbplan.2012.10.017

Keshavarz, N., et al. (2016). *A history of urban gardens in Europe* (pp. 8–32).

https://doi.org/10.4324/9781315686608-2

Knapp, J. L., et al. (2021). Socio-psychological factors, beyond knowledge, predict people's

engagement in pollinator conservation. *People and Nature*, *3*(1), 204–220.

https://doi.org/10.1002/pan3.10168

Krištín, A., Hoi, H., & Kaňuch, P. (2024). Local population decline of the threatened lesser grey shrike

Lanius minor is linked to the modernisation of the rural landscape. *Bird Conservation*

International, *34*, e18. https://doi.org/10.1017/S095927092400011X

Krols, J., et al. (2022). Residential green space, gardening, and subjective well-being: A cross-

sectional study of garden owners in northern Belgium. *Landscape and Urban Planning*, *223*,

104414. https://doi.org/10.1016/j.landurbplan.2022.104414

Kuhn, M., et al. (2023). *caret: Classification and Regression Training* (Version 6.0-94) [Software].

https://cran.r-project.org/web/packages/caret/index.html

Lacey, L. A., et al. (2015). Insect pathogens as biological control agents: Back to the future. *Journal of*

Invertebrate Pathology, *132*, 1–41. https://doi.org/10.1016/j.jip.2015.07.009

Landis, D. A., Wratten, S. D., & Gurr, G. M. (2000). Habitat management to conserve natural enemies

of arthropod pests in agriculture. *Annual Review of Entomology*, *45*(Volume 45, 2000), 175–

201. https://doi.org/10.1146/annurev.ento.45.1.175

Larson, K. L., et al. (2022). Examining the potential to expand wildlife-supporting residential yards

and gardens. *Landscape and Urban Planning*, *222*, 104396.

https://doi.org/10.1016/j.landurbplan.2022.104396

Lerman, S. B., et al. (2018). To mow or to mow less: Lawn mowing frequency affects bee abundance

and diversity in suburban yards. *Biological Conservation*, *221*, 160–174.

https://doi.org/10.1016/j.biocon.2018.01.025

Li, M., Remme, R., Bodegom, P., & van Oudenhoven, A. (2023). *Solution to what? Global review of*

nature-based solutions, urban challenges, and outcomes.

https://doi.org/10.1101/2023.12.07.570577

Liang, H., et al. (2023). The effects of urbanization on pollinators and pollination: A meta-analysis.

Ecology Letters, *26*(9), 1629–1642. https://doi.org/10.1111/ele.14277

Lin, B. B., et al. (2017). How green is your garden?: Urban form and socio-demographic factors

influence yard vegetation, visitation, and ecosystem service benefits. *Landscape and Urban*

Planning, *157*, 239–246. https://doi.org/10.1016/j.landurbplan.2016.07.007

Lin, B. B., Philpott, S. M., & Jha, S. (2015). The future of urban agriculture and biodiversity-

ecosystem services: Challenges and next steps. *Basic and Applied Ecology*, *16*(3), 189–201.

https://doi.org/10.1016/j.baae.2015.01.005

Lindemann-Matthies, P., & Marty, T. (2013). Does ecological gardening increase species richness and

aesthetic quality of a garden? *Biological Conservation*, *159*, 37–44.

https://doi.org/10.1016/j.biocon.2012.12.011

Lundberg, S. M., & Lee, S.-I. (2017). A unified approach to interpreting model predictions. *Advances*

in Neural Information Processing Systems, *30*.

https://proceedings.neurips.cc/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-

Abstract.html

Majewska, A. A., & Altizer, S. (2020). Planting gardens to support insect pollinators. *Conservation*

Biology, *34*(1), 15–25. https://doi.org/10.1111/cobi.13271

Marques, P., et al. (2021). Home gardens can be more important than other urban green infrastructure

for mental well-being during COVID-19 pandemics. *Urban Forestry & Urban Greening*, *64*,

127268. https://doi.org/10.1016/j.ufug.2021.127268

Mayer, M., & Stando, A. (2023). *shapviz: SHAP Visualizations* (Version 0.9.2) [Software].

https://cran.r-project.org/web/packages/shapviz/index.html

Mayer, M., Watson, D., & Biecek, P. (2023). *kernelshap: Kernel SHAP* (Version 0.4.0) [Software].

https://cran.r-project.org/web/packages/kernelshap/index.html

Muller, N., Werner, P., & Kelcey, J. G. (2010). *Urban Biodiversity and Design*. John Wiley & Sons.

Muratet, A., & Fontaine, B. (2015). Contrasting impacts of pesticides on butterflies and bumblebees in

private gardens in France. *Biological Conservation*, *182*, 148–154.

https://doi.org/10.1016/j.biocon.2014.11.045

Pebesma, E. (2018). Simple features for r: Standardized support for spatial vector data. *The R Journal*,

10(1), 439–446. https://doi.org/10.32614/RJ-2018-009

Perpiña Castillo, C., et al. (2019). *Main land-use patterns in the EU within 2015-2030* (JRC115895;

JRC Policy Insights). European Commission, Ispra, 2019,

https://publications.jrc.ec.europa.eu/repository/handle/JRC115895 (accessed on 16 July 2024)

Petzke, N., König, B., & Bokelmann, W. (2021). Plant protection in private gardens in Germany:

Between growing environmental awareness, knowledge and actual behaviour. *European*

Journal of Horticultural Science, *86*, 59–68. https://doi.org/10.17660/eJHS.2021/86.1.7

Philpott, S. et al., (2020). Gardener demographics, experience, and motivations drive differences in

plant species richness and composition in urban gardens. *Ecology and Society*, *25*(4).

https://doi.org/10.5751/ES-11666-250408

Poniży, L., et al. (2021). The rich diversity of urban allotment gardens in Europe: Contemporary

rrends in the context of historical, socio-economic and legal conditions. *Sustainability*, *13*(19),

Article 19. https://doi.org/10.3390/su131911076

Pourias, J., Aubry, C., & Duchemin, E. (2016). Is food a motivation for urban gardeners?

Multifunctionality and the relative importance of the food function in urban collective gardens

of Paris and Montreal. *Agriculture and Human Values*, *33*(2), 257–273.

https://doi.org/10.1007/s10460-015-9606-y

Prendergast, K. S., et al. (2022). Urban native vegetation remnants support more diverse native bee

communities than residential gardens in Australia's southwest biodiversity hotspot. *Biological*

Conservation, *265*, 109408. https://doi.org/10.1016/j.biocon.2021.109408

Proske, A., Lokatis, S., & Rolff, J. (2022). Impact of mowing frequency on arthropod abundance and

diversity in urban habitats: A meta-analysis. *Urban Forestry & Urban Greening*, *76*, 127714.

https://doi.org/10.1016/j.ufug.2022.127714

Python Software Foundation. (2019). *Python Language Reference* (Version 3.8) [Software].

http://www.python.org

Quesada-Moraga, E., et al. (2022). Multitrophic interactions of entomopathogenic fungi in BioControl.

BioControl, *67*(5), 457–472. https://doi.org/10.1007/s10526-022-10163-5

Quistberg, R. D., Bichier, P., & Philpott, S. M. (2016). Landscape and local correlates of bee

abundance and species richness in urban gardens. *Environmental Entomology*, *45*(3), 592–601.

https://doi.org/10.1093/ee/nvw025

R Core Team. (2021). *R: A language and environment for statistical computing* (Version 4.4)

[Software]. https://www.R-project.org/

Ramalho, C. E., & Hobbs, R. J. (2012). Time for a change: Dynamic urban ecology. *Trends in Ecology*

& Evolution, *27*(3), 179–188. https://doi.org/10.1016/j.tree.2011.10.008

Revelle, W. (2021). *psych: Procedures for psychological, psychometric, and personality research*

(Version 2.4.6) [Software]. https://CRAN.R-project.org/package=psych

Samus, A., et al. (2022). Relationships between nature connectedness, biodiversity of private gardens,

and mental well-being during the Covid-19 lockdown. *Urban Forestry & Urban Greening*, *69*,

127519. https://doi.org/10.1016/j.ufug.2022.127519

Seitz, B., et al. (2022). Land sharing between cultivated and wild plants: Urban gardens as hotspots for

plant diversity in cities. *Urban Ecosystems*, *25*(3), 927–939. https://doi.org/10.1007/s11252-

021-01198-0

Shwartz, A., et al. (2012). Urban biodiversity, city-dwellers and conservation: How does an outdoor

activity day affect the human-nature relationship? *PLOS ONE*, *7*(6), e38642.

https://doi.org/10.1371/journal.pone.0038642

Smith, J., & Jehlička, P. (2013). Quiet sustainability: Fertile lessons from Europe's productive

gardeners. *Journal of Rural Studies*, *32*, 148–157.

https://doi.org/10.1016/j.jrurstud.2013.05.002

Soga, M., Gaston, K. J., & Yamaura, Y. (2017). Gardening is beneficial for health: A meta-analysis.

Preventive Medicine Reports, *5*, 92–99. https://doi.org/10.1016/j.pmedr.2016.11.007

Sturm, U., et al. (2021). Fascination and joy: Emotions predict urban gardeners' pro-pollinator

behaviour. *Insects*, *12*(9), Article 9. https://doi.org/10.3390/insects12090785

Tassin de Montaigu, C., & Goulson, D. (2023). Factors influencing butterfly and bumblebee richness

and abundance in gardens. *Science of The Total Environment*, 167995.

https://doi.org/10.1016/j.scitotenv.2023.167995

The pandas development team. (2022). *pandas-dev/pandas: Pandas* (Version v1.5.2) [Software].

Zenodo. https://doi.org/10.5281/zenodo.7344967

Tikka, P. M., Kuitunen, M. T., & Tynys, S. M. (2000). Effects of educational background on students'

attitudes, activity levels, and knowledge concerning the environment. *The Journal of*

Environmental Education, *31*(3), 12–19. https://doi.org/10.1080/00958960009598640

Tóth, A., et al. (2018). Changing patterns of allotment gardening in the Czech Republic and Slovakia.

Nature and Culture, *13*(1), 161–188. https://doi.org/10.3167/nc.2018.130108

Trendov, N. (2018). Comparative study on the motivations that drive urban community gardens in

Central Eastern Europe. *Annals of Agrarian Science*, *16*.

https://doi.org/10.1016/j.aasci.2017.10.003

Tresch, S., et al. (2019). Litter decomposition driven by soil fauna, plant diversity and soil

management in urban gardens. *Science of The Total Environment*, *658*, 1614–1629.

https://doi.org/10.1016/j.scitotenv.2018.12.235

Tresch, S., et al. (2018). A gardener's influence on urban soil quality. *Frontiers in Environmental*

Science, *6*. https://doi.org/10.3389/fenvs.2018.00025

Troumbis, A. Y. (2021). Imbalances in attitudes of European citizens towards biodiversity: Did the

communication of the European Biodiversity Strategy work? *Journal for Nature*

Conservation, *63*, 126041. https://doi.org/10.1016/j.jnc.2021.126041

United Nations. (2018). *World Urbanization Prospects: The 2018 Revision*. United Nations,

Department of Economic and Social Affairs, Population Division.

van der Veen, M. (2005). Gardens and fields: The intensity and scale of food production. *World*

Archaeology, *37*(2), 157–163. https://doi.org/10.1080/004382405000130731

van Elsen, T. (2000). Species diversity as a task for organic agriculture in Europe. *Agriculture,*

Ecosystems & Environment, *77*(1), 101–109. https://doi.org/10.1016/S0167-8809(99)00096-1

Varga-Szilay, Z., et al. (2024). Bridging biodiversity and gardening: Unravelling the interplay of socio-

demographic factors, garden practices, and garden characteristics. *Urban Forestry & Urban*

Greening, *97*(128367). https://doi.org/10.1016/j.ufug.2024.128367

Varga-Szilay, Z., & Pozsgai, G. (2022). Plant growers' environmental consciousness may not be

enough to mitigate pollinator declines: A questionnaire-based case study in Hungary. *Pest*

Management Science, *79*(4), 1284–1294. https://doi.org/10.1002/ps.7277

Vávra, J., Daněk, P., & Jehlička, P. (2018). What is the contribution of food self-provisioning towards

environmental sustainability? A case study of active gardeners. *Journal of Cleaner*

Production, *185*, 1015–1023. https://doi.org/10.1016/j.jclepro.2018.02.261

Vávra, J., Lapka, M., & Cudlínová, E. (2014). *Current challenges of Central Europe: Society and*

environment. Chapter: Challenges of urban agriculture: highlights on the Czech and Slovak

Republic specifics. Filozofická fakulta Univerzity Karlovy.

1093