1   **Scales of marine endemism in oceanic islands and the Provincial-Island endemism**

3   Hudson T. Pinheiro[1], Luiz A. Rocha[2], Juan P. Quimbayo[3]

5   [1] Center for Marine Biology, University of São Paulo, São Sebastião, SP, Brazil
6   [2] California Academy of Sciences, San Francisco, CA, USA
7   [3] Department of Biology, University of Miami, Miami, FL, USA

8
9

Abstract

Oceanic islands are remote environments commonly harboring endemic species, which often are
unique lineages originated and maintained by a variety of ecological, biogeographical and evolutionary
processes. Endemic species are found mostly in a single oceanic island or archipelago, however, a great
number of species can be considered multiple-island endemics, i.e. species found on multiple oceanic
islands that still have a restricted distribution. The geographic criteria chosen to classify endemic species
has a direct impact on the endemism rate of islands, and many studies have used multiple scales of
endemism (single and multiple-island endemics), which has historically influenced wide-scale
comparisons. In this perspective, we accessed the importance of single and multiple-island endemic
species to the biodiversity of oceanic islands, introducing the concept of Provincial-island endemism as an
additional strategy to standardize biogeography studies.

Key-words: marine biodiversity; endemic species; remote islands; speciation; biogeography;
conservation

Main Text

It is well known that many oceanic islands harbor unique biodiversity, and due to their faunal
simplicity and limited distributions, these endemic species have often been the focus of important
evolutionary, ecological and biogeographic studies. The diversification patterns and premises in island
biogeography, however, differ for terrestrial and marine realms. Terrestrial species often diversify
following adaptive radiation, while marine species, which are better dispersers, experience fewer
opportunities for in situ cladogenesis (Pinheiro et al., 2017). Therefore, marine endemism is more often
related to geographical isolation (Ávila et al., 2019, 2018; Hachich et al., 2019, 2015). Accordingly, the
origin of new lineages based on the colonization of islands by immigrants and their subsequent isolation
represents one of the most classical scenarios of evolution, a process known as peripatric speciation. This
process has been suggested to drive the highest levels of endemism in islands of both Pacific (e.g., Hawaii
and Eastern Island) and Atlantic (e.g., Ascension, St Helena and St Paul's) oceans (Delrieu-Trottin et al.,
2019; Floeter et al., 2008; Hourigan and Reese, 1987; Randall, 1998). The colonization process is
suggested to happen stochastically, but also in waves, what is often related to favorable oceanographic

conditions for founder populations to reach the islands or source populations to expand (Delrieu-Trottin
et al., 2017; Hodge et al., 2014; Pinheiro et al., 2017).
42
Recent studies have explored a variety of processes that contribute to the current endemism level of
oceanic islands. Ecological or parapatric speciation has been evoked to explain the high endemism rate in
the Marquesas (Gaither et al., 2015; Randall, 1998), and the evolution of species in the Northeastern
Brazilian oceanic islands (Rocha et al., 2005). Pleistocene sea-level variations have also promoted
vicariant events, driving allopatric speciation (Pinheiro et al., 2017) – the connections between source
and insular populations change as shallow habitats are created or suppressed. In the Vitoria-Trindade
Chain, low sea level stands exposed many seamounts that function as stepping-stones for weak disperser
species to reach remote oceanic islands (Simon et al., 2022). Subsequent sea-level rise isolated their
populations eliminating connecting shallow habitat, closing migration and driving allopatric speciation.
Moreover, phylogenetic studies are also revealing many paleoendemic species in oceanic islands (relict
lineages that likely had wider distributions in the past; Delrieu-Trottin et al., 2019; Pinheiro et al., 2017).
54
In this context, islands function as dynamic biodiversity engines that not only work as cradles for
species origination, but also as museums that contribute to lineages persistence over time. The role of
islands to develop and support new lineages is commonly measured by their endemism rate, which is the
proportion of the local biodiversity that is exclusively found there. Recent empirical work and
biogeographic analyses have shown that many islands share species exclusively found in remote and
isolated oceanic islands (Floeter et al., 2008; Pinheiro et al., 2018; Robertson and Cramer, 2009). These
multiple-island endemic species have their origin or persistence associated with the insular environment,
and in many cases, they can compose a substantial portion of the local biodiversity (Pinheiro et al., 2020).
In this perspective we assess the importance of single and multiple-island endemic species to the
biodiversity of oceanic islands, discuss how the scale of endemism has historically biased large-scale
biogeographic comparisons, and suggest strategies to move forward. We are herein considering as
oceanic islands only remote formations predominantly formed by volcanic activity, and for simplification,
we will hereafter refer to oceanic islands, archipelagoes, atolls, and shallow seamounts, all as "oceanic
islands".
69
We analyzed the distribution of 7,289 fish species associated with reef environments of 87 oceanic
islands and 189 coastal reefs around the world (e.g., Kulbicki et al., 2013), and uncovered that 5,260
(72%) species occur in oceanic islands. From those, 4,346 species occur in both oceanic islands and
continental coastal reefs, and 890 species are exclusively found around oceanic islands. Therefore, 12.2%
of the world´s reef fish biodiversity seems to be endemic to oceanic islands. Regarding the endemic
fishes, 541 (60.7%) are single island endemics, while 349 (39.3%) are multiple island endemics (Figure 1
A). Moreover, some species are genetically connected in distances that are greater than their isolation to
the continent (Rocha et al., 2005; Bernardi et al., 2014). For instance, *Stegastes beebei* is an endemic
species found in Galápagos, Cocos, and Malpelo, in the Eastern Tropical Pacific, but not in the adjacent
coastline (Robertson and Cramer, 2009). Similarly, the oceanic islands in the Mid-Atlantic Ridge, over
1,000 km apart, also share many endemic species (Brown et al., 2019; Pinheiro et al., 2020; Wirtz et al.,
2017). In the Brazilian Province, there is an extreme case where a species (*Opistognathus thionyi*) is
shared between islands almost 2,000 km apart, where the shortest distance to the mainland is only 350
km (Pinheiro et al., 2018) – a connectivity pattern also observed for a coral lineage (Peluso et al., 2018)
and a Brazil-Caribbean link (Rocha et al. 2005). It has been suggested that an oceanic "pathway" driven
by stochastic and ecological factors connects species among oceanic islands at the same time that
constrains their establishment in the continental shelf (Pinheiro et al., 2020, 2018). Indeed, several

species might have the ability to disperse, but the environment is a key factor for their persistence
(Rocha et al., 2005; Mazzei et al., 2021), and can contribute for the establishment of historical refuges in
oceanic islands during unfavorable geological periods (Peluso et al., 2018). Thus, even the most remote
archipelagoes do not constitute dead ends, often exporting new lineages to islands and seamounts
nearby (Simon et al., 2022) and also back to biodiversity centers in a process known as Biodiversity
Feedback (Bowen et al., 2013).
93
From these multiple-island endemic species, 231 (66%) are found in only two or three islands or
archipelagoes, and a strong negative relationship is observed between the number of endemic species
and the number of islands they occur (Figure 1). For instance, very few multiple-island endemic fishes are
found in more than 10 islands, so, in general, they tend to have restricted distributions. This pattern can
be explained by the evolutionary history and life-history of endemic fishes, as well as the geography of
islands. Fishes in general have good dispersal potential due the combination of pelagic larval duration
and swimming abilities, and most species that inhabit oceanic islands or are widespread along the oceans
have large body size, pelagic spawning and larvae, and exhibit rafting abilities (Luiz et al., 2013, 2012;
Mazzei et al., 2021; Pinheiro et al., 2018). The evolution of an endemic lineage often occurs after the
termination of gene flow with source populations, a process commonly observed in vicariant events and
stochastic colonization of weak dispersers, or the establishment of weak and moderate dispersers in
extremely remote locations. This combination of life-history traits and island geography contribute to a
wide variety of morphologies and taxonomic groups among endemic reef fishes (Robertson, 2001), and
to the positive relationship between the number of endemic fishes and the island isolation (Hachich et
al., 2015). Therefore, the same drivers that allow speciation can also contribute to the maintenance of
their restricted distribution. Moreover, many lineages are considered neo-endemics, evolving during the
Pleistocene and possibly not having enough time to disperse or adapt to other oceanic islands and
seamounts. For instance, the evolutionary history of endemic fishes in Rapa Nui (Pacific Ocean) and
Trindade (Atlantic Ocean) islands shows a majority of neo-endemics with restricted geographic range,
while some wider-distributed endemics show older lineages (Delrieu-Trottin et al., 2019; Pinheiro et al.,
2017).
115
The distribution of endemic fishes contributes directly to the endemism rate of the islands. The
Hawaiian archipelago and Rapa Nui have traditionally been considered hotspots of endemism, showing
endemism rates above 20% (Randall, 1998). However, according to our analyses, these localities would
present 10 and 13.4% of endemism, respectively, if only single-island endemics are considered (Figure 2).
The endemism rates of these locations are substantially higher when multiple-island endemics are
considered, rising to 22.2 and 35.9% respectively. Some islands that present high proportions of multiple-
island endemics do not necessarily present high proportions of single-island endemics. Indeed, most of
the islands herein analyzed (66.6%) present more multiple-island than single-island endemics, and 25.4 %
of the islands do not have a single-island endemic fish, some presenting high proportions of multiple-
island endemics, such as Salas y Gomes (32.0%), Johnston Atoll (16.2%), Madeira (7.1%) and Gambier
(7.0%). Accordingly, considering endemic species that are shared among nearby islands, the highest
endemism rates are found in the remote islands of Juan Fernandez and Desventuradas, where 69.5 and
53.3% of their respective biodiversity are composed by island endemics restricted to their biogeographic
province. In the Atlantic, over 24% of the fishes found in each of the islands of St Helena and Ascension
are exclusively found in the Mid-Atlantic Ridge.
131
So, why do the scales of endemism in oceanic islands matter? Some of the reasons include their
importance for evolutionary, ecologic and biogeographic studies, as well as for conservation planning.

Because they contain simplified species communities, oceanic islands have for a long time been
considered natural laboratories to understand the origin of marine biodiversity, and omitting about 40%
of their endemicity represent disregarding a great amount of evolutionary history and diversification
processes that are promoted in the islands. Recent studies that have included multiple island endemics in
their analyses are finding a wide range of genetic connectivity, meaning that, while some are expanding
their distribution, others are under ongoing fragmentation (Bernardi et al., 2014; Delrieu-Trottin et al.,
2017). Moreover, others studies are revealing relict species, remains of apparently previously widely
distributed lineages that only survived in remote locations. Therefore, the inclusion of single and multiple
island endemics in evolutionary and biogeography studies is important to clarify our understanding of the
evolution of marine communities in remote islands.
144
All oceanic islands analyzed showed some degree of endemism (i.e., containing single or multiple
island endemic species), having many species that are exclusively adapted to oceanic environments. The
adaptation process that leads to speciation or distributional expansion is also associated with ecological
advantages. These species might be better competitors and have higher abundances than widespread
species (Hobbs et al., 2012; Macieira et al., 2015). The ecology of the endemic fishes can contribute
insights about their dispersive ability (Robertson, 2001) and be related to their evolutionary history
(Pinheiro et al., 2017). Thus, considering the endemism status of a species could have important
implications on ecological studies that use these remote environments as natural laboratories.
153
In biogeography, overlooking endemics that are shared between islands can downplay the strength
of barriers and/or the ecological factors that drive speciation and taxonomic uniqueness. Endemism
represents singularity, a proportion of the fauna that evolved or has been maintained in a certain
province, region or location. Notwithstanding, studies have traditionally been reporting endemism rates
for islands using multiple scales of endemism (single and multiple-island endemics), which has historically
biased wide-scale contrasting. To move forward in the comparisons among oceanic islands, we suggest
biogeographers to clarify two aspects of endemism: 1) single-island endemics; endemism rate of the local
biodiversity when removing species recorded in different oceanic islands; 2) provincial-island endemics;
local endemism rate considering as endemic those species restricted to oceanic islands of the same
biogeographic province (e.g., provinces in Kulbicki et al., 2013). Following the former criteria, the islands
with highest endemism rates are Juan Fernandez, St. Paul Rocks, St. Helena, Ascension, Eastern Island,
Marquesas, Trindade, Galapagos, Cabo Verde and Hawaii, all with over 10% of single-island endemism
(Figure 2A). According to the proposed novel criteria, the islands of Juan Fernandez, Deventuradas, St
Helena, Ascension, Rapa Nui, Salas y Gomes, Galapagos, St. Paul Rocks, Hawaii, and Marquesas present
the highest endemism rates, ranging from 18.5% to 69.5 , (Figure 2B). Applying these criteria leads to
better assessments of the drivers of speciation and species maintenance.
170
Although reef fish communities in oceanic islands have experimented severe local impacts by both
commercial and recreational fisheries (Guabiroba et al., 2020; Luiz and Edwards, 2011; Pinheiro et al.,
2010; Quimbayo et al., 2019; Tuya et al., 2006), very few endemic species are considered important
fishing resources.. Endemic lineages of angelfishes of unique color morphs from St. Paul's Archipelago
have reached the aquarium trade, but they are now protected. Nevertheless, the restricted distribution
makes endemic fishes vulnerable to population fluctuations and crashes, and global stressors driven by
climate change, ocean warming and acidification are ongoing issues. For instance, *Azurina eupalama*, an
endemic species from Galápagos Archipelago, hasn't been seen since the 1982/83 very strong El Nino
(Edgar et al., 2010). Many endemic species show signs of bottlenecks and decreasing population size
(Pinheiro et al., 2017), which could make them more vulnerable to human derived impacts. The IUCN

threat classification has historically used the distribution of species and their population sizes as a proxy
for vulnerability, and a special attention to single and multiple-island endemics is here supported.
183
Moreover, biodiversity hotspots have been determined based on the endemism rate and threats to
their biodiversity, and have often been used as a proxy for the establishment of priority areas for
conservation (Roberts et al., 2002). This view has evolved to the protection of both connectivity and
genetic diversity as well as peripheral areas of high endemism (Hughes et al., 2002). Many very-large
marine protected areas have been created around oceanic islands throughout the world, however, many
of them do not directly protect coastal reefs and endemic species, contributing to the mismatch between
protection and the conservation priorities (Giglio et al., 2018; Lindegren et al., 2018). Therefore,
understanding how much of the diversity of an island represents species of restricted distribution is the
first step to clarify its vulnerability and priority for conservation. Applying current scientific criteria in
conservation planning for the protection of priority areas is the next step to be followed.
194
195

197 Figure 1 - Distribution of endemic species among islands and provinces. A) Relationship between the
198 number of island endemic species and the number of islands they occur. B) Relationship between the
199 number of island endemic fishes and the number of provinces they occur in.
200

201
202 Figure 2 - Proportion of single-island (a) and province-island (b) endemism among the oceanic islands
203 with highest endemism.
204
205
206
207

Acknowledgements

209   HTP thanks Fundação de Amparo à Pesquisa do Estado de São Paulo for funding (2019/24215-2) and
210   fellowship (2021/07039-6), and the administrative and logistics support of CEBIMar-USP, especially
211   Monica Zinneck, Irineu D. dos Santos, J. Jeronimo Favaro, Eduardo Honuma e Joseilto M. de Oliveira.

Data, scripts, code, and supplementary information availability

213   Data is curated by the senior author of this study JPQ (quimbayo.j.p@gmail.com), who is currently
214   leading a data paper (in preparation). The data used in this article was not a requisite for the conception
215   and development of the ideas presented, but show the importance of them.
216

Conflict of interest disclosure

218   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
219   relation to the content of the article.
220

Funding

222   HTP thanks Fundação de Amparo à Pesquisa do Estado de São Paulo for funding (2019/24215-2) and
223   fellowship (2021/07039-6).

References

Ávila, S.P., Cordeiro, R., Madeira, P., Silva, L., Medeiros, A., Rebelo, A.C., Melo, C., Neto, A.I., Haroun, R.,
Monteiro, A., Rijsdijk, K., Johnson, M.E., 2018. Global change impacts on large-scale biogeographic
patterns of marine organisms on Atlantic oceanic islands. Mar. Pollut. Bull. 126, 101–112.
doi:10.1016/j.marpolbul.2017.10.087
Ávila, S.P., Melo, C., Berning, B., Sá, N., Quartau, R., Rijsdijk, K.F., Ramalho, R.S., Cordeiro, R., Sá, N.C., Pimentel,
A., Baptista, L., Medeiros, A., Gil, A., Johnson, M.E., 2019. Towards a 'Sea-Level Sensitive' dynamic
model: impact of island ontogeny and glacio-eustasy on global patterns of marine island biogeography.
Biol. Rev. doi:10.1111/brv.12492
Bernardi, G., Ramon, M.L., Alva-Campbell, Y., McCosker, J.E., Bucciarelli, G., Garske, L.E., Victor, B.C., Crane,
N.L., 2014. Darwin's fishes: phylogeography of Galápagos Islands reef fishes. Bull. Mar. Sci. 90, 533–549.
doi:10.5343/bms.2013.1036
Bowen, B.W., Rocha, L.A., Toonen, R.J., Karl, S.A., 2013. The origins of tropical marine biodiversity. Trends Ecol.
Evol. 28, 359–66. doi:10.1016/j.tree.2013.01.018
Brown, J., Beard, A., Clingham, E., Fricke, R., Henry, L., Wirtz, P., 2019. The fishes of St Helena Island, central
Atlantic Ocean—new records and an annotated check-list. Zootaxa 4543, 151–194.
Delrieu-Trottin, E., Mona, L.B.S., Neglia, V., Giles, E.C., Pablo, C.R., 2019. Understanding the origin of the most
isolated endemic reef fish fauna of the Indo-Pacific: Coral reef fishes of Rapa Nui. J. Biogeogr. in press.
doi:10.1111/jbi.13531
Delrieu-Trottin, E., Mona, S., Maynard, J., Neglia, V., Veuille, M., Planes, S., 2017. Population expansions
dominate demographic histories of endemic and widespread Pacific reef fishes. Sci. Rep. 7, 40519.
doi:10.1038/srep40519
Edgar, G.J., Banks, S.A., Brandt, M., Bustamante, R.H., Chiriboga, A., Earle, S.A., Garske, L.E., Glynn, P.W.,
Grove, J.S., Henderson, S., Hickman, C.P., Miller, K.A., Rivera, F., Wellington, G.M., 2010. El Niño, grazers
and fisheries interact to greatly elevate extinction risk for Galapagos marine species. Glob. Chang. Biol.
16, 2876–2890. doi:10.1111/j.1365-2486.2009.02117.x
Floeter, S.R., Rocha, L.A., Robertson, D.R., Joyeux, J.C., Smith-Vaniz, W.F., Wirtz, P., Edwards, A.J., Barreiros,

J.P., Ferreira, C.E.L., Gasparini, J.L., Brito, A., Falcón, J.M., Bowen, B.W., Bernardi, G., 2008. Atlantic reef
fish biogeography and evolution. J. Biogeogr. 35, 22–47. doi:10.1111/j.1365-2699.2007.01790.x

Gaither, M.R., Bernal, M.A., Coleman, R.R., Bowen, B.W., Jones, S.A., Simison, W.B., Rocha, L.A., 2015.
Genomic signatures of geographic isolation and natural selection in coral reef fishes. Mol. Ecol. 24, 1543–
1557. doi:10.1111/mec.13129

Giglio, V.J., Pinheiro, H.T., Bender, M.G., Bonaldo, R.M., Costa-lotufo, L. V, Ferreira, C.E.L., Floeter, S.R., Freire,
A., Gasparini, J.L., Joyeux, J., Paulo, J., Lindner, A., Longo, G.O., Lotufo, T.M.C., Loyola, R., Luiz, O.J.,
Macieira, R.M., Magris, R.A., Mello, T.J., Quimbayo, J.P., Rocha, L.A., Segal, B., Teixeira, J.B., Vila-nova,
D.A., Vilar, C.C., Zilberberg, C., Francini-Filho, R., 2018. Large and remote marine protected areas in the
South Atlantic Ocean are flawed and raise concerns: Comments on Soares and Lucas (2018). Mar. Policy
96, 13–17. doi:10.1016/j.marpol.2018.07.017

Guabiroba, H.C., Santos, M.E.A., Pinheiro, H.T., Simon, T., Pimentel, C.R., Vilar, C.C., Joyeux, J.-C., 2020. Trends
in recreational fisheries and reef fish community structure indicate decline in target species population
in an isolated tropical oceanic island. Ocean Coast. Manag. 191, 105194.
doi:10.1016/j.ocecoaman.2020.105194

Hachich, N.F., Bonsall, M.B., Arraut, E.M., Barneche, D.R., Lewinsohn, T.M., Floeter, S.R., 2015. Island
biogeography: patterns of marine shallow-water organisms in the Atlantic Ocean. J. Biogeogr. 42, 1871–
1882. doi:10.1111/jbi.12560

Hachich, N.F., Ferrari, D.S., Quimbayo, J.P., Pinheiro, H.T., Floeter, S.R., 2019. Island Biogeography of Marine
Shallow-Water Organisms, in: Encyclopedia of the World's Biomes. Elsevier Inc., pp. 1–15.
doi:https://doi.org/10.1016/B978-0-12-409548-9.11947-5

Hobbs, J.-P.A., Jones, G.P., Munday, P.L., Connolly, S.R., Srinivasan, M., 2012. Biogeography and the structure
of coral reef fish communities on isolated islands. J. Biogeogr. 39, 130–139. doi:10.1111/j.1365-
2699.2011.02576.x

Hodge, J.R., van Herwerden, L., Bellwood, D.R., 2014. Temporal evolution of coral reef fishes: global patterns
and disparity in isolated locations. J. Biogeogr. 41, 2115–2127. doi:10.1111/jbi.12356

Hourigan, T.F., Reese, E.S., 1987. Mid-ocean isolation and the evolution of Hawaiian reef fishes. Trends Ecol.
Evol. 2, 187–191. doi:10.1016/0169-5347(87)90018-8

Hughes, T.P., Bellwood, D.R., Connolly, S.R., 2002. Biodiversity hotspots , centres of endemicity , and the
conservation of coral reefs. Ecol. Lett. 5, 775–784.

Kulbicki, M., Parravicini, V., Bellwood, D.R., Arias-Gonzàlez, E., Chabanet, P., Floeter, S.R., Friedlander, A.,
McPherson, J., Myers, R.E., Vigliola, L., Mouillot, D., 2013. Global Biogeography of Reef Fishes: A
Hierarchical Quantitative Delineation of Regions. PLoS One 8, e81847.
doi:10.1371/journal.pone.0081847

Lindegren, M., Holt, B.G., MacKenzie, B.R., Rahbek, C., 2018. A global mismatch in the protection of marine
biodiversity and ecosystem services. Sci. Rep. 8, 4099. doi:10.1038/s41598-018-22419-1

Luiz, O.J., Allen, A.P., Robertson, D.R., Floeter, S.R., Kulbicki, M., Vigliola, L., Becheler, R., Madin, J.S., 2013.
Adult and larval traits as determinants of geographic range size among tropical reef fishes. Proc. Natl.
Acad. Sci. U. S. A. 110, 16498–502. doi:10.1073/pnas.1304074110

Luiz, O.J., Edwards, A.J., 2011. Extinction of a shark population in the Archipelago of Saint Paul's Rocks
(equatorial Atlantic) inferred from the historical record. Biol. Conserv. 144, 2873–2881.
doi:10.1016/j.biocon.2011.08.004

Luiz, O.J., Madin, J.S., Robertson, D.R., Rocha, L.A., Wirtz, P., Floeter, S.R., 2012. Ecological traits influencing
range expansion across large oceanic dispersal barriers: insights from tropical Atlantic reef fishes. Proc.
R. Soc. B Biol. Sci. 279, 1033–40. doi:10.1098/rspb.2011.1525

Macieira, R.M., Simon, T., Pimentel, C.R., Joyeux, J.-C., 2015. Isolation and speciation of tidepool fishes as a
consequence of Quaternary sea-level fluctuations. Environ. Biol. Fishes 98, 385–393.
doi:10.1007/s10641-014-0269-0

Mazzei, E.F., Pinheiro, H.T., Simon, T., Moura, R.L., Macieira, R.M., Pimentel, C.R., Teixeira, J.B., Floeter, S.R.,

Ferreira, C.E.L., Ghisolfi, R.D., Francini-Filho, R.B., Quimbayo, J.P., Rocha, L.A., Gasparini, J.L., Joyeux, J.C., 2021. Mechanisms of dispersal and establishment drive a stepping stone community assembly on seamounts and oceanic islands. Mar. Biol. 168, 109. doi:10.1007/s00227-021-03919-7

Peluso, L., Tascheri, V., Nunes, F.L.D., Castro, C.B., Pires, D.O., Zilberberg, C., 2018. Contemporary and historical oceanographic processes explain genetic connectivity in a Southwestern Atlantic coral. Sci. Rep. 8, 1–12. doi:10.1038/s41598-018-21010-y

Pinheiro, H.T., Bernardi, G., Simon, T., Joyeux, J.-C., Macieira, R.M., Gasparini, J.L., Rocha, C., Rocha, L.A., 2017. Island biogeography of marine organisms. Nature 549, 82–85. doi:10.1038/nature23680

Pinheiro, H.T., Macena, B.C.L., Francini-Filho, R.B., Ferreira, C.E.L., Albuquerque, F. V., Bezerra, N.P.A., Carvalho-Filho, A., Ferreira, R.C.P., Luiz, O.J., Mello, T.J., Mendonça, S.A., Nunes, D.M., Pimentel, C.R., Pires, A.M.A., Soares-Gomes, A., Viana, D.L., Hazin, F.H. V., Rocha, L.A., 2020. Fish biodiversity of Saint Peter and Saint Paulʼs Archipelago, Mid-Atlantic Ridge, Brazil: new records and a species database. J. Fish Biol. 97, 1143–1153. doi:10.1111/jfb.14484

Pinheiro, H.T., Martins, A.S., Gasparini, J.L., 2010. Impact of commercial fishing on Trindade Island and Martin Vaz Archipelago, Brazil: characteristics, conservation status of the species involved and prospects for preservation. Brazilian Arch. Biol. Technol. 53, 1417–1423.

Pinheiro, H.T., Rocha, L.A., Macieira, R.M., Carvalho-Filho, A., Anderson, A.B., Bender, M.G., DiDario, F., Ferreira, C.E.L., Francini-Filho, R.B., Gasparini, J.L., Joyeux, J.-C., Luiz, O.J., Mincarone, M., Moura, R.L., Nunes, J. de A.C.C., Quimbayo, J.P., Rosa, R.S., Sampaio, C.L.S.S., Sazima, I., Simon, T., Vila-Nova, D.A., Floeter, S.R., 2018. South-western Atlantic reef fishes: zoogeographic patterns and ecological drivers reveal a secondary biodiversity center in the Atlantic Ocean. Divers. Distrib. 24, 951–965. doi:10.1111/ddi.12729

Quimbayo, J.P., Dias, M.S., Kulbicki, M., Mendes, T.C., Lamb, R.W., Johnson, A.F., Aburto-Oropeza, O., Alvarado, J.J., Bocos, A.A., Ferreira, C.E.L., Garcia, E., Luiz, O.J., Mascareñas-Osorio, I., Pinheiro, H.T., Rodriguez-Zaragoza, F., Salas, E., Zapata, F.A., Floeter, S.R., 2019. Determinants of Reef Fish Assemblages in Tropical Oceanic Islands. Ecography (Cop.). 42, 77–87. doi:10.1111/ecog.03506

Randall, J.E., 1998. Zoogeography of Shore Fishes of the Indo-Pacific Region. Zool. Stud. 37, 227–268.

Roberts, C.M., Mcclean, C.J., Veron, J.E.N., Hawkins, J.P., Allen, G.R., Mcallister, D.E., Mittermeier, C.G., Schueler, F.W., Spalding, M., Wells, F., Vynne, C., Werner, T.B., 2002. Marine Biodiversity Hotspots and Conservation Priorities for Tropical Reefs. Science (80-.). 295, 1280–1284.

Robertson, D., Cramer, K., 2009. Shore fishes and biogeographic subdivisions of the Tropical Eastern Pacific. Mar. Ecol. Prog. Ser. 380, 1–17. doi:10.3354/meps07925

Robertson, D.R., 2001. Population maintenance among tropical reef fishes: inferences from small-island endemics. Proc. Natl. Acad. Sci. U. S. A. 98, 5667–70. doi:10.1073/pnas.091367798

Rocha, L.A., Robertson, D.R., Roman, J., Bowen, B.W., 2005. Ecological speciation in tropical reef fishes. Proc. R. Soc. B Biol. Sci. 272, 573–579. doi:10.1098/2004.3005

Simon, T., Pinheiro, H.T., Santos, S., Macieira, R.M., Ferreira, Y.S.S., Bernardi, G., Rocha, L.A., Floeter, S.R., Ferreira, C.E.L., Joyeux, J.-C., 2022. Comparative phylogeography of reef fishes indicates seamounts as stepping stones for dispersal and diversification. Coral Reefs 41, 551–561. doi:10.1007/s00338-021-02178-8

Tuya, F., Ortega-Borges, L., Sanchez-Jerez, P., Haroun, R.J., 2006. Effect of fishing pressure on the spatio-temporal variability of the parrotfish, Sparisoma cretense (Pisces: Scaridae), across the Canarian Archipelago (eastern Atlantic). Fish. Res. 77, 24–33. doi:10.1016/j.fishres.2005.07.017

Wirtz, P., Bingeman, Jane, Bingeman, John, Fricke, R., Hook, T.J., Young, J., 2017. The fishes of Ascension Island, central Atlantic Ocean-new records and an annotated checklist. J. Mar. Biol. Assoc. United Kingdom 97, 783–798. doi:10.1017/S0025315414001301