Measles virus co-opts epithelial-to-mesenchymal transition and live cell extrusion to exit

human airway epithelia

Camilla E. Hippee[1], Lorellin A. Durnell[1], Justin W. Kaufman[2], Eileen Murray[2], Brajesh K. Singh[2],

and Patrick L. Sinn[1,2†]

[1]Microbiology and Immunology, Carver College of Medicine, The University of Iowa, Iowa City,

IA, 52242, USA

[2]Stead Family Department of Pediatrics, Carver College of Medicine, The University of Iowa,

Iowa City, IA, 52242, USA

[†]Corresponding author: Patrick L. Sinn, Stead Family Department of Pediatrics, 169 Newton Rd,

6318 PBDB, The University of Iowa, Iowa City, IA, 52242, USA, Tel: (319) 335-8190, Fax: 319-

335-9412, Email: patrick-sinn@uiowa.edu

Short title: Measles virus exits airway epithelia via cell detachment pathways

Key words: wound healing; cell detachment; viral infection; primary human cells; sphingosine

kinase; TGF-β

Formatting for JVI

ABSTRACT

Measles virus (MeV) is a highly contagious respiratory virus transmitted via aerosols. To

understand how MeV exits the airways of an infected host, we use unpassaged primary cultures

of human airway epithelial cells (HAE). MeV typically remains cell-associated in HAE and forms

foci of infection, termed infectious centers, by directly spreading cell-to-cell. We previously

described the phenomenon in which infectious centers detach *en masse* from HAE and remain

viable. Here we investigate the mechanism of this cellular detachment. Via immunostaining, we

observed loss of tight junction and cell adhesion proteins within infectious centers. These

morphological changes indicate activation of epithelial-to-mesenchymal transition (EMT). EMT

can contribute to wound healing in respiratory epithelia by mobilizing nearby cells. Inhibiting

TGF-β, and thus EMT, reduced infectious center detachment. Compared to uninfected cells,

MeV-infected cells also expressed increased levels of sphingosine kinase 1 (SK1), a regulator

of live cell extrusion. Live cell extrusion encourages cells to detach from respiratory epithelia by

contracting the actomyosin of neighboring cells. Inhibition or induction of live cell extrusion

impacted infectious center detachment rates. Thus, these two related pathways contributed to

infectious center detachment in HAE. Detached infectious centers contained high titers of virus

that may be protected from the environment, allowing the virus to live on surfaces longer and

infect more hosts. This mechanism may contribute to the highly contagious nature of MeV.

IMPORTANCE

Measles virus (MeV) is an extremely contagious respiratory pathogen that continues to cause

large, disruptive outbreaks each year. Here, we examine a phenomenon that may help explain

the contagious nature of MeV: detachment of MeV-infected cells. MeV spreads cell-to-cell in

human airway epithelial cells (HAE) to form groups of infected cells, termed "infectious centers".

We reported that infectious centers ultimately detach from HAE as a unit, carrying high titers of

virus. Viral particles within cells may be more protected from environmental conditions, such as

ultraviolet radiation and desiccation. We identified two host pathways, epithelial-to-

mesenchymal transition and live cell extrusion, that contribute to infectious center detachment.

Perturbing these pathways altered the kinetics of infectious centers detachment. These

pathways influence one another and contribute to epithelial wound healing, suggesting

infectious center detachment may be a usurped consequence of the host's response to infection

that benefits MeV by increasing its transmissibility between hosts.

INTRODUCTION

Measles virus (**MeV**) surpasses other respiratory viruses in transmissibility, but little evidence
exists to explain the contagious advantage of MeV. MeV is a morbillivirus in the paramyxovirus
family and, like other respiratory viruses, is primarily transmitted via aerosols. In an
unvaccinated population, an individual with RSV may spread virus to 3 others, while an
individual with MeV may spread virus to 12-18 others (1, 2). Vaccination rates must exceed 92%
for herd immunity; however, global vaccination rates have declined to 83% in the past two
years, leading to increased incidence of measles outbreaks and elevating MeV's status to an
imminent threat (3, 4, 5).
65
To model MeV respiratory infections, we use primary human airway epithelial cells (HAE). HAE
are derived from human lung donors and are directly seeded onto transwell filters without
passage (6). These cultures grow at an air-liquid interface, polarize, and differentiate into a
pseudostratified cell layer. HAE recapitulate the airways *in vivo* (7, 8). MeV infects HAE via its
epithelial cell receptor, nectin-4 (9, 10). Immune cells, such as dendritic cells or lymphocytes,
deliver the virus to nectin-4 on the basolateral side of HAE (11, 12). MeV then replicates in HAE
and spreads directly cell-to-cell via intercellular pores (13). MeV typically remains cell-
associated in HAE and forms foci of infection, termed infectious centers (shown schematically,
Fig 1A) (6, 13, 14). Infectious centers are unique to morbilliviruses; viruses outside of this genus
do not form infectious centers on HAE (13). Infectious centers are unique from syncytia and
multinucleated giant cells as the nuclei remain separate and the plasma membranes between
infected cells remain largely intact (6, 13). MeV infectious centers have been observed in HAE,
in human tracheal explants (6), and in the nasal passage, trachea, and bronchi of
experimentally infected macaques (15, 16).
80
Most respiratory viruses leave airway epithelial cells via apical budding, exocytosis, or cell lysis.
Many enveloped RNA respiratory viruses (e.g., IAV (17), RSV (18), and SARS-CoV-2 (19))
utilize apical budding at the host cell's plasma membrane to exit airway epithelial cells. Non-
enveloped viruses, like rhinovirus, exit via vesicles (exocytosis) or by bursting the cell open (cell
lysis) (20). MeV, an enveloped virus, was assumed to exit HAE via apical budding; however, we
found that MeV infection uniquely leads to dislodging of infectious centers containing
concentrated virus (**Fig 1A**) (21). Evidence for MeV-induced cell detachment is also found in
monkey epithelial cultures, *in vivo* monkey experiments, and human observation studies (15, 22,
23). The cellular signals that promote IC detachment are not known; however, cells of detached

infectious centers are metabolically active and do not display markers of cell death pathways,

suggesting live cell extrusion pathways are involved (21, 23).

Here, we investigate the mechanism of infectious center detachment. As will be shown, single-

cell RNA sequencing (21) suggests epithelial-to-mesenchymal transition (EMT) is activated

within infectious centers. Airway epithelial cells are typically quiescent and nonmigratory.

However, certain circumstances necessitate epithelial cells to divide and migrate. EMT is

initiated by diverse stimuli to dissolve cell-cell connections and promote mobilization. EMT is

involved in wound repair, tumor metastasis, and response to viral infection (24, 25, 26).

Importantly, EMT encompasses a spectrum of intermediate cell types.

EMT signaling associates with live cell extrusion (LCE), a pathway driven by sphingosine kinase

1 (SK1). SK1 phosphorylates sphingolipids, which are cell signaling molecules that localize to

membranes. In infected cells, increased SK1 activity drives pro-survival signaling that results in

more virus production (27, 28, 29, 30, 31, 32). In addition, downstream signaling of SK1 directs

neighboring cells to contract their actomyosin, creating a luminal force on the cell to be extruded

(33, 34). Cell extrusion is important for maintaining homeostasis within epithelia and can be

activated by several stimuli, including EMT, apoptosis, overcrowding, or loss of cell polarity (35,

36, 37). LCE and EMT signaling pathways overlap and each can activate the other. Here, we

show that both pathways are active within infectious centers and contribute to detachment.

RESULTS

Single cell RNA-sequencing implicates two detachment pathways.

To investigate what pathways may be responsible for infectious center detachment, we probed

our previously published single-cell RNA sequencing (scRNA-seq) dataset (21). In this dataset,

HAE from 10 human donors were mock-infected or infected with a MeV expressing a GFP

reporter protein (MeV-GFP). At multiple timepoints post-infection, samples were sorted via

fluorescence-activated cell sorting (FACS) to enrich the GFP+ population before performing

scRNA-seq. Lists of genes differentially expressed between mock and MeV infected (both GFP+

and GFP-) HAE were analyzed by gene ontology (**Fig 1B**). Many gene ontology terms from this

analysis pointed to EMT processes, including "positive regulation of EMT". Enrichment of gene

products such as TGFB1, SNAI2, and metalloproteinases (MMPs) are emblematic of EMT (**Fig

1C**).

EMT-related mRNAs were notably increased in a distinct cluster of cells, termed "ISG-high" (**Fig**

1D and 1E). This cluster was exclusively observed in MeV-infected HAE cultures (**S1A Fig**),

contained ~80% GFP+ and ~20% GFP- cells, and the cells expressed unique transcriptomes

that did not align with known HAE cell types (**Fig 1D**). Previously, we erroneously dubbed these

cells as "interferon-high cells" due to elevated interferon-stimulated gene (ISG) transcripts (21).

However, further analysis revealed that expression of these ISGs was not interferon-mediated

(38). As such, we renamed these cells "ISG-high cells" and continue to investigate their

characteristics. We could not identify what cell type the ISG-high cells were before they were

infected possibly because of the transcriptional impact of EMT.

Notably, ISG-high cells had elevated SNAI2 transcripts, a transcriptional regulator of EMT (**Fig**

1E and S1B Fig). Further investigation of the transcriptional profiles of ISG-high cells revealed

a potential second detachment pathway: live cell extrusion (LCE). LCE is induced by

sphingosine kinase 1 (SK1), one of the top enriched gene products within the ISG-high cluster

(**Fig 1F and S1C Fig**). In this manuscript, we will separately consider EMT and LCE for

infectious center detachment.

EMT is activated within infectious centers.

We evaluated the role of EMT in MeV infection. Quantitative Real-Time PCR (qRT-PCR)

confirmed EMT-related gene transcripts are elevated in MeV-infected cultures (**S2A Fig**). In

addition, TGF-β, a potent inducer of EMT, was secreted in higher concentrations following MeV-

infection (**S2B Fig**). EMT is characterized by a loss of epithelial cell markers (39, 40, 41, 42).

Indeed, immunostaining of infectious centers at 3 days post-infection and 7 days post-infection

with MeV-mCherry showed a loss of β-catenin (**Fig 2A**), ZO-1 (**Fig 2B**), and E-Cadherin (**Fig**

2C). In all cases, the loss of the markers was more pronounced toward the center of the

infectious center. Degradation increased over time and ~60% of the infectious center area had

loss of the indicated junctional protein by day 7 (**Fig 2D**). Additionally, cells of infectious centers

may have altered apical-basal polarity, as evidenced by an atypical pattern of expression of the

apically targeted ezrin protein (arrow; **S2C Fig**). These results demonstrate EMT activation in

MeV-infected HAE.

EMT contributes to infectious center detachment.

Infectious center detachment can be tracked using live cell fluorescence microscopy. Images

were collected daily post-infection and the timepoint at which each infectious center detached

was documented (**Fig 3A**). These data were used to calculate a probability of detachment. To

determine the impact of EMT on infectious center detachment, we treated MeV-infected HAE

with a TGF-β inhibitor, SB431542, which blocks downstream transcriptional activation of TGF-β

(43, 44). With TGF-β inhibited, infectious centers detached at a slower rate than vehicle-treated

controls (**Fig 3B**). Titers (**Fig 3C**), infectious center number (**Fig 3D**), infectious center area (**Fig**

3E), and cytotoxicity (**Fig 3F**) were not statistically different between the treatment and control

groups, suggesting that TGF-β inhibition slows IC detachment without impacting viral replication

or cell-to-cell spread.

LCE is activated within infectious centers.

EMT can lead to LCE. Once a cell is fated for extrusion, sphingosine, a cellular lipid, is

phosphorylated by sphingosine kinase 1 (SK1) to produce sphingosine-1-phosphate (S1P) (**Fig**

4A). S1P binds its receptor, S1PR2, on neighboring cells, signaling for the activation of the Rho

pathway. As a result, a ring of cells surrounding the fated cell will contract their actomyosin,

creating an apical force to expel the cell (33, 34, 35, 37). Transcripts for *SK1*, Rho pathway

genes (*RHOA*, *RHOB*, and *RHOD*), and genes responsible for actomyosin contraction (*MYH9*

and *ROCK1*) (37, 45) were elevated within the ISG-high cell cluster of our scRNA-seq dataset

(**Fig 4B**). Immunostaining revealed altered SK1 localization within the cells of infectious centers

(**Fig 4C and 4D**). Specifically, SK1 was most abundant at the nucleus (arrows; **Fig 4D**).

Together, these data suggest SK1 is expressed in MeV-infected cells and initiates LCE

signaling.

LCE contributes to infectious center detachment.

Using the infectious center detachment assay described in Figure 3, we evaluated the impact of

an SK1 inhibitor on infectious center detachment. Treatment with the SK1 inhibitor, SKI-II,

delayed infectious center detachment (**Fig 5A**). SKI-II treatment did not cause a statistically

significant change in titers (**Fig 5B**), infectious center number (**Fig S3A**), or infectious center

area (**Fig S3B**) of MeV-infected HAE. To exacerbate LCE, MeV-infected HAE were treated with

exogenous S1P. The probability of infectious center detachment increased and infectious

centers detached faster (**Fig 5C**), despite viral titers (**Fig 5D**), infectious center number (**Fig**

S3C), infectious center area (**Fig S3D**), and cytotoxicity (**Fig S3E**) remaining statistically

indistinguishable to vehicle-treated controls. These results demonstrate infectious center

detachment utilizes LCE pathways.

DISCUSSION

Here, we showed that two overlapping, wound healing pathways are active within the cells of

infectious centers. Transcriptional regulators of EMT and LCE are elevated in MeV-infected

cells. Physical evidence of EMT and LCE was observed via immunostaining and microscopy.

Additionally, because infectious center detachment could be delayed by inhibiting EMT and LCE

and increased using an LCE agonist, we concluded that both pathways contribute to infectious

center detachment. Although not shown, we induced EMT in HAE with exogenous TGF-β

followed by MeV infection to determine if infectious center detachment increased. However, the

results were uninterpretable because infection of epithelial cells primed for EMT results in

widespread syncytial-like, multi-nucleated giant cells instead of discreet infectious centers. As

we previously reported, infectious center formation requires proper epithelial differentiation (6).

As primary epithelial cell culture has become more accessible, cell detachment has been

observed in several virus infections of epithelia. This includes astrovirus (46), rotavirus (47), and

enterovirus A71 (48). Both EMT and LCE are candidate mechanisms for cell detachment in

these infections and are potentially a common result of epithelial cell viral infection.

EMT and LCE pathways have important roles in maintaining homeostasis of the epithelial layer

and their functions often overlap (35, 45, 49, 50). LCE activates in cases of cell overcrowding

(34, 36, 51), removing damaged cells following injury (51), or following apoptotic signaling (37,

52, 53, 54). EMT is a key tool for wound healing and is active in chronic lung diseases like

idiopathic pulmonary fibrosis (55). In EMT, cells can gain actomyosin contractility, a key

mechanism for LCE (41). Transcriptional regulators of EMT, like SNAI1, impact RhoA signaling

and actomyosin contractility (45). Wee et al. (45) showed that cell-cell adhesion loss works

cooperatively with actomyosin contraction to cause cell extrusion. In a complementary manner,

SK1 promotes EMT by degrading cell-cell adhesions, like E-cadherin, and activating the focal

adhesion kinase pathway, which includes MMPs (56, 57). Inhibiting SK1 blocks signaling of two

EMT activators, NF-κB and TGF-β, whereas exogenous S1P treatment induces EMT (58, 59).

Notably, SK1 and SNAI2 coexpress within human tumors (60). In fact, LCE and EMT appear to

be so intertwined that one does not exist without the other. We postulate that, in the context of

MeV infection of HAE, EMT and LCE activate simultaneously and cooperate to expel infectious

centers. EMT likely promotes infectious center detachment by altering cell-cell adhesions, while

LCE mechanically promotes detachment via actomyosin contraction.

SK1 is involved in several viral infections because of its role in activating cell pro-survival

signaling (29, 30, 61, 62, 63, 64). A stressed cell will accumulate sphingosine in its plasma

membrane, marking the cell for apoptosis; however, in the presence of increased SK1

expression, sphingosine is phosphorylated by SK1 to create S1P, a signal for cell survival (37,

65, 66, 67). S1P prolongs the life of an infected cell and increases viral replication as the virus

has more time to propagate. Viral replication is associated with SK1 expression during infection

with respiratory syncytial virus (32), human cytomegalovirus (68), and influenza virus (28). In

cases of viral infections, SK1 expression is induced via TNF-α signaling and regulated by the

ERK/AKT pathway (69, 70, 71). MeV may also take advantage of SK1's cell survival phenotype

to increase virion production. Indeed, several genes of the TNF-α pathway were upregulated in

our scRNA-seq ISG-high cells (**Fig 4B**).

We observed SK1 expression in the nucleus of infected cells (**Fig 4C and 4D**). While SK1 is

more commonly found in the cytoplasm, nuclear localization is seen in human lungs with

idiopathic pulmonary fibrosis (72) and SK1 translocation to the nucleus is induced with estrogen

treatment of breast cancer cells (73). While the role of nuclear SK1 is still unclear, sphingosine

kinase 2 is known to alter gene expression in the nucleus and may render cells more resistant

to apoptosis (74). In addition, MeV replication was previously shown to be sensitive to SK1

inhibition in HEK 293 cells (30). In that study, NF-κB expression was necessary for MeV

replication and inhibiting SK1 equally inhibited NF-κB. In contrast, we did not find that SKI-II

treatment affected viral titers in HAE. This may be due to differences in cell types used since

HAE are much more resilient than HEK 293 cells to viral infections and are more likely to

survive. Importantly, immortalized tumor cells are likely to be inherently skewed toward a

mesenchymal phenotype even if they were originally derived from an epithelial cell. Thus,

unpassaged, primary epithelial cells may be better suited for studying viral induced EMT and

LCE.

EMT might also explain why infectious centers stop growing. Our recent work determined that

cell-cell spread of MeV is not restricted by interferon-induced innate immune signaling (38).

EMT causes the disassociation of cell-cell junctions, including adherens junctions (24). The MeV

receptor nectin-4 is localized in adherens junctions and is required for cell-cell spread (13). The

loss of adhesion caused by EMT could contribute to halted infectious center growth. This

sequestration of infectious centers from the rest of the epithelia may also explain why the cells

of infectious centers maintain their connections to one another while simultaneously losing their

connection to neighboring cells. While we do not have an answer for why cells of infectious

centers remain attached to one another after they are dislodged, we have two hypotheses: 1)

Cells with similar actin densities group together (75): as MeV infection degrades actin within

infectious centers, the surface tension of the cellular membranes of infectious center cells may

decrease, distinguishing these cells from the uninfected cells around them. 2) While

mesenchymal-like cells do not integrate well within tissues during migration, they can form

transient cell-cell adhesions with each other and nearby stationary cells (76). One such

connection could be formed during the transfer of MeV factors through intercellular pores that

form during cell-cell spread. Future studies identifying cell-cell adhesions that persist in MeV-

infected cells will be key in determining the bottlenecks that impact infectious center growth and

detachment.

We identified a group of "ISG-high cells" in our scRNA-seq that had high transcripts of ISGs as

well as EMT and LCE markers. We postulate that these cells are either within infectious centers

or are uninfected cells near the periphery of infectious centers (**Fig S1A**). These cells are

responding to the infection, seemingly by increasing antiviral gene transcripts. In a recent

publication, we studied interferon stimulation in HAE in the context of MeV infection and

determined interferons are not produced at detectable levels (38). While the mystery of what

initiates ISG transcription remains, we believe the name "ISG-high cells" more accurately

describes this cluster than the previously reported "interferon high cells"(21). Importantly, we are

not suggesting that ISG-high cells are a new epithelial cell type, but cells with an altered

transcriptional profile following MeV infection. ISG-high cells were likely secretory or ciliated

cells before MeV infection, but we cannot be sure because those cell type markers are not

detectable in our samples. Understanding the influences that push cells towards the ISG-high

cell transcriptome profile is of interest.

In summary, our results describe two, related pathways by which MeV extrudes groups of virus-

laden cells. Infectious center detachment was altered by inhibitors and agonists of EMT and

LCE, which supports our conclusion that MeV exits HAE via this strategy. Damage to the cell

during MeV infection likely initiates these pathways and infectious center detachment is an

unintended consequence that gives MeV an advantage. While vaccination remains the best

preventative measure for MeV, understanding MeV infectious center detachment is vital for

investigating the contagious nature of MeV and may prove helpful in healthcare settings.

MATERIALS AND METHODS

Ethical statement. The University of Iowa *In Vitro* Models and Cell Culture Core prepared the

well-differentiated primary cultures of human airway epithelia (HAE) used in these studies.

These cultures are comprised of cells from autopsy, discarded tissue, or surgical specimens. All

human subject studies were conducted with approval from the University of Iowa Institutional

Review Board. We do not receive identifiable information.

Human airway epithelial cells. HAE were cultured and maintained as previously described (6,

7). Briefly, cells from trachea and bronchus epithelia are seeded onto collagen-coated,

polycarbonate transwell inserts (0.4 μm pore size, surface area = 0.33 cm^2; Corning Costar;

#3413) following enzymatic disassociation. Cells were submerged in Ultroser G (USG) medium

at 37°C and 5% CO_2. After 48 hours, apical USG was removed and cells were incubated at the

air-liquid interface and allowed to polarize and differentiate. All HAE cultures used for the

experiments in this study were >3 weeks old and maintained a transepithelial electrical

resistance of at least 500 $\Omega \cdot \mu$m^2.

Bioinformatic analyses. The scRNA-seq dataset was previously described (21) and can be

accessed in the GEO with accession number GSE168775. In brief, MeV-infected or mock-

infected HAE (pooled from 10 human donors) were dissociated using TrypLE and sorted by

fluorescence-activated cell sorting to separate GFP- and GFP+ cells for scRNA-seq. ~30,000

cells make up the dataset. Gene ontology analysis was performed using GOrilla (77, 78). Violin

plots, t-SNE plots, and heatmap data were generated in Loupe Browser version 6.4.1.

Heatmaps were visualized with the R package gplots version 3.1.3.

Measles virus production. Recombinant MeV-GFP and MeV-mCherry viruses are derived

from the IC323 strain and were produced as previously described (79). Briefly, Vero cells that

stably express human SLAMF1 (80) were maintained in Dulbecco modified Eagle medium

(DMEM; Thermo Fisher Scientific) supplemented with 5% newborn calf serum (NCS; Thermo

Fisher Scientific) and penicillin-streptomycin (100 mg/mL; Thermo Fisher Scientific). Vero-

hSLAM cells were infected with either MeV-GFP for MeV-mCherry at an MOI of 0.01 for 2 hours.

2 days post-infection, cells were collected and subjected to several freeze/thaw cycles to induce

cell membrane rupture and virus release. Virus was titrated using TCID$_{50}$ assays with Vero-

hSLAMF1 cells. We have not detected differences in MeV-GFP and MeV-mCherry infections.

We selected MeV-GFP or MeV-mCherry for our experiments due to their compatibility with

reagents and instrumentation.

Infection of HAE. HAE cultures in this study were infected with MeV as previously published

(13, 14). Briefly, HAE cultures were inverted to expose the basolateral surface. 50 μL virus

inoculum, containing MeV virus and serum-free opti-MEM (Thermo Fisher Scientific;

#31985062) was placed on the basolateral side of the cells. Virus inoculum was incubated on

HAE for ~4 hours at 37°C and 5% CO_2. After the infection period, the inoculum was removed

and the cultures were placed back into an upright position with medium in the basolateral

chamber.

Immunostaining and confocal microscopy. All cells prepped for immunofluorescence were

fixed with 2% paraformaldehyde for 15 minutes followed by washing with 1X PBS and

permeabilization with 0.2% Triton X-100 in Superblock (Thermo Fisher Scientific; #37580) for 1

hour. All primary antibodies were used at 1:100 dilution in Superblock overnight at 4°C. After

overnight incubation, cells were washed 3x with 1X PBS and incubated with fluorescently

labelled secondary antibodies at a 1:1000 dilution in Superblock buffer for 1 hour at RT, gentle

rocking. HAE filters were cut from the transwell using a razor blade and mounted onto glass

microscope slides, covered with VECTASHIELD Mounting Medium with DAPI (Vector

Laboratories; #H-1200-10) and a glass coverslip overtop. Coverslips were sealed with clear nail

polish. For confocal analysis, a Leica SP8 STED confocal microscope (Leica Microsystems)

was used. Image processing was performed using ImageJ2 version 2.3.0. Primary antibodies

used: Beta-catenin (Thermo Fisher Scientific; #13-8400), ZO-1 (Thermo Fisher Scientific; #33-

9100), E-cadherin (Cell Signaling Technology; #3195), Ezrin (Cell Signaling Technology; #3145),

SK1 (Invitrogen; #PA5-28584). Secondary antibodies used: Goat Anti-Rabbit IgG Highly Cross-

Adsorbed (H+L) Alexa Fluor 568 (Invitrogen; #A-11036), Goat Anti-Mouse IgG Highly Cross-

Adsorbed (H+L) Alexa Fluor 546 (Invitrogen; #A-11030).

Quantification of cytoskeleton breakdown. Confocal images of 3, randomly selected

infectious centers were collected from 3 human donors for BCAT, ZO-1, and ECAD. Infectious

center area was measured using the freehand tool in ImageJ2 version 2.3.0. Adhesion

disruption area was determined by accumulating the area measurements of all protein

degradation within the infectious center. The data are plotted as a percentage: (total area of

degradation) / (total area of infectious center) x 100.

Quantitative reverse transcriptase polymerase chain reaction (qRT-PCR). At 3 dpi, mock-

and MeV-infected cell lysates were collected with TRIzol reagent (Thermo Fisher Scientific;

#15596026). Cellular RNA was isolated using the Direct-zol RNA Miniprep kit (Zymo Research;

#R2052). After isolation, RNA was transcribed into cDNA with the High-Capacity RNA-to-cDNA

kit and the manufacturer's instructions (Thermo Fisher Scientific; #R2052). qPCR of *MMP1*,

MMP10, *TGFB1*, *SNAI2*, and *SMAD3* was accomplished using primers, *Power*SYBR Green

Real-Time PCR Master Mix (Thermo Fisher Scientific; #4367659), and a QuantStudio 7 Pro

(Applied Biosystems). The housekeeping gene used was *SFRS9*. Primer sequences can be

found in Table S1.

TGF-β ELISA. Apical washes (100 µl Opti-MEM) and basolateral media (100 µl USG medium)

were collected from mock- and MeV-infected HAE. Secretions were measured in duplicate for

TGF-β protein content using a Human TGF-beta 1 ELISA Kit (RayBiotech; #ELH-TGFb1-1) and

following the manufacturer's protocol. ELISAs were read on XXX.

Drug treatments. SKI-II (MedChemExpress; #312636-16-1) and SB431542 (Tocris; #1614)

were diluted to 20 µM with USG medium. Recombinant S1P (Tocris; #1370) was diluted to 10

µM with USG medium. Treatments were placed in the basolateral chamber of the HAE cultures.

Equivalent volume of DMSO (4 µl) was diluted in USG medium for vehicle control cultures. SKI-

II and S1P treatments began 1-day post-infection. SB431542 treatment began 1 day pre-

infection. Treatments remained on the cells for the length of the experiment and

media/treatments were refreshed every 2 days.

Live-image microscopy and infectious center analyses. Image acquisition was performed

using an All-in-one-Fluorescence Microscope BZ-X800E (Keyence Corporation of America) with

a 4X objective. Images from each culture were arranged sequentially by time post-infection. A

blinded individual tracked individual infectious centers over the time course of the experiment

and determined which day post-infection the infectious center detached. These data were

converted to binary and were graphed as a probability of detachment.

Titration of MeV. MeV-infected HAE lysates for titration analysis were scraped and collected in

serum-free medium and 10-fold serially diluted. Serial dilutions were plated in the 96-well flat-

bottom plate containing Vero-hSLAM cells. 3 days post-infection, cells were titrated using

$TCID_{50}$/mL analysis.

LDH cytotoxicity colormetric assay. HAE cultures were infected and supernatants from the

basolateral chamber were collected. Each cultures received fresh medium after collection. LDH

release was detected following the manufacturer's protocol (BioVision; #K311). Briefly,

supernatant and reaction mixture were mixed 1:1 and incubated in a 96-well Maxi-Sorp

Immunoplate (Thermo Fisher Scientific; #439454). After 30 minutes at room temperature,

absorbance was measured at 490_{OD}. Percent cytotoxicity was measured using the following

formula: (test sample – low control) ÷ (high control – low control). Mock-infected supernatants

served as the low controls. High control samples were obtained by incubating cells with media

containing 0.1% Triton X-100 for 30 minutes at 37°C, followed by media collection. This method

allows for complete cell death, and therefore maximum LDH release, to occur.

Sectioning HAE. HAE were fixed with 2% paraformaldehyde for 15 minutes at room

temperature. Filters were excised from the transwell, submerged in Scigen Tissue-Plus O.C.T.

Compound (Thermo Fisher Scientific; #23-730-571), and frozen. 10 μm sections were collected

with a Microm Cryostat HM505E (Leica) and immunostained following the protocols listed

above.

Statistics. Statistical tests listed in the figure legends were performed in GraphPad Prism

version 10.1.1. Alpha values were set at 0.05. Unless otherwise indicated, numerical data are

presented as mean ± standard deviation (SD).

ACKNOWLEDGEMENTS

We thank Roberto Cattaneo for providing fluorescent MeVs and guidance. Additionally, we thank

Jennifer Bartlett for her thoughtful review of this manuscript. We acknowledge the support of the

University of Iowa Central Microscopy Research Facility, the Genomics Division of the Iowa

Institute of Human Genetics, and the In Vitro Models and Cell Culture Core.

FUNDING

This work was supported by the National Institutes of Health: NIH R01 AL-132402 and T32 AL-

007533.

AUTHOR CONTRIBUTIONS

Conceptualization: Camilla E. Hippee, Brajesh K. Singh, Patrick L. Sinn.

Data curation: Camilla E. Hippee.

Formal Analysis: Camilla E. Hippee, Lorellin A. Durnell, Justin W. Kaufman.

Funding Acquisition: Camilla E. Hippee, Patrick L. Sinn.

Investigation: Camilla E. Hippee, Lorellin A. Durnell, Justin W. Kaufman, Eileen Murray.

Visualization: Camilla E. Hippee, Lorellin A. Durnell, Justin W. Kaufman.

Writing – original draft: Camilla E. Hippee.

Writing – review & editing: Camilla E. Hippee, Lorellin A. Durnell, Patrick L. Sinn.

REFERENCES

1.   Guerra FM, Bolotin S, Lim G, Heffernan J, Deeks SL, Li Y, et al. The basic reproduction
number (R(0)) of measles: a systematic review. Lancet Infect Dis. 2017;17(12):e420-e8.
2.   Reis J, Shaman J. Retrospective Parameter Estimation and Forecast of Respiratory
Syncytial Virus in the United States. PLoS Comput Biol. 2016;12(10):e1005133.
3.   Anderson RM, May RM. Directly transmitted infections diseases: control by vaccination.
Science. 1982;215(4536):1053-60.
4.   Anderson RM, May RM. Age-related changes in the rate of disease transmission:
implications for the design of vaccination programmes. J Hyg (Lond). 1985;94(3):365-436.
5.   Partnership MR. Measles Remains a Serious Threat: WHO and UNICEF National
Immunization Coverage Data Shows 2023 [Available from:
https://measlesrubellapartnership.org/measles-remains-a-serious-threat-who-and-unicef-
national-immunization-coverage-data-shows/.
6.   Kaufman JW, Singh BK, Durnell LA, Sinn PL. Representative measles virus infection
requires appropriate airway epithelia culture conditions. J Virol. 2023:e0105123.
7.   Karp PH, Moninger TO, Weber SP, Nesselhauf TS, Launspach JL, Zabner J, et al. An in
vitro model of differentiated human airway epithelia. Methods for establishing primary cultures.
Methods Mol Biol. 2002;188:115-37.
8.   Pezzulo AA, Starner TD, Scheetz TE, Traver GL, Tilley AE, Harvey BG, et al. The air-
liquid interface and use of primary cell cultures are important to recapitulate the transcriptional
profile of in vivo airway epithelia. Am J Physiol Lung Cell Mol Physiol. 2011;300(1):L25-31.
9.   Delpeut S, Noyce RS, Richardson CD. The tumor-associated marker, PVRL4 (nectin-4),
is the epithelial receptor for morbilliviruses. Viruses. 2014;6(6):2268-86.
10.   Muhlebach MD, Mateo M, Sinn PL, Prufer S, Uhlig KM, Leonard VH, et al. Adherens
junction protein nectin-4 is the epithelial receptor for measles virus. Nature.
2011;480(7378):530-3.
11.   Ferreira CS, Frenzke M, Leonard VH, Welstead GG, Richardson CD, Cattaneo R.
Measles virus infection of alveolar macrophages and dendritic cells precedes spread to
lymphatic organs in transgenic mice expressing human signaling lymphocytic activation
molecule (SLAM, CD150). J Virol. 2010;84(6):3033-42.
12.   Tatsuo H, Ono N, Tanaka K, Yanagi Y. SLAM (CDw150) is a cellular receptor for measles
virus. Nature. 2000;406(6798):893-7.
13.   Singh BK, Hornick AL, Krishnamurthy S, Locke AC, Mendoza CA, Mateo M, et al. The
Nectin-4/Afadin Protein Complex and Intercellular Membrane Pores Contribute to Rapid Spread
of Measles Virus in Primary Human Airway Epithelia. J Virol. 2015;89(14):7089-96.

14.   Sinn PL, Williams G, Vongpunsawad S, Cattaneo R, McCray PB, Jr. Measles virus
preferentially transduces the basolateral surface of well-differentiated human airway epithelia. J
Virol. 2002;76(5):2403-9.
15.   Ludlow M, de Vries RD, Lemon K, McQuaid S, Millar E, van Amerongen G, et al.
Infection of lymphoid tissues in the macaque upper respiratory tract contributes to the
emergence of transmissible measles virus. J Gen Virol. 2013;94(Pt 9):1933-44.
16.   Ludlow M, Lemon K, de Vries RD, McQuaid S, Millar EL, van Amerongen G, et al.
Measles virus infection of epithelial cells in the macaque upper respiratory tract is mediated by
subepithelial immune cells. J Virol. 2013;87(7):4033-42.
17.   Leser GP, Lamb RA. Influenza virus assembly and budding in raft-derived
microdomains: a quantitative analysis of the surface distribution of HA, NA and M2 proteins.
Virology. 2005;342(2):215-27.
18.   El Najjar F, Schmitt AP, Dutch RE. Paramyxovirus glycoprotein incorporation, assembly
and budding: a three way dance for infectious particle production. Viruses. 2014;6(8):3019-54.
19.   Pinto AL, Rai RK, Brown JC, Griffin P, Edgar JR, Shah A, et al. Ultrastructural insight into
SARS-CoV-2 entry and budding in human airway epithelium. Nat Commun. 2022;13(1):1609.
20.   Kerr SL, Mathew C, Ghildyal R. Rhinovirus and Cell Death. Viruses. 2021;13(4).
21.   Hippee CE, Singh BK, Thurman AL, Cooney AL, Pezzulo AA, Cattaneo R, et al. Measles
virus exits human airway epithelia within dislodged metabolically active infectious centers. PLoS
Pathog. 2021;17(8):e1009458.
22.   Bischoff WE, McNall RJ, Blevins MW, Turner J, Lopareva EN, Rota PA, et al. Detection
of Measles Virus RNA in Air and Surface Specimens in a Hospital Setting. J Infect Dis.
2016;213(4):600-3.
23.   Lin WW, Tsay AJ, Lalime EN, Pekosz A, Griffin DE. Primary differentiated respiratory
epithelial cells respond to apical measles virus infection by shedding multinucleated giant cells.
Proc Natl Acad Sci U S A. 2021;118(11).
24.   Le Bras GF, Taubenslag KJ, Andl CD. The regulation of cell-cell adhesion during
epithelial-mesenchymal transition, motility and tumor progression. Cell Adh Migr. 2012;6(4):365-
73.
25.   Shirogane Y, Takeda M, Tahara M, Ikegame S, Nakamura T, Yanagi Y. Epithelial-
mesenchymal transition abolishes the susceptibility of polarized epithelial cell lines to measles
virus. J Biol Chem. 2010;285(27):20882-90.
26.   Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. J Clin Invest.
2009;119(6):1420-8.
27.   Olivera A, Kohama T, Edsall L, Nava V, Cuvillier O, Poulton S, et al. Sphingosine kinase
expression increases intracellular sphingosine-1-phosphate and promotes cell growth and
survival. J Cell Biol. 1999;147(3):545-58.
28.   Seo YJ, Blake C, Alexander S, Hahm B. Sphingosine 1-phosphate-metabolizing
enzymes control influenza virus propagation and viral cytopathogenicity. J Virol.
2010;84(16):8124-31.
29.   Seo YJ, Pritzl CJ, Vijayan M, Bomb K, McClain ME, Alexander S, et al. Sphingosine
kinase 1 serves as a pro-viral factor by regulating viral RNA synthesis and nuclear export of viral
ribonucleoprotein complex upon influenza virus infection. PLoS One. 2013;8(8):e75005.
30.   Vijayan M, Seo YJ, Pritzl CJ, Squires SA, Alexander S, Hahm B. Sphingosine kinase 1
regulates measles virus replication. Virology. 2014;450-451:55-63.
31.   Carr JM, Mahalingam S, Bonder CS, Pitson SM. Sphingosine kinase 1 in viral infections.
Rev Med Virol. 2013;23(2):73-84.
32.   Monick MM, Cameron K, Powers LS, Butler NS, McCoy D, Mallampalli RK, et al.
Sphingosine kinase mediates activation of extracellular signal-related kinase and Akt by
respiratory syncytial virus. Am J Respir Cell Mol Biol. 2004;30(6):844-52.

33.   Gu Y, Forostyan T, Sabbadini R, Rosenblatt J. Epithelial cell extrusion requires the
sphingosine-1-phosphate receptor 2 pathway. J Cell Biol. 2011;193(4):667-76.
34.   Gudipaty SA, Rosenblatt J. Epithelial cell extrusion: Pathways and pathologies. Semin
Cell Dev Biol. 2017;67:132-40.
35.   Ohsawa S, Vaughen J, Igaki T. Cell Extrusion: A Stress-Responsive Force for Good or
Evil in Epithelial Homeostasis. Dev Cell. 2018;44(3):284-96.
36.   Eisenhoffer GT, Loftus PD, Yoshigi M, Otsuna H, Chien CB, Morcos PA, et al. Crowding
induces live cell extrusion to maintain homeostatic cell numbers in epithelia. Nature.
2012;484(7395):546-9.
37.   Rosenblatt J, Raff MC, Cramer LP. An epithelial cell destined for apoptosis signals its
neighbors to extrude it by an actin- and myosin-dependent mechanism. Curr Biol.
2001;11(23):1847-57.
38.   Durnell LA, Hippee CE, Cattaneo R, Bartlett JA, Singh BK, Sinn PL. Interferon-
independent processes constrain measles virus cell-to-cell spread in primary human airway
epithelial cells. Microbiol Spectr. 2023:e0136123.
39.   Elzamly S, Badri N, Padilla O, Dwivedi AK, Alvarado LA, Hamilton M, et al. Epithelial-
Mesenchymal Transition Markers in Breast Cancer and Pathological Responseafter
Neoadjuvant Chemotherapy. Breast Cancer (Auckl). 2018;12:1178223418788074.
40.   Scott LE, Weinberg SH, Lemmon CA. Mechanochemical Signaling of the Extracellular
Matrix in Epithelial-Mesenchymal Transition. Front Cell Dev Biol. 2019;7:135.
41.   Yang J, Antin P, Berx G, Blanpain C, Brabletz T, Bronner M, et al. Guidelines and
definitions for research on epithelial-mesenchymal transition. Nat Rev Mol Cell Biol.
2020;21(6):341-52.
42.   Zeisberg M, Neilson EG. Biomarkers for epithelial-mesenchymal transitions. J Clin
Invest. 2009;119(6):1429-37.
43.   Halder SK, Beauchamp RD, Datta PK. A specific inhibitor of TGF-beta receptor kinase,
SB-431542, as a potent antitumor agent for human cancers. Neoplasia. 2005;7(5):509-21.
44.   Hao Y, Baker D, Ten Dijke P. TGF-beta-Mediated Epithelial-Mesenchymal Transition and
Cancer Metastasis. Int J Mol Sci. 2019;20(11).
45.   Wee K, Hediyeh-Zadeh S, Duszyc K, Verma S, B NN, Khare S, et al. Snail induces
epithelial cell extrusion by regulating RhoA contractile signalling and cell-matrix adhesion. J Cell
Sci. 2020;133(13).
46.   Hargest V, Bub T, Neale G, Schultz-Cherry S. Astrovirus-induced epithelial-
mesenchymal transition via activated TGF-beta increases viral replication. PLoS Pathog.
2022;18(4):e1009716.
47.   Zhang Z, Zou J, Shi Z, Zhang B, Etienne-Mesmin L, Wang Y, et al. IL-22-induced cell
extrusion and IL-18-induced cell death prevent and cure rotavirus infection. Sci Immunol.
2020;5(52).
48.   Moshiri J, Craven AR, Mixon SB, Amieva MR, Kirkegaard K. Mechanosensitive extrusion
of Enterovirus A71-infected cells from colonic organoids. Nat Microbiol. 2023;8(4):629-39.
49.   Eisenhoffer GT, Rosenblatt J. Bringing balance by force: live cell extrusion controls
epithelial cell numbers. Trends Cell Biol. 2013;23(4):185-92.
50.   Tada M. The morphogenetic changes that lead to cell extrusion in development and cell
competition. Dev Biol. 2021;477:1-10.
51.   Franco JJ, Atieh Y, Bryan CD, Kwan KM, Eisenhoffer GT. Cellular crowding influences
extrusion and proliferation to facilitate epithelial tissue repair. Mol Biol Cell. 2019;30(16):1890-9.
52.   Katoh H, Fujita Y. Epithelial homeostasis: elimination by live cell extrusion. Curr Biol.
2012;22(11):R453-5.
53.   Kuipers D, Mehonic A, Kajita M, Peter L, Fujita Y, Duke T, et al. Epithelial repair is a two-
stage process driven first by dying cells and then by their neighbours. J Cell Sci. 2014;127(Pt
6):1229-41.

54.   Song L, Xiong H, Li J, Liao W, Wang L, Wu J, et al. Sphingosine kinase-1 enhances
resistance to apoptosis through activation of PI3K/Akt/NF-kappaB pathway in human non-small
cell lung cancer. Clin Cancer Res. 2011;17(7):1839-49.
55.   Kim KK, Kugler MC, Wolters PJ, Robillard L, Galvez MG, Brumwell AN, et al. Alveolar
epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is
regulated by the extracellular matrix. Proc Natl Acad Sci U S A. 2006;103(35):13180-5.
56.   Liu H, Ma Y, He HW, Zhao WL, Shao RG. SPHK1 (sphingosine kinase 1) induces
epithelial-mesenchymal transition by promoting the autophagy-linked lysosomal degradation of
CDH1/E-cadherin in hepatoma cells. Autophagy. 2017;13(5):900-13.
57.   Liu SQ, Xu CY, Wu WH, Fu ZH, He SW, Qin MB, et al. Sphingosine kinase 1 promotes
the metastasis of colorectal cancer by inducing the epithelial-mesenchymal transition mediated
by the FAK/AKT/MMPs axis. Int J Oncol. 2019;54(1):41-52.
58.   He F, Fan M, Jin Y, Wang H, Ding L, Fan J, et al. Sphingosine kinase 1 inhibition
decreases the epithelial-mesenchymal transition and ameliorates renal fibrosis via modulating
NF-kappaB signaling. Am J Transl Res. 2019;11(9):5879-87.
59.   Riemma MA, Cerqua I, Romano B, Irollo E, Bertolino A, Camerlingo R, et al.
Sphingosine-1-phosphate/TGF-beta axis drives epithelial mesenchymal transition in asthma-like
disease. Br J Pharmacol. 2022;179(8):1753-68.
60.   Wang W, Hind T, Lam BWS, Herr DR. Sphingosine 1-phosphate signaling induces
SNAI2 expression to promote cell invasion in breast cancer cells. FASEB J. 2019;33(6):7180-
91.
61.   Aloia AL, Calvert JK, Clarke JN, Davies LT, Helbig KJ, Pitson SM, et al. Investigation of
sphingosine kinase 1 in interferon responses during dengue virus infection. Clin Transl
Immunology. 2017;6(7):e151.
62.   Stewart CM, Bo Y, Fu K, Chan M, Kozak R, Apperley KY, et al. Sphingosine Kinases
Promote Ebola Virus Infection and Can Be Targeted to Inhibit Filoviruses, Coronaviruses, and
Arenaviruses Using Late Endocytic Trafficking to Enter Cells. ACS Infect Dis. 2023;9(5):1064-
77.
63.   Bonica J, Clarke CJ, Obeid LM, Luberto C, Hannun YA. Upregulation of sphingosine
kinase 1 in response to doxorubicin generates an angiogenic response via stabilization of Snail.
FASEB J. 2023;37(3):e22787.
64.   Bonica J, Mao C, Obeid LM, Hannun YA. Transcriptional Regulation of Sphingosine
Kinase 1. Cells. 2020;9(11).
65.   Cannavo A, Liccardo D, Komici K, Corbi G, de Lucia C, Femminella GD, et al.
Sphingosine Kinases and Sphingosine 1-Phosphate Receptors: Signaling and Actions in the
Cardiovascular System. Front Pharmacol. 2017;8:556.
66.   Maceyka M, Harikumar KB, Milstien S, Spiegel S. Sphingosine-1-phosphate signaling
and its role in disease. Trends Cell Biol. 2012;22(1):50-60.
67.   Rutherford C, Childs S, Ohotski J, McGlynn L, Riddick M, MacFarlane S, et al.
Regulation of cell survival by sphingosine-1-phosphate receptor S1P1 via reciprocal ERK-
dependent suppression of Bim and PI-3-kinase/protein kinase C-mediated upregulation of Mcl-
1. Cell Death Dis. 2013;4(11):e927.
68.   Machesky NJ, Zhang G, Raghavan B, Zimmerman P, Kelly SL, Merrill AH, Jr., et al.
Human cytomegalovirus regulates bioactive sphingolipids. J Biol Chem. 2008;283(38):26148-
60.
69.   Adada MM, Orr-Gandy KA, Snider AJ, Canals D, Hannun YA, Obeid LM, et al.
Sphingosine kinase 1 regulates tumor necrosis factor-mediated RANTES induction through p38
mitogen-activated protein kinase but independently of nuclear factor kappaB activation. J Biol
Chem. 2013;288(38):27667-79.

70.   Alvarez SE, Harikumar KB, Hait NC, Allegood J, Strub GM, Kim EY, et al. Sphingosine-
1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2. Nature.
2010;465(7301):1084-8.
71.   Xia P, Wang L, Moretti PA, Albanese N, Chai F, Pitson SM, et al. Sphingosine kinase
interacts with TRAF2 and dissects tumor necrosis factor-alpha signaling. J Biol Chem.
2002;277(10):7996-8003.
72.   Milara J, Navarro R, Juan G, Peiro T, Serrano A, Ramon M, et al. Sphingosine-1-
phosphate is increased in patients with idiopathic pulmonary fibrosis and mediates epithelial to
mesenchymal transition. Thorax. 2012;67(2):147-56.
73.   Sukocheva OA, Hu DG, Meech R, Bishayee A. Divergence of Intracellular Trafficking of
Sphingosine Kinase 1 and Sphingosine-1-Phosphate Receptor 3 in MCF-7 Breast Cancer Cells
and MCF-7-Derived Stem Cell-Enriched Mammospheres. Int J Mol Sci. 2021;22(9).
74.   Hait NC, Allegood J, Maceyka M, Strub GM, Harikumar KB, Singh SK, et al. Regulation
of histone acetylation in the nucleus by sphingosine-1-phosphate. Science.
2009;325(5945):1254-7.
75.   Chugh P, Clark AG, Smith MB, Cassani DAD, Dierkes K, Ragab A, et al. Actin cortex
architecture regulates cell surface tension. Nat Cell Biol. 2017;19(6):689-97.
76.   Kurosaka S, Kashina A. Cell biology of embryonic migration. Birth Defects Res C
Embryo Today. 2008;84(2):102-22.
77.   Eden E, Lipson D, Yogev S, Yakhini Z. Discovering motifs in ranked lists of DNA
sequences. PLoS Comput Biol. 2007;3(3):e39.
78.   Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. GOrilla: a tool for discovery and
visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics. 2009;10:48.
79.   Leonard VH, Sinn PL, Hodge G, Miest T, Devaux P, Oezguen N, et al. Measles virus
blind to its epithelial cell receptor remains virulent in rhesus monkeys but cannot cross the
airway epithelium and is not shed. J Clin Invest. 2008;118(7):2448-58.
80.   Ono N, Tatsuo H, Hidaka Y, Aoki T, Minagawa H, Yanagi Y. Measles viruses on throat
swabs from measles patients use signaling lymphocytic activation molecule (CDw150) but not
CD46 as a cellular receptor. J Virol. 2001;75(9):4399-401.
651

FIGURE LEGENDS

Fig 1. Single-cell RNA sequencing implicates two detachment pathways. (A) Schematic

depiction of infectious center formation and detachment. Green cells represent MeV-infected

cells. Cell-to-cell spread of MeV over 3-5 days results in an infectious center, left. Beginning

around 5 days post-infection, infectious centers begin to detach as a unit from the rest of the

epithelial layer, right. (B) Differentially expressed gene lists of MeV-infected HAE were derived

from a previously published scRNA-seq dataset (21). Genes with increased mRNA levels in

MeV-infected HAE at 3 days post-infection vs. mock-infected HAE were analyzed with biological

process gene ontology. Related gene ontology terms are connected with arrows. (C) Heatmap

of mRNA expression for EMT-related genes in MeV-infected HAE at day 3 post-infection. Gene

expression was normalized to mock-infected HAE and derived from the scRNA-seq dataset. (D)

Cells from the scRNA-seq dataset are visualized via t-SNE plot. Cells are clustered by similar

gene transcription. Cell types were assigned to clusters using cell type markers described in

(21). (E) Cells expressing SNAI2, a transcriptional regulator of EMT, are colored according to

mRNA expression. A cluster of cells with high SNAI2 expression is highlighted in the inset. (F)

SK1, which promotes LCE, is also expressed in this cell cluster (see inset).

Fig 2. EMT is activated within infectious centers. HAE infected with MeV-mCherry were fixed

at 3- and 7-days post-infection. Cultures were immunostained for βCAT (A), ZO-1 (B), or ECAD

(C). Confocal images of 3 infectious centers were collected per donor (n=3). Representative

images are shown. Corresponding z-stacks are shown below each image. Blue = DAPI. (D)

Percent cytoskeleton breakdown was calculated by dividing total area of breakdown for each

cytoskeletal protein by the total area of the infectious center. Multiple t-tests were performed. *

$p < 0.05$.

Fig 3. EMT contributes to infectious center detachment. (A) Graphic representation of

infectious center detachment assay. Images are taken each day post-infection. Individual

infectious centers (green dot) are tracked until they detach (represented by a red X). The day

post-infection at which each infectious center detaches is recorded. (B) HAE were treated with

either SB431542 (SB) or a vehicle control (DMSO) for 24 hours before being infected at an MOI

of 1 (MeV-GFP). Data from the detachment assay are graphed as a probability of detachment

(i.e., percent of infectious centers detached at each time point). A Gehan-Breslow-Wilcoxon test

was performed and indicated that there was a statistical significance between the two curves. *

$p < 0.05$. (C) Whole culture lysates, which may include detached infectious centers in the apical

space, were collected from MeV-infected HAE (with either SB treatment or vehicle) at the

indicated timepoint for titration via $TCID_{50}$. A one-way ANOVA with multiple comparisons was

performed. ns = not significant. (D) Live fluorescent images were collected at 3 days post-

infection. Numbers and size (E) of infectious centers were measured with ImageJ. Counts were

normalized to vehicle controls and displayed as a percentage. Shapes indicate individual donors

(n=5). Student's t-test. (E) Dots indicate individual infectious centers. Median $\pm$ interquartile

range. Student's t-test. (F) Cytotoxicity of SB treatment was determined via LDH assay at 3

days post-infection with MeV. SB and DMSO treated cells had comparable cytotoxicity when

normalized to Triton X-100 treated cells (+ control). One-way ANOVA.

Fig 4. LCE is activated within infectious centers. (A) Diagram of SK1's function. SK1

phosphorylates sphingosine to make S1P. (B) Heatmap of LCE-related gene expression in ISG-

high cells at 3 days post-infection. Shade of red indicates increased fold change over mock-

infected cells. (C) *En face* immunostaining of SK1 in HAE infected with MeV-mCherry at 3 days

post-infection. Scale bar = 50 μm. (D) Immunostained sections of MeV-mCherry infected HAE at

3 days post-infection. Arrows indicate cells with SK1 staining localized to the nucleus. Scale bar

= 25 μm.

Fig 5. LCE contributes to infectious center detachment. (A) An SK1 inhibitor (SKI-II) or (C)

exogenous S1P was applied 1-day post-infection with MeV. The infectious center detachment

assay was performed as described in **Fig 3**. Gehan-Breslow-Wilcoxon tests were performed

and indicated that there were statistical significances between the curves. * $p < 0.05$; ****

$p < 0.0001$. (B) Cell lysates were collected from MeV-infected HAE treated with SKI-II or (D) S1P

at the indicated timepoint for titration via $TCID_{50}$. A one-way ANOVA with multiple comparisons

was performed. ns = not significant.

Fig S1. Characterization of ISG-high cells. (A) The t-SNE plot from Fig 1 with GFP status

overlayed. GFP+ indicates infected cells, GFP- indicates uninfected cells from MeV-infected

HAE. Mock indicates cells mock infected. The box highlights the ISG-high cell cluster. The

adjoining graph identifies the division of ISG-high cells that are GFP+ or GFP-. (B) Violin plot of

SNAI2 expression across cell clusters. (C) Violin plot of SK1 expression across cell clusters.

Fig S2. qRT-PCR confirmation of scRNA-seq. (A) scRNA-seq results were confirmed with

qRT-PCR. Transcript changes of MeV-infected HAE are shown as a fold change over mock-

infected cells. One-way ANOVA was performed. * $p < 0.05$, ** $p < 0.01$. (B) Apical (open shapes)

and basolateral (filled shapes) secretions were collected from mock- and MeV-infected HAE at 3

days post-infection. Protein content was measured using a TGF-β ELISA kit. Shapes indicate

distinct donors (n=3). Paired Student's t-test; * $p < 0.05$. (C) HAE infected with MeV-mCherry

were fixed at 7 days post-infection and immunostained for ezrin. A z-stack image is shown

below. The arrow points to ezrin staining that has been disturbed within the infectious center.

Blue = DAPI. Scale bar = 50 μm.

Fig S3. Impact of LCE treatments on infection kinetics. Live fluorescent images of (A-B)

SKI-II treated or (C-D) S1P treated HAE were collected at 3 days post-infection with MeV.

Numbers (A and C) and size (B and D) of infectious centers were measured with ImageJ.

Counts were normalized to vehicle controls and displayed as a percentage. Shapes indicate

individual donors (n=4 in A; n=5 in C). Student's t-test; ns = not significant. (B and D) Dots

indicate individual infectious centers. Median ± interquartile range. Student's t-test. (E)

Cytotoxicity of SB treatment was determined via LDH assay at 3 days post-infection with MeV.

SKI-II, S1P, and DMSO treated cells had comparable cytotoxicity when normalized to Triton X-

100 treated cells (+ control). One-way ANOVA.

S1 Table. Primer sequences for qRT-PCR analysis.

Gene name	Direction	Sequence
MMP1	Forward	CCTAGTCTATTCATAGCTAATCAAGAGGATGT
	Reverse	AGTGGAGGAAAGCTGTGCATAC
MMP10	Forward	ATCTGAGATGCCAGCCAAGT
	Reverse	AGGGTTCCAGTGGGATCTTC
TGFB1	Forward	GGCCTCCCCACCACACCAG
	Reverse	GCCGCAGCTTGGACAGGATC
SNAI2	Forward	ATCTGCGGCAAGGCGTTTTCCA
	Reverse	GAGCCCTCAGATTTGACCTGTC
SMAD3	Forward	GCCTGTGCTGGAACATCATC
	Reverse	TTGCCCTCATGTGTGCTCTT
SFRS9	Forward	TGCGTAAACTGGATGACACC
	Reverse	CCTGCTTTGGTATGGAGAGTC

Fig 1. Single-cell RNA sequencing implicates two detachment pathways.

Fig 2. EMT is activated within infectious centers.

Fig 3. EMT contributes to infectious center detachment.

Fig 4. LCE is activated within infectious centers.

Fig 5. LCE contributes to infectious center detachment.