Title: Moderate-term dimethyl fumarate treatment reduces pathology of dystrophic skeletal and cardiac muscle in a mouse model

Authors: Stephanie Kourakis[*1,2], Cara A. Timpani[*1,2,3], Ryan M. Bagaric[1,2], Bo Qi[1,2], Benazir A. Ali[1,2], Rebecca Boyer[1], Guinevere Spiesberger[1,2], Nitika Kandhari[4], Amanda L. Peterson[5], Didier Debrincat[1,2], Thomas J. Yates[1], Xu Yan[1,6], Nicole Stupka[3,2,1], Jujiao Kuang[1], Brunda Nijagal[5], Deanna Deveson-Lucas[4], Dirk Fischer[6], Emma Rybalka[1,2,3,7#]

Authors Affiliations:

[1]Victoria University, Institute for Health and Sport (IHeS), Melbourne, Victoria, Australia

[2]Inherited and Acquired Myopathies Program, Australian Institute for Musculoskeletal Science (AIMSS), St Albans, Victoria, Australia

[3]Department of Medicine – Western Health, Melbourne Medical School, The University of Melbourne, St Albans, Victoria, Australia

[4]Monash Genomics and Bioinformatics Platform, Biomedical Discovery Institute, Monash University, Clayton, Victoria, Australia

[5]Metabolomics Australia, Bio21 Institute, The University of Melbourne, Melbourne, Victoria, Australia

[6]Sarcopenia Research Program, Australian Institute for Musculoskeletal Science, Sarcopenia Research Program, St Albans, Victoria, Australia

[7]Division of Neuropediatric and Developmental Medicine University Children's Hospital of Basel (UKBB), Basel, Switzerland

* Authorship note: SK and CAT contributed equally to this work as co–first authors.

[#]Denotes corresponding author

[#]Correspondance: emma.rybalka@vu.edu.au; Tel.: +61-3-8395-8226; Institute for Health and Sport, Victoria University, PO Box 14428, Melbourne City MC, Victoria, Australia, 8001.

Conflict of Interest: ER discloses consultancy work for Santhera Pharmaceuticals and Epirium Bio outside of this project.

Abstract:

There has been a wave of new therapeutics in Duchenne muscular dystrophy (DMD) that target the genetic, (missing) protein and flow-on pathogenic mechanisms emerging recently. These medicines will be vital to extend lifespan complementary to corticosteroids, which have been used as a standard of care tool for more than 30 years. While corticosteroids significantly slow disease progression, they also impart side effects severe enough to preclude use in some patients. We have previously demonstrated that short-term treatment with dimethyl fumarate (DMF), a drug with indication and established safety data in Multiple Sclerosis, more selectively modulates Duchenne (*mdx*) immunology than the frequently used corticosteroid, prednisone. Here, we assess the effect of moderate-term DMF treatment over 5 weeks in exercise-aggravated *mdx* mice. We show that like prednisone, DMF maintains anti-inflammatory, anti-fibrotic, and anti-lipogenic activity on muscle with moderate-term use but does not reduce muscle degeneration per se. This study supports our previous work highlighting DMF as a possible repurposing candidate for DMD, especially for patients who cannot tolerate chronic corticosteroid treatment.

Introduction:

Duchenne muscular dystrophy (DMD) is a severe, progressive, muscle-wasting disease caused by mutations in the *DMD* gene. Dystrophin loss impairs the junction between the intracellular cytoskeleton and extracellular matrix (ECM), which stabilises the muscle fibre during contraction. Lack of dystrophin results in chronic muscle damage, degradation, fibrosis and adiposis (1). Affecting predominantly males, DMD is typically diagnosed by ~5 years of age presenting as delayed motor milestones and frequent falls (2). By early adolescence, loss of ambulation and pulmonary function decline is common (3-5). Pulmonary distress is complicated by diaphragmatic weakness and fibrosis, and scoliosis, which alters the shape of the thoracic cavity and reduces respiratory capacity (6). Natural history data suggest that cardiac symptoms can emerge from as early as 12-14 years leading to dilated cardiomyopathy in almost all patients even with prophylactic treatment with cardiac medication, which can delay the onset (7-9).

The DMD therapeutic landscape has grown significantly over the past decade. Since 2016, four exon-skipping antisense oligonucleotide drugs (Exondys 51, Vyondys 53, Viltepso and Amonndys 45) and two gene replacement therapies (Elevidys and fordadistrogene movaparvovec) have been granted accelerated access (10). However, their efficacy is unclear, and fatalities have occurred with gene replacement strategies raising safety concerns (11, 12). While these therapies have the best chance of providing long-term disease-modifying benefits, there are relatively fewer therapeutics in development to address the secondary pathomechanisms, especially the hyper-inflammation and immune system activation that drives muscle fibrosis (13). Anti-inflammatory corticosteroids have prevailed as the pharmacological standard of care (SOC) for DMD for >3 decades targeting this aspect. With early intervention, they effectively slow disease progression, prolonging ambulation, and pulmonary function (14). Yet, they have an extensive side-effect profile, e.g., weight gain, osteoporosis and metabolic syndrome (14). Approved in 2023, steroid analogue Agamree (vamorolone) appears to reduce some of these side-effects. However, the typical corticosteroid side effects of weight gain, behavioural issues and cushingoid symptoms persist (15, 16)

indicating that non-steroid interventions may be better for long-term use. Recently approved non-steroidal anti-inflammatory, Duvyzat (givinostat) slows loss of muscle strength (17) yet significant side effects were reported in the Phase III trial (e.g., diarrhoea, vomiting, thrombocytopenia) indicating a well-balanced efficacy-to-safety profile is critical.

We recently identified dimethyl fumarate (DMF), a drug prescribed for the autoimmune diseases, relapsing-remitting Multiple Sclerosis (MS) and psoriasis (18), as a repurposing candidate for DMD. DMF is a potent immunoregulatory and anti-inflammatory and -oxidative drug via nuclear factor erythroid 2-related factor 2 (Nrf2) activation and hydroxycarboxylic acid receptor 2 (HCAR2) agonism (18). Our proof-of-concept study in the *mdx* mouse demonstrated short-term DMF treatment was more effective against key disease indices than SOC prednisone (PRED), including histopathology, muscle function, mitochondrial metabolism and disease driving gene signatures (19). In the present study, we aimed to evaluate moderate-term effects in *mdx* mice with an aggravated phenotype. To pre-empt that future clinical trial participants would be steroid treated, we also aimed to assess combinatorial treatment, i.e., DMF+PRED, and potential drug interactions.

Results:

Exercise aggravation increases plasma creatine kinase (CK) levels and worsens dystrophic muscle histopathology.

The dystrophic *mdx* mouse is the most used pre-clinical drug screening tool despite its relatively mild phenotype compared to patients. Phenotype differences between sedentary (SED) wild-type (WT) and *mdx* mice (Figure 1) are obvious at 8 weeks age, including elevated plasma (Figure 1B) and skeletal (Figure 1D-H, J-Q) and cardiac (Figure 1I & R) muscle damage indicators. However, since translational outcomes have been particularly poor for DMD, more rigorous experimental approaches are needed. Implementation of TREAT-NMD's standard operating procedures enables the standardisation of pre-clinical data across laboratory groups for improved translational outcomes. We used twice-weekly forced horizontal treadmill running from 35 days age to aggravate the *mdx* phenotype (Figure 1A) according to TREAT-NMD SOP DMD_M.2.1.001 (20).

Exercise aggravation (AGG) significantly increased plasma creatine kinase (CK) levels in *mdx* mice (Figure 1B) but had no impact on forelimb grip strength (Figure 1C) or other function measures (Supplemental Figure 1A-C). While neither anthropometric changes nor respiratory or *ex vivo* force deficits were observed (see also Supplemental Figure 1 A-L), AGG did exacerbate dystrophic histopathology. We analysed quadriceps, which are particularly damaged during exercise (21, 22), and heart, to assess indices of muscle damage, inflammation, regeneration, immune cell infiltration and fibrosis. AGG increased the unhealthy/healthy tissue ratio (*p*=0.0527; Figure 1D), percentage of regenerating centronucleated fibres (Figure 1E-H), CD68+ macrophage infiltrate (*p*=0.0527; Figure 1J-M), and fibrosis (Figure 1N-Q) in *mdx* quadriceps (our data are consistent with levels of fibrosis in 8 week old *mdx* relative to WT as reported previously (23, 24)) but not hearts (Figure 1I and R) relative to *mdx* SED controls. Perilipin+ fat infiltration was minimal and not impacted by aggravation (Supplemental Figure 1M). We also assessed mitochondrial respiration in flexor digitorum brevis (FDB) fibres using

extracellular flux and a mitochondrial stress test (Figure 1S). *Mdx* AGG FDB fibres had an overall higher energetical state (Figure 1T) in line with increased mechanical damage and regenerative activity relative to SED *mdx* and WT controls. However, AGG had no effect on global cytokines levels (Supplemental Figure 1N) or urinary oxidative stress biomarker, 8-isoprostane (Supplemental Figure 1O).

DMF improves grip strength and transcriptional control of myofibril assembly.

Following 5 weeks of daily treatment, endpoint plasma CK levels were unchanged by DMF or PRED treatment. Surprisingly, combinatorial DMF+PRED increased CK levels (Figure 2A). Both DMF and DMF+PRED treatment increased forelimb grip strength (corrected for body weight) at the experimental endpoint (Figure 2B) (notably, this measure was higher in *mdx* VEH mice compared to WT (Figure 1C)) but there was no effect of treatment on the hang test minimal holding impulse, which assesses both strength and endurance (Figure 2C). DMF treated alone and in the DMF+PRED combination reduced ambulatory distance (assessed over 10-minutes in the open-field; Figure 2E) but had no effect on rotarod performance (Figure 2D). Whilst PRED did not improve ambulatory capacity in the open field, it significantly improved time to fall in the rotarod test suggesting some improvement of neuromotor coordination.

In comparison to *mdx* VEH, all treatments increased the fat mass index (Figure 2F), which was reduced by ~40% in *mdx* relative to WT mice (Supplemental Figure 1D). Since all treatments increased the fat mass index, we probed the muscle (quadriceps) transcriptome for genes associated with fatty acid metabolism. Of the 16 upregulated genes in *mdx* mice (compared to WT), DMF downregulated 9 genes, PRED downregulated 2 genes and DMF+PRED downregulated 10 genes (Figure 2G). Consistent with the known suppressive effects of PRED on muscle and bone growth (14, 25, 26), PRED treated alone and in the DMF+PRED combination, significantly reduced the lean mass index relative to *mdx* VEH (Figure 2H). At the muscle transcriptome level, PRED significantly downregulated many genes involved in branched-chain amino acid (BCAA) catabolism but upregulated *Bcat1* (branched chain amino acid transaminase 1), which was already upregulated by 4-fold in *mdx* AGG relative to WT (Figure 2I). Hydration index was reduced by PRED treatment in line with reduced lean mass (Figure 2J). PRED reduced liver mass and all treatments normalised the higher quadriceps mass noted in *mdx* AGG (and SED) relative to WT (Supplemental Table 1 and 2) suggesting multi-tissue drivers of the overall reduction in lean mass index.

At the experimental endpoint, EDL, and SOL muscle, predominantly fast- and slow-twitch respectively, were harvested for *ex vivo* contractile testing. There was no significant effect of treatments on either EDL or SOL absolute or specific force production (Figure 2, K-N) and only DMF+PRED combination affected the force frequency relationship by increasing force production at 60 and 80Hz (Supplemental Figure 2A). No treatment impacted fatigue or recovery measures (Supplemental Figure 2C-D). Notably, *ex vivo* contraction measures were comparable between WT and *mdx* EDL and SOL indicating no scope for improvement (Supplemental Figure 1K-L). We further probed transcriptional control of muscle contraction pathways for molecular changes in mixed fibre type muscle (i.e., quadriceps) that may be induced by treatments but not captured through *ex vivo* studies using fibre type exclusive/predominant muscles (i.e., EDL and SOL). Slow troponin isoforms, *Tnni1* and

Tnnt1, were significantly upregulated in *mdx* AGG compared to WT muscle (Figure 2O), consistent with previous studies in dystrophic muscle (27). DMF significantly downregulated both slow troponin isoforms, but upregulated *Myh3, Myh8* and *Tnnt2,* genes associated with regenerating DMD muscle (27, 28). Comparatively, PRED only upregulated *Myh8*. The downregulation of *Actn2* by all treatments suggests concomitant sarcomere remodelling but only DMF and DMF+PRED treatment upregulated expression of *Tmod1, Neb* and *Actn3* (and *Mybpc2* for DMF treatment), promoters of sarcomere stability (29, 30) in the absence of dystrophin.

DMF moderates the immune response in *mdx* quadriceps muscle.

Persistent activation of the innate immune system leads to hyperinflammation and tissue damage in dystrophic muscles (13). The unhealthy/healthy tissue ratio was significantly higher in *mdx* AGG quadriceps (Figure 1D) but was not reduced by any drug treatment (Figure 3A, C-F). However, there was a trend (p=0.0734) for DMF to reduce the percentage of centronucleated regenerating fibres (Figure 3B-F) suggesting reduced muscle damage. Further investigation of the myogenesis transcriptional pathway revealed that the upregulated *mdx* genes involved in muscle regeneration, *Myog, Myod1, Tcf4, Mapk11* (Figure 3G), were moderated by treatment. DMF downregulated *Myog* and *Mapk11* while PRED downregulated *Myod1 and Mapk12,* but in combination, DMF+PRED only downregulated *Tcf4*. Pan macrophage marker, CD68, was significantly reduced by DMF and PRED (Figure 3H-L), consistent with their anti-inflammatory profiles. Interestingly, combined delivery of DMF+PRED had no effect on macrophage infiltrate. We probed our transcriptomic dataset for commonly dysregulated inflammatory markers associated with DMD, including *Ccl2, Ccl6, Ccl7, Ccl8, Ccr2, Cx3cr1, Nfkb1* and *Nfkb2* (Figure 3M). 7 genes were significantly upregulated in *mdx* VEH quadriceps and DMF treatment significantly downregulated 5 (*Ccl2, Ccl7, Ccl8, Ccr2* and *Ccl6*) and trended to downregulate *Nfkb2* (p=0.0702), a component of the master regulator of innate immunity. Both PRED and DMF+PRED downregulated fewer inflammatory genes: one (*Cx3cr1*) and three (*Ccl7, Ccl2* and *Ccr2*), respectively.

DMF decreases fibrosis in the quadriceps and downregulates key disease driving gene, *Timp1*.

In a recent meta-analysis, five genes were identified as candidate seed genes driving the DMD disease module (*Timp1, Spp1, Fn1, Mmp2* and *Igf1*) (31). A more recent study identified a *Bmp4*-induced molecular signature in DMD patient muscles involving upregulated expression of *Serpling1, Adamts3, HCAR2, Smad8* and *Unc13c* (32). We show that *Timp1, Spp1* and *Fn1* (Figure 4A) and *Serping1* (Figure 4B) are upregulated in *mdx* AGG quadriceps alongside fibrosis (Figure 4C-D). DMF treatment significantly decreased *Timp1* expression (Figure 4A) and fibrosis of quadriceps (Figure 4C and E) while PRED upregulated *Spp1, Fn1, Igf1* and *Bmp4* transcription and DMF+PRED increased *Spp1* expression (Figure 4A). Neither PRED nor DMF+PRED affected quadriceps fibrosis – in fact, DMF+PRED treatment worsened it (Figure 4C, F-G). We probed 3 Reactome-based fibrosis pathways in our transcriptomics dataset: collagen formation, ECM degradation and ECM organisation. 18 genes associated with collagen formation were upregulated in *mdx* AGG quadriceps and DMF downregulated *Col23a1* (Figure 4H). In contrast, PRED upregulated 15 genes and

DMF+PRED treatment had no effect. Of the 13 upregulated ECM degradation-associated genes in *mdx* quadriceps, DMF downregulated 2 (*Casp3* and *Mmp11*) and PRED, either alone or combined with DMF, downregulated 1 (*Adamts5*) (Figure 4I). PRED and DMF+PRED treatment further upregulated expression of 8 and 5 genes, respectively. The 1 downregulated gene in *mdx* quadriceps, *Capn3*, was upregulated by DMF treatment only. There were 18 upregulated genes in the ECM organisation pathway in *mdx* quadriceps and DMF significantly downregulated 2 (*Ntn4* and *Trappc4*), PRED upregulated 11 and downregulated 1 (*Itgb6*), and DMF+PRED significantly upregulated 2 (*Bgn* and *Itga9*) and downregulated 3 (*Itgb6, Ntn4* and *Trappc4*) (Figure 4J).

DMF reduces muscle lipid content through modulation of the adipogenic and sphingolipid metabolism transcription programs in *mdx* muscle.

Muscle adiposis is prominent in human DMD muscle but minimal in *mdx* muscle, i.e., perilipin-1, a protein that localises to lipid droplets, was comparable between WT and *mdx* quadriceps (Supplemental Figure 1M). Nevertheless, DMF treatment decreased lipid droplet abundance whereas PRED, either alone or in combination with DMF, had no effect (Figure 5A-E). Since DMF modulates lipid metabolism in MS (33), we probed lipid metabolism and fibro-adipogenic progenitor (FAP)-related pathways in our transcriptomics dataset. Of the 3 established FAP-associated marker genes (*Cd34, Pdgfra* and *Dcn*), *Cd34* expression was upregulated in *mdx* quadriceps and was downregulated by DMF (Figure 5F). Reactome pathways analysis revealed the tumour necrosis factor receptor (TNFR)-mediated ceramide production pathway was significantly upregulated in *mdx* quadriceps and was normalised by DMF but not PRED treatment (Figure 7I). We probed genes within this pathway (Figure 5G) and revealed DMF normalised *Tnfrsf1a* and *Nsmaf* expression. Sphingolipid metabolism pathways were also probed since DMF is known to modulate sphingolipids through HCAR2 agonism in the context of MS (33). Of the 11 upregulated sphingolipid metabolism genes identified in *mdx* muscle, DMF downregulated 2 while PRED and DMF+PRED significantly upregulated 9 and 5 genes, respectively (Figure 5H). Furthermore, DMF normalised the expression of *Sgms1* and *Cers1*, which were downregulated in *mdx* quadriceps. 10 adipogenesis genes were dysregulated in *mdx* muscle: 4 upregulated and 6 downregulated (Figure 5I). Of the 4 upregulated genes, DMF downregulated 3 (*Cidea, Fabp4* and *Rbbp4*), PRED downregulated 1 (*Rbbp4*) and DMF+PRED downregulated 2 (*Cidea* and *Fabp4*). Of the 6 downregulated genes, DMF upregulated 4 (*Cdh4, Gatad2b, Carm1* and *Smad4*), PRED further downregulated 1 gene (*Lpl*), and DMF+PRED upregulated 2 (*Chd4* and *Carm1*) and further downregulated *Lpl*. Finally, we used targeted lipidomics to assess the lipid species profile in quadriceps muscles. 20 dysregulated lipid species were detected in *mdx* quadriceps relative to WT (11 upregulated and 9 downregulated) but these were not modulated by any treatment (Figure 5J). Of note, both PRED and DMF+PRED modulated normally expressed lipid species, 1 and 4, respectively.

DMF transcribes a highly specific anti-inflammatory and anti-oxidation signature in muscle.

A well-established mechanism of action (MOA) of DMF is via activation of Nrf2, which initiates a potent cytoprotective response (Figure 6A). Deep transcriptome profiling did not

reveal changes to expression of *Nf2el2* (Nrf2), or its negative repressor, *Keap1*, in response to DMF treatment. However, DMF normalised the otherwise upregulated expression of proteasomal subunit genes, *Psmb1, Psme1, Psma4* and *Psma1* (Figure 6B), which degrade Nrf2 and prevent its activity. DMF also tempered expression of antioxidant/detoxification enzymes *Ccs, Prdx2* and *Txn2* (Figure 6C). Our pathways analysis revealed DMF trended to drive DNA damage reversal pathways (FDR=0.0686; data not shown), thus we further probed this pathway. While *Alkbh5* was the only downregulated gene in *mdx* AGG quadriceps, DMF upregulated expression of both *Alkbh5* and *Fto*, and DMF+PRED upregulated *Alkbh5* expression (Figure 6D). We next probed the interleukin (IL) signalling pathways where 15 genes were dysregulated in *mdx* quadriceps (14 upregulated, 1 downregulated; Figure 6E). Of the 14 upregulated genes, DMF reduced expression of 5, PRED further upregulated 3 whereas DMF+PRED up- and down-regulated 2 each. DMF modulated more dysregulated genes associated with the TNF signalling pathway than either PRED or DMF+PRED (Figure 6F). DMF reduced 4 upregulated genes related to the pro-inflammatory response (*Fadd, Tnfrsf1a, Otulin* and *Rps27a*) and upregulated 5 genes associated with anti-inflammation (*Cyld, Usp4, Spata2, Sppl2a* and *Xiap*). Comparatively, PRED further downregulated *Optn* expression and DMF+PRED normalised *Tnfrsf1a*. To understand whether modulation of inflammatory signalling at the transcriptomic level changed circulating cytokines, we conducted Bio-Plex™ immunoassay on plasma. Only 2 of 23 cytokines were dysregulated in *mdx* plasma and only DMF+PRED treatment modulated them by further upregulating the expression of MCP-1 and normalising expression of G-CSF (Figure 6G). No effect of treatment was observed on urinary oxidative stress biomarker, 8-isoprostane (Figure 6H), albeit 8-isoprostane levels were comparable between *mdx* and WT samples (Supplemental Figure 1O).

DMF reduces purine degradation and mitobiogenic stimuli but has no effect on respiratory flexibility.

Since short-term DMF increased mitochondrial respiration via anaplerosis (19), we assessed metabolic function in animals via a fatigue run and respiratory parameters in response to moderate-term treatment (Figure 7A). DMF had no significant effect on fatigability during running (Supplemental Figure 3A) or basal, ATP-linked, maximal, or non-mitochondrial respiration (Figure 7A and Supplemental Figure 3B-E), PRED decreased coupling efficiency (Supplemental Figure 3F) and spare respiratory capacity (Supplemental Figure 3G), a measure of mitochondrial responsivity to stress, while DMF+PRED reduced maximal respiration and the spare respiratory capacity. Metabolic phenograms were produced to survey proportionate oxidative (OCR) to glycolytic (ECAR) metabolism and revealed *mdx* VEH FDB fibres were more energetic – an effect that was normalised by all treatments.

Untargeted polar metabolomics revealed Coenzyme A and acetyl-CoA (Figure 7C), adenosine monophosphate (AMP; Figure 7D), adenine (Figure 7E), NADP and NADH (Figure 7H) and U5DP (Supplemental Figure 3I) were reduced, while hypoxanthine (Figure 7E) and maltose and maltotriose (Figure 7F) were increased in *mdx* AGG quadriceps indicating the stimuli for bioenergetical demand. No differences in fatty acid (Figure 7G) or polyamine metabolism (Supplemental Figure 3H) were observed. Of the 10 dysregulated metabolites, DMF normalised hypoxanthine, a purine degradation metabolite, while PRED, alone and in DMF+PRED combination, had no effect. The full metabolomic signature is presented in

Supplemental Figure 3J and principal component analysis (PCA) and volcano plots are shown in Supplemental Figure 4.

Transcriptomic pathways analysis revealed pathways associated with mitochondrial termination, elongation, initiation, and translation and TNFR1-mediated ceramide production were upregulated in *mdx* muscle (Figure 7I and Table 1). All treatments downregulated these pathways, except for the TNFR1-mediated ceramide production pathway, against which only DMF was effective. We probed our dataset for mitochondrial biogenesis-related transcripts (Figure 7J) to determine whether reduced mitochondrial number could explain the downregulation. Eight mitochondrial biogenesis genes were upregulated, but transcription of traditional biogenesis enzyme biomarker, *Cs* (citrate synthase) was reduced in *mdx* muscle (Figure 7K). DMF and DMF+PRED downregulated more biogenesis (7 vs. 6 vs. 2, respectively) and TCA cycle (9 with DMF and DMF+PRED) treatment, 2 with PRED treatment. Two uncoupling proteins (UCP) genes were probed: *Ucp2* was significantly upregulated in *mdx* AGG quadriceps and further upregulated by PRED and DMF+PRED treatment, while *Ucp3* was normally expressed in *mdx* muscle, but upregulated by all 3 treatments (Figure 7L). Since fumarate is also endogenously produced by the purine nucleotide cycle (PNC), we probed this pathway for treatment effects. Of the 3 downregulated genes in *mdx* quadriceps, only DMF upregulated *Ampd1* (Figure 7M).

DMF significantly reduces cardiac fibrosis.

While cardiorespiratory decline is mostly evident in older *mdx* mice, we assessed heart histology and respiratory function since we aggravated our model. Cardiomyocyte degeneration was greater in *mdx* hearts, however, DMF was unable to reduce this (Figure 8A-C). In contrast, PRED trended to increase cardiomyocyte degeneration ($p=0.0718$; Figure 8A and D) while DMF+PRED significantly increased this histopathologic feature (Figure 8A and E). DMF was the only treatment to reduce cardiac fibrosis (Figure 8F-J) consistent with its anti-fibrotic effects in quadriceps (Figure 4C). Since there was no scope for modulation of respiratory function in *mdx* mice, as plethysmography readouts were comparable to WT, no treatment had any affect (Figure 8K-M). Similarly, thoracolumbar deformity was comparable between WT and *mdx* AGG and treatments had no effect on kyphosis index (Figure 8N-R).

DMF significantly recovers inflammatory, fibrosis and fat mass associated parameters.
The recovery score is used to express the effect of a treatment, not just by the difference between treated and untreated *mdx* mice, but relative to the extent of the deficiency between WT and *mdx* mice (34). Here, we calculated the recovery score (Figure 9) of parameters that were significantly affected by AGG. CD68 macrophage infiltration of quadriceps was improved by both DMF and PRED to a similar extent (65% vs. 66%, respectively) whilst DMF+PRED showed no improvement (0.78%). Fat mass index was recovered more so by DMF and DMF+PRED (59% and 44%, respectively) than PRED, which had no effect (-1.27%). DMF was the most effective at recovering cardiac fibrosis (46%) followed by PRED and DMF+PRED, which had a comparable recovery score (20% and 24%, respectively). Similarly, DMF was the most effective treatment to recover quadriceps fibrosis (37%) followed by PRED (23%). DMF+PRED resulted in a significant reduction of recovery score (-47%).

Discussion:

The *mdx* mouse is a useful pre-clinical therapeutic screening tool but presents with challenges that may compromise translational outcomes later in the drug development pipeline. After an initial acute degenerative period during rapid maturational growth at ~3-5 weeks age, the disease stabilises into relatively mild cyclical degeneration and regeneration bouts across the lifespan until ~12 months age when disease progression escalates (34). Applied exercise exacerbates muscle dystrophy in the *mdx* mouse (22, 35, 36) and here, we used moderate, twice-weekly forced horizontal treadmill running to aggravate disease to better scope DMF's efficacy and translational potential. Our methodological approach was successful to increase clinically relevant muscle damage biomarker, plasma CK, and histopathological hallmarks in hindlimb quadriceps muscle. Cardiorespiratory phenotypes are usually only evident from ~12-18 months age in *mdx* mice (37) and it was encouraging that cardiac muscle degeneration and fibrosis was induced by exercise aggravation. Increasing the frequency (e.g., thrice weekly running), duration (e.g., >20 mins per session) or difficulty (e.g., downhill running) of the exercise regimen could induce both histopathological and functional phenotypes across all impacted physiological systems in future work.

Nrf2 is a well-established MOA of DMF in the context of MS. In our short-term DMF screen, we demonstrated Nrf2 activation via increased expression of Phase II antioxidant enzymes, NQO1 and SOD1 (19), the biomarkers of DMF's MOA and efficacy in MS (38). However, in this study, moderate-term DMF treatment, at least transcriptionally, did not maintain NQO1 or SOD1 upregulation. DMF likely induces temporal Nrf2 activation in response to environmental (oxidant, inflammation) cues to maintain a proportionate antioxidant defence system. For example, GSK-3β (glycogen synthase kinase 3β) phosphorylates Nrf2 resulting in its nuclear exclusion and degradation independent of Keap1 (39). GSK-3β is possibly an oxidant sensitive override mechanism to avoid hyper-Nrf2-activation. DMF treatment (alone and in combination with PRED) increased GSK-3β transcription indicating this mechanism is predominate and antioxidative protein expression sufficient to handle the myocellular redox climate. Our data highlight that DMF's MOA over the moderate term is to regulate highly specific immune modulation, possibly via HCAR2 agonism. We show that DMF was able to recover histopathological exacerbations induced by aggravation of the *mdx* phenotype by on average, 50% (Figure 9). In contrast, PRED recovered on average, 26%. DMF is twice as effective at treating these measures and from a safety standpoint could be better, since it does not reduce lean tissue mass like PRED.

In the context of human MS, DMF is known to re-program fat metabolism, especially sphingolipid metabolism (33). Our data highlight multiple points of control of systemic and muscle-specific adipose tissue levels. *Mdx* mice (SED and AGG) are particularly lean (low fat mass index relative to WT) and our multi-omic data point to increased metabolic demand as a mechanism. DMF treatment normalised these multi-omic signatures and the fat mass of *mdx* animals. The primary active metabolite of DMF, monomethyl fumarate (MMF), is a ligand for HCAR2 which is highly expressed on adipocytes and immune cells (40). HCAR2 agonism prevents lipolysis through inhibition of adipose triglyceride and hormone sensitive lipases (33). In contrast to these systemic effects, DMF reduced lipid droplet abundance (perilpin-1) at the

muscle level. Via its distinct immunomodulatory effects on macrophage sub-populations, DMF downregulated FAP-associated marker genes, which is important because a subset of FAPs can undergo adipogenic differentiation contributing to fatty infiltration of DMD muscle (41). DMF also reduced *Tnfrsf1a* transcripts which were upregulated in *mdx* muscle. TNFR1 agonism contributes to muscle adiposis by inducing sphingomyelin hydrolysis-mediated ceramide production. Ceramides are bioactive molecules capable of activating apoptotic and pro-inflammatory pathways and their accumulation in muscle is linked to multiple myopathies (42, 43). Consistent with a study by Laurila *et al.*(44) in *mdx* mice, we show that the sphingolipid metabolism pathway is upregulated in dystrophic muscle. Although DMF downregulated only 2 of 10 significantly upregulated genes within this pathway, PRED led to further upregulation of 9 genes. Sphingolipid metabolism is heightened in MS resulting in ceramide production (45) and more recently, sphingosine-1-phosphate (S1P), as well as transcripts of other sphingolipids were shown to be highly upregulated in dystrophic human and *mdx* muscle (46). Muscle adiposity is much more of a problem in human than *mdx* DMD (47, 48), as is obesity (49, 50). Whether DMF will be efficient enough in humans given that the adipose pathology is worse is an important translational question for follow up. If it were to be similarly effective against human DMD muscle adiposis, DMF could be an ideal repurposing candidate.

The most striking effect of DMF in our study was reducing fibrosis of the heart and quadriceps although it is important to note that fibrosis was mild in our 8-week-old mdx mice (~4-5%consistent with others (23)) even with exercise aggravation. Recently, we showed DMF could modulate fibrosis-associated seed genes (*Mmp2, Spp1* and *Timp1*) (19) that drive the DMD disease program described by Lombardo *et al.* (31). Here we show that moderate-term DMF treatment sustains *Timp1* suppression, whilst PRED significantly upregulates expression of 3 seed genes (*Spp1, Fn1* and *Igf1*). DMF downregulated key genes involved in collagen formation, ECM degradation and organisation whilst PRED significantly upregulated over more than half (34 out of 54 dysregulated genes). DMF was previously shown to ameliorate lung (51-53) and renal (54) fibrosis indicating systemic anti-fibrotic function. Of the several possible molecular mechanisms regulating DMFs anti-fibrotic effect, our data indicate reduced (CD68+) macrophage recruitment/accumulation/transition as central. We saw no evidence of DMF-dependent inhibition of TGF-ß/Smad3 signalling or Spp1/osteopontin expression. Notably, when treated in combination DMF+PRED exacerbated exclusively quadriceps fibrosis (-47% recovery score). This effect could be a result of over-immunosuppression, highlighting the intricate symphony of immunology required to maintain muscle over fibrotic tissue. As a weight-bearing and high force generating muscle group, quadriceps are more prone to damage than the heart and therefore seem more sensitive to the detrimental effect of over-immunosuppression. This aspect requires careful consideration in the planning of future clinical trials testing DMF and other immunomodulatory/anti-fibrotic agents for DMD where SOC may need to be withheld or exclusively steroid naïve participants recruited.

A notable effect of short-term but not moderate-term DMF treatment was mitochondrial anaplerosis (19). Endogenous fumarate production is achieved within the mitochondrial TCA cycle via reversible fumarase activity from substrate succinate or malate, and the cytosolic PNC via adenylosuccinate lyase (ADSL) from substrate adenylosuccinate. ADSL-synthesised fumarate can be transported into the mitochondria via the malate-aspartate shuttle (after conversion of fumarate to malate in the cytosol) where exchanged aspartate is redirected into

the PNC to synchronise these two metabolic systems (55). Adding exogenous fumarate to the system, i.e., DMF treatment, at least acutely, amplifies ATP production potential. However, prolonged exposure in the present study, drove transcriptional changes to reduce mitochondrial ATP producing capacity. Historically, purine depletion was linked to DMD pathogenesis, and especially, to deficient purine salvage mechanisms (56). The PNC functions to trap inosine monophosphate (IMP) produced from ATP hydrolysis or *de novo* purine biosynthesis, protecting it from further degradation (IMP>inosine>hypoxanthine>xanthine) and flux from muscle (as uric acid). Our data show transcriptional downregulation of all three PNC enzymes (*Adss1*, *Adsl* and *Ampd1*) in *mdx* muscle as has been reported in human DMD muscle (57), and increased degraded purine metabolites, hypoxanthine, and adenine. DMF treatment normalised *Ampd1* expression and hypoxanthine levels. Mechanistically, defending purine salvage may be a useful strategy to de-burden mitochondria and ultimately to reduce reactive oxygen species production. Mitochondria are important signalling hubs directing cellular metabolism, innate immunity, cell death and the stress response (58). While mild mitochondrial stress may be beneficial to trigger a protective response (mitohormesis) severe and/or prolonged energy stress may contribute to oxidant-induced damage, fibroblast activation and ECM pathology (58).

Our study reveals important translational aspects. Firstly, immune system involvement in DMD progression is complex as its pharmacological control. Timely and successful muscle regeneration and repair is highly dependent on a well-regulated inflammatory cascade (59) – chronic inflammation and cytokine signalling can promote macrophage phenotype shifts and muscle pathology, while over-immunosuppression can dysregulate muscle regeneration. Alone, DMF and PRED share similar anti-inflammatory and immunomodulatory capability (transcriptionally) to effectively reduce macrophage infiltration of muscle. From a translational perspective, DMF could be a relevant PRED surrogate to slow DMD progression on this point. However, our omics studies reveal vastly different molecular mechanisms, and it is currently unclear which mechanism provides the best long-term results. While our data indicate DMF is a more effective modulator of fibrosis, it would be remiss not to point out that we used a higher PRED dose (5 mg.kg^{-1}/day) than commonly used in preclinical studies, yet a clinically precise one (0.75 mg.kg^{-1}/day (60) guide-lined human dose (60) corrected for mouse metabolism using 12.3 multiplier applied as per FDA guidelines (61)), which may have over-immunosuppressed mice. When combined, DMF+PRED exacerbated quadriceps fibrosis highlighting the probability of over-immunosuppression, drug interactions or both. Recent protocols for murine PRED dosing have emerged demonstrating pulsed delivery (1 mg.kg^{-1} once per week) best mimics human comparable benefits of daily treatment (62). Similarly, weekend pulsing (1-4 mg.kg^{-1} on both days) maintains clinical benefits but with reduced side-effects in DMD patients (63). While our PRED dosing strategy somewhat limits the translational relevance of our DMF findings based on head-to-head comparison, it does flag the implications of overdoing immunomodulation. Secondly, while DMF did have clear anti-fibrotic and anti-lipogenic properties, we saw no improvement of plasma CK levels and muscle degeneration/regeneration histology, the most impacted *mdx* AGG variables, or *in vivo* and *ex vivo* muscle function which were remarkably comparable to WT. Model selection is important in pre-clinical DMD studies and we expected AGG would induce a more human comparable phenotype. The use of aged

mdx mice, which do show human relevant characteristics particularly regarding fibrosis and loss of function, will be crucial for well-informed clinical translation decisions.

In summary, our data highlight DMF could be a promising translational candidate for DMD. Our follow-up study due for completion in 2025 intends to confirm longitudinal, disease modifying capacity as a final hurdle to clinical translation. Moderate-term DMF had robust anti-inflammatory, anti-fibrotic and anti-lipogenic effects against the mild DMD phenotype of 8-week-old exercise aggravated *mdx* mice. A strong point of our study was the inclusion of a combinatorial DMF+PRED group to survey potential drug interactions. Worsening of skeletal muscle fibrosis with combined treatment suggests careful consideration of dose and dosing regimen will be needed during clinical trial design. We note the high dose of PRED used in our study (to match our short-term dosing protocol) and daily administration which contradicts current preclinical corticosteroid dosing recommendations (63) as limitations. These aspects have been addressed in our on-going longitudinal study and will better inform anticipated drug interactions and dosages in humans.

Methods:

Animals
Male C57BL/10ScSnJ (WT) and dystrophin-deficient C57BL/10ScSn-Dmd^{mdx}/J (*mdx*) mice were bred at Western Centre for Health, Research and Education (Sunshine Hospital, Victoria, Australia). Animals were on a 12:12 hour light-dark cycle at 20-25°C and 40% humidity. Litters were weaned at 21 days of age and randomly assigned to cages (up to 4/cage). Food and water consumption was monitored weekly (Supplemental Figure 5A-D).

Treatment
Mdx mice were randomly assigned to treatment groups at 21 days of age: (1) 0.5% methylcellulose (VEH), (2) DMF, (3) PRED or (4) a DMF and PRED combination (all treatments suspended in VEH). WT mice were treated with 0.5% methylcellulose VEH. Animals were weighed and treated daily up until 8 weeks of age via oral gavage. Treatments were prepared relative to individual body weights to give a final daily dosage of 100mg.kg^{-1}/day DMF, 5mg.kg^{-1}/day PRED or 100mg.kg^{-1}/day DMF with 5mg.kg^{-1}/day PRED. Selected dosages are consistent with our previous study (19), which aligns with pre-clinical studies assessing DMF for MS (64) and work by Manico *et al.*(65) who pre-clinically used this PRED dosage in *mdx* mice. There was no impact of any treatment on body weight or food and water consumption in *mdx* mice (Supplemental Figure 5A-D).

Treadmill exercise-aggravation protocol
From 4 weeks of age, mice in the aggravated cohort were subjected to a 30 minute run on a horizontal treadmill (PanLab Harvard Bioscience) at 12 m/min, twice-weekly based on the protocol developed by TREAT-NMD (SOP DMD_M.2.1.001) (20). All animals were exercised in a temperature and light controlled environment and the exercise was performed with constant monitoring. A forced treadmill run to fatigue, starting at 5 m/min speed for 5 minutes and increasing by 1 m/min thereafter, was substituted for the final treadmill run and was undertaken 3 days prior to the experimental endpoint.

Functional muscle strength and neuromotor coordination testing

Forelimb and whole-body grip strength was assessed as previously stated by us (19). Neuromotor coordination was measured across 3 trials on a Rotarod (Ugo Basile, VA, Italy) at the experimental endpoint at 8 weeks of age. Mice were acclimatised one day prior to the test and each trial started with a stabilizing period followed by acceleration (up to 45rpm) until either the animal fell from the rotarod, or 600 seconds was reached (21). Time to fall (in seconds) was recorded for each mouse. All functional testing was performed blinded.

Respiratory function

Respiratory function was assessed at the experimental endpoint at 8 weeks of age in conscious, unrestrained mice using a whole-body, non-invasive plethysmograph (WBP; Data Sciences International, USA). The plethysmograph and bias flow regulator was calibrated prior to the run. Mice were placed in one of the four plethysmograph chambers and acclimated for 30 mins. Thereafter, respiratory measurements were collected using Buxco FinePointe software over a 15 min period.

Open field animal activity testing

Open field testing enabled us to test ambulation and locomotor function at the experimental endpoint at 8 weeks of age. The test utilises a square, acrylic box (720 mm x 720 mm and 330 mm high) in a light and temperature-controlled room. At 8 weeks of age, animals were placed in the centre of the box and left to freely roam for 10 minutes. The test is captured from above and is automatically analysed via the TopScan LITE System utilising the animals centre of mass to track it over the course of the experiment.

Body composition analysis

At 8 weeks of age, body composition was analysed using the EchoMRI-100H scanner (EchoMRI, Houston, USA) as conducted by us previously (66, 67).

Kyphosis Index

Prior to endpoint, at 8 weeks of age, animals were lightly anaesthetised (2.5% isoflurane) and X-rayed in an IVIS Lumina X5 (PerkinElmer). Images were acquired with Living Image(R) 4.8.0 (64bit) using photograph (medium binning, F/stop 6) and X-ray (high resolution binning, F/stop 6) overlay. Images were imported into Image J and the kyphosis index was calculated from a line drawn between the C7 vertebra to L6 (usually corresponds with the posterior edge of the iliac wing; line AB) divided by a line perpendicular to this from the dorsal edge of the vertebra at the point of greatest curvature (line CD) (68).

Biomarkers

Urine

Before functional testing at 8 weeks of age, mice were temporarily housed in sterile individual holding cages and urine was collected and snap frozen. Oxidative stress biomarker, 8-

isoprostane, was quantitated via enzyme-linked immunoassay (ELISA; Cayman Chemical Company, USA).

Blood

Blood was collected from the endpoint surgery via cardiac puncture into Greiner Bio-One 0.5ml lithium heparin tubes. Plasma was separated via centrifugation (5 mins, 3000G, 4°C). Creatine kinase units were quantitated spectrophotometrically (CK-NAC kit, Randox Laboratories, UK) (69). Cytokines and chemokines were quantitated using a Bio-Plex™ 200 System (Pro Mouse Cytokine 23-plex Assay #M60009RDPD, BioRad, USA) according to the manufacturers manual.

Surgical Procedure

At the experimental endpoint, animals were weighed, treated via gavage, and then deeply anaesthetised (4% induction and 2.5% maintenance isoflurane). FDBs were removed, the left EDL and SOL were tied tendon to tendon with suture thread and excised for *ex vivo* contractile experimentation. Other muscles and organs were excised and snap frozen to be weighed later (Refer to Supplemental Tables 1 and 2).

Ex vivo Contractile Function

After excision, the proximal ends of the EDL and SOL muscles were placed onto a force transducer and the distal end fixed to a micromanipulator in organ baths (Danish Myo Technology, Hinnerup, Denmark) via knotted loops. In depth information regarding the contractile protocol has previously been described by our lab group (70). Outcome measures included specific/absolute force, force frequency, fatigue, and recovery.

Seahorse Extracellular Flux

Excised FDB's were prepared and incubated as previously described by us (71) using a standard mitochondrial stress test on a Seahorse Bioscience XF24 Analyser (Agilent).

Histological analyses of the quadriceps and heart

The right quadriceps and hearts were initially snap frozen then later slow thawed at 4°C in 10% neutral buffered formalin for 48 hours. Following fixation, quadriceps and hearts were transferred to 70% ethanol until paraffin embedding. Sections were cut at 5 μm thick for the quadriceps and heart. Hearts were cut diagonally at the mid-ventricle. Sections were stained and assessed as detailed below. All assessments were performed on blinded images by the same experimenter.

Haematoxylin and Eosin (H&E)

Quadriceps and hearts were stained with H&E (71). For quantification of healthy versus unhealthy tissue in the quadriceps, the H&E colour deconvolution plugin on ImageJ was used. The eosin component (normal, undamaged myofibers) was measured via thresholding (TREAT-NMD SOP: DMD_M.1.2.007) (72). The proportion of centronucleated fibres in the quadriceps was calculated by analysing ~200 fibres per section (73, 74). For active

cardiomyocyte degeneration of the heart, degenerating tissue and clusters of inflammatory infiltrate were measured and expressed relative to the total area (75).

Masson's Trichrome

Masson's trichrome was used to assess collagen/fibrotic connective tissue content in the quadriceps and heart. Analysis of the percent area of fibrosis in both the quadriceps and heart was conducted using the Masson's colour deconvolution plugin on ImageJ (37).

Immunohistochemistry

To quantify macrophage and adipocyte infiltration, anti-CD68 (ab125212) and perilipin-1 (9349S) primary antibodies were utilised at a dilution of 1:500 and 1:200, respectively and stained as described by us previously (19). CD68 positive macrophages were manually counted using DotDotGoose (version 1.7.0) and expressed as number per square millimetre of muscle cross section as measured via ImageJ. Adipocytes were manually traced using ImageJ and expressed as percentage of the total area. Full cross-sectional areas were imaged at 1X using a Zeiss Axio Imager Z2 microscope (Carl Zeiss MicroImaging, GmbH, Germany).

Transcriptomics

RNA extractions, quantification, and integrity

Frozen quadriceps muscles (10 to 15 mg) (*n*=10 per group) were removed from -80C storage where extractions were performed using the AllPrep DNA/RNA/miRNA Universal Kit (Qiagen, Valencia, USA) which includes a genomic DNA elimination step to remove DNA from total RNA. Aliquots were taken for RNA quantification and integrity testing using a Nanodrop spectrophotometer (Thermo Fisher Scientific). 1 μl of sample was placed on the instrument and readings for RNA concentration (ng/μl), the A260/280 ratio and A260/230 ratio were recorded. Samples were diluted to 100 +/- 20ng/μl and RNA integrity was assessed using an automated microcapillary electrophoresis system (Experion, Bio-Rad Laboratories, Hercules, CA). An RNA quality indicator score of greater than 7 was considered intact and of high integrity (76).

RNA sequencing

The library construction and RNA sequencing was performed by Beijing Genomics Institute (Shenzhen, China) and ~1μg total RNA was used for library construction. For library preparation, samples were denatured, and mRNA was enriched using oligo (dT) attached magnetic beads. The mRNA was fragmented into small pieces and the double strand cDNA was synthesised. These double-stranded cDNA fragments were subjected to end-repair and a single 'A' nucleotide is added to the 3' ends of the blunt fragments and subsequently ligated to the adapter. The ligation product was purified and enriched with PCR amplification to yield the final cDNA library. Sufficient quality DNA nanoballs (DNBs) were loaded into patterned nanoarrays using high-intensity DNA nanochip technique and sequenced through combinatorial Probe-Anchor Synthesis (cPAS).

Bioinformatics

Fastq files were processed using the NfCore/RNAseq (v3.10.1) pipeline (77). Reads were aligned to the *Mus musculus* Ensembl GRCm39.release-109 reference genome using STAR aligner (78) and quantified using Salmon (79) producing raw genes count matrix. Various quality control metrics were generated and summarised in a multiQC report (80). Raw counts were then analysed with Degust (81), a web tool that performs normalisation using trimmed mean of M values (TMM) (82), and differential expression analysis using limma/voom (83). Aggravated *mdx* model was expressed relative to WT exercised control to rule out the effects of exercise-related adaptations. Data presented in heatmaps considered significantly different based on a 2.0-fold change threshold and a *P* value of <0.05.

Metabolomics and Lipidomics

Metabolomics was run and analysed at Metabolomics Australia (Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Australia).

Extraction and sample preparation

Tissue optimisation was performed and 30mg of tissue was determined to be optimal biomass of tissue required to provide the best metabolite coverage in samples. Tissues were pre-weighed and extracted by homogenisation using a hand-held homogenising tool by pulsing (6 x 10 seconds) for liquid chromatography–mass spectrometry (LC–MS) analysis by adding 300 μl of ice-cold 3:1 Methanol:Milli-Q water containing MA-Internal Standards. Extracts were vortexed for 10 seconds, mixed on a thermomixer and centrifuged at 4°C for 10 minutes at maximum speed.

Polar metabolite analysis

100 μl of the supernatant was pipetted into a glass HPLC vial containing an insert. A further 20 μl from each sample was pooled to generate a pooled biological quality control sample (pbQC), which was run after every five-biological samples.

Targeted lipid analysis

To the remaining homogenate and lysate, 300 μl of 100% chloroform (containing MA-lipid Internal Standards) was added to create a ratio of 2:1 chloroform:methanol for the remaining homogenate. Extracts were vigorously vortexed to resuspend the tissue pellet. Samples were mixed on a thermomixer and centrifuged at 10°C for 5 mins at 15,000 rpm. The LCMS methodologies are outlined previously (84).

Bioinformatics

Statistical data analysis and data normalisation of metabolomics and lipidomics was performed in MetaboAnalyst 6.0. Data were normalised by median log transformation without data scaling. 2.0-fold change threshold and a raw *P* value of <0.05 considered significant for data presented in heatmaps.

Statistics

Data are presented as mean ± SEM unless otherwise stated. One-way ANOVAs were performed using Prism software (Graphpad, La Jolla, CA) for all analyses unless otherwise stated to assess either the extent of phenotype or *mdx* VEH relative to treatment groups and Tukey's post hoc test was used for multiple comparisons. For mitochondrial oxygen consumption/extracellular flux and contractile force-frequency experiments, a two-way ANOVA with repeated measures was performed using Sidak posthoc testing to interrogate differences between phenotype/treatment and time/frequency. A $P < 0.05$ was considered significant and trends were reported at $P < 0.1$. Outliers were removed if outside ± 2 standard deviations from the mean.

Study Approval
Ethics approval was granted by Victoria University Animal Ethics Committee. All breeding, handling and animal experiments adhered to the Australian Code of Practice for Care and Use of Animals for Scientific Purposes (National Health and Medical Research Council, Australia, 8th edition). At 14 days of age, mice were transferred from breeding program (AEC# 20-005) to an approved experimental project (AEC# 20-006).

Data availability:
All individual data values represented in graphs and are available in the supporting data values file.

Author contributions: CAT, DF, and ER conceptualised and designed the study. SK, CAT, RMB, BQ, BAA, and ER conducted animal studies. SK, CAT, RMB, GS, DD, and ER conducted experiments. SK, CAT, RMB, RB, GS, NK, ALP, TJY, and ER conducted data analysis. CAT, NK, AP, XY, NS, JH, BN, MVP, AAR, DDL, DF and ER contributed to the methodological and technical know-how, resources, and data interpretation. SK, CAT, and ER drafted the manuscript. All authors reviewed and approved the final manuscript.

Acknowledgments: This work was supported by funding from The Muscular Dystrophy Association U.S.A (Ideas grant: MDA871929) and Australian Physiological Society (AuPS; HDR Student Small Grant). ER's lab is supported by AFM Téléthon (France), The Jack Brockhoff Foundation and Duchenne Parent Project Netherlands. SK, CAT, and ER acknowledge outstanding support from Anne Luxford, Tricia Murphy, and Steve Holloway with animal care and breeding (Victoria University Animal Services, WCHRE, Sunshine Hospital). We gratefully acknowledge the Melbourne Histology Platform, The University of Melbourne, for expert service in tissue embedding, cutting, and staining. We gratefully acknowledge Patricia Quianbao from Metabolomics Australia, University of Melbourne, for assistance with metabolomics studies. We gratefully acknowledge Annemieke Aartsma-Rus and Maaike Van Putten, Leiden University Medical Centre, for their advice on the exercise aggravation protocol and critical reading of the manuscript. Schematics were created with BioRender.com and all figures were constructed in Adobe Illustrator.

References:

1. Duan D, Goemans N, Takeda Si, Mercuri E, and Aartsma-Rus A. Duchenne muscular dystrophy. *Nature Reviews Disease Primers.* 2021;7(1):13.
2. Wong SH, McClaren BJ, Archibald AD, Weeks A, Langmaid T, Ryan MM, et al. A mixed methods study of age at diagnosis and diagnostic odyssey for Duchenne muscular dystrophy. *Eur J Hum Genet.* 2015;23(10):1294-300.
3. Wahlgren L, Kroksmark A-K, Lindblad A, Tulinius M, and Sofou K. Respiratory comorbidities and treatments in Duchenne muscular dystrophy: impact on life expectancy and causes of death. *Journal of Neurology.* 2024.
4. Mhandire DZ, Burns DP, Roger AL, O'Halloran KD, and ElMallah MK. Breathing in Duchenne muscular dystrophy: translation to therapy. *J Physiol.* 2022;600(15):3465-82.
5. Passamano L, Taglia A, Palladino A, Viggiano E, D'Ambrosio P, Scutifero M, et al. Improvement of survival in Duchenne Muscular Dystrophy: retrospective analysis of 835 patients. *Acta Myol.* 2012;31(2):121-5.
6. Archer JE, Gardner AC, Roper HP, Chikermane AA, and Tatman AJ. Duchenne muscular dystrophy: the management of scoliosis. *J Spine Surg.* 2016;2(3):185-94.
7. Theo L-G, Hannah L, Rahul S, and Pradeep PAM. In: Gustav M, and Peter M eds. *Cardiomyopathy*. Rijeka: IntechOpen; 2021:Ch. 14.
8. Adorisio R, Mencarelli E, Cantarutti N, Calvieri C, Amato L, Cicenia M, et al. Duchenne Dilated Cardiomyopathy: Cardiac Management from Prevention to Advanced Cardiovascular Therapies. *J Clin Med.* 2020;9(10).
9. Spurney C, Shimizu R, Morgenroth LP, Kolski H, Gordish-Dressman H, and Clemens PR. Cooperative International Neuromuscular Research Group Duchenne Natural History Study demonstrates insufficient diagnosis and treatment of cardiomyopathy in Duchenne muscular dystrophy. *Muscle Nerve.* 2014;50(2):250-6.
10. Szwec S, Kapłucha Z, Chamberlain JS, and Konieczny P. Dystrophin- and Utrophin-Based Therapeutic Approaches for Treatment of Duchenne Muscular Dystrophy: A Comparative Review. *BioDrugs.* 2024;38(1):95-119.
11. Lek A, Wong B, Keeler A, Blackwood M, Ma K, Huang S, et al. Death after High-Dose rAAV9 Gene Therapy in a Patient with Duchenne's Muscular Dystrophy. *New England Journal of Medicine.* 2023;389(13):1203-10.
12. Deng J, Zhang J, Shi K, and Liu Z. Drug development progress in duchenne muscular dystrophy. *Front Pharmacol.* 2022;13:950651.
13. Tulangekar A, and Sztal TE. Inflammation in Duchenne Muscular Dystrophy-Exploring the Role of Neutrophils in Muscle Damage and Regeneration. *Biomedicines.* 2021;9(10).
14. Kourakis S, Timpani CA, Campelj DG, Hafner P, Gueven N, Fischer D, et al. Standard of care versus new-wave corticosteroids in the treatment of Duchenne muscular dystrophy: Can we do better? *Orphanet Journal of Rare Diseases.* 2021;16(1):117.
15. Smith EC, Conklin LS, Hoffman EP, Clemens PR, Mah JK, Finkel RS, et al. Efficacy and safety of vamorolone in Duchenne muscular dystrophy: An 18-month interim analysis of a non-randomized open-label extension study. *PLoS Med.* 2020;17(9):e1003222.
16. Mah JK, Clemens PR, Guglieri M, Smith EC, Finkel RS, Tulinius M, et al. Efficacy and Safety of Vamorolone in Duchenne Muscular Dystrophy: A 30-Month Nonrandomized Controlled Open-Label Extension Trial. *JAMA Netw Open.* 2022;5(1):e2144178.

17. Mercuri E, Vilchez JJ, Boespflug-Tanguy O, Zaidman CM, Mah JK, Goemans N, et al. Safety and efficacy of givinostat in boys with Duchenne muscular dystrophy (EPIDYS): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *The Lancet Neurology.* 2024;23(4):393-403.

18. Kourakis S, Timpani CA, de Haan JB, Gueven N, Fischer D, and Rybalka E. Targeting Nrf2 for the treatment of Duchenne Muscular Dystrophy. *Redox Biol.* 2021;38:101803.

19. Timpani CA, Kourakis S, Debruin DA, Campelj DG, Pompeani N, Dargahi N, et al. Dimethyl fumarate modulates the dystrophic disease program following short-term treatment. *JCI Insight.* 2023;8(21).

20. De Luca A. Use of treadmill and wheel exercise for impact on mdx mice phenotype (SOP DMD_M.2.1.001). https://www.treat-nmd.org/wp-content/uploads/2023/07/MDX-DMD_M.2.1.001.pdf.

21. Aartsma-Rus A, and van Putten M. Assessing functional performance in the mdx mouse model. *J Vis Exp.* 2014(85).

22. Schill KE, Altenberger AR, Lowe J, Periasamy M, Villamena FA, Rafael-Fortney JA, et al. Muscle damage, metabolism, and oxidative stress in mdx mice: Impact of aerobic running. *Muscle Nerve.* 2016;54(1):110-7.

23. Nakae Y, Dorchies OM, Stoward PJ, Zimmermann BF, Ritter C, and Ruegg UT. Quantitative evaluation of the beneficial effects in the mdx mouse of epigallocatechin gallate, an antioxidant polyphenol from green tea. *Histochemistry and Cell Biology.* 2012;137(6):811-27.

24. Fiore PF, Benedetti A, Sandonà M, Madaro L, De Bardi M, Saccone V, et al. Lack of PKCθ Promotes Regenerative Ability of Muscle Stem Cells in Chronic Muscle Injury. *Int J Mol Sci.* 2020;21(3).

25. Lavi E, Cohen A, Libdeh AA, Tsabari R, Zangen D, and Dor T. Growth hormone therapy for children with Duchenne muscular dystrophy and glucocorticoid induced short stature. *Growth Horm IGF Res.* 2023;72-73:101558.

26. Ward LM, and Weber DR. Growth, pubertal development, and skeletal health in boys with Duchenne Muscular Dystrophy. *Curr Opin Endocrinol Diabetes Obes.* 2019;26(1):39-48.

27. Suárez-Calvet X, Fernández-Simón E, Natera D, Jou C, Pinol-Jurado P, Villalobos E, et al. Decoding the transcriptome of Duchenne muscular dystrophy to the single nuclei level reveals clinical-genetic correlations. *Cell Death Dis.* 2023;14(9):596.

28. Wang Q, Reiter RS, Huang QQ, Jin JP, and Lin JJ. Comparative studies on the expression patterns of three troponin T genes during mouse development. *Anat Rec.* 2001;263(1):72-84.

29. Clarkson PM, Devaney JM, Gordish-Dressman H, Thompson PD, Hubal MJ, Urso M, et al. ACTN3 genotype is associated with increases in muscle strength in response to resistance training in women. *Journal of Applied Physiology.* 2005;99(1):154-63.

30. Pappas CT, Krieg PA, and Gregorio CC. Nebulin regulates actin filament lengths by a stabilization mechanism. *Journal of Cell Biology.* 2010;189(5):859-70.

31. Lombardo SD, Basile MS, Ciurleo R, Bramanti A, Arcidiacono A, Mangano K, et al. A Network Medicine Approach for Drug Repurposing in Duchenne Muscular Dystrophy. *Genes (Basel).* 2021;12(4).

32. Hanna S, Xianzhen H, David KC, Ying S, Matthew SA, Merry-Lynn NM, et al. Late-Stage Skeletal Muscle Transcriptome in Duchenne muscular dystrophy shows a BMP4-Induced Molecular Signature. *bioRxiv.* 2024:2024.04.19.590266.

33. Bhargava P, Fitzgerald KC, Venkata SLV, Smith MD, Kornberg MD, Mowry EM, et al. Dimethyl fumarate treatment induces lipid metabolism alterations that are linked to immunological changes. *Ann Clin Transl Neurol.* 2019;6(1):33-45.

34. Willmann R, De Luca A, Benatar M, Grounds M, Dubach J, Raymackers JM, et al. Enhancing translation: guidelines for standard pre-clinical experiments in mdx mice. *Neuromuscul Disord.* 2012;22(1):43-9.

35. Terrill JR, Radley-Crabb HG, Grounds MD, and Arthur PG. N-Acetylcysteine treatment of dystrophic mdx mice results in protein thiol modifications and inhibition of exercise induced myofibre necrosis. *Neuromuscular Disorders.* 2012;22(5):427-34.

36. De Luca A, Nico B, Liantonio A, Didonna MP, Fraysse B, Pierno S, et al. A multidisciplinary evaluation of the effectiveness of cyclosporine a in dystrophic mdx mice. *Am J Pathol.* 2005;166(2):477-89.

37. Morroni J, Schirone L, Vecchio D, Nicoletti C, D'Ambrosio L, Valenti V, et al. Accelerating the Mdx Heart Histo-Pathology through Physical Exercise. *Life.* 2021;11(7):706.

38. SOGC Guideline Retirement Notice No. 2. *J Obstet Gynaecol Can.* 2022;44(10):1104-12.

39. Culbreth M, and Aschner M. GSK-3β, a double-edged sword in Nrf2 regulation: Implications for neurological dysfunction and disease. *F1000Res.* 2018;7:1043.

40. Mao C, Gao M, Zang S-K, Zhu Y, Shen D-D, Chen L-N, et al. Orthosteric and allosteric modulation of human HCAR2 signaling complex. *Nature Communications.* 2023;14(1):7620.

41. Molina T, Fabre P, and Dumont NA. Fibro-adipogenic progenitors in skeletal muscle homeostasis, regeneration and diseases. *Open Biol.* 2021;11(12):210110.

42. Mandal N, Grambergs R, Mondal K, Basu SK, Tahia F, and Dagogo-Jack S. Role of ceramides in the pathogenesis of diabetes mellitus and its complications. *J Diabetes Complications.* 2021;35(2):107734.

43. Miranda ER, and Funai K. Suppression of de novo sphingolipid biosynthesis mitigates sarcopenia. *Nat Aging.* 2022;2(12):1088-9.

44. Laurila PP, Luan P, Wohlwend M, Zanou N, Crisol B, Imamura de Lima T, et al. Inhibition of sphingolipid de novo synthesis counteracts muscular dystrophy. *Sci Adv.* 2022;8(4):eabh4423.

45. Dasgupta S, and Ray SK. Insights into abnormal sphingolipid metabolism in multiple sclerosis: targeting ceramide biosynthesis as a unique therapeutic strategy. *Ther Targets Neurol Dis.* 2017;4.

46. De la Garza-Rodea AS, Moore SA, Zamora-Pineda J, Hoffman EP, Mistry K, Kumar A, et al. Sphingosine Phosphate Lyase Is Upregulated in Duchenne Muscular Dystrophy, and Its Inhibition Early in Life Attenuates Inflammation and Dystrophy in Mdx Mice. *Int J Mol Sci.* 2022;23(14).

47. Naarding KJ, Reyngoudt H, van Zwet EW, Hooijmans MT, Tian C, Rybalsky I, et al. MRI vastus lateralis fat fraction predicts loss of ambulation in Duchenne muscular dystrophy. *Neurology.* 2020;94(13):e1386-e94.

48. Khattri RB, Batra A, Matheny M, Hart C, Henley-Beasley SC, Hammers D, et al. Magnetic resonance quantification of skeletal muscle lipid infiltration in a humanized mouse model of Duchenne muscular dystrophy. *NMR Biomed.* 2023;36(3):e4869.

49. Houwen-van Opstal SLS, Rodwell L, Bot D, Daalmeyer A, Willemsen MAAP, Niks EH, et al. BMI-z scores of boys with Duchenne muscular dystrophy already begin to increase before losing ambulation: a longitudinal exploration of BMI, corticosteroids and caloric intake. *Neuromuscular Disorders.* 2022;32(3):236-44.

50. Billich N, Adams J, Carroll K, Truby H, Evans M, Ryan MM, et al. The Relationship between Obesity and Clinical Outcomes in Young People with Duchenne Muscular Dystrophy. *Nutrients.* 2022;14(16).

51. Grzegorzewska AP, Seta F, Han R, Czajka CA, Makino K, Stawski L, et al. Dimethyl Fumarate ameliorates pulmonary arterial hypertension and lung fibrosis by targeting multiple pathways. *Scientific Reports.* 2017;7(1):41605.

52. Kato K, Papageorgiou I, Shin YJ, Kleinhenz JM, Palumbo S, Hahn S, et al. Lung-Targeted Delivery of Dimethyl Fumarate Promotes the Reversal of Age-Dependent Established Lung Fibrosis. *Antioxidants (Basel).* 2022;11(3).

53. Liu J, Wu Z, Liu Y, Zhan Z, Yang L, Wang C, et al. ROS-responsive liposomes as an inhaled drug delivery nanoplatform for idiopathic pulmonary fibrosis treatment via Nrf2 signaling. *J Nanobiotechnology.* 2022;20(1):213.

54. Oh CJ, Kim J-Y, Choi Y-K, Kim H-J, Jeong J-Y, Bae K-H, et al. Dimethylfumarate Attenuates Renal Fibrosis via NF-E2-Related Factor 2-Mediated Inhibition of Transforming Growth Factor-β/Smad Signaling. *PLOS ONE.* 2012;7(10):e45870.

55. Rybalka E, Kourakis S, Bonsett CA, Moghadaszadeh B, Beggs AH, and Timpani CA. Adenylosuccinic Acid: An Orphan Drug with Untapped Potential. *Pharmaceuticals.* 2023;16(6):822.

56. Timpani CA, Hayes A, and Rybalka E. Revisiting the dystrophin-ATP connection: How half a century of research still implicates mitochondrial dysfunction in Duchenne Muscular Dystrophy aetiology. *Med Hypotheses.* 2015;85(6):1021-33.

57. Bakay M, Zhao P, Chen J, and Hoffman EP. A web-accessible complete transcriptome of normal human and DMD muscle. *Neuromuscul Disord.* 2002;12 Suppl 1:S125-41.

58. Zhang H, Tsui CK, Garcia G, Joe LK, Wu H, Maruichi A, et al. The extracellular matrix integrates mitochondrial homeostasis. *Cell.* 2024.

59. Muire PJ, Mangum LH, and Wenke JC. Time Course of Immune Response and Immunomodulation During Normal and Delayed Healing of Musculoskeletal Wounds. *Front Immunol.* 2020;11:1056.

60. Gloss D, Moxley RT, 3rd, Ashwal S, and Oskoui M. Practice guideline update summary: Corticosteroid treatment of Duchenne muscular dystrophy: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology.* 2016;86(5):465-72.

61. Administration USFaD. In: Services USDoHaH ed. Rockville, MD 2005.

62. Wintzinger M, Miz K, York A, Demonbreun AR, Molkentin JD, McNally EM, et al. Effects of Glucocorticoids in Murine Models of Duchenne and Limb-Girdle Muscular Dystrophy. *Methods Mol Biol.* 2023;2587:467-78.

63. Quattrocelli M, Barefield DY, Warner JL, Vo AH, Hadhazy M, Earley JU, et al. Intermittent glucocorticoid steroid dosing enhances muscle repair without eliciting muscle atrophy. *J Clin Invest.* 2017;127(6):2418-32.

64. Kourakis S, Timpani CA, de Haan JB, Gueven N, Fischer D, and Rybalka E. Dimethyl Fumarate and Its Esters: A Drug with Broad Clinical Utility? *Pharmaceuticals (Basel).* 2020;13(10).

65. Mâncio RD, Hermes TA, Macedo AB, Mizobuti DS, Rupcic IF, and Minatel E. Dystrophic phenotype improvement in the diaphragm muscle of mdx mice by diacerhein. *PLoS One.* 2017;12(8):e0182449.

66. Campelj DG, Debruin DA, Timpani CA, Hayes A, Goodman CA, and Rybalka E. Sodium nitrate co-supplementation does not exacerbate low dose metronomic doxorubicin-induced cachexia in healthy mice. *Scientific Reports.* 2020;10(1):15044.

67. Sorensen JC, Petersen AC, Timpani CA, Campelj DG, Cook J, Trewin AJ, et al. BGP-15 Protects against Oxaliplatin-Induced Skeletal Myopathy and Mitochondrial Reactive Oxygen Species Production in Mice. *Front Pharmacol.* 2017;8:137.

68. Laws N, and Hoey A. Progression of kyphosis in mdx mice. *J Appl Physiol (1985).* 2004;97(5):1970-7.

69. James C, Dugan CW, Boyd C, Fournier PA, and Arthur PG. Temporal tracking of cysteine 34 oxidation of plasma albumin as a biomarker of muscle damage following a bout of eccentric exercise. *Eur J Appl Physiol.* 2024.

70. Debruin DA, Timpani CA, Lalunio H, Rybalka E, Goodman CA, and Hayes A. Exercise May Ameliorate the Detrimental Side Effects of High Vitamin D Supplementation on Muscle Function in Mice. *J Bone Miner Res.* 2020;35(6):1092-106.

71. Timpani CA, Goodman CA, Stathis CG, White JD, Mamchaoui K, Butler-Browne G, et al. Adenylosuccinic acid therapy ameliorates murine Duchenne Muscular Dystrophy. *Sci Rep.* 2020;10(1):1125.

72. Grounds M. Quantification of histopathology in Haemotoxylin and Eosin stained muscle sections (SOP DMD_M.1.2.007). https://www.treat-nmd.org/wp-content/uploads/2023/07/MDX-DMD_M.1.2.007-28.pdf.

73. Stupka N, Plant DR, Schertzer JD, Emerson TM, Bassel-Duby R, Olson EN, et al. Activated calcineurin ameliorates contraction-induced injury to skeletal muscles of mdx dystrophic mice. *J Physiol.* 2006;575(Pt 2):645-56.

74. Stupka N, Schertzer JD, Bassel-Duby R, Olson EN, and Lynch GS. Calcineurin-Aα activation enhances the structure and function of regenerating muscles after myotoxic injury. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology.* 2007;293(2):R686-R94.

75. Morroni J, Schirone L, Valenti V, Zwergel C, Riera CS, Valente S, et al. Inhibition of PKCθ Improves Dystrophic Heart Phenotype and Function in a Novel Model of DMD Cardiomyopathy. *International Journal of Molecular Sciences.* 2022;23(4):2256.

76. Lin W, Saner NJ, Weng X, Caruana NJ, Botella J, Kuang J, et al. The Effect of Sleep Restriction, With or Without Exercise, on Skeletal Muscle Transcriptomic Profiles in Healthy Young Males. *Front Endocrinol (Lausanne).* 2022;13:863224.

77. Harshil Patel PE, Alexander Peltzer, Rickard Hammarén, Olga Botvinnik, Gregor Sturm, Denis Moreno, Pranathi Vemuri, silviamorins, Lorena Pantano, FriederikeHanssen, Maxime U. Garcia, Chris Cheshire, rfenouil, nf-core bot, marchoeppner, Peng Zhou, Gisela Gabernet, Daniel Straub, … Matthias Hörtenhuber.*;* 2021.

78. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.* 2012;29(1):15-21.

79. Patro R, Duggal G, Love MI, Irizarry RA, and Kingsford C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods.* 2017;14(4):417-9.

80. Ewels P, Magnusson M, Lundin S, and Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics.* 2016;32(19):3047-8.

81. David R. Powell AP, Michael Milton. zenodo; 2019.

82. Robinson MD, and Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology.* 2010;11(3):R25.

83. Law CW, Chen Y, Shi W, and Smyth GK. voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology.* 2014;15(2):R29.

84. Lu T, Freytag L, Narayana VK, Moore Z, Oliver SJ, Valkovic A, et al. Matrix Selection for the Visualization of Small Molecules and Lipids in Brain Tumors Using Untargeted MALDI-TOF Mass Spectrometry Imaging. *Metabolites.* 2023;13(11):1139.

85. Gillis JM. The recovery score to evaluate therapy efficiency in NMD: a common, quantitative and comparative scoring system (SOP DMD_M.1.1_001). https://www.treat-nmd.org/wp-content/uploads/2023/07/MDX-DMD_M.1.1_001-21.pdf.

FIGURE LEGENDS

Figure 1. Twice-weekly forced treadmill running aggravates the *mdx* phenotype. (A) Schematic of treatment and treadmill protocols. *Mdx* aggravation was proved using clinically compatible parameters of **(B)** plasma creatine kinase and **(C)** forelimb grip strength. H & E staining assessed the **(D)** unhealthy/healthy tissue ratio (black arrows) and **(E)** centronucleated fibre proportion of the quadriceps in **(F-H)** (orange arrows) and active cardiomyocyte degeneration in **(I)** hearts. Pan macrophage marker, CD68, and Masson's trichrome staining assessed **(J-M)** immune infiltration and **(N-R)** fibrosis of quadriceps and heart. **(S)** Mitochondrial oxygen consumption was measured in FDB fibres and **(T)** metabolic phenotypes in response to chemical uncoupling are shown. Extent of DMD phenotype was calculated (*mdx* AGG/*mdx* SED*100) to indicate extent of disease aggravation (34). Data in **B-E, I-J**, **N** and **R** are presented as mean ± SEM and *n* are indicated by individual data points. Panel **S** *n*: WT SED = 7, *mdx* SED = 12, *mdx* AGG = 6. Panel **T** *n*: WT = 9, *mdx* SED = 11, *mdx* AGG = 8. Statistical analysis used one-way ANOVA: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$. **F-H**, **K-M** and **O-Q** scale bar = 50μm.

Figure 2. DMF treatment improves grip strength and modulates transcriptional control of myofibril assembly. (A) Clinically compatible biomarker creatine kinase and functional parameters including **(B)** maximal forelimb grip strength, **(C)** minimal holding impulse, **(D)** rotarod time to fall and **(E)** distance travelled in the open field were assessed at the experimental endpoint. Anthropometric measures and their associated transcriptional pathways in muscle were also assessed: **(F)** Fat mass index and **(G)** the fatty acid metabolism pathway, **(H)** lean mass index, **(I)** the BCAA catabolism pathway; and **(J)** hydration index. Absolute force was measured *ex vivo* in **(K)** EDL and **(L)** SOL and specific force was calculated **(M-N)**. The transcriptomics dataset was probed for **(O)** muscle contractile associated genes. Data in **E, G** and **O** are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Data in **A-D**, **F**, and **H-N** are presented as mean ± SEM and *n* are indicated by individual data points. Statistical significance was tested via one-way ANOVA: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

Figure 3. DMF reduces macrophage infiltration of quadriceps. The **(A)** unhealthy to healthy tissue ratio and **(B)** percentage of centronucleated fibres are shown with **(C-F; yellow** arrows indicate centronucleated fibres; black arrows indicate infiltrate**). (G)** Transcriptomic signature of myogenesis. **(H)** Pan macrophage marker CD68 was immunohistochemically assessed and **(I-L)** representative images show CD68+ macrophages (indicated by black arrows). **(M)** The transcriptomic dataset was probed for inflammatory markers commonly upregulated in DMD. Data in heatmaps (**G** and **N**) are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Data in **A-B** and **H** are presented as mean ± SEM and *n* are indicated by individual data points. Statistical significance was tested via one-way ANOVA: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$. **C-F** and **I-L** scale bar = 50μm.

Figure 4. DMF reduces fibrosis of quadriceps. Expression of the proposed **(A)** fibrosis-associated DMD seed genes (31) and **(B)** DMD molecular signature (32) was probed using our transcriptomic dataset. **(C)** Quadriceps fibrosis was quantified using Masson's trichrome staining and **(D-G)** representative images are shown. Transcriptional pathways related to **(H)** collagen formation, **(I)** ECM degradation and **(J)** ECM organisation were analysed to investigate mechanisms of anti-fibrosis. Data in all heatmaps **(A-B** and **H-J)** are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Data in **C** are presented as mean ± SEM and *n* are indicated by individual data

points. Statistical significance was tested via one-way ANOVA: *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001. **D-G** scale bar = 50μm.

Figure 5. DMF decreases muscle lipid content through modulation of adipogenesis and sphingolipid metabolism pathways in quadriceps. (A-E) Percentage of perilipin-1 positive lipid droplets in quadriceps. Lipid associated transcriptional pathways were probed including **(F)** FAPs, **(G)** TNFR mediated ceramide production, which was a significantly dysregulated Reactome pathway in *mdx* AGG relative to WT quadriceps, **(H)** sphingolipid metabolism and **(I)** adipogenesis. **(J)** Targeted lipidomics was performed to reveal the relative abundance of muscle lipid species. Data in all heatmaps **(F-J)** are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Data in **A** is presented as mean ± SEM and *n* are indicated by individual data points. Statistical significance was tested via one-way ANOVA: *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001. **B-E** Scale bar = 50μm.

Figure 6. DMF treatment tempers expression of detoxification enzymes and transcribes a highly specific anti-inflammatory signature in muscle. (A) DMFs MOA involves dual anti-inflammatory and -oxidative function through transcription of Nrf2. The muscle transcriptome was probed for molecular pathways involved in DMF's MOA: **(B)** Keap1-Nrf2, **(C)** Nrf2 regulation of antioxidant/detoxification enzymes and **(D)** DNA damage reversal pathways. DMF's anti-inflammatory capacity was investigated through **(E)** interleukin and **(F)** TNF signalling pathway analysis, and **(G)** plasma cytokines, chemokines and growth factors, and urinary oxidative stress marker **(H)** 8-isoprostane. All heatmap data **(B-G)** are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Data in **H** are presented as mean ± SEM and *n* are indicated by individual data points. Statistical significance was tested via one-way ANOVA: *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001.

Figure 7. DMF reduces purine degradation and mitobiogenic stimuli. (A) Mitochondrial oxygen consumption was measured in FDB fibres and **(B)** metabolic phenotypes in response to chemical uncoupling are shown. Untargeted polar metabolomics was performed on quadriceps muscle and **(C)** mitochondrial, **(D)** PNC, **(E)** purine degradation, **(F)** glycolysis, **(G)** fatty acid metabolism, and **(H)** charge metabolomes are presented. **(I)** Top 5 dysregulated Reactome pathways of the transcriptomics dataset. Reactome pathways associated with **(J)** mitochondrial biogenesis, **(K)** TCA cycle, **(L)** UCP and **(M)** PNC. Data in all heatmaps **C-H** and **J-M** are based on log2 fold change from WT for *mdx* VEH and *mdx* VEH for treatment groups (DMF, PRED and DMF+PRED). Panel **A** *n*: *mdx* VEH = 6, *mdx* DMF = 12, *mdx* PRED = 12, *mdx* DMF+PRED = 12. Panel **B** *n*: *mdx* VEH = 8, *mdx* DMF = 12, *mdx* PRED = 12, *mdx* DMF+PRED = 12. Statistical significance was tested via one-way ANOVA: *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001.

Figure 8. DMF reduces cardiac fibrosis. (A-E) Degenerating and **(F-J)** fibrotic area of the heart was assessed via H&E and Masson's trichrome staining, respectively. **(K)** Minute ventilation, **(L)** frequency and **(M)** tidal volume were measured via plethysmography. **(N)** Data in **A, F** and **K-N** are mean ± SEM and *n* are indicated by individual data points. **(N-R)** The kyphosis index, a measure of thoracolumbar deformity, was calculated. Statistical significance was tested via one-way ANOVA: *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001. **B-E** and **G-J** scale bar = 50μm.

Figure 9. The impact of the treatments on the _mdx_ mouse. The recovery score is used to express the effect of a treatment, not just by the difference between treated and untreated _mdx_ mice, but relative to the extent of the deficiency between WT and _mdx_ mice. A recovery score of 100% indicates the parameter is equal to that of the WT while a score of 0% indicates no improvement has been made. This was calculated in accordance with TREAT-NMD SOP (DMD_M.1.1_001) (85) as follows: (_mdx_ treated)-(_mdx_ untreated)/(WT)-(_mdx_ untreated) x 100.

TABLES

Table 1. Effect of DMF, PRED and DMF+PRED treatment on significantly dysregulated pathways in _mdx_ vs WT. *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001.

	mdx vs WT	DMF vs _mdx_	PRED vs _mdx_	DMF+PRED vs _mdx_
Pathway	Direction/False Discovery Rate (FDR)			
Mitochondrial translation termination	↑ ****	↓ ****	↓ ****	↓ ****
Mitochondrial translation elongation	↑ ****	↓ ****	↓ ****	↓ ****
Mitochondrial translation initiation	↑ ****	↓ ****	↓ ****	↓ ****
Mitochondrial translation	↑ ***	↓ ****	↓ ****	↓ ****
TNRF1-mediated ceramide production	↑ **	↓ ***	-	-
Selenocysteine synthesis	↑ *	↓ **	-	-
Eukaryotic translation elongation	↑ *	↓ **	-	-
Defective ABCB4 causes PFIC3, ICP3 and GBDP1	↓ *	-	-	-
Eukaryotic translation termination	↑ *	↓ **	-	-
Peptide chain elongation	↑ *	↓ **	-	-
SRP-dependent co-translational protein targeting to membrane	↑ *	↓ *	-	-
Viral mRNA translation	↑ *	↓ *	-	-

A

	mdx/ WT	*mdx* DMF/ VEH	*mdx* PRED/ VEH	*mdx* D+P/ VEH	
	****	*			Timp1
	**		**	*	Spp1
	**		**		Fn1
					Mmp2
			*		Igf1

DMD-Associated Seed Genes
(Lombardo et al., 2021)

B

	mdx/ WT	*mdx* DMF/ VEH	*mdx* PRED/ VEH	*mdx* D+P/ VEH	
	*				Serping1
			*		BMP4
					Adamts3

DMD Molecular Signature
(Proposed by Sothers et al., 2024)

C

D *mdx* VEH

E *mdx* DMF

F *mdx* PRED

G *mdx* DMF+PRED

- - - Equivalent to wild-type

H

Collagen Formation

	mdx/ WT	*mdx* DMF/ VEH	*mdx* PRED/ VEH	*mdx* D+P/ VEH	
	****				Col8a1
	***				Col26a1
	**		*		Col14a1
	****		**		Col3a1
	**		**		Col12a1
	**	**			Col23a1
	**		**		Col16a1
	****		**		Lox
	**		**		Adamts14
	***		**		Col5a2
	***		***		Loxl3
	**		***		Plod2
	*		**		Col1a1
	*		**		Col1a2
	**		**		Col6a3
	*		*		Col5a1
	**		*		Bmp1
	*		**		Pcolce
	***				Plec

I

ECM Degradation

	mdx/ WT	*mdx* DMF/ VEH	*mdx* PRED/ VEH	*mdx* D+P/ VEH	
	**		***	**	Mmp12
	****		***	*	Adam8
	****		**	*	Ctss
	****				Mmp14
	****		**		Mmp19
	****		**	*	Ctsk
	****		**	**	Adamts5
	**				Mmp24
	****	*			Casp3
	****		*	*	Ctsl
	**	**			Mmp11
	**		*		Ctsb
	**				Capns1
			**		Mmp9
	****	****			Capn3

J

ECM Organisation

	mdx/ WT	*mdx* DMF/ VEH	*mdx* PRED/ VEH	*mdx* D+P/ VEH	
	***		**		Adam12
	***		**		Tnc
	***		**		Mfap4
	****		*		Tgfb1
	****				Itga5
	**		*		Itga10
	**		**		Adamts14
	**		**	*	Bgn
	***		*		Vcam1
	*				Itgb7
	**				Mfap5
	*		*		Emilin1
	**				Icam1
	**				Pxdn
	***		**		Efemp2
	**		*	***	Itgb6
	*	*		***	Ntn4
	****	*		**	Trappc4
					Itgb1
	*				Tgfb2
	*		*	**	Itga9

Log2 Fold Change (scale from 6 red to -6 blue)

Equivalent to WT (red dashed line)

A — Active Cardiomyocyte Degeneration (%); *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; *p*=0.0718; *; *; **

B — *mdx* VEH

C — *mdx* DMF

D — *mdx* PRED

E — *mdx* DMF+PRED

F — Fibrosis (%); *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; **

G — *mdx* VEH

H — *mdx* DMF

I — *mdx* PRED

J — *mdx* DMF+PRED

K — Minute Ventilation (ml/min); *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; *

L — Frequency (breaths/min); *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; *

M — Tidal Volume (ml); *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; *p*=0.0701; *

N — Kyphosis Index; *mdx* VEH, *mdx* DMF, *mdx* PRED, *mdx* D+P; *

O — *mdx* VEH

P — *mdx* DMF

Q — *mdx* PRED

R — *mdx* DMF+PRED