Title: Dominance hierarchies are linear but shallow in the social amoeba *Dictyostelium discoideum.*

Cathleen M.E. Broersma*[1], Elizabeth A. Ostrowski[2]

[1]Institute of Biology, Leiden University, Leiden, Netherlands

[2]Liggins Institute, University of Auckland, Auckland, New Zealand

*Corresponding author: cathleen.broersma@gmail.com

Abstract

Social groups often form dominance hierarchies, and these hierarchies are almost always linear. However, why

linear dominance hierarchies emerge is not well understood. In the social amoeba *Dictyostelium discoideum*,

cells form a multicellular fruiting body when starved, which consists of a ball of viable spores held aloft by a

stalk of dead cells. In genetically mixed ('chimeric') fruiting bodies, conflicts can arise over the equitable

sacrifice of cells to the dead stalk, and some strains predictably dominate others in the spores. Using pairwise

mixes of strains that co-occurred in small soil samples, we determined the dominance hierarchies in four natural

populations of *Dictyostelium*. These hierarchies were significantly linear in two of four populations, but also

extremely shallow, indicating that co-occurring strains are competitively similar. We used quantitative genetic

analyses to assess the causes of social dominance. Each strain's solo spore production was a significant

predictor of its performance in pairs. However, we detected additional genetic contributions of both the focal

and partner strain, indicating additional cryptic traits that mediate social competitiveness. In contrast to earlier

studies showing strong fitness differences among strains collected over a larger spatial scale, we show that co-

occurring strains are remarkably competitively equivalent, resulting in linear yet shallow hierarchies. Our results

underscore the importance of biologically relevant spatial scales in assessing fitness interactions among

microbes.

Keywords: dominance hierarchy, social amoeba, social evolution, linearity

Introduction

Social interactions within a group frequently result in the formation of a dominance hierarchy, commonly

referred to as a "pecking order" (Schjelderup-Ebbe 1922). These social structures emerge from repeated

antagonistic interactions between individuals that result in the establishment of a clear dominant and subordinate

individual (Drews 1993). For example, fish and rodents establish dominance through aggressive behaviors that

involve chasing and biting, whereas crustaceans exhibit claw-grasping (Chase et al. 2002). Based on the

outcomes of the individual dominance relationships, a ranking of all individuals in the group is established. An

individual's rank can significantly impact their fitness, with higher-ranked individuals obtaining greater access

to limited resources and reproductive opportunities compared to lower-ranked individuals (Wilson 1975;

Clutton-Brock, Albon, and Guinness 1984; Nelissen 1992; Goessmann, Hemelrijk, and Huber 2000).

The structure of a dominance hierarchy can be characterized by two properties: its linearity and steepness (de

Vries, Stevens, and Vervaecke 2006). These properties are assessed by analyzing the collective of pairwise, or

'dyadic', interactions within the group (Drews 1993; de Vries 1995; de Vries, Stevens, and Vervaecke 2006). In

a perfectly linear hierarchy, one individual dominates all others, the second dominates all but the most

dominant, and so on. For example, in a triad of individuals A, B, and C, the hierarchy is linear if the

relationships are transitive, e.g., if A>B, B>C, and A>C, where ">" indicates dominance (Figure 1A). In

contrast, a hierarchy is increasingly non-linear when one or more intransitive triads exist, e.g., if A>B, B>C, but

C>A. The steepness of a dominance hierarchy in turn reflects the differences in dominance ability between

adjacently ranked individuals, with a hierarchy being shallow when the differences are small and steep when the

differences are large (Figure 1B) (Vehrencamp 1983; Barrett et al. 1999; de Vries, Stevens, and Vervaecke

2006).

Figure 1. Two properties of a dominance hierarchy. (A) Linearity. In a perfectly linear dominance hierarchy, all triads are
transitive (e.g., A>B, B>C, and A>C; left side). Alternatively, in a less linear dominance hierarchy, one or more triads are
nontransitive (e.g., A>B, B>C, but C>A; right side). (B) Steepness. A dominance hierarchy is steep if the differences
between adjacently ranked individuals are large and outcomes are highly repeatable (left). A hierarchy is flat if the
differences between adjacently ranked individuals are small and/or less repeatable (right).

The majority of dominance hierarchies examined so far have been linear (Hausfater, Altmann, and Altmann

1982; Barkan et al. 1986; Heinze 1990; LeBrun 2005; Valderrabano-Ibarra, Brumon, and Drummond 2007;

Wittemyer and Getz 2007; Correa et al. 2013; Vullioud et al. 2019). One major hypothesis to explain this

linearity proposes that differences in individuals' pre-existing attributes predict their success in social contests

(Chase et al. 2002; Chase and Seitz 2011). Consequently, the hierarchy can be predicted before it has formed,

and this scenario is therefore referred to as the 'prior attributes' hypothesis (Jackson and Winnegrad 1988;

Drews 1993; Chase et al. 2002). These prior attributes may relate to physical (size and weight), personality

(aggressiveness), genetic (genotype), social (maternal rank), and physiological (hormone levels) characteristics

(Jackson and Winnegrad 1988; Drews 1993; Nakano and Furukawa-Tanaka 1994; Beaugrand and Cotnoir 1996;

Goessmann, Hemelrijk, and Huber 2000; Sundström et al. 2004; Schjolden, Stoskhus, and Winberg 2005; Chase

and Seitz 2011). However, identifying which dominance attributes are of significant influence on dominance

and quantifying their relative importance for the establishment of a hierarchy and its linearity remains

challenging (Appleby 1983; de Vries 1995; Chase and Seitz 2011; Shizuka and McDonald 2012; Schmid and de

Vries 2013).

The social amoeba *Dictyostelium discoideum* serves as an effective model system for investigating social

dominance and dominance hierarchies. In response to starvation, thousands of unicellular amoebae aggregate to

form a multicellular fruiting body, which consists of a stalk of nonviable cells that lifts and supports a sorus of

viable spores (Huss 1989; Kessin 2001). The self-sacrifice of the stalk cells is considered to be an altruistic act,

as it likely benefits the survival and dispersal of the spores (Kuzdzal-Fick et al. 2007; Smith, Queller, and

Strassmann 2014). Importantly, when genetically different individuals aggregate to form a single "chimeric"

fruiting body, some strains dominate by preferentially forming spores and relying on the stalk formation by their

social partner, such that spore production is unfair (Filosa 1962; Hamilton 1964a, 1964b; Buss 1982;

Strassmann, Zhu, and Queller 2000). Consequently, chimeric fruiting body formation can involve a dominance

relationship with a dominant and subordinate strain, offering a basis for constructing a dominance hierarchy

among a group of co-occurring strains (Fortunato, Queller, and Strassmann 2003).

In two prior studies that examined the dominance hierarchy within a group of natural strains of *D. discoideum*,

both revealed linearity (Fortunato, Queller, and Strassmann 2003; Buttery et al. 2009). In support of the prior-

attributes hypothesis, Buttery *et al.* (2009) showed that pre-existing differences in spore production between

strains during clonal development could predict social dominance and linearity in the group. However, both studies used the same seven strains, thus establishing a linear dominance hierarchy only once. They also found a slightly different order of strains, suggesting that the dominance hierarchies may not be repeatable. Moreover, these strains were collected within a 1 km^2 area at a single site in North Carolina. Owing to the large spatial scale over which these strains were sampled, it is unknown whether the strains interact in nature and, therefore, whether the observed dominance hierarchy is ecologically relevant. Indeed, strains collected over more relevant spatial scales might be expected to exhibit lower dominance levels and less linear hierarchies. In other words, co-occurring strains might co-occur *because* they exhibit non-transitive relationships. Thus, assessing the linearity of dominance hierarchies over more realistic spatial scales is a priority.

Here we examined the dominance relationships among strains in four natural populations of *D. discoideum*. The strains that make up a population were isolated from soil taken from the surface of a 10-by-10 cm plot. Thus, the maximum distance between strains at the time of collection was 14 cm (i.e., the diagonal of the square plot). Interactions between strains at this distance are reasonable, given that *D. discoideum* slugs routinely travel multiple centimeters from their aggregation sites (e.g., see Jack et al. (2015)), and moderate gene flow occurs over distances <0.5 km (Kuzdzal-Fick, unpublished). From the observed dominance relationships, we constructed and evaluated the linearity and steepness of the dominance hierarchy. For those populations that did show significant linearity, we asked if pre-existing differences in dominance ability, assessed through the clonal spore production, could predict the dominance outcome and a strain's rank in the hierarchy. Additionally, we performed a quantitative genetic analysis to estimate the genetic basis of variation in social dominance. This analysis examined a focal strain's success as a function of its own genes ('direct genetic effects', or DGEs), those of its partner ('indirect genetic effects', or IGEs), and their interaction (genotype x genotype (G×G) epistasis). IGEs and G×G epistasis are collectively referred to as the social environment, and existing studies have demonstrated that these factors are widespread and can significantly alter evolutionary trait dynamics compared to expectations based solely on DGEs (Moore, Brodie, and Wolf 1997; Wolf et al. 1998; Bijma and Wade 2008).

Of the four natural populations of *D. discoideum* we tested, two exhibited a significantly linear dominance hierarchy. Conversely, the other two populations did not show significant linearity, with one being marginally nonlinear. Notably, in the cases where linearity was observed, the dominance hierarchies were shallow,

indicating minimal fitness differences among co-existing strains, at least in terms of their ability to dominate in

the spores in a chimera. The lack of linearity in two populations, combined with shallow dominance hierarchies

in all populations indicates co-occurring strains that are well-matched for social fitness. Lastly, the quantitative

genetic analyses showed the importance of genetic identity and traits of the focal individual but also that of its

social partner on the level of social dominance of the focal individual. These results demonstrate the importance

of the social genetic environment and demonstrate that linearity can emerge even when social dynamics are

present.

Materials and Methods

Strain collection and selection

The strains of *D. discoideum* used in this study were acquired between 2016 and 2019 by staff and students of

the Ostrowski laboratory. Soil sampling and strain isolation methods are described in detail by Kuzdzal-Fick *et*

al. (2023). The strains and sampling sites are listed in Table S1 and shown in Figure 2A.

To ensure that the chosen strains were not clones of one another, we used Illumina genome sequencing to

determine the pairwise genetic distances among the strains from each site. Illumina sequence data was provided

by Kuzdzal-Fick, and detailed information and methods concerning this data are described in Kuzdzal-Fick *et*

al. (2023). We chose strains with a Euclidean genetic distance of >5 (or 1,756 SNPs), as our analyses indicated

that values below this cut-off are within the margin of error for clones for this sequencing methodology. Figure

2B shows a genetic distance-based tree based on the Euclidean distance.

Experimental design

We examined all possible pairwise combinations of strains for each of the four sites. For three sites with seven

strains each —MLBS, GSNP, and AFCT— this resulted in 21 mixes. For site SMFS, with six strains, this

resulted in 15 mixes. The lower number of strains in site SMFS was a result of having excluded strain EO631

because it did not progress beyond the slug stage during the experiments. We tested all combinations of strains

from GSNP and SMFS on the same day, and from MLBS and AFCT on a different day. We performed three

temporally independent blocks, each initiated from frozen spore stocks.

Figure 2. (A) The 27 strains of *D. discoideum* in this study were sampled from four sites across the US. (B) A distance-based tree showing the genetic similarity among the strains based on genome-wide SNPs.

Cultivation, cell staining, and developmental assays

At the start of each block, we inoculated the relevant strains from the freezer onto SM-agar plates (Formedium Ltd, 2% agar) with *Klebsiella pneumoniae* as a food source. After fruiting bodies had formed, we collected and replated $5x10^5$ spores with *K. pneumoniae* on fresh SM-agar plates for germination and population expansion. After approximately 40 hours, we harvested the cells during the mid-exponential growth phase and washed them three times via differential centrifugation (450 x *g* for 3 min) in KK2 buffer (14.0 mM K_2HPO_4 and 3.4 mM KH_2PO_4, pH 6.4) to remove the bacteria. After washing, the cells were resuspended to a density of $1x10^7$ cells/ml in KK2.

To distinguish the two strains in a mix, we treated one strain in each pair with the fluorescent dye CellHunt Green CMFDA (Molecular Probes), diluted in DMSO according to the manufacturer's specifications. We added the cell dye at a concentration of 20 µM, incubated the cells on a shaking platform in the dark for 30 minutes, washed them once in KK2, and incubated them again for 30 minutes to allow the efflux of excess dye. We then washed the cells twice in cold KK2 and resuspended them at a concentration of $1x10^8$ cells/mL in cold KK2. We treated the unlabeled strain with DMSO only, and these samples underwent the same treatment as the labeled strains.

For each mix, we combined equal volumes of the labeled and unlabeled cells and deposited a 50 µl aliquot of

the mix in a 3-by-3 square (an area of 1 cm^2) of a 47 mm gridded 0.45 µm nitrocellulose filter, resulting in a

total cell number of 5x10^6 cells (Figure 3). We placed the nitrocellulose filter in a 6-cm Petri dish on top of a

Pall filter moistened with 1.5 ml of PDF (per litre: 1.5 g KCl, 1.07 g MgCl·6H$_2$O, 1.8 g KH$_2$PO$_4$, 1.6 g K$_2$HPO$_4$,

0.5 g streptomycin sulphate). We transferred the Petri dishes to a sealed plastic box that contained wet tissues

and placed the box in the dark for 48 hours at 22°C to allow fruiting bodies to develop.

Following the development, we collected the spores in 3 mL of detergent (KK2 buffer + 0.1% IGEPAL and 10

mM EDTA), which dissolved any remaining cells. We measured the fraction of both strains in the spores on a

BD FACSCanto II flow cytometer (488 nm laser, 513/15 GFP filter). We used an automated cell counter (Cell

Countess II FL, Thermo Fisher) to quantify the total number of spores. For each mix, we developed the labeled

and unlabeled cells of each strain by themselves. We also developed mixes that consisted of labeled and

unlabeled cells of the same strain, which we refer to as the 'control mix' (Figure 3). All three types of clonal

controls (100% labeled, 100% unlabeled and the 50-50% control mix) were otherwise treated identically to the

chimeric mixes.

Grow up strains ⟶ Label strains ⟶ Prepare mixes ⟶ Allow development ⟶ Harvest all spores

Figure 3. Overview of the mix experiments to assess social dominance. For each mix, the cells of a labeled and unlabeled
strain were combined at equal cell numbers, spotted on a nitrocellulose filter, and allowed to develop fruiting bodies. After
development, the spores were harvested, and the fractions of the labeled and unlabeled strains were determined using a flow
cytometer. For each mix, the labeled and unlabeled cells of both strains were developed by themselves, as well as in a
'control mix' consisting of labeled and unlabeled cells of the same strain.

 Determination of the winner and loser strain and quantification of social dominance

For each mix, we determined the 'winner' and 'loser' strain based on their average fraction in the spores (across

N=3 blocks). Since the strains were combined at equal starting cell ratios, we identified the winner as the strain

with >50% of the spores, and the loser as the strain with <50% of the spores. We quantified the winner's extent

of social dominance as the magnitude of spore inequity: the increase in the strain's representation in the spores

compared to its initial representation in the cells. This measure ranges from 0 to 0.5, where a value of near 0

indicates roughly equal fractions of the winner and loser in the spores, and a value of 0.5 indicates that the

winner makes up all the spores. For example, a social dominance value of 0.1 indicates that the winner makes

up 60% of the spores in the chimera (i.e., increased by 10% relative to its starting frequency).

Quantification of the linearity of the dominance hierarchy

We calculated the linearity of the dominance hierarchy using standard methods (Appleby 1983). Briefly, we

constructed a win/loss matrix of N-by-N individuals, with the order of the individuals initially being arbitrary.

We assigned an entry in the matrix a value of 0 if the row individual was the 'loser' when mixed with the

column individual (<50% of the spores), and a value of 1 if the row individual was the 'winner' when mixed

with the column individual (>50% of the spores). Using the row sums (S_i) derived from the matrix, we

calculated the observed number of transitive triads (d) using the formula:

207
$$d = \frac{N(N-1)(2N-1)}{12} - \frac{1}{2} \times \sum (S_i)^2$$

209 We calculated the degree of linearity (K) using the formulas:

210
$$K = 1 - \frac{24d}{N^3 - N} \text{ or}$$

212
$$K = 1 - \frac{24d}{N^3 - 4N}$$

214 for even or uneven values of N, respectively. To assess the significance of K, we determined the degrees of

215 freedom (df) and chi-square (χ^2), since the distribution of d approaches that of a chi-square distribution with

216 increasing N. The formulas for these calculations are:

217
$$df = \frac{N(N-1)(N-2)}{(N-4)^2}$$

218 and

219
$$\chi^2 = \frac{8}{N-4}\left[\frac{N(N-1)(N-2)}{24} - d + \frac{1}{2}\right] + df$$

In the given formula, when the group size is $N=6$, the df equals 30, and when $N=7$, the df equals 23.3.

According to a chi-square reference table, K was considered significant at $P<0.05$ if $\chi^2>43.8$ when $df=30$ and if

$\chi^2>35.2$ when $df=23.3$.

In response to observing a very low estimate of steepness (discussed in the Results), we also calculated the

triangle transitivity using the 'transitivity' function from the 'EloRating' package. Briefly, this function

calculates the proportion of transitive triangles (P_t) using the formula:

228
$$P_t = \frac{N_{transitive}}{N_{transitive} + N_{cyclic}}$$

229 in which $N_{transitive}$ is the number of transitive triads in the group, and N_{cyclic} is the number of intransitive triads

230 (Shizuka and McDonald 2012). The triangle transitivity (t_{tri}) was then calculated as:

231
$$t_{tri} = 4(P_t - 0.75)$$

t_{tri} is scaled to run from 0 to 1, where 0 indicates the random expectation (0.75; regardless of the N) and 1

indicates maximum transitivity. Statistical significance of t_{tri} was assessed via randomization. A $P<0.025$ or

$P>0.975$ means that the hierarchy is significantly more or less linear than expected by chance, respectively,

based on 1,000 randomizations that were used to determine the null-distribution.

Quantification of the hierarchy steepness

The hierarchy steepness quantifies the absolute differences in dominance ability between consecutively ranked

individuals in a hierarchy. Steepness therefore also serves as a measure of ranking certainty. To calculate

hierarchy steepness, we used the 'steepness' function from the 'EloRating' package (Neumann et al. 2011). In

contrast to the binary win/loss matrix that we used to calculate the linearity, steepness was calculated using

continuous data—specifically, the fraction of a strain in the spores. Steepness values range from 0 to 1, where 0

indicates a shallow hierarchy with small differences between adjacent ranks, and 1 indicates a steep hierarchy

with large differences between adjacent ranks, and higher-ranked individuals consistently win against adjacent

lower-ranked individuals (Figure 1B). The null hypothesis is that the observed steepness is no greater than a

steepness based on random win probabilities (de Vries, Stevens, and Vervaecke 2006). To test this,

randomization is used. Here the statistical significance (right-tailed P-value) of the observed steepness is

determined by calculating the proportion of times that the steepness generated randomly under the null

hypothesis is greater than or equal to the observed steepness (based on N=1000 iterations).

Determining the ranking of strains in a hierarchy

For those populations that demonstrated significant linearity, we obtained the strain ranking using the 'isi13'

function from the 'compete' package (Curley 2016). This function uses randomization to identify the optimal

order of strains in a linear hierarchy (based on N=1000 iterations). The resulting linear order is that with the

fewest inconsistencies (I) and the smallest inconsistencies (SI) among all potential orders. An inconsistency

describes the situation where a lower-ranked individual dominates a higher-ranked individual in the assumed

linear order (Schmid and de Vries 2013).

Do strains vary in their prior attribute clonal spore production?

To assess if strains within a site vary in their inherent ability to dominate, we used a generalized linear mixed

model (GLMM) that examined the clonal spore production as a function of strain ID and block. Because the

response variable was a count, we used a Poisson error structure with a log link function. We included both

terms as random effects and assessed their significance through single deletions of terms, comparing the reduced

models with the full model using a likelihood ratio test.

Does linearity arise from differences between strains in their ability to produce spores?

To test the hypothesis that linearity arises from a pre-existing hierarchy based on differences between

individuals in their inherent ability to sporulate, we first ranked strains within each site based on their clonal

spore production. The strain that produced the most spores clonally was ranked first, and so on. We then

compared this 'prior attribute ranking' with the ranking based on the observed dominance relationships

determined from the mixes above. The expectation, under the prior attribute hypothesis, was that the rankings in

the two hierarchies would align.

 Quantitative genetic effects analysis

We performed two GLMMs to investigate the effects of clonal spore production, the genotypes of the focal and

partner strain, and the interaction between them, on the focal strain's relative (fraction; Model 1) and absolute

(total; Model 2) number of spores in a mix.

In Model 1, we used beta regression since the response variable was a fraction between 0 and 1. Given the

paired data (with fractions summing to 1), we randomly selected one strain from each pair as the focal strain and

ran the model on this subset. We modelled the *clonal spore production* of the focal and partner strains and

assessed their significance through type II Wald chi-square tests. We modelled the *genotype* of the focal strain,

the partner strain, and their interaction as random effects. In addition, we included block as a random effect. We

assessed the significance of the random effects through single deletions of terms, comparing the reduced models

and the full model using a likelihood ratio test. Finding a significant effect of the focal genotype would indicate

the presence of direct genetic effects (DGEs), a significant effect of partner genotype would indicate the

presence of indirect genetic effects (IGEs), and a significant effect of the interaction between the focal and

partner genotype would indicate the presence of genotype-by-genotype (G×G) interactions, or epistasis. To

quantify the proportional contributions of the DGE, IGE and G×G interaction terms, we divided the variance

components of each random term by the sum of the variance components of all random terms.

Model 2 incorporated several modifications compared to Model 1. Specifically, as the response variable in

Model 2 was a count (i.e., the number of spores of the focal strain) and showed overdispersion, we used a

negative binomial error structure (family=nbinom2). In addition, we included genetic distance between two

strains in a chimera as a covariate and we used the full data set because the total spore production of both strains

in a pair was independent. We analyzed all other terms and performed tests of significance as described in

Model 1.

In summary, with these models, we aimed to *i.* assess if the prior attribute 'spore production during clonal

development' serves as a significant predictor of a strain's relative (Model 1) and absolute (Model 2) dominance

over spore production during chimeric development, and *ii.* quantify the relative importance of direct genetic

effects (DGEs), indirect genetic effects (IGEs) and genotype-by-genotype (G×G) interactions on the focal

strain's spore production during chimeric development.

Results

Dominance hierarchies are mostly linear but shallow

Tables 1-4 show the dominance matrices for the sites AFCT, GSNP, MLBS, and SMFS, respectively. Each matrix value indicates the average spore fraction obtained by the strain in that row when co-developed with the strain in the corresponding column (based on N=3 blocks). Table 5 compares the linearity and steepness estimates for the four sites, along with the previously studied North Carolina (NC) site by Fortunato *et al.* (2003). Fortunato's study, which tested all pairwise combinations of seven strains collected from Little Butt's Gap (NC), demonstrated significant linearity, though steepness was not determined. We calculated the steepness for the NC hierarchy using their reported values (Table 2 from Fortunato *et al.* 2003). Similar to the NC site, sites AFCT and MLBS showed significant linearity based on Kendall's K metric (Table 5). Sites GSNP and SMFS did not show significant linearity. Steepness for each hierarchy, including NC, did not significantly differ from zero. To address potential issues that may result from calculating linearity at a low steepness, we additionally calculated the triangle transitivity (t_{tri}) (Sánchez-Tójar, Schroeder, and Farine 2018). Using this method, we observed a lower degree of linearity, but the hierarchies of sites AFCT and MLBS remained significantly linear (Table 5). In addition, using this metric, the hierarchy in site GSNP was marginally nonlinear (observed value is in the 3.4th percentile of the null distribution—i.e., P=0.068; explained in the Methods).

Table 1-4. The dominance hierarchy matrix for sites SMFS, AFCT, MLBS, and GSNP respectively. Each entry indicates the mean fraction of the spores achieved by the strain in that row when co-developed with the strain listed in the corresponding column, based on N=3 blocks.

SMFS	EO620	EO622	EO626	EO633	EO634	EO638
EO620	-	0.526	0.515	0.489	0.488	0.518
EO622		-	0.417	0.513	0.461	0.443
EO626			-	0.388	0.527	0.481
EO633				-	0.514	0.520
EO634					-	0.477
EO638						-

AFCT	EO676	EO677	EO679	EO681	EO684	EO686	EO690
EO676	-	0.631	0.596	0.586	0.475	0.450	0.424
EO677		-	0.479	0.505	0.478	0.391	0.335
EO679			-	0.532	0.544	0.508	0.409
EO681				-	0.498	0.458	0.461
EO684					-	0.446	0.493
EO686						-	0.505
EO690							-

MLBS	EO1273	EO1274	EO1275	EO1277	EO1279	EO1282	EO1283
EO1273	-	0.450	0.493	0.478	0.389	0.470	0.461
EO1274		-	0.527	0.533	0.413	0.537	0.512
EO1275			-	0.485	0.405	0.469	0.442
EO1277				-	0.417	0.394	0.464
EO1279					-	0.475	0.610
EO1282						-	0.509
EO1283							-

GSNP	EO600	EO602	EO603	EO604	EO605	EO611	EO615
EO600	-	0.577	0.537	0.628	0.495	0.476	0.501
EO602		-	0.504	0.623	0.572	0.421	0.478
EO603			-	0.623	0.572	0.421	0.478
EO604				-	0.502	0.506	0.467
EO605					-	0.515	0.485
EO611						-	0.603
EO615							-

Table 5. Summary of the properties of the four dominance hierarchies from this study and the NC population tested in Fortunato *et al.* (2003). K is a metric of linearity, P_t is the proportion of transitive triads, and t_{tri} is the triangle transitivity. See main text for explanations of terms. For site GSNP, t^{tri} is strongly negative and in the 3rd percentile of the distribution based on randomization (i.e., 966/1000 randomizations were more positive than the observed value, and thus $P=0.034*2=0.068$ (the P-value is doubled because significance is now tested in the opposite direction, i.e., two-tailed test).

	K	χ^2	df	$P(K)$	$P_{t \text{ (expect 0.75 if random)}}$	t_{tri}	$P(t_{tri})$	steepness	$p_{steepness}$
SMFS	0.250	38.7	30	<0.90	0.700	-0.200	0.777	0.030	1.00
AFCT	0.786	40.0	23.3	<0.025	0.914	0.657	0.029	0.064	1.00
MLBS	0.929	45.3	23.3	<0.005	0.971	0.886	0.009	0.060	1.00
GSNP	0.143	16.0	23.3	<0.90	0.657	-0.371	0.966[*]	0.040	1.00
NC (Fortunato *et al.* 2003)	0.929	45.3	23.3	<0.005	0.971	0.886	0.009	0.304	0.36

Are dominance hierarchies generally linear in populations of D. discoideum?

The North Carolina (NC) site, as tested by Fortunato, exhibited the same linearity as site MLBS (Table 5; K=0.929, $\chi^2_{23.3}$=45.3, P<0.005). Despite this similarity, MLBS displayed a very shallow hierarchy, with a steepness approaching zero, a pattern also observed in AFCT, GSNP, and SMFS. Notably, all four steepness values were substantially lower than that of the NC population. The lack of steepness in our study indicates much lower social dominance and, therefore, more equitable spore production. Specifically, in the Fortunato study, the average social dominance was 0.26 (across N=21 mixes), indicating a shift from a 50-50% distribution in the cells to 76-34% in the spores. In contrast, in this study, the average social dominance was only 0.045 (across N=78 mixes; Figure 4). Significant social dominance was observed in only 9% of all mixes: in 3 out of 21 mixes from GSNP, 2 out of 21 mixes from AFCT and MLBS, and none of the 15 mixes from SMFS. Significance was assessed based on a one-sample t-test, where the null hypothesis is that the observed social dominance is equal to that of the control mixes (Figure 4). In those mixes where we observed significant social dominance, the magnitude was 0.094±0.011 (mean±se, N=7), indicating a shift from a 50-50% distribution in the cells to 59-41% in the spores.

Figure 4. The magnitude of social dominance in control mixes and mixes of genetically different strains. Each point indicates the mean social dominance in a mix across *N*=3 blocks. For example, a value of 0.10 indicates a shift of 10%: from 50-50% in the cells to 60-40% in the spores. The control mixes are 'mixes' that consist of labeled and unlabeled cells of the same strain; these mixes serve as a baseline against which the mixes of genetically different strains can be compared.

Can differences in the inherent ability to dominate in the spores explain linearity?

Prior work found that social dominance and linearity resulted from inherent differences between strains in their

spore production (Buttery et al. 2009). The strains used in this study likewise showed significant variation in

spore production (SMFS: χ^2=3.8e6, df=1, *P*<0.001, GSNP: χ^2=2.9e6, df=1, *P*<0.001, AFCT: χ^2=1.4e6, df=1,

P<0.001, MLBS: χ^2=4.4e5, df=1 *P*<0.001), allowing us to test this hypothesis for the two sites with significant

linearity. To do so, we ranked the strains from most to least dominant based on their clonal spore production

(termed 'prior attribute' ranking) and compared it to the ranking derived from the outcomes of the pairwise

strain interactions (termed 'observed' ranking) (Figure 5). If the two rankings are similar, spore production is a

robust predictor of social dominance and linearity. Conversely, if the rankings differ substantially, then spore

production of the two strains is not a good predictor of social dominance and linearity. Figure 5 shows that in

both populations the two rankings exhibited multiple differences in order, suggesting that the differences

between strains in their clonal spore production are not strong predictors of the dominance hierarchy.

Figure 5. Comparison of the expected dominance rank (left), based on the prior attribute of a strain's solo spore production, and the observed dominance rank (right) for the two populations (AFCT and MLBS) that exhibit significant linearity. To visualize the changes in the order of the ranking, a single strain in the two rankings is connected by a line, and the crossing of two lines indicates changes in rank order.

Evidence that prior attributes, direct genetic effects, and social genetic effects influence social dominance

So far, we have examined the variation in social dominance exclusively as a product of a strain's own predetermined dominance ability—in this case, its ability to sporulate. However, dominance is likely also determined by the traits of the social partner. The relevant traits might not have been identified yet or might only be expressed during the social interaction. The influence of a social partner's genes on the phenotype of the focal individual is called indirect genetic effects (IGEs) (Wolf et al. 1998). Conversely, direct genetic effects (DGEs) describe the direct influence of an individual's own genes on its phenotype. In addition, when the interaction between genotypes is dependent on the specific combination of genotypes (i.e., social context) these effects are called genotype-by-genotype (G×G) interactions. Collectively, IGEs and G×G interactions are referred to as the social environment: the environment created by genes present in social partners (Wolf 2003).

To estimate the relative contributions of these different genetic effects to social dominance, we performed a quantitative genetic analysis (Bijma 2014). Since prior work showed that clonal spore production significantly influences a strain's spore fraction (Buttery et al. 2009), the model also included both the focal and partner strain's clonal spore production as explanatory factors. Lastly, we included site and genetic distance between strains in a mix. We performed two models: model 1 tested the focal strain's spore *fraction* in a mix as the response variable, and model 2 tested the focal strain's *total* spore production in a mix (discussed below).

The results of model 1 showed that a focal strain's spore fraction in a chimera was significantly influenced by its clonal spore production, as well as that of its partner strain (Table 6). There was no significant effect of genetic distance, and so this term was removed from the final model (χ^2=0.039, df=1, *P*=0.84). The focal strain's spore

fraction was significantly influenced by the direct genetic effect (DGE) of the focal strain, but not by the indirect

genetic effect (IGE) of the partner strain and the interaction between the genotypes of the focal and partner

strain (G×G interaction). Analysis of the partitioned genetic effects revealed that 53.1% of the random effects

variance was explained by DGEs, 28.5% by IGEs, and 17.4% by G×G interactions (Table 6). Because the focal

and partner genotypes can influence dominance through two different terms (their clonal spore production and

respective random effects), we assessed their relative contributions using an R^2 metric appropriate for mixed

models (Nakagawa and Schielzeth 2013). We found that 11% of the total variance could be explained by the

fixed effects (i.e., clonal spore production) and 63.7% by the fixed and random effects. Thus, although highly

significant, the clonal spore production of the two strains explains only a small fraction of the total variance.

This result indicates that the prior attributes were not the main drivers of the outcome: rather, these strains must

have other, unidentified attributes that largely determine their ability to dominate the spores.

Table 6. The summary of the model that tested the focal strain's fraction in the spores in chimera as a function of prior attribute clonal spore production and genetic identity of the focal and partner strain and their interaction. The variance explained by the individual genetic effects (DGE, IGE, and G×G interaction) and block were calculated by dividing each term's variance by the sum of the variances of all random effects.

	Term	df	χ^2	P	Variance	% Variance explained
Fixed	Focal clonal spore production	1	11.464	<0.001		
	Partner clonal spore production	1	8.551	0.003		
Random	Focal genotype (DGE)	1	14.925	<0.001	0.016	53.1
	Partner genotype (IGE)	1	5.268	0.022	0.009	28.5
	Focal-by-partner interaction (G×G)	1	0.422	0.52	0.005	17.4
	Block	1	0.036	0.85	0.0003	1.0
	Total				0.0303	

We found that total spore production in chimeras is often greater than what is expected based on the average

spore production of both strains when developed clonally, consistent with other studies (Buttery et al. 2009;

Kuzdzal-Fick et al. 2023). The excess spore production found in this study shows, however, no relationship with

the spore inequity observed in a mix (Figure S1; Spearman's rank correlation: r=-0.08, df=76, P=0.50).

Therefore, modelling variation in a strain's fraction in the spores as a response variable may ignore important

G×G interactions. Specifically, if both strains increase their spore allocation, but do so *equally*, then each

strain's spore fraction would remain at 0.5 despite a strong change in behavior in response to one another. For

this reason, we examined not only a strain's spore *fraction* (i.e., its dominance, as tested in Model 1) in chimera

but also its *total* spore production (Model 2).

The results of Model 2 showed that the focal strain's total spore production in chimera was significantly

influenced by the clonal spore production of the partner strain, but not that of the focal strain (Table 7). In

addition, there was a significant positive relationship between the genetic distance between the two strains and

spore production (estimate=0.003; P<0.001; 95% CI: 0.002-0.004). In other words, the more genetically

different two strains are, the more spores the focal strain produces.

In contrast to the earlier model, the focal strain's spore production was significantly influenced by IGEs, but not

by DGEs. Similar to before, there was no significant influence of the GxG interactions. Analysis of the

partitioned genetic effects revealed that 9.3% of the total variation in spore production was explained by DGEs,

82.4% by IGEs, 5.4% by G×G interactions, and 2.9% by block (Table 7). We again estimated the R^2 for the

fixed and random effects, which showed that 35.8% of the variance in outcome was attributable to the fixed

effects, and 43.5% was attributable to the fixed and random effects together. The small difference means that the

outcome is largely explained by the fixed effects—the clonal spore production of the two strains and how

genetically different they are. While significant, the random effects are small in magnitude. Finally, the total R^2

is also lower compared to the earlier model (43.5% versus 63.7%), indicating that the total spore production of

two strains together is less predictable than their percentages.

Table 7. Results of the model that tested the focal strain's total spore production in chimera as a function of prior

attribute clonal spore production, genetic distance, genetic identity of the focal and partner strain, and their

interaction. The variance explained by the individual genetic effects (DGEs, IGEs, and G×G interactions) and block were

calculated by dividing each term's variance by the sum of the variances of all random effects.

	Term	df	χ^2	P		
Fixed	Focal clonal spore production	1	63.602	<0.001		
	Partner clonal spore production	1	7.772	0.005		
	Genetic distance	1	39.862	<0.001		
	Term	df	χ^2	P	Variance	% Variance explained
Random	Focal genotype (DGE)	1	0.484	0.49	0.001	9.3
	Partner genotype (IGE)	1	17.49	<0.001	0.006	82.4
	Focal-by-partner interaction (G×G)	1	0.017	0.90	0.0004	5.4
	Block	1	0.150	0.70	0.0002	2.9
	Total				0.0076	

Discussion

Here we examined the structure of dominance hierarchies, focusing on their linearity and steepness, in four natural populations of the social amoeba *Dictyostelium discoideum*. In two populations, we identified significantly linear dominance hierarchies. In the third population, the hierarchy was not significantly linear, whereas in the fourth, it was marginally non-linear. Despite this variation, all four hierarchies were shallow, with relatively few significant deviations from spore equity. Together, these results indicate strains with a high degree of competitive similarity for social fitness.

Our finding of less linear and relatively flat social hierarchies (or: *less inequity*) differs from two earlier studies, that used the same seven isolates, which were collected at varying distances over a ~1 km^2 area of North Carolina (Fortunato, Queller, and Strassmann 2003; Buttery et al. 2009). Notably, our strains co-occurred in small soil samples of 10 g or less, and the tested strains were no more than 14 cm from each other at the time of sampling—and presumably, often much less. They were thus much more likely to consist of strains that do interact in nature. Our results underscore the importance of an appropriate spatial scale for the study of microbial cooperation in nature.

We expect co-occurring strains to be well-matched for important fitness characteristics, for several reasons. First, a large variance in social fitness can arise when selection is relaxed in the laboratory (Larsen et al. 2023). By similar logic, low variance is expected for traits subject to strong past selection, and so our observation of little social fitness variation is an important line of evidence that this trait could be experiencing selection in nature. In other words, co-occurring strains might co-occur *because* they do not have large fitness differences. Second, previous studies have shown that spore inequality will drive counter-evolutionary changes to restore fairness (Khare et al. 2009; Hollis 2012; Levin et al. 2015; Miller, Sidell, and Ostrowski 2023). Thus, both ecological and evolutionary processes should promote the co-existence of well-matched competitors.

The level of spore inequity we found in this study is similar in magnitude to what has been observed previously in laboratory screens for cheating behaviors (Santorelli et al. 2008, 2013; Miller, Sidell, and Ostrowski 2023) and another study using a different set of natural strains (Broersma and Ostrowski, in prep.), suggesting that

these results are robust and general. In addition, it implies that relatively few genetic differences could underlie

this level of spore inequity.

In animal societies, dominance hierarchies are often linear, and much work has been devoted to understanding

the determinants of a dominance hierarchy and whether it arises from 'prior attributes' (i.e., genetic attributes)

or emerges from 'social dynamics', which may include winner or loser effects (Chase and Seitz 2011). In this

study, we did not allow for winner/loser effects, as we re-grew the strains from frozen for each replicate. That

said, the variance attributed to block effects was *more than an order of magnitude smaller* than that attributed to

the different genetic effects (Table 6). Overall, these results suggest that the dominance hierarchies, despite

being shallow, were repeatable.

In the populations that did exhibit significant linearity, we could investigate the causes of that linearity. We

hypothesized that the difference in representation of the strains in the spores could reflect inherent differences in

their spore-stalk allocation—their 'prior attribute'. However, we observed many shifts in rank between the

expected performance of strains based on this prior attribute and their actual performance in the mix. Despite

this result, the focal and partner strains' clonal spore production were significant predictors of their percentage

of the spores, indicating that these prior attributes do contribute to the establishment of the linear dominance

hierarchy, at least in part. Our statistical analyses also indicate that social dominance was significantly

influenced by direct and indirect genetic effects, but not genotype x genotype interactions. In addition, we

detected these significant genetic effects *after* accounting for the clonal sporulation of the two strains. These

results suggest that the linearity results, in part, from additional, non-quantified dominance traits.

A major focus in this paper was on the percentage of the two strains in the spores—the reason for this focus is

clear: it reflects *relative* fitness differences between two strains. These differences are paramount for the

evolutionary dynamics of allele frequency changes. However, we also considered the *absolute* number of spores

produced by each strain. In prior work, we showed that a near-universal response to nonself strains is a

facultative increase in the differentiation to spore cells (Kuzdzal-Fick et al. 2023). Moreover, this increase likely

results from non-matching *tgrB1-tgrC1* alleles that mediate kin discrimination and influence cell type

proportioning responses downstream of this locus (Benabentos et al. 2009; Hirose et al. 2011, 2017; Kuzdzal-

Fick et al. 2023). These findings suggested that strains sense nonself and respond selfishly, increasing their

investment in the spores relative to the dead stalk. We again observed this response in this study. Taken together, our results are consistent with two types of selfishness, described previously (Liu et al. 2022). On one hand, we see that strains increase their allocation to the spores in the presence of non-identical strains. In the framework of Liu et al., this behaviour corresponds to 'selfishness' (i.e., a strategy of cooperating more or less depending on the relatedness of social interactants). On the other hand, we also see significant genetic effects of both the focal and partner strains as determinants of the outcome of a mix. This finding suggests that some of what determines social success may involve manipulative cheating and/or its suppression (Liu et al. 2022).

In summary, our study shows that linearity is indeed a common attribute of social amoeba hierarchies, as determined by four populations at large geographic distances from one another. However, the social amoeba social hierarchies tend to be shallow, indicating low levels of inequity. Nevertheless, there is significant inequity present in natural populations, given that the experimental mixes differed from the clonal controls. We show that prior attributes contribute to the linearity of the dominance hierarchy. However, we also find substantial contributions of DGEs and IGEs, where the traits were not specifically quantified in advance or predicted. Our study adds to a large body of work that indicates social dominance hierarchies are frequently linear. However, the shallowness indicates a relatively fair distribution of the spores among cooperating strains. Whether this substantial cooperation reflects that cheating has not arisen or that it is successfully suppressed is a crucial topic for future study.

Acknowledgements

The authors would like to thank Smith MacLeish Field Station, Proctor Academy, Mountain Lake Lodge, and Mountain Lake Biological Station for allowing us to collect soil samples. Soil collection in Great Smoky National Park was done under permit GRSM-2017-SCI-2014. We thank Michael Miller, Maria Polo Prieto, Rafael Polo, Jennie Kuzdzal-Fick, Armando Moreno, and Scott Clark for assistance with soil collection and strain isolation. We thank David Fisher for his advice on the statistical models. This work was funded by grants to EAO from the US National Science Foundation DEB-1557023 and the Marsden Fund, administered by Te Apārangi Royal Society of New Zealand, as well as a Massey University Doctoral Scholarship to CMB.

Conflict of interest

None declared.

Author contributions

CMB and EAO conceived the idea and designed the experiments. CMB collected the data. CMB and EAO

analyzed the data, wrote the manuscript, and approved the final version.

Data accessibility

Data and scripts are available upon request.

Literature cited

Appleby, Michael C. 1983. "The Probability of Linearity in Hierarchies." *Animal Behavior* 31: 600–608.

Barkan, C. P. L., J. L. Craig, S. D. Strahl, A. M. Stewart, and J. L. Brown. 1986. "Social Dominance in Communal Mexican
Jays Aphelocoma Ultramarina." *Animal Behaviour* 34 (February): 175–87.

Barrett, L., S. P. Henzi, T. Weingrill, J. E. Lycett, and R. A. Hill. 1999. "Market Forces Predict Grooming Reciprocity in
Female Baboons." *Proceedings of the Royal Society of London. Series B: Biological Sciences* 266 (1420): 665–70.

Beaugrand, J. P., and P. A. Cotnoir. 1996. "The Role of Individual Differences in the Formation of Triadic Dominance
Orders of Male Green Swordtail Fish (Xiphophorus Helleri)." *Behavioural Processes* 38 (3): 287–96.

Benabentos, Rocio, Shigenori Hirose, Richard Sucgang, Tomaz Curk, Mariko Katoh, Elizabeth A. Ostrowski, Joan E.
Strassmann, et al. 2009. "Polymorphic Members of the Lag Gene Family Mediate Kin Discrimination in
Dictyostelium." *Current Biology: CB* 19 (7): 567–72.

Bijma, P. 2014. "The Quantitative Genetics of Indirect Genetic Effects: A Selective Review of Modelling Issues." *Heredity*
112 (1): 61–69.

Bijma, P., and M. J. Wade. 2008. "The Joint Effects of Kin, Multilevel Selection and Indirect Genetic Effects on Response
to Genetic Selection." *Journal of Evolutionary Biology* 21 (5): 1175–88.

Buss, L. W. 1982. "Somatic Cell Parasitism and the Evolution of Somatic Tissue Compatibility." *Proc. Natl. Acad. Sci. U S
A* 79: 5337–41.

Buttery, N. J., D. E. Rozen, J. B. Wolf, and C. R. L. Thompson. 2009. "Quantification of Social Behavior in *D. Discoideum*
Reveals Complex Fixed and Facultative Strategies." *Curr. Biol.* 19 (16): 1373–77.

Chase, I. D., and K. Seitz. 2011. "Self-Structuring Properties of Dominance Hierarchies: A New Perspective." *Advances in
Genetics* 75: 51–81.

Chase, I. D., C. Tovey, D. Spangler-Martin, and M. Manfredonia. 2002. "Individual Differences versus Social Dynamics in
the Formation of Animal Dominance Hierarchies." *Proceedings of the National Academy of Sciences of the United
States of America* 99 (8): 5744–49.

Clutton-Brock, T. H., S. D. Albon, and F. E. Guinness. 1984. "Maternal Dominance, Breeding Success and Birth Sex Ratios in Red Deer." *Nature* 308 (5957): 358–60.

Correa, Loreto A., Beatriz Zapata, Horacio Samaniego, and Mauricio Soto-Gamboa. 2013. "Social Structure in a Family Group of Guanaco (Lama Guanicoe, Ungulate): Is Female Hierarchy Based on 'prior Attributes' or 'Social Dynamics'?" *Behavioural Processes* 98 (September): 92–97.

Curley, J. P. 2016. *Compete: Analyzing Social Hierarchies: R Package* (version 0.1). https://github.com/jalapic/compete.

Drews, Carlos. 1993. "The Concept and Definition of Dominance in Animal Behaviour." *Behaviour* 125 (3): 283–313.

Filosa, M. F. 1962. "Heterocytosis in Cellular Slime Molds." *The American Naturalist* 96 (887): 79–91.

Fortunato, A., D. C. Queller, and J. E. Strassmann. 2003. "A Linear Dominance Hierarchy among Clones in Chimeras of the Social Amoeba Dictyostelium Discoideum." *Journal of Evolutionary Biology* 16 (3): 438–45.

Goessmann, C., C. Hemelrijk, and R. Huber. 2000. "The Formation and Maintenance of Crayfish Hierarchies: Behavioral and Self-Structuring Properties." *Behavioral Ecology and Sociobiology* 48 (6): 418–28.

Hamilton, W. D. 1964a. "The Genetical Evolution of Social Behaviour. I." *Journal of Theoretical Biology* 7 (1): 1–16.

———. 1964b. "The Genetical Evolution of Social Behaviour. II." *Journal of Theoretical Biology* 7 (1): 17–52.

Hausfater, G., J. Altmann, and S. Altmann. 1982. "Long-Term Consistency of Dominance Relations Among Female Baboons (*Papio Cynocephalus*)." *Science* 217 (4561): 752–55.

Heinze, J. 1990. "Dominance Behavior among Ant Females." *Die Naturwissenschaften* 77 (1): 41–43.

Hirose, Shigenori, Rocio Benabentos, Hsing-I Ho, Adam Kuspa, and Gad Shaulsky. 2011. "Self-Recognition in Social Amoebae Is Mediated by Allelic Pairs of Tiger Genes." *Science* 333 (6041): 467–70.

Hirose, Shigenori, Gong Chen, Adam Kuspa, and Gad Shaulsky. 2017. "The Polymorphic Proteins TgrB1 and TgrC1 Function as a Ligand-Receptor Pair in Dictyostelium Allorecognition." *Journal of Cell Science* 130 (23): 4002–12.

Hollis, Brian. 2012. "Rapid Antagonistic Coevolution between Strains of the Social Amoeba Dictyostelium Discoideum." *Proceedings. Biological Sciences / The Royal Society* 279 (1742): 3565–71.

Huss, Martin J. 1989. "Dispersal of Cellular Slime Molds by Two Soil Invertebrates." *Mycologia* 81 (5): 677–82.

Jack, Chandra N., Neil Buttery, Boahemaa Adu-Oppong, Michael Powers, Christopher R. L. Thompson, David C. Queller, and Joan E. Strassmann. 2015. "Migration in the Social Stage of Dictyostelium Discoideum Amoebae Impacts Competition." *PeerJ* 3 (October): e1352.

Jackson, Wendy M., and Robin L. Winnegrad. 1988. "Linearity in Dominance Hierarchies: A Second Look at the Individual Attributes Model." *Animal Behaviour* 36 (4): 1237–40.

Kessin, R. H. 2001. *Dictyostelium. Evolution, Cell Biology, and the Development of Multicellularity*. Edited by J. B. L. Bard, P. W. Barlow, and D. L. Kirk. Developmental and Cell Biology. Cambridge University Press.

Khare, Anupama, Lorenzo A. Santorelli, Joan E. Strassmann, David C. Queller, Adam Kuspa, and Gad Shaulsky. 2009. "Cheater-Resistance Is Not Futile." *Nature* 461 (7266): 980–82.

Kuzdzal-Fick, J. J., K. R. Foster, D. C. Queller, and J. E. Strassmann. 2007. "Exploiting New Terrain: An Advantage to

Sociality in the Slime Mold Dictyostelium Discoideum." *Behavioral Ecology: Official Journal of the International*

Society for Behavioral Ecology 18 (2): 433–37.

Kuzdzal-Fick, J. J., A. Moreno, C. M. E. Broersma, T. F. Cooper, and E. A. Ostrowski. 2023. "From Individual Behaviors to

Collective Outcomes: Fruiting Body Formation in Dictyostelium as a Group-Level Phenotype." *Evolution;*

International Journal of Organic Evolution 77 (3): 731–45.

Larsen, Tyler J., Israt Jahan, Debra A. Brock, Joan E. Strassmann, and David C. Queller. 2023. "Reduced Social Function in

Experimentally Evolved Dictyostelium Discoideum Implies Selection for Social Conflict in Nature." *Proceedings.*

Biological Sciences / The Royal Society 290 (2013): 20231722.

LeBrun, Edward G. 2005. "Who Is the Top Dog in Ant Communities? Resources, Parasitoids, and Multiple Competitive

Hierarchies." *Oecologia* 142 (4): 643–52.

Levin, S. R., D. A. Brock, D. C. Queller, and J. E. Strassmann. 2015. "Concurrent Coevolution of Intra-Organismal Cheaters

and Resisters." *Journal of Evolutionary Biology* 28 (4): 756–65.

Miller, M. E., S. Sidell, and E. A. Ostrowski. 2023. "Costs of Resistance Limit the Effectiveness of Cooperation

Enforcement." *BioRxiv*. https://doi.org/10.1101/2023.08.29.555259.

Moore, A. J., E. D. Brodie, and J. B. Wolf. 1997. "Interacting Phenotypes and the Evolutionary Process: I. Direct and

Indirect Genetic Effects of Social Interactions." *Evolution; International Journal of Organic Evolution* 51 (5):

1352–62.

Nakagawa, Shinichi, and Holger Schielzeth. 2013. "A General and Simple Method for ObtainingR2from Generalized Linear

Mixed-Effects Models." *Methods in Ecology and Evolution* 4 (2): 133–42.

Nakano, Shigeru, and Tetsuo Furukawa-Tanaka. 1994. "Intra- and Interspecific Dominance Hierarchies and Variation in

Foraging Tactics of Two Species of Stream-Dwelling Chars." *Ecological Research* 9 (1): 9–20.

Nelissen, Mark H. J. 1992. "Does Body Size Affect the Ranking of a Cichild Fish in a Dominance Hierarchy?" *Journal of*

Ethology 10 (2): 153–56.

Neumann, Christof, Julie Duboscq, Constance Dubuc, Andri Ginting, Ade Maulana Irwan, Muhammad Agil, Anja Widdig,

and Antje Engelhardt. 2011. "Assessing Dominance Hierarchies: Validation and Advantages of Progressive

Evaluation with Elo-Rating." *Animal Behaviour* 82 (4): 911–21.

Sánchez-Tójar, Alfredo, Julia Schroeder, and Damien Roger Farine. 2018. "A Practical Guide for Inferring Reliable

Dominance Hierarchies and Estimating Their Uncertainty." *The Journal of Animal Ecology* 87 (3): 594–608.

Santorelli, Lorenzo A., Adam Kuspa, Gad Shaulsky, David C. Queller, and Joan E. Strassmann. 2013. "A New Social Gene

in Dictyostelium Discoideum, ChtB." *BMC Evolutionary Biology* 13 (January): 4.

Santorelli, Lorenzo A., Christopher R. L. Thompson, Elizabeth Villegas, Jessica Svetz, Christopher Dinh, Anup Parikh,

Richard Sucgang, et al. 2008. "Facultative Cheater Mutants Reveal the Genetic Complexity of Cooperation in

Social Amoebae." *Nature* 451 (7182): 1107–10.

Schjelderup-Ebbe, T. 1922. "Beiträge Zur Sozialpsychologie Des Haushuhns." *Zeitschrift Für Psychologie Und Physiologie*

Der Sinnesorgane 88: 225–52.

Schjolden, Joachim, Argaudas Stoskhus, and Svante Winberg. 2005. "Does Individual Variation in Stress Responses and

Agonistic Behavior Reflect Divergent Stress Coping Strategies in Juvenile Rainbow Trout?" *Physiological and*

Biochemical Zoology: PBZ 78 (5): 715–23.

Schmid, V. S., and H. de Vries. 2013. "Finding a Dominance Order Most Consistent with a Linear Hierarchy: An Improved

Algorithm for the I&SI Method." *Animal Behaviour* 86 (5): 1097–1105.

Shizuka, Daizaburo, and David B. McDonald. 2012. "A Social Network Perspective on Measurements of Dominance

Hierarchies." *Animal Behaviour* 83: 925–34.

Smith, J., D. C. Queller, and J. E. Strassmann. 2014. "Fruiting Bodies of the Social Amoeba Dictyostelium Discoideum

Increase Spore Transport by Drosophila." *BMC Evolutionary Biology* 14 (May): 105.

Strassmann, J. E., Y. Zhu, and D. C. Queller. 2000. "Altruism and Social Cheating in the Social Amoeba Dictyostelium

Discoideum." *Nature* 408 (6815): 965–67.

Sundström, L. Fredrik, Erik Petersson, Johan Höjesjö, Jörgen I. Johnsson, and Torbjörn Järvi. 2004. "Hatchery Selection

Promotes Boldness in Newly Hatched Brown Trout (Salmo Trutta): Implications for Dominance." *Behavioral*

Ecology: Official Journal of the International Society for Behavioral Ecology 15 (2): 192–98.

Valderrabano-Ibarra, C., I. Brumon, and H. Drummond. 2007. "Development of a Linear Dominance Hierarchy in Nestling

Birds." *Animal Behaviour* 74 (6): 1705–14.

Vehrencamp, S. L. 1983. "Optimal Degree of Skew in Cooperative Societies." *American Zoologist* 23 (2): 327–35.

Vries, Han de. 1995. "An Improved Test of Linearity in Dominance Hierarchies Containing Unknown or Tied

Relationships." *Animal Behaviour*. https://doi.org/10.1016/0003-3472(95)80053-0.

Vries, Han de, Jeroen M. G. Stevens, and Hilde Vervaecke. 2006. "Measuring and Testing the Steepness of Dominance

Hierarchies." *Animal Behaviour* 71 (3): 585–92.

Vullioud, Colin, Eve Davidian, Bettina Wachter, François Rousset, Alexandre Courtiol, and Oliver P. Höner. 2019. "Social

Support Drives Female Dominance in the Spotted Hyaena." *Nature Ecology & Evolution* 3 (1): 71–76.

Wilson, E. O., ed. 1975. *Sociobiology: The New Synthesis*. The Belknap Press of Harvard University Press Cambridge,

Massachusetts, and London, England.

Wittemyer, G., and W. M. Getz. 2007. "Hierarchical Dominance Structure and Social Organization in African Elephants,

Loxodonta Africana." *Animal Behaviour* 73 (4): 671–81.

Wolf, J. B. 2003. "Genetic Architecture and Evolutionary Constraint When the Environment Contains Genes." *Proceedings*

of the National Academy of Sciences of the United States of America 100 (8): 4655–60.

Wolf, J. B., E. D. Brodie I, J. M. Cheverud, A. J. Moore, and M. J. Wade. 1998. "Evolutionary Consequences of Indirect

Genetic Effects." *Trends in Ecology & Evolution* 13 (2): 64–69.