**Title: Happiness promotes global processing in haptic perception**

Müge Cavdan[§,†*1], Aycan Kapucu[‡], Katja Doerschner[§,†+], & Knut Drewing[§+]

[§]Experimental Psychology Department, Justus Liebig University Giessen; Otto-Behaghel

Strasse 10F, 35394, Giessen, Germany

[†]Center for Mind, Brain, and Behavior (CMBB), Philipps University Marburg and Justus

Liebig University Giessen; Hans-Meerwein-Strasse 6, 35032, Marburg, Germany

[‡] Psychology Department, Ege University, Izmir, Türkiye

*Correspondence : Muege.Cavdan@psychol.uni-giessen.de ; X Handle – @mugecavdan

[+]Equal contribution

10                                    **Summary**

Happy and sad mood promote global and local visual processing, respectively. However, it is

unclear whether mood also affects the processing level in haptic perception. Here, we used

classical music to induce happy and sad mood in blindfolded participants before they scanned

3D-printed configurations with their fingers. Global shapes were triangles, circles, or  squares

(33mm) composed of smaller local relief shapes (3 mm): either triangles, circles, or squares.

Participants explored a probe stimulus with identical local and global shapes, and two

comparison stimuli, matching the probe in local, or global shape. They reported which

comparison stimulus appeared more similar to the probe. In the 'sad' group, participants chose

the locally-matching comparison more frequently than in the 'happy' group, revealing that sad

mood promotes local processing also in touch. Overall, participants chose the globally-

matching comparison more often, suggesting that global processing is more prominent in touch

than assumed.

---

[1] lead contact: Muege.Cavdan@psychol.uni-giessen.de

## Introduction

Feelings modulate how we perceive and interact with the world. Mood modulates cognitive processes and social behavior, e.g., happy feelings promote reliance on accessible general information such as stereotypes[1,2] and scripts[3] while sad feelings have been associated with focusing on details, avoiding stereotypes, which often leads to an improved accuracy of judgements[4]. One explanation for this shift of focus is offered by the affect-as-information hypothesis[5,6]. According to this theory, mood is a signal that can be used to guide behavior: happy mood signals a low-threat situation that affords global processing whereas sad mood may signal that a situation is potentially threatening and therefor require more detailed processing. An alternative explanation for the association between mood and level of processing suggests that motivation may act as a mediator. According to this idea detailed processing is effortful and may result in a reduction of mood[7]. Consequently, when happy, people may be less motivated to perform effortful, detailed processing, in order to maintain their positive mood.

Interestingly, this shift from general, global to detailed, local processing as the mood shifts from happy to sad, has not only been observed in social and cognitive studies but also emerges robustly in visual shape perception[3,8,9]. For example, in an experiment by Gasper and Clore[9], 'happy' and 'sad' participants indicated the similarity of a given shape (triangle or square) that consisted of local elements (triangle or square, also see Kimchi & Palmer[10] to two figures that either matched the global or the local shape of the stimulus. Their results showed that individuals in the sad mood condition more often classified shapes based on local features compared to 'happy'. In line with this, Curby and collegues[11] showed that negative mood decreased the holistic processing of faces. The attentional narrowing of attention when mood is low[8,12–16] is interesting, as it appears to 'override' the default holistic processing visual shape and scene perception[17,18].

In general, the haptic system applies more weight to local features than global ones

when judging shapes[19]. Thus, it is not clear how and in what direction changes in mood would

modulate the haptic perception of shape. To address these questions, we investigated how

positive and negative mood influence the level of haptic shape processing. Upon mood

induction, blindfolded participants scanned a haptic shape stimulus with their fingers and

indicated which one of the two comparison stimuli was more similar to the probe. Comparisons

matched the probe in either global or local shape. Our working hypothesis was that happy mood

would promote global processing while sad mood would promote a focus on local shape

features. Our results provide a deeper understanding to what extend mood-induced changes in

attentional focus might be a general perceptual phenomenon.

**Methods**

**Participants**

The required sample size was calculated based on a large effect (Cohen's $d = 0.8$), power of

80%, and alpha 5%. The projected sample size was 42 for one-sided between-subjects $t$-test

(G*Power[33]). Accordingly, 42 participants (9 males, $M_{age} = 22.8$, age range: 18-30) were

randomly assigned to one of the two mood conditions (happy or sad). They were recruited

through the university email system and compensated with 8 €/hour or course credit. None of

the participants reported sensory, motor, or cutaneous impairments. The two-point

discrimination threshold at the index finger of the dominant hand was 3 mm or better. The

study was ethically approved by a local ethics committee LEK FB06 (2017-0034) in

accordance with the Declaration of Helsinki[34] excluding the preregistration. Informed consents

were obtained prior to the experiment.

**Stimuli and materials**

*Haptic Stimuli.* Haptic stimuli were modeled using the OpenSCAD software and 3D printed

(Objet30Pro, StratasysLtd., USA) with model photopolymer material (VeroClear) at a

resolution of $600 \times 600 \times 1.600$ dpi (x, y, and z axes respectively). As previous research has

shown that the haptic system weighs local shape features more than global features[19], we first

conducted a pilot experiment where we investigated the detection thresholds of a small triangle,

a circle, and a square in order to determine the shape size that would allow us to reduce the

natural saliency of the local features relative to the global ones in the main experiment. Stimuli

were two-dimensional (2D) triangles, squares, and circles ranging from 1 mm to 10 mm (size:

diameter of a circle, one side of a square, and triangle base; shapes embossed with a height of

1 mm on a plane plate). Blindfolded participants touched the shapes and named the shape they

perceive. For each shape mean value, the sizes where a person recognized the shapes correctly

were calculated. These values were taken as a threshold and then used as the local shape sizes

for the later stimuli production in the main experiment.

Nine stimuli consisting of local and global features modelled using OpenSCAD

software and 3D printed at a high resolution as in the pilot experiment. The global shapes were

printed on a 40 mm (x-axis) $\times$ 40 mm (y-axis) long and 5 mm (z-axis) thick base. The stimuli

were mounted on a 3D printed plate which has three squares. This ensured that the stimuli

remained fixed on the table during the exploration. Local shape sizes were 3 mm (both x and

y-axes) and global shapes size was 33 (both x and y-axes) for square, circle, and triangle with

the height of 1 mm (z-axes). The distance between each adjacent local shape was kept constant

at 1 mm. The shapes consisted of a circle of circles, a triangle of triangles, and a square of

squares as probes while comparison stimuli were a circle of triangles, a square of triangles, a

triangle of squares, circle of squares, a square of circles, and a triangle of circles. The stimuli

were combined with each other to make triads (six triads in total). Each triad consisted of one

probe (i.e., standard) and two comparison stimuli (see Figure 1 for an example). Probes had

the same global and local shapes such as square of squares. Comparison stimuli on the other

hand, matched either local or global shape of the probe such as square of circles (global match)

and circle of squares (local match).

*Mood induction stimuli and manipulation check.* Classical music pieces from Ackerley et al.[35]

were tested in a pilot experiment in order to affirm their effectiveness in our population[2]. The

pieces that evoked strongest happiness and sadness were selected for mood induction in the

current study. *Kol Nidrei* by *Max Bruch (1938-1920)* and *Carmen Suite Nr. 1: Les Toréadors*

by *Georges Bizet (1838-1875)* were used to induce sad and happy mood, respectively.

In order to measure the efficacy of mood induction, two different scales were used

before and after the mood induction, the Self-Assessment Manikin (SAM[36] and the Differential

Emotions Scale – DES[37,38]). The SAM measured participants' valence and arousal levels on a

9-point pictorial depiction (*valence*: 1= very negative to 9= very positive; *arousal*: 1= very

calm to 9= very excited). Participants were asked to rate the SAM based on their emotional

state. The DES is widely used in psychological research and a reliable method to assess

emotions. It consists of several subscales with various adjectives. We selected the sadness and

happiness subscales with their corresponding three adjectives per subscale; happiness: *happy,*

*joyful, amused*; sad: *sad, downhearted*, and *blue*. The adjectives were presented with a 9-point

---

[2] Total of 20 participants participated in the between-subjects pilot experiment. Participants were randomly assigned to either happy or sad condition. Participants in the happy condition listened to the happy classical music pieces while participants in the sad condition listened to the sad classical music. Although more than one classical piece was presented in the experiment, we will be focusing on the pieces subject to the current article. Upon listening to either *Carmen Suite Nr.1 : Les Toréadors* or *Kol Nidrei* participants indicated the strongest level of each emotion they experienced while listening to the music piece on a 9-point scale (0: not at all, 8: extremely). The following emotion words were rated in a random order; happiness: *happy, joyful, amused*; sad: *sad, downhearted*, and *blue*. The average scores across happiness and sadness were calculated. Two-independent samples t-test were used to test the efficacy of the classical music pieces. For happiness, the happy music piece yielded a higher rating than the sad piece, $t$ (18) = 4.65 , $p$ < .01, $d$ = 2.08. For sadness, the sad music piece yielded a higher rating than the happy piece, $t$ (18) = -5.40 , $p$ < .01, $d$ = 2.41. These results clearly show the efficacy of both classical music pieces in inducing the desired target moods.

114    scale (0: *not at all*, 8: *extremely*). Participants rated these adjectives based on how much they

115    felt each emotion.

**Procedure**

A between-subjects design with two moods – happy and sad – was used. Participants

were randomly assigned to either the happy or the sad mood conditions. Upon arrival in the

lab, a participant received information about the experiment and signed an informed consent.

This was followed by a two-point touch discrimination threshold test to measure any tactile

deficiencies[39]. Next, participants responded to the mood questions of SAM and DES, presented

on a monitor screen, using a numpad. The presentation order of the questionnaires was

randomized. After completion of the questionnaires, a classical music piece, either happy or

sad - depending on the condition, was played. After this  SAM and DES questionnaires were

completed    a second time. Finally, participants performed the Kimchi-Palmer task[10], as

described next.

On each trial, a probe (i.e., same global and local shape) was always presented in the

first square of the mounted plate (Figure 1). The two comparison shapes (one matching the

global shape of the probe and the other matching the local shape) were presented in the next

two squares of the mounted plate. The presentation order of the second and third gaps were

counterbalanced. Blindfolded participants were asked to explore each stimulus of the triad

sequentially (from first gap to third) using lateral movements with the index finger of the

dominant hand. Participants were able to re-explore the probe and the comparison stimuli as

many times as wanted. Their task was to indicate which of the two comparison stimuli was

more similar to the probe.

Each of the six triads was repeated 6 times which resulted in 36 trials in total. The

overall procedure lasted 30-45 minutes.

**Data analysis**

All analyses were conducted in SPSS 25 software (SPSS inc.) and figures are plotted in

MATLAB R2021b (MathWorks inc.).

Cronbach's alpha was used to calculate reliability per happiness and sadness subscales (DES)

for pre- and post-induction. All four analyses showed good to excellent consistency

(Cronbach's $a$ = .70 to .92) allowing for averaging across adjectives. Next, we calculated the

average values across the three adjectives per mood (i.e., happiness: happy, joy, and amused;

sadness: sad, downhearted, and blue) which were then submitted to an ANOVA.

For the Kimchi-Palmer task, the frequency of selecting the local comparison shape over

the global one was calculated from the raw data of each participant. Selecting the local match

indicated that the participant relied on local shape processing. The percentage of local

processing across all trials was calculated per participant in order to conduct a t-test analysis.

Greenhouse-Geisser correction was applied when the sphericity assumption was violated.

Similarly, degrees of freedom were adjusted when Levene's test indicated unequal variances.

**Results**

**Global and local processing**

An independent-samples $t$-test was used to determine the effect of mood on local processing:

Local processing was significantly less frequent after listening to the happy music piece, than

after listing to the sad music piece: $t$ (25.92) = -2.30 , $p$ = .015 (one-tailed), $d$ = .71 (Figure 2).

Overall the global match was selected six times more often than the local one (Figure 2).

Next, we tested whether there was an effect of global-local shape combination in level

of processing. To this end, we conducted a repeated-measures ANOVA with the variable shape

triad (6 levels). The main effect of shape triad was not statistically significant, $F$(3.87, 158.63)

$= 1.44$, $p = .21$, $\eta_p{}^2 = .03$, showing that the percentage of local match choice was not dependent

on a specific shape-combination. Finally, to test the possible effects of presentation orders on

the level of processing, 6 paired-samples t-tests were conducted (i.e., one per shape-

combination). None of the t-test were statistically significant showing that there was no effect

of presentation order on the percentage of local match (all $p >.05$).

**Mood manipulation check**

The efficacy of mood induction was examined using 4 mixed-design ANOVAs to test the effect

of music piece (levels: *happy* and *sad*) and time point (levels: *pre-induction* and *post-induction*)

separately on mean ratings for Self-assessment Manikin (SAM) valence, SAM arousal,

Differential Emotion Scale (DES) happiness, and DES sadness (see Table 1 for the summary

of the results).

*SAM.*  As expected, listening to the happy music piece increased **valence** towards

positive, while listening to the sad music piece decreased it:  The main effects of mood and

time point were not statistically significant for valence, $F(1,40) = 2.21$, $p = .15$, $\eta_p{}^2 = .05$,

$F(1,40) = 0.17$, $p = .90$, $\eta_p{}^2 = 00$, respectively. However, the interaction between time and mood

was statistically significant, $F(1,40) = 8.94$, $p < .01$, $\eta_p{}^2 = .18$. Follow-up analyses showed that,

in the happy mood condition, participants were more positive in the post-test compared to the

pre-test, $t(20) = -2.33$, $p = .015$ (one-sided). Also expectedly, after listening to the sad musical

piece, participants reported to be in a more negative mood than during the pre-test phase, $t(20)$

$= 1.98$, $p = .03$ (one-sided, Figure 3).

The sad music piece decreased **arousal**, as expected: The main effect of mood, $F(1,40) = 9.81$,

$\eta_p{}^2 = .20$ and the interaction between time point and mood, $F(1,40) = 11.24$, $\eta_p{}^2 = .22$, were

statistically significant (both $p < .01$). However, the main effect of time point was not

statistically significant for arousal, $F(1,40) = .70$, $p = .41$, $\eta p{}^2 = .02$. Follow-up analyses showed

that for happy mood, there was no significant arousal difference between pre-test and post-test,

$t(20) = -1.92$, $p = .07$. As expected , for sad mood, arousal decreased on post-test compared to

pre-test, $t(20) = 2.78$, $p < .01$ (Figure 3).

*DES*. Participants in the happy mood were happier upon mood induction. However,

participants in sad mood did not feel significantly sadder after manipulation which will be

discussed further.

For the happiness subscale, neither the main effect of time point, $F(1,40) = .28$, $\eta_p{}^2 =$

.01, nor the main effect of mood, $F(1,40) = 2.34$, $\eta_p{}^2 = .06$ were statistically significant.

However, the interaction between time point and mood was statistically significant $F(1,40) =$

11.04, $p < .01$, $\eta_p{}^2 = .06$. In the happy mood condition happiness increased between pre- and

post-test, whereas in the sad mood condition happiness decreased between pre- and post-test:

$t(20) = -3.11$, $p < .01$, $t(20) = 1.78$, $p < .05$ (both one-sided).

For the sadness subscale, although people in the sad mood felt less happy and more sad,

this did not reach significance: The main effect of time point: $F(1,40) = .01$, $\eta_p{}^2 = .00$ , mood:

$F(1,40) = .00$, $\eta_p{}^2 = .00$ and their interaction: $F(1,40) = 1.74$ , $\eta_p{}^2 = .04$ (all $p > 0.05$) were not

statistically significant.

**Discussion**

When we look at a scene, we have the ability to focus on a single element or to take in the

entire shape formed by these individual elements. Gasper and Clore[9] suggest that whether we

perceive the entire shape or the single element depends on our mood. Namely, while happy

mood fosters the entire shape view (i.e., global processing), sad mood fosters the single element

view (i.e., local processing). Here, we showed that this effect is not limited to vision but a

broader mechanism that operates at the perceptual level, and is therefore also present in the

haptic modality. Individuals in the happy mood condition matched the probe with the global

HAPPINESS PROMOTES GLOBAL PROCESSING IN HAPTICS

shape more often than individuals in the sad mood condition. Interestingly, individuals,

independent of the mood condition, more often matched the global target to the probe – the

global match was selected overall six times more often than the local one. This overall global

perception preference is interesting considering that typically people focus on local details

when exploring objects with their hands[19–21]. However, the objects used in these earlier studies

were larger than a typical hand, which might have pushed individuals to do sequential

exploration, which might have resulted in increased focus on local detail. Thus, the current

global precedence finding might be due to the overall size of the global form[22,23], as well as

the density[22–24] of the local features (see[25] for a meta-analysis on the environmental factors

moderating global precedence. Although local-global focus preferences did not receive the

same attention in haptic perception as in vision, a recent haptic study reported a similar global

preference as we observed in our experiment: the smaller the local shape the more global

processing was observed[23]. Therefore, the level of processing seems to critically depend on the

number and relative size of the elements[25,26]. It would be interesting, in future experiments, to

identify the local-global relative size relationship where the perceptual focus switches.

The general observed global preference suggests that the most available and easily

accessible information was the global form of our stimuli. The usage of global information,

however, was modulated by the mood: Individuals in the happy mood condition selected the

global match to the probe more frequently than individuals in the sad mood condition, and this

effect was independent of the presentation order or the geometrical features (circle, triangle,

square). Our finding is in line with the levels-of-focus hypothesis arguing that individuals in

happy mood condition are more likely to perceive the easily accessible knowledge resulting in

broader attentional processing[9]. In contrast, sad mood would result in increased detailed (local)

processing. Consistent with this idea, we observed (informally[3]) that participants in the sad

mood condition in our experiment spent more time when exploring the stimuli.

Not only mood influences the processing of the stimuli but also using a certain type of

processing (global or local) might facilitate a mood[27]. This has been also observed for face

identification – global processing facilitated happy face identification while local processing

facilitated sad face identification[28,29]. Future research should investigate the possible

bidirectionality between mood and level of processing in haptics.

Our work might have potential clinical relevance. For example, obsessive compulsive

individuals tend to focus on details[30,31], a cognitive style, that has been associated with local

interference of small details when the task was the identification of global information[32]. Future

studies should test whether individuals with obsessive compulsive cognitive style use more

local processing compared to a control group also in haptic tasks. Insights from such studies

potentially yield the development of novel diagnosis tools for this group of patients.

In conclusion, individuals in happy mood are more likely to *feel* the entire shape. Using

a stimulus set consisting of local and global features and a haptic shape perception task, we

found that happy mood fosters global processing compared to sad mood. To the best of our

knowledge, this is the first study that extended the levels-of-processing effect found in vision

to the haptic domain. Unlike previous research in haptic perception, we also found global

precedence when people perceive different geometrical shapes. Therefore, more research is

needed to understand the different components of global and local processing in human haptic

perception.

---

[3] Note, that timing of the trials was not recorded and the differences in timing between conditions was only an informal observation by the experimenter.

**Author contributions**

Conceptualization: M.C., A.K., K.D., K.Dr; Methodology: M.C., K.Dr.; Investigation: M.C.; Data curation: M.C.; Formal analysis: M.C., A.K.; Visualization: M.C., K.D.; Writing— original draft: M.C.; Writing—reviewing and editing: M.C., A.K., K.D., K.Dr; Supervision: M.C., A.K., K.D., K.Dr.; Funding Acquisition: K.D., K.Dr.; Project Administration: M.C., K.Dr.

**Acknowledgement**

Authors thank Hatice Dokumaci for the haptic stimuli and data collection, Rochelle Ackerley for sharing the classical music pieces. This research was supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 222641018 – SFB/TRR 135 (A5 and B8) and EU Marie Curie Initial Training Network "DyVito" (H2020-ITN, Grant Agreement: 765121). K.Dr and M.C. were supported by DFG-project no. 502774891-ORA project "UNTOUCH", K.Do was supported by Research Cluster "The Adaptive Mind", funded by the Excellence Program of the Hessian Ministry of Higher Education, Science, Research and the Arts. Artificial intelligence.

279    **Figures**

Probe                Global match                Local match

284    **Figure 1.** Illustration of an example triad and the plate used in the experiment. Probe: square

285    of squares, global match: circle of squares, local match: squares of circle. The three squares

286    on which the stimuli are placed illustrate the layout of the plate.

301     **Figure 2.**  Mean frequency of choosing local match. Error bars correspond to one standard

302     error of the mean.

309 **Figure 3.** Mean valence (left panel) and arousal (right panel) during pre-test and post-test for

310 happy and sad mood induction conditions. Error bars correspond to one standard error of the

311 mean.

323    **Table 1.** SAM Means (with standard deviations) for happiness, sadness, valence, and arousal

324    ratings across happy and sad mood before and after mood induction. Bold numbers in the post

325    test indicate the effect is in the expected direction. * indicates the effect was statistically

326    significant.

| | Happy mood | | Sad mood | |
|---|---|---|---|---|
| | pre-test | post-test | pre-test | post-test |
| Scales | M (SD) | M (SD) | M (SD) | M (SD) |
| Valence (SAM) | 7.00 (1.10) | **7.52** (1.54)* | 7.05 (.80) | **6.48** (1.40)* |
| Arousal(SAM) | 5.52 (2.02) | **6.24** (2.30) | 4.86 (1.71) | **3.67** (1.53)* |
| Happiness(DES) | 5.92 (1.45) | **6.56** (1.14)* | 5.97 (1.03) | **5.51** (1.07)* |
| Sadness(DES) | 2.33 (1.37) | **2.09** (1.22) | 2.13 (1.51) | **2.33** (1.24) |

**Lead contact**

Further information and requests for the resources should be directed to lead contact, Müge

Cavdan (Muege.Cavdan@psychol.uni-giessen.de).

**Data and code availability**

Data have been deposited on the Open Science Framework and will be publicly available upon

publication. Any additional information required to reanalyze the data reported in this work is

available from the lead contact upon request.

**References**

1. Isen, A.M., Niedenthal, P.M., and Cantor, N. (1992). An influence of positive affect on social categorization. Motiv. Emot. *16*, 65–78. 10.1007/BF00996487.

2. Isen, A.M., and Daubman, K.A. (1984). The influence of affect on categorization. J. Pers. Soc. Psychol. *47*, 1206–1217. 10.1037/0022-3514.47.6.1206.

3. Bless, H., Clore, G.L., Schwarz, N., Golisano, V., Rabe, C., and Wölk, M. (1996). Mood and the use of scripts: Does a happy mood really lead to mindlessness? J. Pers. Soc. Psychol. *71*, 665–679. 10.1037/0022-3514.71.4.665.

4. Bodenhausen, G.V., Kramer, G.P., and Süsser, K. (1994). Happiness and stereotypic thinking in social judgment. J. Pers. Soc. Psychol. *66*, 621–632. 10.1037/0022-3514.66.4.621.

5. Clore, G.L. (1992). Cognitive phenomenology: Feelings and the construction of judgment. In The construction of social judgments. (Lawrence Erlbaum Associates, Inc), pp. 133–163.

6. Schwarz, N., and Clore, G.L. (1983). Mood, misattribution, and judgments of well-being: Informative and directive functions of affective states. J. Pers. Soc. Psychol. *45*, 513–523. 10.1037/0022-3514.45.3.513.

7. Clore, G.L., Wyer Jr., R.S., Dienes, B., Gasper, K., Gohm, C., and Isbell, L. (2001). Affective feelings as feedback: Some cognitive consequences. In Theories of mood and cognition: A user's guidebook. (Lawrence Erlbaum Associates Publishers), pp. 27–62.

8. Gasper, K. (2004). Do you see what I see? Affect and visual information processing. Cogn. Emot. *18*, 405–421. 10.1080/02699930341000068.

9. Gasper, K., and Clore, G.L. (2002). Attending to the big picture: Mood and global versus local processing of visual information. Psychol. Sci. *13*, 34–40. 10.1111/1467-9280.00406.

10. Kimchi, R., and Palmer, S.E. (1982). Form and texture in hierarchically constructed
patterns. J. Exp. Psychol. Hum. Percept. Perform. *8*, 521–535. 10.1037/0096-1523.8.4.521.

11. Curby, K.M., Johnson, K.J., and Tyson, A. (2011). Face to face with emotion: Holistic face
processing is modulated by emotional state. Cogn. Emot. *26*, 93–102.
10.1080/02699931.2011.555752.

12. Derryberry, D., and Tucker, D.M. (1994). Motivating the focus of attention. In The heart's
eye: Emotional influences in perception and attention. (Academic Press), pp. 167–196.
10.1016/B978-0-12-410560-7.50014-4.

13. Easterbrook, J.A. (1959). The effect of emotion on cue utilization and the organization of
behavior. Psychol. Rev. *66*, 183–201. 10.1037/h0047707.

14. Fredrickson, B.L. (1998). What good are positive emotions? Rev. Gen. Psychol. *2*, 300–
319. 10.1037/1089-2680.2.3.300.

15. Hartikainen, K.M., Ogawa, K.H., and Knight, R.T. (2010). Trees over forest: unpleasant
stimuli compete for attention with global features. NeuroReport *21*, 344–348.
10.1097/WNR.0b013e328336eeb3.

16. Von Mühlenen, A., Bellaera, L., Singh, A., and Srinivasan, N. (2018). The effect of sadness
on global-local processing. Atten. Percept. Psychophys. *80*, 1072–1082. 10.3758/s13414-
018-1534-7.

17. Navon, D. (1977). Forest before trees: The precedence of global features in visual
perception. Cognit. Psychol. *9*, 353–383. 10.1016/0010-0285(77)90012-3.

18. Schyns, P.G., and Oliva, A. (1994). From blobs to boundary edges: Evidence for time- and
spatial-scale-dependent scene recognition. Psychol. Sci. *5*, 195–200. 10.1111/j.1467-
9280.1994.tb00500.x.

19. Lakatos, S., and Marks, L.E. (1999). Haptic form perception: Relative salience of local and
global features. Percept. Psychophys. *61*, 895–908. 10.3758/BF03206904.

20. Lederman, S.J., and Klatzky, R.L. (1987). Hand movements: A window into haptic object
recognition. Cognit. Psychol. *19*, 342–368. 10.1016/0010-0285(87)90008-9.

21. Lederman, S.J., and Klatzky, R.L. (1993). Extracting object properties through haptic
exploration. Acta Psychol. (Amst.) *84*, 29–40. 10.1016/0001-6918(93)90070-8.

22. Kinchla, R.A., and Wolfe, J.M. (1979). The order of visual processing: "Top-down,"
"bottom-up," or "middle-out." Percept. Psychophys. *25*, 225–231. 10.3758/BF03202991.

23. Tortelli, C., Senna, I., Binda, P., and Ernst, M.O. (2023). Development of local-global
preference in vision and haptics. J. Vis. *23*, 6. 10.1167/jov.23.4.6.

24. Blanca, M.J., and López-Montiel, G. (2009). Hemispheric differences for global and local
processing: Effect of stimulus size and sparsity. Span. J. Psychol. *12*, 21–31.
10.1017/S113874160000144X.

25. Rezvani, Z., Katanforoush, A., and Pouretemad, H. (2020). Global precedence changes by
environment: A systematic review and meta-analysis on effect of perceptual field variables
on global-local visual processing. Atten. Percept. Psychophys. *82*, 2348–2359.
10.3758/s13414-020-01997-1.

26. Kimchi, R. (1992). Primacy of wholistic processing and global/local paradigm: A critical
review. Psychol. Bull. *112*, 24–38. 10.1037/0033-2909.112.1.24.

27. Ji, L.-J., Yap, S., Best, M.W., and McGeorge, K. (2019). Global processing makes people
happier than local processing. Front. Psychol. *10*, 670. 10.3389/fpsyg.2019.00670.

28. Srinivasan, N., and Gupta, R. (2011). Rapid communication: Global–local processing
affects recognition of distractor emotional faces. Q. J. Exp. Psychol. *64*, 425–433.
10.1080/17470218.2011.552981.

29. Srinivasan, N., and Hanif, A. (2010). Global-happy and local-sad: Perceptual processing
affects emotion identification. Cogn. Emot. *24*, 1062–1069. 10.1080/02699930903101103.

30. Shapiro, D. (1965). Neurotic styles. (Basic Books).

31. Witkin, H.A., Goodenough, D.R., and Oltman, P.K. (1979). Psychological differentiation:
Current status. J. Pers. Soc. Psychol. *37*, 1127–1145. 10.1037/0022-3514.37.7.1127.

32. Yovel, I., Revelle, W., and Mineka, S. (2005). Who sees trees before forest?: The
obsessive-compulsive style of visual attention. Psychol. Sci. *16*, 123–129. 10.1111/j.0956-
7976.2005.00792.x.

33. Faul, F., Erdfelder, E., Lang, A.-G., and Buchner, A. (2007). G*Power 3: A flexible
statistical power analysis program for the social, behavioral, and biomedical sciences.
Behav. Res. Methods *39*, 175–191. 10.3758/BF03193146.

34. World medical association declaration of helsinki: Ethical principles for medical research
involving human subjects (2013). JAMA *310*, 2191. 10.1001/jama.2013.281053.

35. Ackerley, R., Aimonetti, J.-M., and Ribot-Ciscar, E. (2017). Emotions alter muscle
proprioceptive coding of movements in humans. Sci. Rep. *7*, 8465. 10.1038/s41598-017-
08721-4.

36. Bradley, M.M., and Lang, P.J. (1994). Measuring emotion: The Self-Assessment Manikin
and the semantic differential. J. Behav. Ther. Exp. Psychiatry *25*, 49–59. 10.1016/0005-
7916(94)90063-9.

37. Izard, C.E. (1977). Differential emotions theory. In Human Emotions (Springer US), pp.
43–66. 10.1007/978-1-4899-2209-0_3.

38. Renaud, D., and Unz, D. (2006). Die M-DAS - eine modifizierte version der differentiellen
affekt skala zur erfassung von emotionen bei der mediennutzung. Z. Für Medien. *18*, 70–
75. 10.1026/1617-6383.18.2.70.

39. Johnson, K.O., and Phillips, J.R. (1981). Tactile spatial resolution. I. Two-point
discrimination, gap detection, grating resolution, and letter recognition. J. Neurophysiol.
*46*, 1177–1192. 10.1152/jn.1981.46.6.1177.

450