Title: Happiness promotes global processing in haptic perception

Müge Cavdan[§,†*1], Aycan Kapucu[‡], Katja Doerschner[§,†+], & Knut Drewing[§+]

[§]Experimental Psychology Department, Justus Liebig University Giessen; Otto-Behaghel Strasse 10F, 35394, Giessen, Germany

[†]Center for Mind, Brain, and Behavior (CMBB), Philipps University Marburg and Justus Liebig University Giessen; Hans-Meerwein-Strasse 6, 35032, Marburg, Germany

[‡]Psychology Department, Ege University, Izmir, Türkiye

*Correspondence : Muege.Cavdan@psychol.uni-giessen.de ; X Handle – @mugecavdan

[+]Equal contribution

---

[1] lead contact: Muege.Cavdan@psychol.uni-giessen.de

Abstract

Happy and sad mood promote global and local visual processing, respectively. However, it is unclear whether mood also affects the processing level in haptic perception. Here, we used classical music to induce happy and sad mood in blindfolded participants before they scanned 3D-printed configurations with their fingers. Additionally, we included a neutral group who did not listen to any music. Global shapes were triangles, circles, or squares (33mm) composed of smaller local relief shapes (3 mm): either triangles, circles, or squares. Participants explored a probe stimulus with identical local and global shapes, and two comparison stimuli, matching the probe in local, or global shape. They reported which comparison stimulus appeared more similar to the probe. In the 'sad' group, participants chose the locally matching comparison more frequently than in the 'happy' and 'neutral' groups, revealing that sad mood promotes local processing also in touch. Overall, participants chose the globally matching comparison more often, suggesting that global processing is more prominent in touch than previously assumed.

Keywords: mood, level of processing, haptics, shape perception

Introduction

Feelings modulate how we perceive and interact with the world. Mood modulates cognitive processes and social behavior, e.g., happy feelings promote reliance on accessible general information such as stereotypes (1,2) and scripts (3) while sad feelings have been associated with focusing on details, avoiding stereotypes, which often leads to an improved accuracy of judgements (4). One explanation for this shift of focus is offered by the affect-as-information hypothesis (5,6). According to this theory, mood is a signal that can be used to guide behavior: happy mood signals a low-threat situation that affords global processing whereas sad mood may signal that a situation is potentially threatening and therefor require more detailed processing. An alternative explanation for the association between mood and level of processing suggests that motivation may act as a mediator. According to this idea detailed processing is effortful and may result in a reduction of mood (7). Consequently, when happy, people may be less motivated to perform effortful, detailed processing, in order to maintain their positive mood.

Interestingly, this shift from general, global to detailed, local processing as the mood shifts from happy to sad, has not only been observed in social and cognitive studies but also emerges robustly in visual shape perception (3,8,9). For example, in an experiment by Gasper and Clore (9), 'happy' and 'sad' participants indicated the similarity of a given shape (triangle or square) that consisted of local elements (triangle or square, also see Kimchi & Palmer (10) to two figures that either matched the global or the local shape of the stimulus. Their results showed that individuals in the sad mood condition more often classified shapes based on local features compared to 'happy'. In line with this, Curby and colleagues (11) showed that negative mood decreased the holistic processing of faces. The attentional narrowing of attention when mood is low (8,12–16) is interesting, as it appears to 'override' the default holistic processing visual shape and scene perception (17,18).

In general, the haptic system applies more weight to local features than global ones

when judging shapes (19). Thus, it is not clear how and in what direction changes in mood

would modulate the haptic perception of shape. To address these questions, we investigated

how positive and negative mood influences the level of haptic shape processing. Upon mood

induction, blindfolded participants scanned a haptic shape stimulus with their fingers and

indicated which one of the two comparison stimuli was more similar to the probe. Comparisons

matched the probe in either global or local shape. A neutral condition with no mood induction

(i.e., no music) was included to assess baseline haptic processing. Our working hypothesis was

that happy mood would promote global processing while sad mood would promote a focus on

local shape features. Our results provide a deeper understanding of to what extent mood-

induced changes in attentional focus might be a general perceptual phenomenon.

Methods

Participants

The required sample size was calculated based on a large effect (Cohen's $d = 0.8$), power of

80%, and alpha 5%. The projected sample size was 42 for one-sided between-subjects t-test

(G*Power(20)). Accordingly, 42 participants (9 males, $M_{age} = 22.8$, age range: 18-30) were

randomly assigned to one of the two mood conditions (happy or sad). An additional 21

participants were recruited to the control condition (7 males, $M_{age} = 24.14$, age range: 20-34).

Eligibility criteria for the study required that participants have no diagnosed mental disorders,

as such conditions might influence processing levels (21) and introduce potential confounds in

the experiment. They were recruited through the university email system and compensated with

8 €/hour or course credit. None of the participants reported sensory, motor, or cutaneous

impairments. The two-point discrimination threshold at the index finger of the dominant hand

was 3 mm or better. The study was ethically approved by a local ethics committee LEK FB06

(2017-0034) in accordance with the Declaration of Helsinki(22) excluding the preregistration.

Informed consents were obtained prior to the experiment.

Stimuli and materials

Haptic Stimuli. Haptic stimuli were modeled using the OpenSCAD software and 3D printed

(Objet30Pro, StratasysLtd., USA) with model photopolymer material (VeroClear) at a

resolution of $600 \times 600 \times 1.600$ dpi (x, y, and z axes respectively). As previous research has

shown that the haptic system weighs local shape features more than global features(19), we

first conducted a pilot experiment where we investigated the detection thresholds of a small

triangle, a circle, and a square in order to determine the shape size that would allow us to reduce

the natural saliency of the local features relative to the global ones in the main experiment.

Stimuli were two-dimensional (2D) triangles, squares, and circles ranging from 1 mm to 10

mm (size: diameter of a circle, one side of a square, and triangle base; shapes embossed with a

height of 1 mm on a plane plate). Blindfolded participants touched the shapes and named the

shape they perceive. For each shape mean value, the sizes where a person recognized the shapes

correctly were calculated. These values were taken as a threshold and then used as the local

shape sizes for the later stimuli production in the main experiment.

Probe   Global match   Local match

Figure 1. Illustration of an example triad and the plate used in the experiment. Probe: square

of squares, global match: circle of squares, local match: squares of circle. The three squares

on which the stimuli are placed illustrate the layout of the plate.

Nine stimuli consisting of local and global features modelled using OpenSCAD

software and 3D printed at a high resolution as in the pilot experiment. The global shapes were

printed on a 40 mm (x-axis) × 40 mm (y-axis) long and 5 mm (z-axis) thick base. The stimuli

were mounted on a 3D printed plate which has three gaps. This ensured that the stimuli

remained fixed on the table during the exploration. Local shape sizes were 3 mm (both x and

y-axes) and global shapes size was 33 (both x and y-axes) for square, circle, and triangle with

the height of 1 mm (z-axes). The distance between each adjacent local shape was kept constant

at 1 mm. The shapes consisted of a circle of circles, a triangle of triangles, and a square of

squares as probes while comparison stimuli were a circle of triangles, a square of triangles, a

triangle of squares, circle of squares, a square of circles, and a triangle of circles. The stimuli

were combined with each other to make triads (six triads in total). Each triad consisted of one

probe (i.e., standard) and two comparison stimuli (see Figure 1 for an example). Probes had

the same global and local shapes such as square of squares. Comparison stimuli on the other

hand, matched either local or global shape of the probe such as square of circles (global match)

and circle of squares (local match).

Mood induction stimuli and manipulation check. Classical music pieces from Ackerley et al.

(23) were tested in a pilot experiment in order to affirm their effectiveness in our population[2].

The pieces that evoked strongest happiness and sadness were selected for mood induction in

---

[2] Total of 20 participants participated in the between-subjects pilot experiment. Participants were randomly assigned to either happy or sad condition. Participants in the happy condition listened to the happy classical music pieces while participants in the sad condition listened to the sad classical music. Although more than one classical piece was presented in the experiment, we will be focusing on the pieces subject to the current article. Upon listening to either *Carmen Suite Nr.1 : Les Toréadors* or *Kol Nidrei* participants indicated the strongest level of each emotion they experienced while listening to the music piece on a 9-point scale (0: not at all, 8: extremely). The following emotion words were rated in a random order; happiness: *happy*, *joyful*, *amused*; sad: *sad*, *downhearted*, and *blue*. The average scores across happiness and sadness were calculated. Two-independent samples t-test were used to test the efficacy of the classical music pieces. For happiness, the happy music piece yielded a higher rating than the sad piece, $t(18) = 4.65$, $p < .01$, $d = 2.08$. For sadness, the sad music piece yielded a higher rating than the happy piece, $t(18) = -5.40$, $p < .01$, $d = 2.41$. These results clearly show the efficacy of both classical music pieces in inducing the desired target moods.

the current study. *Kol Nidrei* by *Max Bruch (1938-1920)* and *Carmen Suite Nr. 1: Les*

Toréadors by *Georges Bizet (1838-1875)* were used to induce sad and happy mood,

respectively.

In order to measure the efficacy of mood induction, two different scales were used

before and after the mood induction, the Self-Assessment Manikin (SAM (24) and the

Differential Emotions Scale – DES (25,26)). The SAM measured participants' valence and

arousal levels on a 9-point pictorial depiction (*valence*: 1= very negative to 9= very positive;

arousal: 1= very calm to 9= very excited). Participants were asked to rate the SAM based on

their emotional state. The DES is widely used in psychological research and a reliable method

to assess emotions. It consists of several subscales with various adjectives. We selected the

sadness and happiness subscales with their corresponding three adjectives per subscale;

happiness: *happy, joyful, amused*; sad: *sad, downhearted*, and *blue*. The adjectives were

presented with a 9-point scale (0: *not at all*, 8: *extremely*). Participants rated these adjectives

based on how much they felt each emotion.

Procedure

A between-subjects design with two moods – happy and sad – and a neutral condition

(no mood manipulation) was used. Participants were randomly assigned to either the happy,

the sad mood, and neutral conditions. Upon arrival in the lab, a participant received information

about the experiment and signed an informed consent. This was followed by a two-point touch

discrimination threshold test to measure any tactile deficiencies (27). Next, participants

responded to the mood questions of SAM and DES, presented on a monitor screen, using a

numpad. The presentation order of the questionnaires was randomized. After completion of the

questionnaires, a classical music piece, either happy or sad - depending on the condition, was

played while no music was played in the neutral condition. After this, SAM and DES

questionnaires were completed a second time in happy and sad conditions. Participants then

performed the Kimchi-Palmer task (10), as described next. In the neutral condition, participants

completed the SAM and DES questionnaires a second time after approximately 15-20 minutes

had passed.

On each trial, a probe (i.e., same global and local shape) was always presented in the

first gap of the mounted plate (Figure 1). The two comparison shapes (one matching the global

shape of the probe and the other matching the local shape) were presented in the next two gaps

of the mounted plate. The presentation order of the second and third gaps were

counterbalanced. Blindfolded participants were asked to explore each stimulus of the triad

sequentially (from first gap to third) using lateral movements with the index finger of the

dominant hand. Participants were able to re-explore the probe and the comparison stimuli as

many times as wanted. Their task was to indicate which of the two comparison stimuli was

more similar to the probe.

Each of the six triads was repeated 6 times which resulted in 36 trials in total. The

overall procedure lasted 30-45 minutes in happy and sad music conditions while it lasted 20-

30 minutes in the neutral condition.

Data analysis

All analyses were conducted in SPSS 25 software (SPSS inc.) and figures are plotted in

MATLAB R2022b (MathWorks inc.).

Cronbach's alpha was used to calculate reliability per happiness and sadness subscales (DES)

for pre- and post-induction in happy and sad conditions and pre- and post-Kimchi-Palmer task

in neutral condition. All eight analyses showed good to excellent consistency (Cronbach's $a =$

.70 to .96) allowing for averaging across adjectives. Next, we calculated the average values

across the three adjectives per mood (i.e., happiness: happy, joy, and amused; sadness: sad,

downhearted, and blue) which were then submitted to an ANOVA to compare happy and sad

conditions. Additionally, first mood measurements were submitted to an ANOVA to test

whether the baseline mood was comparable across conditions.

For the Kimchi-Palmer task, the frequency of selecting the local comparison shape over

the global one was calculated from the raw data of each participant. Selecting the local match

indicated that the participant relied on local shape processing. The percentage of local

processing across all trials was calculated per participant in order to conduct a one-way analysis

of variances (ANOVA) to examine the effect of condition on the level of processing. This was

followed by Bonferroni-corrected multiple comparisons to further investigate group

differences.

Greenhouse-Geisser correction was applied when the sphericity assumption was violated.

Similarly, degrees of freedom were adjusted when Levene's test indicated unequal variances.

Results

Global and local processing

A one-way ANOVA showed a significant main effect of condition: $F(2, 62) = 7.92$, $p < .001$,

$\eta^2 = .21$. Specifically, local processing was significantly less frequent after listening to the

happy music piece, than after listing to the sad music piece ($M_{diff} = -11.24$, $SE = 4.08$, $p = .023$)

(Figure 2). Overall, the global match was selected six times more often than the local one

(Figure 2). Also, no music condition ($M = 3.44$, $SE = 1.05$) yielded similar local processing

frequency as the happy music piece ($M_{diff} = 4.50$, $SE = 4.08$, $p = .82$) while it resulted in less

frequent than sad music condition ($M_{diff} = -15.74$, $SE = 4.08$, $p < .001$).

Next, we tested whether there was an effect of global-local shape combination in level

of processing. To this end, we conducted a repeated-measures ANOVA with the variable shape

triad (6 levels). The main effect of shape triad was not statistically significant, $F(4.05, 251.31)$

$= 1.02, p = .40, \eta_p{}^2 = .02$, showing that the percentage of local match choice was not dependent

on a specific shape-combination. Finally, to test the possible effects of presentation orders on

the level of processing, 6 paired-samples t-tests were conducted (i.e., one per shape-

combination). None of the t-test were statistically significant showing that there was no effect

of presentation order on the percentage of local match (all $p > .05$).

Figure 2. Mean frequency of choosing local math. Error bars correspond to one standard

error of the mean.

Mood manipulation check

The efficacy of mood induction was examined using 4 mixed-design ANOVAs to test the effect

of music piece (levels: *happy* and *sad*) and time point (levels: *pre-induction* and *post-induction*)

separately on mean ratings for Self-assessment Manikin (SAM) valence, SAM arousal,

Differential Emotion Scale (DES) happiness, and DES sadness (see Table 1 for the summary

of the results).

Additionally, we tested whether the individuals in all conditions had a comparable

baseline (i.e., before mood induction) mood with one-way ANOVA on SAM valence, SAM

arousal, and DES happiness and sadness subscales. We expected all conditions to have a

comparable baseline.

SAM. As expected, listening to the happy music piece increased **valence** towards

positive, while listening to the sad music piece decreased it: The main effects of mood and

time point were not statistically significant for valence, $F(1,40) = 2.21$, $p = .15$, $\eta_p^2 = .05$,

$F(1,40) = 0.17$, $p = .90$, $\eta_p^2 = 00$, respectively. However, the interaction between time and mood

was statistically significant, $F(1,40) = 8.94$, $p < .01$, $\eta_p^2 = .18$. Follow-up analyses showed that,

in the happy mood condition, participants were more positive in the post-test compared to the

pre-test, $t(20) = -2.33$, $p = .015$ (one-sided). Also expectedly, after listening to the sad musical

piece, participants reported to be in a more negative mood than during the pre-test phase, $t(20)$

$= 1.98$, $p = .03$ (one-sided, Figure 3).

The sad music piece decreased **arousal**, as expected: The main effect of mood, $F(1,40)$

$= 9.81$, $\eta_p^2 = .20$ and the interaction between time point and mood, $F(1,40) = 11.24$, $\eta_p^2 = .22$,

were statistically significant (both $p < .01$). However, the main effect of time point was not

statistically significant for arousal, $F(1,40) = .70$, $p = .41$, $\eta p^2 = .02$. Follow-up analyses showed

that for happy mood, there was no significant arousal difference between pre-test and post-test,

$t(20) = -1.92$, $p = .07$. As expected, for sad mood, arousal decreased on post-test compared to

pre-test, $t(20) = 2.78$, $p < .01$ (Figure 3).

Finally and expectedly, prior to listening to the music, there was no difference in

valence, $F(2,60) = .54$, $p = .54$, $\eta_p^2 = .01$ and arousal, $F(2,60) = 2.31$, $p = .11$, $\eta_p^2 = .07$ across

conditions.

Figure 3. Mean valence (left panel) and arousal (right panel) during pre-test and post-test for

happy and sad mood induction conditions. Error bars correspond to one standard error of the

mean.

DES. Participants in the happy mood were happier upon mood induction. However,

participants in sad mood did not feel significantly sadder after manipulation which will be

discussed further.

For the happiness subscale, neither the main effect of time point, $F(1, 40) = .28$, $\eta_p^2 =$

.01, nor the main effect of mood, $F(1, 40) = 2.34$, $\eta_p^2 = .06$ were statistically significant.

However, the interaction between time point and mood was statistically significant $F(1, 40) =$

11.04, $p < .01$, $\eta_p^2 = .06$. In the happy mood condition happiness increased between pre- and

post-test, whereas in the sad mood condition happiness decreased between pre- and post-test:

$t(20) = -3.11$, $p < .01$, $t(20) = 1.78$, $p < .05$ (both one-sided).

For the sadness subscale, although people in the sad mood felt less happy and more sad,

this did not reach significance: The main effect of time point: $F(1, 40) = .01$, $\eta_p^2 = .00$, mood:

$F(1, 40) = .00$, $\eta_p{}^2 = .00$ and their interaction: $F(1, 40) = 1.74$, $\eta_p{}^2 = .04$ (all $p > 0.05$) were not

statistically significant.

Additionally, prior to listening to the music pieces, individuals had comparable

happiness, $F(2, 60) = .30$, $p = .75$, $\eta_p{}^2 = .01$, and sadness, $F(2, 60) = .15$, $p = .86$, $\eta_p{}^2 = .01$

levels.

Table 1. SAM Means (with standard deviations) for happiness, sadness, valence, and arousal

ratings across happy and sad mood before and after mood induction. Bold numbers in the post

test indicate the effect is in the expected direction. * indicates the effect was statistically

significant.

	Happy mood		Sad mood	
	pre-test	post-test	pre-test	post-test
Scales	M (SD)	M (SD)	M (SD)	M (SD)
Valence (SAM)	7.00 (1.10)	**7.52** (1.54)*	7.05 (.80)	**6.48** (1.40)*
Arousal(SAM)	5.52 (2.02)	**6.24** (2.30)	4.86 (1.71)	**3.67** (1.53)*
Happiness(DES)	5.92 (1.45)	**6.56** (1.14)*	5.97 (1.03)	**5.51** (1.07)*
Sadness(DES)	2.33 (1.37)	**2.09** (1.22)	2.13 (1.51)	**2.33** (1.24)

Discussion

When we look at a scene, we have the ability to focus on a single element or to take in the

entire shape formed by these individual elements. Gasper and Clore (9) suggest that whether

we perceive the entire shape or the single element depends on our mood. Namely, while happy

mood fosters the entire shape view (i.e., global processing), sad mood fosters the single element

view (i.e., local processing). Here, we showed that this effect is not limited to vision but a

broader mechanism that operates at the perceptual level, and is therefore also present in the

haptic modality. Individuals in the happy mood condition matched the probe with the global

shape more often than individuals in the sad mood condition. Interestingly, individuals,

independent of the mood condition, more often matched the global target to the probe – the

global match was selected overall six times more often than the local one. This overall global

perception preference is interesting considering that typically people focus on local details

when exploring objects with their hands (19,28,29). However, the objects used in these earlier

studies were larger than a typical hand, which might have pushed individuals to do sequential

exploration, which might have resulted in increased focus on local detail. Thus, the current

global precedence finding might be due to the overall size of the global form (30,31), as well

as the density (30–32) of the local features (see (33) for a meta-analysis on the environmental

factors moderating global precedence. Although local-global focus preferences did not receive

the same attention in haptic perception as in vision, a recent haptic study reported a similar

global preference as we observed in our experiment: the smaller the local shape the more global

processing was observed (31). Therefore, the level of processing seems to critically depend on

the number and relative size of the elements (33,34). It would be interesting, in future

experiments, to identify the local-global relative size relationship where the perceptual focus

switches.

The general observed global preference suggests that the most available and easily

accessible information was the global form of our stimuli. The usage of global information,

however, was modulated by the mood: Individuals in the happy mood condition selected the

global match to the probe more frequently than individuals in the sad mood condition, and this

effect was independent of the presentation order or the geometrical features (circle, triangle,

square). Our finding is in line with the levels-of-focus hypothesis arguing that individuals in

happy mood condition are more likely to perceive the easily accessible knowledge resulting in

broader attentional processing (9). In contrast, sad mood would result in increased detailed

(local) processing.

Not only mood influences the processing of the stimuli but also using a certain type of

processing (global or local) might facilitate a mood (35). This has been also observed for face

identification – global processing facilitated happy face identification while local processing

facilitated sad face identification (36,37). Future research should investigate the possible

bidirectionality between mood and level of processing in haptics.

Our work might have potential clinical relevance. For example, obsessive compulsive

individuals tend to focus on details (38,39), a cognitive style, that has been associated with

local interference of small details when the task was the identification of global information

(40). Future studies should test whether individuals with obsessive compulsive cognitive style

use more local processing compared to a control group also in haptic tasks. Insights from such

studies potentially yield the development of novel diagnosis tools for this group of patients.

In conclusion, individuals in happy mood are more likely to *feel* the entire shape. Using

a stimulus set consisting of local and global features and a haptic shape perception task, we

found that happy mood fosters global processing compared to sad mood. To the best of our

knowledge, this is the first study that extended the levels-of-processing effect found in vision

to the haptic domain. Unlike previous research in haptic perception, we also found global

precedence when people perceive different geometrical shapes. Therefore, more research is

needed to understand the different components of global and local processing in human haptic

perception.

Author contributions

Conceptualization: M.C., A.K., K.D., K.Dr; Methodology: M.C., K.Dr.; Investigation: M.C.;

Data curation: M.C.; Formal analysis: M.C., A.K.; Visualization: M.C., K.D.; Writing—

original draft: M.C.; Writing—reviewing and editing: M.C., A.K., K.D., K.Dr; Supervision:

M.C., A.K., K.D., K.Dr.; Funding Acquisition: K.D., K.Dr.; Project Administration: M.C.,

K.Dr.

Acknowledgement

Authors thank Hatice Dokumaci for the haptic stimuli and data collection, Yeter Nazlıcan Kara

for the data collection, Rochelle Ackerley for sharing the classical music pieces. This research

was supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) –

project number 222641018 – SFB/TRR 135 (A5 and B8) and EU Marie Curie Initial Training

Network "DyVito" (H2020-ITN, Grant Agreement: 765121). K.Dr and M.C. were supported

by DFG-project no. 502774891-ORA project "UNTOUCH", K.Do was supported by Research

Cluster "The Adaptive Mind", funded by the Excellence Program of the Hessian Ministry of

Higher Education, Science, Research and the Arts. Artificial intelligence.

Funding

This research was supported by Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) – project number 222641018 – SFB/TRR 135 (A5 and B8) and EU Marie Curie

Initial Training Network "DyVito" (H2020-ITN, Grant Agreement: 765121). K.Dr and M.C.

were supported by DFG-project no. 502774891-ORA project "UNTOUCH", K.Do was

supported by Research Cluster "The Adaptive Mind", funded by the Excellence Program of

the Hessian Ministry of Higher Education, Science, Research and the Arts.

Lead contact

Further information and requests for the resources should be directed to lead contact, Müge

Cavdan (Muege.Cavdan@psychol.uni-giessen.de).

Data and code availability

Data have been deposited on the Open Science Framework and will be publicly available upon

publication. Any additional information required to reanalyze the data reported in this work is

available from the lead contact upon request.

References

1. Isen AM, Niedenthal PM, Cantor N. An influence of positive affect on social categorization. Motiv Emot. 1992 Mar;16(1):65–78.

2. Isen AM, Daubman KA. The influence of affect on categorization. J Pers Soc Psychol. 1984 Dec;47(6):1206–17.

3. Bless H, Clore GL, Schwarz N, Golisano V, Rabe C, Wölk M. Mood and the use of scripts: Does a happy mood really lead to mindlessness? J Pers Soc Psychol. 1996;71(4):665–79.

4. Bodenhausen GV, Kramer GP, Süsser K. Happiness and stereotypic thinking in social judgment. J Pers Soc Psychol. 1994;66(4):621–32.

5. Clore GL. Cognitive phenomenology: Feelings and the construction of judgment. In: The construction of social judgments. Hillsdale, NJ, US: Lawrence Erlbaum Associates, Inc; 1992. p. 133–63.

6. Schwarz N, Clore GL. Mood, misattribution, and judgments of well-being: Informative and directive functions of affective states. J Pers Soc Psychol. 1983 Sep;45(3):513–23.

7. Clore GL, Wyer Jr. RS, Dienes B, Gasper K, Gohm C, Isbell L. Affective feelings as feedback: Some cognitive consequences. In: Theories of mood and cognition: A user's guidebook. Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers; 2001. p. 27–62.

8. Gasper K. Do you see what I see? Affect and visual information processing. Cogn Emot. 2004;18(3):405–21.

9. Gasper K, Clore GL. Attending to the big picture: Mood and global versus local processing of visual information. Psychol Sci. 2002 Jan;13(1):34–40.

10. Kimchi R, Palmer SE. Form and texture in hierarchically constructed patterns. J Exp Psychol Hum Percept Perform. 1982;8(4):521–35.

11. Curby KM, Johnson KJ, Tyson A. Face to face with emotion: Holistic face processing is
modulated by emotional state. Cogn Emot. 2011 May;26(1):93–102.

12. Derryberry D, Tucker DM. Motivating the focus of attention. In: The heart's eye:
Emotional influences in perception and attention. San Diego, CA, US: Academic Press;
1994. p. 167–96.

13. Easterbrook JA. The effect of emotion on cue utilization and the organization of
behavior. Psychol Rev. 1959;66(3):183–201.

14. Fredrickson BL. What good are positive emotions? Rev Gen Psychol. 1998
Sep;2(3):300–19.

15. Hartikainen KM, Ogawa KH, Knight RT. Trees over forest: unpleasant stimuli compete
for attention with global features. NeuroReport. 2010 Mar 31;21(5):344–8.

16. Von Mühlenen A, Bellaera L, Singh A, Srinivasan N. The effect of sadness on global-
local processing. Atten Percept Psychophys. 2018 Jul;80(5):1072–82.

17. Navon D. Forest before trees: The precedence of global features in visual perception.
Cognit Psychol. 1977;9(3):353–83.

18. Schyns PG, Oliva A. From blobs to boundary edges: Evidence for time- and spatial-
scale-dependent scene recognition. Psychol Sci. 1994 Jul;5(4):195–200.

19. Lakatos S, Marks LE. Haptic form perception: Relative salience of local and global
features. Percept Psychophys. 1999 Jan;61(5):895–908.

20. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. Behav Res
Methods. 2007 May;39(2):175–91.

21. De Fockert JW, Cooper A. Higher levels of depression are associated with reduced global
bias in visual processing. Cogn Emot. 2014 Apr 3;28(3):541–9.

22. World medical association declaration of helsinki: Ethical principles for medical research
involving human subjects. JAMA. 2013 Nov 27;310(20):2191.

23. Ackerley R, Aimonetti JM, Ribot-Ciscar E. Emotions alter muscle proprioceptive coding
of movements in humans. Sci Rep. 2017 Aug 16;7(1):8465.

24. Bradley MM, Lang PJ. Measuring emotion: The Self-Assessment Manikin and the
semantic differential. J Behav Ther Exp Psychiatry. 1994;25(1):49–59.

25. Izard CE. Differential emotions theory. In: Human Emotions [Internet]. Boston, MA:
Springer US; 1977 [cited 2023 Aug 21]. p. 43–66. Available from:
http://link.springer.com/10.1007/978-1-4899-2209-0_3

26. Renaud D, Unz D. Die M-DAS - eine modifizierte version der differentiellen affekt skala
zur erfassung von emotionen bei der mediennutzung. Z Für Medien. 2006 Apr;18(2):70–
5.

27. Johnson KO, Phillips JR. Tactile spatial resolution. I. Two-point discrimination, gap
detection, grating resolution, and letter recognition. J Neurophysiol. 1981 Dec
1;46(6):1177–92.

28. Lederman SJ, Klatzky RL. Hand movements: A window into haptic object recognition.
Cognit Psychol. 1987 Jul;19(3):342–68.

29. Lederman SJ, Klatzky RL. Extracting object properties through haptic exploration. Acta
Psychol (Amst). 1993 Oct;84(1):29–40.

30. Kinchla RA, Wolfe JM. The order of visual processing: "Top-down," "bottom-up," or
"middle-out." Percept Psychophys. 1979 May;25(3):225–31.

31. Tortelli C, Senna I, Binda P, Ernst MO. Development of local-global preference in vision
and haptics. J Vis. 2023 Apr 25;23(4):6.

32. Blanca MJ, López-Montiel G. Hemispheric differences for global and local processing:
Effect of stimulus size and sparsity. Span J Psychol. 2009 May;12(1):21–31.

33. Rezvani Z, Katanforoush A, Pouretemad H. Global precedence changes by environment:
A systematic review and meta-analysis on effect of perceptual field variables on global-
local visual processing. Atten Percept Psychophys. 2020 Jul;82(5):2348–59.

34. Kimchi R. Primacy of wholistic processing and global/local paradigm: A critical review.
Psychol Bull. 1992;112(1):24–38.

35. Ji LJ, Yap S, Best MW, McGeorge K. Global processing makes people happier than local
processing. Front Psychol. 2019 Mar 26;10:670.

36. Srinivasan N, Gupta R. Rapid communication: Global–local processing affects
recognition of distractor emotional faces. Q J Exp Psychol. 2011 Mar;64(3):425–33.

37. Srinivasan N, Hanif A. Global-happy and local-sad: Perceptual processing affects
emotion identification. Cogn Emot. 2010 Sep;24(6):1062–9.

38. Shapiro D. Neurotic styles. Oxford, England: Basic Books; 1965. (Neurotic styles.).

39. Witkin HA, Goodenough DR, Oltman PK. Psychological differentiation: Current status. J
Pers Soc Psychol. 1979 Jul;37(7):1127–45.

40. Yovel I, Revelle W, Mineka S. Who sees trees before forest?: The obsessive-compulsive
style of visual attention. Psychol Sci. 2005 Feb;16(2):123–9.

457