Efficacy and mechanism of actions of cipargamin as an antibabesial drug candidate

Authors

Hang Li[1#], Shengwei Ji[1,2#], Nanang R. Ariefta[1], Eloiza May S. Galon[3], Shimaa AES El-Sayed[1,4],

Lijun Jia[2], Yoshifumi Nishikawa[1], Mingming Liu[1,5]*, Xuenan Xuan[1]*

Affiliations

[1]National Research Center for Protozoan Diseases, Obihiro University of Agriculture

Veterinary Medicine, Inada-cho, Obihiro, Japan

[2]Department of Veterinary Medicine, Agriculture College of Yanbian University, Yanji, China

[3]College of Veterinary Medicine and Biomedical Sciences, Cavite State University, Indang,

Philippines

[4]Department of Biochemistry and Molecular Biology, Faculty of Veterinary Medicine,

Mansoura University, Egypt

[5]School of Basic Medicine, Hubei University of Arts and Science, Xiangyang, China

* Correspondence: Mingming Liu; Xuenan Xuan

Address: Longzhong Road No.296, Xiangyang, China; Nishi 11 Minami 39 Chome, Obihiro,

Hokkaido

Telephone number: +8618904416800; +8109079036369

E-mail address: lmm_2010@hotmail.com; gen@obihiro.ac.jp

Competing interest: The authors have declared that no conflict of interest exists.

# These authors contributed equally to this work.

Abstract

Babesiosis is a disease brought on by intraerythrocytic parasites of the genus *Babesia*. Current

chemotherapies are accompanied by side effects and parasite relapse. Therefore, it is crucial to

develop highly effective drugs against *Babesia*. Cipargamin (CIP) has shown inhibition against

apicomplexan parasites, mainly *Plasmodium* and *Toxoplasma*. This study evaluated the

growth-inhibiting properties of CIP against *Babesia* spp. and investigated the mechanism of

CIP on *B. gibsoni*. The half inhibitory concentration (IC_{50}) values of CIP against the *in vitro*

growth of *B. bovis* and *B. gibsoni* were 20.2 ± 1.4 nM and 69.4 ± 2.2 nM, respectively. CIP

significantly inhibited the growth of *B. microti* and *B. rodhaini in vivo.* Resistance was

conferred by L921V and L921I mutations in *Bg*ATP4, which reduced the sensitivity to CIP by

6.1- and 12.8-fold. An *in silico* investigation revealed reductions in affinity for CIP binding to

*Bg*ATP4^{L921V} and *Bg*ATP4^{L921I} compared to *Bg*ATP4^{WT}. In this study, we characterized the

efficacy of CIP against *Babesia* spp. by investigating the mechanistic basis for the resistance

to CIP conferred by mutations in the *Bg*ATP4. Our findings present a promising starting point

for the establishment of new therapeutic interventions against *Babesia* infection.

Keywords: Cipargamin; *Babesia* spp.; drug resistance; mutation; ATP4

Introduction

Babesia is an apicomplexan tick-transmitted hemoparasite, which not only impacts the livestock economy but also causes an emerging disease in humans (Jalovecka et al., 2019). *Babesia* is a global pathogen that is more prevalent in certain regions such as Asia, Europe, and North America. However, as climate change brings with it higher temperatures and humidity, ticks and their reservoir hosts are anticipated to expand northward for survival and activity (Drews et al., 2023). The disease it causes is known as babesiosis, commonly characterized by fever and hemolytic anemia, but chronic infections are asymptomatic (Almazán et al., 2022). In acute cases, up to 20% of patients die, particularly individuals with compromised immune systems (Krause, 2019).

Currently, the most common drugs for the treatment of human babesiosis include a combination of atovaquone (ATQ) plus azithromycin (AZM) or clindamycin plus quinine (Holbrook et al., 2023). Still, these recommended treatments often result in various problems. For instance, multiple mutations in the *B. microti* cytochrome b, which was associated with the atovaquone-target region, were discovered in an atovaquone-treated relapse patient (Holbrook et al., 2023). The combination of clindamycin and quinine is the last resort for patients with severe symptoms. Despite its efficacy, some had adverse drug reactions after taking these drugs (Vannier and Krause, 2012). The 8-aminoquinoline analog, tafenoquine (TAF), was found to be effective in curing the patient's *B. microti* infection that was resistant to azithromycin and atovaquone with a 200 mg weekly dose (Marcos et al., 2022). However, the efficacy of TAF treatment may vary between individuals and cases and its contraindication in G6PD-deficient patients limits its clinical use. Due to the aforementioned issues, it is undoubtedly urgent to search for compounds that treat infections caused by *Babesia* spp. while simultaneously minimizing the detrimental effects of antibabesial drugs.

Spiroindolone cipargamin (CIP), a promising antimalarial, has been found to effectively

suppress the growth of all strains of *Plasmodium falciparum* and *P. vivax* with potency in the

low nanomolar ranges (Rosling et al., 2018; Rottmann et al., 2010). The assessment of the

pharmacokinetic characteristics of the drug indicated that oral administration of CIP had good

absorption, a prolonged half-life, and exceptional bioavailability (Schmitt et al., 2022).

Following oral CIP treatment (30 mg daily for three days) in adults with simple *P. falciparum*

or *P. vivax* malaria, the parasitemia was rapidly cleared in a phase II trial (Schmitt et al., 2022).

Currently, a clinical trial is being conducted for the intravenous administration of CIP as a

treatment for severe malaria patients (ClinicalTrials, 2024). CIP was also evaluated for treating

toxoplasmosis as a second-line medicine in cases where there are intolerable toxicity issues or

allergies to the currently used treatments. Five days post-*Toxoplasma gondii* infection, CIP-

treated mice (100 mg/kg/day on the day of and the day after infection) had 90% fewer parasites

than the control mice (Zhou et al., 2014).

Due to its excellent efficacy against other apicomplexan parasites, we hypothesize that CIP

may also be effective as an antibabesial drug. Therefore, the objectives of this study were to

determine whether CIP could inhibit the growth of *Babesia* spp., namely *B. bovis* and *B. gibsoni*

in vitro and *B. microti* and *B. rodhaini in vivo*, and identify the inhibitory mechanisms of CIP

on *Babesia* parasites (*Bg*ATP4) by monitoring the Na^+ and H^+ balance.

Results

Inhibitory efficacy of CIP on *B. bovis* and *B. gibsoni in vitro*

The *in vitro* activity of CIP against *B. bovis* and *B. gibsoni* showed a steep growth inhibition

curve with half inhibitory concentration (IC_{50}) values of 20.2 ± 1.4 nM (Figure 1A) and $69.4 \pm$

2.2 nM (Figure 1B), respectively. The CC_{50} value of CIP on MDCK cells was 38.7 ± 2.0 μM

(Figure supplement 1), resulting in a predicted selectivity index of 557.6. Furthermore, at a

concentration of 100 μM, CIP exhibited a low erythrocyte hemolysis rate of 0.11 ± 0.03 % (not

shown).

CIP effect on *B. microti* and *B. rodhaini* infections *in vivo*

Concurrently, CIP showed effective inhibition on *B. microti* and *B. rodhaini in vivo*. The

parasitemia of *B. microti*-infected BALB/c mice grew dramatically in the vehicle-treated

control group and peaked at 10 DPI (35.88 ± 4.32%) (Figure 1C). On the other hand, the

treatment with CIP (20 mg/kg of body weight) or atovaquone (ATO) plus azithromycin (AZI)

administered orally resulted in a significantly lower peak parasitemia, 1.06 ± 0.20% and 1.61

± 0.20%, respectively. Hematocrit (HCT) variations were tracked every 4 days as an indicator

of anemia in *B. microti*-infected mice. The vehicle-treated group showed a drop in HCT levels

from 8 DPI to 24 DPI, with statistically significant decreases at 12 and 16 DPI ($P < 0.01$). No

significant reduction in HCT levels was observed in the CIP-treated group or the ATO plus

AZI-treated group (Figure 1D). This suggests that the administration of CIP could control *B.*

microti infection and prevent anemia from developing in *B. microti*-infected mice. BALB/c

mice infected with *B. rodhaini* treated with sesame oil or ATO plus AZI showed high

parasitemia, 90.73 ± 1.97%, and 86.23 ± 3.06%, respectively (Figure 1E), and all mice died

within 10 DPI (Figure 1F). CIP treatment in *B. rodhaini*-infected mice precluded the emergence

of parasitemia for the following eight days (Figure 1E), which led to 66.67% of mice surviving

the challenge infection (Figure 1F). At 14 DPI, parasites had recurred in all CIP-treated *B.*

rodhaini-infected mice (10.32 ± 15.51%), which were eventually cleared as indicated by

undetectable parasites at 18 DPI (Figure 1E).

Identification of *B. gibsoni* ATP4 mutation in CIP-resistant strains

We sequenced the *B. gibsoni* ATP4 gene from two resistant parasite wells. A single nucleotide

variant (SNV) in *Bg*ATP4 with a substitution at position 2,761 (from C to G) was found − a

nonsynonymous coding change from leucine to valine (L921V) (Figure 2B). In another

resistant strain, the mutation showed in the same position. However, the nucleotide substitution

was from C to A, and the coding changed from leucine to isoleucine (L921I) (Figure 2C).

BgATP4^{L921V} and BgATP4^{L921I} lines were tested for their susceptibility to CIP, and had IC$_{50}$

values of 421.0 ± 15.88 nM and 887.9 ± 61.97 nM, respectively (Figure 2D). These findings

demonstrate a 6.1- and 12.8-fold reduction in CIP sensitivity of resistant parasite lines

BgATP4^{L921V} and BgATP4^{L921I}.

The effect of CIP on BgATP4^{WT}, BgATP4^{L921V} and BgATP4^{L921I} function

Microscopic observation of thin blood smears was performed to determine the morphological

changes of *B. gibsoni* exposed to CIP. The CIP-treated parasites became swollen after

incubation with the drug for 72 hr (Figure 3A). For both the treatment and control groups, one

hundred parasites were measured. The mean size of treated parasites was notably bigger than

the parasites in the untreated group ($P < 0.0001$) (Figure 3B). The addition of the ATP4

inhibitor CIP resulted in a time-dependent increase in the concentrations of $[Na^+]_i$ in wild-type

B. gibsoni (Figure 3C), with improved signal-to-noise ratios at the higher drug concentration

of 20 nM. We observed that the Na$^+$ concentrations in both BgATP4^{L921V} and BgATP4^{L921I} lines

decreased compared to those of the control BgATP4^{WT} line after being exposed to 20 nM CIP

for 20 min, with a significantly lower Na$^+$ concentration in BgATP4^{L921I} ($P = 0.0087$) (Figure

3E). We also demonstrated here that the addition of CIP in wild-type *B. gibsoni* caused an

increase in the cytosolic pH (Figure 3D). Specifically, 4 min after the drug was added, the

average pH of 20 nM CIP group reached as high as 7.278, while the 1 nM CIP group reached

7.089 and the untreated group reached 7.062. The pH values of the 20 nM CIP group were

consistently higher than those of the other two groups, although declining with time. In resistant

lines, a 20-min exposure to 20 nM CIP caused small changes in the pH values compared to the

wild-type line (Figure 3F), with the BgATP4^{L921I} line (7.048 ± 0.042) having a notably lower

pH value ($P = 0.0229$).

Multiple sequence alignment of *Babesia* ATP4 and molecular docking

The whole amino acid sequence of BgATP4 (GenBank: KAK1443404.1) shared identity values

of 29.75%, 49.40%, 49.67%, 62.21%, and 52.47% with *Homo sapiens* ATP4 (GenBank:

NM_000704.3), *P. falciparum* ATP4 (GenBank: PF3D7_1211900), *T. gondii* ATP4 (GenBank:

XP_018635122.1), *B. bovis* ATP4 (PiroplasmaDB: BBOV_IV010020), and *B. microti* ATP4

(GenBank: BMR1_03g01005), respectively (Figure supplement 2).

The pLDDT value of BgATP4^{WT} prediction was 80.7 using Colab-fold. Multiple potential

binding sites for CIP were revealed by blind docking throughout the whole protein surface

(Figure supplement 3). CIP binds in close proximity to L921, as demonstrated by focused

docking on this area (Figure 4A). The contribution of each residue to the predicted binding

affinity in either mutant structure was reduced; the precise values of BgATP4^{WT} (Figure 4B),

BgATP4^{L921V} (Figure 4C) and BgATP4^{L921I} (Figure 4D) were -6.43, -6.40 and -6.26 kcal/mol,

respectively. The interactions of CIP from each docking simulation are shown in Table

supplement 2.

Discussion

The repositioning of antimalarial drugs is critical in developing novel strategies for treating

babesiosis. CIP is a novel compound that inhibits *Plasmodium* development and has been

extensively tested in Phase 1 and Phase 2 clinical trials by targeting *Pf*ATP4 (Qiu et al., 2022).

Therefore, our study aimed to repurpose the antimalarial CIP by assessing its efficacy on

Babesia species. The IC$_{50}$ values of CIP against *B. bovis* and *B. gibsoni in vitro* were lower

than the previously reported IC$_{50}$ value of TAF (Ji et al., 2022b) and ATO against *B. gibsoni*

(Matsuu et al., 2004). The present investigation also demonstrated that the inhibitory effects of

CIP on *B. microti*-infected BALB/c mice were comparable to that of ATO plus AZI, the

medication prescribed by the U.S. CDC. CIP was also proven to protect mice from the deadly

B. rodhaini infection, with protection limited to 66.67% as recorded in the current trial. These

results suggest that CIP may be a potential chemotherapy candidate for babesiosis.

In a previous study, the *P. falciparum* Dd2 parent line initially had a subnanomolar IC_{50} value,

after the CIP-selected cultures to limiting dilution and selected clonal lines, the highly resistant

strain reported IC_{50} values 4200-4280 fold higher than those of wild-type Dd2 parasites with a

G358S mutation in the *Pf*ATP4 protein, a *P. falciparum* P-type Na^+ ATPase (Qiu et al., 2022).

The vital protein ATP4 is found in the parasite plasma membrane and is distinctive to the

subclass of the apicomplexan parasite (Mohring et al., 2022). To date, *in vitro* evolution

experiments using CIP have produced at least 18 parasite lines with various *Pf*ATP4 mutations

(Lee and Fidock, 2016; Rottmann et al., 2010). Another finding showed that the ATP4 activity

associated with $TgATP4^{G419S}$ was 34 times less susceptible to CIP than that of $TgATP4^{WT}$ (Qiu

et al., 2022). It follows that there is a significant chance that the resistant *Babesia* parasites will

also emerge from CIP exposure. In this study, we successfully produced two CIP-resistant

strains using six independent selections. The newly discovered L921V and L921I mutations in

*Bg*ATP4 decreased the CIP sensitivity by 6.1 and 12.8 times, respectively. We provided

compelling evidence that natural variety mutations L921V and L921I in *Bg*ATP4 significantly

affected the proteins' susceptibility to CIP inhibition, albeit the mutational sites were different

from those of *P. falciparum* and *T. gondii*.

Although ATP4 was initially recognized as a Ca^{2+} transporter (Krishna et al., 2001), current

evidence suggests that ATP4 functions as an ATPase for exporting Na^+ while importing H^+

(Mohring et al., 2022), as well as causing a variety of other physiological perturbations

including an increase in the volume of parasites and infected erythrocytes, due to the osmotic

impact of the $[Na^+]_{cyt}$ increase (Dennis et al., 2018), a decline in cholesterol extrusion from the

parasite plasma membrane as a result of the increase in $[Na^+]_{cyt}$ (Das et al., 2016), and an

intensified rigidity of erythrocytes infected with ring-stage parasites (Zhang et al., 2016). In

this study, the CIP-exposed wild-type *B. gibsoni* became swollen, which was identical to a prior

study on *P. falciparum* (Dennis et al., 2018). We hypothesize that the swelling of the isolated

parasites can be ascribed to the osmotic consequences of Na^+ uptake and is contingent upon

the presence of Na^+ in the external environment. This hypothesis is supported by the following

experimental results.

The output of Na^+ and input of H^+ diminished upon CIP-induced inhibition of *Pf*ATP4, and

the continued outflow of H^+ via V-type H^+-ATPase led to an alkalinization that ultimately killed

the parasites (Spillman et al., 2013). Our capacity to measure a time-dependent increase in the

concentration of $[Na^+]_i$ and pH value for wild-type *B. gibsoni* facilitated us to gain insight into

the basic mechanisms of CIP on *Bg*ATP4^{WT} function. Furthermore, our results validate that

internal alkalinization was the main factor in *Babesia* death and support our hypothesis that the

swollen isolated parasites were produced by Na^+ absorption. To explore further how mutations

in *Bg*ATP4 are associated with the upregulation of the parasite's $[Na^+]_i$ and $[H^+]_i$, we tested two

*Bg*ATP4-mutant lines that were chosen previously with *Bg*ATP4 inhibitors (*Bg*ATP4^{L921V} and

*Bg*ATP4^{L921I}). Due to the presence of L921V and L921I mutations in drug-resistant strains of

*Bg*ATP4, the concentration of Na^+ did not increase as much as it would have in the wild-type

strain following the addition of 20 nM CIP for 20 min. As intraerythrocytic alkalinization rises,

the same outcome happens. From these results, we deduced how natural mutations in *Bg*ATP4

may affect ATP4 inhibitor susceptibility by dysregulating H^+ and Na^+ balance, which helped

parasites survive in a relatively high concentration of CIP. An illustration of the putative

processes for regulating Na^+ and H^+ in erythrocytes infected with the *Babesia* parasite is

presented in Figure supplement 4.

According to studies using the *Pf*ATP4 model for molecular docking, the G358S mutation

results in a steric clash that lowers CIP's binding affinity (Qiu et al., 2022). The results from

the current study were similar to the *Pf*ATP4 model. The molecular docking was constructed

using a ColabFold model of wild-type *Bg*ATP4, which predicted the binding mode and affinity

between ATP4 protein and the ligand to provide a possible mechanistic explanation. It

suggested that the L921V mutation caused changes at the atomic level, whereas the L921I

mutation created a steric clash that reduced the binding affinity of CIP. The predicted affinity

score of the L921I mutation was lower than that of the L921V mutation. Thus, it is possible

that BgATP4^{L921I} has a weaker binding to the drug compared to BgATP4^{L921V}. These findings

are consistent with the results obtained from measuring the IC$_{50}$ of the drug against parasites

with the L921V and L921I mutations.

Materials and methods

Parasite culture

The parasites *B. gibsoni* (Oita strain) and *B. bovis* Texas strain were *in vitro* cultured in 24-well

plates and maintained in an atmosphere of 5% CO_2 and 5% O_2 at 37 °C (Liu et al., 2018). For

the *in vivo* studies, *B. microti* Peabody mjr strain-(ATCC PRA-99) and *B. rodhaini* Australia

strain-infected RBCs (iRBCs), which were collected and diluted with phosphate-buffered

saline (PBS) when the parasitemia levels in the donor mice reached ~20% and 50%,

respectively, and were intraperitoneally injected into BALB/c mice. Each BALB/c mouse was

infected with 1.0×10^7 *B. microti* or *B. rodhaini* iRBCs for the *in vivo* trials (Ji et al., 2022a).

***In vitro* cytotoxicity of CIP in canine cells and hemolysis rate in canine erythrocytes**

Cultures of Madin-Darby canine kidney (MDCK) cells were maintained at 37 °C under an

atmosphere of 5% CO_2 and 5% O_2 and the cytotoxic effect of CIP (MedChem Express, Tokyo,

Japan) was assessed using a cell viability assay by CCK-8 (Dojindo, Japan) as described

previously (Li et al., 2023). The selectivity index is calculated as the ratio between the half

maximal inhibitory concentration (IC$_{50}$) and the 50% cytotoxic concentration (CC$_{50}$) values.

Canine erythrocytes were collected from healthy beagle dogs raised in NRCPD and stocked in

Vega y Martinez (VYM) phosphate-buffered saline solution at 4 °C (Vega et al., 1985). A

canine erythrocyte hemolysis assay was performed at concentrations of 0.1, 1, 5, 10, 25, 50,

and 100 µM as previously described (Ariefta et al., 2022).

Evaluation of the efficacy of CIP against *Babesia* parasites *in vitro*

The efficacy of CIP against *B. gibsoni* and *B. bovis* was determined using a fluorescence assay,

as previously described (Guswanto et al., 2014). The IC_{50} values were determined from the

fluorescence values and by non-linear regression analysis (curve fit) in GraphPad Prism 9

(GraphPad Software Inc., USA).

Chemotherapeutic effects of CIP against *Babesia* infections *in vivo*

CIP was evaluated on *B. microti*- and *B. rodhaini*-infected mice, as previously described

(Ndayisaba et al., 2021; Tuvshintulga et al., 2022). When all mice had a 1% average parasitemia

at 4 days postinfection (DPI), the drug treatments were administered, which continued for

seven days. Three groups of *B. microti*-infected mice were administered different treatments.

The CIP group (n = 6) and the ATO plus AZI group (n = 6) were orally treated with 20 mg/kg

CIP and 20 mg/kg ATO plus AZI (Sigma, Tokyo, Japan), respectively. The infected mice of the

vehicle group (n = 6) orally received 0.2 mL of sesame oil as the control. Eighteen mice infected

with *B. rodhaini* were likewise placed into three groups for treatment, and they received the

same care as the mice infected with *B. microti*. A light microscope (Nikon, Japan) and a

hematology analyzer (Celltac α MEK-6450, Nihon Kohden Corporation, Tokyo, Japan) were

used to observe the parasitemia and hematocrit levels every two and four days, respectively.

Selection of CIP-resistant *B. gibsoni in vitro*

Selections were initiated by exposing six independent flasks, each containing 10 μL *B. gibsoni*

iRBCs mixed with 40 μL RBC into a 450 μL culture medium, which contained increasing

concentrations of CIP: 5, 10, 20, 30 to 694 nM ($10 \times IC_{50}$). The medium containing CIP was

replaced daily until parasites treated with $10 \times IC_{50}$ CIP reached multiplication rates that were

approximately comparable to those of the untreated controls (Hwang et al., 2010). Then, to

evaluate the decreased sensitivity to CIP, IC_{50} values of resistant strains were determined by

nonlinear regression using the GraphPad Prism software.

Detection of *B. gibsoni* ATP4 gene mutations

The genomic DNA of the mutant parasites was extracted and sequenced (Ji et al., 2022a). The

278   primer sets used for sequencing are listed in Table supplement 1. The single nucleotide variants

279   were confirmed by pairwise alignment to the BgATP4^{WT} sequence (GenBank:

280   JAVEPI010000002.1).

Morphological changes in CIP-treated *in vitro* cultured *B. gibsoni*

282   A microscopy assay was used to detect the morphological changes of wild-type *B. gibsoni* after

283   exposure to 50 nM CIP for three consecutive days (Liu et al., 2021; Tayebwa et al., 2018). At

284   4 days post-treatment, ImageJ software was used to measure the sizes of 100 randomly selected

285   parasites in the CIP-treated group and the control group on Giemsa-stained blood smears.

Parasites [Na$^+$]$_i$ and pH$_i$ measurements

For both [Na$^+$]$_i$ and pH$_i$ measurements, the wild-type and mutant parasites were initially

separated from erythrocytes by treatment with saponin (0.05% [wt/vol]) for 5 min (Saliba and

Kirk, 1999). The Na$^+$-sensitive fluorescent dye SBFI (Thermo Fisher Scientific; product

S1263) was used to quantify intracellular sodium ([Na$^+$]$_i$). Saponin-isolated parasites (at 1×10^8

parasites/mL) were loaded with SBFI (5 μM; in the presence of 0.02% w/v Pluronic F127) for

1 hr at room temperature (RT) (Spillman et al., 2013). Thereafter, SBFI-loaded parasites were

resuspended in physiological saline (120 mM NaCl, 5 mM KCl, 25 mM HEPES, 20 mM D-

glucose, and 1 mM MgCl$_2$ [pH 7.1]) at RT in the presence or absence of CIP. A 96-well

microtiter plate was filled with around 200 μL of parasite suspension per well. The dye-loaded

cells fluoresced at 515 nm after being stimulated at 340 and 380 nm. Parasites loaded with

SBFI were suspended in calibration buffers containing [Na$^+$] values ranging from 0 to 140 mM

(pH 7.1) to establish calibration curves (Diarra et al., 2001).

The cytosolic pH of wild-type and mutant strains was measured using the pH-sensitive

fluorescent dye BCECF [2',7'-bis-(2-carboxyethyl)-5-(and-6)- carboxyfluorescein] (Biotium;

product 51011). The BCECF was added to saponin-isolated parasites by suspension (1×10^8

parasites/mL) and incubated for 20 minutes at 37 °C in RPMI-1640 culture medium (Gibco,

USA) (Saliba and Kirk, 1999). Thereafter, the parasites loaded with dye were rinsed thrice

(12,000 × g, 1 min) in RPMI-1640 culture medium, then resuspended in physiological saline

(as previously mentioned) with or without various concentrations of CIP. A 96-well microtiter

plate was filled with around 200 μL of parasite suspension per well. The dye-loaded cells

fluoresced at 520 nm after being stimulated at 440 and 490 nm. A pH calibration was carried

out for each experiment (Mohring et al., 2022).

Multiple ATP4 sequence alignment and molecular docking

B. gibsoni ATP4 (GenBank: KAK1443404.1), *B. bovis* ATP4 (PiroplasmaDB:

BBOV_IV010020), *B. microti* ATP4 (GenBank: BMR1_03g01005), *T. gondii* ATP4

(GenBank: XP_018635122.1), and *Homo sapiens* ATP4 (GenBank: NM_000704.3) sequences

were obtained by a homology search using *P. falciparum* ATP4 (GenBank: PF3D7_1211900).

Sequence alignment was analyzed using MUSCLE in Jalview v2.11.3.2 software and BLAST

(http://www.ncbi.nlm.nih.gov/BLAST/).

AlphaFold was used to predict the structure of BgATP4^{WT} (Jumper et al., 2021). Using the

GROMACS 2021 Molecular Dynamics package, the energy minimization was carried out

following the model's generation (Lindahl et al., 2021). The mutations were produced by using

Charmm-GUI PDB reader (Jo et al., 2014). PyMol (version 2.0 Schrödinger, LLC) was used

to confirm the position of the mutation site (center: -20.367, 10.904, 9.435; size: 30×30×30)

(Trott and Olson, 2010). The ligand molecules CIP was downloaded from PubChem (CID

44469321 (https://pubchem.ncbi.nlm.nih.gov/compound/ 44469321). The ligand was used to

dock with Gnina (Eberhardt et al., 2021). The affinity score and binding pose were chosen only

from the highest CNN (convolutional neural network) score results from docking simulations.

The models were visualized using the PyMOL Molecular Graphic System and Discovery

Studio.

Statistical analysis

Data analysis, namely one-way analysis of variance (ANOVA) and two-tailed unpaired t-tests,

was performed using GraphPad Prism (La Jolla, CA, USA) version 9. A P value of <0.05 was

considered a statistically significant result.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research (22H02509), the JSPS

Core-to-Core program, both from the Ministry of Education, Culture, Sports, Science, and

Technology of Japan, and a grant from the Strategic International Collaborative Research

Project (JPJ008837) promoted by the Ministry of Agriculture, Forestry, and Fisheries of Japan,

and Natural Science Foundation of Hubei Province (2023AFC001).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Author contributions

Hang Li, Investigation, Data curation, Formal analysis, Writing – original draft, Writing –

review & editing; Shengwei Ji, Methodology, Investigation, Software, Validation; Nanang R.

Ariefta, Methodology, Software, Formal analysis; Eloiza May S. Galon, Visualization, Writing

– review & editing; Shimaa AES El-Sayed, Methodology, Visualization; Lijun Jia, Data

curation, Visualization; Yoshifumi Nishikawa, Methodology, Resources; Mingming Liu,

Conceptualization, Methodology, Investigation, Resources, Data curation, Funding acquisition,

Supervision, Writing – review & editing; Xuenan Xuan, Conceptualization, Methodology,

Validation, Formal analysis, Supervision, Project administration, Writing – review & editing.

Ethics

All animal experiments were performed according to the Guide for the Care and Use of Laboratory Animals of Obihiro University of Agriculture and Veterinary Medicine, which was also approved by the Committee on the Ethics of Animal Experiments at the Obihiro University of Agriculture and Veterinary Medicine, Japan (permit numbers: animal experiment, 22-145 and 23-132; DNA experiment, 2207 and 2208; pathogen, 202308 and 202306).

References

Almazán C, Scimeca RC, Reichard MV, Mosqueda J. 2022. Babesiosis and Theileriosis in North America. *Pathogens* **11**:168. doi:10.3390/pathogens11020168

Ariefta NR, Pagmadulam B, Nihei C, Nishikawa Y. 2022. Sparsomycin exhibits potent antiplasmodial activity *in vitro* and *in vivo*. *Pharmaceutics* **14**:544. doi:10.3390/pharmaceutics14030544

ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). Identifier NCT04675931, To evaluate efficacy, safety, tolerability and PK of intravenous cipargamin in participants with severe *Plasmodium falciparum* malaria; 2024 Feb 29 [cited 2024 Apr 9]. Available from: https://clinicaltrials.gov/study/NCT04675931

Das S, Bhatanagar S, Morrisey JM, Daly TM, Burns JM, Coppens I, Vaidya AB. 2016. Na^+ influx induced by new antimalarials causes rapid alterations in the cholesterol content and morphology of *Plasmodium falciparum*. *PLoS Pathog* **12**:e1005647. doi:10.1371/journal.ppat.1005647

Dennis ASM, Lehane AM, Ridgway MC, Holleran JP, Kirk K. 2018. Cell swelling induced by the antimalarial KAE609 (cipargamin) and other *Pf*ATP4-associated antimalarials. *Antimicrob Agents Chemother* **62**:e00087-18. doi:10.1128/AAC.00087-18

Diarra A, Sheldon C, Church J. 2001. In situ calibration and $[H^+]$ sensitivity of the fluorescent Na^+ indicator SBFI. *American Journal of Physiology-Cell Physiology* **280**:C1623–C1633. doi:10.1152/ajpcell.2001.280.6.C1623

Drews SJ, Kjemtrup AM, Krause PJ, Lambert G, Leiby DA, Lewin A, O'Brien SF, Renaud C, Tonnetti L, Bloch EM. 2023. Transfusion-transmitted *Babesia* spp.: a changing landscape of epidemiology, regulation, and risk mitigation. *J Clin Microbiol* **61**:e01268-22. doi:10.1128/jcm.01268-22

Eberhardt J, Santos-Martins D, Tillack AF, Forli S. 2021. AutoDock Vina 1.2.0: new docking methods, expanded force field, and python bindings. *J Chem Inf Model* **61**:3891–3898.

doi:10.1021/acs.jcim.1c00203

Guswanto A, Sivakumar T, Rizk MA, Elsayed SAE, Youssef MA, ElSaid EES, Yokoyama N,
Igarashi I. 2014. Evaluation of a fluorescence-based method for antibabesial drug screening.
Antimicrob Agents Chemother **58**:4713–4717. doi:10.1128/AAC.00022-14

Holbrook NR, Klontz EH, Adams GC, Schnittman SR, Issa NC, Bond SA, Branda JA, Lemieux
JE. 2023. *Babesia microti* variant with multiple resistance mutations detected in an
immunocompromised patient receiving atovaquone prophylaxi. *Open Forum Infectious*
Diseases **10**:ofad097. doi:10.1093/ofid/ofad097

Hwang S-J, Yamasaki M, Nakamura K, Sasaki N, Murakami M, Wickramasekara
Rajapakshage BK, Ohta H, Maede Y, Takiguchi M. 2010. Development and characterization
of a strain of *Babesia gibsoni* resistant to diminazene aceturate *in vitro*. *J Vet Med Sci*
72:765–771. doi:10.1292/jvms.09-0535

Jalovecka M, Sojka D, Ascencio M, Schnittger L. 2019. *Babesia* life cycle-when phylogeny
meets biology. *Trends in Parasitology* **35**:356–368. doi:10.1016/j.pt.2019.01.007

Ji S, Galon EM, Rizk MA, Yi Y, Zafar I, Li H, Ma Z, Iguchi A, Asada M, Liu M, Xuan X. 2022.
Efficacy of the antimalarial MMV390048 against *Babesia* infection reveals
phosphatidylinositol 4-kinase as a druggable target for babesiosis. *Antimicrob Agents*
Chemother **66**:e00574-22. doi:10.1128/aac.00574-22

Ji S, Liu M, Galon EM, Rizk MA, Tuvshintulga B, Li J, Zafar I, Hasegawa Y, Iguchi A,
Yokoyama N, Xuan X. 2022. Inhibitory effect of naphthoquine phosphate on *Babesia*
gibsoni in vitro and *Babesia rodhaini in vivo*. *Parasites Vectors* **15**:10. doi:10.1186/s13071-
021-05127-0

Jo S, Cheng X, Islam SM, Huang L, Rui H, Zhu A, Lee HS, Qi Y, Han W, Vanommeslaeghe K,
MacKerell AD, Roux B, Im W. 2014. CHARMM-GUI PDB manipulator for advanced
modeling and simulations of proteins containing nonstandard residues. *Adv Protein Chem*
Struct Biol **96:** 235–265. doi:10.1016/bs.apcsb.2014.06.002

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
Paredes B, Nikolov S, Jain R, Adler J, Back T, et al. 2021. Highly accurate protein structure
prediction with AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-021-03819-2

Krause PJ. 2019. Human babesiosis. *International Journal for Parasitology* **49**:165–174.
doi:10.1016/j.ijpara.2018.11.007

Krishna S, Woodrow C, Webb R, Penny J, Takeyasu K, Kimura M, East JM. 2001. Expression
and functional characterization of a *Plasmodium falciparum* Ca^{2+}-ATPase (*Pf*ATP4)
belonging to a subclass unique to apicomplexan organisms. *Journal of Biological Chemistry*
276:10782–10787. doi:10.1074/jbc.M010554200

Lee AH, Fidock DA. 2016. Evidence of a mild mutator phenotype in cambodian *Plasmodium*
falciparum malaria parasites. *PLoS ONE* **11**:e0154166. doi:10.1371/journal.pone.0154166

Li H, Galon EM, Ji S, Zafar I, Ma Z, Do T, Amer MM, Ma Y, Liu M, Xuan X. 2023. *In vitro*
screening of compounds from the Food and Drug Administration-approved library identifies
anti-*Babesia gibsoni* activity of idarubicin hydrochloride and vorinostat. *Parasitology*
International **96**:102774. doi:10.1016/j.parint.2023.102774

Lindahl, Abraham, Hess, Spoel VD. 2021. GROMACS 2021 Manual.
doi:10.5281/ZENODO.4457591

Liu M, Adjou Moumouni PF, Cao S, Asada M, Wang G, Gao Y, Guo H, Li J, Vudriko P,
Efstratiou A, Ringo AE, Lee S-H, Hakimi H, Masatani T, Sunaga F, Kawazu S, Yamagishi J,
Jia L, Inoue N, Xuan X. 2018. Identification and characterization of interchangeable cross-
species functional promoters between *Babesia gibsoni* and *Babesia bovis*. *Ticks and Tick-*
borne Diseases **9**:330–333. doi:10.1016/j.ttbdis.2017.11.008

Liu M, Ji S, Kondoh D, Galon EM, Li J, Tomihari M, Yanagawa M, Tagawa M, Adachi M,
Asada M, Igarashi I, Iguchi A, Xuan X. 2021. Tafenoquine is a promising drug candidate for
the treatment of babesiosis. *Antimicrob Agents Chemother* **65**:e00204-21.
doi:10.1128/AAC.00204-21

Marcos LA, Leung A, Kirkman L, Wormser GP. 2022. Use of tafenoquine to treat a patient
with relapsing babesiosis with clinical and molecular evidence of resistance to azithromycin
and atovaquone. *IDCases* **27**:e01460. doi:10.1016/j.idcr.2022.e01460

Matsuu A, Koshida Y, Kawahara M, Inoue K, Ikadai H, Hikasa Y, Okano S, Higuchi S. 2004.
Efficacy of atovaquone against *Babesia gibsoni in vivo* and *in vitro*. *Veterinary Parasitology*
124:9–18. doi:10.1016/j.vetpar.2004.07.005

Mohring F, Van Schalkwyk DA, Henrici RC, Blasco B, Leroy D, Sutherland CJ, Moon RW.
2022. Cation ATPase (ATP4) orthologue replacement in the malaria parasite *Plasmodium*
knowlesi reveals species-specific responses to ATP4-targeting drugs. *mBio* **13**:e01178-22.
doi:10.1128/mbio.01178-22

Ndayisaba G, Yeka A, Asante KP, Grobusch MP, Karita E, Mugerwa H, Asiimwe S, Oduro A,
Fofana B, Doumbia S, Jain JP, Barsainya S, Kullak-Ublick GA, Su G, Schmitt EK, Csermak
K, Gandhi P, Hughes D. 2021. Hepatic safety and tolerability of cipargamin (KAE609), in
adult patients with *Plasmodium falciparum* malaria: a randomized, phase II, controlled,
dose-escalation trial in sub-Saharan Africa. *Malar J* **20**:478. doi:10.1186/s12936-021-
04009-1

Qiu D, Pei JV, Rosling JEO, Thathy V, Li D, Xue Y, Tanner JD, Penington JS, Aw YTV, Aw
JYH, Xu G, Tripathi AK, Gnadig NF, Yeo T, Fairhurst KJ, Stokes BH, Murithi JM,
Kümpornsin K, Hasemer H, Dennis ASM, et al. 2022. A G358S mutation in the *Plasmodium*
falciparum Na$^+$ pump *Pf*ATP4 confers clinically-relevant resistance to cipargamin. *Nat*
Commun **13**:5746. doi:10.1038/s41467-022-33403-9

Rosling JEO, Ridgway MC, Summers RL, Kirk K, Lehane AM. 2018. Biochemical
characterization and chemical inhibition of *Pf*ATP4-associated Na$^+$-ATPase activity in

Plasmodium falciparum membranes. *Journal of Biological Chemistry* **293**:13327–13337. doi:10.1074/jbc.RA118.003640

Rottmann M, McNamara C, Yeung BKS, Lee MCS, Zou B, Russell B, Seitz P, Plouffe DM, Dharia NV, Tan J, Cohen SB, Spencer KR, González-Páez GE, Lakshminarayana SB, Goh A, Suwanarusk R, Jegla T, Schmitt EK, Beck H-P, Brun R, et al. 2010. Spiroindolones, a potent compound class for the treatment of malaria. *Science* **329**:1175–1180. doi:10.1126/science.1193225

Saliba KJ, Kirk K. 1999. pH regulation in the intracellular malaria parasite, *Plasmodium falciparum*. *Journal of Biological Chemistry* **274**:33213–33219. doi:10.1074/jbc.274.47.33213

Schmitt EK, Ndayisaba G, Yeka A, Asante KP, Grobusch MP, Karita E, Mugerwa H, Asiimwe S, Oduro A, Fofana B, Doumbia S, Su G, Csermak Renner K, Venishetty VK, Sayyed S, Straimer J, Demin I, Barsainya S, Boulton C, Gandhi P. 2022. Efficacy of cipargamin (KAE609) in a randomized, phase II dose-escalation study in adults in Sub-Saharan Africa with uncomplicated *Plasmodium falciparum* malaria. *Clinical Infectious Diseases* **74**:1831–1839. doi:10.1093/cid/ciab716

Spillman NJ, Allen RJW, McNamara CW, Yeung BKS, Winzeler EA, Diagana TT, Kirk K. 2013. Na$^+$ regulation in the malaria parasite *Plasmodium falciparum* involves the cation ATPase *Pf*ATP4 and is a target of the spiroindolone antimalarials. *Cell Host & Microbe* **13**:227–237. doi:10.1016/j.chom.2012.12.006

Tayebwa DS, Tuvshintulga B, Guswanto A, Nugraha AB, Batiha GE-S, Gantuya S, Rizk MA, Vudriko P, Sivakumar T, Yokoyama N, Igarashi I. 2018. The effects of nitidine chloride and camptothecin on the growth of *Babesia* and *Theileria* parasites. *Ticks and Tick-borne Diseases* **9**:1192–1201. doi:10.1016/j.ttbdis.2018.04.019

Trott O, Olson AJ. 2010. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* **31**:455–461. doi:10.1002/jcc.21334

Tuvshintulga B, Sivakumar T, Nugraha AB, Ahedor B, Batmagnai E, Otgonsuren D, Liu M, Xuan X, Igarashi I, Yokoyama N. 2022. Combination of clofazimine and atovaquone as a potent therapeutic regimen for the radical cure of *Babesia microti* infection in immunocompromised hosts. *The Journal of Infectious Diseases* **225**:238–242. doi:10.1093/infdis/jiab537

Vannier E, Krause PJ. 2012. Human Babesiosis. *N Engl J Med* **366**:2397–2407. doi:10.1056/NEJMra1202018

Vega CA, Buening GM, Green TJ, Carson CA. 1985. *In vitro* cultivation of *Babesia bigemina*. *Am J Vet Res* **46**:416–420.

Zhang R, Suwanarusk R, Malleret B, Cooke BM, Nosten F, Lau Y-L, Dao M, Lim CT, Renia L, Tan KSW, Russell B. 2016. A basis for rapid clearance of circulating ring-stage malaria

500   parasites by the spiroindolone KAE609. *J Infect Dis* **213**:100–104. doi:10.1093/infdis/jiv358

501 Zhou Y, Fomovska A, Muench S, Lai B-S, Mui E, McLeod R. 2014. Spiroindolone that inhibits

502   *Pf*ATPase4 is a potent, cidal inhibitor of *Toxoplasma gondii* tachyzoites *in vitro* and *in vivo*.

503   *Antimicrob Agents Chemother* **58**:1789–1792. doi:10.1128/AAC.02225-13

Figure 1. CIP demonstrates potent inhibition on *Babesia* spp.. (A) and **(B)** Dose-dependent

growth curve of CIP on *B. bovis* and *B. gibsoni in vitro*. Each value represents the mean ±

standard deviation (SD) of three independent experiments carried out in triplicate. **(C)**

Inhibitory effects of CIP and atovaquone (ATO) plus azithromycin (AZI) on the proliferation

of *B. microti* in BALB/c mice. **(D)** Hematocrit (HCT) values in mice treated with CIP or ATO

plus AZI compared with vehicle-treated mice. **(E)** Inhibitory effects of CIP and atovaquone

(ATO) plus azithromycin (AZI) on the proliferation of *B. rodhaini* in BALB/c mice. **(F)**

Survival rates of CIP-treated, ATO plus AZI-treated, and vehicle-treated mice. The treatment

time is shown by arrows, and significant differences ($P < 0.01$) between the drug-treated groups

and the vehicle-treated control group are indicated by asterisks. The data from one of six

individual experiments are expressed as means ± SD. *, $P < 0.05$; **, $P < 0.01$.

Figure 2. Mutations in *Bg*ATP4 mediate CIP resistance. (A-C) Representative sequencing chromatogram of resistant parasites from CIP-treated *B. gibsoni*. The resistant parasite genomic DNA is extracted from blood samples after a 60 day-treatment. The *Bg*ATP4 gene was amplified and sequenced using the DNA. **(D)** Dose-dependent growth curve of *Bg*ATP4^{WT}, *Bg*ATP4^{L921V}, and *Bg*ATP4^{L921I} *in vitro*. Each value represents the mean ± standard deviation (SD) of three independent experiments carried out in triplicate.

Figure 3. Effect of CIP on $[Na^+]_i$ and pH regulation in *B. gibsoni* lines. (A) Untreated and CIP-treated parasite morphology after incubation for 72 h. Scale bar: 5 μm. **(B)** Sizes of 100 parasites in two groups measured with ImageJ software in panels A. Statistically significant differences between the means of variables determined by t-test. ****, $P < 0.0001$. **(C)** $[Na^+]_i$ concentrations after addition of CIP in BgATP4^{WT} line. Representative traces from the experiment that highlight the impact of adding 20 nM CIP (green), 1 nM CIP (blue), or 0 nM CIP (grey) on the concentration $[Na^+]_i$ of the BgATP4^{WT} line. **(D)** Alkalinization of pH$_i$ in BgATP4^{WT} line upon addition of the ATP4 inhibitor. **(E)** Addition of 20 nM CIP to the wild-type and resistant parasite lines results in different $[Na^+]_i$ concentrations. **(F)** Addition of 20 nM CIP to the wild-type and resistant parasite lines results in different pH$_i$ concentrations. Experiments were performed in technical duplicates for at least three biological repeats. ns, non significant, $p > 0.05$; *, $P < 0.05$; **, $P < 0.01$.

Figure 4. Binding sites proximal to *Bg*ATP4 residue 921 predicted by molecular docking.

(A) The lowest energy poses for CIP were located in reference to the whole protein structure,

docking against the WT (green), L921V (yellow), and L921I (pink) mutant *Bg*ATP4. The side

chain of L921 is also shown in a red stick at its position. **(B-D)** The zoomed views of the

binding locations of CIP.

Supplementary Figures

Figure supplement 1. Cytotoxicity assay of CIP on the MDCK cell line. The MTP-500

microplate reader is utilized to detect the absorbance at 450 nm. The results are displayed as

the mean ± standard deviation of three separate tests.

Figure supplement 2. Multiple sequence alignment of ATP4 in different species. A yellow square denotes the *Bg*ATP4 mutation site discovered in this investigation; red squares represent sites linked to *P. falciparum* CIP resistance, and a green square represents sites associated with *T. gondii*.

Figure supplement 3. Binding sites for CIP found by Gnina search across the entire surface of the protein. (A-C) The binding space of WT, L921V, and L921I mutants in *Bg*ATP4 are labeled in green, yellow and pink, respectively.

Figure supplement 4. Proposed mechanism of inhibition of CIP on wild-type and mutant parasite-infected erythrocytes. CIP disrupts the *Bg*ATP4 function of wild-type parasites, which causes a net influx of Na$^+$ and efflux of H$^+$ from the parasite. The osmotic load imposed on the influx of Na$^+$ further brings about parasite swelling and internal alkalinization, which are the main factors in *Babesia* death. Mutations in ATP4 minimize the susceptibility to ATP4 inhibitors by recovering H$^+$ and Na$^+$ balance.

575 **Supplemental Tables**

576 **Table supplement 1.** Primer sets of *B. gibsoni* ATP4

Gene	Primer (5'-3')
*Bg*ATP4 1F	ACGTAGTTGTCGAACATCTTACACCAATGC
*Bg*ATP4 1R	CAGAGGTAGCATAGTGAGATCGCCCCGA
*Bg*ATP4 2F	GAGATTGCAAGGTTTACGGATGAGTCTATGG
*Bg*ATP4 2R	ATGACAACGTCACCGGGTACAACACGCTTG
*Bg*ATP4 3F	GAAAAGCTTGCTCAAATGTCTTCACCGACAAC
*Bg*ATP4 3R	GTTTCTGGACGAGTTGGATCCTCATAACC
*Bg*ATP4 4F	AAATTGGGGGGGATAATTGGCATTATCTCC
*Bg*ATP4 4R	ACCTTATCAAAGGGTACCTCTTGCTTTG
*Bg*ATP4 5F	TCAATCCATTCGGTGGAATTTTCAAAAAGG
*Bg*ATP4 5R	GTGTTTGAATAACATATCTGGTGCCCCCTT
*Bg*ATP4 6F	GGTAAAGAACAGATTTGCTGACATAGATCT
*Bg*ATP4 6R	ACCCTATTTCTTGTGCGATGGCCGTAGCTG
*Bg*ATP4 7F	AGGGACGCTATAAATACTTGTGGAAAGGC
*Bg*ATP4 7R	GTACGCCCAGACTCGATTGAGTTAACTATTG
*Bg*ATP4 8F	GGGTATTAATGGCACAGATGTTGCAAAAGG
*Bg*ATP4 8R	GACAGTTCCCACTAGCATCATTAAGTGTAA
*Bg*ATP4 9F	ATCATCATTGCGCTATATGTTTCTACCGG
*Bg*ATP4 9R	CTAATGGGGTAGCATTTCCATTCTCAACAA
*Bg*ATP4 10F	AGGGAGTTCCTCCCATGTATAAAGATACCT
*Bg*ATP4 10R	TTAGGCTTCTGCCCTTTTCATTTTTCGCTT

 Table supplement 2. Interactions of CIP from docking simulations

Receptor	Name	Distance	Category	Types
*Bg*ATP4^{WT}	GLY407	2.67648	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	CYS959	3.2396	Halogen	Halogen (Cl, Br, I)
	VAL962	2.66882	Hydrophobic	Pi-Sigma
	CYS959	4.38742	Hydrophobic	Alkyl
	VAL962	4.29249	Hydrophobic	Alkyl
	ALA963	3.8231	Hydrophobic	Alkyl
	ALA1202	3.67015	Hydrophobic	Alkyl
	MET211	5.06565	Hydrophobic	Alkyl
	VAL962	4.25668	Hydrophobic	Alkyl
	ILE414	4.70605	Hydrophobic	Alkyl
	CYS959	4.83891	Hydrophobic	Pi-Alkyl
	VAL962	4.98402	Hydrophobic	Pi-Alkyl
	VAL411	5.43239	Hydrophobic	Pi-Alkyl
	VAL962	5.05537	Hydrophobic	Pi-Alkyl
	PHE954	4.07586	Hydrophobic	Pi-Alkyl
*Bg*ATP4^{L921V}	GLY407	2.69013	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	CYS959	3.22066	Halogen	Halogen (Cl, Br, I)
	VAL962	2.66391	Hydrophobic	Pi-Sigma
	CYS959	4.35422	Hydrophobic	Alkyl
	VAL962	4.31879	Hydrophobic	Alkyl
	ALA963	3.8209	Hydrophobic	Alkyl
	ALA1202	3.63038	Hydrophobic	Alkyl
	MET211	5.0878	Hydrophobic	Alkyl
	VAL962	4.24186	Hydrophobic	Alkyl
	ILE414	4.71819	Hydrophobic	Alkyl
	CYS959	4.82036	Hydrophobic	Pi-Alkyl
	VAL962	4.99879	Hydrophobic	Pi-Alkyl
	VAL411	5.4334	Hydrophobic	Pi-Alkyl
	VAL962	5.04532	Hydrophobic	Pi-Alkyl
	PHE954	4.07482	Hydrophobic	Pi-Alkyl
*Bg*ATP4^{L921I}	GLY958	2.92346	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	GLY958	3.3301	Halogen	Halogen (Fluorine)
	MET410	5.88505	Other	Pi-Sulfur
	VAL955	4.31406	Hydrophobic	Alkyl

MET410	3.69023	Hydrophobic	Alkyl
VAL411	4.83101	Hydrophobic	Alkyl
VAL955	4.79513	Hydrophobic	Alkyl
VAL955	4.84729	Hydrophobic	Alkyl
CYS959	4.1269	Hydrophobic	Alkyl
ILE1205	4.82529	Hydrophobic	Alkyl
CYS959	5.21399	Hydrophobic	Alkyl
VAL955	4.90527	Hydrophobic	Pi-Alkyl
PHE954	5.03012	Hydrophobic	Pi-Alkyl