Efficacy and mechanism of action of cipargamin as an antibabesial drug candidate

Authors

Hang Li[1#], Shengwei Ji[1,2#], Nanang R. Ariefta[1], Eloiza May S. Galon[3], Shimaa AES

El-Sayed[1,4], Thom Do[1], Lijun Jia[2], Miako Sakaguchi[5], Masahito Asada[1], Yoshifumi

Nishikawa[1], Xin Qin[6], Mingming Liu[6]*, Xuenan Xuan[1,7]*

Affiliations

[1]National Research Center for Protozoan Diseases, Obihiro University of Agriculture

Veterinary Medicine, Inada-cho, Obihiro, Japan

[2]Department of Veterinary Medicine, Agriculture College of Yanbian University, Yanji,

China

[3]College of Veterinary Medicine and Biomedical Sciences, Cavite State University, Indang,

Philippines

[4]Department of Biochemistry and Molecular Biology, Faculty of Veterinary Medicine,

Mansoura University, Egypt

[5]Central Laboratory, Institute of Tropical Medicine (NEKKEN), Nagasaki University,

Nagasaki, Japan

[6]School of Basic Medicine, Hubei University of Arts and Science, Xiangyang, China

[7]Research Center for Asian Infectious Diseases, The University of Tokyo, Tokyo, Japan

*Correspondence: Mingming Liu; Xuenan Xuan

Address: Longzhong Road No.296, Xiangyang, China; Nishi 11 Minami 39 Chome, Obihiro,

Hokkaido

Telephone number: +8618904416800; +8109079036369

E-mail address: lmm_2010@hotmail.com; gen@obihiro.ac.jp

Competing interest: The authors have declared that no conflict of interest exists.

# These authors contributed equally to this work.

Abstract

Babesiosis is a disease brought on by intraerythrocytic parasites of the genus *Babesia*.

Current chemotherapies are accompanied by side effects and parasite relapse. Therefore, it is

crucial to develop highly effective drugs against *Babesia*. Cipargamin (CIP) has shown

inhibition against apicomplexan parasites, mainly *Plasmodium* and *Toxoplasma*. This study

evaluated the growth-inhibiting properties of CIP against *Babesia* spp. and investigated the

mechanism of CIP on *B. gibsoni*. The half inhibitory concentration (IC_{50}) values of CIP

against the *in vitro* growth of *B. bovis* and *B. gibsoni* were 20.2 ± 1.4 nM and 69.4 ± 2.2 nM,

respectively. CIP significantly inhibited the growth of *B. microti* and *B. rodhaini in vivo.*

Resistance was conferred by L921V and L921I mutations in BgATP4, which reduced the

sensitivity to CIP by 6.1- and 12.8-fold. The inhibitory potency of CIP against

BgATP4-associated ATPase activity was moderately reduced in mutant strains, with a

1.3-fold and 2.4-fold decrease in $BgATP4^{L921V}$ and $BgATP4^{L921I}$, respectively, compared to

that of $BgATP4^{WT}$. An *in silico* investigation revealed reductions in affinity for CIP binding to

$BgATP4^{L921V}$ and $BgATP4^{L921I}$ compared to $BgATP4^{WT}$. Resistant strains showed no

significant cross-resistance to atovaquone (ATO) or tafenoquine succinate (TQ), with less

than a onefold change in IC_{50} values. Combining CIP with TQ effectively eliminated *B.*

microti infection in SCID mice with no relapse, and parasite DNA was not detected by qPCR

within 90 days post-infection. Our findings reveal the efficacy of CIP as an anti-babesial

agent, its limitations as a monotherapy due to resistance development, and the potential of

combination therapy with TQ to overcome said resistance and achieve complete parasite

clearance.

Keywords: Cipargamin; *Babesia* spp.; drug resistance; ATP4; mutation

Introduction

Babesia is an apicomplexan tick-transmitted hemoparasite, which not only impacts the livestock economy but also causes an emerging disease in humans (Jalovecka et al., 2019). *Babesia* is a global pathogen that is more prevalent in certain regions such as Asia, Europe, and North America. However, as climate change brings with it higher temperatures and humidity, ticks and their reservoir hosts are anticipated to expand northward for survival and activity (Gray and Ogden, 2021). The disease is known as babesiosis, commonly characterized by fever and hemolytic anemia, but chronic infections can be asymptomatic (Almazán et al., 2022). The fatality rate ranges from 1% among all cases to 3% among hospitalized cases, and as high as 20% in immunocompromised patients (Krause, 2019).

Currently, the most common drugs for the treatment of human babesiosis include a combination of atovaquone (ATO) plus azithromycin (AZI) or clindamycin (CLN) plus quinine (QUI) (Krause et al., 2021). Still, these recommended treatments often result in various problems. For instance, multiple mutations in the *B. microti* cytochrome b, which is targeted by ATO, were identified in patients with relapsing babesiosis (Holbrook et al., 2023; Krause et al., 2024; Lemieux et al., 2016; Marcos et al., 2022; Rogers et al., 2023; Rosenblatt et al., 2021; Simon et al., 2017). The combination of CLN and QUI is the last resort for patients with severe symptoms (Kletsova et al., 2017). Despite its efficacy, this combination can elicit adverse drug reactions (Vannier and Krause, 2012). The 8-aminoquinoline analog tafenoquine (TQ) was found to be effective in curing immunocompromised patients experiencing relapsing babesiosis caused by *B. microti* (Rogers et al., 2023). However, the efficacy of TQ treatment may vary between individuals and cases and its contraindication in patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency limits its clinical use (Chu and Freedman, 2019). Due to the aforementioned issues, it is undoubtedly urgent to search for compounds that treat infections caused by *Babesia* spp. while simultaneously

minimizing the detrimental side effects of antibabesial drugs.

Spiroindolone cipargamin (CIP), a promising antimalarial, has been found to effectively

suppress the growth of all strains of *Plasmodium falciparum* and *P. vivax* with potency in the

low nanomolar ranges (Rosling et al., 2018; Rottmann et al., 2010). The assessment of the

drug indicated that CIP taken orally had good absorption, a long half-life, and exceptional

bioavailability (Schmitt et al., 2022). Following oral CIP treatment (30 mg daily for three

days) in adults with simple *P. falciparum* or *P. vivax* malaria, parasitemia was rapidly cleared

in a phase II trial (Schmitt et al., 2022). Currently, a clinical trial is being conducted for the

intravenous administration of CIP as a treatment for severe malaria patients (ClinicalTrials,

2024). CIP was also evaluated for treating toxoplasmosis as a second-line medicine in cases

where there are intolerable toxicity issues or allergies to the currently used treatments. Mice

infected with *Toxoplasma gondii* that were treated with CIP on the day of infection and the

following day had 90% fewer parasites five days post-infection (Zhou et al., 2014).

*Pf*ATP4, a P-type ATPase in *P. falciparum*, functions as a Na^+/H^+ transporter critical for

maintaining ionic balance (Spillman et al., 2013). Mutations in *Pf*ATP4 confer resistance to

CIP, which disrupts Na^+ homeostasis, leading to increased cytosolic Na^+ concentration and

various physiological changes, such as cytosolic alkalinization, osmotic swelling, etc.

(Mohring et al., 2022). These mutations reduce the Na^+-dysregulating effects of CIP and the

resting Na^+ levels. While *Pf*ATP4 is essential for *P. falciparum* survival, its homolog in *T.*

gondii (*Tg*ATP4) is less critical, as *T. gondii* experiences brief, high Na^+ exposure and can

survive without *Tg*ATP4 expression (Lehane et al., 2019).

Due to its excellent efficacy against other apicomplexan parasites, including *Plasmodium*

spp., and its ability to target parasite ATP4—a conserved protein essential for ion homeostasis

in apicomplexan organisms—we hypothesize that CIP could also effectively inhibit *Babesia*

spp., a related apicomplexan parasite. Therefore, the objectives of this study were to

determine whether CIP could inhibit the growth of *Babesia* spp., namely *B. bovis* and *B.*

gibsoni in vitro and *B. microti* and *B. rodhaini in vivo*, and identify the inhibitory mechanisms

of CIP on *Babesia* parasites.

Results

Inhibitory efficacy of CIP on *B. bovis* and *B. gibsoni in vitro*

In vitro efficacy of CIP against *B. bovis* and *B. gibsoni* showed a steep growth inhibition

curve with half inhibitory concentration (IC_{50}) values of 20.2 ± 1.4 nM (Figure 1A) and 69.4

± 2.2 nM (Figure 1B), respectively. The 50% cytotoxic concentration (CC_{50}) value of CIP on

Madin-Darby canine kidney (MDCK) cells and human foreskin fibroblasts (HFF) were 38.7

± 2.0 μM and 70.8 ± 4.9 μM (Figure supplement 1), respectively. Based on these values, the

predicted selectivity indices (SIs), which reflect the drug's safety and specificity, were

calculated to be greater than 500. Furthermore, at a concentration of 100 μM, CIP exhibited a

low erythrocyte hemolysis rate of 0.11 ± 0.03 % (data not shown).

CIP effect on *B. microti* and *B. rodhaini* infections *in vivo*

Concurrently, CIP showed effective inhibition on *B. microti* and *B. rodhaini in vivo*. The

parasitemia of *B. microti*-infected BALB/c mice increased dramatically in the vehicle-treated

control group and peaked at 10 days postinfection (DPI) ($38.55 \pm 4.32\%$) (Figure 1C). On the

other hand, seven days of treatment with CIP (20 mg/kg) or atovaquone (ATO) plus

azithromycin (AZI) administered orally resulted in a significantly lower peak parasitemia,

$1.06 \pm 0.20\%$ and $1.61 \pm 0.20\%$, respectively (Figure 1C). Hematocrit (HCT) variations were

tracked every 4 days as an indicator of anemia in *B. microti*-infected mice. The

vehicle-treated group showed a drop in HCT levels at 12 DPI and 16 DPI ($P < 0.01$) (Figure

1D). No significant reduction in HCT levels was observed in the CIP-treated group or the

ATO plus AZI-treated group (Figure 1D). This indicates that the administration of CIP could

control *B. microti* infection and prevent anemia from developing in *B. microti*-infected mice.

BALB/c mice infected with *B. rodhaini* treated with sesame oil or ATO plus AZI showed high

parasitemia, $90.73 \pm 1.97\%$, and $86.23 \pm 3.06\%$, respectively (Figure 1E), and all mice died

within 7 DPI (Figure 1F). CIP treatment in *B. rodhaini*-infected mice precluded the

emergence of parasitemia for the following eight days (Figure 1E), which led to 66.67% of

mice surviving the challenge infection (Figure 1F). At 12 DPI, parasites had recurred in all

CIP-treated *B. rodhaini*-infected mice ($10.32 \pm 15.51\%$), which were eventually cleared as

indicated by undetectable parasites at 18 DPI (Figure 1E).

Identification of *B. gibsoni* ATP4 mutation in CIP-resistant strains

After being exposed to CIP at increasing concentrations up to 10 times the IC_{50}, the resistant

parasites in two of the culture wells were able to regrow. We sequenced the *B. gibsoni* ATP4

gene from the wild-type and two resistant strains. The wild-type strain has a C at nucleotide

2,761, which translates to leucine (Figure 2A). In one resistant strain, a single nucleotide

variant (SNV) in BgATP4 with a substitution at position 2,761 (from C to G) was found − a

nonsynonymous coding change from leucine to valine (L921V) (Figure 2B). In another

resistant strain, the mutation occurred in the same position. However, the nucleotide

substitution was from C to A, and the coding changed from leucine to isoleucine (L921I)

(Figure 2C). Next-generation sequencing (NGS) revealed that for $Bgatp4^{2761C>G}$, 99.97% of

7,960 reads were G at nucleotide 2,761, and for $Bgatp4^{2761C>A}$, 99.92% of 7,862 reads were A

at nucleotide 2,761 (Figure 2D). $BgATP4^{L921V}$ and $BgATP4^{L921I}$ lines were tested for their

susceptibility to CIP, and had IC_{50} values of 421.0 ± 15.9 nM and 887.9 ± 62.0 nM,

respectively (Figure 2E). These findings demonstrate a 6.1- and 12.8-fold reduction in CIP

sensitivity of the resistant parasite lines $BgATP4^{L921V}$ and $BgATP4^{L921I}$.

The effect of CIP on $BgATP4^{WT}$, $BgATP4^{L921V}$ and $BgATP4^{L921I}$ function

Microscopic observation of thin blood smears was performed to determine the morphological

changes of *B. gibsoni* exposed to CIP. The CIP-treated parasites became swollen after incubation with the drug for 72 h (Figure 3A). For both the treatment and control groups, one hundred parasites were measured. The mean size of treated parasites was notably bigger than the parasites in the untreated group ($P < 0.0001$) (Figure 3B). Significant vacuolization was observed in the cytoplasm of parasites in the CIP-treated group, as revealed by transmission electron microscopy (TEM). Despite this, the nuclear membrane structure and parasitic membranes remained intact until the parasites were completely destroyed (Figure 3C). The addition of the ATP4 inhibitor CIP resulted in a time-dependent increase in the concentrations of $[Na^+]_i$ in wild-type *B. gibsoni*, with improved signal-to-noise ratios at the higher drug concentration of 20 nM (Figure 3D). We observed that the Na^+ concentrations in both BgATP4^{L921V} and BgATP4^{L921I} lines were lower when compared with those of the control BgATP4^{WT} line after being exposed to 20 nM CIP for 20 min, with a significantly lower Na^+ concentration in BgATP4^{L921I} ($P = 0.0087$) (Figure 3F). We also demonstrated here that the addition of CIP in wild-type *B. gibsoni* caused an increase in the cytosolic pH (Figure 3E). Specifically, 4 min after the drug was added, the average pH of the 20 nM CIP group reached as high as 7.278, while the 1 nM CIP group reached 7.089 and the untreated group reached 7.062 (Figure 3E). The pH values of the 20 nM CIP group were consistently higher than those of the other two groups, although declining with time (Figure 3E). In resistant lines, a 20-min exposure to 20 nM CIP caused small changes in the pH values compared to the wild-type line, with the BgATP4^{L921I} line (7.048 ± 0.042) having a notably lower pH value ($P = 0.0229$) (Figure 3G).

Sensitivity of BgATP4-associated ATPase activity to CIP in BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I}

The BgATP4-associated ATPase activity in erythrocytes infected with BgATP4^{WT} (6.31 ± 1.20 nmol Pi/mg protein/min), measured in the presence of 150 mM Na^+, was higher than

those observed in BgATP4^{L921V} (5.11 ± 0.50 nmol Pi/mg protein/min) and BgATP4^{L921I} (4.58

± 0.53 nmol Pi/mg protein/min) ($P = 0.04$) (Figure 3H).

We further investigated the concentration-dependent inhibition of BgATP4-associated ATPase

activity by CIP in wild-type and mutant parasites. In membranes prepared from *B.gibsoni*,

CIP inhibited BgATP4-associated ATPase activity with IC$_{50}$ values of 111.4 ± 31.8 nM for

BgATP4^{WT}, 149.8 ± 21.7 nM for BgATP4^{L921V}, and 269.5 ± 29.8 nM for BgATP4^{L921I}. The

potency of CIP in inhibiting BgATP4-associated ATPase activity was reduced by 1.3-fold and

2.4-fold in membranes prepared from BgATP4^{L921V} and BgATP4^{L921I}, respectively, compared

to BgATP4^{WT} (Figure 3I).

Multiple sequence alignment of *Babesia* ATP4 and molecular docking

The whole amino acid sequence of *B. gibsoni* ATP4 (GenBank: KAK1443404.1) shared

identity values of 29.75%, 49.40%, 49.67%, 62.21%, and 52.47% with *Homo sapiens* ATP4

(GenBank: NM_000704.3), *P. falciparum* ATP4 (GenBank: PF3D7_1211900), *T. gondii*

ATP4 (GenBank: XP_018635122.1), *B. bovis* ATP4 (PiroplasmaDB: BBOV_IV010020), and

B. microti ATP4 (GenBank: BMR1_03g01005), respectively (Figure supplement 2).

The pLDDT (predicted Local Distance Difference Test) value of BgATP4^{WT} prediction was

80.7 using Colab-fold. Multiple potential binding sites for CIP were revealed by blind

docking throughout the whole protein surface (Figure supplement 3). CIP binds in close

proximity to L921, as demonstrated by focused docking on this area (Figure 4A). The

contribution of each residue to the predicted binding affinity in either mutant structure was

reduced; the precise values of BgATP4^{WT} (Figure 4B), BgATP4^{L921V} (Figure 4C), and

BgATP4^{L921I} (Figure 4D) were -6.43, -6.40 and -6.26 kcal/mol, respectively. The interactions

of CIP from each docking simulation are shown in Table supplement 3.

Cross-resistance of BgATP4^{L921V} and BgATP4^{L921I} mutants to ATO and TQ

The BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I} lines were tested for their susceptibility to

ATO and TQ. The IC_{50} values for these lines were 380.1 ± 3.1 nM, 412.9 ± 5.4 nM, and 360.3

± 7.9 nM, respectively, for ATO (Figure 5A), and 39.5 ± 0.4 μM, 29.9 ± 1.4 μM, and $39.3 \pm$

0.3 μM, respectively, for TQ (Figure 5B). The IC_{50} values for both ATO and TQ in the

resistant strains showed only slight changes compared to the wild-type strain, with less than a

onefold difference. This minimal variation suggests that the resistant strain has a mild

alteration in susceptibility to ATO and TQ, but not enough to indicate strong resistance or

significant cross-resistance.

Combination treatment of CIP plus TQ in SCID mice with *B. microti* infection

The parasitemia of *B. microti*-infected SCID mice in the vehicle group increased dramatically,

peaking at 10 DPI ($77.03 \pm 2.45\%$) (Figure 5C). Although the trend showed a subsequent

decline, the parasitemia remained around 50% until the mice were euthanized at 90 DPI.

Treatment with CIP (20 mg/kg) initially reduced parasitemia to undetectable by 18 DPI.

However, a relapse occurred, with parasitemia increasing rapidly and stabilizing at levels

comparable to the vehicle group during the subsequent observation period. Notably, no

mutations were detected in the relapsed *B. microti* parasites from SCID mice (data not

shown). In contrast, parasitemia was completely cleared in the TQ and CIP plus TQ groups

by 8 DPI and 6 DPI, respectively, with no parasites observed under the microscope thereafter.

qPCR analysis at 90 DPI detected parasite DNA in all groups except the CIP plus TQ group

(Figure 5D).

Discussion

The repositioning of antimalarial drugs is critical in developing novel strategies for treating

babesiosis. CIP is a novel compound that inhibits *Plasmodium* development by targeting

ATP4 and has been extensively tested in Phase 1 and Phase 2 clinical trials (Qiu et al., 2022).

Since ATP4 is conserved across apicomplexans, including *Babesia* species, our study aimed

to repurpose the antimalarial CIP by assessing its efficacy on *Babesia* species (Lehane et al.,

2019). The IC_{50} values of CIP against *B. bovis* and *B. gibsoni in vitro* were lower than the

previously reported IC_{50} value of TQ (Carvalho et al., 2020; Ji et al., 2022b) and ATO against

B. gibsoni (Matsuu et al., 2004). The present investigation also demonstrated that the

inhibitory effects of CIP on *B. microti*-infected BALB/c mice were comparable to that of

ATO plus AZI, the combination recommended by the CDC in the United States. CIP was also

proven to protect mice from the deadly *B. rodhaini* infection, with a survival rate of up to

67%, as recorded in the current trial. These results suggest that CIP may be a potential

chemotherapy candidate for babesiosis.

In a previous study, the *P. falciparum* Dd2 strain that acquired resistance to CIP carried the

G358S mutation in the *Pf*ATP4 protein, a *P. falciparum* P-type Na^+ ATPase (Qiu et al., 2022).

The vital protein ATP4 is found in the parasite plasma membrane and is specific to the

subclass of apicomplexan parasites (Mohring et al., 2022). To date, *in vitro* evolution

experiments using CIP have produced at least 18 parasite lines with various *Pf*ATP4

mutations (Lee and Fidock, 2016; Rottmann et al., 2010). In another study involving *T.*

gondii, a cell line that carried the mutation G419S in the *Tg*ATP4 gene was 34 times less

susceptible to CIP than that of *Tg*ATP4[WT] (Qiu et al., 2022). It follows that there is a

significant possibility that resistant *Babesia* parasites will emerge from CIP exposure. In this

study, we successfully produced two CIP-resistant strains using six independent selections.

The newly discovered L921V and L921I mutations in BgATP4 decreased the CIP sensitivity

by 6.1 and 12.8 times, respectively. We provided compelling evidence that natural variety

mutations L921V and L921I in BgATP4 significantly affected the protein's susceptibility to

CIP inhibition, albeit the mutational sites were different from those of *P. falciparum* and *T.*

gondii. Based on our observation, the growth and generation rates of the mutant strains are

comparable to those of the wild-type strain.

Although ATP4 was initially recognized as a Ca^{2+} transporter (Krishna et al., 2001), current

evidence suggests that ATP4 functions as an ATPase for exporting Na^+ while importing H^+

(Mohring et al., 2022), as well as causing a variety of other physiological perturbations

including an increase in the volume of parasites and infected erythrocytes due to the osmotic

impact of the $[Na^+]_{cyt}$ increase (Dennis et al., 2018), a decline in cholesterol extrusion from

the parasite plasma membrane as a result of the increase in $[Na^+]_{cyt}$ (Das et al., 2016), and an

intensified rigidity of erythrocytes infected with ring-stage parasites (Zhang et al., 2016). In

this study, the CIP-exposed wild-type *B. gibsoni* became swollen, which was identical to a

prior study on *P. falciparum* (Dennis et al., 2018). Interestingly, transmission electron

microscopy (TEM) analysis of parasites incubated with CIP revealed ultrastructural

alterations characterized by significant vacuole formation in the cytoplasm, a hallmark of

stress or early stages of cell death. These changes were similar to those observed in *B. bovis*

treated with clotrimazole and ketoconazole, which ultimately led to growth inhibition or

parasite death (Bork et al., 2003). Despite the vacuolization, the structural integrity of the

nucleus and membranes was maintained, suggesting a gradual process where vacuolization

precedes complete destruction of the parasite. One explanation is the swelling of the isolated

parasites, which can be ascribed to the osmotic consequences of Na^+ uptake and is contingent

upon the presence of Na^+ in the external environment.

The output of Na^+ and input of H^+ diminished upon CIP-induced inhibition of *Pf*ATP4, and

the continued outflow of H^+ via V-type H^+-ATPase led to an alkalinization that ultimately

killed the parasites (Spillman et al., 2013). Our capacity to measure a time-dependent

increase in the concentration of $[Na^+]_i$ and pH value for wild-type *B. gibsoni* facilitated us to

gain insight into the basic mechanisms of CIP on BgATP4^{WT} function. Furthermore, our

results corroborate internal alkalinization as the main factor in *Babesia* death and support our

hypothesis that the swollen isolated parasites were produced by Na^+ absorption. To explore

further how mutations in BgATP4 are associated with the upregulation of the parasite's $[Na^+]_i$

and $[H^+]_i$, we tested two BgATP4-mutant lines that were chosen previously with BgATP4

inhibitors (BgATP4^{L921V} and BgATP4^{L921I}). Due to the presence of L921V and L921I

mutations in drug-resistant strains of BgATP4, the concentration of Na^+ did not increase as

much as it would have in the wild-type strain following the addition of 20 nM CIP for 20 min.

As intraerythrocytic alkalinization rises, the same outcome happens. From these results, we

deduced how natural mutations in BgATP4 may affect ATP4 inhibitor susceptibility by

dysregulating H^+ and Na^+ balance, which helped parasites survive in a relatively high

concentration of CIP. An illustration of the putative processes for regulating Na^+ and H^+ in

erythrocytes infected with the *Babesia* parasite is presented in Figure supplement 4.

In a previous study's isolated membrane assay, CIP was found to be more effective at

inhibiting parasite development than *Pf*ATP4-associated ATPase activity and reduced

sensitivity of *Pf*ATP4-associated ATPase activity to the medication was associated with a

decrease in parasite susceptibility to CIP (Rosling et al., 2018). In our study, the higher

resistance to CIP-mediated inhibition of parasite proliferation observed in BgATP4^{L921I}

compared to BgATP4^{L921V} correlated with a higher resistance of BgATP4-associated ATPase

activity to the drug. These findings provide additional evidence that the ATPase activity

measured in this study corresponds to the activity of the BgATP4 protein.

According to studies using the *Pf*ATP4 model for molecular docking, the G358S mutation

results in a steric clash that lowers CIP's binding affinity (Qiu et al., 2022). The results from

the current study were similar to the *Pf*ATP4 model. The molecular docking was constructed

using a ColabFold model of wild-type BgATP4, which predicted the binding mode and

affinity between ATP4 protein and the ligand to provide a possible mechanistic explanation. It

suggested that the L921V mutation caused changes at the atomic level, whereas the L921I

mutation created a steric clash that reduced the binding affinity of CIP. The predicted affinity

score of the L921I mutation was lower than that of the L921V mutation. Thus, it is possible

that CIP had a weaker binding to BgATP4^{L921I} than to BgATP4^{L921V}. These findings are

consistent with the results obtained from measuring the IC$_{50}$ of the drug against parasites with

the L921V and L921I mutations.

While CIP shows promise as an antibabesial agent, the emergence of resistance emphasizes

the importance of rational drug use and the development of combination therapies. Our

results demonstrate that CIP did not exhibit cross-resistance with ATO or TQ, as IC$_{50}$ values

for ATO and TQ in resistant strains showed minimal changes (<1-fold) compared to the

wild-type strain. This observation suggests that CIP could be effectively combined with other

antibabesial drugs to reduce the risk of resistance development. Previous studies have also

reported that a single dose of TQ may not be sufficient to prevent parasite relapse in

immunocompromised hosts, recommending either repeated TQ administration or

combination therapy with other antibabesial agents to address this issue (Mordue et al., 2019).

Supporting this approach, our study demonstrates that the combination of CIP plus TQ

effectively cleared *B. microti* infections in SCID mice, with no detectable parasitemia or

DNA observed at 90 DPI, indicating complete parasite elimination.

This preclinical finding underscores the potential of the CIP plus TQ combination as a

viable treatment strategy for babesiosis, especially in cases where monotherapy is ineffective.

Further studies are needed to fully explore the pharmacokinetics, long-term efficacy, and

potential side effects of such combinations in clinical settings, ultimately facilitating

optimized therapeutic strategies.

In summary, our findings provide a comprehensive understanding of CIP's efficacy,

mechanism of resistance, and identifying strategies to enhance its therapeutic potential. The

combination of CIP with drugs like TQ offers a promising approach to combat babesiosis and

mitigate the development of resistance.

Materials and methods

Parasite culture

The parasites *B. gibsoni* Oita strain and *B. bovis* Texas strain were *in vitro* cultured in 24-well plates and maintained in an atmosphere of 5% CO_2 and 5% O_2 at 37 °C (Liu et al., 2018). For the *in vivo* studies, *B. microti* Peabody mjr strain-(ATCC PRA-99™) and *B. rodhaini* Australia strain-infected RBCs (iRBCs), which were collected and diluted with phosphate-buffered saline (PBS) when the parasitemia levels in the donor mice reached ~20% and 50%, respectively, and were intraperitoneally injected into BALB/c mice. Each BALB/c mouse was infected with 1.0×10^7 *B. microti* or *B. rodhaini* iRBCs for the *in vivo* trials (Ji et al., 2022a).

In vitro cytotoxicity of CIP and hemolysis rate in canine erythrocytes

Cultures of Madin-Darby canine kidney (MDCK) cells and human foreskin fibroblasts (HFF) were maintained at 37 °C under an atmosphere of 5% CO_2 and 5% O_2 and the cytotoxic effect of CIP (MedChem Express, Tokyo, Japan) was assessed using a cell viability assay by CCK-8 (Dojindo, Japan) as described previously (Li et al., 2023). The selectivity index is calculated as the ratio between the half maximal inhibitory concentration (IC_{50}) and the 50% cytotoxic concentration (CC_{50}) values.

Canine erythrocytes were collected from healthy beagle dogs raised in NRCPD and stocked in Vega y Martinez (VYM) phosphate-buffered saline solution at 4 °C (Vega et al., 1985). A canine erythrocyte hemolysis assay was performed at concentrations of 0.1, 1, 5, 10, 25, 50, and 100 μM as previously described (Ariefta et al., 2022).

Evaluation of the efficacy of CIP against *Babesia* parasites *in vitro*

The efficacy of CIP against *B. gibsoni* and *B. bovis* was determined using a fluorescence assay, as previously described (Guswanto et al., 2014). The IC_{50} values were determined from the fluorescence values and by non-linear regression analysis (curve fit) in GraphPad Prism 9

351 (GraphPad Software Inc., USA).

Chemotherapeutic effects of CIP against *Babesia* infections *in vivo*

CIP was evaluated on *B. microti*- and *B. rodhaini*-infected mice, as previously described

(Ndayisaba et al., 2021; Tuvshintulga et al., 2022). When *B. microti*- and *B. rodhaini*-infected

mice had a 1% average parasitemia at 4 and 2 days postinfection (DPI), respectively, the drug

treatments were administered and continued for seven days. Three groups of *B.*

microti-infected mice were administered different treatments. The CIP group (n = 6) and the

ATO plus AZI group (n = 6) were orally treated with 20 mg/kg CIP and 20 mg/kg ATO plus

20 mg/kg AZI (Sigma, Tokyo, Japan), respectively. All drugs were prepared in sesame oil.

The infected mice of the vehicle group (n = 6) orally received 0.2 mL of sesame oil as the

control. Eighteen mice infected with *B. rodhaini* were likewise placed into three groups for

treatment, and they received the same treatments as the mice infected with *B. microti*. A light

microscope (Nikon, Japan) and a hematology analyzer (Celltac α MEK-6450, Nihon Kohden

Corporation, Tokyo, Japan) were used to assess the parasitemia and hematocrit levels every

two and four days, respectively.

Selection of CIP-resistant *B. gibsoni in vitro*

Selections were initiated by exposing six independent flasks, each containing 10 μL (5×10^6)

B. gibsoni iRBCs mixed with 40 μL (4×10^8) RBCs into a 450 μL culture medium, which

contained increasing concentrations of CIP: 5, 10, 20, 30 to 694 nM ($10\times IC_{50}$). The medium

containing CIP was replaced daily until parasites treated with $10\times IC_{50}$ CIP reached

multiplication rates that were approximately comparable to those of the untreated controls

(Hwang et al., 2010). Then, to evaluate the decreased sensitivity to CIP, IC_{50} values of

resistant strains were determined by nonlinear regression using the GraphPad Prism software.

Detection of *B. gibsoni* ATP4 gene mutations

The genomic DNA of the mutant parasites was extracted and sequenced (Ji et al., 2022a). The

primer sets used for sequencing are listed in Table supplement 1. The single nucleotide

variants were identified by pairwise alignment to the BgATP4^{WT} sequence (GenBank:

JAVEPI010000002.1). Next-generation sequencing (NGS) was performed to detect the ratio

of wild-type to mutant parasites by using the Illumina NovaSeq6000 sequencing platform

(Jeon et al., 2021).

Morphological changes in CIP-treated *in vitro* cultured *B. gibsoni*

A microscopy assay was used to detect the morphological changes of wild-type *B. gibsoni*

after exposure to 50 nM CIP for three consecutive days (Liu et al., 2021; Tayebwa et al.,

2018). At 4 days post-treatment, ImageJ software was used to measure the sizes of 100

randomly selected parasites in the CIP-treated group and the control group on Giemsa-stained

blood smears. After treating the parasites as described above, iRBCs were fixed in GA

fixation buffer (2% glutaraldehyde in 0.1 M sodium cacodylate buffer containing 1 mM

$CaCl_2$ and 1 mM $MgCl_2$), then stored in rinse buffer (0.1 M sodium cacodylate buffer

containing 1 mM $CaCl_2$ and 1 mM $MgCl_2$) at 4 ºC for transmission electron microscopy

(TEM) analysis (Hidayati et al., 2023).

Parasites $[Na^+]_i$ and pH_i measurements

For both $[Na^+]_i$ and pH_i measurements, the wild-type and mutant parasites were initially

separated from erythrocytes by treatment with saponin (0.05% [wt/vol]) for 5 min (Saliba and

Kirk, 1999). The Na^+-sensitive fluorescent dye SBFI (Thermo Fisher Scientific; product

S1263) was used to quantify intracellular sodium ($[Na^+]_i$). Saponin-isolated parasites (at

1×10^8 parasites/mL) were loaded with SBFI (5 μM; in the presence of 0.02% w/v Pluronic

F127) for 1 hr at room temperature (RT) (Spillman et al., 2013). Thereafter, SBFI-loaded

parasites were resuspended in physiological saline (120 mM NaCl, 5 mM KCl, 25 mM

HEPES, 20 mM D-glucose, and 1 mM $MgCl_2$ [pH 7.1]) at RT in the presence or absence of

CIP. A 96-well microtiter plate was filled with around 200 μL of parasite suspension per well.

The dye-loaded cells fluoresced at 515 nm after being stimulated at 340 and 380 nm.

Parasites loaded with SBFI were suspended in calibration buffers containing [Na^+] values

ranging from 0 to 140 mM (pH 7.1) to establish calibration curves (Diarra et al., 2001).

The cytosolic pH of wild-type and mutant strains was measured using the pH-sensitive

fluorescent dye BCECF [2',7'-bis-(2-carboxyethyl)-5-(and-6)- carboxyfluorescein] (Biotium;

product 51011). The BCECF was added to saponin-isolated parasites by suspension (1×10^8

parasites/mL) and incubated for 20 minutes at 37 °C in RPMI-1640 culture medium (Gibco,

USA) (Saliba and Kirk, 1999). Thereafter, the parasites loaded with dye were rinsed thrice

(12,000 × g, 1 min) in RPMI-1640 culture medium, then resuspended in physiological saline

(as previously mentioned) with or without various concentrations of CIP. A 96-well microtiter

plate was filled with around 200 μL of parasite suspension per well. The dye-loaded cells

fluoresced at 520 nm after being stimulated at 440 and 490 nm. A pH calibration was carried

out for each experiment (Mohring et al., 2022).

Measurements of membrane ATPase activity

Membranes from isolated *B. gibsoni* were prepared by lysing the parasites in ice-cold

deionized water containing 7 × Protease Inhibitor Cocktail Tablets (Roche, Germany). The

membrane preparation was then washed three times with ice-cold deionized water, with

protease inhibitors included in the first two washes (Qiu et al., 2022). Protein concentrations

in the membrane samples were determined using a Bradford assay. The production of

inorganic phosphate (Pi) from ATP hydrolysis was measured using the Malachite Green

Phosphate Assay (BioAssay Systems, USA). Membrane preparations were diluted in either a

high Na^+ solution (final reaction conditions: 150 mM NaCl, 20 mM KCl, 2 mM $MgCl_2$, 50

mM Tris, pH 7.2) or a Na^+-free solution (final reaction conditions: 150 mM choline chloride,

20 mM KCl, 2 mM $MgCl_2$, 50 mM Tris, pH 7.2) to achieve a final protein concentration of

40 μg/mL. CIP was added at the concentrations specified in the respective figure legends.

Reactions were conducted according to the manufacturer's protocol provided with the assay

kit.

Multiple ATP4 sequence alignment and molecular docking

B. gibsoni ATP4 (GenBank: KAK1443404.1), *B. bovis* ATP4 (PiroplasmaDB:

BBOV_IV010020), *B. microti* ATP4 (GenBank: BMR1_03g01005), *T. gondii* ATP4

(GenBank: XP_018635122.1), and *Homo sapiens* ATP4 (GenBank: NM_000704.3)

sequences were obtained by a homology search using *P. falciparum* ATP4 (GenBank:

PF3D7_1211900). Sequence alignment was analyzed using MUSCLE in Jalview v2.11.3.2

software and BLAST (http://www.ncbi.nlm.nih.gov/BLAST/).

AlphaFold was used to predict the structure of BgATP4^{WT} (Jumper et al., 2021). Using the

GROMACS 2021 Molecular Dynamics package, the energy minimization was carried out

following the model's generation (Lindahl et al., 2021). The mutations were produced by

using Charmm-GUI PDB reader (Jo et al., 2014). PyMOL (version 2.0 Schrödinger, LLC)

was used to confirm the position of the mutation site (center: -20.367, 10.904, 9.435; size:

30×30×30) (Trott and Olson, 2010). The ligand molecules CIP was downloaded from

PubChem (CID 44469321 (https://pubchem.ncbi.nlm.nih.gov/compound/44469321). The

ligand was used to dock with Gnina (Eberhardt et al., 2021). The affinity score and binding

pose were chosen only from the highest CNN (convolutional neural network) score results

from docking simulations. The models were visualized using the PyMOL Molecular Graphic

System and Discovery Studio.

Cross-resistance to TQ and ATO in mutant parasites

447 The efficacy of TQ and ATO against BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I} was

448 evaluated using a fluorescence assay, as described above. The IC$_{50}$ values were calculated

449 from the fluorescence values using non-linear regression analysis (curve fitting) in GraphPad

450 Prism 9 (GraphPad Software Inc., USA).

Efficacy of CIP plus TQ combination therapy in SCID mice infected with *B. microti*

Twenty 6-week-old female immunocompromised mice (C.B-17/IcrJcl-Prkdcscid strain, CLEA Japan, Inc.) were intraperitoneally injected with 1.0×10^7 of *B. microti*-infected blood cells (Tuvshintulga et al., 2022). When the average parasitemia across all mice reached 1% (at 4 DPI), the drug treatments were initiated. The CIP group (n = 5) was orally administered 20 mg/kg CIP for seven days. The TQ group (n = 5) was orally administered a single dose of 20 mg/kg TQ. The CIP plus TQ group (n = 5) received the combination treatment, following the dosage and administration methods described above. The vehicle group (n = 5) orally received 0.2 mL of sesame oil as the control. Thin blood smears were prepared every other day and examined under a light microscope to determine parasitemia levels.

qPCR analysis

To further verify the presence or absence of *B. microti* DNA in treated SCID mice, real-time quantitative PCR analysis was performed as described previously (Vydyam et al., 2024). Briefly, genomic DNA was extracted from blood samples (200 μl) collected at 90 DPI and subjected to qPCR analysis using SYBR Green I, targeting a highly conserved region of *Babesia* mitochondrial genome (mtDNA). Primers used were: Bmic-F 5′-TTGCGATAGTAATAGATTTACTGC-3′ and B-lsu-R2 5′-TCTTAACCCAACTCACGTACCA-3′ (Qurollo et al., 2017). The reaction mixture consisted of: $1 \times$ Advanced Universal SYBR Green Super Mix ($2 \times$) (1725270; Bio-Rad); 0.5 μM of each primer, and 100 ng of genomic DNA.

Detection of *B. microti* ATP4 gene mutations in relapsed infected SCID mice

The genomic DNA of the mutant parasites was extracted and sequenced (Ji et al., 2022a). The primer sets used for sequencing are listed in Table supplement 2. The single nucleotide variants were confirmed by pairwise alignment to the *Bm*ATP4^{WT} sequence (GenBank: BMR1_03g01005).

Statistical analysis

Data analysis, namely one-way analysis of variance (ANOVA) and two-tailed unpaired t-tests, was performed using GraphPad Prism (La Jolla, CA, USA) version 9. A P value of <0.05 was considered a statistically significant result.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research (22H0250906), the JSPS Core-to-Core program, both from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and a grant from the Strategic International Collaborative Research Project (JPJ008837) promoted by the Ministry of Agriculture, Forestry, and Fisheries of Japan, and International Science and Technology Cooperation Project of Hubei Province (2024EHA018).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author contributions

Hang Li, Investigation, Data curation, Formal analysis, Writing – original draft, Writing – review & editing; Shengwei Ji, Methodology, Investigation, Software, Validation; Nanang R. Ariefta, Methodology, Software, Formal analysis; Eloiza May S. Galon, Visualization, Writing – review & editing; Shimaa AES El-Sayed, Methodology, Visualization; Thom Do, Methodology, Investigation; Lijun Jia, Data curation, Visualization; Miako Sakaguchi, Methodology, Visualization; Masahito Asada, Investigation, Visualization; Yoshifumi

Nishikawa, Methodology, Resources; Xin Qin, Methodology, Data curation; Mingming Liu, Conceptualization, Methodology, Investigation, Resources, Data curation, Funding acquisition, Supervision, Writing – review & editing; Xuenan Xuan, Conceptualization, Methodology, Validation, Formal analysis, Supervision, Project administration, Writing – review & editing.

Ethics

All animal experiments were performed according to the Guide for the Care and Use of Laboratory Animals of Obihiro University of Agriculture and Veterinary Medicine, which was also approved by the Committee on the Ethics of Animal Experiments at the Obihiro University of Agriculture and Veterinary Medicine, Japan (permit numbers: animal experiment, 22-145 and 23-132; DNA experiment, 2207 and 2208; pathogen, 202308 and 202306).

References

Almazán C, Scimeca RC, Reichard MV, Mosqueda J. 2022. Babesiosis and Theileriosis in North America. *Pathogens* **11**:168. doi:10.3390/pathogens11020168

Ariefta NR, Pagmadulam B, Nihei C, Nishikawa Y. 2022. Sparsomycin exhibits potent antiplasmodial activity *in vitro* and *in vivo*. *Pharmaceutics* **14**:544. doi:10.3390/pharmaceutics14030544

Bork S, Yokoyama N, Matsuo T, Claveria FG, Fujisaki K, Igarashi I. 2003. Clotrimazole, ketoconazole, and clodinafop-propargyl inhibit the *in vitro* growth of *Babesia bigemina* and *Babesia bovis* (Phylum Apicomplexa). *Parasitology* **127**:311–315. doi:10.1017/S0031182003003895

Carvalho LJM, Tuvshintulga B, Nugraha AB, Sivakumar T, Yokoyama N. 2020. Activities of artesunate-based combinations and tafenoquine against *Babesia bovis in vitro* and *Babesia microti in vivo*. *Parasites Vectors* **13**:362. doi:10.1186/s13071-020-04235-7

Chu CS, Freedman DO. 2019. Tafenoquine and G6PD: a primer for clinicians. *J Travel Med* **26**:taz023. doi:10.1093/jtm/taz023

ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). Identifier

NCT04675931, To evaluate efficacy, safety, tolerability and PK of intravenous cipargamin in participants with severe *Plasmodium falciparum* malaria; 2024 Feb 29 [cited 2024 Apr 9]. Available from: https://clinicaltrials.gov/study/NCT04675931

Das S, Bhatanagar S, Morrisey JM, Daly TM, Burns JM, Coppens I, Vaidya AB. 2016. Na^+ influx induced by new antimalarials causes rapid alterations in the cholesterol content and morphology of *Plasmodium falciparum*. *PLoS Pathog* **12**:e1005647. doi:10.1371/journal.ppat.1005647

Dennis ASM, Lehane AM, Ridgway MC, Holleran JP, Kirk K. 2018. Cell swelling induced by the antimalarial KAE609 (cipargamin) and other *Pf*ATP4-associated antimalarials. *Antimicrob Agents Chemother* **62**:e00087-18. doi:10.1128/AAC.00087-18

Diarra A, Sheldon C, Church J. 2001. In situ calibration and [H^+] sensitivity of the fluorescent Na^+ indicator SBFI. *American Journal of Physiology-Cell Physiology* **280**:C1623–C1633. doi:10.1152/ajpcell.2001.280.6.C1623

Eberhardt J, Santos-Martins D, Tillack AF, Forli S. 2021. AutoDock Vina 1.2.0: new docking methods, expanded force field, and python bindings. *J Chem Inf Model* **61**:3891–3898. doi:10.1021/acs.jcim.1c00203

Gray JS, Ogden NH. 2021. Ticks, human babesiosis and climate change. *Pathogens* **10**:1430. doi:10.3390/pathogens10111430

Guswanto A, Sivakumar T, Rizk MA, Elsayed SAE, Youssef MA, ElSaid EES, Yokoyama N, Igarashi I. 2014. Evaluation of a fluorescence-based method for antibabesial drug screening. *Antimicrob Agents Chemother* **58**:4713–4717. doi:10.1128/AAC.00022-14

Hidayati AR, Melinda, Ilmi H, Sakura T, Sakaguchi M, Ohmori J, Hartuti ED, Tumewu L, Inaoka DK, Tanjung M, Yoshida E, Tokumasu F, Kita K, Mori M, Dobashi K, Nozaki T, Syafruddin D, Hafid AF, Waluyo D, Widyawaruyanti A. 2023. Effect of geranylated dihydrochalcone from *Artocarpus altilis* leaves extract on *Plasmodium falciparum* ultrastructural changes and mitochondrial malate: Quinone oxidoreductase. *International Journal for Parasitology: Drugs and Drug Resistance* **21**:40–50. doi:10.1016/j.ijpddr.2022.12.001

Holbrook NR, Klontz EH, Adams GC, Schnittman SR, Issa NC, Bond SA, Branda JA, Lemieux JE. 2023. *Babesia microti* variant with multiple resistance mutations detected in an immunocompromised patient receiving atovaquone prophylaxi. *Open Forum Infectious Diseases* **10**:ofad097. doi:10.1093/ofid/ofad097

Hwang S-J, Yamasaki M, Nakamura K, Sasaki N, Murakami M, Wickramasekara Rajapakshage BK, Ohta H, Maede Y, Takiguchi M. 2010. Development and characterization of a strain of *Babesia gibsoni* resistant to diminazene aceturate *in vitro*. *J Vet Med Sci* **72**:765–771. doi:10.1292/jvms.09-0535

Jalovecka M, Sojka D, Ascencio M, Schnittger L. 2019. *Babesia* life cycle-when phylogeny meets biology. *Trends in Parasitology* **35**:356–368. doi:10.1016/j.pt.2019.01.007

Jeon SA, Park JL, Park S-J, Kim JH, Goh S-H, Han J-Y, Kim S-Y. 2021. Comparison
between MGI and Illumina sequencing platforms for whole genome sequencing. *Genes*
Genom **43**:713–724. doi:10.1007/s13258-021-01096-x

Ji S, Galon EM, Rizk MA, Yi Y, Zafar I, Li H, Ma Z, Iguchi A, Asada M, Liu M, Xuan X.
2022a. Efficacy of the antimalarial MMV390048 against *Babesia* infection reveals
phosphatidylinositol 4-kinase as a druggable target for babesiosis. *Antimicrob Agents*
Chemother **66**:e00574-22. doi:10.1128/aac.00574-22

Ji S, Liu M, Galon EM, Rizk MA, Tuvshintulga B, Li J, Zafar I, Hasegawa Y, Iguchi A,
Yokoyama N, Xuan X. 2022b. Inhibitory effect of naphthoquine phosphate on *Babesia*
gibsoni in vitro and *Babesia rodhaini in vivo*. *Parasites Vectors* **15**:10.
doi:10.1186/s13071-021-05127-0

Jo S, Cheng X, Islam SM, Huang L, Rui H, Zhu A, Lee HS, Qi Y, Han W, Vanommeslaeghe
K, MacKerell AD, Roux B, Im W. 2014. CHARMM-GUI PDB manipulator for advanced
modeling and simulations of proteins containing nonstandard residues. *Adv Protein Chem*
Struct Biol **96**: 235–265. doi:10.1016/bs.apcsb.2014.06.002

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A,
Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, et al. 2021. Highly accurate protein
structure prediction with AlphaFold. *Nature* **596**:583–589.
doi:10.1038/s41586-021-03819-2

Kletsova EA, Spitzer ED, Fries BC, Marcos LA. 2017. Babesiosis in Long Island: review of
62 cases focusing on treatment with azithromycin and atovaquone. *Ann Clin Microbiol*
Antimicrob **16**:26. doi:10.1186/s12941-017-0198-9

Krause PJ. 2019. Human babesiosis. *International Journal for Parasitology* **49**:165–174.
doi:10.1016/j.ijpara.2018.11.007

Krause PJ, Auwaerter PG, Bannuru RR, Branda JA, Falck-Ytter YT, Lantos PM, Lavergne V,
Meissner HC, Osani MC, Rips JG, Sood SK, Vannier E, Vaysbrot EE, Wormser GP. 2021.
Clinical practice guidelines by the infectious diseases society of America (IDSA): 2020
guideline on diagnosis and management of babesiosis. *Clinical Infectious Diseases*
72:e49–e64. doi:10.1093/cid/ciaa1216

Krause PJ, Rogers R, Shah MK, Kang H, Parsonnet J, Kodama R, Vannier E. 2024.
Tafenoquine for relapsing babesiosis: a case series. *Clinical Infectious Diseases*
79:130–137. doi:10.1093/cid/ciae238

Krishna S, Woodrow C, Webb R, Penny J, Takeyasu K, Kimura M, East JM. 2001.
Expression and functional characterization of a *Plasmodium falciparum* Ca^{2+}-ATPase
(*Pf*ATP4) belonging to a subclass unique to apicomplexan organisms. *Journal of*
Biological Chemistry **276**:10782–10787. doi:10.1074/jbc.M010554200

Lee AH, Fidock DA. 2016. Evidence of a mild mutator phenotype in cambodian *Plasmodium*

607  *falciparum* malaria parasites. *PLoS ONE* **11**:e0154166. doi:10.1371/journal.pone.0154166

608 Lehane AM, Dennis ASM, Bray KO, Li D, Rajendran E, McCoy JM, McArthur HM,
609  Winterberg M, Rahimi F, Tonkin CJ, Kirk K, Van Dooren GG. 2019. Characterization of
610  the ATP4 ion pump in *Toxoplasma gondii*. *Journal of Biological Chemistry* **294**:5720–5734.
611  doi:10.1074/jbc.RA118.006706

612 Lemieux JE, Tran AD, Freimark L, Schaffner SF, Goethert H, Andersen KG, Bazner S, Li A,
613  McGrath G, Sloan L, Vannier E, Milner D, Pritt B, Rosenberg E, Telford S, Bailey JA,
614  Sabeti PC. 2016. A global map of genetic diversity in *Babesia microti* reveals strong
615  population structure and identifies variants associated with clinical relapse. *Nat Microbiol*
616  **1**:16079. doi:10.1038/nmicrobiol.2016.79

617 Li H, Galon EM, Ji S, Zafar I, Ma Z, Do T, Amer MM, Ma Y, Liu M, Xuan X. 2023. *In vitro*
618  screening of compounds from the Food and Drug Administration-approved library
619  identifies anti-*Babesia gibsoni* activity of idarubicin hydrochloride and vorinostat.
620  *Parasitology International* **96**:102774. doi:10.1016/j.parint.2023.102774

621 Lindahl, Abraham, Hess, Spoel VD. 2021. GROMACS 2021 Manual.
622  doi:10.5281/ZENODO.4457591

623 Liu M, Adjou Moumouni PF, Cao S, Asada M, Wang G, Gao Y, Guo H, Li J, Vudriko P,
624  Efstratiou A, Ringo AE, Lee S-H, Hakimi H, Masatani T, Sunaga F, Kawazu S, Yamagishi
625  J, Jia L, Inoue N, Xuan X. 2018. Identification and characterization of interchangeable
626  cross-species functional promoters between *Babesia gibsoni* and *Babesia bovis*. *Ticks and*
627  *Tick-borne Diseases* **9**:330–333. doi:10.1016/j.ttbdis.2017.11.008

628 Liu M, Ji S, Kondoh D, Galon EM, Li J, Tomihari M, Yanagawa M, Tagawa M, Adachi M,
629  Asada M, Igarashi I, Iguchi A, Xuan X. 2021. Tafenoquine is a promising drug candidate
630  for the treatment of babesiosis. *Antimicrob Agents Chemother* **65**:e00204-21.
631  doi:10.1128/AAC.00204-21

632 Marcos LA, Leung A, Kirkman L, Wormser GP. 2022. Use of tafenoquine to treat a patient
633  with relapsing babesiosis with clinical and molecular evidence of resistance to
634  azithromycin and atovaquone. *IDCases* **27**:e01460. doi:10.1016/j.idcr.2022.e01460

635 Matsuu A, Koshida Y, Kawahara M, Inoue K, Ikadai H, Hikasa Y, Okano S, Higuchi S. 2004.
636  Efficacy of atovaquone against *Babesia gibsoni in vivo* and *in vitro*. *Veterinary*
637  *Parasitology* **124**:9-18. doi:10.1016/j.vetpar.2004.07.005

638 Mohring F, Van Schalkwyk DA, Henrici RC, Blasco B, Leroy D, Sutherland CJ, Moon RW.
639  2022. Cation ATPase (ATP4) orthologue replacement in the malaria parasite *Plasmodium*
640  *knowlesi* reveals species-specific responses to ATP4-targeting drugs. *mBio* **13**:e01178-22.
641  doi:10.1128/mbio.01178-22

642 Mordue DG, Wormser GP. 2019. Could the drug tafenoquine revolutionize treatment of
643  *Babesia microti* infection? *J Infect Dis* **220**:442-447. doi: 10.1093/infdis/jiz119

644 Ndayisaba G, Yeka A, Asante KP, Grobusch MP, Karita E, Mugerwa H, Asiimwe S, Oduro A,

Fofana B, Doumbia S, Jain JP, Barsainya S, Kullak-Ublick GA, Su G, Schmitt EK, Csermak K, Gandhi P, Hughes D. 2021. Hepatic safety and tolerability of cipargamin (KAE609), in adult patients with *Plasmodium falciparum* malaria: a randomized, phase II, controlled, dose-escalation trial in sub-Saharan Africa. *Malar J* **20**:478. doi:10.1186/s12936-021-04009-1

Qiu D, Pei JV, Rosling JEO, Thathy V, Li D, Xue Y, Tanner JD, Penington JS, Aw YTV, Aw JYH, Xu G, Tripathi AK, Gnadig NF, Yeo T, Fairhurst KJ, Stokes BH, Murithi JM, Kümpornsin K, Hasemer H, Dennis ASM, et al. 2022. A G358S mutation in the *Plasmodium falciparum* Na$^+$ pump *Pf*ATP4 confers clinically-relevant resistance to cipargamin. *Nat Commun* **13**:5746. doi:10.1038/s41467-022-33403-9

Qurollo BA, Archer NR, Schreeg ME, Marr HS, Birkenheuer AJ, Haney KN, Thomas BS, Breitschwerdt EB. 2017. Improved molecular detection of *Babesia* infections in animals using a novel quantitative real-time PCR diagnostic assay targeting mitochondrial DNA. *Parasites Vectors* **10**:128. doi:10.1186/s13071-017-2064-1

Rogers R, Krause PJ, Norris AM, Ting MH, Nagami EH, Cilley B, Vannier E. 2023. Broad antimicrobial resistance in a case of relapsing babesiosis successfully treated with tafenoquine. *Clinical Infectious Diseases* **76**:741–744. doi:10.1093/cid/ciac473

Rosenblatt J, Leung A, Baneman E, Fuller R, Taimur S, Paniz-Mondolfi AE, Malone AK, Kirkman L, Jacobs SE. 2021. Relapsed *Babesia microti* infection following allogeneic hematopoietic cell transplantation in a patient with B-cell acute lymphoblastic leukemia: case report and review of the literature. *Open Forum Infectious Diseases* **8**:ofab323. doi:10.1093/ofid/ofab323

Rosling JEO, Ridgway MC, Summers RL, Kirk K, Lehane AM. 2018. Biochemical characterization and chemical inhibition of *Pf*ATP4-associated Na$^+$-ATPase activity in *Plasmodium falciparum* membranes. *Journal of Biological Chemistry* **293**:13327–13337. doi:10.1074/jbc.RA118.003640

Rottmann M, McNamara C, Yeung BKS, Lee MCS, Zou B, Russell B, Seitz P, Plouffe DM, Dharia NV, Tan J, Cohen SB, Spencer KR, González-Páez GE, Lakshminarayana SB, Goh A, Suwanarusk R, Jegla T, Schmitt EK, Beck H-P, Brun R, et al. 2010. Spiroindolones, a potent compound class for the treatment of malaria. *Science* **329**:1175–1180. doi:10.1126/science.1193225

Saliba KJ, Kirk K. 1999. pH regulation in the intracellular malaria parasite, *Plasmodium falciparum*. *Journal of Biological Chemistry* **274**:33213–33219. doi:10.1074/jbc.274.47.33213

Schmitt EK, Ndayisaba G, Yeka A, Asante KP, Grobusch MP, Karita E, Mugerwa H, Asiimwe S, Oduro A, Fofana B, Doumbia S, Su G, Csermak Renner K, Venishetty VK, Sayyed S, Straimer J, Demin I, Barsainya S, Boulton C, Gandhi P. 2022. Efficacy of cipargamin (KAE609) in a randomized, phase II dose-escalation study in adults in Sub-Saharan Africa

with uncomplicated *Plasmodium falciparum* malaria. *Clinical Infectious Diseases* **74**:1831–1839. doi:10.1093/cid/ciab716

Simon MS, Westblade LF, Dziedziech A, Visone JE, Furman RR, Jenkins SG, Schuetz AN, Kirkman LA. 2017. Clinical and molecular evidence of atovaquone and azithromycin resistance in relapsed *Babesia microti* infection associated with rituximab and chronic lymphocytic leukemia. *Clinical Infectious Diseases* **65**:1222–1225. doi:10.1093/cid/cix477

Spillman NJ, Allen RJW, McNamara CW, Yeung BKS, Winzeler EA, Diagana TT, Kirk K. 2013. Na$^+$ regulation in the malaria parasite *Plasmodium falciparum* involves the cation ATPase *Pf*ATP4 and is a target of the spiroindolone antimalarials. *Cell Host & Microbe* **13**:227–237. doi:10.1016/j.chom.2012.12.006

Tayebwa DS, Tuvshintulga B, Guswanto A, Nugraha AB, Batiha GE-S, Gantuya S, Rizk MA, Vudriko P, Sivakumar T, Yokoyama N, Igarashi I. 2018. The effects of nitidine chloride and camptothecin on the growth of *Babesia* and *Theileria* parasites. *Ticks and Tick-borne Diseases* **9**:1192–1201. doi:10.1016/j.ttbdis.2018.04.019

Trott O, Olson AJ. 2010. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* **31**:455–461. doi:10.1002/jcc.21334

Tuvshintulga B, Sivakumar T, Nugraha AB, Ahedor B, Batmagnai E, Otgonsuren D, Liu M, Xuan X, Igarashi I, Yokoyama N. 2022. Combination of clofazimine and atovaquone as a potent therapeutic regimen for the radical cure of *Babesia microti* infection in immunocompromised hosts. *The Journal of Infectious Diseases* **225**:238–242. doi:10.1093/infdis/jiab537

Vannier E, Krause PJ. 2012. Human babesiosis. *N Engl J Med* **366**:2397–2407. doi:10.1056/NEJMra1202018

Vega CA, Buening GM, Green TJ, Carson CA. 1985. *In vitro* cultivation of *Babesia bigemina*. *Am J Vet Res* **46**:416–420.

Vydyam P, Pal AC, Renard I, Chand M, Kumari V, Gennaro JC, Mamoun CB. 2024. Tafenoquine-atovaquone combination achieves radical cure and confers sterile immunity in experimental models of human babesiosis. *The Journal of Infectious Diseases* **229**:161–172. doi:10.1093/infdis/jiad315

Zhang R, Suwanarusk R, Malleret B, Cooke BM, Nosten F, Lau Y-L, Dao M, Lim CT, Renia L, Tan KSW, Russell B. 2016. A basis for rapid clearance of circulating ring-stage malaria parasites by the spiroindolone KAE609. *J Infect Dis* **213**:100–104. doi:10.1093/infdis/jiv358

Zhou Y, Fomovska A, Muench S, Lai B-S, Mui E, McLeod R. 2014. Spiroindolone that inhibits *Pf*ATPase4 is a potent, cidal inhibitor of *Toxoplasma gondii* tachyzoites *in vitro* and *in vivo*. *Antimicrob Agents Chemother* **58**:1789–1792. doi:10.1128/AAC.02225-13

Figure Legends

Figure 1. CIP demonstrates potent inhibition on *Babesia* spp. (A) and **(B)** Dose-dependent

growth curve of CIP on *B. bovis* and *B. gibsoni in vitro*. Each value represents the mean ±

standard deviation (SD) of three independent experiments carried out in triplicate. IC_{50}: the

half maximal inhibitory concentration. **(C)** Inhibitory effects of CIP and atovaquone (ATO)

plus azithromycin (AZI) on the proliferation of *B. microti* in BALB/c mice. **(D)** Hematocrit

(HCT) values in mice treated with CIP or ATO plus AZI compared with vehicle-treated mice.

(E) Inhibitory effects of CIP and ATO plus AZI on the proliferation of *B. rodhaini* in BALB/c

mice. **(F)** Survival rates of CIP-treated, ATO plus AZI-treated, and vehicle-treated mice. The

treatment time is shown by two-way arrows, and significant differences ($P < 0.01$) between

the drug-treated groups and the vehicle-treated control group are indicated by asterisks. The

data from one of six individual experiments are expressed as means ± SD. [**], $P < 0.01$.

Figure 2. Mutations in BgATP4 mediate CIP resistance. (A-C) Representative sequencing

chromatogram of wild-type and resistant parasites from CIP-treated *B. gibsoni*. The resistant

parasite genomic DNA is extracted from blood samples after a 60 day-treatment. The

BgATP4 gene was amplified and sequenced using the DNA. **(D)** Genes of high-frequency

sequence variants detected by NGS. **(E)** Dose-dependent growth curve of BgATP4[WT],

BgATP4[L921V], and BgATP4[L921I] *in vitro*. Each value represents the mean ± standard deviation

(SD) of three independent experiments carried out in triplicate.

Figure 3. Mechanistic basis for resistance to CIP conferred by the L921V and L921I

mutations in BgATP4. (A) Untreated and CIP-treated parasite morphology after incubation

for 72 h. Scale bar: 5 μm. **(B)** Sizes of 100 parasites in two groups measured with ImageJ

software in panels A. Statistically significant differences between the means of variables determined by t-test. [****], $P < 0.0001$. **(C)** TEM of untreated and CIP-treated parasite. N, Nucleus; SB, spherical body; V, vacuole. Scale bar: 500 nm. **(D)** $[Na^+]_i$ concentrations after the addition of CIP in the BgATP4^{WT} line. Representative traces from the experiment that highlight the impact of adding 20 nM CIP (blue), 1 nM CIP (green), or 0 nM CIP (grey) on the concentration $[Na^+]_i$ of the BgATP4^{WT} line. **(E)** Alkalinization of pH$_i$ in BgATP4^{WT} line upon addition of the ATP4 inhibitor. **(F)** Addition of 20 nM CIP to the wild-type and resistant parasite lines results in different $[Na^+]_i$ concentrations. **(G)** Addition of 20 nM CIP to the wild-type and resistant parasite lines results in different pH$_i$ concentrations. Experiments were performed in technical duplicates for at least three biological repeats. **(H)** Data acquired in the low Na$^+$ condition (containing only the 2 mM Na$^+$ introduced upon the addition of 1 mM Na$_2$ATP) was subtracted from data obtained in the high Na$^+$ condition to determine the ATPase activity related to the BgATP4 proteins. The results are presented as the average of data from three independent tests. **(I)** Dose-dependent BgATP4-associated ATPase activity curve of BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I} *in vitro*. ATPase activity was determined at pH 7.2 in the presence of 150 mM Na$^+$ and 1 mM Na$_2$ATP. Each value represents the mean ± standard deviation (SD) of three independent experiments carried out in triplicate. [*], $P < 0.05$; [**], $P < 0.01$.

Figure 4. Binding sites proximal to BgATP4 residue 921 predicted by molecular docking. **(A)** The lowest energy poses for CIP were located in reference to the whole protein structure, docking against the WT (green), L921V (yellow), and L921I (pink) mutant BgATP4. The side chain of L921 is also shown in a red stick at its position. **(B-D)** The zoomed views of the binding locations of CIP.

Figure 5. Cross-resistance between ATO and TQ in resistant parasites and combination therapy based on CIP plus TQ. (A) Dose-dependent growth curve of BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I} by ATO treatment *in vitro*. **(B)** Dose-dependent growth curve of BgATP4^{WT}, BgATP4^{L921V}, and BgATP4^{L921I} by TQ treatment *in vitro*. Each value represents the mean ± standard deviation (SD) of three independent experiments carried out in triplicate. **(C)** Inhibitory effects of CIP plus TQ on the proliferation of *B. microti* in SCID mice. Treatment started at 4 DPI: CIP was given at 20 mg/kg once daily for 7 days, TQ was administered as a single 20 mg/kg dose, and the combination group received both treatments (CIP at 20 mg/kg once daily for 7 days plus a single dose of TQ at 20 mg/kg). The data are presented as the means from one of five independent experiments. **(D)** Parasite DNA was detected by qPCR on genomic DNA extracted from blood collected from untreated and treated SCID mice infected with *B. microti* at 90 DPI. A dotted grey line across the graph represents the average cut-off Cq value. Cut-off Cq ≤ 35 was considered positive, while Cq > 35 or no amplification was considered negative. Each sample was analyzed in duplicate, and two independent experiments were conducted. NA indicates no amplification.

Supplementary Materials

Figure supplement 1. Cytotoxicity assay of CIP on the Madin-Darby canine kidney (MDCK) cells and human foreskin fibroblasts (HFF). The MTP-500 microplate reader was utilized to detect the absorbance at 450 nm. The results are displayed as the mean ± standard deviation of three separate tests. CC_{50}: the 50% cytotoxic concentration.

Figure supplement 2. Multiple sequence alignment of ATP4 in different species. A yellow square and arrow denotes the BgATP4 mutation site discovered in this investigation; purple squares and arrows represent sites linked to *P. falciparum* CIP resistance, and a gray square and arrow represent sites associated with *T. gondii*.

Figure supplement 3. Binding sites for CIP found by Gnina search across the entire surface of the protein. (**A-C**) The binding space of WT, L921V, and L921I mutants in BgATP4 are labeled in green, yellow, and pink, respectively.

Figure supplement 4. Proposed mechanism of inhibition of CIP on wild-type and mutant parasite-infected erythrocytes. CIP disrupts the BgATP4 function of wild-type parasites, which causes a net influx of Na^+ and efflux of H^+ from the parasite. The osmotic load imposed on the influx of Na^+ further brings about parasite swelling and internal alkalinization, which are the main factors in *Babesia* death. Mutations in ATP4 minimize the susceptibility to ATP4 inhibitors by recovering H^+ and Na^+ balance, as indicated by the dotted arrows.

 Supplemental Tables

807 **Table supplement 1.** Primer sets of *B. gibsoni* ATP4

Gene	Primer (5'-3')
Bgatp4-1F	ACGTAGTTGTCGAACATCTTACACCAATGC
Bgatp4-1R	CAGAGGTAGCATAGTGAGATCGCCCCGA
Bgatp4-2F	GAGATTGCAAGGTTTACGGATGAGTCTATGG
Bgatp4-2R	ATGACAACGTCACCGGGTACAACACGCTTG
Bgatp4-3F	GAAAAGCTTGCTCAAATGTCTTCACCGACAAC
Bgatp4-3R	GTTTCTGGACGAGTTGGATCCTCATAACC
Bgatp4-4F	AAATTGGGGGGGATAATTGGCATTATCTCC
Bgatp4-4R	ACCTTATCAAAGGGTACCTCTTGCTTTG
Bgatp4-5F	TCAATCCATTCGGTGGAATTTTCAAAAAGG
Bgatp4-5R	GTGTTTGAATAACATATCTGGTGCCCCCTT
Bgatp4-6F	GGTAAAGAACAGATTTGCTGACATAGATCT
Bgatp4-6R	ACCCTATTTCTTGTGCGATGGCCGTAGCTG
Bgatp4-7F	AGGGACGCTATAAATACTTGTGGAAAGGC
Bgatp4-7R	GTACGCCCAGACTCGATTGAGTTAACTATTG
Bgatp4-8F	GGGTATTAATGGCACAGATGTTGCAAAAGG
Bgatp4-8R	GACAGTTCCCACTAGCATCATTAAGTGTAA
Bgatp4-9F	ATCATCATTGCGCTATATGTTTCTACCGG
Bgatp4-9R	CTAATGGGGTAGCATTTCCATTCTCAACAA
Bgatp4-10F	AGGGAGTTCCTCCCATGTATAAAGATACCT
Bgatp4-10R	TTAGGCTTCTGCCCTTTTCATTTTTCGCTT

809   **Table supplement 2.** Primer sets of *B. microti* ATP4

Gene	Primer (5'-3')
Bmatp4-1F	ATGAGTAGATCTCAGTCTAACATTTC
Bmatp4-1R	TGCTATTAGTAGTGCTATAACAAAG
Bmatp4-2F	CAAGTATGGCGAAAATGCTATAAAC
Bmatp4-2R	CATTCCGTTTGTTACGCTGGTGC
Bmatp4-3F	CTTACAGGTGAGAGTGAAGATGTAAAG
Bmatp4-3R	ACAAATTGCTGAGCAGCAACCCAGAG
Bmatp4-4F	CTATTTCATTGTCATTGGGAGCTAA
Bmatp4-4R	CAAATCCGGCCTTAGCTGCGCCAA
Bmatp4-5F	TCATACGGTTCAACGTTATCACATG
Bmatp4-5R	TCTATCCGACGACTCCAACCTTTC
Bmatp4-6F	GTTGTCCAAGAATGCTTATAGGGTG
Bmatp4-6R	CATTGCCACGCCAATGTCCGCCTGTTT
Bmatp4-7F	GTTAACTCGTTGAAGAGACAAGGTTAC
Bmatp4-7R	CACAATTGTACCGACAACTGCCGCTG
Bmatp4-8F	GGCCAAAATCTCAGCCTTTAATGAC
Bmatp4-8R	CAGCTCCAATGGGTTGGTAGTAACCATT
Bmatp4-9F	CGTTATTGACGGCGTTGAGAATGATG
Bmatp4-9R	CCCAAAACACCTTTGCCATTTAACTC
Bmatp4-10F	GAAAATTGTAGAGCATTTCAAAACCAG
Bmatp4-10R	TTATTGCAGGGCAGCCTGCTTAGA

 Table supplement 3. Interactions of CIP from docking simulations

Receptor and binding energy	Name	Distance	Category	Types
BgATP4^{WT} -6.43 kcal/mol	GLY407	2.67648	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	CYS959	3.2396	Halogen	Halogen (Cl, Br, I)
	VAL962	2.66882	Hydrophobic	Pi-Sigma
	CYS959	4.38742	Hydrophobic	Alkyl
	VAL962	4.29249	Hydrophobic	Alkyl
	ALA963	3.8231	Hydrophobic	Alkyl
	ALA1202	3.67015	Hydrophobic	Alkyl
	MET211	5.06565	Hydrophobic	Alkyl
	VAL962	4.25668	Hydrophobic	Alkyl
	ILE414	4.70605	Hydrophobic	Alkyl
	CYS959	4.83891	Hydrophobic	Pi-Alkyl
	VAL962	4.98402	Hydrophobic	Pi-Alkyl
	VAL411	5.43239	Hydrophobic	Pi-Alkyl
	VAL962	5.05537	Hydrophobic	Pi-Alkyl
	PHE954	4.07586	Hydrophobic	Pi-Alkyl
BgATP4^{L921V} -6.40 kcal/mol	GLY407	2.69013	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	CYS959	3.22066	Halogen	Halogen (Cl, Br, I)
	VAL962	2.66391	Hydrophobic	Pi-Sigma
	CYS959	4.35422	Hydrophobic	Alkyl
	VAL962	4.31879	Hydrophobic	Alkyl
	ALA963	3.8209	Hydrophobic	Alkyl
	ALA1202	3.63038	Hydrophobic	Alkyl
	MET211	5.0878	Hydrophobic	Alkyl
	VAL962	4.24186	Hydrophobic	Alkyl
	ILE414	4.71819	Hydrophobic	Alkyl
	CYS959	4.82036	Hydrophobic	Pi-Alkyl
	VAL962	4.99879	Hydrophobic	Pi-Alkyl
	VAL411	5.4334	Hydrophobic	Pi-Alkyl
	VAL962	5.04532	Hydrophobic	Pi-Alkyl
	PHE954	4.07482	Hydrophobic	Pi-Alkyl
	GLY958	2.92346	Hydrogen Bond; Halogen	Carbon Hydrogen Bond; Halogen (Fluorine)
	GLY958	3.3301	Halogen	Halogen (Fluorine)
	MET410	5.88505	Other	Pi-Sulfur

	VAL955	4.31406	Hydrophobic	Alkyl
	MET410	3.69023	Hydrophobic	Alkyl
BgATP4^{L921I}	VAL411	4.83101	Hydrophobic	Alkyl
-6.26 kcal/mol	VAL955	4.79513	Hydrophobic	Alkyl
	VAL955	4.84729	Hydrophobic	Alkyl
	CYS959	4.1269	Hydrophobic	Alkyl
	ILE1205	4.82529	Hydrophobic	Alkyl
	CYS959	5.21399	Hydrophobic	Alkyl
	VAL955	4.90527	Hydrophobic	Pi-Alkyl
	PHE954	5.03012	Hydrophobic	Pi-Alkyl

A $CC_{50}: 38.7 \pm 2.0\ \mu M$

B $CC_{50}: 70.8 \pm 4.9\ \mu M$

```
Homo_sapiens/1-1035   233  SEPQTRSPECTHESPLETRNIAFFSTMCLEGTVQGLVVNTGDRTIIGRIASL--ASGVENEKTFIAIEIEHFVDIIAGLAILFGATFFIVAMCIGYT---  327
P._falciparum/1-1264  286  SEDIKKTIVADNLSTPFATNLCFATTSVTSGSGKGIVISTGLDTQVGKIASQLKKSSKGSKLTPLQVALNKLGGLIGLIAIIVLVVIISLAVIIKYRDPA  385
T._gondii/1-1338      347  SEDISKTLRAKDYDTPFATNLCFASTIVTNGSGRGLVFATGMETQVGRIAQQLKKAGEGSRLTPLQRGLNRLGSMIGLAICVLIIVVVVAILTGYRDPA  446
B._gibsoni/1-1264     283  SDAVRKQLVAADLDSPFSTNLCFASTSITSAGSKAVVVKTGMETQVGCIAKQLQAATHRSTMTPLQLGLNKLGGIIGISIGILITIVLVAVFSGYEDPT  382
B._bovis/1-1261       283  SEPVKKHLVADDLNAPFATNMCFASTSVVCGSAKGIVVTTGMQTQVGHIATQLKSALSGSDVTPLQSSLNSLGGIIGISIFVLITIIVVAIVTKYEDPT  382
B._microti/1-1234     273  SEDVKKTLLSPDPDSPFSSNLVFSSTSVTNGMGKAIVTHVGMDTQVGKIAHQLKKASQKSKMTPLQRGLNKLGGLIGIISVIILVVVIVLVALIFNYSDPT  372

Homo_sapiens/1-1035   328  ------FLRAMVFFMAIVVAYVPEGLLATVTVCLSLTAKRLASKNCVVKNLEAVETLGSTSVICSDKTGTLTQNRMTVSHLW-------------------  403
P._falciparum/1-1264  386  HADKDPTFVIIIIGVGFAVSSIPEGLPMVVTITLSAGAKDMVKKNANVRKLPAVETLGCCSVICSDKTGTLTEGKMTIANAVTICKNSSLSDENNKLTKT  485
T._gondii/1-1338      447  HPDADPVFTIVLVAVGFAVSSIPEGLPMVVTICLSLGARDMVKRKANVRKLPAVETLGCCSVICSDKTGTLTEGKMTAVRLVTVCRNGKVVDADG-LTKS  545
B._gibsoni/1-1264     383  RPETNRILSIVLLAVGFAVSSIPEGLPMVVTISLSLGAKDMARRNANIRKLPAVETLGCCSIICSDKTGTLTEGKMTATTIITFNSVG-----GDCTSEE  477
B._bovis/1-1261       383  RPDTNRVLAIVLLAVGFAVSAIPEGLPMVVTISLSLGAKDMRDNANIRKLPAVETLGCCSIICSDKTGTLTEGKMTSISAMIFHNED-----DGWKASE  477
B._microti/1-1234     373  KPNDNRILTILLLAVGFAVSSIPEGLPMIVTISLSLGANDMAACNANIRKLPAVETLGCCSAICSDKTGTLTQGKMTVVSLITFVTDS--LSSKSPVARH  470

Homo_sapiens/1-1035   404  ------------------FDNHIHTADTTED-----QSG--QTFDQSSETWRA--------LCRVLTLCNRAAFKSGQDAV-----PVPKRIVIGDASETAL  467
P._falciparum/1-1264  486  FDFYPTKGFEFCGGLFDSNELTSEKKKEIVIAKNQN---TSYDKVLYNYGNPSNKSV---IVDKTRSLMFAAYLNSYDTTLSRDPKTLKWGIHGNMSEGPI  580
T._gondii/1-1338      546  FGFYPTKGFDPNGGIFDYNALDEKTKSNLMLQYRDGAFQDFDAVCYNYGNPANKDP---TTKLVRSVMLSGYLNSHATTLSRDPDTNRWLAKGNMSEGAI  642
B._gibsoni/1-1264     478  LLLFPTLGFNPFGGIFKKENLTEGLKADLLEMAKQE--VPFDKVKENLCSAKNKSA---VSRQVKACLLSAYLNSRGTKIERNENHTDWKGVGNMTECPL  572
B._bovis/1-1261       478  LQFYPTMGFNPYGGVFKPEHLTDDFKAELQKQTHAA--ASFDNVTNNVCAKGNTSS---DSKQIRACMLSAYLNSNSTRVEQDEASKAWVAMGNMTECPI  572
B._microti/1-1234     471  FKFFPTLGFDPRGGIFPEQYLTPATTKELLEAAEGG--DNLRAHEQNLGSPSYAGPDIIDSKRVRLAMLASYLNSYGSTLSHEDG--KYKVVGNMSEGAL  566

Homo_sapiens/1-1035   468  L-------KFSELTLGNAMG-----YRDRFPKVC--EIPFNSTNKFQLSIHTLE-------------DPRDPRHLLVMKGAPERVLERC-SSILIKGQ----  537
P._falciparum/1-1264  581  VVAAAKVGYSFINNPNHKS----YLDNFQRLDDLEVTFNSSRKMKITFYKLKTVNVFENVYLDK-PGKVYTHVALIKGAPDRLLDR--STHLLEETSMKK  673
T._gondii/1-1338      643  VVGAAKARFGETVAGQEMCGMHDAKADFPRVQELEVPFNSSRKMMMTVHQLPAVNYFGDICLNNTTGTKYTHCAIVKGAPDRVLQH--VRYTVRE-GISG  739
B._gibsoni/1-1264     573  IVAAAKCGIGKTVNPDDKT----CSD-YPSIDALEVPFNSSRKMMCTVHKLKVKNRFADIDLTS-DGTTYTHVATIKGAPDMLFKH--ISATIKRDN-DS  663
B._bovis/1-1261       573  IVASAKCGIGSTINKSDAT----YDD-HPMLKDLEVPFNSGRKMMVTIHKLKTENMFGDLNTIN-GFKTYTHVATIKGAPDMLFKY--RIVTVKRNG-EE  663
B._microti/1-1234     567  VVGAAKAGFGALEDG--IY----AHDIYPRLSRLEIPFSSSRKMMITVHAIK--GSFGDL-----GLSGFSGIALIKGAPDRLFSYCNQVISVDGKG-KL  652

Homo_sapiens/1-1035   538  -------ELPLDEQWREAFQTAYLSLGGLGERVLGFCQLYLNEKDYPP--------GYAFDVEAMNFPSSGLCFAGLVSMIDPPRATVPDAVLKCRTA  620
P._falciparum/1-1264  674  VQVSW----NSTITQEERNVLIKKNLELSQKALRVLSICIKPLTDQNIEELKKLEDADERLKYVN---YDENGGFIPMGYVASFDPPRPGVKEAIQTCREA  767
T._gondii/1-1338      740  PSVEW----EKQMTPEEIMKVEAVNLELSEQALRVLALTFRPLTDADVAALRRQAGADERLKFAL---GETREELVLLGVIGSVDPPRVGVREAIDRCGEA  833
B._gibsoni/1-1264     664  CTIDWNM-DINESSKIMEKFLKANDELSSRALRVLMVGMFPLTDTDVESLKRCDDSDQRLTWLLTGKGMGGCQLVLLGFIGSLDPPRFGVRDAINTCGKA  762
B._bovis/1-1261       664  CVIDWASGDIENNETLLNVMKKANNDLSSRALRVLLVGMFPLTAEDITALEGCEDSDSRLNWLLNGHEHGSSPLVLLGFIGNLDPPRFGVKEAVATCGKA  763
B._microti/1-1234     653  C------HMGVDDNLAKIMSDANSLLSKNAYRVLLLAICPLIAEDLENLERLESSD-RQEYIL------GKKLIVLGLTGSVDPPRSGVKEAIVSFSQA  738

Homo_sapiens/1-1035   621  GIRVIMVTGDHPITAKAIAASVGIISEGSETVEDIAARLRVPVDQVNRKDARACVINGMQLKDMDPSELVEALRTHPEMVFARTSPQQKLVIVESCQRLG  720
P._falciparum/1-1264  768  QVKVIMITGDQKPTAVAISKLIGLIEEKSEQVEDINSLAI----------ECSELHINKNPNEPILPNDQLDEF-IDKILIYSRAQPEDKITIVQSLKRKG  857
T._gondii/1-1338      834  GIRVIMITGDQRPTAVAIAKDIGLLTSQ----DDPEQQSI----------QCSGLHVDDDPMNEHLPEEELDEI-IARVKVFSRAQPEDKIAIVEALKRQG  919
B._gibsoni/1-1264     763  GVRVIMITGDQKATATAIAQEIGLFNQF-DYLADAKSGAI----------ECNSLHVNGEHFREYLPDDIIDGY-TSKASVFCRAQPEDKVVVESLKRQG  851
B._bovis/1-1261       764  GVRVVMITGDQKPTATAIGKDIGLFNQF-DYLGDENSGVI----------ECSQMHINNDHFREYLPDGVIDSF-TSRVSVFCRAQPEDKVVIVKSLKRQG  852
B._microti/1-1234     739  GVRVLMITGDQKDTATAIAKQIGIIAE-----TEDSLSR----------ECRELHIGGDHSMPYLEDAEIDEI-TSKSNVFSRAQPEDKVVINSLKRQG  822

Homo_sapiens/1-1035   721  AIVAVTGDGVNDSPALKKADIGVAMGIAGSDAAKNAADMILLDDNFASIVTGVEQGRLIFDNLKKSIAYTLTKNIPELTPYLIYITVSVPLPLGCITILF  820
P._falciparum/1-1264  858  YLVAMTGDGVNDAPALKAADIGVAMGINGTEVAKGASEMILLDDNFCTVSAIDVGRTIFSNIQKFVCFLLGTNIGEIIYLSVAIVAQMPFPLEALQILF  957
T._gondii/1-1338      920  HTVAMTGDGVNDAPALKAADIGVAMGIAGTDVAKGASEMVLLDDNFVTIVAAVEEGRKIYSNIQKFVCFLLGTNIGEIIYLTIAIAASMPLPLEALQVLF  1019
B._gibsoni/1-1264     852  YLTAMTGDGVNDAPALKAANIGIAMGINGTDVAKGAADMVLLDDNFCTIVNSIESGRTIYANIQKFVCFLGTNIGEILYLTTSIMMRTLPPVEALQILY  951
B._bovis/1-1261       853  YLTAMTGDGVNDAPALKTADIGVAMGINGTDVAKGAADMVLLDDNFCTIVNSIESGRTIYANIQKFVCFLGTNIGEIFYLTTALLMKTLPPVEALQILF  952
B._microti/1-1234     823  YLTAMTGDGVNDAPALKAADIGVAMGINGTDVAKGAADMVLLDDNFCTIVSAIKCGRIIYSNIQKFVSFLLGTNIGEILYLTIAVIISTLPPVEALQILF  922

Homo_sapiens/1-1035   821  IELCTDIFPSVSLAYEKAESDIMHLRPRNPKRDRLVNE--------PLAAY----------------SYFQIGAIQS-------------------FA  875
P._falciparum/1-1264  958  LNLMTDGCPAVALSREPPNDDNMKTPPRPKKQPIMTKRWWFYGILPHTIFEALCVLLSLAFSLYICTGFYNLNGIHNLCKTVNLVDVNDANVYHEYKYFC  1057
T._gondii/1-1338      1020 LNLMSDGCPAVALAKEPSDDENMKIPPRPRKQPIMTRDWWLYGNLPHTIFEAGCVLMSLALGLYLCTGVVQLNPLHEQCSYFTATQLSH-NKDIDYRYFC  1118
B._gibsoni/1-1264     952  LNFVTDGCPAVALSREPPEEYTMTTKPRRPKQPIMTLDWWIYGNLPHAIFEAMMVICSIIIALYVSTGVIFMQDLRNQGSTVTLNDAS--GNCHDFIYFC  1049
B._bovis/1-1261       953  LNFVTDGCPAVALSREPPEDDVLTTKPRDPGQPIMTRNWWLYGNIPHTIFQGIFVIGSILTALYISTGVIFVKDIHEQCKYVELANGA--GTKYRYTYFC  1052
B._microti/1-1234     923  LNLMTDGCPAVALSREPADENIMKIPPRPKSQPLMTKDWWIYGNLPHTIFEAAAVVGTIVLGLYVCTGVYTIGDLHSQCKYVTFKHD---NVETKYMYYC  1019
```