Unaware Processing of Words Activates Experience-Derived Information in Conceptual-Semantic Brain Networks

Marta Ghio[1,2,*], Barbara Cassone[3,*], Marco Tettamanti[2,3]

[1] Faculty of Mathematics and Natural Sciences, Heinrich Heine University Düsseldorf, Düsseldorf, Germany.

[2] CIMeC - Center for Mind/Brain Sciences, University of Trento, Italy.

[3] Department of Psychology, University of Milan-Bicocca, Milano, Italy.

[*] **Equal contribution**

Corresponding author:

Marco Tettamanti

Department of Psychology, University of Milan-Bicocca, Milano, Italy

Piazza dell'Ateneo Nuovo 1, I-20126 Milano, Italy

Email: marcodante.tettamanti@unimib.it

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgments: M.G. was supported by a postdoctoral researcher grant by the University of Trento, Italy (Decree No. 48, 23.05.2019) awarded to M.T.

Author contributions: M.G., B.C., and M.T. designed the experiment; M.G. and B.C. collected the data; M.G, B.C., M.T. analyzed the data; M.G., B.C., M.T. organized and wrote the manuscript.

Abstract

The recognition of manipulable objects results from the encoding of sensory input in

combination with predictive decoding of experience-derived visuomotor information stored in

conceptual-semantic representations. This grounded interpretive processing was previously found to

subsist even under unaware perception of manipulable object pictures. In this fMRI study, we first

aimed to extend this finding by testing whether experientially grounded visuomotor representations

are unawarely recruited when manipulable objects are not visually depicted, but only referred to by

words presented subliminally through continuous flash suppression. Second, we assessed the

generalizability of decoding experience-derived conceptual information to other semantic

categories, by extending our investigation to subliminally presented emotion words and testing for

unaware recruitment of grounded emotion representations in the limbic system. Univariate analysis

of data sampled from 21 human participants (14 females) showed that manipulable object words

selectively activated a left-lateralized visuomotor network, both when words were presented below

perceptual threshold and when participants subjectively reported lack of stimulus awareness.

Emotion words selectively engaged the bilateral limbic network, although univariate analysis did

not provide evidence for its recruitment under subliminal perceptual conditions. In turn,

multivariate pattern analysis showed that neural codes associated with both manipulable object and

emotion words could be decoded even in the absence of perceptual awareness. These findings

suggest that the brain automatically engages in conceptual-semantic decoding of experience-derived

information not only when circumstances require to interact with manipulable objects and emotions,

but also when these referents are dislocated in time and space and only referred to by words.

Keywords

Introduction

Our brain constantly receives sensory input from around and within us and reacts to it, also

in anticipatory mode, by activating conceptual-semantic representations derived from previous

experience with similar percepts. For example, visual perception of manipulable objects (MO)

engages not only bottom-up visual encoding but also top-down conceptual-semantic decoding based

on stored MO concepts (Noppeney et al., 2006). These are considered to encompass high-level

semantic (e.g., categorical) as well as visual, motor, and functional information derived from our

previous experience with MO exemplars (Binder & Desai, 2011; Lambon Ralph et al., 2017). The

representation, storage and retrieval of visual properties rely on bilateral ventral occipito-temporal

areas, while motor and functional properties draw on portions of the left-hemispheric parietal and

premotor cortices that support the motor programming of actions related to the object function

(Ishibashi et al., 2016; Chen et al., 2017). This MO-specific visuomotor network (MOVN) interacts

with the general semantic brain network (Binder et al., 2009) to form inferences about the incoming

visual input that allow us to ultimately identify the object and possibly to plan a motor response

(Binder & Desai, 2011; Lambon Ralph et al., 2017).

Notably, MO conceptual-semantic representations seem to be activated also under unaware

visual perception conditions (Ludwig et al., 2016; Fahrenfort et al., 2017) such as those instantiated

by Continuous Flash Suppression (CFS, Tsuchiya & Koch, 2005). By applying CFS in combination

with fMRI, we showed the recruitment and effective connectivity of the MOVN both under aware

and unaware processing of MO pictures (Tettamanti et al., 2017; Ghio et al., 2022). A possible

concern, however, is that the MOVN was recruited due to the visual processing of irrelevant

perceptual properties such as the prototypical elongated shape of MO, rather than to the retrieval of

semantic features such as manipulability and affordance. The first aim of the current study was to

address this concern by evaluating the subliminally-induced activation of the MOVN in processing

MO stimuli that lack a visual depiction of the objects, as is the case for words. While both pictorial

and verbal stimuli trigger conceptual-semantic processing, the latter are arbitrarily associated with

their referents through learning and do not carry any object-related visuomotor information. A

recruitment of the MOVN under subliminal processing of MO words would thus speak for the

hypothesis that conceptual-semantic processing activating experience-derived MO information

occurs automatically.

The second aim was to examine the generalizability of the hypothesis that experience-

derived information constituting conceptual-semantic representations is automatically recruited in

subliminal perception, by extending our investigation to emotions (EM). Conceptual representations

of discrete EM (e.g., *happiness, anger*) are characterized by valence, arousal, interoceptive and

somatic information derived from our previous experience of EM instances (Ghio & Tettamanti,

2016; Borghi et al., 2018; Conca et al., 2021). This information is supposed to be represented and

stored in the same limbic network that supports the experience of EM and comprises, bilaterally, the

amygdala, insula, anterior cingulate, and orbitofrontal cortices, putamen, and midbrain (Lindquist et

al., 2012). This EM-specific limbic network (EMLN) has been shown to be activated when we

visually perceive an instance of an EM in, e.g., a face picture, even presented below the perceptual

threshold (Jiang & He, 2006; Lindquist et al., 2012; Dahlén et al., 2022). As for MO, however, to

overcome the potential criticism that subliminal activation is due to visual properties of the pictures,

we used subliminally presented EM words. Previous behavioural and ERP studies using CFS

proved that EM word meaning is also accessed under subliminal conditions (Lei et al., 2017; Sklar

et al., 2012; Vinson et al., 2011; Yang & Yeh, 2011), without providing any insight, however, into

the automatic recruitment of the EMLN.

In the present fMRI study, we tested the aforementioned hypothesis and its generalization

with CFS of MO and EM words. Univariate and multivariate pattern analyses showed automatic

activation of MOVN and EMLN during, respectively, MO and EM word processing.

Materials and Methods

Experimental design and hypotheses

This fMRI study aimed at assessing the hypothesis that experience-derived information

constituting MO (e.g., *hammer*) and EM (e.g., *envy*, *joy*) conceptual-semantic representations is

automatically recruited in subliminal perception. To this purpose we tested whether the subliminal

processing of MO and EM words specifically engages, respectively, the MOVN and the EMLN (see

Fig. 1A and the "Data analysis" section below for the definition of the specific regions of interest

and coordinates). Because valence is a dominant feature for EM conceptual-semantic

representations (Conca et al., 2021), with respect to which EM words are consistently characterized

across languages (Jackson et al., 2019), EM words were parametrically distributed along the

negative-to-positive valence dimension, including an equal number of negative (EMneg) and

positive (EMpos) words. In addition, neutral words NEU (e.g., *thought*) served as a control

condition to test the effect of valence within the EMLN.

To instantiate subliminal word processing we applied a CFS paradigm (Fig. 1B). After

determining the perceptual threshold for each participant, the word stimuli (Word Type: MO,

EMneg, EMpos, NEU) were presented at five incremental visual contrast levels (Word Contrast:

two levels below, one at participant's perceptual threshold, and two above), with void stimuli as

perceptual control. Our CFS paradigm also required to indicate subjective ratings along a perceptual

awareness scale (PAS; Ramsøy and Overgaard, 2004), which allowed us to determine the presence

or absence of subjective perceptual awareness, independently of word contrast.

We first applied univariate analyses (Fig. 1C) to assess subliminal fMRI activation (1) as a

function of Word Contrast (reflecting the subliminal vs. supraliminal conditions defined based on

participant's perceptual threshold), and (2) as a function of PAS ratings (reflecting the subjective

perceptual awareness). In both analyses we tested the hypothesis that the MOVN and the EMLN

specifically respond to, respectively, MO and EM words following a "subliminal" activation

amplitude profile, which was defined as stable activation either (1) across the five incremental

contrast levels or (2) with and without subjective awareness. The subliminal profile, which would

speak for unaware processing, was compared against a "linear" profile, i.e., activation increase with

increasing word contrast (only in 1) and a "stepwise" profile, i.e., drop of activation either (1) for

word contrast below perceptual threshold or (2) without subjective awareness. In addition, for EM

we also tested the effects of valence by distinguishing EMneg, EMpos and NEU.

Secondly, we applied multivariate pattern analyses (Fig. 1D), again (1) as a function of

Word Contrast and (2) as a function of PAS ratings. We hypothesized that the involvement of

MOVN and EMLN for the representation and subliminal processing of, respectively, MO and EM

words would be reflected by the successful decoding of the word types (1) not only above but also

below perceptual threshold and (2) not only with but also without subjective perceptual awareness.

Participants

Twenty-five Italian native speakers voluntarily participated in the study. Two participants

were excluded due to the finding of brain structural anomalies. Two other participants did not

comply with the CFS task and were discarded from the analyses. The sample of the remaining 21

subjects comprised 7 male and 14 female participants (mean age = 22.31 years, SD = 4.01; mean

years of education = 15.76, SD = 3.25). All participants reported to be right-handed, which was in

line with the Edinburgh inventory assessment (Oldfield, 1971; mean score = 0.80, SD = 0.16, with

20 participants scoring in the range from 0.5 to 1 and thus classified as right-handed and one

participant scoring 0.4 and classified as ambidextrous). The Miles test (Miles, 1929) was

administered for evaluating eye dominance for the CFS task: 8 participants were left, and 13 right

eye dominant. All participants reported neither sensory deficits nor developmental, neurological, or

psychiatric diagnoses. No participants had reading deficits according to their performance with real

word (mean score = 100%, SD = 0%) and pseudoword (mean score = 99%, SD = 1%) reading tests

included in the Italian battery for assessing aphasic deficits (Miceli, 1994). All participants gave

written consent and received a reimbursement of 30 euro for their participation in the study. The

study was approved by the ethics committee of the University of Trento, Italy.

Stimuli

Word stimuli for the fMRI CFS task

As experimental word stimuli we used 75 nouns referring to MO and 75 to EM. As

described above, EM nouns were parametrically distributed along the negative-to-positive valence

dimension, including 25 EMneg and 25 EMpos nouns, with additional 25 NEU nouns as control

condition for testing the valence effects. EMneg and EMpos were defined as referring to "a mental

state that could be felt" (e.g., *anger* or *joy*; Jackson et al., 2019), also known as emotion-label nouns

(Conca et al., 2021). Emotion-laden nouns (e.g., *war* or *party*) were avoided. NEU were defined as

referring to "neutral mental states" (Jackson et al., 2019) that "can not be felt" (e.g., *thought*). In

line with these definitions, we checked that in the lexical database Multiwordnet (Pianta et al.,

2002) all EMneg, EMpos, and NEU nouns are classified as psychological features or factotum

(semantic domain level), but only EMneg and EMpos are defined as emotions or traits/attitudes

(affective level). The complete list of stimuli is available as Extended Data supporting Table 1 and

labeled as Table 1-1A.

To validate the variability of EM words along the negative-to-positive valence dimension,

all nouns were rated on a bipolar Likert scale (-3 = negative, 0 = neutral, +3 = positive) by an

independent group of 16 volunteers (10 female and 5 male participants, 1 did not specify the sex;

mean age = 29.25 years, SD = 5.48). Results showed that EMneg, EMpos, and NEU nouns were

clearly distributed along the negative-to-positive valence dimension, and significantly differed from

MO (Table 1A for summary statistics and Extended Data Table 1-1B for single word rating scores).

Ratings for arousal (1 = calm or relaxation, 7 = high excitement or agitation) were also collected, as

valence and arousal are considered to be independent dimensions of the semantic space (Osgood &

Suci, 1955). Results showed that EMpos, NEU, and MO did not differ in arousal, although all were

lower that EMneg (Table 1A). Finally, ratings on a unipolar Likert scale for imageability (1 =

barely imaginable, 7 = easily imaginable), which is considered as a proxy for concreteness

(Hoffman, 2016), showed that EM and NEU were less imageable than MO words. Experimental

stimuli were rated for valence, arousal and imageability also by all (but one) fMRI participants (on

average 8.55 days, SD = 9.97, after the MRI session). The results showed a high correlation

between the ratings of the fMRI and independent participant samples (Spearman's correlation

coefficient: valence: rho = 0.971; arousal: rho = 0.937; imageability: rho = 0.994, all p values

< .001).

Experimental stimuli were balanced for relevant sub-lexical and lexical variables (Table 1B

for detailed descriptive and inferential statistics). Lexical frequency, which has been demonstrated

to influence speed in word recognition (Brysbaert et al., 2016; but see Heyman & Moors, 2014),

was extracted from the SUBTLEX-IT database (Crepaldi et al., 2015) and matched across the four

types of words. Moreover, since sub-lexical processing has also been found to affect word

recognition (Westbury & Buchanan, 2002), MO, EMneg, EMpos, and NEU were matched for the

frequency of the first, last, and average medial bigrams of each word, plus the overall average

bigram frequency. Bigram frequencies were extracted from the PhonItalia database (Goslin et al.,

2014). Stimuli were also matched for their number of orthographic neighbours (again extracted

from the PhonItalia database, Goslin et al., 2014), for it has been demonstrated that a large

orthographic neighbourhood size makes words easier to read (Duñabeitia et al., 2008). Finally,

stimuli were balanced for their number of syllables, number of letters, and visual length calculated

in millimeters using the software GIMP 2.10.4 (www.gimp.org).

Mask stimuli for the CFS task

We used MATLAB R2017b to create 80 mask images with cyan background to make the

mask visible only to the dominant eye when wearing the red-cyan anaglyph goggles (see below).

Over the cyan background we displayed black patterns, which were similar in shape (e.g., closed

unfilled geometrical shapes such as ellipses and rectangles, and open u-shaped figures with different

orientations) and perceptual features (e.g., line width, dimension) to the Latin alphabet letters of the

word stimuli (Kido & Makioka, 2014; Axelrod et al., 2015). Then we overlayed a black frame and a

black fixation cross at the centre of every mask, using GIMP 2.10.4 (www.gimp.org).

Experimental procedure

Participants first performed a training session to familiarize with the CFS task outside the

MRI scanner room. Then, they entered into the MRI scanner and performed a pre-fMRI behavioral

session of the CFS task to assess their perceptual threshold in the same contextual and perceptual

conditions as for the fMRI CFS task. Finally, they performed the experimental CFS task session,

while their brain activity was measured with fMRI. In all these three sessions, participants wore

MRI-compatible red-cyan anaglyph goggles, with the cyan lens placed above the dominant eye. The

CFS trial structure was identical across the three sessions, as well as the use of Psychtoolbox-3

(v3.0.14, running in MATLAB 2017b) for stimulus delivery and behavioural response collection.

CFS trial

Each CFS trial began with the 200-ms presentation of a fixation cross on a red background

overlaid by an empty cyan image with 50% transparency (alpha blending). Then, a single word was

presented in black on a red background to make it visible only to the non-dominant eye when

wearing the red-cyan anaglyph goggles. Each word was written in uppercase (font: Helvetica, font

dimension: 80) and displayed in the center of the screen at a trial-specific (see below) visual

contrast level, for a duration of 1600 ms. In the same 1600-ms time interval, a set of 16 different

masks presented at 10 Hz was overlaid on the word with 50% transparency (Fig. 1B). For every

trial, the 16 masks were randomly extracted from the whole pool of 80 mask images and then

presented in a randomized order. To reduce after image effects (Hong & Blake, 2009), which can

occur at the offset of the last mask, we displayed for 1000 ms an image with multiple white capital

X letters on a black background. After a variable within-trial delay (pseudo-randomized duration of

either 1425, 1925, or 2425 ms in 4:2:1 proportion), participants were shown the display "1 2 3 4"

(duration 1000 ms), which represented the four level of the PAS (Ramsøy & Overgaard, 2004): 1 =

no experience; 2 = brief glimpse without recognition of any letter; 3 = almost clear perception,

recognition of one or more letters, but not of the entire word; 4 = absolutely clear perception and

recognition of the entire word. Participants were asked to press one of the four buttons of a response

pad according to their subjective level of visual perception of the word. To prevent explicit

semantic processing, questions assessing the objective perception of stimuli were avoided. The

variable within-trial delay was introduced in order to reduce the temporal correlation between the

stimulus visual presentation and the motor response to the PAS scale, thus preventing possible

biases in interpreting BOLD signal fluctuations. At the end of the trial, a delay of variable duration

(short: 2925 or 3425 ms, medium: 4075 or 4575 ms, long: variable from 5075 to 10575 ms, in 4:2:1

proportion) was included before the start of the next trial. Within-trial and between-trial delays were

determined by OPTseq2 (surfer.nmr.mgh.harvard.edu/optseq) to maximize the sensitivity of the

hemodynamic signal in the fMRI CFS event-related design (Dale, 1999). Trial timings were kept

identical also for the training and the pre-fMRI behavioral sessions.

Training session

Before entering the MRI scanner, participants were trained to perform the CFS task on a

laptop computer. After performing 6 practice trials, participants started the training session, which

comprised 105 consecutive trials. For 100 trials, we used nouns referring to jobs and careers (e.g.,

biologist), which were matched with the experimental stimuli with respect to lexical frequency,

number of syllables, and number of letters (all $p > .072$; Extended Data Table 1-2). These nouns

were presented at a specific contrast level according to the following procedure (similarly to

Tettamanti et al., 2017). For each participant, the 100 nouns were randomly ordered in a list and

then sequentially drawn without replacement from such list and assigned, in batches of 5, to 1 of 20

transparency levels (alpha blending value ranging from 8 to 255 = fully opaque, by steps of 13

units). In addition, for 5 trials we applied a transparency level with alpha = 0, which resulted in 5

empty trials (i.e., trials devoid of words, only including the red background and the flickering cyan

masks). The presentation order of the 105 trials was pseudo-randomized across participants.

Pre-fMRI behavioral session: psychophysical assessment of perceptual threshold

The perceptual threshold of each participant was assessed before the fMRI data acquisition

session. Participants were positioned in the MRI scanner and performed the CFS task on 105

consecutive trials, including 5 empty trials and 100 trials, during which 100 nouns referring to

living/natural items (e.g., *elephant*, *sunflower*, *storm*, *forest)* were presented at 20 different alpha

transparency levels (defined as in the training session described above). This set of nouns was

matched with the experimental stimuli with respect to lexical frequency, number of syllables,

number of letters, bigram frequency (initial, final, medial, average), and number of orthographic

neighbours (all $p > .103$; Extended Data Table 1-3).The same presentation procedure as in the

training session was applied.

To estimate the subjective perceptual threshold, we adopted the same procedure as in

Tettamanti et al. (2017). First, for each participant we plotted the number of responses with a PAS

score equal to 3 or 4 (indicating high to maximal subjective perceptual awareness) for each of the

20 transparency levels plus the empty trials. By using nonlinear least-squares estimation in

MATLAB 2017b, we fitted these values with a psychometric sigmoid function $f = L + (R − L) / \{1$

$+ \exp [− (x − i) / W]\}$, where x = transparency level, L = left horizontal asymptote (initial value =

0); R = right horizontal asymptote (initial value = 5), i = point of inflection (initial value = 2.5), and

W = width of the rising interval (initial value = 1). We thus defined the subjective perceptual

threshold (PT) as the first transparency level above the estimated point of inflection. Based on the

subjective perceptual threshold, five transparency levels were individually tailored to be used in the

fMRI session (and henceforth defined as Word Contrast levels). Specifically, two tailored contrast

levels were below PT: one defined as PT -25% (PT-2) and the other as PT -10% (PT-1). One

contrast level was at PT. Two tailored contrast levels were above PT: one defined as PT +10%

(PT+1), while the other was set for all participants at the maximum available contrast (Full, i.e.,

corresponding to alpha = 255). In addition, for all participants, we included a null contrast control condition (Empty), consisting of a red square devoid of any words. For participants with a subjective PT equal or lower than the transparency alpha level value of 69, the contrast levels below PT had to be tailored to a narrower range. Specifically, for participants (n = 9) with a subjective PT between alpha = 36 and 69, we set PT-2 to alpha = 5. For participants (n = 4) with a subjective PT lower than alpha = 36, we set PT-2 to alpha = 5 and PT-1 to alpha = 10.

fMRI CFS task

The fMRI experiment consisted in a within-subject event-related experimental design, where Word Type (MO, EMneg, EMpos, NEU) and Word Contrast (PT-2, PT-1, PT, PT+1, Full) were factorially manipulated. All participants completed 6 runs. Every run comprised 55 trials: 5 empty, 25 MO and 25 EM trials. All the 25 EM words in every run belonged to the same valence category (either EMneg, or EMpos, or NEU), and the EM category assignment to the runs was counterbalanced for every participant on the basis of a Latin square, such that all words of the three categories occurred once in the first three runs before being extracted again in the second three runs. Each of the 75 MO words were presented once in the first three runs and a second time in the last three runs. Thus, every word was presented twice throughout the experiment. Within every run, 5 MO and 5 EM were presented at one of the 5 word contrast levels. The presentation order of the different Word Type × Word Contrast combinations was unique for every participant and run, and determined by OPTseq2. The assignment of the specific word stimuli to the Word Type × Word Contrast combinations was pseudo-randomized for every participant and run, in order to minimize the number of stimuli presented at the same contrast in their first occurrence (during one of the first three runs) versus their second occurrence (during one of the last three runs) for a given participant.

MRI data acquisition

MRI scans were acquired with a 3 Tesla Siemens Magnetom Prisma whole body MRI scanner using a 64-channels head coil. During the PT estimation session, we obtained a T1-

weighted MPRAGE anatomical scan (288 slices, TR = 3280 ms, TE = 2.01 ms, flip angle = 7°;

FOV = 252 × 252 mm, slice thickness = 0.7 mm). During the experimental CFS task session,

whole-brain functional images were acquired with a T2*-weighted gradient-echo, EPI pulse

sequence, using BOLD contrast (TR = 1000 ms, TE = 28 ms, flip angle = 59°, multi-band

acceleration factor = 5). Each functional image comprised 65 transversal slices (2 mm thick),

acquired in interleaved mode (FOV = 200 x 200 mm). Each participant underwent 6 consecutive

functional scanning sessions, each comprising 520 scans and lasting 8 min and 50 s.

The stimuli were projected on a computer monitor located in the back of the scanner and

visible to the participants through a mirror placed on the head coil above their eyes. An MRI-

compatible fibre-optic response box with four buttons allowed the participants to give PAS score

responses using their left hand (index finger on the "4" button). The left hand was chosen to avoid

contamination of the BOLD signal acquired in left hemisphere where language processing is

lateralized, even though the overlapping between motor responses and word presentation was

anyway minimized thank to the within-trial delay.

Data analysis

Statistical analysis of behavioural data

To evaluate the effects of Word Type and Word Contrast on the PAS scores during fMRI,

we ran a series of Generalized Linear Mixed Models using R 4.2.2 (R Core Team, 2022). The first

simplest model comprised PAS scores as dependent variable, the six runs as fixed-effect confounds,

and the participants as random-effect factor. Word stimuli were also modelled as random-effect

factor, although we note that, since their variance equaled zero (SD = 0), their inclusion or omission

from the model yields identical results (Pasch et al., 2013). We then specified a series of models

with increasing complexity by adding Word Type (MO, EMneg, EMpos, NEU), Word Contrast

(PT-2, PT-1, PT, PT+1, Full), and the Word Type by Word Contrast interaction as fixed-effect

factors. All models were fit to a Poisson distribution, with a log-link function. To evaluate whether

each additional predictor significantly increased the model fit, we ran a χ2 loglikelihood ratio test

(declared significance α level: .05) between each hierarchically more complex model and the

simpler one.

Univariate statistical analyses of fMRI data

For MRI data preprocessing and univariate statistical analyses, we used SPM12 v7771

(www.fil.ion.ucl.ac.uk/spm). Structural MRI images were segmented and registered to the Montreal

Neurological Institute (MNI) standard space. Functional images were corrected for geometrical

distortions, spatially realigned, corrected for slice timing, and normalized to the MNI space by

using the segment procedure with the subject-specific segmented structural images as customized

segmentation priors. Functional images were spatially smoothed with a 4 mm FWHM Gaussian

kernel.

By applying a two-stage random-effects statistical approach, we performed univariate

statistical analyses of fMRI activation (1) as a function of Word Contrast and (2) as a function of

PAS rating. For all the reported univariate analyses, the significance threshold was declared at peak

level p = .05, using a small volume familywise error (FWE) type correction for multiple

comparisons, with spherical small volumes of 6 mm radius centered on the coordinates of interest

(x, y, z cartesian values in mm referring to the MNI space) within the MOVN and the EMLN (Fig.

1A). The coordinates for the MOVN were taken from the meta-analysis by Ishibashi et al. (2016),

and included the left ventral premotor cortex (LVPM; x = -50.3, y = 6.3, z = 30.7), the left inferior

(LIPL; x = -45.7, y = -31.7, z = 44.8) and superior parietal lobules (LSPL; x = -23.4, y = -61.1, z =

60.9), and the left middle temporal gyrus (LMTG; x = 17.8, y = 65.1, z = 1.8). For EMLN, the

coordinates were taken from Lindquist et al. (2012), and included the left (LAMY; x = -20, y = -4, z

= -16) and right amygdala (RAMY; x = 22, y = -4, z = -16), the left (LOFC; x = -40, y = 26, z = -6)

and right orbito-frontal cortex (ROFC; x = 42, y = 24, z = -4), the left anterior insula (LINS; x = -

38, y = 8, z = -10) and the sub-/pre-genual anterior cingulate cortex (sACC; x = 0, y = 38, z = 6).

(1) Univariate statistical analysis as a function of Word Contrast. At the first level, for each

participant we specified a general linear model, which comprised six separate runs, each including

11 regressors of interest: Empty, MOPT-2, MOPT-1, MOPT, MOPT+1, MOFull, EMPT-2, EMPT-

1, EMPT, EMPT+1, EMFull, with EM corresponding to either EMneg, EMpos, or NEU depending

on the run. Evoked responses were aligned to the onset of each trial (appearance of the fixation

cross), and their duration was fixed to 2800 ms (comprising 200 ms for the presentation of the

fixation cross, 1600 ms for the presentation of the masked stimuli, and 1000 ms for the presentation

of the anti-after effect image). Experimental confounds, namely PAS responses (for each trial, onset

aligned to the display of the PAS scale and duration equal to the interval between this time point

and the participant's response), task instructions, and head movement parameters yielded by image

realignment, were modelled as separate regressors. If there were trials in which participants did not

answer to the PAS scale (misses) or answered before the PAS question was displayed, these were

modelled in the same confound regressor with task instructions. For all experimental conditions, we

modelled the evoked responses as canonical haemodynamic response functions. The time series

were high-pass filtered at 128 s and pre-whitened by means of an autoregressive model AR(1).

Global normalization was performed to account for possible differences in signal due to the

presentation of EMneg, EMpos, and NEU stimuli in different runs. Statistical analyses were

restricted by using an explicit mask including only the voxels with gray matter tissue probability

> .1, as determined based on the segmented structural images of each participant.

We defined two different random-effects, full factorial designs. The first one was aimed at

testing the main effect of Word Type. Accordingly, within the estimated first-level general linear

model we defined a series of 20 condition-specific contrasts with the Empty baseline subtracted

(e.g., MOPT-2 − Empty). Namely, we defined a weight of +1 for the regressor of the specific

condition of interest (e.g., MOPT-2), a weight of -1 for the Empty baseline and a weight of 0 for all

other regressors. At the second-level, we entered the resulting 20 first-level contrasts into a model

including Word Type (4 levels: MO, EMneg, EMpos, NEU) and Word Contrast (5 levels: PT-2,

PT-1, PT, PT+1, Full) as within-subject factors. We assumed dependence and equal variance

between the levels of both factors. Within this full factorial design, we specified a series of T-

contrasts. Specifically, to examine the effects of MO we tested (i) MO − (EMneg+EMpos), whereas

to examine the effects of EM, we tested: (ii) (EMneg+EMpos) − MO. In addition for EM we tested

(iii) the simple effect of valence (i.e., with linear contrast weights: -1 for EMneg, 0 for NEU, and +1

for EMpos); (iv) the quadratic effect of valence (i.e., with non-linear weights: +1 for EMneg, -2 for

NEU, and +1 for EMpos).

The second random-effects, full factorial design aimed at testing whether the effects of MO

and EM as a function of Word Contrast followed either a subliminal (i.e., activation both above and

below perceptual threshold), a linear (i.e., gradual activation increase with contrast), or a stepwise

(i.e., activation only above perceptual threshold) profile (Fig. 1C). In order to model these three

different profiles, within the estimated first-level General Linear Model, we defined 24 condition-

specific contrasts with the Empty baseline not subtracted but explicitly modeled. Namely, we

defined a weight of +1 for the regressor of the specific condition of interest (e.g., MOPT-2) and a

weight of 0 for all the other regressors. At the second level, we entered the resulting 24 contrasts

into a model including the within-subject factors Word Type (4 levels: MO, EMneg, EMpos, NEU)

and Word Contrast (6 levels: Empty, PT-2, PT-1, PT, PT+1, Full), with dependence and equal

variance assumed between the levels of both factors. Then, we investigated whether the Word

Contrast level modulated the [MO − (EMneg+EMpos)] and [(EMneg+EMpos) − MO] activation

effects, following either a subliminal (weights: [-5 +1 +1 +1 +1 +1]; order: [Empty PT-2 PT-1 PT

PT+1 Full]), a linear [-2.5 -1.5 -0.5 +0.5 +1.5 +2.5] or a stepwise [-1 -1 -1 +1 +1 +1] BOLD

amplitude model.

(2) Univariate statistical analysis as a function of PAS. To test the MO and EM effects

without subjective awareness, but independently of Word Contrast, we further analyzed BOLD

responses as a function of PAS ratings. Specifically, for each type of words, all trials in which the

reported PAS score was either 1 or 2 (PAS{1,2}) were modelled as indicating no subjective

awareness, whereas all trials in which the PAS score was either 3 or 4 (PAS{3,4}) were modelled as

indicating subjective awareness. We restricted this analysis to only those subjects that produced

comparable response frequencies across all PAS levels. To exclude participants who did not satisfy

this criterion, we applied the same procedure as in Tettamanti et al. (2017): first we estimated the

maximum likelihood of the individual Contrast-to-PAS linear function to a Contrast-to-PAS linear

model function with R. Then we ranked participants based on the resulting Akaike information

criterion. None of the participants markedly departed from a linear increase of PAS score with

increasing contrast level. However, one participant was excluded from the following analyses

because of an exceedingly low number of PAS{3,4} responses, even for the PT+1 and Full contrast

levels (see Extended Data supporting Figure 2D and labeled as Figure 2-1). Thus, these analyses

were performed on a sub-sample of 20 participants.

At the first level, we specified a general linear model with the same parameters as defined

above for the analysis as a function of word contrast. However, here we modelled the six runs by

assigning the CFS trial events on the basis of PAS score and including the Empty baseline

condition. Specifically, each run included 5 regressors of interest: Empty, MOPAS{1,2},

MOPAS{3,4}, EMPAS{1,2}, EMPAS{3,4}, with EM being either EMneg, EMpos, or NEU

depending on the run. All the confound regressors were defined as in the analysis as a function of

contrast (see above). Within the estimated first-level general linear model, we defined a set of 12

condition-specific contrasts by specifying a weight of +1 for the regressor of interest (e.g.,

MOPAS{1,2}) and a weight of 0 for all the other regressors.

For the second-level, random-effects full factorial design, we entered the 12 condition-

specific contrasts of each participant into a model including the within-subject factors Word Type

(4 levels: MO, EMneg, EMpos, NEU) and PAS (3 levels: Empty, PAS{1,2}, PAS{3,4}), with

dependence and equal variance assumed between the levels of both factors. In this full factorial

design, we investigated whether the absence or presence of subjective perceptual awareness (as

reflected by the PAS score) modulated the activation effects of [MO − (EMneg+EMpos)] and

[(EMneg+EMpos) – MO], following a subliminal (weights: [-2 +1 +1]; order: [Empty PAS{1,2}

PAS{3,4}]) or a stepwise [-1 -1 2] BOLD amplitude model.

Multivariate pattern analyses (MVPA)

MVPA was carried out using the PyMVPA 2.6.5 toolbox (www.pymvpa.org) running under

Python 3.9.2. We first re-estimated the SPM12 univariate first-level analyses on preprocessed but

spatially unsmoothed fMRI data. We then entered the resulting spmT map images into the MVPA

(Misaki et al., 2010). All the included spmT maps incorporated the Empty baseline subtraction (e.g.,

[MOPT-2 – Empty]). We applied no Z-scoring or averaging within the PyMVPA toolbox. We

restricted the analysis to voxels with gray matter tissue probability > .1, which were comprised

within 12-mm radius spherical regions of interest in the MOVN and EMLN (Fig. 1A).

We trained a Gaussian Naive Bayes classifier on spmT maps to distinguish the levels of the

Word Type factor (MO, EMneg, EMpos) either between all three levels (3 x 3 classification

problems) or between pairwise combinations of these levels (2 x 2 classification problems), based

on a leave-one-subject-out cross-validation procedure, analogous to a second-level, random effects

analysis (Ghio et al., 2016). More specifically, we performed a searchlight analysis (Kriegeskorte et

al., 2006) with 4-mm radius spheres, and for each sphere the mean classification accuracy, as well

as the confusion matrix of predicted against actual classes, were calculated across leave-one-

subject-out folds.

A Monte Carlo procedure was applied to assess the significance of the classification

accuracies. In particular, the spmT map labels were randomly permuted over the factor levels in

each sphere 10000 times (Stelzer et al., 2013), and the actual classification accuracy was then

compared against the random permutation distribution with a declared P < .0001 threshold.

The mean classification accuracies, together with the corresponding confusion matrices are

reported for the significant searchlight spheres, resulting from the analysis of 2 different spmT map

sets. A first set entered the 3 x 3 and 2 x 2 multivariate pattern analyses as a function of Word

Contrast, with each classification problem assessed: i) across the 5 Word Contrast levels (PT-2, PT-

1, PT, PT+1, Full); ii) across levels below perceptual threshold (PT-2, PT-1); iii) across levels

above perceptual threshold (PT+1, Full). The second set entered the 3 x 3 and 2 x 2 multivariate

pattern analyses as a function of PAS ratings, with each classification problem assessed: i) across all

PAS scores (PAS{1,2}, PAS{3,4}); ii) across PAS{1,2}; ii) across PAS{3,4}.

For the 2 x 2 classification problems, we assessed the average spmT value across the voxels

of any significant searchlight spheres, and counted in each of the regions of interest within the

MOVN and the EMLN the presence of spheres whose average estimate reflected a greater

activation for condition A (e.g., MO) minus B (e.g., EMneg), or vice versa for condition B minus A

(note that each region of interest may contain spheres with average activation in just one of these

two directions, in both, or no significant spheres at all).

Code accessibility

The data and code related to the data analysis described above is available at

https://doi.org/10.17632/gfb3xb5yty.1 (Ghio et al., 2024).

Results

Behavioural results

The Generalized Linear Mixed Model analysis on the PAS behavioural responses during the

fMRI session showed that the Word Contrast level significantly modulated the level of subjective

awareness, $\chi 2(4) = 2286.3$, $p = 2.2 \times 10\text{-}16$ (Fig. 2A). Post-hoc multiple comparison tests (Tuckey's

corrected) revealed that every pairwise difference between levels of Word Contrast was significant

(all $p < .001$). The Word Type main effect ($\chi 2(3) = 4.107$, $p = 0.250$) and the Word Type by Word

Contrast interaction ($\chi 2(12) = 0.847$, $p = 1$) were not significant predictors of the PAS scores.

Univariate fMRI analyses

Effects of Word Type

We found that processing MO elicited greater activations than EM (i.e., EMneg+EMpos) in

all the regions of the MOVN (Table 2; Fig. 2B). In turn, when considering EM against MO, we

found activations in the sACC, the left OFC, the LAMY and, at trend level, the RAMY (Table 2;

Fig. 2B). We found no specific activation for neither the simple nor the quadratic effect of valence,

which therefore were not further investigated.

Effects of Word Type as a function of Word Contrast

We examined whether the levels of Word Contrast modulated activations within the MOVN

and the EMLN following a subliminal profile (i.e., activation both above and below perceptual

threshold), according to our experimental hypothesis. For comparison, we also tested a linear (i.e.,

gradual activation increase with contrast) and a stepwise (i.e., activation only above perceptual

threshold) BOLD amplitude model.

Subliminal BOLD modulations for MO compared to EM were found in three regions of the

MOVN, i.e., LVPM, LIPL, LSPL (Table 3; Fig. 2C). Linear and stepwise modulations for MO

compared to EM were found in all regions of the MOVN (Table 3). In turn, we did not find any

subliminal, linear, nor stepwise BOLD modulations for EM compared to MO.

Effects of Word Type as a function of PAS

To test the presence of Word Type effects in the absence of subjective awareness,

independently of Word Contrast, we further analyzed BOLD responses as a function of PAS

ratings. We tested whether activations within the regions of interest conformed to a subliminal

and/or a stepwise BOLD amplitude model.

We found stronger subliminal and stepwise BOLD modulations for MO compared to EM in

all regions of the MOVN (Table 4; Fig. 2D; Extended Data Figure 2-1). We did not found any

stronger subliminal nor stepwise BOLD modulations for EM compared to MO.

Multivariate classification of fMRI activation patterns

To evaluate whether activation differences between MO and EM, as resulting from the

univariate fMRI analyses, reflected distinct category-specific neural codes that could be

distinguished by a machine learning classifier, we also ran a searchlight MVPA on the same fMRI

data.

MVPA as a function of Word Contrast

In the 3 x 3 analyses as a function of Word Contrast (Fig. 3A), reflecting the main effect of

Word Type, significant spheres distinguishing among the 3 types of words with accuracy above

chance (chance level = 33.3%) were found, either when we considered all Word Contrast levels

(173 spheres distributed over all regions of interest except LAMY; mean classification accuracy =

44.4%), just those above PT (36 spheres distributed over RAMY, LOFC, ROFC, LINS, sACC,

LVPM, LIPL, LSPL, LMTG; mean accuracy = 49.8%), or, most importantly, just the Word

Contrast levels below PT (3 spheres over ROFC and LINS; mean accuracy = 50.8%).

In the 2 x 2 analyses as a function of Word Contrast (Table 5), reflecting the pairwise post-

hoc comparisons between the different word types, the MVPA classifier was able to distinguish

significantly above chance (chance level = 50%) among all word types pairs, again either when we

considered all Word Contrast levels, or just those above or just those below perceptual threshold.

Significant classification spheres were variably distributed across the 10 regions of interest, but with

a relative greater proportion in regions of the MOVN of spheres with greater activation for MO

(particularly the LSPL and LMTG), and a relatively greater proportion of spheres whose average

BOLD estimate reflected a greater activation for EM words (EMneg and EMpos) in the EMLN

regions of interest.

MVPA as a function of PAS

Also in the 3 x 3 analyses as a function of PAS ratings (Fig. 3B), the three word types where

significantly distinguished above chance (chance level = 33.3%), either when we considered all

PAS levels (33 spheres distributed over LOFC, ROFC, LINS, sACC LVPM, LSPL, LMTG; mean

classification accuracy = 49.6%), or just those indicating presence of subjective awareness (11

spheres distributed over LOFC, LINS, LVPM, and LSPL; mean accuracy = 55.2%). When we

considered just the PAS levels indicating absence of subjective awareness, significant spheres were

only found at a lower significance threshold (P < 0.001 against 10000 permutations; 9 spheres over

LOFC, ROFC, LINS, sACC, and LMTG; mean accuracy = 54.8%).

In the 2 x 2 analyses as a function of PAS ratings (Table 6), the MVPA classifier

distinguished above chance all word types pairs and for all three PAS rating level combinations (all

levels, just those indicating presence, or just those indicating absence of subjective awareness),

except for the pair EMneg versus EMpos (with and without subjective awareness), which was only

distinguished at a lower significance threshold (P < 0.001 against 10000 permutations).

Classification spheres were variably distributed across the 10 regions of interest, but with a slightly

greater proportion in MOVN regions of spheres with greater activation for MO, and a slightly

greater proportion of spheres whose average BOLD estimate reflected a greater activation for EM

(EMneg and EMpos) in EMLN regions of interest.

Discussion

The present fMRI study aimed at testing whether brain regions coding for experience-
derived information that constitutes conceptual representations are automatically recruited during
the subliminal perception of MO- and EM-related words. Univariate analyses of BOLD signal
showed a MO word-specific activation of the MOVN, a left-lateralized visuomotor network
involved in object-directed action and object identification (Caspers et al., 2010; Ishibashi et al.,
2016), and supposed to represent and process visual and motor-related features such as
manipulability and affordances (Buxbaum and Kalénine, 2010; Sakreida et al., 2016). Notably, this
activation was found even under subliminal unaware processing conditions, both when the words
were presented below perceptual threshold and when participants subjectively reported a lack of
stimulus awareness. EM word processing specifically engaged the EMLN, a bilateral limbic
network involved in emotion perception and recognition (Dalgleish, 2004; Dahlén et al., 2022), and
supposed to represent affective properties (Kober et al., 2008; Lindquist et al, 2012). Univariate
analyses, however, did not provide evidence for EM word-specific effects under subliminal
unaware processing conditions. MVPA, instead, showed that distinct MO and EM word-specific
neural codes could be distinguished above chance in MOVN and EMLN, when words were
presented both above and below the perceptual threshold, as well as with and without subjective
perceptual awareness.

Previous CFS studies have provided conflicting evidence for the recruitment of the MOVN
as a function of awareness when processing tool pictures subliminally: whereas some works found
activations in some but not all regions of this left-hemispheric premotor-parietal-temporal network
(Fang & He, 2005; Almeida et al., 2010; Suzuki et al., 2014), other studies did not report any
significant effects (Hesselmann & Malach, 2011; Hesselmann et al., 2011; Fogelson et al., 2014;
Ludwig et al., 2015, 2016). In contrast, we previously showed the recruitment and effective
connectivity of the MOVN both under aware and unaware processing of MO pictures (Tettamanti et
al., 2017; Ghio et al., 2022). Here, we confirmed and expanded these findings, in line with the

hypothesis that semantic features are automatically retrieved also during MO words processing,

thus despite the lack of a visual or other sensory depiction of tools, and so excluding the potentially

confounding effect of perceptual properties, such as the prototypical elongated shape of MO.

Indeed, we found that MO elicited a subliminal BOLD response profile as a function of Word

Contrast in LVPM, LIPL, and SPL of the MOVN, that is a stable activation regardless of stimulus

perception being under aware or unaware conditions. These three regions also exhibited a linear and

a stepwise response profile, in concomitance with the subliminal one. Taken together, the

significance of all three modulations indicates activation both above and below perceptual

threshold, with a relatively stronger activation with than without awareness, and with relatively

higher response amplitudes as the word contrast level increased. A fourth MOVN area, namely

LMTG, mainly implied in object identification (Bracci et al., 2012), showed only linear and

stepwise haemodynamic modulations. However, this effect does not contradict our hypothesis, as it

still suggests a significant activation of this region below the perceptual threshold, although weaker

than above. Furthermore, assessing fMRI activation as a function of PAS ratings revealed

significant subliminal and stepwise modulations for all MOVN regions, indicting that these regions

are activated both in the presence and, although to a lesser extent, in the absence of subjective

awareness. Consistently with the univariate analyses, MVPA showed that a supporting vector

machine classifier trained on the activity in MOVN regions was able to successfully tell apart MO

and EM activation maps, not only above but also below perceptual threshold, and both with and

without subjective perceptual awareness. Taken together, our results suggest that subliminal

perception of visually presented MO words automatically activate conceptual-semantic decoding of

visuomotor and functional MO-related information that is likely encoded during previous

experiences with tool exemplars (Binder & Desai, 2011; Lambon Ralph et al., 2017).

Regarding EM word processing, we found that it specifically involved the EMLN, a

network engaged in what Lindquist and colleagues (2012) defined as "core affect" domain. This

term refers to the mental representations sustaining the semantic processing of discrete emotions,

and comprising bodily sensations – somatovisceral, kinesthetic, proprioceptive, and neurochemical

– experienced along two main dimensions: valence and arousal. Previous CFS studies have

suggested that valence plays an important role in an EM stimulus emergence to consciousness

(Jiang & He, 2006; Vizueta et al., 2012), even though evidence is inconsistent. While some studies

reported a negativity bias (i.e., negatively-valenced words emerging from suppression faster than

positively-valenced and neutral stimuli; Yang & Yeh, 2011; Sklar et al., 2012), other works

reported the opposite pattern (Vinson et al., 2011) or no effect of valence (Rabagliati et al., 2018;

Cheng et al., 2019). Our univariate analysis revealed no subliminal response profile, either as a

function of Word Contrast or as a function of PAS scores, and no parametric effects of valence,

either simple or quadratic. However, the entire EMLN was consistently implicated in the

multivariate searchlight decoding of EMneg versus EMpos activation patterns, both when

considering contrast levels below perceptual threshold and the absence of subjective awareness.

Considering the theoretical and methodological differences between univariate and multivariate

analysis (Hebart & Baker, 2018) which make the latter more sensitive to detecting mental

representations (Norman et al., 2006), our findings suggest that voxels within EMLN regions are

likely not more activated for positively than negatively valenced EM words (or vice versa), but

rather exhibit patterns of distributed activations or de-activations which give rise to a clearly

identifiable EM-specific neural coding in the EMLN, even under unaware processing.

Our results about the subliminal activation of the MOVN and the EMLN suggest that

unaware processing of visually presented MO and EM words automatically elicits the retrieval of,

respectively, tool-related visuomotor properties (e.g., features useful to identify objects as well as to

functionally interact with them, such as manipulability and affordance) and core affect (e.g.,

interoceptive and somatic) information. Since (i) no overt task implying, for instance, imagery or

deep semantic processing, was required to participants, and (ii) neither operationalized nor

subjective awareness was implicated in word processing, we suggest that the category-specific

features encoded by MOVN and EMLN regions are constitutively integrated into the conceptual

representation of MO and EM. These findings are consistent with grounded cognition theories of

conceptual knowledge, according to which concepts rely, at least partially, on the reuse of specific

information derived by previous experience with similar percepts (Meteyard et al., 2012; Lambon

Ralph et al., 2017; Kiefer & Harpaintner, 2020). This information refers to sensory-motor

experiences in the case of MO (see Willems & Casasanto, 2011), and to internal (i.e.,

proprioceptive, somatovisceral) states in the case of EM (Winkielman et al., 2018). The hypothesis

that these category-specific features are encoded over multiple experiences of EM and MO

instances, represented in conceptual representations, and recruited every time the corresponding

concepts are semantically processed, is consistent with the view of the brain as a predictive system,

evolved to efficiently meet the ongoing needs of the body (Adams et al., 2013; Clark, 2013).

Whether the circumstances require to promptly interact with physical objects outside the body, or to

cope with physiological states inside the body, knowing how to plan an appropriate response is

essential for survival (Frye & Lindquist, in press). In this scenario, language plays an important

role: words rapidly and automatically activate properties of the entities they refer to, which overlap

with those elicited during the perception of and the experience with such entities. By considering a

broader perspective on language and semantic processing, this mechanism can be considered in line

with the idea suggested by Lupyan and colleagues that language acts as a flexible "artificial"

context, according to which the brain unconsciously generates predictive beliefs about incoming

inputs, possibly through a fast re-tuning of the organism's entire semantic network , thereby helping

constraining which type of top-down information is useful to recruit and what impact it has on

lower-level processes as well as on inference and reasoning (Lupyan & Ward, 2013; Lupyan &

Clark, 2015).

In conclusion, previous evidence has demonstrated that visually perceiving objects in the

absence of awareness elicits a comparable neurophysiological response signature to that evoked by

their conscious perception, thus reflecting not only a visual but also a semantic processing

(Fahrenfort et al., 2017; Tettamanti et al., 2017). We extend these findings by providing evidence

that also tool words, which are arbitrarily associated with their referents through learning and do not

carry any object-related visuomotor information, recruit a manipulable object-specific visuomotor

network. Additionally, we generalize these results to the domain of discrete emotions, by

demonstrating that processing emotion label words activates portions of the core affect limbic

network. Crucially, MO and EM specific neural activations are decoded even in the absence of

perceptual awareness, suggesting that experience-derived information constituting conceptual

representations is automatically recruited.

References

Adams, R.A., Shipp, S. & Friston, K.J. (2013). Predictions not commands: active inference in the motor system. Brain Structure and Function, 218(3), 611–643.

Almeida, J., Mahon, B. Z., & Caramazza, A. (2010). The role of the dorsal visual processing stream in tool identification. Psychological science, 21(6), 772-778.

Axelrod, V., Bar, M., Rees, G., & Yovel, G. (2015). Neural correlates of subliminal language processing. Cerebral Cortex, 25(8), 2160-2169.

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. Trends in cognitive sciences, 15(11), 527-536.

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. Cerebral cortex, 19(12), 2767-2796.

Borghi, A. M., Barca, L., Binkofski, F., & Tummolini, L. (2018). Varieties of abstract concepts: development, use and representation in the brain. Philosophical Transactions of the Royal Society B: Biological Sciences, 373(1752), 20170121.

Bracci, S., Cavina-Pratesi, C., Ietswaart, M., Caramazza, A., & Peelen, M. V. (2012). Closely overlapping responses to tools and hands in left lateral occipitotemporal cortex. Journal of neurophysiology, 107(5), 1443-1456.

Buxbaum, L. J., & Kalénine, S. (2010). Action knowledge, visuomotor activation, and embodiment in the two action systems. Annals of the New York Academy of Sciences, 1191(1), 201-218.

Brysbaert, M., Stevens, M., Mandera, P., & Keuleers, E. (2016). The impact of word prevalence on lexical decision times: Evidence from the Dutch Lexicon Project 2. Journal of Experimental Psychology: Human Perception and Performance, 42(3), 441.

Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action observation and imitation in the human brain. Neuroimage, 50(3), 1148-1167.

Chen, L., Lambon Ralph, M. A., & Rogers, T. T. (2017). A unified model of human semantic knowledge and its disorders. Nature human behaviour, 1(3), 0039.

Cheng, K., Ding, A., Jiang, L., Tian, H., & Yan, H. (2019). Emotion in Chinese words couldn't be extracted in continuous flash suppression. Frontiers in human neuroscience, 13, 309.

Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive science. Behavioral and Brain Sciences, 36(3), 181–204.

Conca, F., Borsa, V. M., Cappa, S. F., & Catricalà, E. (2021). The multidimensionality of abstract concepts: A systematic review. Neuroscience & Biobehavioral Reviews, 127, 474-491.

Crepaldi, D., Amenta, S., Pawel, M., Keuleers, E., & Brysbaert, M. (2015). SUBTLEX-IT. Subtitle-based word frequency estimates for Italian. In Proceedings of the Annual Meeting of the Italian Association For Experimental Psychology (pp. 10-12).

Dahlén, A. D., Schofield, A., Schiöth, H. B., & Brooks, S. J. (2022). Subliminal emotional faces elicit predominantly right-lateralized amygdala activation: a systematic meta-analysis of fMRI studies. Frontiers in neuroscience, 16, 868366.

Dale, A. M. (1999). Optimal experimental design for event-related fMRI. Human brain mapping, 8(2-3), 109-114.

Dalgleish, T. (2004). The emotional brain. Nature Reviews Neuroscience, 5(7), 583-589.

Diano, M., Celeghin, A., Bagnis, A., & Tamietto, M. (2017). Amygdala Response to Emotional Stimuli without Awareness: Facts and Interpretations. Frontiers in psychology, 7, 2029.

Duñabeitia, J. A., & Vidal-Abarca, E. (2008). Children like dense neighborhoods: Orthographic neighborhood density effects in novel readers. The Spanish journal of psychology, 11(1), 26-35.

Fahrenfort, J. J., Van Leeuwen, J., Olivers, C. N., & Hogendoorn, H. (2017). Perceptual integration without conscious access. Proceedings of the National Academy of Sciences, 114(14), 3744-3749.

Fang, F., & He, S. (2005). Cortical responses to invisible objects in the human dorsal and ventral pathways. Nature neuroscience, 8(10), 1380-1385.

Fogelson, S. V., Kohler, P. J., Miller, K. J., Granger, R., & Tse, P. U. (2014). Unconscious neural processing differs with method used to render stimuli invisible. Frontiers in psychology, 5, 601.

Frye, N.G. & Lindquist, K.A. (in press). Language and emotion concepts in the predictive brain. In J. Armony and P. Vuilleumier (Eds). 2nd edition. The Handbook of Affective Neuroscience: Cambridge University Press.

Ghio, M., & Tettamanti, M. (2016). Grounding sentence processing in the sensory-motor system. In Neurobiology of language (pp. 647-657). Academic Press.

Ghio, M., Vaghi, M. M. S., Perani, D., & Tettamanti, M. (2016). Decoding the neural representation of fine-grained conceptual categories. Neuroimage, 132, 93-103.

Ghio, M., Conca, F., Bellebaum, C., Perani, D., & Tettamanti, M. (2022). Effective connectivity within the neural system for object-directed action representation during aware and unaware tool processing. Cortex, 153, 55-65.

Ghio, M., Cassone, B., & Tettamanti, M. (2024). Unaware Processing of Words Activates Experience-Derived Information in Conceptual-Semantic Brain Networks. Bicocca Open Archive Research Data, V1, https://doi.org/10.17632/gfb3xb5yty.1.

Goslin, J., Galluzzi, C., & Romani, C. (2014). PhonItalia: a phonological lexicon for Italian. Behavior research methods, 46(3), 872-886.

Hebart, M. N., & Baker, C. I. (2018). Deconstructing multivariate decoding for the study of brain function. Neuroimage, 180, 4-18.

Hesselmann, G., & Malach, R. (2011). The link between fMRI-BOLD activation and perceptual awareness is "stream-invariant" in the human visual system. Cerebral Cortex, 21(12), 2829-2837.

Hesselmann, G., Hebart, M., & Malach, R. (2011). Differential BOLD activity associated with subjective and objective reports during "blindsight" in normal observers. Journal of Neuroscience, 31(36), 12936-12944.

Heyman, T., & Moors, P. (2014). Frequent words do not break continuous flash suppression differently from infrequent or nonexistent words: Implications for semantic processing of words in the absence of awareness. PLoS One, 9(8), e104719.

Hoffman, P. (2016). The meaning of 'life'and other abstract words: Insights from neuropsychology. Journal of neuropsychology, 10(2), 317-343.

Hong, S. W., & Blake, R. (2009). Interocular suppression differentially affects achromatic and chromatic mechanisms. Attention, Perception, & Psychophysics, 71(2), 403-411.

Ishibashi, R., Pobric, G., Saito, S., & Lambon Ralph, M. A. (2016). The neural network for tool-related cognition: an activation likelihood estimation meta-analysis of 70 neuroimaging contrasts. Cognitive neuropsychology, 33(3-4), 241-256.

Jackson, J. C., Watts, J., Henry, T. R., List, J. M., Forkel, R., Mucha, P. J., ... & Lindquist, K. A. (2019). Emotion semantics show both cultural variation and universal structure. Science, 366(6472), 1517-1522.

Jiang, Y., & He, S. (2006). Cortical responses to invisible faces: dissociating subsystems for facial-information processing. Current Biology, 16(20), 2023-2029.

Kido, K., & Makioka, S. (2014). Priming effects under continuous flash suppression: An examination on subliminal bottom-up processing. Japanese Psychological Research, 56(2), 126-138.

Kiefer, M., & Harpaintner, M. (2020). Varieties of abstract concepts and their grounding in perception or action. Open Psychology, 2(1), 119-137.

Kober, H., Barrett, L. F., Joseph, J., Bliss-Moreau, E., Lindquist, K., & Wager, T. D. (2008). Functional grouping and cortical–subcortical interactions in emotion: a meta-analysis of neuroimaging studies. Neuroimage, 42(2), 998-1031.

Kriegeskorte, N., Goebel, R., & Bandettini, P. (2006). Information-based functional brain mapping. Proceedings of the National Academy of Sciences, 103(10), 3863-3868.

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and computational bases of semantic cognition. Nature reviews neuroscience, 18(1), 42-55.

Lei, Y., Dou, H., Liu, Q., Zhang, W., Zhang, Z., & Li, H. (2017). Automatic processing of emotional words in the absence of awareness: The critical role of P2. Frontiers in Psychology, 8, 592.

Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F. (2012). The brain basis of emotion: a meta-analytic review. Behavioral and brain sciences, 35(3), 121-143.

Ludwig, K., Kathmann, N., Sterzer, P., & Hesselmann, G. (2015). Investigating category- and shape-selective neural processing in ventral and dorsal visual stream under interocular suppression. Human Brain Mapping, 36(1), 137-149.

Ludwig, K., Sterzer, P., Kathmann, N., & Hesselmann, G. (2016). Differential modulation of visual object processing in dorsal and ventral stream by stimulus visibility. Cortex, 83, 113-123.

Lupyan, G., & Clark, A. (2015). Words and the world: Predictive coding and the language-perception-cognition interface. Current Directions in Psychological Science, 24(4), 279-284.

Lupyan, G., & Ward, E. J. (2013). Language can boost otherwise unseen objects into visual awareness. Proceedings of the National Academy of Sciences, 110(35), 14196-14201.

Meteyard, L., Cuadrado, S. R., Bahrami, B., & Vigliocco, G. (2012). Coming of age: A review of embodiment and the neuroscience of semantics. Cortex, 48(7), 788-804.

Miceli, G., Laudanna, A., Burani, C., & Capasso, R. (1994). Batteria per l'analisi dei deficit afasici—BADA CEPSAG. Universita Cattolica del Sacro Cuore, Rome.

Miles, W. R. (1929). Ocular dominance deMWnstrated by unconscious sighting. Journal of experimental psychology, 12(2), 113.

Nakamura, K., Inomata, T., & Uno, A. (2020). Left amygdala regulates the cerebral reading network during fast emotion word processing. Frontiers in Psychology, 11, 1.

Noppeney, U., Price, C. J., Penny, W. D., & Friston, K. J. (2006). Two distinct neural mechanisms for category-selective responses. Cerebral Cortex, 16(3), 437e445.

Norman, K. A., Polyn, S. M., Detre, G. J., & Haxby, J. V. (2006). Beyond mind-reading: multi-voxel pattern analysis of fMRI data. Trends in Cognitive Sciences, 10(9), 424-430.

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. Neuropsychologia, 9(1), 97-113.

Osgood, C. E., & Suci, G. J. (1955). Factor analysis of meaning. Journal of experimental psychology, 50(5), 325.

Pasch, B., Bolker, B. M., & Phelps, S. M. (2013). Interspecific dominance via vocal interactions mediates altitudinal zonation in neotropical singing mice. The American Naturalist, 182(5), E161-E173.

Pianta, E., Bentivogli, L., & Girardi, C. (2002). MultiWordNet: developing an aligned multilingual database. In First international conference on global WordNet (pp. 293-302).

R Core Team. (2022). A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rabagliati, H., Robertson, A., & Carmel, D. (2018). The importance of awareness for understanding language. Journal of Experimental Psychology: General, 147(2), 190.

Ramsøy, T. Z., & Overgaard, M. (2004). Introspection and subliminal perception. Phenomenology and the cognitive sciences, 3(1), 1-23.

Sakreida, K., Effnert, I., Thill, S., Menz, M. M., Jirak, D., Eickhoff, C. R., ... & Binkofski, F. (2016). Affordance processing in segregated parieto-frontal dorsal stream sub-pathways. Neuroscience & Biobehavioral Reviews, 69, 89-112.

Sklar, A. Y., Levy, N., Goldstein, A., Mandel, R., Maril, A., & Hassin, R. R. (2012). Reading and doing arithmetic nonconsciously. Proceedings of the National Academy of Sciences, 109(48), 19614-19619.

Stelzer, J., Chen, Y., & Turner, R. (2013). Statistical inference and multiple testing correction in classification-based multi-voxel pattern analysis (MVPA): random permutations and cluster size control. Neuroimage, 65, 69-82.

Suzuki, M., Noguchi, Y., & Kakigi, R. (2014). Temporal dynamics of neural activity underlying unconscious processing of manipulable objects. Cortex, 50, 100-114.

Tettamanti, M., Conca, F., Falini, A., & Perani, D. (2017). Unaware processing of tools in the neural system for object-directed action representation. Journal of Neuroscience, 37(44), 10712-10724.

Tsuchiya, N., & Koch, C. (2005). Continuous flash suppression reduces negative afterimages. Nature neuroscience, 8(8), 1096-1101.

Vinson, D., Anderson, A., Ratoff, W., Bahrami, B., & Vigliocco, G. (2011). Slow to anger: emergence of emotionally loaded words and faces from interocular suppression. In Proceedings of the Annual Meeting of the Cognitive Science Society (Vol. 33, No. 33).

Vizueta, N., Patrick, C. J., Jiang, Y., Thomas, K. M., & He, S. (2012). Dispositional fear, negative affectivity, and neuroimaging response to visually suppressed emotional faces. NeuroImage, 59(1), 761–771.

Westbury, C., & Buchanan, L. (2002). The probability of the least likely non-length-controlled bigram affects lexical decision reaction times. Brain and Language, 81(1-3), 66-78.

Willems, R. M., & Casasanto, D. (2011). Flexibility in embodied language understanding. Frontiers in psychology, 2, 116.

Winkielman, P., Coulson, S., & Niedenthal, P. (2018). Dynamic grounding of emotion concepts. Philosophical Transactions of the Royal Society B: Biological Sciences, 373(1752), 20170127.

Yang, Y. H., & Yeh, S. L. (2011). Accessing the meaning of invisible words. Consciousness and cognition, 20(2), 223-233.

Table 1. *Descriptive and inferential statistics for rating scores and (sub)lexical variables characterizing the experimental stimuli*

Variable	MO	EMneg	EMpos	NEU	Statistics	*p* value
A. Rating scores						
Valence	0.01 (0.43)	-2.11 (0.35)	2.02 (0.35)	0.49 (0.53)	$\chi = 109.32$	2.2×10^{-16}
Arousal	3.17 (0.44)	4.97 (0.74)	3.02 (1.23)	2.84 (0.66)	$\chi = 59.476$	7.6×10^{-13}
Imageability	6.89 (0.14)	3.25 (0.61)	3.29 (0.58)	2.06 (0.33)	$\chi = 120.39$	2.2×10^{-16}
B. (Sub)lexical variables						
Lexical frequency	975.4 (1663.0)	879.7 (949.1)	1109.9 (1319.3)	966.3 (1462.2)	$\chi = 4.0229$	.259
First bigram frequency	36399.6 (32386.9)	51354.0 (47659.7)	37908.6 (35402.5)	47136.0 (42814.6)	$\chi = 2.3598$	.501
Last bigram frequency	70599.6 (49678.5)	76472.7 (55314.3)	75793.5 (60578.2)	79331.6 (40676.1)	$\chi = 2.4059$	.493
Medial bigram frequency	74687.2 (22995.9)	67007.0 (25336.2)	75560.5 (21534.3)	81195.7 (20707.4)	$F = 1.635$	.184
Mean bigram frequency	60562.1 (20598.2)	64944.6 (27122.5)	63087.5 (29200.5)	69221.1 (23766.0)	$\chi = 2.4212$	.490
N. orthographic neighbours	1.51 (1.51)	1.32 (0.94)	1.36 (1.58)	0.92 (1.52)	$\chi = 6.2852$	.098
N. syllables	3.48 (0.67)	3.56 (0.87)	3.80 (0.87)	3.80 (0.91)	$\chi = 5.3522$	.148
N. letters	8.76 (1.80)	8.32 (1.55)	8.72 (2.23)	9.20 (1.91)	$\chi = 2.7892$	.425
Visual length (mm)	31.04 (6.44)	28.89 (5.29)	29.59 (8.50)	31.88 (6.23)	$F = 1.169$	.324

Note. In the first four columns we report the mean and standard deviation (in brackets) for each Word Type (manipulable object words = MO, emotion words = EM, including negative = EMneg, positive = EMpos, and neutral words = NEU). In the last two columns, we report the results of the one-way ANOVA (*F* value, when data were normally distributed) or Kruskal-Wallis rank sum test (χ value, when data were not normally distributed) applied to test the main effect of Word Type. In table 2A, the main effect of Word Type resulted significant for valence, arousal, and imageability, and in what follows we therefore report the results of the post-hoc comparisons. For valence, we found a significant difference between all four types of words (all p values < .001). For arousal, EMneg showed significantly higher mean scores compared to all the other three categories (all p values < .001), which in turn did not differ from each other. For imageability, MO were significantly higher than EM (both EMneg and EMpos, both p-values < .001) and NEU (p value < .001). There was no significant difference between EMneg and EMpos, whereas all other post-

hoc comparisons revealed significant differences (all p values < .001). In Table 2B, the main effect of Word Type was never significant, indicating that the four types of words did not differ for any of the lexical and sub-lexical variables (see also the Materials and Methods section).

Table 2. *Effects of Word Type*

	Voxels	*p* value	Z score	x, y, z (mm)
MO > [EMneg+EMpos]				
LVPM	57	<.001	6.11	-46 8 28
LIPL	63	<.001	6.23	-42 -34 44
LSPL	57	<.001	5.86	-24 -64 58
LMTG	46	.003	4.95	-46 -62 -6
[EMneg+EMpos] > MO				
LAMY	6	.012	3.96	-22 -8 -16
RAMY	1	.042*	3.60	24 0 -14
LOFC	3	.010	4.02	-44 26 -10
sACC	2	.005	4.20	2 42 10

Note. MO = manipulable object words; EMneg = emotion words with negative valence; EMpos = emotion words with positive valence; LVPM = left ventral premotor cortex; LIPL= left inferior parietal lobule; LSPL = left superior parietal lobule; LMTG = left middle temporal gyrus; LAMY = left amygdala; RAMY = right amygdala, LOFC = left orbito-frontal cortex; sACC = sub-/pre-genual anterior cingulate cortex; x, y, z cartesian values in mm referring to the MNI space. *Trend level, assuming a Bonferroni correction and an alpha = .025.

Table 3. *Effects of Word Type as a function of Word Contrast*

	Voxels	*p* value	Z score	x, y, z (mm)
A. Subliminal				
MO > [EMneg+EMpos]				
LVPM	15	.001	4.51	-46 8 28
LIPL	23	.005	4.12	-42 -36 44
LSPL	34	.002	4.31	-24 -64 58
B. Linear				
MO > [EMneg+EMpos]				
LVPM	71	<.001	6.72	-46 8 28
LIPL	47	<.001	5.77	-44 -36 42
LSPL	57	<.001	4.70	-22 -62 56
LMTG	37	<.001	5.88	-44 -66 -6
C. Stepwise				
MO > [EMneg+EMpos]				
LVPM	84	<.001	6.40	-46 8 28
LIPL	60	<.001	6.82	-44 -36 42
LSPL	71	<.001	5.59	-28 -60 62
LMTG	24	<.001	5.70	-44 -66 -6

Note. MO = manipulable object words; EMneg = emotion words with negative valence; EMpos = emotion words with positive valence; LVPM = left ventral premotor cortex; LIPL= left inferior parietal lobule; LSPL = left superior parietal lobule; LMTG = left middle temporal gyrus; x, y, z cartesian values in mm referring to the MNI space.

Table 4. *Effects of Word Type as a function of PAS score*

	Voxels	*p* value	Z score	x, y, z (mm)
A. Subliminal				
MO > [EMneg+EMpos]				
LVPM	39	<.001	5.58	-46 8 28
LIPL	46	.001	4.51	-44 -34 44
LSPL	24	.002	4.02	-24 -64 60
LMTG	1	.001	3.62	-46 -68 -6
B. Stepwise				
MO > [EMneg+EMpos]				
LVPM	82	<.001	8.0	-46 8 28
LIPL	81	<.001	5.85	-44 -36 44
LSPL	34	.001	4.52	-28 -60 62
LMTG	33	<.001	5.13	-46 -68 -6

Note. MO = manipulable object words; EMneg = emotion words with negative valence; EMpos = emotion words with positive valence; LVPM = left ventral premotor cortex; LIPL= left inferior parietal lobule; LSPL = left superior parietal lobule; LMTG = left middle temporal gyrus; LAMY = left amygdala; x, y, z cartesian values in mm referring to the MNI space.

Table 5. *MVPA: 2 x 2 analyses as a function of Word Contrast*

classification	PT	Mean Acc	N	MOVN				EMLN					
				LVPM	LSPL	LIPL	LMTG	LAMY	RAMY	LOFC	ROFC	LINS	sACC
MO vs EMneg	All	63.4%	169	X	X		X	X	X X	X X	X X	X X	X
MO vs EMneg	aPT	69.4%	35	X	X X	X	X X		X	X	X X	X	X
MO vs EMneg	bPT	71.6%	11							X	X	X	
MO vs EMpos	All	62.6%	110	X	X X	X	X	X	X	X X	X X	X X	
MO vs EMpos	aPT	68.5%	17	X X	X		X		X				
MO vs EMpos	bPT	70.0%	5	X	X		X						
EMneg vs EMpos	All	64.0%	117	X		X	X	X	X X	X	X	X X	X
EMneg vs EMpos	aPT	70.1%	9			X			X	X		X	X
EMneg vs EMpos	bPT	71.4%	1									X	

Note. The "X" indicates the successful classification of the pair of word types within the region of interest, with the colour indicating the direction of associated mean spmT values. PT = perceptual threshold, aPT = above perceptual threshold, bPT = below perceptual threshold, N = number of voxels in the cluster. MO = manipulable object words; EMneg = emotion words with negative valence; EMpos = emotion words with positive valence. MOVN = MO-specific visuomotor network, comprising: LVPM = left ventral premotor cortex, LIPL= left inferior parietal lobule, LSPL = left superior parietal lobule, LMTG = left middle temporal gyrus. EMLN = EM-specific limbic network, comprising: LAMY = left amygdala, RAMY = right amygdala, LOFC = left orbito-frontal cortex, ROFC = right orbito-frontal cortex, LINS = left anterior insula, sACC = sub-/pre-genual anterior cingulate cortex.

Table 6. *MVPA: 2 x 2 analyses as a function of PAS*

classification	SA	Mean Acc	N	MOVN				EMLN					
				LVPM	LSPL	LIPL	LMTG	LAMY	RAMY	LOFC	ROFC	LINS	sACC
MO vs EMneg	All	70.5%	20			X	X	X X	X		X X		
MO vs EMneg	wSA	79.6%	6	X					X	X		X	
MO vs EMneg	w/oSA	81.3%	2							X		X	
MO vs EMpos	All	71.1%	8						X	X	X		
MO vs EMpos	wSA	76.0%	5			X	X			X			
MO vs EMpos	w/oSA	76.3%	2	X X									
EMneg vs EMpos	All	71.0%	18				X	X	X	X	X X	X	X
EMneg vs EMpos	wSA	75.0%	6	(X)		(X)	(X)			(X)		(X)	
EMneg vs EMpos	w/oSA	76.9%	4								(X)(X)	(X)	

Note. The "X" indicates the successful classification of the pair of word types within the ROI, with the colour indicating the direction of associated mean spmT values. SA = subjective awareness, wSA = with subjective awareness, w/oSA = without subjective awareness, N = number of voxels in the cluster. MO = manipulable object words; EMneg = emotion words with negative valence; EMpos = emotion words with positive valence. MOVN = MO-specific visuomotor network, comprising: LVPM = left ventral premotor cortex, LIPL= left inferior parietal lobule, LSPL = left superior parietal lobule, LMTG = left middle temporal gyrus. EMLN = EM-specific limbic network, comprising: LAMY = left amygdala, RAMY = right amygdala, LOFC = left orbito-frontal cortex, ROFC = right orbito-frontal cortex, LINS = left anterior insula, sACC = sub-/pre-genual anterior cingulate cortex.

Figure legends

Fig. 1. **Experimental hypotheses and procedures**. A) We tested the hypothesis (H1) that, in

comparison to emotion (EM) words with either a negative (EMneg) or a positive (EMpos) valence,

subliminally perceived manipulable object (MO) words selectively activated the MO-specific

visuomotor network (MOVN), constituted by the left ventral premotor cortex (LVPM), the left

inferior parietal lobule (LIPL) the left superior parietal lobule (LSPL), and the left middle temporal

gyrus (LMTG). Conversely, our second hypothesis (H2) predicted that stronger activations for the

subliminal perception of EMneg and EMpos compared to MO would be found in the EM-specific

limbic network (EMLN), comprising the left amygdala (LAMY), the right amygdala (RAMY), the

left orbito-frontal cortex (LOFC), the right orbito-frontal cortex (ROFC), the left anterior insula

(LINS), and the sub-/pre-genual anterior cingulate cortex (sACC). For both the H1 and H2

hypotheses, the alternative hypothesis (H0) was that a lack of selective activation in the MOVN or

EMLN networks for subliminal word processing would undermine the view that conceptual

semantic representations encompass experience-derived information. Anatomical spheres of interest

of 6 mm radius are rendered on a volumetric average brain image of all participants. B)

Experimental trials consisted in the presentation of a red-filtered written word (word types: MO,

EMneg, EMpos), which was only visible through the red anaglyph lense positioned on the non-

dominant eye. The word was presented either with null visual contrast (Empty baseline), at

subjective perceptual threshold (PT), and below (PT-2 and PT-1) or above (PT+1, Full) that

threshold. These subliminal or supraliminal visual contrast levels resulted from Continuous Flash

Suppression, instantiated through cyan-filtered mask overlays flickered at 10 Hz, which were only

visible through the cyan anaglyph lense positioned on the dominant eye. C) fMRI data were

inspected through univariate analysis for the presence of, respectively, MO-specific BOLD

responses in the MOVN or EM-specific responses in the EMLN, fitting a "subliminal", as opposed

to a "stepwise" or "linear", profile of activation as a function of Word Contrast (i.e., below or above

PT), and as a function of PAS rating (i.e., with or without subjective awareness (SA)). D)

Multivariate analysis of brain activation patterns sought for accurate (i.e., above chance) decoding

of Word Types, following a "subliminal" as opposed to a "stepwise" decoding profile; that is, a

successful decoding both below and above PT, and both with and without SA.

Fig. 2. **BOLD as a function of Word Type, Word Contrast, or PAS rating**. A) Mean (n = 21)

behavioral PAS rating scores for the experimental word types as a function of Word Contrast level.

Vertical bars indicate 95 % confidence intervals. B) Word Type-specific brain activation in,

respectively, the MOVN and the LOFC (as one of the significantly modulated EMLN brain

regions). Significant ($p < .05$, small volume FWE correction) brain activations are overlaid (MO:

purple clusters; EMneg/EMpos: blue clusters) on the small volumes of interest (MOVN: orange

spheres; EMLN: yellow spheres) and displayed on a volumetric rendering of the average anatomical

image of all participants. Dot-plots illustrate mean BOLD responses across all significant voxels.

Vertical bars indicate 95 % confidence intervals. C) Subliminal MO-specific brain activation in the

MOVN as a function of Word Contrast (all conventions are identical to panel B). D) Subliminal

MO-specific brain activation in the MOVN as a function of PAS rating (all conventions are

identical to panel B).

Fig. 3. **Multivariate pattern analysis**. Confusion matrices (labels for all matrices correspond to

those displayed on the top right matrix) representing the percent classification accuracy across all

significant ($p < .0001$; 10000 permutations) searchlight spheres. A) 3 x 3 classification of Word

Type as a function of Word Contrast for all levels above (left) or below perceptual threshold (right).

B) 3 x 3 classification of Word Type as a function of PAS rating for all levels indicating presence

(left) or absence of subjective awareness (right).

Figure 1. *MV*

Figure 2. *MV*

Figure 3. *MV*

Word Contrasts above PT   **Word Contrasts below PT**

PAS levels with SA   **PAS levels without SA**

Extended Data Table 1-1A. Lexical and sublexical variables of experimental stimuli.

Word stimuli	Word Type	N. orth. neigh.	Lex. freq.	First bigram freq.	Medial bigrams freq.	Last bigram freq.	Mean bigram freq.	N. syllables	N. letters	Visual length (mm)
accendino	MO	2	557	12476	81489.50	133168	75711.17	4	9	32.68
anello	MO	5	5515	42551	82939	34588	53359.33	3	6	21.51
annaffiatoio	MO	NaN	12	42551	52265	52265	49027	5	12	40.72
apriscatole	MO	0	62	10873	80625.25	73827	55108.42	5	11	38.44
spada	MO	4	4367	19719	20830.50	10909	17152.83	2	5	18.21
bastone	MO	5	1654	14724	101660.25	70433	62272.42	3	7	26.16
bicchiere	MO	1	3888	6547	80262.67	133974	73594.56	3	9	29.72
bisturi	MO	0	447	6547	64315	35748	35536.67	3	7	21.93
bollitore	MO	1	121	6010	97511.83	133974	79165.28	4	9	30.73
bottiglia	MO	1	4283	6010	63278.67	57191	42159.89	3	9	28.87
bottone	MO	3	952	6010	102066.50	70433	59503.17	3	7	26.16
braccialetto	MO	1	751	5011	76116.78	223642	101589.9267	4	12	41.99
bullone	MO	2	78	4458	92656.50	70433	55849.17	3	7	25.32
cacciavite	MO	1	312	46052	54964	86456	62490.67	4	10	33.1
caffettiera	MO	1	107	46052	70069.75	65653	60591.58	4	11	37.59
calcolatrice	MO	0	118	46052	88551.67	18061	50888.22	5	12	41.83
campanello	MO	2	877	46052	77158.71	34588	52599.57	4	10	37.17
cavatappi	MO	1	102	46052	75488.67	3639	41726.56	4	9	30.82
cellulare	MO	1	7074	14848	92517	133168	80177.67	4	9	31.58
cerniera	MO	1	213	14848	103106.60	65653	61202.53	3	8	28
chitarra	MO	1	1421	81739	87567.40	65653	78319.80	3	8	27.86
coltello	MO	3	4712	135447	86921.80	34588	85652.27	3	8	28.36
compasso	MO	2	35	135447	53220.20	40259	76308.73	3	8	31.5
computer	MO	0	8130	135447	63069	48415	82310.33	3	8	31.07
coperchio	MO	0	293	135447	61852.17	52265	83188.06	3	9	32.94
cucchiaio	MO	0	575	10568	39132.50	52265	33988.50	3	9	30.65
fazzoletto	MO	1	708	33631	55289.29	223642	104187.43	4	10	35.9
fiammifero	MO	1	330	25684	59668.43	48641	44664.48	4	10	34.46
forbici	MO	1	677	17526	69147.25	14195	33622.75	3	7	22.44
forchetta	MO	1	394	17526	70944.17	93005	60491.72	3	9	33.36
fucile	MO	5	3590	9500	40289	73827	41205.33	3	6	19.3
grattugia	MO	0	45	14632	88826.67	57191	53549.89	3	9	32
grilletto	MO	0	1584	14632	85315.83	223642	107863.28	3	9	30.65
interruttore	MO	1	652	107539	93962.89	133974	111825.30	5	12	43.01
manette	MO	2	1368	56674	97576	86456	80235.33	3	7	25.91
manganello	MO	1	99	56674	89747.43	34588	60336.48	4	10	37.76
maniglia	MO	2	306	56674	71121.60	57191	61662.20	3	8	26.33
mannaia	MO	1	70	56674	66843.75	57191	60236.25	3	7	24.8
manopola	MO	2	111	56674	69067	78467	68069.33	4	8	30.9
martello	MO	5	1141	56674	98616.60	34588	63292.87	3	8	29.38
matita	MO	4	559	56674	129059.33	93005	92912.78	3	6	20.24
mestolo	MO	2	63	28080	90120	34588	50929.33	3	7	26.59
orologio	MO	0	4034	13108	48697.60	52265	38023.53333	4	8	29.72
padella	MO	6	368	45016	69713.50	78467	64398.83	3	7	22.5

pennarello	MO	1	159	74312	105621.86	34588	71507.29	4	10	36.49
pennello	MO	3	335	74312	95097.40	34588	67999.13	3	8	29
pentola	MO	5	492	74312	126565.25	78467	93114.75	3	7	25.5
pettine	MO	4	233	74312	94042	70433	79595.67	3	7	23.28
pinzatrice	MO	NaN	47	33910	91263.14	18061	47744.71	4	10	32.6
portafoglio	MO	0	1513	42994	58623.13	52265	51294.04	4	11	39.54
pugnale	MO	1	859	13830	45160	73827	44272.33	3	7	25.82
pulsante	MO	1	1066	13830	88011	86456	62765.67	3	8	28.79
racchetta	MO	3	148	16575	57153.33	93005	55577.78	3	9	33.3
rasoio	MO	0	569	16575	23392.67	52265	30744.22	3	6	21.25
rastrello	MO	1	72	16575	92033.83	34588	47732.28	3	9	32.26
righello	MO	0	69	51082	53767.80	34588	46479.27	3	8	27.6
schiaccianoci	MO	0	72	25029	62087.80	14195	33770.60	4	13	43.6
scolapasta	MO	0	8	25029	56137.86	93005	58057.29	4	10	36.49
siringa	MO	1	417	50691	64356.75	5231	40092.92	3	7	22.8
spatola	MO	1	63	19719	92287	78467	63491	3	7	25.3
spazzola	MO	2	248	19719	36734.20	78467	44973.40	3	8	28.62
spazzolino	MO	1	479	19719	51548.43	133168	68145.14	4	10	34.71
spremiagrumi	MO	NaN	15	19719	37193.33	9105	22005.78	5	12	42.84
strofinaccio	MO	0	38	49172	51829.56	52265	51088.85	4	12	41.57
tagliaunghie	MO	NaN	16	10981	35408.22	12081	19490.07	4	12	40.72
tamburello	MO	0	73	10981	60408	34588	35325.67	4	10	36.91
tastiera	MO	2	259	10981	97769.80	65653	58134.60	3	8	27
telecomando	MO	1	551	25753	69158.88	36447	43786.29	5	11	41.91
temperino	MO	1	40	25753	87376.17	133168	82099.06	4	9	32.43
termometro	MO	0	154	25753	71501	48641	48631.67	4	10	39.45
trapano	MO	2	254	29265	84975.25	133168	82469.42	3	7	26.33
tubetto	MO	2	96	16862	52790.75	223642	97764.92	3	7	25.48
ventaglio	MO	0	139	25262	108094.67	52265	61873.89	3	9	31.41
volante	MO	1	1900	15502	122185	86456	74714.33	3	7	25.4
zuccheriera	MO	1	14	333	85319.88	65653	50435.29	4	11	39.45
angoscia	EMneg	1	401	42551	41736	57191	47159.33	3	8	28.87
antipatia	EMneg	1	61	42551	98048.50	57191	65930.17	5	9	28.79
avversione	EMneg	1	104	32484	92905.29	70433	65274.10	4	10	35.48
disagio	EMneg	0	1521	174105	45165.25	52265	90511.75	3	7	23.03
disgusto	EMneg	1	231	174105	44772.60	223642	147506.53	3	8	28.62
gelosia	EMneg	1	782	8939	61503.50	57191	42544.50	4	7	23.96
imbarazzo	EMneg	2	1989	14207	60728.67	7038	27324.56	4	9	31.92
infelicità	EMneg	0	212	107539	41897	93005	80813.67	5	10	29.8
inquietudine	EMneg	1	53	107539	36583.67	70433	71518.56	5	12	39.45
invidia	EMneg	3	517	107539	41897	57191	68875.67	3	7	20.15
malinconia	EMneg	1	181	56674	91237.57	57191	68367.52	5	10	34.12
nostalgia	EMneg	1	566	52903	77439.33	57191	62511.11	4	9	31.41
ostilità	EMneg	0	368	3806	82699.40	93005	59836.80	4	8	24.81
panico	EMneg	2	2866	45016	107347	26810	59724.33	3	6	21
rabbia	EMneg	3	3431	16575	23626.67	57191	32464.22	2	6	20.07
rammarico	EMneg	1	191	16575	77057.17	26810	40147.39	4	9	33.7
rancore	EMneg	1	895	16575	104412	133974	84987	3	7	27.09
repulsione	EMneg	0	65	23590	55560.14	70433	49861.05	4	10	35.22

rimorso	EMneg	3	679	51082	57015.50	40259	49452.17	3	7	26.08
rimpianto	EMneg	2	459	51082	92641.33	223642	122455.11	3	9	30.56
sdegno	EMneg	1	90	170	34449.33	133168	55929.11	2	6	23.45
sofferenza	EMneg	1	1445	57963	77515.57	25784	53754.19	4	10	36.66
terrore	EMneg	2	1148	25753	108271.80	133974	89332.93	3	7	26.59
tristezza	EMneg	1	1109	29265	67533.17	25784	40860.72	3	9	30.06
vergogna	EMneg	3	2628	25262	53131.80	61022	46471.93	3	8	31.41
affetto	EMpos	6	1602	5695	56892.50	223642	95409.83	3	7	24.89
allegria	EMpos	1	371	68073	65022.40	57191	63428.80	4	8	26.84
ammirazione	EMpos	1	272	11832	74967.13	70433	52410.71	5	11	38.69
divertimento	EMpos	1	2717	174105	99486.56	223642	165744.52	5	12	41.57
entusiasmo	EMpos	2	853	4561	62046.86	27069	31225.62	4	10	36
estasi	EMpos	2	311	26683	98875	28994	51517.33	3	6	19.47
euforia	EMpos	1	128	1839	63701.25	57191	40910.42	4	7	24.3
felicità	EMpos	1	2929	8694	87007.20	93005	62902.07	4	8	24.13
fiducia	EMpos	0	5456	25684	26870.25	57191	36581.75	3	7	21.84
gioia	EMpos	4	3090	28311	35936	57191	40479.33	2	5	15.66
gradimento	EMpos	1	257	14632	93770.29	223642	110681.43	4	10	37
gratitudine	EMpos	1	775	14632	86536.13	70433	57200.38	5	11	37.25
ilarità	EMpos	0	20	67973	98161.75	93005	86379.92	4	7	21
letizia	EMpos	0	62	38814	57952.75	57191	51319.25	3	7	20.48
meraviglia	EMpos	3	1939	28080	73575.57	57191	52948.86	4	10	33.6
ottimismo	EMpos	0	200	2888	61817.17	27069	30591.39	4	9	31.16
passione	EMpos	1	2456	45016	99446	70433	71631.67	3	8	28.11
riconoscenza	EMpos	1	163	51082	82940.33	25784	53268.78	5	12	44
serenità	EMpos	0	246	58313	123363	93005	91560.33	4	8	27.52
simpatia	EMpos	1	440	50691	79781	57191	62554.33	4	8	25.57
soddisfazione	EMpos	1	805	57963	54687.30	70433	61027.77	5	13	44.79
sollievo	EMpos	0	1189	57963	77574	12497	49344.67	3	8	27.09
stima	EMpos	5	742	49172	63435.50	22625	45077.50	2	5	16.93
tenerezza	EMpos	1	271	25753	88882	25784	46806.33	4	9	32.43
tranquillità	EMpos	0	454	29265	76285.11	93005	66185.04	4	12	39.54
abitudine	NEU	1	1455	9465	51751.50	70433	43883.17	5	9	29.38
buonsenso	NEU	0	187	4458	82890.33	40259	42535.78	3	9	34.71
cautela	NEU	1	426	46052	54334.50	78467	59617.83	3	7	24.98
contegno	NEU	2	135	135447	104467	133168	124360.67	3	8	31.41
criterio	NEU	0	117	9576	115488.80	52265	59109.93	3	8	26.33
decenza	NEU	3	269	116530	70689.25	25784	71001.08	3	7	25.99
discrezione	NEU	0	656	174105	73872.25	70433	106136.75	4	11	37.51
giudizio	NEU	0	2352	28311	110561	52265	63712.33	3	8	24.55
imparzialità	NEU	0	53	14207	71884.78	93005	59698.93	5	12	37.76
impressione	NEU	2	2782	14207	89438.50	70433	58026.17	4	11	38.02
memoria	NEU	2	5175	28080	74617.50	57191	53296.17	3	7	26
moderazione	NEU	0	106	29591	96005.88	70433	65343.29	5	11	40.47
neutralità	NEU	0	34	40713	76388.43	93005	70035.48	4	10	34.21
normalità	NEU	1	620	52903	78091.17	93005	74666.39	4	9	32.09
obiettività	NEU	0	66	1240	64806.50	93005	53017.17	5	11	33.95
opinione	NEU	1	3404	4461	94509	70433	56467.67	4	8	26.8
parsimonia	NEU	0	24	45016	78901.43	57191	60369.48	4	10	34.46

pensiero	NEU	1	4320	74312	102951	48641	75301.33	3	8	28.28
presentimento	NEU	1	785	66996	115006.20	223642	135214.73	5	13	47.07
prudenza	NEU	1	520	66996	53890.60	25784	48890.20	3	8	29.89
quotidianità	NEU	0	21	59294	68089.22	93005	73462.74	5	12	39.2
raziocinio	NEU	0	21	16575	62232.60	52265	43690.87	4	10	32
regolarità	NEU	0	89	23590	77927.71	93005	64840.90	5	10	35.22
senno	NEU	7	416	58313	113335	133168	101605.33	2	5	19
sobrietà	NEU	0	125	57963	47763.20	93005	66243.73	3	8	27.77

Extended Data Table 1-1B. Descriptive statistics of the rating scores for valence (val.), arousal (aro.), and imaginability (ima.) for every experimental stimulus obtained in the preliminary rating study on independent sample of 16 participants.

Word stimuli	Word Type	Mean val.	Sd val.	Mean aro.	Sd aro.	Mean ima.	Sd ima.
accendino	MO	0.06	0.44	3.25	1.24	7.00	0.00
anello	MO	0.56	0.63	2.81	1.38	6.88	0.34
annaffiatoio	MO	0.00	0.00	2.88	1.50	6.94	0.25
apriscatole	MO	0.13	0.34	3.00	1.41	6.81	0.75
spada	MO	0.0625	0.77	3.75	1.29	7.00	0.00
bastone	MO	-0.25	0.45	3.56	1.31	6.94	0.25
bicchiere	MO	0.38	0.62	3.00	1.41	7.00	0.00
bisturi	MO	-0.81	0.98	4.13	1.75	6.81	0.75
bollitore	MO	NaN	NaN	NaN	NaN	NaN	NaN
bottiglia	MO	0.19	0.54	3.00	1.37	7.00	0.00
bottone	MO	0.06	0.25	2.94	1.44	6.94	0.25
braccialetto	MO	NaN	NaN	NaN	NaN	NaN	NaN
bullone	MO	0.06	0.25	3.00	1.41	7.00	0.00
cacciavite	MO	0.13	0.34	3.06	1.44	6.88	0.50
caffettiera	MO	NaN	NaN	NaN	NaN	NaN	NaN
calcolatrice	MO	0.38	0.62	2.94	1.44	7.00	0.00
campanello	MO	0.06	0.25	3.19	1.47	6.94	0.25
cavatappi	MO	0.13	0.34	2.88	1.50	7.00	0.00
cellulare	MO	0.25	0.77	3.06	1.48	7.00	0.00
cerniera	MO	0.13	0.50	3.00	1.41	6.81	0.75
chitarra	MO	0.63	0.50	2.94	1.53	7.00	0.00
coltello	MO	-0.38	0.62	4.13	1.31	7.00	0.00
compasso	MO	0.19	0.54	2.88	1.50	7.00	0.00
computer	MO	0.50	1.03	3.13	1.54	7.00	0.00
coperchio	MO	0.06	0.25	2.88	1.50	6.94	0.25
cucchiaio	MO	0.06	0.25	3.06	1.44	7.00	0.00
fazzoletto	MO	0.13	0.34	2.94	1.34	6.81	0.75
fiammifero	MO	0.13	0.34	3.19	1.05	6.63	1.50
forbici	MO	0.00	0.37	3.25	1.18	6.75	1.00
forchetta	MO	0.19	0.40	2.88	1.50	6.94	0.25
fucile	MO	-1.31	1.14	4.19	1.56	6.94	0.25
grattugia	MO	0.06	0.57	2.88	1.50	7.00	0.00
grilletto	MO	-0.81	1.11	4.13	1.67	6.44	1.26
interruttore	MO	0.06	0.25	2.94	1.44	6.88	0.34
manette	MO	-1.00	0.82	4.25	1.39	6.94	0.25
manganello	MO	-1.50	1.10	4.38	1.31	6.88	0.50
maniglia	MO	0.06	0.25	3.06	1.44	6.81	0.75
mannaia	MO	-0.81	0.83	3.94	1.24	6.44	1.36
manopola	MO	0.00	0.00	3.00	1.41	6.69	0.79
martello	MO	0.13	0.50	3.13	1.41	6.94	0.25
matita	MO	0.69	0.70	2.69	1.54	7.00	0.00
mestolo	MO	0.19	0.54	2.88	1.50	7.00	0.00
orologio	MO	NaN	NaN	NaN	NaN	NaN	NaN

padella	MO	0.31	0.48	3.00	1.41	7.00	0.00
pennarello	MO	0.44	0.63	2.81	1.42	6.56	1.50
pennello	MO	0.31	0.48	2.88	1.50	6.69	1.25
pentola	MO	0.19	0.40	2.88	1.50	7.00	0.00
pettine	MO	-0.13	0.81	3.00	1.51	7.00	0.00
pinzatrice	MO	0.06	0.25	3.00	1.41	6.81	0.54
portafoglio	MO	0.06	0.44	3.13	1.41	6.81	0.75
pugnale	MO	-0.94	0.93	4.38	1.36	6.56	1.26
pulsante	MO	0.06	0.25	3.06	1.48	6.81	0.54
racchetta	MO	0.50	0.63	2.94	1.57	6.94	0.25
rasoio	MO	0.00	0.37	3.38	1.09	7.00	0.00
rastrello	MO	0.00	0.00	2.94	1.44	7.00	0.00
righello	MO	0.06	0.25	2.81	1.47	6.94	0.25
schiaccianoci	MO	0.00	0.37	2.81	1.47	6.94	0.25
scolapasta	MO	NaN	NaN	NaN	NaN	NaN	NaN
siringa	MO	-1.06	0.93	4.19	1.47	7.00	0.00
spatola	MO	0.19	0.54	2.88	1.50	6.63	0.89
spazzola	MO	0.06	0.68	3.00	1.41	6.81	0.75
spazzolino	MO	0.13	0.50	2.94	1.44	7.00	0.00
spremiagrumi	MO	0.38	0.62	2.75	1.39	6.81	0.54
strofinaccio	MO	0.00	0.52	2.88	1.50	6.88	0.50
tagliaunghie	MO	0.13	0.34	3.00	1.41	6.81	0.75
tamburello	MO	0.38	0.62	3.00	1.37	7.00	0.00
tastiera	MO	0.25	0.58	3.13	1.54	7.00	0.00
telecomando	MO	0.19	0.54	3.13	1.36	7.00	0.00
temperino	MO	0.19	0.40	2.88	1.50	7.00	0.00
termometro	MO	-0.19	0.83	3.25	1.48	7.00	0.00
trapano	MO	-0.13	0.62	3.63	1.86	7.00	0.00
tubetto	MO	0.00	0.00	2.88	1.50	6.75	1.00
ventaglio	MO	0.31	0.48	2.94	1.34	6.81	0.54
volante	MO	0.31	0.79	2.75	1.34	7.00	0.00
zuccheriera	MO	NaN	NaN	NaN	NaN	NaN	NaN
angoscia	EMneg	-2.81	0.40	5.81	1.52	3.13	1.89
antipatia	EMneg	-1.75	0.45	4.81	0.98	3.06	1.53
avversione	EMneg	-1.81	1.11	4.75	0.93	3.06	1.61
disagio	EMneg	-2.38	0.72	4.75	1.53	2.75	1.81
disgusto	EMneg	-2.25	0.58	4.56	0.81	3.75	1.39
gelosia	EMneg	-1.75	1.00	5.19	1.17	3.81	1.97
imbarazzo	EMneg	-1.38	1.26	4.94	1.29	3.50	1.83
infelicità	EMneg	-2.44	0.63	4.56	1.82	3.75	1.69
inquietudine	EMneg	-2.31	0.79	5.38	1.50	2.56	1.75
invidia	EMneg	-2.19	0.98	5.25	1.06	3.31	1.82
malinconia	EMneg	-1.81	0.91	3.44	1.67	2.81	1.47
nostalgia	EMneg	-1.81	0.98	3.81	1.33	2.63	1.82
ostilità	EMneg	-2.06	0.85	5.69	1.25	3.00	1.75
panico	EMneg	-2.63	0.62	6.63	0.81	4.13	1.78
rabbia	EMneg	-1.88	1.31	5.75	1.18	4.31	1.66
rammarico	EMneg	-1.69	0.70	4.44	0.89	2.38	1.54
rancore	EMneg	-2.25	1.13	5.44	1.15	2.75	1.73

repulsione	EMneg	-1.88	0.72	4.75	1.00	3.38	1.67
rimorso	EMneg	-2.19	0.83	4.75	1.06	2.19	1.52
rimpianto	EMneg	-2.19	0.83	4.44	1.21	2.50	1.75
sdegno	EMneg	-1.69	1.14	4.75	0.45	3.00	1.67
sofferenza	EMneg	-2.50	0.82	5.13	1.41	4.06	1.65
terrore	EMneg	-2.63	0.62	6.25	1.18	4.06	1.34
tristezza	EMneg	-2.19	0.75	3.81	1.64	3.88	1.26
vergogna	EMneg	-2.19	0.83	5.25	1.06	3.56	1.75
affetto	EMpos	2.38	0.81	2.25	1.44	3.44	2.06
allegria	EMpos	2.38	0.72	3.94	1.98	4.00	1.46
ammirazione	EMpos	1.75	0.86	2.94	1.34	2.56	1.59
divertimento	EMpos	2.31	0.79	4.31	1.49	3.88	1.96
entusiasmo	EMpos	2.19	0.75	4.75	1.81	3.75	1.84
estasi	EMpos	1.75	1.29	3.00	1.71	3.63	1.86
euforia	EMpos	1.75	1.18	5.38	1.54	3.88	1.67
felicità	EMpos	2.56	0.51	3.75	1.88	4.06	1.88
fiducia	EMpos	2.25	0.77	1.88	1.26	2.44	1.63
gioia	EMpos	2.50	0.73	4.06	1.88	4.19	1.80
gradimento	EMpos	1.44	0.96	2.50	1.26	2.81	1.76
gratitudine	EMpos	1.94	0.77	2.44	1.36	2.56	1.75
ilarità	EMpos	1.38	1.41	4.25	1.53	3.25	2.08
letizia	EMpos	1.38	1.09	2.38	1.36	2.88	1.71
meraviglia	EMpos	2.63	0.62	4.00	1.55	3.50	1.51
ottimismo	EMpos	1.81	1.22	2.44	1.21	2.69	1.96
passione	EMpos	2.19	0.83	5.44	1.63	4.13	1.71
riconoscenza	EMpos	1.88	0.81	2.25	1.06	2.63	2.00
serenità	EMpos	2.25	0.68	1.44	0.89	3.31	1.89
simpatia	EMpos	2.00	0.82	3.19	1.17	2.81	1.68
soddisfazione	EMpos	2.13	0.81	2.06	1.12	3.38	2.25
sollievo	EMpos	2.00	0.73	1.50	0.73	3.25	2.11
stima	EMpos	1.88	0.89	2.56	1.46	2.44	1.90
tenerezza	EMpos	2.13	0.62	1.50	0.63	3.81	1.64
tranquillità	EMpos	1.75	0.86	1.25	0.77	3.06	1.44
abitudine	NEU	0.00	1.10	2.13	1.45	2.38	1.96
buonsenso	NEU	1.38	0.96	2.38	1.15	1.69	1.30
cautela	NEU	0.25	0.77	3.13	1.36	2.13	1.26
contegno	NEU	0.25	0.86	2.81	1.22	2.00	1.15
criterio	NEU	0.69	0.95	2.88	1.31	1.44	0.73
decenza	NEU	0.13	1.15	2.69	1.20	2.06	1.44
discrezione	NEU	0.88	0.81	2.44	1.21	1.75	1.29
giudizio	NEU	-0.25	1.24	3.94	1.77	2.44	1.71
imparzialità	NEU	0.81	0.98	2.81	1.47	2.00	1.41
impressione	NEU	0.25	0.68	3.88	1.41	1.69	1.40
memoria	NEU	1.06	1.29	2.88	1.31	2.25	1.61
moderazione	NEU	0.38	0.96	2.38	1.31	1.81	1.17
neutralità	NEU	-0.13	0.96	3.00	1.32	1.81	1.05
normalità	NEU	0.44	0.73	2.31	1.30	2.00	1.32
obiettività	NEU	0.88	0.89	2.31	1.25	1.81	1.47
opinione	NEU	0.94	1.12	3.75	1.53	2.31	2.09

parsimonia	NEU	0.13	1.09	2.81	1.28	1.81	1.11
pensiero	NEU	1.13	1.41	3.13	1.89	2.38	1.67
presentimento	NEU	-0.94	0.77	4.75	1.24	2.13	1.67
prudenza	NEU	0.56	0.63	3.06	1.34	2.13	1.31
quotidianità	NEU	0.56	0.81	2.31	1.20	2.75	2.08
raziocinio	NEU	1.25	1.06	2.56	1.21	1.88	1.31
regolarità	NEU	0.38	0.89	1.75	1.06	2.44	1.55
senno	NEU	1.00	1.03	2.44	1.21	1.69	1.08
sobrietà	NEU	0.31	0.95	2.50	1.21	2.69	1.82

Extended Data Table 1-2A. 100 words referring to jobs and careers used in the training session.

1.	allenatore	51.	ginecologa
2.	archeologa	52.	giornalista
3.	architetto	53.	giudice
4.	astronauta	54.	hostess
5.	astronomo	55.	idraulico
6.	attore	56.	imbianchino
7.	bagnina	57.	infermiera
8.	balia	58.	ingegnere
9.	ballerina	59.	insegnante
10.	banchiere	60.	macchinista
11.	barbiere	61.	macellaio
12.	barista	62.	magistrato
13.	benzinaio	63.	maresciallo
14.	biologa	64.	minatore
15.	calciatore	65.	modella
16.	calzolaio	66.	neurologa
17.	cameriera	67.	notaio
18.	camionista	68.	oste
19.	cantante	69.	ostetrica
20.	carabiniere	70.	parrucchiera
21.	cardiologa	71.	pasticcere
22.	carpentiere	72.	pediatra
23.	casalinga	73.	pescatore
24.	cassiere	74.	pilota
25.	centralinista	75.	pittore
26.	chef	76.	poeta
27.	chitarrista	77.	pompiere
28.	ciclista	78.	postino
29.	commessa	79.	prestigiatore
30.	contabile	80.	prete
31.	contadino	81.	psicologa
32.	coreografa	82.	ragioniere
33.	dentista	83.	redattore
34.	deputato	84.	regista
35.	disinfestatore	85.	rettore
36.	economista	86.	ricercatore
37.	elettricista	87.	sarta
38.	estetista	88.	scenografo

39. fabbro
40. falegname
41. farmacista
42. filosofa
43. fioraia
44. fisioterapista
45. fornaio
46. fotografa
47. fruttivendolo
48. gelataio
49. geologa
50. giardiniere

89. sceriffo
90. scrittrice
91. scultore
92. segretaria
93. senatore
94. sindaco
95. soldato
96. stilista
97. tassista
98. tipografo
99. truccatrice
100. usciere

Extended Data Table 1-2B. Lexical and sublexical variables of training stimuli referring to jobs and careers, matched with manipulable objects (MO) and emotion (EM, with negatively valenced [EMneg], positively valenced [EMpos], and neutral [NEU]) word stimuli, mean (sd).

Variable	MO	EMneg	EMpos	NEU	Career	Kruskal-Wallis	*p* value
Lexical frequency	975.4 (1663)	879.7 (949.1)	1109.9 (1319.3)	966.3 (1462.2)	770.43 (1254.557)	5.6171	0.2296
N. syllables	3.48 (0.67)	3.56 (0.87)	3.8 (0.87)	3.8 (0.91)	3.69 (0.87)	6.8592	0.143
N. letters	8.76 (1.8)	8.32 (1.55)	8.72 (2.23)	9.2 (1.91)	8.8 (2.01)	2.7892	0.56

Extended Data Table 1-3A. 100 words referring to living/natural entities used to assess participants' perceptual threshold.

1.	acquazzone	51.	ippopotamo
2.	agnello	52.	lavanda
3.	agrifoglio	53.	libellula
4.	albero	54.	lucertola
5.	alligatore	55.	magnolia
6.	anatra	56.	maiale
7.	anguilla	57.	margherita
8.	aragosta	58.	marmotta
9.	armadillo	59.	montagna
10.	avvoltoio	60.	nebulosa
11.	balena	61.	oceano
12.	barboncino	62.	orchidea
13.	camaleonte	63.	ornitorinco
14.	cammello	64.	pantera
15.	campagna	65.	pappagallo
16.	canarino	66.	pecora
17.	canguro	67.	pellicano
18.	castoro	68.	pettirosso
19.	cavalletta	69.	pianeta
20.	cavallo	70.	pinguino
21.	cespuglio	71.	pioggia
22.	ciliegio	72.	pipistrello
23.	cinghiale	73.	procione
24.	civetta	74.	pterodattilo
25.	coccinella	75.	pulcino
26.	coccodrillo	76.	quadrifoglio
27.	colomba	77.	quercia
28.	coniglio	78.	rinoceronte
29.	cornacchia	79.	rosmarino
30.	costellazione	80.	salamandra
31.	criceto	81.	sanguisuga
32.	crisantemo	82.	scarabeo
33.	delfino	83.	scarafaggio
34.	deserto	84.	sciacallo
35.	dinosauro	85.	scimmia
36.	elefante	86.	scoiattolo
37.	farfalla	87.	serpente
38.	foresta	88.	spiaggia
39.	formica	89.	squalo
40.	furetto	90.	tacchino
41.	gabbiano	91.	tartaruga
42.	galassia	92.	tempesta
43.	gallina	93.	temporale
44.	gamberetto	94.	testuggine
45.	gelsomino	95.	tirannosauro
46.	girasole	96.	tricheco
47.	gorilla	97.	uccello
48.	granchio	98.	universo
49.	insalata	99.	usignolo
50.	insetto	100.	vitello

Extended Data Table 1-3B. Lexical and sublexical and psycholinguistic variables of the stimuli referring living/natural items used for assessing the participants' perceptual threshold, matched with manipulable objects (MO) and emotion (EM, with negatively valenced [EMneg], positively valenced [EMpos], and neutral [NEU]) word stimuli, mean (sd).

Variable	MO	EMneg	EMpos	NEU	Living	Kruskal-Wallis	*p* value
Lexical frequency	975.4 (1663)	879.7 (949.1)	1109.9 (1319.3)	966.3 (1462.2)	844.74 (1372)	5.2586	0.262
First bigram frequency	36399.6 (32386.9)	51354 (47659.77)	37908.56 (35402.45)	47136 (42814.64)	40578.68 (38356.7)	2.3387	0.6737
Last bigram frequency	70599.56 (49678.53)	76472.72 (55314.3)	75793.48 (60578.25)	79331.6 (40676.11)	72701.65 (48623.17)	2.3375	0.674
Medial bigrams frequency	74687.16 (22995.85)	67007 (25336.21)	75560.52 (21534.27)	81195.73 (20707.4)	71242.46 (23138.97)	6.4627	0.1672
Mean bigram frequency	60562.12 (20598.21)	64944.58 (27122.55)	63087.52 (29200.52)	69221.12 (23766.04)	61507.6 (23470.87)	2.6986	0.601
N. orthographic neighbours	1.51 (1.51)	1.32 (0.94)	1.36 (1.58)	0.92 (1.52)	1.4 (1.188)	7.7002	0.103
N. syllables	3.48 (0.67)	3.56 (0.87)	3.8 (0.87)	3.8 (0.91)	3.5 (0.76)	6.2544	0.181
N. letters	8.76 (1.8)	8.32 (1.55)	8.72 (2.23)	9.2 (1.91)	8.43 (1.55)	4.7944	0.309

Extended Data Figure 2-1.

Legend to Extended Data Figure 2-1. **Selection of participants for the BOLD analysis as a function of PAS**. The participants were ranked according to the Akaike information criterion (AIC; bottom right inset), obtained by Maximum Likelihood Estimation of the fit of the individual Word Contrast-to-PAS linear function (red dashed line; calculated on the average PAS ratings across all stimulus types (Empty, MO, EMneg, EMpos, NEU)) to a Word Contrast-to-PAS linear model function (gray line). The conventions of the dot-plots of all participants are as displayed for the "Participant020" on top left. None of the participants markedly departed from a linear increase of PAS score with increasing contrast level. However, "Participant012" was excluded from the analyses of BOLD activation as a function of PAS rating, because of an exceedingly low number of PAS{3,4} responses. This left the sub-sample for these analyses with 20 participants.