Title: Actin dynamics sustains spatial gradients of membrane tension in adherent cells.

Authors: Juan Manuel García-Arcos[1], Amine Mehidi[1,#], Julissa Sánchez Velázquez[2], Pau Guillamat[1,&], Caterina Tomba[1,¶], Laura Houzet[1], Laura Capolupo[3,†], Giovanni D'Angelo[3], Adai Colom[4,5], Elizabeth Hinde[2,6], Charlotte Aumeier[1], Aurélien Roux[1,*].

[1]Department of Biochemistry, University of Geneva, CH-1211 Geneva, Switzerland.
[2]School of Physics, University of Melbourne, Parkville, VIC 3010, Australia.
[3]School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland.
[4]Biofisika Institute (CSIC, UPV/EHU) and Department of Biochemistry and Molecular Biology, University of the Basque Country, ES-48940 Leioa, Spain.
[5]Ikerbasque, Basque Foundation for Science, ES-48013 Bilbao, Spain.
[6]Department of Biochemistry and Pharmacology, University of Melbourne, Parkville, VIC 3010, Australia.
[#]Present address: Interdisciplinary Institute for Neuroscience, Université de Bordeaux, UMR 5297, 33000 Bordeaux, France; Interdisciplinary Institute for Neuroscience, CNRS, UMR 5297, FR-33000 Bordeaux, France.
[&]Present address: Institute for Bioengineering of Catalonia, The Barcelona Institute for Science and Technology, ES-08028 Barcelona, Spain.
[¶]Present address: Univ Lyon, CNRS, INSA Lyon, Ecole Centrale de Lyon, Université Claude Bernard Lyon 1, CPE Lyon, INL, UMR5270, FR-69622 Villeurbanne, France.
[†]Present address: Friedrich Miescher Institute for Biomedical Research (FMI), CH-4056 Basel, Switzerland.
*Correspondence: aurelien.roux@unige.ch

Keywords: Membrane tension, cell mechanics, actin dynamics, actomyosin, cytoskeleton, cell migration, biophysics.

Abstract

Tension propagates extremely fast in lipid bilayers, precluding the formation of tension gradients. Nevertheless, plasma membrane tension gradients have been evidenced in migrating cells and along axons. Here, using a fluorescent membrane tension probe, we show that membrane tension gradients exist in all adherent cells, whether they migrate or not. Non-adhering cells do not display tension gradients. We further show that branched actin increases tension, while membrane-to-cortex attachments facilitate its propagation. Tension is the lowest at the edge of adhesion sites and highest at protrusions, setting the boundaries of the tension gradients. By providing a quantitative and mechanistic basis behind the organization of membrane tension gradients, our work explains how they are actively sustained in adherent cells.

Introduction

Cells and organelles are separated from their environment by lipid membranes, which are self-healing fluid films,[1] conferring them with unique mechanical properties. The main modulator of membrane mechanics is membrane tension, defined as the stress resulting from changing the apparent surface area of the membrane.[2] In cells, plasma membrane tension coordinates endo- and exocytosis rates both temporally and spatially,[3–6] determines cell shape during single-cell migration,[7–13] and collective migration,[14] and regulates membrane signaling,[15,16] actin polymerization,[17] axon growth,[18,19] and cell spreading.[20,21]

Among the open questions about cellular membrane mechanics, the propagation of membrane tension in the plasma membrane is still under debate.[22] Simply put, if membrane tension propagates too fast, tension gradients cannot arise. In pure lipid membranes (i.e.: giant unilamellar vesicles, GUVs), because of their fluidity, tension is expected to propagate at the speed of sound waves[23] (>0.1 m/s), causing equilibration within tens of milliseconds and preventing observable tension gradients upon local perturbations.[24] However, membrane tension gradients have been observed in supported lipid bilayers adhering to a solid substrate during expansion, which dramatically reduces the diffusion of lipids.[1,25] In cells, early studies of the membrane tension of axons showed that the membrane flowed from the growth cone to the soma driven by a membrane tension gradient.[3] While this gradient could be specific to the unique strong membrane-cytoskeleton interactions of axons,[26–29] more recent studies have established that similar gradients can spread from the front to the rear of migrating cells.[8,12] During cell migration, tension propagation is considered to be fast (< 1s across the cell) and long-ranged (at the cellular scale) and was proposed to act as a global signal transducer.[30] But the maintenance of the gradient in this case is proposed to be linked to the high dynamics of actin forces. The current understanding is that the propagation of tensile stresses throughout the cell membrane depends on the cell type and the kind of perturbation, but that at rest, no cell exhibits tension gradients.[19,31–35] In analogy to *in vitro* experiments, these divergences might originate from strong mechanical interactions between the membrane and the underlying actomyosin cortex or the substrate.[31,36,37] However, whether these gradients can be generated in every cell is unknown, and a formal quantitative assessment is required to establish how they are formed.

Another layer of complexity in cellular membranes is their lateral separation into nanodomains.[38–43] The lateral heterogeneity of cell membranes stems from a complex lipid composition constantly stirred by the dynamics of the underlying actomyosin cortex.[44,45] Domain formation depends on temperature and lipid composition, but mechanical factors can play a role as well: studies *in vitro* have shown that, around a critical point,[46] adhesion[47] or increasing membrane tension by hypotonic shocks drives lateral phase separation.[48–53] Interestingly, in model membranes far from critical points, increasing tension leads to an increased lipid spacing (less packing), as reported by Laurdan[54,55] or Flipper-TR probes.[24] In cells, results suggest membranes are near a critical transition point:[46] increasing plasma membrane tension by hypotonic shocks globally promotes domain formation[56] and increased lipid packing reported with Flipper-TR[24] and Laurdan membrane probes.[57] This link between membrane mechanics and molecular organization allows for proxy measurements of membrane tension using lipid packing-sensitive probes such as Flipper-TR, by determining the probe conformation by fluorescence lifetime imaging microscopy (FLIM).[24,58] Importantly, the relation between

lipid packing and tension does not depend on whether the experiments are performed with
model membranes or in cells, but on the critical state of the membrane.

Membrane tension gradients were evidenced by differences in the force required to hold
plasma membrane tubes (tether force) pulled from different parts of the cell. Tether force
includes contributions not only from plasma membrane tension but also from membrane-
cytoskeleton interactions and bilayer bending rigidity,[37,59–63] which is composition-
dependent.[64] Therefore it remains unclear to what extent the reported changes in tether
force stem from actual membrane tension gradients, from changes in membrane-cortex
binding energy, or from lipid composition changes. Moreover, limited by discrete spatial
measurements of the dorsal cellular plasma membrane, tube pulling cannot provide
insights into the shape of the gradients nor investigate the tension distributions of the
ventral cell membrane in contact with the substrate. To visualize membrane tension
gradients and investigate the mechanisms that could sustain them, we used the Flipper-
TR probe.

Results

Flipper-TR reports membrane tension gradients in reconstituted membranes.

To test whether Flipper-TR can report membrane tension gradients, we aimed at
reconstituting tension gradients in model lipid membranes *in vitro*, consisting of expanding
glass-supported supported lipid bilayers (SLB) with well-controlled lipid composition and
stable tension gradients.[1,25] In brief, lipids were dried over large silica beads acting as lipid
carriers and placed over plasma-cleaned glass. Upon hydration, lipids wet the clean glass
and expand from the bead outwards as a single bilayer. SLB expansion is counteracted
by the friction of the membrane onto the glass, allowing the formation of a tension gradient
(Figure 1A).[1] SLBs expand rapidly at first and slow down following a power-law while the
tension gradient relaxes (Figure S1A),[1,25] allowing us to sample a range of tension
gradients (see Methods).

We performed confocal FLIM of the SLBs with Flipper-TR using 5 different lipid
compositions   containing   Dioleoylphosphatidylcholine/serine   (DOPC/S),   brain
sphingomyelin (bSM), and cholesterol (Chol): DOPC:DOPS 60:40; DOPC:DOPS:Chol
42:28:30; DOPC:bSM:Chol 1:1:1, DOPC:bSM:Chol 1:1:2, and DOPC:bSM:Chol 2:2:1.
Flipper-TR lifetime distributions qualitatively showed pronounced spatial gradients (Figure
1B), particularly evident in fast SLBs spreading at velocities comparable to cell migration
(>5µm/min) (Supplementary Movie 2). As expected, more ordered membranes led to
higher Flipper-TR lifetimes (Figure 1C). Interestingly, single-phase compositions display a
lower average Flipper-TR lifetime at the front (leading edge) than the rear, closer to the
lipid source, while some compositions containing phase-separating lipids (bSM and Chol)
display an inverted lifetime gradient (Figure 1D). Thus, considering the gradient from the
inner to the outer part of the membrane patch, compositions around a critical point display
a positive slope while others display a negative slope.

Importantly, only bilayers that are expanding – thereby having a tension gradient – display
a lifetime gradient (Figure S1A). To better understand the interplay between Flipper-TR
lifetime and membrane tension, we plotted the slope of gradients as a function of the SLB
spreading speed. For long time points, the speed reduces to zero as well as the slope
value, confirming that when SLBs stop spreading and dissipate their tension gradient, the
Flipper-TR lifetime gradient also vanishes (Supplementary Movie 2). For SLBs with

compositions that cannot phase-separate, the slope of the Flipper-TR lifetime gradient (see Methods) increased from negative values to zero roughly proportionally to expansion speed (Figure 1E, DOPC:DOPS 60:40 and DOPC:bSM:Chol 2:2:1). For SLBs with compositions close to phase separation, the slope decreased instead from positive values to zero, also roughly proportionally to expansion speed (Figure 1E, DOPC:bSM:Chol 1:1:1 and DOPC:bSM:Chol 1:1:2). This effect is robust across a wide range of spreading speeds and lifetime gradients (Figure S1A).

Raedler and colleagues described that tracer fluorescent molecules display an exponentially decaying concentration profile in spreading SLBs with a tension gradient[1,25]. The decay length depends on the bulkiness of the tracer relative to the other lipids of the mix, probes with bulky headgroups such as DOPE-atto647 localize at areas with lower lipid packing, a visual evidence of membrane tension gradients.[1] DOPE-atto647 localized to the edge of the spreading SLBs in single-phase lipid composition, but it was excluded from the edge of SLBs composed of phase-separating composition (Figure 1SB). This confirmed that in spreading SLBs under tension made of phase-separating compositions, a more ordered phase was colocalized with areas of higher tension.

Flipper-TR distribution also displayed fluorescence intensity profiles, inverted compared to the DOPE-A647 profiles (Figure 1SB). It is however more difficult to interpret these profiles in terms of concentration since the quantum efficiency of Flipper-TR varies with lipid order.[65] However, the higher Flipper-TR fluorescence intensity at the edge of SLBs with phase-separating compositions is consistent with the presence of more ordered domains in these regions. Overall, these data showed the correspondence between tension and lipid packing and confirmed that Flipper-TR can report tension gradients in model membranes.

Leading-edge extension increases Flipper-TR lifetime in migrating cells

Membrane tension gradients in migrating cells were first characterized in the fish keratocyte.[12] Fish keratocytes have a very characteristic crescent shape and perform fast and persistent migration driven by a large lamellipodium at the leading edge.[10] We investigated whether we could visualize such tension differences during keratocyte migration using Flipper-TR. Keratocytes were labeled with Flipper-TR following previously reported procedures[66] and their ventral plasma membrane was imaged using confocal FLIM (see Methods). Our results confirmed that cells robustly display large lifetime differences between the lamellipodium leading edge and the cell rear (Figure 2A): $\langle \text{Lifetime}_{front} \rangle = 4.85 \pm 0.13$ns; $\langle \text{Lifetime}_{rear} \rangle = 4.65 \pm 0.16$ns; $\langle \Delta \text{Lifetime} \rangle = 0.20 \pm 0.12$ns. This is consistent with the literature, reporting higher lipid packing and higher tension values at the front of migrating cells.[12] At the level of single cells, front and rear values fluctuate over time, but the single-cell time average displays a robust gradient in Flipper-TR lifetime (Figure 2B).

To extend these observations, different cell lines were selected to cover a wide range of migratory behaviors: U2OS and RPE1, which migrate efficiently, and HeLa and Cos7 which appear slow and less persistent (Figure S2A-B). Qualitatively, as for keratocytes, the cells showed large heterogeneities of Flipper-TR lifetime (Figure 2C), being higher at cell edges and decaying toward the cell center. Contrary to keratocytes, these cell lines do not maintain shape during migration, allowing us to manually sort cells into polarized and non-polarized categories based on their shape. While cells displaying a highly polarized shape showed clear differences between front and rear spatial decays (Figure 2C-D),

non-polarized cells did not (Figure S2C-D). Lifetime values decayed linearly over more
than 10 microns from the cell leading edge, while they decayed exponentially within a few
microns to the baseline levels from the rear (Figure 2D-F). Lifetime differences between
front and rear were more pronounced in cells with elongated morphologies (Figure 2F).
Overall, the cell lines that displayed a more persistent migration (Figure S2A-B) were the
ones that displayed the most robust Flipper-TR gradients. Interestingly, U2OS cells
displayed overall lower lifetime values than other cell types but had the same gradient as
the other persistent cell type RPE1. Since plasma membrane lipidomes are cell-type
specific, offsets in the average lifetime may be due to a specific lipid composition in this
cell line.[67]

We next investigated the mechanism leading to the observed differences in Flipper-TR
lifetime. As the actin cortex shapes migrating cells, we wondered how the actin
cytoskeleton would colocalize with Flipper-TR lifetime gradients. For this, following
Flipper-TR lifetime imaging, U2OS cells were fixed and labeled for actin and vinculin (see
Methods). In polarized cells, actin and vinculin had a stereotypical localization pattern
composed of a dense, narrow actin strip marking the lamellipodium edge and a less
dense actin region with small elongated focal adhesions corresponding to the lamella. At
the rear, we observed low cortical actin and large stress fibers originating from large
mature focal adhesions (Figure 2G). Regions of highest and lowest lifetime coincided with
characteristic features in the actin organization. Flipper-TR lifetime increased both in the
lamellipodium and lamella regions (Figure 2G, Flipper-TR panel, and Figure 2H). We used
actin images to train a machine learning algorithm using Ilastik[68] to segment cells into
lamellipodium, lamella, and cell body regions, and calculate their average Flipper-TR
lifetime values. This confirmed that the differences in actin cytoskeleton morphology led to
different Flipper-TR lifetimes.

Actin polymerization extends the leading edge of lamellipodium and thereby could
increase plasma membrane tension.[13,20,32,69] To study this, we performed live FLIM on
RPE1 and U2OS cells, to track cell edge dynamics simultaneously with Flipper-TR
lifetime. FLIM time-lapses on migrating cells confirmed that, in the event of cell
repolarization, the region of high lifetime dynamically changes localization to the new
leading edge (Supplementary Movie 1). For every point of the cell edge, the gradient of
Flipper-TR lifetime (from the cell edge inwards) and the edge velocity were measured in
RPE1 and U2OS cells. Plotting the spatial lifetime gradient binned by edge velocity
revealed that the protrusion speed is positively correlated with increased Flipper-TR
lifetime (Figure 2J). The protruding parts of the cell display higher Flipper-TR lifetime
values than the retracting ones (Figure 2K): $\langle$U2OS Lifetime$_{protrusion}\rangle$ = 4.63±0.16ns; $\langle$U2OS
Lifetime$_{retraction}\rangle$ = 4.46±0.21ns; $\langle$RPE1 Lifetime$_{protrusion}\rangle$ = 4.76±0.36ns; $\langle$RPE1
Lifetime$_{retraction}\rangle$ = 4.47±0.22ns. These results are consistent with our hypothesis, thereby
confirming that protrusions in migratory cells increase Flipper-TR lifetime.

Since the lowest lifetimes often correspond to the central region of the cells, coinciding
with the nucleus position, we tested whether the presence of the nucleus could influence
the reported plasma membrane Flipper-TR lifetime gradients. To test this, we
characterized the lifetime of the ventral membrane of enucleated cells (Methods).
Enucleated U2OS and RPE1 cells maintained similar Flipper-TR lifetime distributions
(Figure S2E-F) supporting that the nucleus is not generating lifetime gradients within the
plasma membrane.

Membrane tension gradients are present in non-migrating cells

After confirming that Flipper-TR can reveal membrane tension gradients in expanding membranes both in migrating cells and *in vitro*, we tested if plasma tension gradients also exist in non-migrating cells. To this end, we cultured single HeLa cells on fibronectin-coated micropatterns[71,72] and labeled them with Flipper-TR. Controlling cell shape allowed us to extract average lifetime maps from large sets of cells with the same shape, thus with comparable spatial organization of intracellular components, including the nuclei, the actin cytoskeleton, and the focal adhesions (Figure S3A). Flipper-TR lifetime showed a heterogeneous spatial distribution at the ventral plasma membrane (Figure 3A), as previously observed[5], while its distribution appeared homogeneous on the dorsal membrane (Figure 3B). Despite no significant differences were observed in the average Flipper-TR lifetimes between different cell shapes, neither on the ventral (Figure 3A) nor dorsal planes (Figure 3B), we found a significant shift in the average Flipper-TR lifetime value between the dorsal and ventral planes (Figure 3C, S3B): $\langle Lifetime_{dorsal} \rangle$ = 4.98±0.05ns; $\langle Lifetime_{ventral} \rangle$ = 4.71±0.06ns; such as $\langle \Delta Lifetime \rangle$ = 0.26±0.07ns.

Consistent with the previous results, cellular membrane protrusions on the non-adherent areas exhibited higher lifetime values (Figure 3C). The average lifetime at the ventral membrane was comparable between adhesive and non-adhesive regions (Figure 3C). The spatial probability maps of Flipper-TR lifetime (Figure 3E) suggested that, while the gradients along the cell contour or across the cell diameter differ between shapes (Figure 3F), the average Flipper-TR lifetime increases at the cell edges and decreases over the boundaries of the adhesive micropatterns, regardless of shape. The lateral lifetime gradients between the regions of high and low lifetime were: $\langle Lifetime_{high} \rangle$ = 4.85±0.13ns; $\langle Lifetime_{low} \rangle$ = 4.67±0.13ns; such as $\langle \Delta Lifetime \rangle$ = 0.19±0.09ns. We further analyzed this by assigning two spatial descriptors to each lifetime value, namely the shortest distance cell edge and the shortest distance to the micropattern boundary. Note that negative distances account for pixels inside the cell or the micropattern. By plotting the average lifetime as a function of these two distances we can compare the lifetime gradients we obtained from the three different pattern shapes (Figure 3D). The fact that the gradients exhibited the same distribution regardless of the shape confirmed that the positions of the cell edge and the micropattern boundary are the main determinants of the Flipper-TR lifetime distribution (Figure 3G-H).

Epithelial RPE1 cells displayed similar spatial lifetime gradients decreasing from the cell edges to the cell center. Dorsal to ventral lifetime shifts were also conserved (Figure S3C-D). However, the increase of lifetime at plasma membrane regions over non-adhesive regions seems to be cell-line specific. This could be attributed to protrusive activity and ruffle formation over non-adhesive areas in HeLa (Supplementary movie 3), not present in RPE1. This was confirmed by visualizing average membrane tension gradients on different radial designs for HeLa cells, including full discs and double-ring patterns (Figure S3E). Cells on wider ring pattern lifetime decrease at the inner boundary of the pattern (Figure S3E). In full disc patterns, lifetime was distributed homogeneously (Figure S3E). Importantly, in all cases, Flipper-TR lifetime increased over non-adhesive regions in HeLa.

Despite the standardization of cell shape, cells still exhibit significant differences in vinculin and actin organization leading to two main protrusive behaviors: "low protrusion" cells display a very dense vinculin recruitment on focal adhesions and thick stress fibers and no protrusions; "high protrusion" cells display homogeneous vinculin localization and

271 no stress fibers, with numerous protrusions projecting outwards (Figure S4A). By plotting
272 the lifetime gradients of the 10% cells with the highest protrusion area and the 10% cells
273 with the lowest we confirmed that cells with more protrusive behavior had higher lifetime
274 values overall, suggesting, as in migrating cells, that actin protrusions play a fundamental
275 role in the tension increase at the cells' edges (Figure S4B).

Main lipid components are uniformly distributed in patterned cells

While our *in vitro* data showed that Flipper-TR can report membrane tension gradients in
reconstituted membranes, we wondered whether the lifetime gradients we observed in
cells had the same origin or if they could originate from lipid composition variations. To
investigate this, we studied the spatial distribution of lipid composition by using Matrix-
Assisted Laser Desorption Ionization Mass Spectrometry Imaging (MALDI-MSI). This
technique allows the mass-spectrometry analysis of the chemical content of each pixel in
an image providing a detailed analysis of most lipid species with a subcellular resolution.[73]
The subcellular resolution is achieved by averaging over many images, which was made
possible using standardized cell shapes (Figure S3F and Methods).

We measured the fraction of different lipid species in cross-shaped micropatterned cells:
namely phosphatidylcholine (PC), phosphatidylethanolamine (PE), ceramide (Cer),
sphingomyelin (SM), and globotriaosylceramide (Gb3) (Figure S3G). From our results on
model membranes, we expect that composition gradients can be linked to tension
gradients only in lipids in trace amounts with extreme bulkiness (Figure S1B). Consistent
with our hypothesis, lipid species that are abundant in these membranes (PC, PE) display
homogeneous distributions while less abundant lipid species (Cer, SM, and Gb3) display
heterogeneous spatial distributions in the form of lipid concentration gradients (Figure
S3H). The spatial gradient of Gb3, which accumulates at the cell membrane protrusions is
consistent with the tension gradient previously reported (Figure 3E), as tension is maximal
at the cell edges (Figure S3H). Interestingly, the ceramide and sphingomyelin distribution
showed a moderate preference for focal adhesion sites which do not correlate with an
increase in Flipper-TR lifetime (Figure S3H). This supports that the Flipper-TR lifetime
differences are not due to differences in lipid composition, as we could expect
sphingolipids to increase membrane order. To further explore the spatial distribution of
lipid composition throughout the plasma membrane, we used fluorescent dyes, Filipin,
Cholera toxin, Equina toxin, and Shiga toxin that specifically bind to cholesterol, GM1,
sphingomyelin, and Gb3 lipids, respectively (Figure S3I). Qualitatively, the concentration
of all these lipid species tends to increase at the vertices of the cells on cross-shaped
patterns, but the lipid packing effects for sphingolipids associated with lipid raft[74] cannot
explain the spatial variations observed in the average lifetime maps. Overall, these
observations support that gradients reported by Flipper-TR lifetime are due to membrane
tension gradients and not lipid composition.

Actin polymerization and adhesions scale tension gradients

To test the role of actomyosin cytoskeleton in setting membrane tension gradients, we
characterized the effect of several pharmacological treatments on the Flipper-TR lifetime
gradients. We used 1µM Latrunculin A ('LatA') for inhibiting actin polymerization, 85 µM
CK-666 for inhibiting Arp2/3-mediated actin nucleation, 10µM NSC668394 ('NSC66') for
inhibiting ezrin – a major actin-membrane linker protein – and 20µM Y-27632 for inhibiting
Rho kinase, responsible for myosin contractility. Moreover, we used the JLY cocktail
consisting of 20µM Y-27632, 5µM Latrunculin B, and 8µM Jasplakinolide to block actin

dynamics[75]. For each treatment and each shape, we quantified: i) the spatial lifetime
distribution (Figure 4A top panels), ii) the radial lifetime distribution for the ring pattern
(Figure 4A, bottom panels), iii) the average lifetime per cell (Figure 4B), and iv) the
maximal amplitude of the gradients (Figure 4C).

Both LatA and JLY treatments inhibited actin dynamics. As in control experiments, we
observed a significant shift in the average lifetime between the ventral and dorsal
membranes (JLY on Figure S4E). However, the increase in lifetime in non-adhesive areas
and the pattern boundary effect were abolished. As a result, membrane tension became
more homogeneously distributed at the ventral membrane, remaining high at the edges
(Figure 4A and 4C). The linear decay of the Flipper-TR lifetime at the edge became
exponential, decaying quickly towards the center. Therefore, LatA and JLY transformed
front-like tension gradients to rear-like tension gradients through actin depolymerization.
This suggests that the linear gradients observed in the lamellipodial regions of migrating
cells (Figure 2) depend on the actin organization at the lamellipodium. F-actin was almost
absent in both these conditions, although some focal adhesions remained in place (Figure
4D). These results confirm that actin dynamics is essential to generate membrane tension
gradients.

In contrast with the LatA or JLY treatments, inhibiting the Arp2/3 complex with CK-666 did
not decrease F-actin contents. Arp2/3 complexes nucleate about half of the cortical F-
actin content in HeLa cells in the shape of branched networks.[76] As expected, Arp2/3
inhibition led to the formation of more filopodia, larger lamellipodia, and thicker focal
adhesions and stress fibers (Figure 4D). Surprisingly, the inhibition of Arp2/3 reduced
considerably the average Flipper-TR lifetime value (Figure 4B). It nevertheless maintained
both the higher values at the cell edges (Figure 4A and 4C) and the shift between the
ventral and dorsal planes (Figure S4E). The lifetime drastically decreased at the lamellar
regions, resulting in a steeper gradient than in control conditions (Figure 4A). This
suggests that, in control conditions, lamellipodial extension – promoted by Arp2/3
branching – is generating higher membrane tension, and/or reducing the diffusion of
lipids, setting a longer characteristic decay length of the gradient. The role of branched
actin networks was also supported by Rho kinase inhibitor Y-27632 treatment, which
significantly extended lamellipodia (Figure 4D). This resulted in a global increase in the
average lifetime (Figure 4B), particularly in the numerous lamellipodia extending outwards
(Figure 4A), and reduced the amplitude of the lifetime gradients (Figure 4C).

Finally, to inhibit membrane-to-cortex attachment, we used NSC668394, which inhibits
ezrin activity by preventing its T567 phosphorylation, a member of the ERM protein family
that acts as a major membrane-cortex linker.[77] Cells under ezrin inhibition displayed
similar protrusive morphology as the control cells but displayed smaller focal adhesions
with less vinculin (Figure 4D). This resulted in a similar lifetime in ruffled areas, but a
larger decay of lifetime in adhesive areas (Figure 4C). This indicates that with less
membrane-to-cortex attachment can increase membrane tension at protrusions, but this
tension can dissipate faster away from protrusive regions. We, therefore, concluded that
actin-membrane attachment may be involved in reducing the local diffusion of membrane
components, thereby reducing tension propagation and gradients. Overall, these results
confirm that tension generation is controlled by the dynamics of the actin cytoskeleton,
specifically of branched actin, via membrane-to-cortex attachment.

Lipid diffusion and flows are limited in patterned cells

Previous studies have shown that gradients in membrane tension are associated with lipid
flows,[18] with lipids flowing from low to higher tension regions of neuronal growth cones.
On the other hand, other studies on migrating cells have shown either no flow[78] or
rearward flow[79] in the cell reference frame. We therefore wanted to investigate lipid
diffusion and flows in our system. Fluorescence Recovery After Photobleaching (FRAP)
assays revealed no significant difference in the recovery times across different cell
regions at the ventral membrane (Figure S5A, note that FRAP $t_{1/2}$ does not change on the
adhesive and non-adhesive areas of the edges). We further tested the presence of flows
using FRAP, bleaching a line tangential to the cell edge and tracking if the fluorescence
minima moved during recovery. These assays showed that intensity minima moved
towards the cell edge (Figure S5B). However, this apparent lipid flow did not change in the
absence of actin dynamics (Figure S5B, JLY treatment). FRAP data can be interpreted as
resulting from either a ventral plasma membrane flow directed towards the cell edges that
is non-dependent on actin dynamics, or from the existence of a diffusion barrier at the
leading edge. While the presence of a diffusion barrier at the leading edge was proposed
by previous studies,[80] we aimed at discriminating between flows and diffusion barriers with
other means than bleaching.

For this, we employed fluorescence fluctuation spectroscopy (FFS) and spatiotemporally
investigated lipid dynamics without photobleaching by concomitant autocorrelation
function (ACF) and pair correlation function (pCF) analysis across line scan FFS
measurements.[81–83] In particular, single-channel line scans across the ventral membrane
of patterned cells stained with the plasma membrane dye CellMask were acquired at a
rate faster than the rate at which the membrane dye molecules diffuse (Figure S5C-D).
Then the local versus long range mobility of the diffusing dye molecules across the
membrane was extracted by an ACF and pCF analysis of the acquired fluorescence
fluctuations with themselves versus at specific distances in the forward or reverse
direction (Figure S5E, see Methods). We positioned the line scans crossing the pattern
boundaries, thus across regions with a sharp decrease of the Flipper-TR fluorescence
lifetime (Figure S5G). The ACF analysis showed that lipid mobility does not change
between the adhesive and non-adhesive areas (Figure S5G-H). The diffusion values are
consistent with point FCS measurements of the same lipid probe from the literature ($\langle D_{cross}$
$_{control}\rangle$ = 2.2±2.7µm^2/s; $\langle D_{ring\ control}\rangle$ = 1.9±2.6µm^2/s)[84]. Under JLY treatment, we noticed a
weak trend decreasing overall diffusion and transport ($\langle D_{cross\ JLY}\rangle$ = 1.7±2.0µm^2/s; $\langle D_{ring\ JLY}\rangle$
= 1.5±2.0µm^2/s). The pCF analysis revealed an absence of membrane lipid flow (Figure
S5G, no significant differences between forward and retrograde transport). From these
data, we concluded that patterned cells do not display a steady-state lipid flow at their
ventral membrane, despite the presence of membrane tension gradients. This is
consistent with the fact that membrane tension gradients are maintained through time, as
they do not dissipate by lipid flows.

402   *Substrate rigidity and adhesiveness modulate membrane tension*

Our results in patterned cells indicate that tension gradients can exist in non-migrating
cells due to adhesion patterns and actin polymerization. Therefore, this proves that
tension gradients are not exclusive to migrating cells. We next wondered if they were
exclusive to adherent cells. To address this, we examined whether membrane tension
gradients are present in non-adherent migrating cells which usually migrate through bleb
expansion. We confined HeLa cells to trigger leader bleb migration—a mode of migration

independent of specific adhesion, but where a gradient of cortical actin orients blebbing at
the leading edge of the cell.[85,86] Cells were confined for over an hour to ensure the
predominance of the leader bleb phenotype (Figure 5A).[87] Our findings indicated that
Flipper-TR lifetime gradients are absent at the protruding front of leader bleb migrating
cells (Figures 5B and 5C). The lifetime was only marginally increased at the rear of the
leader bleb, adjacent to the uropod (i.e. the adhesive rear of the cell training behind the
leader bleb). Nevertheless, the magnitude of these lifetime changes was negligible
compared to the gradients observed in adherent cells. This shows that adhesion is
necessary to create a tension gradient, suggesting that tension gradients are not essential
for migration.

Since adhesion is essential to create tension gradients in the plasma membrane, we
hypothesized that the modulation of adhesion strength could change those gradients.
Specifically, we wondered if adhesion could be responsible for the differences in lifetime
observed between the bottom and middle planes of the cells (Figures S3B-C and S4E).
This difference was persistent despite various pharmacological treatments perturbing the
actin cytoskeleton, indicating that it was not driven by actin dynamics. Considering leader
bleb cells exhibit the same shift between bottom and middle plane lifetimes as cells
adhering to fibronectin-coated glass, we assume the ventral-to-dorsal shift in average
lifetime does not depend on specific molecular interactions. As substrate stiffness is
known to strongly affect the adhesion strength of cells,[88] we plated HeLa cells on
substrates of different stiffnesses (Figure 5D) and measured the average Flipper-TR
lifetime at the ventral and dorsal membranes, i.e., bottom and middle planes (Figure 5E).
Interestingly, cells plated on fibronectin-coated soft polyacrylamide (PAA) hydrogels (E =
0.5 and 4.0 kPa, Methods), did not exhibit any differences in the average Flipper-TR
lifetime between middle and bottom planes (Figure 5E). This was due to the lifetime at the
bottom membrane of cells increasing on soft substrates with respect to the bottom
membranes of cells plated on glass. Moreover, we observed an attenuation of lateral
lifetime gradients on soft hydrogels. These data confirmed not only that adhesion per se is
essential for the generation of tension gradients, but also that adhesion strength, which
scales with substrate stiffness,[88] is also critical.

To further test the role of adhesion strength in the generation and sustaining of plasma
membrane tension gradients, we next aimed at pharmacologically reducing adhesion.
Because tension gradients did not correlate with focal adhesions, and because removing
focal adhesions may lead to complete detachment of cells, we investigated other adhesive
structures. Clathrin plaques, distinct from transient clathrin-coated pits, are adhesive
structures that rely on αvβ5 integrin-ECM interactions and are present in some cell lines
when bound to high-rigidity substrates.[89] We patterned HeLa cells transiently expressing
clathrin light chain fused with GFP and quantified the localization of the stable clathrin
population using TIRF microscopy. Clathrin plaques colocalized with low-lifetime zones,
appearing at the inner edge and at the edges of the cross pattern (Figure 5F). Treatment
with cilengitide, which inhibits αvβ5-integrin-mediated adhesions, resulted in an increase
of Flipper-TR lifetime at the areas surrounding the patterns (Figure 5G), like the effect of
plating cells on low-rigidity substrates. Thus, reducing adhesion strength by releasing
molecular bounds flattens membrane tension gradients.

Discussion

By using Flipper-TR as a mechano-responsive molecular probe, here we established how branched actin networks and cellular adhesions play a critical role in generating and sustaining spatial tension gradients within the ventral plasma membrane of adherent cells. Also, employing *in vitro* experiments with model membranes, we could establish the exact nature of the coupling between Flipper-TR lifetime and membrane tension. Therefore, this work validates the use of Flipper-TR as an indicator of membrane tension and shows the influence of lipid composition on the relation between tension and lipid packing. In particular, we describe how branched actin networks and membrane-to-cortex attachment are essential for tension buildup and tension propagation, respectively.

The differences between the upper and lower values of Flipper-TR lifetime gradients are in the range of 0.20 ns. From the calibration between flipper-TR lifetime and effective tension in HeLa,[24,90] we can evaluate $\Delta\sigma_{eff}$ to be in the range of 0.25 mN.m^{-1}. Therefore, we estimate the tension to drop to be about 30-40% from the resting σ_{eff} of 0.1-0.6 mN.m^{-1} found in all cell types studied, which is far from being a negligible tension gradient. Within the same range, an increase of 30% higher at the leading edge compared to the trailing edge was previously measured in fish keratocytes by other means.[12]

Some studies proposed that membrane tension propagates very fast, therefore homogenizing tension values,[32] while others have proposed that tension gradients can be maintained in polarized cells.[12,91] Our work reconciles both ideas by proposing that tension propagation depends on the degree of membrane-substrate interaction. Therefore, contrary to what has been suggested,[35] the existence of tension gradients does not depend on the cell type, but on the migration mode.[87] At the ventral membrane, attached to the coverslip, we find strong inhomogeneities, which disappear at the apical membrane not attached to the substrate (Figure 5G, top panel). This allows us to propose a mechanism by which membrane tension gradients can be sustained in adherent cells (Figure 5G). Dynamic extension of the membrane by actomyosin turnover at the cell edge locally increases the tension. The propagation of tension towards the cell center is hindered by adhesion to the substrate and between the membrane and the actin cortex. The constant pulling force at protrusion sites combined with the decrease of tension at adhesions set the shape of the tension gradient. In the following, we discuss the consistency of this model with previous literature.

We propose that cells must adhere to generate and sustain tension gradients. A remaining question is which are the molecular principles at play to limit lipid diffusion. While rapid lipid diffusion can relax tension gradients, tension could be relaxed by other mechanisms, such as membrane unfolding or osmotic pressure release.[21,90] Therefore, a reduction of lipid diffusion is not sufficient to create tension gradients, and assuming that hindered propagation of membrane flows is a signature of tension gradient may be fallacious.[34] Also, lipid diffusion depends on which scale it is measured. At the nanometer scale, no major difference in mobility is observed in lipid diffusion[92] between the ventral and dorsal membrane. At the submicron scale, lipid diffusion may be constrained by 'corrals' created by the mesh of cortical actin,[93] by transient actin asters driven by Arp2/3[94,95], or by lipid nanodomains. The two last modes are very dynamic, so their contribution to the long-range flow barrier is less obvious. But it is really at the cell scale that lipid flows may counteract or create membrane tension gradients. This may explain discrepancies found in the literature on how lipid flows and membrane tension gradients

are coupled, as the mobility of lipids is usually investigated at a molecular scale, while gradients are always evidenced at the cellular scale.

Regarding the role of actin, our data shows that branched actin both increases membrane tension at the leading edge and plays a role in its propagation. There is extensive evidence linking branched actin networks to membrane tension increase, often via ERM proteins.[13,21,32,35] What molecular mechanism mediates branched actin tension increase? On one hand, the branched actin nucleator Arp2/3 must bind to membrane-bound WAVE to add new filaments,[96] linking actin polymerization and membrane binding. However, the specific properties of branched actin networks that promote tension increase might instead originate from their architecture, which confers them the ability to adapt density and pushing angle to the load to maintain polymerization.[17,97,98]

Overall, our study demonstrates that membrane tension gradients in adherent cells arise from the interplay between actin dynamics, lipid diffusion/flow constraints, and cell-to-substrate adhesion. By visualizing tension gradients, our work links the physical properties of the plasma membrane with cytoskeletal dynamics and understands how cell membrane tension gradients are formed, opening new avenues for investigating their role in cellular processes such as receptor signaling, phagocytosis, cell polarity or cell differentiation. For example, we speculate that membrane partitioning and self-organized criticality may allow lipid organization to respond to small perturbations like tension changes,[46] impacting allosteric interactions at the membrane and cell polarity signaling.

Limitations of the study

The limitations of our study stem from the use of fluorescence lifetime imaging microscopy (FLIM) on a scanning confocal setup. This technique is slow compared to the speed of actin dynamics, and especially to membrane diffusion (the acquisition time of single cells was ≈ 30s), which means that images of single cells are in fact averaged over times during which both actin and lipids move. Other FLIM techniques such as gated and modulated CCD image sensors might enable one to go faster and obtain images with higher time resolution.[99] While our results from MALDI-MSI on micropattern cells agree with lipid distributions expected from membrane tension gradients *in vitro*, it is important to note that MALDI-MSI requires fixation and quantifies lipids from all membranes, not just plasma membrane, factors that may interfere with the experimental interpretation. These factors may limit the use of MALDI-MSI to study lipid compositional gradients in live cells.

Acknowledgments

We thank Javier Espadas for training on supported lipid bilayers and useful feedback, Martin Lenz for discussions on membrane flows, Julie Miesch and Anne-Laure Boinet for TIRF microscope training, and Vincent Mercier and Chloé Roffay for FLIM confocal microscope training. Thanks to Rafael Caetano, Frédéric Humbert, and Roux lab members for technical assistance and feedback. Special thanks to Mireia Andreu Calvo, Guillaume Pernollet, Vincent Mercier, and the ACCESS platform for micropatterning training. We thank Henry De Belly, Matthieu Piel, Mathieu Dedenon, and Larisa Venkova for critical reading. A.R. acknowledges funding from the Swiss National Fund for Research grant numbers #CRSII5_189996 and #310030_200793 and the European Research Council Synergy grant number #951324-R2-TENSION. A.M. acknowledges funding from the Human Frontiers of Science Program (grant number LT000762/2020-L). P.G. acknowledges support from the Human Frontiers of Science Program (grant number LT-

000793/2018-C). C.A. acknowledges funding from the DIP of the Canton of Geneva,
SNSF (31003A_182473 and TMSGI3_211433).

Author contributions

Conceptualization, J.M.G.A., A.R.; Methodology, J.M.G.A., A.R; Software, J.M.G.A., A.M.,
E.H.; Formal Analysis, J.M.G.A., A.M., E.H., J.S.; Investigation, J.M.G.A., P.G., C.T. A.C.,
A.M.; Resources, ; Data Curation, J.M.G.A.; Writing – Original Draft, J.M.G.A., A.R.;
Writing – Review & Editing, J.M.G.A., A.R., and P.G.; Visualization, J.M.G.A. and A.M.;
Supervision, A.R., G.D., E.H., C.A.; Project Administration, J.M.G.A. and A.R.; Funding
Acquisition, A.R., G.D., E.H., C.A.

Declaration of interests

The University of Geneva has licensed Flipper-TR probes to Spirochrome for
commercialization.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used Grammarly and ChatGPT in order to
improve readability in minor parts of the text. After using this tool, the authors reviewed
and edited the content and take full responsibility for the content of the publication.

Figures Titles and Legends

Figure 1: Flipper-TR reports membrane tension gradients in reconstituted
membranes. A: Schematics of supported lipid bilayer experiments. **B**: Examples of
expanding supported lipid bilayers with different compositions. Color indicates average
Flipper-TR fluorescence lifetime (ns); black mask shows average photon count. **C**:
Average Flipper-TR fluorescence lifetime in supported lipid bilayers with different
compositions, spatially averaged over the field of view. **D**: Front (expanding edge) and
rear (closest to lipid source) average Flipper-TR fluorescence lifetime in supported lipid
bilayers with different compositions. Lines represent data pairing. Right, front-rear
differences for different lipid compositions. **E**: Left, representative kymographs of
expanding supported lipid bilayers of different compositions. Right, linear fits of spatial
Flipper-TR gradients (ns/µm) over time. **B,E**: Scale bar, 10 µm.

Figure S1: Flipper-TR reports membrane tension gradients in reconstituted
membranes. A: Top left, bilayer spreading speed (µm/min) over time. Red line for visual
reference. Rest, bilayer spreading speed (µm/min) as a function of linear fits of spatial
Flipper-TR gradients (ns/µm) for different lipid compositions. Histograms of binned data
overlaid in black. **B**: Left, fluorescence confocal images of Flipper-TR and DOPE-atto647
in spreading bilayers of different compositions. Right, spatial profile of Flipper-TR (blue)
and DOPE-atto647 (red) fluorescence intensity, and Flipper-TR average fluorescence
lifetime (black, ns). Exponential (for fluorescence intensity, λ) or linear fits (for lifetime,
slope) overlaid.

Figure 2: Leading-edge extension increases Flipper-TR lifetime in migrating cells.
A: Left, representative live confocal FLIM microscopy image of fish keratocytes labeled
with Flipper-TR. Color indicates average Flipper-TR fluorescence lifetime (ns); black mask
shows average photon count. Right, average Flipper-TR fluorescence lifetime at the front
and rear 20% areas of migrating fish keratocytes (n=27, Welch's $P<10^{-4}$), with lifetime
profile along the normalized position (mean ± standard deviation). Front and rear regions
are shaded in red and blue respectively. **B**: Left, representative time-lapse image of

migrating fish keratocytes. Right, average Flipper-TR fluorescence lifetime at the front
(red) and rear (blue) 20% areas during the time lapse. **C**: Representative image of
polarized U2OS, RPE1, HeLa, and Cos7 cells labeled with Flipper-TR. Color indicates
average Flipper-TR fluorescence lifetime (ns); black mask shows average photon count.
D: Average Flipper-TR fluorescence lifetime as a function of distance from the edge at the
front (red) and rear (blue) of polarized U2OS (n=90), RPE1 (n=37), HeLa (n=11), and
Cos7 (n=15) cells. Line represents mean ± standard deviation. **E**: Average Flipper-TR
fluorescence lifetime as a function of distance from the edge and radial position (front at
0°) of polarized U2OS (n=90), RPE1 (n=37), HeLa (n=11), and Cos7 (n=15) cells. Color
indicates average Flipper-TR fluorescence lifetime (ns). **F**: Difference of average Flipper-
TR fluorescence lifetime between front and rear 10-μm regions as a function of cell aspect
ratio for U2OS (blue, n=132), RPE1 (magenta, n=43), HeLa (orange, n=63), and Cos7
(grey, n=32) cells. **G**: Left, representative image of fixed U2OS cells labeled with
phalloidin-Alexa647, Hoechst, and anti-vinculin-alexa488. Right, live image of U2OS
labeled with Flipper-TR. Dotted region indicates line scan region for panel H. **H**: Average
Flipper-TR fluorescence lifetime along a line scan marked in panel G. Dotted lines indicate
lamellipodium, lamella (Lm.), and cell body (Cb.). **I**: Left, segmentation of the
representative cell in panel G into lamellipodium (red, Lp.), lamella (dark grey, Lm.), and
cell body (light grey, Cb.). Right, average Flipper-TR fluorescence lifetime in these regions
(n=26, paired Welch's P). **J**: Average Flipper-TR fluorescence lifetime as a function of
distance from the edge and edge velocity in U2OS (n=37), RPE1 (n=56) cells. Color
indicates average Flipper-TR fluorescence lifetime (ns). **K**: Average Flipper-TR
fluorescence lifetime at protruding (P) and retracting (R) regions in U2OS (n=37), RPE1
(n=56) cells (paired Welch's P). **A,B,C,G**: Scale bar, 10 μm.

Figure S2: Leading-edge extension increases Flipper-TR lifetime in migrating cells.
A: Mean square displacement (MSD, μm²) and directionality ratio as a function of elapsed
time for RPE1 (magenta), U2OS (blue), HeLa (orange), and Cos7 (grey) cells, n=53. **B**:
Instantaneous speed (μm/min) and trajectories over 180 minutes for RPE1 (magenta),
U2OS (blue), HeLa (orange), and Cos7 (grey) cells, n=53. **C**: Representative image of
non-polarized U2OS, RPE1, HeLa, and Cos7 cells labeled with Flipper-TR. **D**: Average
Flipper-TR fluorescence lifetime as a function of distance from the edge at the front (red)
and rear (blue) of non-polarized U2OS (n=90), RPE1 (n=37), HeLa (n=11), and Cos7
(n=15) cells. Line represents mean ± standard deviation. **E**: Representative image of
U2OS and RPE1 cytoplasts labeled with Flipper-TR and Hoechst. **F**: Average Flipper-TR
fluorescence lifetime as a function of distance from the edge at the front (red) and rear
(blue) of U2OS (n=10) and RPE1 (n=14) cytoplasts. Line represents mean ± standard
deviation. **G**: Average Flipper-TR fluorescence lifetime as a function of distance from the
edge and radial position (front at 0°) of U2OS (n=10) and RPE1 (n=14) cytoplasts. Color
indicates average Flipper-TR fluorescence lifetime (ns). **A,C,E**: Scale bar, 10 μm.

Figure 3: Tension gradients are also present in non-migrating cells. A: Left,
representative images of ventral membrane (bottom plane) HeLa cells labeled with
Flipper-TR on cross, crossbow, and ring micropatterns. Right, average Flipper-TR
fluorescence lifetimes of cross, crossbow, and ring micropatterned cells (n=130, 52, 161,
Welch's P). Data distribution in black. **B**: Same as panel A, but on the dorsal plane (3 μm
above the glass surface). **C**: Average Flipper-TR fluorescence lifetime (ns) in cell regions
of cross (green), crossbow (blue), and ring (orange) micropatterned cells. Left, ventral
(bottom) vs. dorsal (middle) planes. Center, cell body (Cb) vs. protrusive areas (Prot).

Right, adhesive vs. non-adhesive areas of the cell body membrane. Data distribution in
black. **D**: Average Flipper-TR fluorescence lifetime as a function of distance from
micropattern edge and cell body edge. Color indicates average Flipper-TR fluorescence
lifetime (ns); black mask shows data point density. **E**: Top, cell averaging of Flipper-TR
photon counts in ventral membrane HeLa cells on micropatterns. Bottom, average Flipper-
TR lifetime. **F**: Top, average Flipper-TR fluorescence lifetime along the cell contour of
cross (green), crossbow (blue), and ring (orange) micropatterned cells. Bottom, lifetime
distribution across the cell. **G**: Average Flipper-TR lifetime along pattern edges in
crossbow (left, blue) and ring-patterned cells (orange). Shaded lines show individual
traces. **H**: Average Flipper-TR lifetime across pattern in crossbow (left, blue) and cross-
patterned cells (green). Shaded lines show individual traces. **A,B,E**: Scale bar, 10 μm.

Figure S3: Adhesion and cell morphology shape spatial gradients in membrane
tension but not major lipid components. A: Average fluorescence images at basal
plane of cross (top row, n=14), crossbow (middle row, n=13), and ring (bottom row, n=19)
micropatterned cells stained with phalloidin for actin (first column), vinculin antibody
(middle column). Last column, color overlay of Hoechst (yellow), vinculin (cyan), and actin
(magenta). **B**: Average Flipper-TR fluorescence lifetime (ns) at ventral plane vs. dorsal
plane of cross (green), crossbow (blue), and ring (orange) micropatterned cells. Labels
refer to r and p values from linear fit. **C**: Average Flipper-TR lifetime (ns) at ventral and
dorsal planes (bottom/middle), and bottom and top quintiles (low/high) of cross (green,
n=27), crossbow (blue, n=16), and ring (orange, n=22) micropatterned RPE1 cells. Labels
represent difference in average values and Welch's test P value. Data distribution in
black. **D**: Cell averaging of average Flipper-TR lifetime in ventral membrane of RPE1 cells
on micropatterns. **E**: Top, cell averaging of average Flipper-TR lifetime in ventral
membrane of patterned HeLa cells on different motifs. Bottom, average Flipper-TR lifetime
vs. distance from pattern center. Shaded areas represent adhesive micropattern location.
F: Schematics representing MALDI-MSI procedure. **G**: Histogram of pixel-wise lipid
fractions, and average lipid fractions for different lipid species: sphingomyelin (SM,
magenta), ceramide (Cer, grey), phosphatidylethanolamine (PE, blue),
phosphatidylcholine (PC, orange), and globotriaosylceramide (Gb3, green). **H**: Cell
averaging of spatial MALDI-MSI of different lipid species. Color scale represents local
differences from mean value normalized in terms of standard deviations. Arrows indicate
relative lipid accumulation. **I**: Average fluorescence images at basal plane of HeLa cells in
cross (left, n=10) and ring (right, n=10) micropatterns stained with filipin, Cholera toxin,
Equina toxin, and Shiga toxin targeting cholesterol, GM1, sphingomyelin, and
globotriaosylceramide, respectively. **A,D,I**: Scale bar, 10 μm.

Figure 4: Actin polymerization and adhesions scale tension gradients. A: Top, cell
averaging of average Flipper-TR lifetime in ventral membrane of HeLa cells on cross,
crossbow, and ring micropatterns under different drug treatments. Bottom, average
Flipper-TR lifetime vs. distance from center of ring micropattern. Shaded areas represent
adhesive micropattern location. **B**: Average Flipper-TR lifetime (ns) per cell of cross
(green), crossbow (blue), and ring (orange) micropatterned cells under different drug
treatments. Labels represent Welch's test P value ($P<10^{-3}$) and difference from control in
ns. Data distribution in black. **C**: Average Flipper-TR lifetime (ns) per cell of bottom and
top quintiles (low/high) of cross (green), crossbow (blue), and ring (orange)
micropatterned HeLa cells under different drug treatments. Labels display difference from
control in ns. Lines represent data pairing. Data distribution in black. **D**: Representative

fluorescence images at basal plane of cross, crossbow, and ring micropatterned cells
stained with phalloidin (red) and vinculin antibody (green) under different drug treatments.
A,D: Scale bar, 10 µm.

Figure S4: Effect of cell morphology on tension gradients. A: Representative
fluorescence images at basal plane of cross, crossbow, and ring micropatterned cells
stained with phalloidin (magenta), vinculin antibody (cyan), and Hoechst (yellow) showing
low (left column) or high number of protrusions (right). **B**: Cell averaging of average
Flipper-TR lifetime in ventral membrane of HeLa cells on micropatterns showing high (top)
or low (bottom) number of protrusions. **C**: Average Flipper-TR lifetime (ns) per cell at
ventral membrane vs. protrusion area (µm²) of cross (green), crossbow (blue), and ring
(orange) micropatterned HeLa cells. **D**: Histogram of protrusion areas (µm²) of cross
(green), crossbow (blue), and ring (orange) micropatterned HeLa cells. **E**: Average
Flipper-TR lifetime (ns) at ventral and dorsal planes (bottom/middle) of cross (green),
crossbow (blue), and ring (orange) micropatterned HeLa cells under different drug
treatments. Labels display difference from control in ns. Lines represent data pairing. Data
distribution in black. **F**: Left, representative images at basal plane of cross micropatterned
actin-GFP HeLa cells under control (top) and JLY (bottom) conditions. Blue line indicates
kymograph position. Blue arrows mark retrograde flow events. **G**: Cell averaging of
particle image velocimetry analysis of actin-GFP HeLa cells on cross, crossbow, and ring
micropatterns under control (top, n=10,10,12 respectively) and JLY treatments (bottom
n=4,4,4 respectively). Color and arrow lengths indicate cortical flow magnitude. **A,B,F,G**:
Scale bar, 10 µm.

Figure 5: Substrate rigidity and adhesiveness modulate membrane tension. A:
Diagram of confinement experiment showing cell morphology and bleb formation before
polarization into a leader bleb. **B**: Representative FLIM image of 3µm-confined HeLa cells
on pll-g-PEG (non-adhesive) stained with Flipper-TR. Lifetime is shown as $\Delta\tau$ from the
middle plane average of cells on glass. **C**: Median, quartile distribution, and individual
traces of Flipper-TR lifetime in stable blebs at different sections. Welch's p: 0.013 or n.s.
D: Representative FLIM image of HeLa cells on different substrates stained with Flipper-
TR. Lifetime shown as $\Delta\tau$ from the middle plane average of cells on glass. **E**: Median and
quartile distribution of Flipper-TR lifetime in HeLa cells on non-adhesive confinement,
fibronectin-coated glass, and fibronectin-coated hydrogels (4.0 and 0.5 kPa). Welch's
$p < 10^{-3}$ or n.s. **F**: Average clathrin light chain intensity and line scans (blue lines) for cross
(top, n=25) and ring (bottom, n=27) patterned cells. Patterned area in grey. **G**: Difference
in Flipper-TR lifetime of cilengitide-treated cells from control average for cross, crossbow,
and ring-patterned cells. **H**: Conceptual model of membrane tension gradient
maintenance. The diagram shows tension variation across the cell, highlighting high
tension on the dorsal side and gradients on the ventral side. Key factors include branched
actin filaments (red), cellular adhesions, and clathrin plaques (blue) acting as barriers,
creating tension gradients and directing lipid flow, crucial for cellular processes like
movement and signaling. **B, D, F, G**: Scale bar, 10 µm. A: 1 min.

Figure S5: Lipid diffusion and flows are limited in patterned cells. A: Left, FRAP
experimental design overlaid on images of HeLa MyrPalm-GFP on cross and ring
micropatterns. Line FRAP bleaching sites marked with red dotted lines; recovery tracked
along black, magenta, and blue arrows. Middle, normalized fluorescence intensity over

time after FRAP for spots at cross tip (grey), cross side (magenta), or ring side (blue).
Shaded lines represent individual time traces. Right, FRAP recovery constant (s^{-1}). **B**: Line
FRAP kymograph (inverted LUT) of ring cell from panel A and intensity vs. distance from
edge, colored by time elapsed from FRAP. Right, velocity (μm/min) of fluorescence
minima vs. distance to cell edge. Statistical test: Welch's *P* value=0.04. **C-D**: Cell mask
and SiR-actin images of HeLa cells on cross (panel C) or ring (panel D) pattern.
Kymographs from blue arrows are shown on the right. **E**: Parameters calculated from pair-
correlation function (pCF, left) and autocorrelation function fits (ACF, right). **F:** Schematics
of the acquisition of line scans, and representative line scan from a cell mask-stained
HeLa cell. **G-H**: Schematics of line scan acquisition and fluorescence correlation analysis
of patterned HeLa cells stained with CellMask dye under control (black) and JLY (red)
conditions. Mean values ± standard deviation. **G**: pCF analysis of cross (top) or ring
(bottom) patterned cells. First column, percentage of molecules arrived. Second column,
arrival time (s). **H**: ACF analysis of cross (top) or ring (bottom) patterned cells. First
column, diffusion coefficient (μm²/s). Second column, number of moving molecules (in
thousands). **A-D**: Scale bar, 10 μm

STAR Methods text

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will
be fulfilled by the lead contact, Aurélien Roux (aurelien.roux@unige.ch).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All original microscopy images and datasets shown in the figures, and all original code
and associated datasets have been deposited at an open repository and will be made
publicly available as of the date of publication (Figshare). Any additional data related to
this paper will be shared by the lead contact upon request.

Any additional information required to reanalyze the data reported in this paper is
available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Mammalian cell lines

Human cervical adenocarcinoma cells HeLa-Kyoto stably expressing myosin IIA (Myh9)-
GFP and LifeAct-mCherry or Myh9-GFP, or LifeAct-mCherry and the plasma membrane-
targeting CAAX box fused to GFP, or TALEN-edited ActB fused with GFP (Cellectis,
Paris, France), or HeLa, Cos7 and U2OS cells with no stable marker were cultured in
DMEM GlutaMAX medium (Gibco, #61965-026) supplemented with 10% FBS (Gibco,
#P30-193306A) and Penicillin Streptomycin (Thermofisher, #15140-122) at 37°C and 5%
CO2. RPE1 cells were cultured in DMEM / F-12 (Gibco) supplemented with 10% FBS and
1% Pen-Strep at 37°C in a 5% CO_2 incubator.

All cell lines were tested for mycoplasma contamination using Mycoplasmacheck PCR
Detection (Eurofins, #50400400), and treated once a year with Plasmocin profilactic

(InvivoGen, #ant-mpp) and treatment Mycoplasma Elimination Reagents (InvivoGen,
#ant-mpt).

Zebrafish keratocytes

Keratocyte cells were extracted from scales of zebrafish (*Danio Rerio*) while anesthetized
with Ethyl 3-aminobenzoate methanesulfonate (Tricaine, SIGMA, E10521). 3-5 scales
were extracted from each fish and placed on a glass-bottom dish containing Leibovitz's
medium (L-15, ThermoFisher #21083027), supplemented with 10% fetal bovine serum
and 1% penicillin/streptomycin antibiotic solution. The scales were subsequently
sandwiched between the dish' glass and a coverslip and placed in an incubator at 28⁰C.
Keratocytes were allowed to spread out from the scales for 12h. Subsequently, cells were
washed twice with PBS and detached from the plates by treating them with trypsin-EDTA
for 2-5 minutes. In order to remove the excess trypsin, cells were centrifuged in culture
medium at 1000 rpm for 3 minutes. After discarding the supernatant, cells were
resuspended in medium and plated in glass-bottom dishes for imaging. This protocol was
carried out by trained personnel at the Zebrafish Core Facility of the medical center of the
University of Geneva.

METHOD DETAILS

DNA constructs and transfections

Cells were transfected with plasmid DNA using Lipofectamine 3000 reagent (Thermofisher
#L3000001) transiently, according to manufacturer's protocol, and imaged from 24 to 72
hours after transfection. The following expression vectors were used for plasmid DNA
transfections: VAMP7-pHLuorin-pcDNA3 (gift from Thierry Galli, Inserm, France), eGFP-
Clathrin Light Chain C1 (gift from P. De Camilli, Yale University, USA).

Drug treatments and chemical probes

The following pharmacological inhibitors and chemical compounds were used: 10 µM of
ezrin inhibitor NSC668394 (Sigma #341216), 20 µM of Rho Kinase inhibitor Y-27632
(Sigma #Y0503), 8 µM of actin filament stabilizer Jasplakinolide (Cayman, #CAY-11705-
100), 5 µM of actin polymerization inhibitor Latrunculin B (Sigma, #428020), 1 µM of actin
polymerization inhibitor Latrunculin A (Sigma, #L5163), 85 µM of Arp2/3 actin
polymerization inhibitor CK-666 (Sigma #SML0006), 10 µM of α_V integrins inhibitor
Cilengitide (Sigma #SML1594), and 10 µg/ml actin inhibitor Cytochalasin B for cell
enucleation experiments (Sigma, #250255). Cells were imaged 30 minutes after
treatment. CellMask Green plasma membrane stain (Sigma, #C37608) and Flipper-TR
live cell fluorescent membrane tension probe (Lubio Science and Spirochrome, #SC020)
were used according to manufacturer's specifications at 1:1000 from a stock dilution in
DMSO, to a final concentration of 50 nM. SiR-actin staining (Spirochrome #CY-SC001)
was performed at 1 µM during 30 minutes with 10 µM verapamil following the
manufacturer's instructions.

Cell fixation and staining

Adherent cells were fixed for 15 minutes with 4% paraformaldehyde in PBS. Cells were
then permeabilized and blocked with the blocking buffer [5% Bovine Serum Albumin and
0.1% Saponin in PBS] for 30 min. Cells were washed three times with PBS before a 1 h
incubation with Phalloidin-AlexaFluorPlus647 at 1:400 from a DMSO stock according to
manufacturer's instructions (Thermofisher #A30107), Anti-Vinculin Vinculin Monoclonal

Antibody (7F9)-Alexa Fluor488 at 1:25 dillution (eBioscience Thermofisher #53-9777-82)
on 1% Gelatin in PBS. Cells were then washed three times with PBS, labeled by NucBlue
Fixed Cell Stain (DAPI, Thermofisher #R37606) for 10 min, and washed again three times
with PBS. Once stained, the cells were imaged directly on the microscope via the saved
localization.

For lipid localization, we stained cells with fluorescently-labelled bacterial toxins that
recognize different sphingolipid headgroups: Shiga toxin 1a-Cy3 (ShTxB1a, at 1:50) binds
to Gb3[100], Cholera toxin-AlexaFluor488 (ChTxB, at 1:800) binds the ganglioside GM1[101],
Equinatoxin II-AlexaFluor647 (EqTx, at 1:200) binds sphingomyelins[102] (all gifts from
Giovanni D'Angelo, EPFL, Switzerland), and Filipin III at 25 µg/ml (Sigma, #F4767). The
cells were fixed with 4% PFA, blocked in PBS containing 5% bovine serum albumin (BSA)
without detergent, and incubated with fluorescently labelled B-subunit toxins for 1h.

Cell enucleation

The day before enucleation, U2OS and RPE1 cells detached with trypsin/EDTA (0.05%
for 2 min, Gibco), and spread on a plastic polystyrene slide covered with bovine
Fibronectin (FN, 10µg/ml, Sigma-Aldrich #F4759) in their respective culture medium.

During enucleation, plastic slides seeded with cells were incubated at 37°C for 15 minutes
in the presence of 10 µg/ml Cytochalasin B (Sigma, #250255), then centrifuged at 15000
g for 40 minutes at 37°C in the presence of Cytochalasin. After centrifugation, to recover
from the effects of Cytochalasin, the cells on plastic slides were washed with PBS, and
transferred back to their respective drug-free culture media for 4 h at 37°C before use.

Migration time lapses of different mammalian cell lines

Mammalian cells were detached with trypsin/EDTA (0.05% for 2 min), then spread at a
density not exceeding 10% confluence into 8-well chambered cover glass (Lab-Tek II)
coated with bovine FN (10µg/ml) in their respective culture medium, 1 to 3 hours before
the experiments.

Time-Lapse Phase-Contrast Microscopy Multisite microscopy of cells in 8-well chambered
cover glass was performed in a humidified CO_2 chamber with an Axio Observer Inverted
TIRF microscope (Zeiss, 3i) and a Prime BSI (Photometrics) using a 10X objective (Zeiss,
10X). SlideBook 6 X64 software (version 6.0.17) was used for image acquisition. Cells
were imaged using phase-contrast imaging, with a time-lapse of 5 minutes between
frames for over 10 hours.

Confocal FLIM live imaging of Flipper-TR

During live FLIM acquisition, a corresponding phenol-red-free alternative of each culture
medium was used. U2OS, HeLa and Cos7 cells were imaged in FluoroBrite DMEM
(Gibco) during migration live experiments or Leibovitz's L-15 (Thermo Fisher Scientific,
#21083027) during micropatterning experiments in incubation at 37°C but without CO_2.
Media was supplemented with 10 % FBS and 1 % Pen-Strep while RPE1 cells were
imaged DMEM/F-12 without phenol red (Gibco) supplemented with 10 % FBS and 1 %
Pen-Strep.

Generally, cells were labelled for at least 10 minutes with FBS-free 50 medium containing
50 nM Flipper-TR. For supported lipid bilayers, the medium consisted on an aqueous
buffer of 10mM HEPES at pH 7.4. After labeling with Flipper-TR or Flipper-TR and
Hoechst, cells were imaged directly in glass-bottom microwell dishes at 37°C and 5%

CO2. For migration assays, untreated and enucleated cells were detached with
trypsin/EDTA (0.05% for 2 min), then spread at a density not exceeding 20% confluence
into 35 mm glass-bottom microwell dishes (MatTek) coated with bovine FN (10µg/ml) in
their respective culture medium, 1 to 3 hours before the experiments. After spreading, the
untreated cells were labeled with Flipper-TR (Spirochrome) in their respective acquisition
medium and incubated at 37°C for 10 min without probe washing. The Flipper-TR stock
solution was composed by dilution of a 1 mM Flipper-TR in DMSO as previously
described [24]. After spreading, the enucleated cells were labeled by 1:1000 dilution of a
stock solution of Flipper-TR and labeled by NucBlue Fixed Cell Stain (DAPI, Thermofisher
#R37606) in their respective acquisition medium and incubated at 37°C for 10 min without
probe washing.

The microscope was an inverted motorized microscope Eclipse Ti2 (Nikon) with a point
scanning A1 confocal system from Nikon, equipped with a Plan Apo λ 100x oil objective
with a NA 1.45 (Nikon, # MRD01905), a perfect focus system, a stage-top incubation
chamber (Okolab) allowing long acquisitions. The imaging on hydrogels and comparing
experiments was done instead with a water immersion 40x objective NA 1.15 (Nikon, #
MRD77410) to allow for a higher working distance.　The A1 confocal system was
equipped with 4 excitation lasers at 405, 488, 561 and 640 nm. For fluorescence lifetime
imaging (FLIM), we used a time-correlated photon counting system from PicoQuant
integrated on Nikon Elements software. Excitation was performed using a pulsed 485 nm
laser (PicoQuant #LDH-D- C-485) mounted on a laser-coupling unit, and a computer-
controlled multichannel picosecond diode laser driver (PicoQuant, Sepia PDF 828)
operating at 20 MHz. The emission signal was collected through a 600/50 nm bandpass
filter using a gated PMA hybrid 40 detector and a multichannel time-correlated single
photon counting system (PicoQuant, MultiHarp 150).

Single frames resulted from the integration of 2-10 frames depending on the sample and
the confocal scanning area, to a total count between 10 and 50 photons per pixel. Frame
rate during time lapse imaging in mammalian cells had to be set at 10 minutes to avoid
phototoxicity and arresting actin dynamics.

Confocal live imaging for fluctuation correlation spectroscopy

Regarding fluctuation correlation spectroscopy analysis, the microscope setup was the
same as for FLIM. For both line scanning used for autocorrelation function (ACF) and
paired-correlation function (pCF) analysis, pixel dwell time was set at 5.2 microseconds,
line time at 960 microseconds, pixel size was set at 160 nm, giving a line length of 10
micrometers. We positioned the 10-micrometer confocal line scans crossing the pattern
boundaries, thus across regions with a sharp decrease of the Flipper-TR fluorescence
lifetime. For the ring pattern, the line scan sits at the cell edge in the region where
retrograde flows are dissipating. The adhesive area is located on the outer side of the line
scan. For the cross pattern, the line scan is located closer to the center of the cell, and the
adhesive pattern is on the inner side of the line.

Preparation of supported lipid bilayers

For supported lipid bilayer experiments, we used: 18:1 (Δ9-Cis) PC (DOPC) (Avanti
#850375), 18:1 PS (DOPS) (Avanti #840035), porcine brain SM (Avanti #860062), plant
cholesterol (Avanti #700100) at the specified mol% ratios. Some lipid mixtures included
0.02 mol% Atto 647 1,2-Dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) (Sigma

#67335). Phospholipids or cholesterol stocks were dissolved in chloroform at
concentrations ranging from 1-10 mg/ml and stored at -80°C in Argon atmosphere. Lipid
mixes were calculated for a final mass of 0.25 mg. The chloroform in the vials was
evaporated under an Argon flow in a fume hood and lipid films were formed. The vials
were then stored in a vacuum oven at room temperature overnight to remove any residual
chloroform.

To prepare liposomes, the vials were brought to room temperature and rehydrated by
adding aqueous buffer HEPES 10mM at pH 7.4 to a final concentration of 0.5 mg/ml of
lipids and mixing. The liposome mixtures were placed on a parafilm surface. 50-micron
silica particles (Sigma #904384-2G) were first diluted 3x with distilled water, washed three
times, and deposed on the lipid droplets on parafilm. The liposome-beads aqueous
mixtures were then stored in a vacuum oven at room temperature overnight to remove
any residual chloroform. To prepare the spreading bilayer assay, culture-grade 35-mm
glass-bottom dishes (Mattek #P35G-1.5-14-C) were plasma cleaned (Harrick Plasma
#PDC-32G) for 2 minutes at high power. Immediately after, dried liposome-bead mixtures
were scrapped from the parafilm surface, deposed on the plasma-cleaned surface, and
rehydrated with 1ml of 10 mM HEPES pH7.4 buffer containing 50 nM Flipper-TR (Lubio
Science and Spirochrome, #SC020)

TIRF-FRAP experiments on patterned cells

All total internal reflection fluorescence microscopy (TIRF) movies were recorded on an
Olympus IX83 widefield microscope equipped with a 150×/NA1.45 objective and an
ImageEM X2 EM-CCD camera (Hamamatsu) under the control of the VisiView software
(Visitron Systems). The 488 nm and 561 nm laser lines were used for illumination of GFP-
and mCherry-tagged proteins. Excitation and emission were filtered using a TRF89902
405/488/561/647 nm quad-band filter set (Chroma). Laser angles were controlled by iLas2
(Roper Scientific).

Fluorescence recovery after photobleaching (FRAP) experiments were performed using a
custom-built set-up that focuses a 488-nm laser beam at the sample plane, on the
Olympus IX81 widefield microscope described above. The diameter of the bleach spot or
width of the bleach line was approximately 0.5 µm.

Chemical passivation and micropatterning

Glass surfaces passivated with Polyethyleneglycol (PEG) were prepared for protein
micropatterning. First, glass bottom dishes (Mattek) were activated by exposing them to
air plasma (Harrick Plasma, PDC-32G) for 3 minutes. Subsequently, the dishes were
treated with a 0.1 mg/ml poly-lysine (PLL) (Sigma) solution for 30 min and washed with
10mM HEPES buffer (pH 8.4). By using this same buffer, a solution of 50 mg/ml
polyethylene glycol (PEG) (molecular weight 5,000) linked to a succinimidyl valerate group
(SVA, Laysan Bio) was prepared and applied to passivate the PLL-coated surface for
1.5 h. Finally, dishes were washed with PBS 3 times.

Micropatterns were generated by using the system PRIMO (Alvéole), mounted on an
inverted microscope Nikon Eclipse Ti-2. In the presence of a photo-initiator compound
(PLPP, Alvéole) and DMD-generated patterns of UV light (375 nm), PEG is degraded.
After illumination (1200 mJ/mm) through a 20x objective, PLL is exposed. After rinsing
with PBS, fibronectin (Calbiochem) was incubated at 50 µg/ml at room temperature for
5 min to coat the PEG-free motifs with the cell-adhesive protein. The excess of

fibronectin was washed out with PBS. PBS was finally replaced by medium, and a
suspension of cells was added at densities of roughly 10^5 cells per cm^2. Samples were
kept in an incubator at 37 °C and 5% CO2. After 10–30 min, non-adhered cells were
washed out.

MALDI-MSI sample preparation

Cells were directly seeded on micropatterns with cross shape in complete media. After
aspiration of media, cells were washed twice with PBS, followed by fixation in 0.25%
glutaraldehyde for 15 minutes. For MALDI-MSI analyses, 150 µL of 2,5-dihydroxybenzoic
acid (DHB), (30 mg/mL in 50:50 acetonitrile/water/0.1% TFA), were deposited on the
surface of the samples using the automatic SMALDIPrep (TransMIT GmbH, Giessen,
Germany).

Cell culture on hydrogels

Commercially-available EasyCoat hydrogels (Cell Guidance Systems, # SV3510-EC-0.5-
EA and # SV3510-EC-4-EA) were used to control substrate rigidity. The hydrogels were
pre-coated with fibronectin to facilitate cell adhesion. Briefly, hydrogels of specified rigidity
(0.5 kPa and 4.0 kPa) were equilibrated in PBS-fibronectin solution (1 µg/mL, Sigma
#F1141) for 1 hour to ensure uniform coating. Excess fibronectin was washed off with
PBS before seeding the cells onto the hydrogels. Cells were cultured on these substrates
under standard conditions, allowing us to investigate the effects of substrate rigidity on
membrane tension and cell behavior.

Cell confinement

Cell confinement was achieved using a PDMS micropillar array mounted on a coverslip
mounted on a PDMS device (named 'suction cup') trigged by a vacuum pump and
pressure controller, following established methods[85,103]. Briefly, a SU8 photolithography
mold was prepared with micropillars. This mold is used to create a PDMS confinement
chamber of 3 micrometers in height, plasma-bonded to a 12-mm glass coverslip. The
microfabricated confiner coverslips were treated with plasma for 1 min, and incubated with
0.5 mg/mL pLL-*g*-PEG (SuSoS, PLL(20)-g[3.5]-PEG(2)) in 10 mM pH 7.4 HEPES buffer
for 1h at room temperature. Cells were trypsinized and seeded in a glass-bottom dish
coated with PLL-g-PEG and the PDMS device was placed on top, creating a confined
environment between the glass and the PDMS coverslip. This setup allowed us to trigger
leader-bleb migration by confining the cells to a 3-micrometer height, promoting a non-
adhesive environment conducive to the desired migration behavior.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis of experimental data

982 Statistical analyses for all experiments were performed in Python 3 (NumPy and SciPy
983 libraries) or Microsoft Excel. Statistical data are presented as median or mean ± standard
984 deviation or interquartile range. For each panel, sample size (*n*), experiment count (*N*)
985 statistical tests used, and *P* values are specified in the figure panel and/or legends. Plots
986 were made using the SciPy and matplotlib Python libraries.

Basic image processing

988 Basic image analysis and format conversion was performed on ImageJ/Fiji software.
989 Images were imported from NIS Elements (.nd2 format) using the BioFormats plugin.

 Migration parameters and cell morphology of mammalian cell lines

Individual cells were tracked semi-automatically by random selection of cells from video
images and manual tracking of migration pathways using the Manual Tracking function in
ImageJ. At least 50 cells were tracked for each cell type (n ≥ 50). The cell migration
tracking data were analyzed as described previously[104].

Cell aspect ratio was calculated from binary masks using Image/Fiji by fitting ellipses to
the cell shape.

Protrusion area in control patterned cells was segmented manually from intensity images.

General considerations about the different Flipper-TR lifetime estimates

Flipper-TR is a mechano-responsive molecular probe that changes its fluorescence
properties in response to changes in membrane packing, making it an invaluable tool for
studying the mechanical properties of the plasma membrane. Upon insertion to the
membrane, Flipper-TR's fluorescence lifetime changes with the tension in the lipid bilayer.
Generally, the fluorescence lifetime is the average time the molecule spends in the
excited state before emitting a photon and returning to the ground state. To calculate the
fluorescence lifetime of Flipper-TR, we used a time-correlated single photon counting
(TCSPC) device as described above. This technique measures the time between the
excitation pulse and the subsequent photon emission, producing as an output the
histogram of photon arrival times. The distribution of these arrival times can be analyzed
to determine the fluorescence lifetime.

Due to the photochemistry of the molecule, Flipper-TR photon arrival time distribution
follows a bi-exponential decay $I(t)$, in the form: $I(t) = A_1 e^{-t/\tau_1} + A_2 e^{-t/\tau_2}$, where A_i and
τ_i represent the amplitude and the decay of a given exponential i, respectively. Previous
studies have employed different methods to estimate the Flipper-TR lifetime. One
common approach is to fit the fluorescence decay curve to a biexponential decay model,
which accounts for the presence of two distinct exponential decay components [24]. From
this model, researchers can extract either the larger exponent (i.e.: τ_1) or calculate an
intensity-weighted average of the two exponents, $\bar{\tau} = \sum_i \tau_i \alpha_i$ [90]. These methods can
provide detailed insights into the Flipper-TR lifetime, including a χ^2 used to judge the
goodness, a greater precision, and the ability to decompose complex lifetime distributions,
but they require more computational resources and can be more sensitive to noise and
initial parameter estimates. Moreover, even at high photon counts, the ability to determine
the precise values of α_i and τ_i by a biexponential fit can be hindered by parameter
correlation [105]. To avoid problems involved with fitting, other works have used phasor plots
have been used to measure changes in Flipper-TR lifetime [106,107]. Phasor plots offer a
graphical representation of the fluorescence decay characteristics, simplifying the analysis
of complex decay patterns. By mapping each pixel to specific G/S coordinates on the
phasor plot, one can visually assess changes in membrane tension and identify different
lifetime components without assuming any feature of the fluorescence decay, such as the
number of exponents. This is particularly useful when using complex samples or Flipper
variants where the photochemical properties have not been fully characterized.

For this work we have instead relied on the barycenter of the lifetime distribution (i.e.:
mean photon arrival time or "fast lifetime" calculation) to estimate fluorescence lifetime. As
the instrumentation induces a delay in the photon arrival, the average lifetime equals time
span from the barycenter of the instrument response function (IRF) to the barycenter of

the decay in a pixel-wise manner, as defined by our TCSPC manufacturer (PicoQuant).
This method is simple, straightforward, and computationally efficient, especially reliable at
low photon counts. The calculation yields a single lifetime exponent, that we refer to as
"average Flipper-TR lifetime" $\bar{\tau}$, numerically equivalent to the intensity-weighted average
of the two exponents coming from a fit. The need for a robust lifetime estimator at low
photon counts was very important for this work as we were highly limited by sample
illumination. Protrusion dynamics are particularly sensitive to phototoxicity and even
relatively low illumination can arrest actin polymerization, so the photon counts were
generally too low to be able to fit in a robust manner.

Flipper-TR lifetime estimation

Processing and plotting of FLIM images. FLIM images acquired on NIS software consisted
of a channel containing average lifetime values and another with photon counts per pixel.
The images were exported as .tif and processed on a custom-made Python script.
Lifetime images were applied to a 2D median filter with a kernel size of 3 pixels
(corresponding to 0.75 μm). In the exported lifetime images, the rainbow color scale in
each pixel represents the values of the decay times. On top of the rainbow color scale, a
dark mask is applied to represent photon counts (i.e.: black for no counts). This allows for
a proper visual representation of the data because it masks the background where lifetime
values are just due to noise.

Flipper-TR lifetime region-of-interest averages. When averaging pixels, the lifetimes of
each pixel τ_i were weighted by the photon count n_i, so that $\bar{\tau} = [\sum_{i=1}^{n}(n_i\tau_i)]/N$. Different
regions corresponded to specific criteria, explained as follows. In keratocytes, front/rear
regions were manually segmented to cover approximately one-fourth of the cell each. In
migrating mammalian cells, front/rear regions were defined as 60° angular sections
extending 0-10 μm from the edge using a custom python script. Front and rear polarity
angles were manually determined based on cell morphology, not lifetime values. For
supported lipid bilayers, only the 100 μm closest to the advancing edge in the field of view
was considered, using a custom python script. Front and rear regions were defined as the
20% closest and most distant areas from the advancing edge, respectively.

Flipper-TR lifetime distribution and ROIs in patterned cells. Generally, FLIM images were
filtered and values under a threshold photon count of 20 photons were defined as NaN
and not considered for analysis. For overall distribution, i.e. the lifetime spatial probability
map, cells were aligned according to the patterned regions as previously described[108], and
a pixel-wise average was calculated using a custom python script. For cells adhered on
micropatterns, different regions were defined to report spatial lifetime gradients:
bottom/middle planes; cell body/protrusions; adhesive/non-adhesive; high/low. For the
bottom plane there was no further segmentation, but for the middle plane (3 microns
above the glass surface) a manual selection was used to segment the plasma membrane.
Cell body and protrusions were manually segmented. Adhesive and non-adhesive regions
were defined based on the fluorescence from micropatterns. Finally, to calculate the
distance map 2D histogram, distances to the selected boundaries were defined per pixel
using binary masks and the 'distance map' function in ImageJ/Fiji. Then, pixels were
ordered and averaged to form a 2D histogram. To define the high and low regions we first
defined the pixels were 99% of control cells were present. From this, we defined a 'high'
region, containing the highest two lifetime deciles, and a 'low' region, containing the

lowest 2 lifetime deciles of the control lifetime spatial distribution. These two regions of
interest were applied to other conditions to represent lifetime gradients.

Flipper-TR lifetime spatial decays in mammalian cell lines were calculated using a custom
python script based on distances to the cell edge and position in the micropattern. For
radial/linear gradients, a 5 micron-wide region was defined and the average lifetime
calculated per micron.

Flipper-TR lifetime as a function of edge position in migrating cells was calculated using a
cell mask based on photon counts and, from a mask, a distance map to the edge. Then,
pixels were ordered and averaged to form a 2D histogram.

Flipper-TR lifetime as a function of edge velocity. The edge velocity was calculated using
a custom python script. For mammalian cells, the edge displacement was calculated by
segmenting the cell contours at each timepoints and calculating the differences between
the distance maps. For supported lipid bilayers, the images were manually cropped into
rectangular regions. The velocity at each timepoint (µm/min) was defined as the net
increase of area (µm^2), divided by the frame rate (min) and the crop width (µm).

Flipper-TR lifetime in protruding/retracting cell regions. The condition to classify edges
intro protruding/retracting was a velocity higher/lower than 0.2 µm/min. From this, angular
sections extending 0-10 µm from the edge were defined.

Flipper-TR lifetime differences to control distributions. For some drug treatments, instead
of the absolute Flipper-TR distributions, the difference to the control was instead plotted.
First, the average control distribution was calculated on a pixel-wise basis. Then, these
control values were subtracted to the drug-treated images on a cell-by-cell basis and the
average differences plotted.

Flipper-TR lifetime in hydrogels and non-adhesive migration was instead calculated using
the SymPhoTime 64 software (PicoQuant) to fit fluorescence decay data (from manual
regions of interest) to a dual exponential model (n-exponential reconvolution) after
deconvolution for the instrument response function calculated by the software. Lifetime
was then expressed in difference in nanoseconds from the reference value, which is the
lifetime at the middle plane of cells adhered to glass.

Particle image velocimetry of actin flows

PIV analysis. Movies were acquired at a 2-frame-per-second rate and segmented using
Ilastik[68] to define cell boundaries. Pixels outside cell boundaries were not considered for
the PIV calculation. PIV vectors were calculated on a custom macro based on the
'Iterative PIV' ImageJ/Fiji plugin[109], using approximately an XY grid of 1.25 µm and a
search area of 1.5 µm. Importantly, even though we used the 'Iterative PIV' plugin, the PIV
was calculated on a single window basis, so there is no a priori correlation between
neighbor vectors.

Quantification of membrane events

Stable clathrin coverage was calculated from a minimum time projection of TIRF movies
over 5 minutes at a 1 frame/second rate. This projection was then binarized to display the
sites where the clathrin signal was stable over 5 minutes on a cell-by-cell basis.

Dynamic clathrin coverage was calculated in an analogous manner. First, we obtained a
binary mask resulting from a maximum projection of the clathrin signal. Then we

subtracted to this mask the stable clathrin sites. This yields a mask containing the sites
where the clathrin signal was dynamic at some point over 5 minutes on a cell-by-cell
basis, excluding the clathrin plaques where the signal was stable.

VAMP7 exocytosis events were manually selected from 5-minute TIRF movies of VAMP7
transfected and patterned cells.

Finally, the probability density distributions were plotted as a function of the distance to
the center of the pattern and normalized to integrate to 1.

Actin-based segmentation of cell regions

To determine the effect of actin organization on Flipper-TR lifespan. Phalloidin-A647
images of each cell were segmented into three regions; Lamellipodium, Lamella, and Cell
body regions. Binary segmentations were generated using the pixel classification process
in Ilastik[68]. Segmentation masks were then applied to the Flipper-TR lifetime image
acquired before fixation, to calculate the average Flipper-TR lifetime in each type of region
for each cell.

Fluctuation correlation spectroscopy analysis

To study lipid diffusion in our system, we employed fluorescence fluctuation spectroscopy
(FFS). This involved first construction of single-channel FFS line scans acquired across
the ventral membrane of patterned cells stained with the plasma membrane dye CellMask
into kymographs (x-dimension represents distance (i.e., 64 pixels), y-dimension
represents time (i.e., 100,000 lines)) and application of a detrending algorithm to remove
slow timescale artefacts (e.g., photobleaching). Then application of the autocorrelation
function (ACF) on a pixel-by-pixel basis ($\delta r = 0$) to three spatially distinct sections across
the kymograph (i.e., pixels 1-21, 22-42, and 43-63), which alongside fitting of the resulting
ACFs to a 2-component diffusion model, enabled recovery of the average number of
moving molecules (G_0) as well as their local diffusion coefficient (D_0) to be mapped across
the non-adhesive versus adhesive coating and the interface in between (Figure 5E, right
panel). And finally, application of the pair correlation function (pCF) on a pixel-by-pixel
basis ($\delta r = 5$) across the entire kymograph (i.e., pixels 1-59 from left to right and pixels 5-
64 from right to left), which alongside fitting of the resulting pCFs to a Gaussian function,
enabled recovery of the molecules direction dependent arrival time (τ) as well as their
transport efficiency (i.e., G_τ/G_0) (Figure 5E, left panel).

Analysis of TIRF-FRAP experiments on patterned cells

FRAP experiments performed on MyrPalm-GFP patterned HeLa cells were analyzed as
previously described[111]. Mean fluorescence values were measured from regions of
interest representing the background, the cell, and the membrane. A custom-written
Python script was used to subtract background fluorescence, correct for photobleaching,
and normalize the values between 0 (corrected fluorescence immediately after
photobleaching) and 1 (mean corrected fluorescence of 5 s before photobleaching). The
recovery curves of individual experiments were aligned to bleach time and averaged. The
average was fitted to a single exponential equation from which the mobile fraction and
recovery half-time were calculated.

To calculate the velocity of lipid flows on line-FRAP experiments, fluorescence intensity
was integrated over 11 microns (100 pixels) parallel to the bleaching line, and the minima
were tracked for 12 frames (0.5 seconds per frame) using a custom-made Python script.

MALDI-MSI data analysis

MSI experiments were performed using AP-SMALDI5 AF systems that couple a Q Exactive orbital trapping mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) with an atmospheric-pressure scanning-microprobe MALDI imaging source (AP-SMALDI, TransMIT GmbH, Giessen, Germany). The MALDI laser focus was optimized manually using the source cameras aiming at a diameter of the focused beam of 7 µm. For each pixel, the spectrum was accumulated from 50 laser shots at 100 Hz. MS parameters in the Tune software (Thermo Fisher Scientific) were set to the spray voltage of 4 kV, S-Lens 100 eV, and capillary temperature of 250°C. The step size of the sample stage was set to 5 µm. Positive ion mode measurements were performed in full scan mode in the mass range m/z 400-1600 with a resolving power set to R = 240000 at m/z = 200. Mass spectra were internally calibrated using the lock mass feature of the instrument.

The output of the MALDI-MSI was processed using a custom-made Python script. The output consists of a multi-dimensional .tif with 389 channels representing the peak intensity at each m/z bin and XY spatial coordinates at 0.5 microns per pixel. First, an integration of all counts allowed to identify pattern positions with cells and to align patterns with one another. Then, the relative intensity of peaks from specific lipid species was calculated over the total intensity of only the known peaks (112/389). For lipid species present in more than one peak, the relative abundances were added together. This yields a lipid fraction per pixel, per cell, of each lipid species. Lipid fractions were also normalized to analyze their spatial heterogeneity. To do so, lipid fractions from a given lipid species from all cells, all pixels were pooled together, and their mean and standard deviation were calculated. Lipid fractions were then subtracted from the mean and divided by the standard deviation.

Supplementary video titles and legends

Supplementary movie 1, corresponding to Fig. 1. Expanding supported lipid bilayers (SLBs) with different compositions.

Supplementary movie 2, corresponding to Fig. 2. Confocal FLIM time lapses of migrating U2OS cells stained with Flipper-TR.

Supplementary movie 3, corresponding to Fig. 5. Actin dynamics in patterned HeLa Actin-GFP cells, examples of lamellipodium and stress fiber dominated phenotypes.

Supplementary movie 4, corresponding to Fig. 5. Actin dynamics in patterned HeLa Actin-GFP cells in control (+DMSO) versus JLY-treated conditions.

References

1. Nissen, J., Gritsch, S., Wiegand, G., and Rädler, J.O. (1999). Wetting of phospholipid membranes on hydrophilic surfaces - Concepts towards self-healing membranes. Eur. Phys. J. B - Condens. Matter Complex Syst. *10*, 335–344. https://doi.org/10.1007/s100510050862.
2. Helfrich, W. (1973). Elastic properties of lipid bilayers: theory and possible experiments. Z. Naturforschung Teil C Biochem. Biophys. Biol. Virol. *28*, 693–703. https://doi.org/10.1515/znc-1973-11-1209.
3. Dai, J., and Sheetz, M.P. (1995). Regulation of endocytosis, exocytosis, and shape by membrane tension. Cold Spring Harb. Symp. Quant. Biol. *60*, 567–571. https://doi.org/10.1101/sqb.1995.060.01.060.
4. Gauthier, N.C., Fardin, M.A., Roca-Cusachs, P., and Sheetz, M.P. (2011). Temporary increase in plasma membrane tension coordinates the activation of exocytosis and contraction during cell spreading. Proc. Natl. Acad. Sci. *108*, 14467–14472. https://doi.org/10.1073/pnas.1105845108.

5.   Lachuer, H., Le, L., Lévêque-Fort, S., Goud, B., and Schauer, K. (2023). Spatial organization
of lysosomal exocytosis relies on membrane tension gradients. Proc. Natl. Acad. Sci. U. S. A.
120, e2207425120. https://doi.org/10.1073/pnas.2207425120.

6.   Wu, X.-S., Elias, S., Liu, H., Heureaux, J., Wen, P.J., Liu, A.P., Kozlov, M.M., and Wu, L.-G.
(2017). Membrane Tension Inhibits Rapid and Slow Endocytosis in Secretory Cells. Biophys.
J. *113*, 2406–2414. https://doi.org/10.1016/j.bpj.2017.09.035.

7.   Gabella, C., Bertseva, E., Bottier, C., Piacentini, N., Bornert, A., Jeney, S., Forró, L.,
Sbalzarini, I.F., Meister, J.-J., and Verkhovsky, A.B. (2014). Contact angle at the leading
edge controls cell protrusion rate. Curr. Biol. CB *24*, 1126–1132.
https://doi.org/10.1016/j.cub.2014.03.050.

8.   Hetmanski, J.H.R., de Belly, H., Busnelli, I., Waring, T., Nair, R.V., Sokleva, V., Dobre, O.,
Cameron, A., Gauthier, N., Lamaze, C., et al. (2019). Membrane Tension Orchestrates Rear
Retraction in Matrix-Directed Cell Migration. Dev. Cell *51*, 460-475.e10.
https://doi.org/10.1016/j.devcel.2019.09.006.

9.   Houk, A.R., Jilkine, A., Mejean, C.O., Boltyanskiy, R., Dufresne, E.R., Angenent, S.B.,
Altschuler, S.J., Wu, L.F., and Weiner, O.D. (2012). Membrane tension maintains cell polarity
by confining signals to the leading edge during neutrophil migration. Cell *148*, 175–188.
https://doi.org/10.1016/j.cell.2011.10.050.

10.   Keren, K., Pincus, Z., Allen, G.M., Barnhart, E.L., Marriott, G., Mogilner, A., and Theriot, J.A.
(2008). Mechanism of shape determination in motile cells. Nature *453*, 475–480.
https://doi.org/10.1038/nature06952.

11.   Kozlov, M.M., and Mogilner, A. (2007). Model of Polarization and Bistability of Cell
Fragments. Biophys. J. *93*, 3811–3819. https://doi.org/10.1529/biophysj.107.110411.

12.   Lieber, A.D., Schweitzer, Y., Kozlov, M.M., and Keren, K. (2015). Front-to-Rear Membrane
Tension Gradient in Rapidly Moving Cells. Biophys. J. *108*, 1599–1603.
https://doi.org/10.1016/j.bpj.2015.02.007.

13.   Lieber, A.D., Yehudai-Resheff, S., Barnhart, E.L., Theriot, J.A., and Keren, K. (2013).
Membrane Tension in Rapidly Moving Cells Is Determined by Cytoskeletal Forces. Curr. Biol.
23, 1409–1417. https://doi.org/10.1016/j.cub.2013.05.063.

14.   Chakraborty, M., Mukherjee, B., Nalinakshan, N., Biswas, A., Nayak, R.K., and Sinha, B.
(2022). Effect of heterogeneous substrate adhesivity of follower cells on speed and tension
profile of leader cells in primary keratocyte collective cell migration. Biol. Open *11*, bio058893.
https://doi.org/10.1242/bio.058893.

15.   Bergert, M., Lembo, S., Sharma, S., Russo, L., Milovanović, D., Gretarsson, K.H., Börmel, M.,
Neveu, P.A., Hackett, J.A., Petsalaki, E., et al. (2021). Cell Surface Mechanics Gate
Embryonic Stem Cell Differentiation. Cell Stem Cell *28*, 209-216.e4.
https://doi.org/10.1016/j.stem.2020.10.017.

16.   De Belly, H., Stubb, A., Yanagida, A., Labouesse, C., Jones, P.H., Paluch, E.K., and Chalut,
K.J. (2021). Membrane Tension Gates ERK-Mediated Regulation of Pluripotent Cell Fate.
Cell Stem Cell *28*, 273-284.e6. https://doi.org/10.1016/j.stem.2020.10.018.

17.   Mueller, J., Szep, G., Nemethova, M., de Vries, I., Lieber, A.D., Winkler, C., Kruse, K., Small,
J.V., Schmeiser, C., Keren, K., et al. (2017). Load Adaptation of Lamellipodial Actin
Networks. Cell *171*, 188-200.e16. https://doi.org/10.1016/j.cell.2017.07.051.

18.   Dal, J., and Sheetz, M.P. (1995). Axon membrane flows from the growth cone to the cell
body. Cell *83*, 693–701. https://doi.org/10.1016/0092-8674(95)90182-5.

19.   Shi, Z., Innes-Gold, S., and Cohen, A.E. (2022). Membrane tension propagation couples
axon growth and collateral branching. Sci. Adv. *8*, eabo1297.
https://doi.org/10.1126/sciadv.abo1297.

20.   Raucher, D., and Sheetz, M.P. (2000). Cell Spreading and Lamellipodial Extension Rate Is
Regulated by Membrane Tension. J. Cell Biol. *148*, 127–136.
https://doi.org/10.1083/jcb.148.1.127.

21.   Venkova, L., Vishen, A.S., Lembo, S., Srivastava, N., Duchamp, B., Ruppel, A., Williart, A.,
Vassilopoulos, S., Deslys, A., Garcia Arcos, J.M., et al. (2022). A mechano-osmotic feedback
couples cell volume to the rate of cell deformation. eLife *11*, e72381.
https://doi.org/10.7554/eLife.72381.

22.   Yan, Q., Gomis Perez, C., and Karatekin, E. (2024). Cell Membrane Tension Gradients,
Membrane Flows, and Cellular Processes. Physiology *39*, 000–000.
https://doi.org/10.1152/physiol.00007.2024.

23.   Mussel, M., and Schneider, M.F. (2021). Sound pulses in lipid membranes and their potential
1273   function in biology. Prog. Biophys. Mol. Biol. *162*, 101–110.
1274   https://doi.org/10.1016/j.pbiomolbio.2020.08.001.
1275   24.   Colom, A., Derivery, E., Soleimanpour, S., Tomba, C., Molin, M.D., Sakai, N., González-
1276   Gaitán, M., Matile, S., and Roux, A. (2018). A fluorescent membrane tension probe. Nat.
1277   Chem. *10*, 1118–1125. https://doi.org/10.1038/s41557-018-0127-3.
1278   25.   Raedler, J., Strey, H., and Sackmann, E. (1995). Phenomenology and Kinetics of Lipid
1279   Bilayer Spreading on Hydrophilic Surfaces. Langmuir *11*, 4539–4548.
1280   https://doi.org/10.1021/la00011a058.
1281   26.   Bernal, R., Pullarkat, P.A., and Melo, F. (2007). Mechanical Properties of Axons. Phys. Rev.
1282   Lett. *99*, 018301. https://doi.org/10.1103/PhysRevLett.99.018301.
1283   27.   Fernández, P., and Pullarkat, P.A. (2010). The role of the cytoskeleton in volume regulation
1284   and beading transitions in PC12 neurites. Biophys. J. *99*, 3571–3579.
1285   https://doi.org/10.1016/j.bpj.2010.10.027.
1286   28.   Ghose, A., and Pullarkat, P. (2023). The role of mechanics in axonal stability and
1287   development. Semin. Cell Dev. Biol. *140*, 22–34.
1288   https://doi.org/10.1016/j.semcdb.2022.06.006.
1289   29.   Leterrier, C., and Pullarkat, P.A. (2022). Mechanical role of the submembrane spectrin
1290   scaffold in red blood cells and neurons. J. Cell Sci. *135*, jcs259356.
1291   https://doi.org/10.1242/jcs.259356.
1292   30.   Diz-Muñoz, A., Fletcher, D.A., and Weiner, O.D. (2013). Use the force: membrane tension as
1293   an organizer of cell shape and motility. Trends Cell Biol. *23*, 47–53.
1294   https://doi.org/10.1016/j.tcb.2012.09.006.
1295   31.   Cohen, A.E., and Shi, Z. (2020). Do Cell Membranes Flow Like Honey or Jiggle Like Jello?
1296   BioEssays *42*, 1900142. https://doi.org/10.1002/bies.201900142.
1297   32.   De Belly, H., Yan, S., Borja da Rocha, H., Ichbiah, S., Town, J.P., Zager, P.J., Estrada, D.C.,
1298   Meyer, K., Turlier, H., Bustamante, C., et al. (2023). Cell protrusions and contractions
1299   generate long-range membrane tension propagation. Cell *186*, 3049-3061.e15.
1300   https://doi.org/10.1016/j.cell.2023.05.014.
1301   33.   Gomis Perez, C., Dudzinski, N.R., Rouches, M., Landajuela, A., Machta, B., Zenisek, D., and
1302   Karatekin, E. (2022). Rapid propagation of membrane tension at retinal bipolar neuron
1303   presynaptic terminals. Sci. Adv. *8*, eabl4411. https://doi.org/10.1126/sciadv.abl4411.
1304   34.   Shi, Z., Graber, Z.T., Baumgart, T., Stone, H.A., and Cohen, A.E. (2018). Cell Membranes
1305   Resist Flow. Cell *175*, 1769-1779.e13. https://doi.org/10.1016/j.cell.2018.09.054.
1306   35.   Sitarska, E., and Diz-Muñoz, A. (2020). Pay attention to membrane tension: Mechanobiology
1307   of the cell surface. Curr. Opin. Cell Biol. *66*, 11–18. https://doi.org/10.1016/j.ceb.2020.04.001.
1308   36.   Barnoy, A., Tsaturyan, A.K., and Kozlov, M.M. (2023). Mechanism of tension propagation in
1309   cell membranes. Preprint at bioRxiv, https://doi.org/10.1101/2023.03.22.533804
1310   https://doi.org/10.1101/2023.03.22.533804.
1311   37.   Dai, J., and Sheetz, M.P. (1999). Membrane Tether Formation from Blebbing Cells. Biophys.
1312   J. *77*, 3363–3370. https://doi.org/10.1016/S0006-3495(99)77168-7.
1313   38.   Eggeling, C., Ringemann, C., Medda, R., Schwarzmann, G., Sandhoff, K., Polyakova, S.,
1314   Belov, V.N., Hein, B., von Middendorff, C., Schönle, A., et al. (2009). Direct observation of the
1315   nanoscale dynamics of membrane lipids in a living cell. Nature *457*, 1159–1162.
1316   https://doi.org/10.1038/nature07596.
1317   39.   He, H.-T., and Marguet, D. (2011). Detecting Nanodomains in Living Cell Membrane by
1318   Fluorescence Correlation Spectroscopy. Annu. Rev. Phys. Chem. *62*, 417–436.
1319   https://doi.org/10.1146/annurev-physchem-032210-103402.
1320   40.   Honigmann, A., Mueller, V., Ta, H., Schoenle, A., Sezgin, E., Hell, S.W., and Eggeling, C.
1321   (2014). Scanning STED-FCS reveals spatiotemporal heterogeneity of lipid interaction in the
1322   plasma membrane of living cells. Nat. Commun. *5*, 5412.
1323   https://doi.org/10.1038/ncomms6412.
1324   41.   Lasserre, R., Guo, X.-J., Conchonaud, F., Hamon, Y., Hawchar, O., Bernard, A.-M., Soudja,
1325   S.M., Lenne, P.-F., Rigneault, H., Olive, D., et al. (2008). Raft nanodomains contribute to
1326   Akt/PKB plasma membrane recruitment and activation. Nat. Chem. Biol. *4*, 538–547.
1327   https://doi.org/10.1038/nchembio.103.
1328   42.   Sharma, P., Varma, R., Sarasij, R.C., Ira, Gousset, K., Krishnamoorthy, G., Rao, M., and
1329   Mayor, S. (2004). Nanoscale Organization of Multiple GPI-Anchored Proteins in Living Cell
1330   Membranes. Cell *116*, 577–589. https://doi.org/10.1016/S0092-8674(04)00167-9.

43. Varma, R., and Mayor, S. (1998). GPI-anchored proteins are organized in submicron
domains at the cell surface. Nature *394*, 798–801. https://doi.org/10.1038/29563.

44. Köster, D.V., and Mayor, S. (2016). Cortical actin and the plasma membrane: inextricably
intertwined. Curr. Opin. Cell Biol. *38*, 81–89. https://doi.org/10.1016/j.ceb.2016.02.021.

45. Schneider, F., Waithe, D., Clausen, M.P., Galiani, S., Koller, T., Ozhan, G., Eggeling, C., and
Sezgin, E. (2017). Diffusion of lipids and GPI-anchored proteins in actin-free plasma
membrane vesicles measured by STED-FCS. Mol. Biol. Cell *28*, 1507–1518.
https://doi.org/10.1091/mbc.e16-07-0536.

46. Shaw, T.R., Ghosh, S., and Veatch, S.L. (2021). Critical Phenomena in Plasma Membrane
Organization and Function. Annu. Rev. Phys. Chem. *72*, 51–72.
https://doi.org/10.1146/annurev-physchem-090419-115951.

47. Gordon, V.D., Deserno, M., Andrew, C.M.J., Egelhaaf, S.U., and Poon, W.C.K. (2008).
Adhesion promotes phase separation in mixed-lipid membranes. Europhys. Lett. *84*, 48003.
https://doi.org/10.1209/0295-5075/84/48003.

48. Akimov, S.A., Kuzmin, P.I., Zimmerberg, J., and Cohen, F.S. (2007). Lateral tension
increases the line tension between two domains in a lipid bilayer membrane. Phys. Rev. E
75, 011919. https://doi.org/10.1103/PhysRevE.75.011919.

49. Hamada, T., Kishimoto, Y., Nagasaki, T., and Takagi, M. (2011). Lateral phase separation in
tense membranes. Soft Matter *7*, 9061–9068. https://doi.org/10.1039/C1SM05948C.

50. Oglęcka, K., Rangamani, P., Liedberg, B., Kraut, R.S., and Parikh, A.N. (2014). Oscillatory
phase separation in giant lipid vesicles induced by transmembrane osmotic differentials. eLife
3, e03695. https://doi.org/10.7554/eLife.03695.

51. Rathe, V., Kuckla, D., and Monzel, C. (2021). Chapter Two - Phase separation in biological
membranes: An overview with focus on experimental effects of illumination and osmotic
pressure changes. In Advances in Biomembranes and Lipid Self-Assembly, A. Iglič, M.
Rappolt, and A. J. García-Sáez, eds. (Academic Press), pp. 31–66.
https://doi.org/10.1016/bs.abl.2021.11.002.

52. Wongsirojkul, N., Shimokawa, N., Opaprakasit, P., Takagi, M., and Hamada, T. (2020).
Osmotic-Tension-Induced Membrane Lateral Organization. Langmuir *36*, 2937–2945.
https://doi.org/10.1021/acs.langmuir.9b03893.

53. Yanagisawa, M., Imai, M., and Taniguchi, T. (2008). Shape Deformation of Ternary Vesicles
Coupled with Phase Separation. Phys. Rev. Lett. *100*, 148102.
https://doi.org/10.1103/PhysRevLett.100.148102.

54. Boyd, M.A., and Kamat, N.P. (2018). Visualizing Tension and Growth in Model Membranes
Using Optical Dyes. Biophys. J. *115*, 1307–1315. https://doi.org/10.1016/j.bpj.2018.08.021.

55. Zhang, Y.-L., Frangos, J.A., and Chachisvilis, M. (2006). Laurdan fluorescence senses
mechanical strain in the lipid bilayer membrane. Biochem. Biophys. Res. Commun. *347*, 838–
841. https://doi.org/10.1016/j.bbrc.2006.06.152.

56. Ayuyan, A.G., and Cohen, F.S. (2008). Raft composition at physiological temperature and pH
in the absence of detergents. Biophys. J. *94*, 2654–2666.
https://doi.org/10.1529/biophysj.107.118596.

57. Zapata-Mercado, E., Azarova, E.V., and Hristova, K. (2022). Effect of osmotic stress on live
cell plasma membranes, probed via Laurdan general polarization measurements. Biophys. J.
121, 2411–2418. https://doi.org/10.1016/j.bpj.2022.05.016.

58. Dal Molin, M., Verolet, Q., Colom, A., Letrun, R., Derivery, E., Gonzalez-Gaitan, M., Vauthey,
E., Roux, A., Sakai, N., and Matile, S. (2015). Fluorescent Flippers for Mechanosensitive
Membrane Probes. J. Am. Chem. Soc. *137*, 568–571. https://doi.org/10.1021/ja5107018.

59. Bozic, B., Svetina, S., and Zeks, B. (1997). Theoretical analysis of the formation of
membrane microtubes on axially strained vesicles. Phys. Rev. E *55*, 5834–5842.
https://doi.org/10.1103/PhysRevE.55.5834.

60. Cuvelier, D., Derényi, I., Bassereau, P., and Nassoy, P. (2005). Coalescence of Membrane
Tethers: Experiments, Theory, and Applications. Biophys. J. *88*, 2714–2726.
https://doi.org/10.1529/biophysj.104.056473.

61. Evans, E., Bowman, H., Leung, A., Needham, D., and Tirrell, D. (1996). Biomembrane
Templates for Nanoscale Conduits and Networks. Science *273*, 933–935.
https://doi.org/10.1126/science.273.5277.933.

62. Heinrich, V., Božič, B., Svetina, S., and Žekš, B. (1999). Vesicle Deformation by an Axial
Load: From Elongated Shapes to Tethered Vesicles. Biophys. J. *76*, 2056–2071.
https://doi.org/10.1016/S0006-3495(99)77362-5.

63. Hochmuth, F.M., Shao, J.Y., Dai, J., and Sheetz, M.P. (1996). Deformation and flow of
membrane into tethers extracted from neuronal growth cones. Biophys. J. *70*, 358–369.

64. Sayem Karal, M.A., Masum Billah, M., Ahmed, M., and Kabir Ahamed, M. (2023). A review on
the measurement of the bending rigidity of lipid membranes. Soft Matter *19*, 8285–8304.
https://doi.org/10.1039/D3SM00882G.

65. Bayard, F., Chen, X.-X., García-Arcos, J.M., Roux, A., Sakai, N., and Matile, S. (2023).
Hydrophobic Interfacing of Fluorescent Membrane Probes. ChemistryEurope *1*, e202300041.
https://doi.org/10.1002/ceur.202300041.

66. Roffay, C., García-Arcos, J.M., Chapuis, P., López-Andarias, J., Schneider, F., Colom, A.,
Tomba, C., Meglio, I.D., Dunsing, V., Matile, S., et al. (2023). Technical insights into
fluorescence lifetime microscopy of mechanosensitive Flipper probes. Preprint at bioRxiv,
https://doi.org/10.1101/2022.09.28.509885 https://doi.org/10.1101/2022.09.28.509885.

67. L. Symons, J., Cho, K.-J., T. Chang, J., Du, G., Neal Waxham, M., F. Hancock, J., Leventhal,
I., and R. Leventhal, K. (2021). Lipidomic atlas of mammalian cell membranes reveals
hierarchical variation induced by culture conditions, subcellular membranes, and cell
lineages. Soft Matter *17*, 288–297. https://doi.org/10.1039/D0SM00404A.

68. Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C., Schiegg, M., Ales,
J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine learning for (bio)image
analysis. Nat. Methods *16*, 1226–1232. https://doi.org/10.1038/s41592-019-0582-9.

69. Batchelder, E.L., Hollopeter, G., Campillo, C., Mezanges, X., Jorgensen, E.M., Nassoy, P.,
Sens, P., and Plastino, J. (2011). Membrane tension regulates motility by controlling
lamellipodium organization. Proc. Natl. Acad. Sci. *108*, 11429–11434.
https://doi.org/10.1073/pnas.1010481108.

70. Goldman, R.D., Pollack, R., and Hopkins, N.H. (1973). Preservation of Normal Behavior by
Enucleated Cells in Culture. Proc. Natl. Acad. Sci. U. S. A. *70*, 750–754.

71. Azioune, A., Carpi, N., Tseng, Q., Théry, M., and Piel, M. (2010). Protein Micropatterns. In
Methods in Cell Biology (Elsevier), pp. 133–146. https://doi.org/10.1016/S0091-
679X(10)97008-8.

72. Théry, M., Pépin, A., Dressaire, E., Chen, Y., and Bornens, M. (2006). Cell distribution of
stress fibres in response to the geometry of the adhesive environment. Cell Motil.
Cytoskeleton *63*, 341–355. https://doi.org/10.1002/cm.20126.

73. Taylor, M.J., Lukowski, J.K., and Anderton, C.R. (2021). Spatially Resolved Mass
Spectrometry at the Single Cell: Recent Innovations in Proteomics and Metabolomics. J. Am.
Soc. Mass Spectrom. *32*, 872–894. https://doi.org/10.1021/jasms.0c00439.

74. Sezgin, E., Leventhal, I., Mayor, S., and Eggeling, C. (2017). The mystery of membrane
organization: composition, regulation and roles of lipid rafts. Nat. Rev. Mol. Cell Biol. *18*, 361–
374. https://doi.org/10.1038/nrm.2017.16.

75. Peng, G.E., Wilson, S.R., and Weiner, O.D. (2011). A pharmacological cocktail for arresting
actin dynamics in living cells. Mol. Biol. Cell *22*, 3986–3994. https://doi.org/10.1091/mbc.e11-
04-0379.

76. Bovellan, M., Romeo, Y., Biro, M., Boden, A., Chugh, P., Yonis, A., Vaghela, M., Fritzsche,
M., Moulding, D., Thorogate, R., et al. (2014). Cellular Control of Cortical Actin Nucleation.
Curr. Biol. *24*, 1628–1635. https://doi.org/10.1016/j.cub.2014.05.069.

77. Fritzsche, M., Thorogate, R., and Charras, G. (2014). Quantitative Analysis of Ezrin Turnover
Dynamics in the Actin Cortex. Biophys. J. *106*, 343–353.
https://doi.org/10.1016/j.bpj.2013.11.4499.

78. Lee, J., Gustafsson, M., Magnusson, K.-E., and Jacobson, K. (1990). The Direction of
Membrane Lipid Flow in Locomoting Polymorphonuclear Leukocytes. Science *247*, 1229–
1233. https://doi.org/10.1126/science.2315695.

79. Tanaka, M., Kikuchi, T., Uno, H., Okita, K., Kitanishi-Yumura, T., and Yumura, S. (2017).
Turnover and flow of the cell membrane for cell migration. Sci. Rep. *7*, 12970.
https://doi.org/10.1038/s41598-017-13438-5.

80. Weisswange, I., Bretschneider, T., and Anderson, K.I. (2005). The leading edge is a lipid
diffusion barrier. J. Cell Sci. *118*, 4375–4380. https://doi.org/10.1242/jcs.02551.

81. Digman, M.A., and Gratton, E. (2011). Lessons in Fluctuation Correlation Spectroscopy.
Annu. Rev. Phys. Chem. *62*, 645–668. https://doi.org/10.1146/annurev-physchem-032210-
103424.

82. Digman, M.A., and Gratton, E. (2009). Imaging Barriers to Diffusion by Pair Correlation
Functions. Biophys. J. *97*, 665–673. https://doi.org/10.1016/j.bpj.2009.04.048.

83. Hinde, E., Thammasiraphop, K., Duong, H.T.T., Yeow, J., Karagoz, B., Boyer, C., Gooding,
J.J., and Gaus, K. (2017). Pair correlation microscopy reveals the role of nanoparticle shape
in intracellular transport and site of drug release. Nat. Nanotechnol. *12*, 81–89.
https://doi.org/10.1038/nnano.2016.160.

84. Colin-York, H., Heddleston, J., Wait, E., Karedla, N., deSantis, M., Khuon, S., Chew, T.-L.,
Sbalzarini, I.F., and Fritzsche, M. (2022). Quantifying Molecular Dynamics within Complex
Cellular Morphologies using LLSM-FRAP. Small Methods *6*, 2200149.
https://doi.org/10.1002/smtd.202200149.

85. García-Arcos, J.M., Ziegler, J., Grigolon, S., Reymond, L., Shajepal, G., Cattin, C.J.,
Lomakin, A., Müller, D., Ruprecht, V., Wieser, S., et al. (2022). Advected percolation in the
actomyosin cortex drives amoeboid cell motility. Preprint at bioRxiv,
https://doi.org/10.1101/2022.07.14.500109 https://doi.org/10.1101/2022.07.14.500109.

86. Liu, Y.-J., Le Berre, M., Lautenschlaeger, F., Maiuri, P., Callan-Jones, A., Heuzé, M., Takaki,
T., Voituriez, R., and Piel, M. (2015). Confinement and Low Adhesion Induce Fast Amoeboid
Migration of Slow Mesenchymal Cells. Cell *160*, 659–672.
https://doi.org/10.1016/j.cell.2015.01.007.

87. García-Arcos, J.M., Jha, A., Waterman, C.M., and Piel, M. (2024). Blebology: principles of
bleb-based migration. Trends Cell Biol. *0*. https://doi.org/10.1016/j.tcb.2024.02.009.

88. Ghibaudo, M., Saez, A., Trichet, L., Xayaphoummine, A., Browaeys, J., Silberzan, P., Buguin,
A., and Ladoux, B. (2008). Traction forces and rigidity sensing regulate cell functions. Soft
Matter *4*, 1836–1843. https://doi.org/10.1039/B804103B.

89. Baschieri, F., Dayot, S., Elkhatib, N., Ly, N., Capmany, A., Schauer, K., Betz, T., Vignjevic,
D.M., Poincloux, R., and Montagnac, G. (2018). Frustrated endocytosis controls contractility-
independent mechanotransduction at clathrin-coated structures. Nat. Commun. *9*, 3825.
https://doi.org/10.1038/s41467-018-06367-y.

90. Roffay, C., Molinard, G., Kim, K., Urbanska, M., Andrade, V., Barbarasa, V., Nowak, P.,
Mercier, V., García-Calvo, J., Matile, S., et al. (2021). Passive coupling of membrane tension
and cell volume during active response of cells to osmosis. Proc. Natl. Acad. Sci. U. S. A.
118, e2103228118. https://doi.org/10.1073/pnas.2103228118.

91. Fogelson, B., and Mogilner, A. (2014). Computational Estimates of Membrane Flow and
Tension Gradient in Motile Cells. PLOS ONE *9*, e84524.
https://doi.org/10.1371/journal.pone.0084524.

92. Barbotin, A., Urbančič, I., Galiani, S., Eggeling, C., Booth, M., and Sezgin, E. (2020). z-STED
Imaging and Spectroscopy to Investigate Nanoscale Membrane Structure and Dynamics.
Biophys. J. *118*, 2448–2457. https://doi.org/10.1016/j.bpj.2020.04.006.

93. Kusumi, A., Nakada, C., Ritchie, K., Murase, K., Suzuki, K., Murakoshi, H., Kasai, R.S.,
Kondo, J., and Fujiwara, T. (2005). Paradigm Shift of the Plasma Membrane Concept from
the Two-Dimensional Continuum Fluid to the Partitioned Fluid: High-Speed Single-Molecule
Tracking of Membrane Molecules. Annu. Rev. Biophys. Biomol. Struct. *34*, 351–378.
https://doi.org/10.1146/annurev.biophys.34.040204.144637.

94. Fritzsche, M., Li, D., Colin-York, H., Chang, V.T., Moeendarbary, E., Felce, J.H., Sezgin, E.,
Charras, G., Betzig, E., and Eggeling, C. (2017). Self-organizing actin patterns shape
membrane architecture but not cell mechanics. Nat. Commun. *8*, 14347.
https://doi.org/10.1038/ncomms14347.

95. Köster, D.V., Husain, K., Iljazi, E., Bhat, A., Bieling, P., Mullins, R.D., Rao, M., and Mayor, S.
(2016). Actomyosin dynamics drive local membrane component organization in an in vitro
active composite layer. Proc. Natl. Acad. Sci. *113*, E1645–E1654.
https://doi.org/10.1073/pnas.1514030113.

96. Welf, E.S., Miles, C.E., Huh, J., Sapoznik, E., Chi, J., Driscoll, M.K., Isogai, T., Noh, J.,
Weems, A.D., Pohlkamp, T., et al. (2020). Actin-Membrane Release Initiates Cell Protrusions.
Dev. Cell *55*, 723-736.e8. https://doi.org/10.1016/j.devcel.2020.11.024.

97. Garner, R.M., and Theriot, J.A. (2022). Leading edge maintenance in migrating cells is an
emergent property of branched actin network growth. eLife *11*, e74389.
https://doi.org/10.7554/eLife.74389.

98. Li, T.-D., Bieling, P., Weichsel, J., Mullins, R.D., and Fletcher, D.A. (2022). The molecular
mechanism of load adaptation by branched actin networks. eLife *11*, e73145.
https://doi.org/10.7554/eLife.73145.

99. Mitchell, A.C., Wall, J.E., Murray, J.G., and Morgan, C.G. (2002). Measurement of
nanosecond time-resolved fluorescence with a directly gated interline CCD camera. J.
Microsc. *206*, 233–238. https://doi.org/10.1046/j.1365-2818.2002.01030.x.

100. Jacewicz, M., Clausen, H., Nudelman, E., Donohue-Rolfe, A., and Keusch, G.T. (1986).
Pathogenesis of shigella diarrhea. XI. Isolation of a shigella toxin-binding glycolipid from
rabbit jejunum and HeLa cells and its identification as globotriaosylceramide. J. Exp. Med.
163, 1391–1404. https://doi.org/10.1084/jem.163.6.1391.
101. van Heyningen, S. (1974). Cholera Toxin: Interaction of Subunits with Ganglioside GM1.
Science *183*, 656–657. https://doi.org/10.1126/science.183.4125.656.
102. Schön, P., García-Sáez, A.J., Malovrh, P., Bacia, K., Anderluh, G., and Schwille, P. (2008).
Equinatoxin II permeabilizing activity depends on the presence of sphingomyelin and lipid
phase coexistence. Biophys. J. *95*, 691–698. https://doi.org/10.1529/biophysj.108.129981.
103. Le Berre, M., Zlotek-Zlotkiewicz, E., Bonazzi, D., Lautenschlaeger, F., and Piel, M. (2014).
Methods for Two-Dimensional Cell Confinement. In Methods in Cell Biology (Elsevier), pp.
213–229. https://doi.org/10.1016/B978-0-12-800281-0.00014-2.
104. Gorelik, R., and Gautreau, A. (2014). Quantitative and unbiased analysis of directional
persistence in cell migration. Nat. Protoc. *9*, 1931–1943.
https://doi.org/10.1038/nprot.2014.131.
105. Lakowicz, J.R. (2007). Principles of Fluorescence Spectroscopy (Springer Science &
Business Media).
106. Puji Pamungka, K.K., Fureraj, I., Assies, L., Sakai, N., Mercier, V., Chen, X.-X., Vauthey, E.,
and Matile, S. (2024). Core-Alkynylated Fluorescent Flippers: Altered Ultrafast Photophysics
to Track Thick Membranes. Angew. Chem. Int. Ed Engl., e202406204.
https://doi.org/10.1002/anie.202406204.
107. Ragaller, F., Sjule, E., Urem, Y.B., Schlegel, J., El, R., Urbancic, D., Urbancic, I., Blom, H.,
and Sezgin, E. (2024). Quantifying Fluorescence Lifetime Responsiveness of Environment-
Sensitive Probes for Membrane Fluidity Measurements. J. Phys. Chem. B *128*, 2154–2167.
https://doi.org/10.1021/acs.jpcb.3c07006.
108. Schauer, K., Duong, T., Bleakley, K., Bardin, S., Bornens, M., and Goud, B. (2010).
Probabilistic density maps to study global endomembrane organization. Nat. Methods *7*, 560–
566. https://doi.org/10.1038/nmeth.1462.
109. Martiel, J.-L., Leal, A., Kurzawa, L., Balland, M., Wang, I., Vignaud, T., Tseng, Q., and Théry,
M. (2015). Chapter 15 - Measurement of cell traction forces with ImageJ. In Methods in Cell
Biology Biophysical Methods in Cell Biology., E. K. Paluch, ed. (Academic Press), pp. 269–
287. https://doi.org/10.1016/bs.mcb.2014.10.008.
110. Wohland, T., Maiti, S., and Macháň, R. (2020). An Introduction to Fluorescence Correlation
Spectroscopy (IOP Publishing).
111. Phair, R.D., Gorski, S.A., and Misteli, T. (2004). Measurement of dynamic protein binding to
chromatin in vivo, using photobleaching microscopy. Methods Enzymol. *375*, 393–414.
https://doi.org/10.1016/s0076-6879(03)75025-3.
1546

Figure 1: Flipper-TR reports membrane tension gradients in reconstituted membranes

Figure S1: Flipper-TR reports membrane tension gradients in reconstituted membranes.

Figure 2: Leading-edge extension increases Flipper-TR lifetime in migrating cells.

Figure S2: Leading-edge extension increases Flipper-TR lifetime in migrating cells.

Figure 3: Tension gradients are also present in non-migrating cells.

Figure S3: Adhesion and cell morphology shape spatial gradients in membrane tension but not major lipid components

Figure 4: Actin polymerization and adhesions scale tension gradients.

Figure S4: Effect of cell morphology on tension gradients.

Figure 5: Substrate rigidity and adhesiveness modulate membrane tension.

A Non-adherent cell
↓ 3μm confinement ~minute
Elongated blebs + Round blebs
↓ Polarization ~hour

B Non-adhesive 3μm confinement
ΔF-TR lifetime (ns): +1.5 / +1.0 / +0.5 / 0 / -0.5 / -1.0 / -1.5
Rel. photon count: 1.0 / 0.8 / 0.6 / 0.4 / 0.2 / 0.0

C ΔF-TR lifetime (ns)
p=0.01   n.s.
Rear — Front
Position in bleb
n=39

D 0.5 kPa + fibronectin   4 kPa + fibronectin   Glass + fibronectin
ΔF-TR lifetime (ns): +1.5 / +1.0 / +0.5 / 0 / -0.5 / -1.0 / -1.5
Rel. photon count: 1.0 / 0.8 / 0.6 / 0.4 / 0.2 / 0.0

E ΔF-TR lifetime (ns)
n=14   n=29   n=17   n=15
p<10⁻⁴   p<10⁻⁴   p<10⁻⁴   n.s.   n.s.
B M   B M   B M   B M
Non-adh. Adhes.   4 kPa   0.5 kPa
Glass surface   Hydrogel
Bottom / Middle

F Stable Clathrin LC   Line scan
n=25
n=27
Norm. intensity (AU): 0.0 / 0.5 / 1.0
Intensity (AU) vs Δx (μm)

G + cilengitide 10μM
n=22   n=27   n=68
ΔF-TR lifetime (ns): +0.3 / +0.2 / +0.1 / 0 / -0.1 / -0.2 / -0.3
Cell probability: 1.0 / 0.8 / 0.6 / 0.4 / 0.2 / 0.0

H Dorsal membrane: homogenous high tension
Actin bundles + Adhesions   Branched actin
Low tension — High tension
Ventral membrane: Tension gradients
Branched actin   Lipid order Tension
1 Branched actin polymerization → High tension
Branched actin   Plasma membrane   Tension   via ERMs
2 Adhesions and clathrin plaques → Low tension
Patterned area   Lipid diffusion   Adhesions
3 Tension gradients
Low tension   Retrograde actin flow   High tension

Figure S5: Lipid diffusion and flows are limited in patterned cells.