Article

Title:
Ancient tree-topologies and gene-flow processes among human lineages in Africa

Short Title:
Ancient demographic histories of African genetic lineages

Authors:
Gwenna Breton 1,2*
Per Sjödin 1
Panagiotis I. Zervakis 1
Romain Laurent 3
Alain Froment 4
Agnès E. Sjöstrand 1
Barry S. Hewlett 5
Luis B. Barreiro 6
George H. Perry 7,8
Himla Soodyall 9,10
Evelyne Heyer 3
Carina M. Schlebusch 1,11,12,*
Mattias Jakobsson 1,11,12,*
Paul Verdu 3,*

Authors affiliations:
Human Evolution, Department of Organismal Biology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18C, SE-752 36 Uppsala, Sweden
Centre for Medical Genomics, Department of Clinical Genetics and Genomics, Sahlgrenska University Hospital, Gothenburg, Sweden
UMR7206 Eco-anthropology, CNRS-MNHN-Université Paris Cité, Paris, France
UMR 208 Patrimoines locaux, IRD-MNHN, Paris, France
Department of Anthropology, Washington State University, Vancouver, Washington, USA
Department of Medicine, Section of Genetic Medicine, University of Chicago, Chicago, IL, USA
Department of Anthropology, Pennsylvania State University, University Park, PA 16801, USA
Department of Biology, Pennsylvania State University, University Park, PA 16801, USA
Division of Human Genetics, School of Pathology, Faculty of Health Sciences, University of the Witwatersrand and National Health Laboratory Service, Johannesburg, South Africa
Academy of Science of South Africa
Palaeo-Research Institute, University of Johannesburg, P.O. Box 524, Auckland Park, 2006, South Africa
SciLifeLab, Uppsala, Sweden

* Authors for correspondence:
gwenna.breton@gu.se,   paul.verdu@mnhn.fr,   mattias.jakobsson@ebc.uu.se, carina.schlebusch@ebc.uu.se

Keywords:

Abstract

The deep history of human evolution in Africa remains intensely debated with increasingly complex models being proposed. To investigate human evolutionary history in Africa, we sequenced and investigated 73 novel high-quality whole genomes from 14 Central and Southern African populations with diverse cultural practices. With extensive simulations and machine-learning Approximate Bayesian Computation inferences, we reconstruct jointly their complex history of divergences and migrations. We find vast genome-wide diversity within and among populations, often uncorrelated with geography and cultural practices. This shows the necessity to explicitly consider local genomic patterns, without merging samples from differentiated populations indiscriminately into larger *a priori* categories to reconstruct human evolutionary histories. We find that tree-like population-histories with long periods of drift between short pulses of unidirectional gene-flow best explain the data, in contrast to continuous gene-flow. Without the need to invoke archaic admixture, the detailed patterns of genomic variation can be accurately fitted to a model where the lineage ancestral to Khoe-San populations diverged around 300,000 years ago from a lineage ancestral to rainforest hunter-gatherers and neighboring agriculturalists. Finally, throughout history, the short periods of gene-flow each coincide with epochs of major ecological and cultural changes in Sub-Saharan Africa.

Introduction

Unraveling genetic structure and gene flow among human lineages in Africa is crucial to the understanding of the biological evolution and diversity of *Homo sapiens* throughout the continent [1–3].

Population geneticists largely agree today that *Homo sapiens* spent a large part of its genetic evolution within Africa only, between its gradual emergence from anatomically archaic forms 600,000-200,000 years ago and the beginning of the Out-of-Africa expansions to the rest of the world around 100,000-50,000 years ago [2]. For the past 20 years, in various regions of Africa, the detailed demographic and evolutionary history that shaped the genetic diversity of extant populations has been the subject of numerous investigations [3–18].

However, how ancient genetic divergences and the dynamics of admixture and/or migration events among ancient human lineages influenced the genetic landscape observed today throughout Africa largely remains to be assessed [19]. Classically tested demographic models [1,20–22], range from a unique ancestral *Homo sapiens* genetic population having recently and rapidly diverged into extant African populations via series of founding events and/or multifurcations (often referred to as tree-like models), to models where extant populations diverged in a remote past and remained isolated over long periods of time until relatively recently (often referred to as within Africa multiregional models). Moreover, each model may encompass possible gene exchanges among lineages via migration or admixture processes. Importantly, the timing, duration, and/or intensity of such gene-flow events may fundamentally change the most likely topology of bifurcating tree-like models compared to results obtained without gene flows, and may also create reticulations among pairs of lineages throughout history [22]. Finally, these classical albeit highly complex models have recently been enriched with possible introgression events from now extinct, unknown, or unsampled non-*Homo sapiens* or ancient "ghost" *Homo-sapiens* lineages [3,22–26]. Such latter models became plausible in Africa in analogy to the likely events of introgressions from now extinct hominid species into *Homo sapien*s lineages unveiled outside of Africa by the major advances of paleogenomics [27,28].

In this context, several recent investigations tested a variety of the above models with maximum-likelihood approaches relying on whole genome sequences from several extant populations sampled throughout the continent. They often reached highly contrasted results, with different ancient tree-topologies between different numbers of ancient lineages [2,22,24–26]. Furthermore, they found that reticulations among a limited number of ancient lineages explained the observed genetic patterns without the necessity of archaic introgressions [22]; or, instead, that admixture with other non-*Homo sapiens* or ancient *Homo sapiens* "ghost" populations most satisfactorily explained the results [23–26].

A possible source of such vast differences among obtained results is likely the variety of population sets considered in each study at a continental scale. Indeed, genetic diversity and differentiation among populations is maximal in Africa [3,29,30], and previously shown to be due to substantial differences in local demographic histories (e.g. [4,6,9,11]). Therefore, representing vast regions of the continent by a single population [22,24], or merging data from differentiated groups [26], likely influenced results obtained across studies and their interpretations.

In addition, and most importantly, all above models are highly nested, i.e. overlapping and genetically largely indistinguishable for vast spaces of model-parameter values. Indeed, depending on values of divergence times, effective population sizes, timing, duration and intensity of migration or admixture events among lineages, tree-like, reticulated, and even multiregional models, with or without ancient admixture, may be hardly identifiable. This essential methodological difficulty is well illustrated by the often relatively similar values of posterior likelihoods of the various competing models, thus indeed proved empirically hard to

discriminate with the maximum-likelihood approaches deployed, even more so when competing models differ in the number and specifications of parameters explored with different statistics. These fundamental statistical issues remain despite the use of high-quality whole genomes, an increasing number of populations considered at once, and increasingly powerful methods based on innovative statistics [22,24,26,31–33]. Finally, the lack of ancient DNA data older than a few thousand years in Africa, strongly hampers the empirical testing of the possible occurrence of ancient *Homo sapiens* "ghost" or non-*Homo sapiens* introgression events in *Homo sapiens* lineages in Africa [22–25,30].

In this context, we investigated 74 high coverage (>30x) whole genome sequences from an anthropologically well-characterized sample of hunter-gatherer, herder, and agricultural neighboring populations from Central and Southern Africa (**Figure1**, **Table1, TableS1**), merged together with 105 previously published high-quality whole genomes [27,34–37]. In particular, we investigated genomic diversity patterns and aimed at reconstructing ancient demographic histories among lineages having led to different groups of Khoe-San hunter-gatherer or herder populations from Southern Africa [6], and to different groups of both Eastern or Western Congo Basin hunter-gatherer populations (often historically designated "Pygmies" by Europeans) and their agriculturalist ("non-Pygmy") neighbors today with whom they share complex socio-economic interactions [4,5,38]. Investigating detailed ancient demography and gene-flow among these three groups of populations, taking explicitly into account local differentiations and possible gene-flow among and within groups, is of particular interest as they have previously been identified to be among the most deeply diverged lineages in our species [2,3,29,39,40].

We conducted formal statistical choice of competing scenarios and subsequent joint estimation of parameters under the best scenario using machine-learning Approximate Bayesian Computation (ABC) [41–45], a methodological approach fundamentally differing from all maximum-likelihood methods previously deployed, and only rarely explored in previous studies reconstructing ancient demography in Africa [24]. ABC allows us to compare a range of complex demographic scenarios presumed to have led to observed genetic patterns in numerous population samples, and to estimate posterior parameter distributions best mimicking the data under the best scenario (e.g. [4,18,24]). Briefly, this is achieved based on informative summary-statistics computed on the observed data and on numerous explicit genetic simulations for which scenario-parameters are drawn randomly in large prior distributions set by the user. We thus aimed at jointly inferring competing tree-topologies, divergence times, effective population size changes, and timing, duration and intensity of possible asymmetric gene-flow events, to reconstruct the detailed evolutionary mechanisms underlying the genomic diversity of extant Central and Southern African populations.

<u>Figure1</u>: Sampling location and number of whole-genome sequenced individuals.
Individuals and populations originally sequenced in this work are indicated with filled circles. Individuals and populations
from previously published data merged with the original data set for comparison purposes are indicated with open circles.
Circles' diameters are proportional to the number of individuals as indicated in the top row of the legend in the bottom left
corner. Note that "Ju|'hoansi_comp" refers to the genomic dataset for the Ju|'hoansi previously published by the SGDP and
HGDP and used for comparison to the original data generated here for another sample of the Ju|'hoansi ethno-linguistic group.
Detailed information about population samples is provided in **Table1**, **TableS1**, and **Material and Methods**.

173
174

175 --
176 **Table1: Population table**
177 Geographical locations of population samples are indicated in **Figure1**. Complementary information for genetic-geography
178 and ANOVA analyses are provided in **TableS1**.
179

Population Name[1]	N[2]	Dataset[3]	Sampling Location	Language Family (language)	Close socio-economic interactions with	Samples included in the ROH-ASD-ADMIXTURE analyses (Figure2, Figure3, Figure4)	Samples included in the ABC analyses (Figure5, Figure6, Figure7, Figure8, Figure9)
Original dataset							
Baka	7	this study	Bosquet (Cameroon)	Niger-Congo non Bantu Adamawa Ubangian Gbanzili (Baka)	Nzime	Yes	wRHG, 5 individuals with highest coverage
Nzime	5	this study	Messea (Cameroon)	Bantu A842 (Nzime)	Baka	Yes	RHGn (West)
Ba.Kola	5	this study	Dispersed between Lolodorf and Kribi (Cameroon)	Bantu A80 (Kola)	Ngumba	Yes	wRHG
Ngumba	5	this study	Dispersed between Lolodorf and Kribi (Cameroon)	Bantu A80 (Ngumba)	Ba.Kola	Yes	RHGn (West)
Bi.Aka_Mbati	5	this study	Bombeketi section of Bagandou (Central African Republic)	Bantu C10 (Mbati)	Mbati farmers (not in the dataset)	Yes	wRHG
Ba.Kiga	5	this study	Mukono (Uganda)	Bantu J10 (Kiga)	Ba.Twa	Yes	RHGn (East)
Ba.Twa	6	this study	Kebiremu, Byumba, Kitariro, Mgungu, Nteko (Uganda)	Bantu J11 (Twa)	Ba.Kiga	Yes	eRHG, 5 individuals with highest coverage
Nsua	5	this study	Bundimassoli (Uganda)	Sudanic Mangbutu (Efe)	Ba.Konjo	Yes	eRHG
Ba.Konjo	5	this study	Mulimassenge (Uganda)	Bantu J40 (Konjo)	Nsua	Yes	RHGn (East)
!Xun	5	this study	Omega camp (Namibia) and Schmidtsdrift (Sout Africa)[4]	Khoisan (Ju)	*Not applicable*	Yes	nKS
Ju|'hoansi	5	this study	Tsumkwe (Namibia)	Khoisan (Ju)	*Not applicable*	Yes	nKS
Nama	5	this study	Windhoek (Namibia)	Khoisan (KhoeKhoe)	*Not applicable*	Yes	sKS
Karretjie People	5	this study	Colesberg (South Africa)	Khoisan (Tuu ancestral language) / Indo-European (Afrikaans current language)	*Not applicable*	Yes	sKS
Khutse-San	5	this study	Kutse Game reserve (Botswana)	Khoisan (Khoe Kalahari)	*Not applicable*	Yes	No
Comparative dataset							
Bantu_Herero	2	SGDP	Namibia	Bantu R30 (Herero)	*Not applicable*	Yes	No
Bantu_Kenya	2	SGDP	Kenya	*No data*	*Not applicable*	Yes	No
Bantu_Tswana	2	SGDP	South Africa	Bantu S31 (Tswana)	*Not applicable*	Yes	No
Biaka	2	SGDP	Central African Republic	Bantu C10 (Aka)	*Not applicable*	Yes	No
Coloured	8	SAHGP	South Africa	Indo-European (Afrikaans)	*Not applicable*	Yes	No
Dinka	4	SGDP, HGDP	Sudan	Nilo-Saharan (Dinka)	*Not applicable*	Yes	No
Esan	3	SGDP, KGP	Nigeria	Niger-Congo non Bantu (Esan)	*Not applicable*	Yes	No
Gambian	2	SGDP	Gambia	Niger-Congo non Bantu (Mandinka)	*Not applicable*	Yes	No
Igbo	2	SGDP	Nigeria	Niger-Congo non Bantu (Igbo)	*Not applicable*	Yes	No
Ju|'hoansi_comp	4	SGDP, HGDP	Namibia	Khoisan (Ju)	*Not applicable*	Yes	No
#Khomani	2	SGDP	South Africa	Khoisan (Tuu)	*Not applicable*	Yes	No
Kongo	1	SGDP	Cameroon	*No data*	*Not applicable*	Yes	No
Lemande	2	SGDP	Cameroon	Bantu A46 (Lemande)	*Not applicable*	Yes	No
Luhya	3	SGDP, KGP	Kenya	Bantu JE32 (Luhya)	*Not applicable*	Yes	No
Luo	2	SGDP	Kenya	Nilo-Saharan (Dholuo)	*Not applicable*	Yes	No
Maasai	2	SGDP	Kenya	Nilo-Saharan (Maasai)	*Not applicable*	Yes	No

Mandenka	4	SGDP, HGDP	Senegal	Niger-Congo non Bantu (West Maninkakan)	*Not applicable*	Yes	No
Mandinka	1	KGP	Gambia	Niger-Congo non Bantu (Mandinka)	*Not applicable*	Yes	No
Mbuti	5	SGDP, HGDP	Democratic Republic of Congo	Bantu D30 (Bambuti)	*Not applicable*	Yes	eRHG
Mende	3	SGDP	Sierra Leone	Niger-Congo non Bantu (Mende)	*Not applicable*	Yes	No
Mozabite	2	SGDP	Algeria	Afro-Asiatic (Mozabite)	*Not applicable*	Yes	No
Saharawi	2	SGDP	Western Sahara	Afro-Asiatic (Saharawi)	*Not applicable*	Yes	No
Somali	1	SGDP	Kenya	Afro-Asiatic (Somali)	*Not applicable*	Yes	No
Sotho	7	SAHGP	South Africa	Bantu S30 (Sotho)	*Not applicable*	Yes	No
Xhosa	8	SAHGP	South Africa	Bantu S41 (Xhosa)	*Not applicable*	Yes	No
Yoruba	4	SGDP, HGDP	Nigeria	Niger-Congo non Bantu (Yoruba)	*Not applicable*	Yes	No
Zulu	1	SAHGP	South Africa	Bantu S42 (Zulu)	*Not applicable*	Yes	No
French	4	SGDP, HGDP	France	Indo-European (French)	*Not applicable*	Yes	No
CEU	2	KGP	United States of America	Indo-European (English)	*Not applicable*	Yes	No
Dai	6	SGDP, KGP	China	Tai (Dai)	*Not applicable*	Yes	No
Papuan	6	SGDP, HGDP	Papua New Guinea	Papuan (no data)	*Not applicable*	Yes	No
Karitiana	5	SGDP, HGDP	Brazil	Tupian (Karitiana)	*Not applicable*	Yes	No

[1]Population Names are self-reported for the original dataset presented in this study.
[2]Number of unrelated individuals considered in all analyses in this study (see **Material and Methods**).
[3]SGDP [34]; KGP [35]; HGDP [27,36]; SAHGP [37].
[4]The place of origin of the !Xun is around Menongue in Angola.

Results

We generated high-coverage whole genome sequences (>30X) in 74 individuals from 14 Central and Southern African populations for whom detailed ethno-anthropological information was gathered in the field jointly with DNA samples (**Figure1** and **Table1**). We merged this original dataset with a comparative dataset comprising 105 individuals from 27 African and five non-African populations for whom similar high-quality raw whole genome sequencing data were made available to the community [27,34–37]. After quality-control, variant-calling, and relatedness filtering procedures conducted on all 179 individuals together, we retained 73 and 104 unrelated individuals in our original and comparative datasets, respectively, for all subsequent analyses (see **Material and Methods**).

Considering only the 73 Central and Southern African unrelated individuals newly sequenced here, we identify a total of 26,780,319 biallelic SNPs (241,428 multiallelic SNPs), and 2,454,965 simple insertions/deletions (969,025 complex indels), compared to the reference sequence of the human genome GRCh38 for autosomes (chromosomes 1 to 22), out of which 854,114 (3.1893%) and 114,362 (4.6584%), respectively, were not previously reported in dbSNP 156 (**TableS2**). Among the 177 unrelated worldwide individuals including our original dataset, we identify 36,272,545 biallelic SNPs (257,906 multiallelic SNPs), and 3,159,306 simple insertions/deletions (977,542 complex indels), out of which 1,055,245 (2.9092%) and 189,124 (5.9863%), respectively, were not previously reported in dbSNP 156 (**TableS2**).

Furthermore, we find a large variability in the mean number of biallelic SNPs across populations (**Figure2A, TableS3**). In particular, we find substantially more biallelic SNPs within African populations and more variation of the mean number of SNPs across African populations (from mean=3,726,018; SD=27,793 across 2 individuals in the Saharawi from Western Sahara, to mean=4,628,957; SD=8,139 across 5 individuals in the Ju|'hoansi from Namibia), than within and among non-African populations (from mean=3,214,086; SD=69,241 across 5 individuals in the Karitiana from Brazil, to mean=3,551,792; SD=7,819 across 2 individuals from the USA CEU population). Overall, Southern African Khoe-San populations exhibit the highest mean number of biallelic SNPs as well as numbers of previously unreported SNPs, followed by Central African Rainforest Hunter-Gatherer populations from the Congo Basin, and then by all other African populations in our dataset. These results show that a substantial number of previously unknown variants can still be found when investigating high-quality whole genome sequences from relatively under-studied Sub-Saharan African populations, consistent with previous studies [22,26,27,34–37,40,46].

Figure2: Whole autosomal genome bi-allelic SNPs counts, unbiased heterozygosities, and distributions of Runs of Homozygosity by bins of length, in 37 African and 5 non-African populations.

(**A**) Numbers of bi-allelic SNPs across individuals within populations of more than one individual, compared to the reference sequence of the human genome GRCh38 and to previously reported variants in dbSNP 156. Detailed variant-counts are provided in **TableS2 and TableS3**. (**B**) Unbiased estimates of multi-locus heterozygosities [47], averaged across all variable (bi-allelic SNPs only) and non-variable autosomal sites with no-missing genotype within each population of more than one individual, corrected then for haploid population sample sizes. (**C**) Mean total ROH lengths in four bins of length categories for each population of more than one individual, separately. For each panel, we present a box zooming specifically on the results obtained for the original dataset of 73 unrelated individuals whole-genome sequenced from Central and Southern Africa.

See **Material and Methods** for filtering and calculation details and the software packages used. For (**A**) and (**B**), box-plots indicate the population median in between the first and third quartiles of the box-limits, whiskers extending to data points no more than 1.5 times the interquartile range of the distribution, and empty circles for all more extreme points beyond this limit, if any. Population geographical location, categorization, and descriptions are provided in **Figure1**, **Table1**, and **TableS1**.

----

238 **Figure2**: Whole autosomal genome bi-allelic SNPs counts, unbiased heterozygosities, and distributions of Runs of
239 Homozygosity by bins of length, in 37 African and 5 non-African populations.
240

1. Large genetic diversities and differentiations in Central and Southern Africa

1.A. Heterozygosities and Runs of Homozygosity

We observe significant differences across populations at a local scale in Central and Southern Africa in unbiased heterozygosities [47], calculated on both varying and non-varying autosomal sites (**Figure2A**, **Figure2B**, and **TableS3**. We find overall more variation in Rainforest Hunter-Gatherer (RHG) populations from the western part of the Congo Basin than in RHG populations from the East (Wilcoxon one-sided test p-value=0.00328 for all RHG populations in our dataset and 1.641×10^{-5} considering only those with original data). Note that the Baka and Ba.Kola Western RHG populations exhibited significantly larger genetic diversities than their Nzime and Ngumba RHG neighbors (Wilcoxon one-sided test p-value=0.04637).). Finally, observed genetic variation across Northern Khoe-San (KS) populations (!Xun, Ju|'hoansi, Ju|'hoansi_comp) and Southern KS populations (Nama, Karretjie People and #Khomani) was not significant (Wilcoxon two-sided test p-value=0.7726). Note that unbiased estimates of heterozygosities significantly corelate positively (Spearman rho=0.8490, p-value=2.2×10^{-16}) with genome-wide numbers of autosomal biallelic SNPs (**FigureS1**).

In addition, we investigated the distributions of Runs of Homozygosity (ROH) across individuals within each of the 46 populations in our dataset. In general (**Figure2C**), individuals from Eastern RHG populations (Nsua, Mbuti, Ba.Twa), exhibit the longest total proportion of their autosomal genome in ROH across Sub-Saharan African populations, for all ROH length-classes (Class 0.2-0.5 median total length: 71.28 Mbp; Class 0.5-1: 35.97 Mbp; Class 1-2: 15.79 Mbp; Class 2-4: 9.72 Mbp), with the exception of #Khomani individuals from Southern Africa, and that of Coloured individuals for short ROH only (Class 0.2-0.5 median total length= 98.15 Mbp and 108.18 Mbp, respectively). Interestingly, we find that Western RHG populations (Ba.Kola, Baka, Biaka, Bi.Aka_Mbati) have significantly less total length of ROH of short and medium classes than Eastern RHG (Class 0.2-0.5: Wilcoxon one-sided test p-value= 4.8×10^{-8}; Class 0.5-1: p-value=0.0055; Class 1-2: p-value=0.1284). Furthermore, we find significantly less short ROH (Class 0.2-0.5: Wilcoxon one-sided test p-value=1.4×10^{-5}), and significantly more long ROH (Class 1-2: p-value=2.4×10^{-7}), in Western RHG than in all RHG neighboring populations across the Congo Basin except for the Ba.Kiga RHGn from Uganda. Finally, we find that all Northern and Southern Khoe-San populations have relatively similar total lengths of short ROH compared to that of RHG neighbors (Class 0.2-0.5 Wilcoxon one-sided p-value=0.01675), with the notable exception of the !Xun, who have among the smallest proportion of their genomes in short ROH, worldwide (Class 0.2-0.5 median total length=42.59 Mb). Nevertheless, we find significantly more ROH of longer class in KS populations than in RHG neighbors (Class 1-2: Wilcoxon one-sided p-value=3.7×10^{-5}), a relative pattern similar to what was observed for Western RHG. The distribution of ROH sizes has been shown to be highly informative about important demographic processes such as levels of inbreeding or endogamy within populations, and reproductive isolation or recent admixture among populations [18,48–50]. Thus, our ROH results highlight possibly vast differences in recent demographic processes and levels of endogamy across Central and Southern African populations. Finally, note that total ROH lengths significantly negatively correlated (Spearman rho=-0.5912, p-value=5.243×10^{-5}) with unbiased estimates of heterozygosities (**FigureS1**).

Altogether, our heterozygosity and ROH results show the vast genetic diversity of Sub-Saharan African populations, as well as substantially diverging genomic patterns at a local geographical scale, in particular among and within groups of Rainforest Hunter-Gatherers and RHG neighbors.

1.B. Individual pairwise genetic differentiation

We investigated individual pairwise genomic differentiation across the 177 worldwide unrelated individuals in our dataset, including the 73 unrelated novel Central and Southern African individuals, using Allele-Sharing Dissimilarities (ASD, [51]). The Neighbor Joining Tree (NJT, [52,53]) representation of the pairwise ASD matrix shows three main clusters among all African individuals, separated by a longer genomic distance from all non-African individuals (**Figure3A**, **FigureS2**). One cluster corresponds to all Southern African Khoe-San populations (represented by the pink and red symbols, see **Figure3B**), largely separated from a cluster corresponding to Rainforest Hunter-Gatherer populations from the Congo Basin (represented by the blue symbols); itself also largely separated from a third cluster grouping all other Western, Eastern, Southern and Central African populations in our data set, the latter group including Rainforest Hunter-Gatherer neighboring populations in green symbols (see, **Figure3B)**.

<u>Figure3</u>: **Individual pairwise genetic differentiation patterns.**
Neighbor-Joining Tree (NJT) [52,53] and Multi-Dimensional scaling (MDS) representation of genome-wide Allele Sharing Dissimilarities (ASD) [51] among pairs of individuals at worldwide and regional African scales. We considered the 14,182,615 genome-wide autosomal SNPs pruned for low LD (r^2 threshold 0.1) to calculate ASD between all pairs of individuals. (**A**) NJT computed for all 177 worldwide individuals in the ASD matrix. For easing the visualization of the internal branches of the NJT, all terminal edges are represented in dotted lines each measuring 1/10th of their true size. See **FigureS2** for a MDS version of this ASD matrix. (**B**) Individual symbols and colors identifying the 73 Central and Southern African individuals originally whole-genome sequenced with filled symbols, and the 104 individuals from worldwide populations, including Africa, merged with the original data with open symbols. (**C**) and (**D**) First three axes of ASD-MDS computed on a subset of the full ASD matrix comprising only the individuals sequenced anew, whose symbols are provided in (**B**). (**E**) NJT computed on the same individual subset of the ASD matrix used for (**C**) and (**D**). Terminal edges of this latter tree are represented with dotted lines each measuring 1/30th of their true size.
----

318   **Figure3: Individual pairwise genetic differentiation patterns.**
319

Note, importantly, that individual pairwise dissimilarities at the genome-wide scale do not reflect geography (**Figure3**, **FigureS2**, **TableS4**), highlighting the large genetic differentiation found across African populations and the complexity of its distribution at both a local and continental scale. As can be seen in **Figure3**, from West to East of the Congo Basin throughout Central Africa, RHG populations cluster separately from RHG geographic neighbors with whom they nevertheless share close socio-economic interactions. Furthermore, Central African RHGn are more genetically resembling certain other Western, Eastern, and Southern Africans than their immediate RHG geographical neighbors. Analogously, Southern African Khoe-San populations are also clustering together and much more distant from other geographically neighboring Southern Africans.

Furthermore (**TableS5**, **TableS6**, **TableS7**), ANOVA analyses show that linguistic families, sub-linguistic families, and traditional food-production strategies (**TableS1**), respectively explained a significant proportion of the variance in average population pairwise ASD throughout Africa (partial r^2 = 15.064%, 26.109%, and 9.021%, respectively, each with ANOVA permutation p <0.001), and throughout Sub-Saharan Africa only (partial r^2 = 14.811%, 22.759%, and 10.010%, respectively, each with ANOVA permutation p <0.001). However, most interestingly, we find more complex ANOVA results at a regional scale. Indeed, we find that linguistic families and sub-linguistic families do not significantly explain average population pairwise ASD patterns in Central Africa nor in the Congo Basin, while traditional food-production strategies do so in both regions (**TableS5**, **TableS6**, **TableS7**). Conversely, we find that traditional food-production strategies do not significantly explain population differentiation patterns among Southern African populations, while, here, linguistic and sub-linguistic families do. Finally, and further exemplifying the complexity of genomic differentiation patterns at a local scale in Africa, note that neither linguistic categories nor food-production strategies significantly explain average population pairwise ASD variation among Eastern African populations in our dataset.

The Multidimensional Scaling (MDS) projection of the subsampled ASD matrix for the 73 novel individuals from Central and Southern Africa, further shows substantial differentiation among groups of individuals within each one of the three clusters identified at the continental scale. Indeed, **Figure3C**, **Figure3D** and **Figure3E** show substantial differentiation between Western and Eastern Central African Rainforest Hunter-Gatherer populations, and even substantial differentiation between Ba.Twa and Nsua RHG very locally in Uganda. Conversely, we find relatively much smaller genetic differentiation among pairs of RHG neighbors from West to East of the Congo Basin. Finally, results also show substantial differentiation across Southern African Khoe-San populations, albeit these populations are relatively closer from one another than the different RHG populations.

NJT and MDS based on the ASD pairwise matrix provide an important view of the major axes of genetic differentiation and variation across samples, but do not easily allow to visually describe higher-order axes of genomic variations. To do so, we conducted an ADMIXTURE analysis [54], on the same entire worldwide dataset for increasing values of K (**Figure4**, **FigureS3**). This descriptive method is known to capture the same information as ASD-MDS and ASD-NJT, but allows to explore multiple axes of variation at the same time [18,54–59].

Figure4: Worldwide interindividual genetic diversity patterns with ADMIXTURE.
Each individual is represented by a single vertical line divided in *K* colors each proportional to an individual's genotype
membership proportion assigned by ADMIXTURE [54] into the virtual cluster of that color. Two individuals showing the same
relative proportions of each color at a given value of *K* are genetically more resembling one another than two individuals with
different relative proportions of the same colors. Population and regional geographic groupings are not considered for the
calculations, individuals are *a posteriori* grouped by populations and ordered by the user. Each population is separated by a
thin vertical black line. Unsupervised clustering conducted with 840,031 genome-wide SNPs pruned for LD (r^2 threshold 0.1)
and minor allele frequency (0.1) using ADMIXTURE for values of *K* ranging from 2 to 10, considering 177 worldwide
individuals. For each value of *K* separately, we performed 20 independent runs and used PONG [60], to identify groups of
resembling results (called "modes") based on Symmetric Similarity Coefficient measures of individual results. The number of
runs among the 20 independent runs belonging to the same mode result (i.e. that have pairwise SSC above 0.998), is indicated
below the number of *K* on the left of each barplot separately. Only those runs are averaged per individual to provide the barplot
result for each value of *K* separately. For each value of *K*, all alternative modes identified among the 20 runs are provided in
FigureS3.

380
381 **Figure4: Worldwide interindividual genetic differentiation patterns with ADMIXTURE.**
382
383
384

At K=2 in **Figure4**, individual genotype memberships to the green virtual genetic cluster
are maximized in individuals from Northern KS populations while those for the dark gray
cluster are maximized in non-African individuals, consistent with largest genetic distances in
the ASD-NJT in **Figure3**, with all other individuals in our dataset presenting intermediate,
unresolved, genotype membership proportions to either cluster.
At K=3, a novel red cluster is maximized in all Southern KS populations separating them
from the rest of African individuals, also consistent with the ASD-NJT. At K=4, the novel blue
cluster differentiates Central African RHG populations, and in particular certain Eastern RHG
populations who maximize their membership proportions to this cluster.
The new gray and lighter gray clusters respectively at K=5 and K=6 differentiate Oceanian
Papuan, South American Karitiana, and Western European French and North American USA-
CEU individuals from one another and from all other non-African individuals. These two new
clusters do not substantially affect genotype membership proportions in African individuals,
except for the Southern African Coloured who show substantial membership to the light gray
"European" cluster as well as reduced memberships to this cluster in Nama individuals and one
Xhosa individual.
At K=7, the novel light blue cluster differentiates Eastern and Western RHG populations,
as observed in **Figure3**, with Ba.Twa Eastern RHG showing largely unresolved clustering at
this value of K with intermediate membership proportions to the light blue, the medium blue,
and the green clusters.
At K=8, the novel dark blue cluster is now maximized in Ba.Twa Eastern RHG individuals
who exhibited unresolved clustering at previous values of K. Note that this novel dark blue
cluster is present only to very limited extents in some other RHG and RHGn populations.
At K=9, the novel white cluster is maximized in Dinka populations, located North-East of
Central Africa and the Congo Basin, and substantially represented in other, but not all, East
African populations.
Finally, at K=10, the novel orange cluster differentiates mainly among Northern and
Southern Khoe-San populations, with the Central Khutse San showing substantial membership
proportions to both the red and the orange clusters, respectively maximized in Northern and
Southern KS. Note that, at K=10, the green cluster is maximized in all non RHG and non KS
Sub-Saharan Africa populations except certain East African populations and the Southern
African Coloured population, albeit note that this cluster is not fully resolved at this value of
K since no individual exhibits 100% "green" genotype membership.
In summary, these ADMIXTURE results recapitulate and further dissect the pairwise
genetic differentiation visually observed with ASD-MDS and ASD-NJT in **Figure3**. Most
importantly, note that numerous Central and Southern African individuals retain, from K=2 to
10, intermediate genotype membership proportions in between certain clusters. This should be
interpreted primarily as these individuals being at intermediate genetic distance between
individuals presenting 100% membership to either cluster, which is consistent with ASD-MDS
and ASD-NJT projections (**Figure3**). In turn, such intermediate distances may be either due to
yet-unresolved ADMIXTURE clustering, which may appear at higher values of K, or can be
due to admixture having occurred between ancestors respective to each cluster, the latter
interpretation being likely but not formally tested by the ADMIXTURE method. The possible
occurrence of gene-flow events across pairs of lineages and their influence on tree-topologies
are incorporated explicitly in our Approximate Bayesian Computation (ABC) inferences
detailed below [41–45,61,62].
431
Altogether, our descriptive results highlight the vast genetic diversity and differentiation
of African populations, maximized in certain groups of Central and Southern African
populations, as previously reported (e.g. [17,22,26,27,34,36,37]). Finally, and most importantly, our

results highlight in particular the vast genetic diversity and differentiation of populations at a
local scale in Africa, including among immediate neighbors, not trivially correlated with
geographical distances among populations (**TableS4**), linguistic family classification
(**TableS5**), sub-linguistic family classification (**TableS6**), nor traditional mode of subsistence
(**TableS7**). In fact, these results anticipate that historical and demographic inferences in Africa
may substantially differ when considering different population samples across the continent
and even at a local scale, as well as advocate for extreme caution when grouping individuals
and populations samples into larger categories.

2. Demographic and migration histories in Central and Southern African

2.A. Summary of the Approximate Bayesian Computation inference design
We aimed at reconstructing the demographic and migration history that produced extant
genomic patterns observed within and among Central and Southern African populations
(**Figure2**, **Figure3**, **Figure4**), with machine-learning ABC scenario-choices followed by
posterior-parameter estimations. To do so, we considered eight competing possible tree-
topologies among five extant Northern and Southern Khoe-San, Western and Eastern
Rainforest Hunter-Gatherer, and Rainforest Hunter-Gatherer neighboring populations
(**Figure5**). Furthermore, for each topology, we considered two possible gene-flow processes
among recent and ancient genetic lineages: instantaneous asymmetric gene-flow corresponding
to instantaneous unidirectional introgression events between pairs of lineages; and recurring
asymmetric gene-flow corresponding to unidirectional recurring migrations among pairs of
lineages (scenarios "i" and "r" respectively, **Figure5**). Finally, we considered, for each
topology and each gene-flow process, three nested gene-flow intensities for each event among
pairs of lineages separately: no to very high gene-flow rates (scenarios "i1" and "r1"); no to
moderate rates (scenarios "i2" and "r2"); and no to limited rates (scenarios "i3" and "r3"). This
design allowed us to evaluate fairly the classes of migration-rate scenarios that best explained
the data, without having numerous a priori unrealistic groups of scenarios (e.g. very strong
migration everywhere or no-migration anywhere), directly challenging more intermediate
realistic scenarios.
This led to 8x2x3=48 competing scenarios in total which can be grouped in different ways
to address formally two nested major questions:

i) which ancestral lineages to extant populations diverged first from the others and when did
they do so, when considering complex gene-flow events among recent and ancient lineages?

ii) did gene-flow events occur recurrently or more instantaneously among pairs of lineages
during the evolutionary history of Central and Southern African populations? When did these
events occur? How intense were they?

Importantly, we aimed at considering the vast genetic diversity and differentiation within
and among populations and groups of populations at a local and regional scale (**Figure2**,
Figure3, **Figure4**, and [3,29,30]). Therefore, we replicated the same ABC scenario-choice and
posterior-parameter estimation procedures considering, in turn, 54 different combinations of
five sampled populations of five sampled-individuals using large sets of autosomal genome-
wide summary-statistics (**Figure5**).
We thus conducted a total of 240,000 coalescent simulations under these 48 competing
scenarios (5000 simulations per scenario), by drawing parameter values from prior
distributions set by the user (**Figure5**, **TableS8**). For each simulation, we then calculated a
vector of 337 summary-statistics (**TableS9**), within and among the five simulated populations;

each vector of summary-statistics thus corresponds to a vector of parameter values drawn randomly from prior distributions and used for one simulation. We then employed Random Forest ABC procedures [45,63], to identify the best scenario or group of scenarios for each 54 combinations of five sampled "real" populations separately. Finally, under the best scenario, we produced 100,000 simulations, computed 202 summary-statistics for each simulation (**TableS9**), and performed Neural Network ABC posterior parameter joint-estimation [43,44], providing posterior-parameter distributions most likely to have produced observed genomic data, for each scenario-parameter.

Figure5: 48 competing scenarios for the history of Central and Southern African populations.

(**A**) Eight competing topologies for the demographic history of five Central and Southern African extant lineages. The Northern Khoe-San lineage (nKS) is represented by either the Ju|'hoansi or the !Xun populations. The Southern Khoe-San lineage (sKS) is represented by either the Karretjie People or the Nama populations. The Rainforest Hunter-Gatherer Neighbors lineage (RHGn) is represented either by the Western Congo Basin Nzime or Ngumba populations, or the Eastern Ba.Konjo or Ba.Kiga populations. The Western Rainforest Hunter-Gatherer lineage (wRHG) is represented by either the Baka, the Ba.Kola, or the Bi.Aka_Mbati populations. The Eastern Rainforest Hunter-Gatherer lineage (eRHG) is represented by either the Nsua, the Ba.Twa, or the Mbuti populations. As indicated in legend in the bottom right corner of the panel, the eight topologies can be grouped according to i) which ancestral lineage (Ancestral RHG or Ancestral KS) diverged first from the two others, or ii) whether the nKS and sKS lineages split earlier or later than the split of the wRHG and eRHG lineages. For each topology, we consider possible changes in lineages' constant effective population size, N, after each divergence event, t. For each topology, we considered possible gene-flow events among ancestral lineages and among recent lineages, represented as horizontal uni-directional arrows (forward in time). (**B**) Example for a given topology (1a), of the fact that each one of the eight topologies are considered under two alternative competing "instantaneous" (Scenario i-1a) or "recurring" (Scenario r-1a) gene-flow processes. Instantaneous gene-flow processes consider that genes can be exchanged among pairs of ancestral or recent lineages at a single parameterized time, tad, with independent parameters of gene-flow intensity, m, from lineage "A" to "B" and from lineage "B" to "A". Recurring gene-flow processes consider instead that gene-flow occur at each generation between pairs of lineages with a constant independent rate, m, from lineage "A" to "B" and from lineage "B" to "A". Note that for all eight topologies and for both instantaneous and recurring gene-flow processes, we parameterized separately the gene-flow from lineage "A" into lineage "B", and the gene-flow from lineage "B" into "A", in order to allow for possibly asymmetrical gene-flow events among pairs of lineages. For both instantaneous and recurring gene-flow processes, we considered three nested intensities of gene-flow parameters indicated at the bottom of (**B**). Scenarios i1 or r1 consider that the gene-flow parameters m in each topology are independently drawn in Uniform distributions between 0 and 1 or 0 and 0.05, respectively, thus allowing for the possibility of no to high gene-flows from one lineage to the other. Scenarios i2 or r2 consider that the gene-flow parameters m in each topology are independently drawn in Uniform distributions between 0 and 0.25 or 0 and 0.0125, respectively, thus allowing only for the possibility of no to intermediate gene-flow from one lineage to the other. Finally, scenarios i3 or r3 consider that the gene-flow parameters m in each topology are independently drawn in Uniform distributions between 0 and 0.01 or 0 and 0.001, respectively, thus only allowing for the possibility of no to reduced gene-flow from one lineage to the other. Altogether we thus considered 8x2x3=48 competing scenarios for the demographic and migration history of Central and Southern African populations inferred with machine-learning Approximate Bayesian Computation. We conducted Random-Forest ABC scenario choice procedures among groups of these 48 scenarios in turn for 54 different sets of five observed populations each, represented as grey lines between population names, each population comprising five whole-genome sequenced individuals. Sampled-population names considered in the 54 combinations are given below each tree-leave, and combinations are limited to those pairing one RHGn with the specific Eastern or Western RHG population with whom they share complex socio-economic interactions, as indicated in **Table1**. See the detailed description of scenarios and their grouping, their parameters and their respective prior distributions used for ABC inference in **Material and Methods** and **TableS8**.

----

534 **Figure5: 48 competing scenarios for the history of Central and Southern African populations.**
535
536

2.B. Are summary-statistics representative of large portions of the genome and are we able to mimic observed data with simulations under the 48 competing scenarios?

We first empirically checked that the observed data for each one of the 54 combinations of five sampled populations used for all ABC inferences hereafter –which considered 100 high-quality 1 Mb-long genomic windows pruned from the full genomes in our working dataset (see **Material and Methods**)– reasonably echoed the statistical descriptions obtained above from our full genomic SNP set. We thus compared standardized individual pairwise ASD matrices for each one of the 54 sets of population-combinations with the corresponding individual subsets of the standardized ASD matrix used for MDS and NJT analyses in **Figure3**. We found that each 54 sets of population-combinations and associated pruned SNP sets for ABC inferences highly resembled the genomic individual pairwise differentiation patterns reported above using the entire SNP set. Indeed, we found significant (1000 Mantel permutation two-sided p < 0.002), Pearson r correlations ranging from 0.945 to 0.974 across the 54 sets of five population combinations (**TableS10**).

Furthermore, we compared entire vectors of summary-statistics computed on the 100 1Mb windows for each 54 combinations of sampled-populations separately, with the same summary-statistics computed on 724 other 1Mb such genomic regions throughout the genome. We found that the 100 windows used for ABC inferences hereafter provided summary-statistics not significantly different from- (Wilcoxon two-sided p-values between 0.6497 and 0.9995 across 54 combinations of five sampled populations), and highly correlated with- (Spearman rho between 0.8566 and 0.99637, all p-values <2.2x10^{-16}), those computed on much larger portions of the genome (**TableS11**).

In order to conduct ABC inferences, we first checked whether we were able to simulate data, under the 48 competing scenarios, for which vectors of summary statistics could mimic those obtained separately from the 54 different combinations of five sampled populations each.

We first conducted goodness-of-fit permutation tests, separately for the 54 combinations of five populations, and found that the observed vectors of 337 summary statistics were never significantly different (54 goodness-of-fit permutation p-values > 0.56), from the 240,000 vectors of 337 summary statistics computed from simulations under the 48 competing scenarios. Second, we computed a two-dimensional PCA on the vectors of summary-statistics obtained from simulations on which we projected, separately, the 54 vectors of these summary-statistics computed on observed data, and found that observations each fell well within the space of simulations (**FigureS4**).

Both results show that we were able to simulate data for which summary-statistics were reasonably close to the observed ones, at least in parts of the parameter space used for simulations from the 48 scenarios and for each one of the 54 separate combinations of five sampled populations. We could therefore conclude that the observed patterns of variation could be generated under some of the 48 scenarios, and we proceeded with Random Forest ABC (RF-ABC) scenario-choice and Neural Network ABC (NN-ABC) posterior-parameter estimation procedures.

Note that the prior error in RF-ABC scenario-choice, using simulations as pseudo-observed data and considering 5000 simulations under each one of the 48 scenarios in competition, converged towards a minimum when using 1000 trees for all RF-ABC scenario-choice analyses conducted hereafter (**FigureS5**).

Importantly, the *a priori* discriminatory powers of the RF, estimated as cross-validation procedures based on all simulations used in-turn as pseudo-observed data, were overall good across all analyses described hereafter, despite increased confusion among scenarios based on gene-flow intensity within groups of instantaneous or recurring processes (**FigureS6, FigureS7, FigureS8, FigureS9**). This was expected as classes of gene-flow intensities distinguishing among scenarios with, respectively, no-to-low, no-to-moderate, or no-to-high

gene-flow intensities, were designed to have nested parameter priors (**Figure5**, **TableS8**). Note
that this confusion among gene-flow intensities due to nestedness was more acute for recurring
gene-flow scenarios than for instantaneous gene-flow scenarios (**FigureS8**).
Finally, we evaluated separately for each analysis how each one of the 337 summary-
statistics (**TableS9**), contributed *a priori* to the RF-ABC scenario-choice in the entire space of
simulations, by considering each simulation "out-of-bag" as pseudo-observed data and without
considering observed data. Most interestingly, we found that individual-pairwise ASD-MDS
summary-statistics describing how individuals are genetically differing from one another
within and among populations and population-pairwise F_{ST}, in particular, were always among
the 50 most informative summary-statistics for RF-ABC scenario-choice *a priori*.
Alternatively, note that we found that all statistics summarizing the 1D unfolded standardized
Site Frequency Spectrum considered here were never among the 50 most informative statistics
a priori using the Gini index of classification, for any of the RF-ABC scenario-choice analyses
(**FigureS10**, **FigureS11**, **FigureS12**, **FigureS13**, **FigureS14**, **FigureS15**, **FigureS16**,
FigureS17, **FigureS18**, **FigureS19**).
602
2.C. Which historic scenario best explains genomic patterns in extant African
populations?
2.C.1. Instantaneous or recurring gene-flows of which intensity among African lineages?
Figure6A shows that gene-flow processes among recent or ancient genetic lineages most likely
occurred during very limited "instantaneous" periods of time, rather than recurrently,
throughout the entire evolutionary history of Central and Southern African populations. Indeed,
for each one of the 54 different combinations of five sampled populations, the group of 24
scenarios considering instantaneous gene-flow processes provides vectors of summary
statistics systematically closest to the observed ones, whichever the tree-topology or intensity
of the gene-flow considered. Note that we find that posterior probabilities of choosing the
correct group of scenarios for each 54 sets of population combinations range from 0.97828 to
0.99510, compared to a prior probability of 1/2 in this particular test.
Furthermore, we find that we imperatively need these gene-flow processes to be
potentially high in order to best explain the observed data for each one of the 54 combinations
of population samples, whichever the tree-topology and gene-flow process (**Figure6B**);
whichever the tree-topology but considering gene-flow processes separately (**Figure6C**); or
even considering 24 competing scenarios under instantaneous gene-flow processes only,
whichever the tree-topology (**FigureS7**). Note that, for these three particular tests, we find that
posterior probabilities of choosing the correct group of scenarios [64], for each 54 sets of
population combinations range, respectively, from 0.66318 to 0.87138 (**Figure6B**), from
0.67765 to 0.83957 (**Figure6C**), and from 0.63935 to 0.87000 (**FigureS7**), compared to a prior
probability of 1/3, 1/6, and 1/3 in these three particular tests, respectively.
Finally, note that when considering the 48 competing-scenarios separately, a very
challenging task *a priori* given the number of competing scenarios and the high level of
nestedness among scenarios, the best scenarios predicted by the RF-ABC procedure is, among
the 54 population-combination tests conducted, systematically found for scenarios considering
an instantaneous gene-flow process allowing for possibly highly intense gene-flow rates
(**FigureS8, FigureS9**).
631

632 --

Figure6: Random-Forest ABC scenario-choices among 48 competing scenarios.
Random Forest Approximate Bayesian Computation scenario-choice [45,63], are conducted, for each analysis in each panel, separately for 54 different combinations of five Central and Southern African sampled populations each. Posterior proportions of votes obtained for each 54 sampled-populations combinations, and for RF-ABC analysis in each panel respectively, are provided as box-plots indicating the median in between the first and third quartile of the box-limits, whiskers extending to data points no more than 1.5 times the interquartile range of the distribution, and empty circles for all more extreme points beyond this limit, if any. In each RF-ABC analysis in each panel, proportion of votes (indicated in the y-axis) are obtained using 1,000 decision trees in the random forest for each group of competing scenarios (indicated in the x-axis table and labels), using 5,000 simulations per scenario and 337 summary-statistics (see **Material and Methods** and **TableS9** for details). Detailed competing-scenarios grouped in each group are indicated in the two bottom lines of the x-axis table, with topology and gene-flow processes codes provided in **Figure5** and explained in detail in **Material and Methods**. The top-line of the x-axis table indicates, for each group of competing scenarios in each analysis, the number of times the corresponding group of scenarios was predicted as the best one among the 54 different combinations of five sampled-populations. (**A**) Scenario-choice results for the two groups of instantaneous or recurring gene-flow processes (24 scenarios and 120,000 simulations in each two competing groups), all gene-flow intensities and all topologies "being equal". Posterior probabilities of choosing the correct group of scenarios [64], for each 54 sets of population combinations ranged from 0.97828 to 0.99510, compared to a prior probability of 1/2 in this particular test. (**B**) Scenario-choice results for the three groups of intensities for gene-flow processes (16 scenarios and 80,000 simulations in each three competing groups), all topologies and both instantaneous and recurring gene-flow processes "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.66318 to 0.87138, compared to a prior probability of 1/3 in this particular test. (**C**) Scenario-choice results for the six groups of gene-flow processes and intensities separately (8 scenarios and 40,000 simulations in each six competing groups), all topologies "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.67765 to 0.83957, compared to a prior probability of 1/6 in this particular test. (**D**) Scenario-choice results for the three groups of scenario topologies differing in the order of ancient lineages divergence (differing number of scenarios in each three competing groups are evened by randomly sampling the same number of simulations, equal to 60,000, in each group, see **Material and Methods**), all gene-flow processes and intensities "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.39862 to 0.48967, compared to a prior probability of 1/3 in this particular test. (**E**) Scenario-choice results for the same test as (**D**) restricted to the 24 competing scenarios considering instantaneous gene-flow processes only (differing number of scenarios in each three competing groups are evened by randomly sampling the same number of simulations, equal to 30,000, in each group, see **Material and Methods**), all gene-flow intensities "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.43333 to 0.51857, compared to a prior probability of 1/3 in this particular test. (**F**) Scenario-choice results for the two groups of scenario topologies differing in the relative order of divergences between Northern and Southern KS, and Western and Eastern RHG lineages, respectively (24 competing scenarios and 120,000 simulations in each two groups), all gene-flow processes and intensities "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.62115 to 0.72015, compared to a prior probability of 1/2 in this particular test. (**G**) Scenario-choice results for the same test as (**F**) restricted to the 24 competing scenarios considering instantaneous gene-flow processes only (12 competing scenarios and 60,000 simulations in each two groups), all gene-flow intensities "being equal". Posterior probabilities of choosing the correct group of scenarios for each 54 sets of population combinations ranged from 0.62033 to 0.71883, compared to a prior probability of 1/2 in this particular test.

674 --

677
678
679

Figure6: Random-Forest ABC scenario-choices among 48 competing scenarios.

2.C.2. Ancient tree-topology among African lineages?

Figure6D shows that, overall, the RF-ABC scenario-choice procedures vote in favor of an
ancient tree-topology where the ancestral Khoe-San lineage diverged first followed by the
divergence between the ancient Rainforest Hunter-Gatherer lineage and that of their neighbors
(Scenarios tree-topologies 1a-1b-1c, **Figure5**), in a majority (35/54) of the 54 combinations of
observed populations tested here, whichever the gene-flow process and intensity. Note that we
find that posterior probabilities of choosing the correct group of scenarios for each 54 sets of
population combinations range from 0.39862 to 0.48967, compared to a prior probability of
1/3 in this particular test (**Figure6**).

Furthermore, we find that this majority of votes is much larger (50/54), when considering
instantaneous gene-flow processes only, whichever their intensities (**Figure6E**). Note that we
find that posterior probabilities of choosing the correct group of scenarios for each 54 sets of
population combinations range from 0.43333 to 0.51857, compared to a prior probability of
1/3 in this particular test (**Figure6**).

Finally, note also that the tree-topologies where the RHGn lineage diverge first from the
lineage ancestral to that of the RHG and the KS (Scenarios tree-topologies 3a-3b, **Figure5**),
are predicted to be the best ones in a minority of the tests (19 out of 54 combinations of sampled
populations, **Figure6D**), whichever the gene-flow process and/or intensity. Furthermore, they
do so in an even smaller minority (4/54) for instantaneous gene-flow processes only, whichever
the intensities (**Figure6E**). Interestingly, we found that tree-topologies where the ancient
lineage for RHG populations diverges first from the two others (Scenarios tree-topologies 2a-
2b-2c, **Figure5**), is never favored in our analyses, whichever the gene-flow process and/or
intensity.

2.C.3. Recent tree-topology among African lineages?

Finally, **Figure6** shows that Northern and Southern Khoe-San extant lineages likely
diverged from one-another before the divergence between Eastern and Western Rainforest
Hunter-Gatherer lineages (Scenarios tree-topologies 1b-1c-2b-3b, **Figure5**), whichever the
ancient tree-topology and gene-flow process and intensity (**Figure6F**), and whichever the
ancient tree-topology for instantaneous gene-flow processes only, all classes of gene-flow
intensities "being equal" (**Figure6G**). Interestingly, the "b" group of tree-topologies is found
to best-describe observed summary-statistics for 34 out of 54 combinations of sampled
populations and this majority of RF-ABC votes is even larger (49/54) when considering only
the group of 24 competing scenarios for which gene-flows are instantaneous rather than
recurring. Note that, for these two particular tests, we find that posterior probabilities of
choosing the correct group of scenarios for each 54 sets of population combinations range,
respectively, from 0.62115 to 0.72015 (**Figure6F**), and from 0.62033 to 0.71883 (**Figure6G**),
compared to a prior probability of 1/2 in both these two particular tests.

2.C.4. Conclusion: intersecting scenario-choice results for the history of Africa.

Altogether, when intersecting RF-ABC scenario-choice inferences in groups of gene-flow
processes and tree-topologies (**Figure6**), with inferences considering all competing-scenarios
separately (**FigureS8, FigureS9**), and among 54 combinations of five sampled populations,
we find that Scenario i1-1b (**Figure5**), systematically best-describes the observed data in all
conformations of scenario-choice tests for 25 combinations of populations (**TableS12**), while
the second scenario best explaining the data (Scenario i1-3b) does so consistently across all
tests for only four combinations of population out of 54.

Therefore, our results point to a scenario for the history of Central and Southern African
populations where the ancestral KS lineage diverged first, followed by the divergence between
the ancestral RHG and the RHGn lineages, with a divergence between the extant Northern and

Southern KS lineages occurring independently before that of the extant Eastern and Western
RHG lineages. Most importantly, the best scenario necessarily involves gene-flow events
among all pairs of lineages throughout history which occurred during relatively short
"instantaneous" periods of time, rather than recurrently over longer periods of time.
Furthermore, these instantaneous gene-flow events must each have been possibly of high
intensity, rather than all limited to moderate or low intensities, and possibly asymmetrical
between pairs of lineages, to best explain the observed data. Finally, our results also highlight
that considering different combinations of extant populations separately in the analyses may
provide contrasted results where alternative evolutionary scenarios may best explain
observations.
For conservativeness and to further consider the genetic diversity of extant Central and
Southern African populations at a local or regional scale, we will henceforth conduct all Neural
Network ABC posterior-parameter estimations separately for the 25 combinations of sampled
populations systematically providing the Scenario i1-1b as the most-likely one among the 48
competing scenarios. **TableS12** shows no specific statistical or anthropological pattern among
the specific sampled-populations in the 25 combinations providing consistent results across
RF-ABC analyses. Indeed, the least represented sampled-population, the Ngumba RHGn,
appears in 20% of the combinations (four RHGn sampled populations are considered in turn);
the most represented sampled-population, the Ju|'hoansi nKS, appears in 76% of the
combinations (two nKS sampled populations are considered in turn); and all 14 sampled-
populations considered in each RF-ABC scenario-choice analysis are present in the 25
combinations.
752
2.D. Which scenario-parameters best explain genomic patterns in extant African
populations?
2.D.1. Divergence times.
Divergence times are most often well estimated by our Neural Network ABC posterior
parameter inferences (**Figure7** and **TableS13**), with relatively narrow 90% Credibility
Intervals (CI), posterior distributions very often largely departing from the priors, and low
cross-validation posterior parameter estimation errors in the vicinity of the observed data
(**TableS14**). Interestingly, we find relatively consistent posterior estimates across the 25
different combinations of five populations for, separately, the Eastern and Western RHG
divergence (t_{RHG}), between 355 generations before present (gbp) (mode point estimate,
90%CI=[203-1,220]) and 1,518 gbp (90%CI=[805-3,016], **TableS13**). We find divergences
between the Northern and Southern KS (t_{KS}), between 2.1 and 8.0 times older than the
corresponding t_{RHG} for the 25 combinations of sampled-populations. These correspond in
absolute to t_{KS} between 1,882 gbp (mode point estimate, 90%CI=[1,114-3,270]) and 3,823 gbp
(90%CI=[2,336-5,669], **TableS13**). Combining (concatenating) the posterior distributions for
these two parameters (**Figure7A, Figure7B**, **TableS15**), respectively, provides, synthetically,
an absolute modal point-estimate divergence time between eRHG and wRHG of 751 gbp
(90%CI =[370-2,222]), and one between nKS and sKS, largely more ancient, having occurred
some 2,857 gbp (90%CI=[1,624-4,662]).
In a more remote past, the divergence time between the lineage ancestral to all RHG
populations and the RHG neighbors' lineage is also most often well estimated, based on CI-
width and departure from prior distributions, and cross-validation posterior parameter
estimation errors in the vicinity of the observed data, for each 25 combinations of populations
respectively, albeit results are more variable across sets of population combinations than for
more recent divergence times (**Figure7C**, **TableS13, TableS14**). The ancient RHG and RHGn
lineages' divergences ($t_{RHG-RHGn}$) are between 4.5 and 22.3 times older than the corresponding
t_{RHG} for the 25 combination of sampled-populations. This translates in absolute between 3,595

gbp (mode point estimate, 90%CI=[2,571-7,368]) and 8,199 gbp (90%CI =[6,083-10,682], **TableS13**). Synthetically combining results together provides an absolute divergence time between ancestors of extant RHGn and extant RHG of 6,347 gbp (90%CI=[4,098-9,734], **Figure7C** and **TableS15**).

Finally, we also find relatively variable posterior estimates for the most ancient divergence time in our tree-topology, between the ancestral lineages to all extant KS lineages and the ancestral lineages to all RHG and the RHG neighbors; with, again, posterior-parameter distributions across 25 combinations of sampled populations very often satisfactorily estimated, with relatively reduced 90%CI and substantial departure from the priors, and relatively low cross-validation posterior parameter estimation errors in the vicinity of the observed data (**Figure7D**, **TableS13**, **TableS14**). We find the original most ancient divergence ($t_{KS-RHG-RHGn}$) in our tree-topology between 8.6 and 33.6 times older than the corresponding most recent divergence t_{RHG} for the 25 combinations of sampled-populations. This translates in absolute to a modal ancestral divergence among all investigated Central and Southern African lineages that occurred some 13,254 gbp (90%CI=[8,044-14,397]), when all results are combined (concatenated) together synthetically (**Figure7D** and **TableS15**).

Figure7: Neural Network ABC posterior parameter distributions of divergence times.

Neural Network Approximate Bayesian Computation posterior parameter joint estimations [43,44], of topological divergence-times (in generations before present) for 25 sets of five Central and Southern African populations for which the best scenario identified by RF-ABC was Scenario i1-1b (**Figure5**). NN ABC posterior parameter estimation procedures were conducted using 100,000 simulations under Scenario i1-1b, each simulation corresponding to a single vector of parameter values drawn randomly from prior distributions provided in **TableS8**. We considered 43 neurons in the hidden layer of the NN and a tolerance level of 0.01, corresponding to the 1,000 simulations providing summary-statistics closest to the observed one. NN posterior estimates are based on the logit transformation of parameter values using an Epanechnikov kernel between the corresponding parameter's prior bounds (see **Material and Methods** and **TableS8**). Posterior parameter densities are represented with solid blue lines. 50% Credibility Intervals are represented as the light blue area under the density. The median and mode values are represented as a solid and dotted blue vertical line, respectively. Parameter prior distributions are represented as dotted gray lines. For all panels, the left plots represent the NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern African populations best explained under Scenario i1-1b, separately (**TableS12** and **TableS13**). For all panels, the right plots represent a single parameter posterior distribution obtained from combining (concatenating) the 25 posterior distributions together. (**A**) Results for parameter t_{RHG} corresponding to the split time between the Western and Eastern Rainforest Hunter-Gather (RHG) lineages (**Figure5**). (**B**) Results for parameter t_{KS} corresponding to the split time between Northern Khoe-San (nKS) and Southern Khoe-San (sKS) lineages (**Figure5**). (**C**) Results for parameter $t_{RHG-RHGn}$ corresponding to the split time between the Rainforest Hunter-Gather neighboring population lineage (RHGn) and the lineage ancestral to Western and Eastern RHG (**Figure5**). (**D**) Results for parameter $t_{KS-RHG-RHGn}$ corresponding to the split time between the lineage ancestral to KS populations and the lineage ancestral to all RHG and RHGn lineages; this event thus corresponding to the split time of all lineages in the history of Central and Southern African populations (**Figure5**). Results for all left panels are summarized in **TableS15** and cross-validation posterior parameter estimation errors in the vicinity of the observed data in **TableS14**.

Figure7: Neural Network ABC posterior parameter distributions of divergence times.

2.D.2. Effective population sizes.
Effective population sizes (Ne) for all recent Northern and Southern KS, Western and Eastern
RHG, and RHG neighbors' lineages are often reasonably well estimated with relatively
reduced 90%CI, substantial departure from their priors, and low cross-validation posterior
parameter estimation errors in the vicinity of the observed data (**TableS14**), for each 25 sets of
sampled population combinations (**FigureS20**, **TableS13**). Combining posterior distributions
among the 25 separate tests, we find Ne posterior estimates ranging from modal point-estimates
of 10,702 diploid effective individuals (90%CI=[5,769-67,086]) in the eRHG to 17,083
(90%CI=[7,606-71,133]) in the wRHG extant lineages (**FigureS20**, **TableS15**). Importantly,
we were unable to satisfactorily estimate, for almost all 25 combinations of sampled
populations, the three effective population sizes for, respectively, the ancient KS lineage, the
ancient RHG lineage, and the lineage ancestral to all RHG and RHG neighbors' extant lineages
(**TableS15**, **FigureS20**, **TableS13**, **TableS14**).
Conversely, our posterior estimates of the effective size of the lineage most ancestral to all
our populations was satisfactorily estimated, with relatively narrow 90%CI, substantial
departure from the prior distributions, and among the lowest cross-validation posterior
parameter estimation errors in the vicinity of the observed data (**TableS14**), for almost all 25
sets of population combinations. Concatenating posterior distributions, despite a noteworthy
variation across the 25 tests (**FigureS20**, **TableS13**), we find an ancestral effective size for the
lineage ancestral to all our Central and Southern African extant populations of 17,005 diploid
effective individuals (90%CI=[9,890-69,775]), the second largest posterior estimate across all
recent and ancient lineages in our analyses (**TableS15**). Note that we considered large priors
(Uniform[10-100,000] diploid individuals) for constant effective population sizes in all
lineages with possible changes at each divergence time, for simplicity. Therefore, the posterior
estimates here found may be different in future procedures considering more complex effective
demographic regimes likely to have occurred in certain lineages, such as possible bottlenecks
and/or population expansions (e.g. [8,40,65]).
855
2.D.3. Instantaneous gene-flow times.
In between each lineage divergence times, we estimated the time of occurrence of
instantaneous and potentially asymmetric gene-flow exchanges across pairs of lineages
(**Figure5**), for each 25 combinations of sampled populations separately. For four out of the six
recent gene-flow times between pairs of extant lineages ($tad_{nKS-sKS}$, $tad_{wRHG-eRHG}$, $tad_{wRHG-RHGn}$,
$tad_{eRHG-RHGn}$, **Figure5**, **TableS8**), we find relatively consistent posterior estimates across the
25 tests, almost all with relatively reduced 90%CI, substantial departure from the priors, and
low cross-validation posterior parameter estimation errors in the vicinity of the observed data
(**Figure8A – Figure8D**, **TableS13**, **TableS14**, **TableS15**). Note that, while we also obtain
similarly satisfactory posterior distributions among the 25 tests for gene-flows among Northern
KS and RHGn lineages ($tad_{nKS-RHGn}$), and among Southern KS and RHGn lineages ($tad_{sKS-RHGn}$), respectively, we found much larger variance across the 25 different population sets
(**Figure8E – Figure8F**, **TableS13**).
Interestingly, we also obtain satisfactory posterior estimates of gene-flow timing among
ancient lineages for almost all 25 sets of population combinations ($tad_{ARHG-RHGn}$, $tad_{AKS-RHGn}$,
$tad_{AKS-ARHG}$, $tad_{AKS-ARHG-RHGn}$, **Figure5**, **TableS8**). Nevertheless, note that posterior estimates
are increasingly variable from one set of sampled populations to the other as the estimated
parameter is further back in time in the tree-topology (**Figure8G – Figure8J**, **TableS15**).
874

<u>**Figure8**</u>**: ABC posterior parameter distribution of gene-flow times.**
Neural Network Approximate Bayesian Computation [43,44], posterior parameter joint estimations of gene-flow instantaneous
times *tad* (in generations before present) for 25 sets of five Central and Southern African populations for which the best
scenario identified by RF-ABC was Scenario i1-1b (**Figure5**, **TableS12**). Methodological details are provided in **Figure7**
caption and detailed in **Material and Methods**.

2.D.4. Gene-flow intensities.
We obtain overall unsatisfactory posterior estimates of the 20 separate unidirectional gene-
flow rates for the 10 separate instantaneous gene-flow events, for a majority of the 25 sets of
population combinations (**FigureS21, TableS13**), with relatively large 90%CI, reduced
posterior distributions' departure from priors, and high cross-validation posterior parameter
estimation errors in the vicinity of the observed data (**TableS14**). Considering only the several
satisfactory posterior estimates for each gene-flow parameter separately, we notice their very
large variability across the sets of population combinations for each parameter, making it
unreasonable to combine posterior distributions to obtain a synthetic value (**TableS15**). The
strong parameter-estimation limitation encountered here is extensively discussed in the
Discussion section below.
895
2.D.5. A synthesis of the demographic history of Central and Southern African populations.
We propose, in **Figure9**, a schematic synthesis of all the above inference results for the
demographic history of 25 out of 54 combinations of five Central and Southern African
populations, for which all our RF-ABC scenario-choice procedures provided systematically
consistent results. **Figure9** synthesizes mode estimates of all reliably inferred parameters
obtained from the combined (concatenated) posterior distributions for the 25 combinations of
sampled populations provided in **TableS15** and presented in **Figure7**, **Figure8**, **FigureS20**,
and **FigureS21**.
Posterior checks show that 25 separate pseudo-observed simulations computed under the
mode point-estimates for this best synthetic demographic scenario are reasonably resembling
those from the 25 observed combinations of five sampled populations in the first two axes of a
PCA computed on the 202 summary-statistics used for all NN-ABC inferences (**FigureS22**).
Furthermore, **FigureS22** shows that, as expected, most variance in summary-statistics is
observed among the 25 observed combinations of sampled populations and that the 25 pseudo-
observed simulations provided relatively less variable vectors of 202 summary-statistics.
This shows that, while the synthetic demographic history proposed here results from
artificially combining the diversity of demographic histories inferred separately for each
combination of sampled-populations, this synthesis provides reasonably similar genomic
patterns as those observed in real data.
We conducted NN-ABC posterior-parameter estimations as for the observed datasets on
the 25 pseudo-observed simulations. We find substantially more variance in combined
posterior-parameter estimates for most parameters that were reasonably accurately estimated
across the 25 combinations of five Central and Southern African sampled-populations,
compared to the variance of combined (concatenated) posterior-parameter distributions
obtained for 25 pseudo-observed simulations under the proposed synthetic demographic
scenario (**FigureS23**, **FigureS24**, **FigureS25**, **FigureS26**, and **TableS16**). Indeed, **TableS16**
shows that 15 out of the 20 demographic parameters posterior combined distributions had a
significantly larger variance with observed datasets than with pseudo-observed simulations;
and an additional two parameters exhibit the same pattern but only marginally significant.
In particular (**TableS16**), the divergence times obtained with observed data are between
22% (t_{RHG}; Fligner test p-value=5.46×10^{-146}) and 154% ($t_{KS-RHG-RHGn}$; Fligner test p-value=0),
more variable than those obtained with pseudo-observed simulations. Furthermore, the
admixture times obtained with observed data are between 16% ($tad_{AKS-RHGn}$; Fligner test p-
value=4.67×10^{-25}) and 211% ($tad_{eRHG-RHGn}$; Fligner test p-value=0), more variable than those
obtained with pseudo-observed simulations, excluding the two parameters with the marginally
significant patterns.
Altogether, these results demonstrate empirically that the large variation in divergence and
instantaneous gene-flow time posterior estimates obtained across combinations of sampled

populations largely depend on the specific samples considered when investigating highly
differentiated Sub-Saharan populations, rather than on variation in inference performances.
Our results thus explicitly advocate for caution when summarizing results obtained across
population sets and further likely explain the, sometimes, apparent discrepancies that arise
across studies considering different sampling designs prior to conducting demographic
evolutionary reconstructions (e.g. [22,26]).
940
941

Figure9: Schematic inferred demographic and migration history of Central and Southern African populations.
Schematic representation of Scenario i1-1b (**Figure5**), and Neural Network ABC posterior parameter mode estimates summarizing results obtained separately for 25 sets of five Central and Southern African populations (**TableS12**), represented by the gray lines in between population names. Gene-flow arrows are indicated forward in time. For the time of each divergence and gene-flow event, mode point estimates are provided in generations before present (gbp) in bold, and 90% Credibility Intervals are provided between parentheses (**TableS15**). We provide two estimates of the divergence times estimates in years before present (ybp), one (upper) corresponding to 30 years per generation and the other (lower) to 20 years per generation. Mode point estimates of effective population sizes *Ne* are provided in numbers of diploid effective individuals and width of lineages are proportional to the estimated *Ne* (**TableS15**, **FigureS20**). NN-ABC posterior distributions for the effective population sizes of the ancestral Khoe-San lineage (AKS), for the ancestral RHG lineage (ARHG), and for the lineage ancestral to RHGn and ARHG, were all three poorly distinguished from their respective prior distributions and with high cross-validation posterior parameter estimation errors (**FigureS20**, **TableS14**, **TableS15**), as indicated by the question marks. NN-ABC posterior distributions for instantaneous asymmetric gene-flow rates were overall poorly departing from their priors and with high cross-validation posterior parameter estimation errors (**FigureS21**, **TableS13**, **TableS14**), as indicated by the doted arrows. All posterior distributions are shown graphically in **Figure7**, **Figure8, FigureS20, FigureS21**, and detailed in **TableS13** and **TableS15**, with cross-validation posterior errors for each parameter in **TableS14**.

Discussion

We aimed at inferring jointly the tree-topology, instantaneous or recurring gene-flow processes, and asymmetric gene-flow intensities across pairs of recent and ancient lineages that most likely produced genome-wide genetic patterns in 54 different combinations of five Central and Southern African populations. Therefore, we explicitly considered the large genetic differentiation observed across African populations at regional and local scales, and the possible confounding effects of complex gene-flow processes on tree-topologies' predictions among 48 complex scenarios in formal statistical competition.

First, our results altogether show that, in fact, considering different sets of populations at a regional and local scale in Africa highlights different aspects of the demographic histories of African populations, sometimes substantially divergent. This demonstrates in practice that apparently discrepant results obtained across previous studies may not necessarily be absolutely reconciled, but rather interpreted as separate illustrations of the large diversity of demographic histories experienced by our species in Africa since its emergence in a remote past. This further strongly advocates for avoiding merging individual samples among differentiated populations in un-discriminated *a priori* categories based on geography, subsistence-strategy, and/or linguistics criteria. Instead, we recommend developing novel approaches, probably at a large computational cost as experienced here, to explicitly take into account African genetic diversity and genetic structure at the continental, regional, and/or local scales, and *a posteriori* interpreting the diversity of results obtained separately.

Short periods of gene-flow rather than recurring migrations in the ancient history of Africa

As pointed out here and in numerous previous theoretical or empirical studies (e.g. [22–24,26,31,66–70]), whether ancient gene-flow processes occurred recurrently or more instantaneously across lineages has fundamental consequences on the biological and cultural evolution of our species. Indeed, recurrent gene-flow processes allow for continuous allelic exchanges across populations -thus never reproductively isolated throughout entire periods of evolution-, which may strongly influence the relative influence of drift and selection across populations and the notion of ancient tree-like evolution in human populations in Africa. We found that instantaneous gene-flow processes systematically vastly outperformed recurring migration processes, whichever the tree-topology for ancient and recent Central and Southern African lineages divergences, and for all combinations of five observed populations. Therefore, our results unambiguously favor an evolutionary history of African lineages ancestral to a variety of Central and Southern African populations where *Homo sapiens* populations experienced long periods of isolation and drift, followed by short periods of possibly asymmetric gene-flow, which may in turn have induced some reticulations among lineages [22,71]. Notably, this scenario further allows for differential selection and adaptation processes across lineages (e.g. [6,10,23]), and for ancient admixture-related selection processes within Africa, analogous to previous findings of such processes having occurred more recently throughout the continent [7,9,72,73]).

Nevertheless, it is certain that *Homo sapiens* evolutionary history of ancient and recent gene-flow in Africa has involved more complex processes among pairs of lineages than the ones here identified. Previous studies considered arguably more realistic models to explain their data, with either a conjunction of recurring symmetric migrations among pairs of lineages with possible instantaneous introgressions among certain pre-specified recent and/or ancient lineages [22,24], or only pre-specified instantaneous introgressions without recurring gene-flows [23,26], including specific events involving the contributions from *Homo sapiens* or non-*Homo*

sapiens unsampled lineages. In this context, our results highlight that complex instantaneous
introgression scenarios should probably be preferred as a starting-point from where one could
then increase complexity, for instance by considering more than a single instantaneous gene-
flow event among pairs of ancient lineages.
Moreover, our results advocate for the possibility of at least some such introgressions to
be relatively intense, as scenarios considering low or moderate gene-flows, whether
instantaneous or recurring, are systematically poorly mimetic of observed genomic patterns.
Echoing this result, a previous study identified high levels of instantaneous introgressions
across African lineages as well [26], albeit the authors considered very different maximum-
likelihood approaches based on different statistics and scenario specifications, and different
population samples, sometimes merged in larger categories at a continental scale.
Nevertheless, our approach failed to provide satisfactory posterior estimates of gene-flow
intensities, a serious limitation which we discuss extensively in the methodological limits and
perspectives section below. Importantly, previous studies investigating conjunctions of
recurring and more instantaneous migrations overall estimated that symmetric recurring gene-
flows were weak, whether recent or ancient (e.g. [22]). Although our results are difficult to
compare with previous studies due to strong differences in model specifications, statistics used
and methodological approaches as well as population samples, this latter previous result may
be reflected in ours that disfavor recurring migration models as being poorly explicative of our
data.
1029
Inferred demographic history in Central and Southern Africa
Extensive previous work agrees that divergences among lineages ancestral respectively to
extant Khoe-San, Rainforest Hunter-Gatherers, and Rainforest Hunter-Gatherer neighbors
were among the most ancient in *Homo sapiens* evolution (e.g. reviewed in [2,3]). However, the
relative order of their divergences as well as their timing has been a matter of extensive debate
[25,26,40]. This is due in part to split-time inferences conducted on tree-like topologies considering
gene-flow or no gene-flow, using ancient DNA data or not, and to differing methods, statistics,
and population samples.
1038
Most ancient divergences in Africa
Here, we formally tested, with Random-Forest ABC scenario-choice, which ancient tree-
topology best explained extant genomic patterns, whichever the gene-flow processes and
intensities across pairs of recent and ancient lineages, for 54 different combinations of five
different Khoe-San, Rainforest Hunter-Gatherer, and Rainforest Hunter-Gatherer neighboring
populations. It is the first time to our knowledge that such formal comparison of numerous
competing-scenarios is conducted systematically for a variety of sets of population samples at
a regional and local scale, rather than comparing maximum-likelihood values from vastly
differing models, each obtained separately using differing population samples [22,25,26].
Importantly, note that [24] also explored highly complex demographic scenarios in Africa with
ABC, focusing on varying unidirectional introgression processes among certain lineages for a
fixed tree-shape among, in particular, three extant African populations at the continental scale.
Our results show that whichever the gene-flow processes considered, the lineage ancestral
to extant KS populations diverged first from the lineage ancestral to RHG and RHGn, for
almost all combinations of sampled populations. Furthermore, we estimated with Neural
Network ABC posterior-parameter inferences that this original divergence likely occurred in a
remote past ~330,000 years ago, followed by the divergence between ancestral RHG and
RHGn lineages ~158,000 years ago, considering 20 or 30 years per generations. Recent paleo-
genomic data from ancient southern African individuals also demonstrate deep population
structure and long-term isolation in southern Africa, and show that present-day Khoe-San

populations retain only part of this ancient structure following later gene-flow [74]. Importantly,
the ABC framework employed here explicitly models gene-flow processes when inferring past
population structure, allowing deep divergence signals to be evaluated in the presence of
migration.
Both our estimates fall within the upper bound of previously obtained results considering
a variety of similar population samples, genetic data, inference methods, and statistics, albeit
scenarios may not have always been specified in analogous ways [4–6,8,25,26,40,75]. Note however,
that considering different combinations of populations at the regional and local scales allowed
us to identify a small minority (4/54) of such combinations for which an alternative scenario
(Scenario i1-3b), where RHGn lineages diverged first from a lineage ancestral to all RHG and
KS extant populations, somewhat analogous to those proposed in [26], would better explain the
data consistently across our analyses, again whichever the gene-flow processes and range of
intensities considered. This latter result may explain apparently discrepant results across some
previous studies, which would then be due to differing population samples used for
demographic inferences and/or to artificial merging samples from differentiated populations
into larger groups [22,24–26], thus advocating for further accounting for the vast genetic diversity
among African populations even at a local scale in future studies [70].
Note that, in [22], the fundamental divergence between their Nama Khoe-San sample and
other Sub-Saharan populations was dated to ~110,000-135,000 years ago under their two best-
fitting models, thus strongly discrepant with our findings. However, these important results are
very difficult to compare with ours, since the authors did not consider any Central African
Rainforest Hunter-Gatherers nor their neighbors in their inferences, and since they investigated
highly complex models specified completely differently from those here envisioned. In
particular, Ragsdale and colleagues included in their models possible very ancient genetic
structures, long before *Homo sapiens* emergence, a feature that is unspecified in our scenarios
which considered simply a single ancestral population in which all extant lineages ultimately
coalesce. Nevertheless, note that substructure and reticulation within the ancestral population
is not *per se* incompatible with our scenarios. In fact, it may be compatible with our posterior
estimates of a large effective population ancestral to all extant populations here investigated,
among the largest of inferred ancient and recent Central and Southern African effective
population sizes. Therefore, it will be reasonable in future work to complexify the scenarios
here proposed to evaluate whether very ancient substructures and reticulations within our
ancestral population, prior to the original divergence between Southern and Central African
populations, may improve the fit to the observed genomic data, as proposed by [22].
1093
More recent divergences in Central and Southern Africa
More recently during the evolutionary history of Sub-Saharan Africa, we found that the
divergence among Northern and Southern Khoe-San populations largely pre-dated the
divergence of Western and Eastern Congo Basin Rainforest Hunter-Gatherers, a question rarely
addressed to our knowledge. First, we found that KS divergence dated sometime between
57,000 and 85,000 years ago, thus substantially more recently than estimates previously
proposed [2]. Beyond vast differences in models, methods, and statistics used to provide either
inference, note that our results for this divergence time were substantially variable across pairs
of sampled populations used in each analysis, some specific sets of populations providing
posterior estimates consistent with previous results. Thus, it further highlights that complex
inferences in Africa imperatively need to explicitly consider population variation at a local
scale.
Interestingly, our synthesized estimates for the Northern and Southern Khoe-San
population divergence were relatively synchronic to the genetic onset of the Out-of-Africa (e.g.
[2]). Population genetics inferences only provide possible mechanisms to explain observed

genetic patterns and are, in essence, not addressing the possible causes underlying the inferred mechanisms. In this context, we may hypothesize that the global climatic shifts inducing massive ecological changes that have occurred in Africa at that time (e.g. [76]), sometimes proposed to have triggered ancient *Homo sapiens* movements Out-of-Africa, may also have triggered, independently, the genetic isolation among ancestral Khoe-San populations. Nevertheless, where the ancestors of extant Khoe-San populations lived at that time remains unknown and is nevertheless crucial to further elaborate possible scenarios for the causes of the genetic divergence here inferred.

We found that Rainforest Hunter-Gatherer populations across the Congo Basin diverged long after this divergence, roughly between 15,000 and 23,000 years ago, relatively consistently across pairs of sampled populations used for inferences; estimates highly consistent with previous studies [8,75,77], despite major differences in gene-flow specifications across RHG groups between studies. Interestingly, this divergence time is relatively synchronic with absolute estimates for the Last-Glacial Maximum in Sub-Saharan Africa (e.g. [78]). The fragmentation of the rainforest massif during this period in the Congo Basin may have induced isolation between Eastern and Western RHG extant populations, as plausibly previously proposed [75]. However, similarly as above for the Northern and Southern Khoe-San populations divergence, where the ancestors of extant Eastern and Western Rainforest Hunter-Gatherers lived remains unknown, which prevents us from formally testing this hypothesis [79].

Altogether, these results show that Central African Rainforest Hunter-Gatherer and Southern African Khoe-San populations have had, respectively, extensive time for selection processes, including adaptive introgression processes, to have influenced independently both groups of populations as well as populations within each group separately (e.g. [6,7,9,10]).

Ancient and recent instantaneous gene-flow times in Africa

We obtained reasonably well estimated instantaneous asymmetric gene-flow times among almost all pairs of ancient and recent lineages, albeit the variation of estimates across sets of Central and Southern African sampled populations increased substantially with most ancient times. In this context, we deem it hard to confidently try to interpret the most ancient event of instantaneous gene-flow between the two most ancestral lineages in our tree-topology. However, other instantaneous gene-flow time estimates throughout the topology were more consistently estimated overall, and showed relative synchronicity in some cases, which has never been reported before to our knowledge, even if they were specified independently in our models and drawn from large distributions *a priori*.

Interestingly, we found strong indications for almost synchronic events of introgressions having occurred during the Last Interglacial Maximum in Africa [71], between ~85,000 and ~135,000 years ago (when considering 20 or 30 years per generation). They involved gene-flow between lineages ancestral to Khoe-San populations and ancestors of Rainforest Hunter-Gatherer neighbors on the one hand and, on the other hand, between lineages ancestral to Khoe-San populations and the lineage ancestral to all Rainforest Hunter-Gatherers. An increase in material-based culture diversification and innovation, possibly linked to climatic and environmental changes locally, has previously been observed during this period of the Middle Stone Age in diverse regions of continental Africa; prompting a long-standing debate as to its causes if human populations were subdivided and isolated biologically and culturally at the time [80–83].

In this context, our results instead may suggest that population movements at that time among previously isolated populations may itself have triggered the observed increased cultural diversification locally, even if obvious signs of pan-African cultural spread at the time are difficult to assess [82]. In turn, it would interestingly echo the known effect of within-

population genetic diversity increase induced by genetic admixture between previously isolated populations (e.g. [18,31,66,84]).

Then, note that we estimated that the instantaneous gene-flow event between the ancestral Rainforest Hunter-Gatherers lineage and that of their extant neighbors seemingly occurred synchronically to the genetic Out-of-Africa ([76]; see above). This would imply that possible climatic and ecological shifts at that time may not have only induced population divergences and displacement, but may also have triggered population gene-flow.

Relatively more recently, around 30,000 years ago, we found two loosely synchronic gene-flow events between ancestors to extant Central African Rainforest Hunter-Gatherer neighbors' lineages and, separately, Northern and Southern Khoe-San lineages. This corresponds to the end of the Interglacial Maximum and a period of major cultural changes and innovations during the complex transition from Middle Stone Age to Late Stone Age in Central and Southern Africa [80,83,85–87]. Nevertheless, connecting the two lines of genetic and archaeological evidence to conclude for increased population movements at the time and their possible causes should be considered with caution. Indeed, in addition to genetic-dating credibility-intervals being inherently much larger than archaeological dating, this period remains highly debated in paleoanthropology mainly due to the scarcity and complexity of the material-based culture records, and that of climatic and ecological changes locally, across vast regions going from the Congo Basin to the Cape of Good Hope [80,83,85–87].

Finally, we found strong signals for multiple instantaneous gene-flow events having occurred between almost all five recent Central and Southern African lineages between 6000 and 15,000 years ago, during the onset of the Holocene in that region, shortly before or during the beginning of the last Post Glacial Maximum climatic crisis in Western Central Africa [88], the emergence and spread of agricultural techniques [89], and the demic expansion of now-Bantu-speaking populations from West Central Africa into the rest of Central and Southern Africa [9,17,90]. These results are consistent with previous investigations that demonstrated the determining influence of Rainforest Hunter-Gatherer neighboring populations' migrations through the Congo Basin in shaping complex socio-culturally determined admixture patterns [4,5,8,75], including admixture-related natural selection processes [9,10,77,91]. As our estimates for introgression events are in the upper bound of previous estimates for the onset of the so-called "Bantu expansion" throughout Central and Southern Africa, we may hypothesize here that major climatic and ecological changes that have occurred at that time may have triggered increased population mobility and gene-flow events between previously isolated populations, rather than consider that the Bantu-expansions themselves were the cause for all the gene-flow events here identified.

Finally, we did not find signals of more recent introgression events from Bantu-speaking agriculturalists populations into Northern or Southern Khoe-San populations, in particular among the !Xun, albeit such events have been identified in several previous studies (see [2]). This is likely due to the fact that we considered only a limited number of individual samples from each population with low apparent levels of recent admixture, and therefore may lack power to detect these very recent events with our data and approach.

Conceptual, methodological, and empirical limitations and perspectives for inferring ancient histories from observed genomic data

All previous attempts at reconstructing the human evolutionary histories which led to genomic patterns observed today across African populations have faced major conceptual, methodological, and empirical challenges. Conceptually, large amounts of more-or-less nested scenarios can be envisioned *a priori* to explain extant genetic diversity, based on previous results from paleo-anthropology, population genetics, and paleogenomics. These scenarios may range from tree-like models without gene-flow events among ancient or recent lineages

to complex networks of weakly differentiated populations exchanging migrants over large periods of time, with or without the contribution of ancient *Homo* or non-*Homo* now extinct or unsampled lineages. Systematically exploring all possible models is often methodologically out of reach due to differing fundamental scenario-specifications or, when scenarios are specified and parameterized in analogous ways, due to scenarios' nestedness and un-identifiability in certain parts of their parameter spaces (e.g. [92]). Empirically, formal scenario comparisons are first hampered by necessarily limited amounts of genomic data representative, at continental, regional, and local scales, of the known diversity and differentiation of human populations in Africa; in addition to yet limited amounts of ancient DNA data throughout the continent. Finally, empirical limitations also emerge from the use of different statistics to explore genomic diversity patterns, which possibly each capture different facets of human evolutionary histories, thus providing discrepant results and interpretations only in appearances. While machine-learning ABC procedures provide significant advantages over other maximum-likelihood approaches, in particular concerning the formal exploration of competing scenarios' fit to observed data across numerous highly complex sometimes nested scenarios and using numerous summary-statistics [43,45,63,92], the above challenges are also largely faced in this study.

While most divergence and gene-flow times, as well as effective sizes, were inferred satisfactorily in our results, the lack of satisfactory posterior estimates for all 20 gene-flow rates parameters consistent among the 25 sets of population combinations likely stems from different limitations. First, considering only five individual genomes per population is inherently limiting when trying to estimate gene-flow rates from inter-population summary-statistics in ABC [93]. In future studies, increasing sample sizes and adding statistics based on the inter- and intra-individual distribution of admixture fractions [31,32,66,93], will likely improve the posterior estimation of these parameters using machine-learning ABC approaches [24]. Note, however, that these statistics require non-trivial computation time in ABC frameworks comprising hundreds of thousands of simulations [94,95]. Furthermore, in all cases, considering multiple sets of different and substantially genetically diverse populations as the ones here considered from Sub-Saharan Africa may inevitably lead to large variation across gene-flow rates' posterior estimates depending on the specific population-combinations.

Second, for simplicity and as a starting point, we considered a single instantaneous time for the two separate unidirectional gene-flow events, for each gene-flow event between pairs of lineages separately. Therefore, while our RF-ABC scenario-choice results strongly support instantaneous asymmetric gene-flow events rather than recurring asymmetric gene-flows to best explain extant genomic patterns, it is plausible that scenarios where each unidirectional gene-flow event may occur at a different time [24], will more realistically explain the observed data. If this is the case, it is possible that the choice of a unique time for two separate gene-flow events rendered the corresponding gene-flow rates harder to identify with our joint NN-ABC posterior parameter estimation procedure. Furthermore, we explored two extreme gene-flow processes, establishing an open competition between only instantaneous gene-flow and only recurring gene-flow processes. While we demonstrated that only instantaneous gene-flow processes vastly outperformed only recurring gene-flow processes, whichever the range of intensities for each event, we only established here a starting point for the future complexification of evolutionary scenarios to be tested. For instance, it will be of natural interest to consider, next, more than a single "pulse" of gene-flow between any two ancient or recent lineage.

Third, to reduce the number of free parameters and provide a starting point for future, more complex and realistic, inferences, we considered constant effective population sizes in the terminal branches of our topologies, albeit extensive previous work have inferred diverse recent demographic dynamics across Sub-Saharan African populations (e.g. [17,65,96–98]).

Importantly, how these recent demographic changes may influence divergence times posterior
estimates is not trivial in models as complex as the ones here investigated. For instance,
previous studies considered ABC frameworks and a variety of autosomal and uniparental
genetic data [4,5,9,75,99]. They investigated divergence times among sampled lineages assuming
possible gene-flow processes and very different population size changes models. Overall, while
informing a variety of bottlenecks, constant, or increases in effective population sizes,
divergence times remained largely consistent across studies that considered the same sets of
sampled populations, and consistent with our genome-wide results for comparable sets of
sampled populations. Therefore, it will be crucial in future studies to further complexify
effective population sizes changes in the proposed topologies to empirically evaluate how they
affect or not divergence times estimates; and further disentangle the complex evolutionary and
migration history of African populations using increasingly realistic scenarios.
Fourth, the neural networks used here may have been unable to satisfactorily identify the
20 gene-flow parameters among the 43 jointly-estimated parameters, due to their lack of
complexity when considering a unique layer of hidden neurons. For instance, and as a first
step, considering Multilayer-Perceptrons in the future, *i.e.* more complex feed-forward neural
networks with additional layers of hidden neurons, may help posterior estimations of these
parameters (e.g. [100]), at the known cost of non-trivial parameterization and optimization of the
neural networks themselves (e.g. [95,101,102]).
Altogether, joint posterior estimation of numerous gene-flow rates parameters and the
timing of their occurrence under highly complex demographic scenarios remains one of the
most challenging tasks in population genetics. It will unquestionably benefit from future
analytical theoretical developments [103,104], and the improvement of machine-learning-based
inference procedures (e.g. [102,105–107]).
1282
Archaic admixture with *Homo sapiens* or non-*Homo sapiens* unsampled lineages?
Our results demonstrate that explicitly considering archaic admixture from unsampled
populations is not necessary to satisfactorily explain the observed genomic diversity of extant
Central and Southern African populations when considering jointly the 337 relatively classical
population genetics summary-statistics used here for demographic inferences, consistently
with a previous study [22], and conversely to others [3,25,26]. We did not explore possible
contributions from unsampled lineages, whether from non-*Homo sapiens* or from ancient
"ghost" human populations, and therefore cannot formally reject such possibilities. However,
by parsimony, our results show that the simpler explanation (without significant archaic
admixture signature at the genomic level), may possibly be more likely.
Our set-up formally comparing competing-scenarios rather than comparing posterior
likelihoods of highly complex yet vastly differing models, provide a clear and reasonable
starting point for future complexification of scenarios comprising possible contributions from
ancient or ghost unsampled populations; where the burden of proof lies on showing that
scenarios with archaic admixture fit the data significantly better than scenarios without such
archaic admixture (despite the increased complexity and more parameters). Such endeavors
will also benefit from the explicit use of additional novel summary-statistics ([22,26,32]; see also
above).
In any case, the complexification of scenario-specifications to account for possible past
"archaic" or "ancient" introgressions will not fundamentally solve the issue of the current lack
of reliable ancient genomic data older than a few hundreds or thousands of years from Sub-
Saharan Africa [3,108,109]. Indeed, analogously to archaic admixture signals that were
unambiguously identified outside Africa only when ancient DNA data were made available for
Neanderthals and Denisovans (e.g. [27,28]), we imperatively need to overcome this lack of

empirical ancient DNA data in Africa to formally test whether, or not, ancient human or non-
human now extinct lineages have significantly contributed to shaping extant African diversity.
1309

1310

Material and Methods

Population samples

Central and Southern Africa dataset

We investigated high-coverage whole genomes newly generated for 74 individual samples (73 after relatedness filtering, see below), from 14 Central and Southern African populations (**Figure1**, **Table1, TableS1**). Based on extensive ethno-anthropological data collected from semi-directed interviews with donors and their respective communities, we grouped *a posteriori* the 73 individual samples from 14 populations in three larger categories.

The Baka, Ba.Kola, Bi.Aka_Mbati, Ba.Twa, and Nsua individuals were categorized as Rainforest Hunter-Gatherers (RHG), based on several criteria including self-identification and relationships with other-than-self, socio-economic practices and ecology, mobility behavior, and musical practices [4,38].

The Nzime, Ngumba, Ba.Kiga, and Ba.Konjo individuals were categorized as Rainforest Hunter-Gatherer neighbors (RHGn), based on the same set of categorization-criteria as RHG. Indeed, RHG and RHGn populations in Central Africa are known to identify themselves separately and to live in different ways in the Central African rainforest, whilst often sharing languages locally as well as complex socio-economic relationships and interactions that may include intermarriages [5]. Historically, RHG populations have been designated as "Pygmies" by European colonists, an exogenous term from ancient Greek that is sometimes used derogatorily by the RHGn.

Finally, the Nama, Ju|'hoansi, Karretjie People, !Xun, and Khutse_San individuals were categorized as Khoe-San populations (KS) based primarily on self-identification and relationships with other-than-self locally, lifestyle, and languages [110]. Note that research was approved by the South African San Council. Individuals from these populations were whole-genome sequenced anew with improved methods (see below), compared to previous work [40].

Note that the three groups can be further subdivided based on geography into Western and Eastern RHG and RHGn, and into Northern, Central, and Southern KS (**Figure1**, **Table1**).

Comparative dataset

The 74 genomes were analyzed together with 105 (104 after relatedness filtering, see below) high-coverage genomes from 32 worldwide populations (**Figure1**, **Table1, TableS1**). 64 samples from the Simon's Genome Diversity Project (SGDP) [34]; 9 HGDP samples [27]; 7 samples from the 1000 Genomes project (KGP) [35]; one Karitiana sample [36]; 24 samples from the South African Human Genome Project (SAHGP) [37]. Selection criteria were: Illumina paired-end reads, high-coverage (>30X), access to raw data and no (known) genetic substructure within population.

Note that, although samples from these datasets have not been collected by us for the purpose of this study, based on previous knowledge and publications, the Biaka and Mbuti HGDP samples can be categorized as RHG and the #Khomani and Ju|'hoansi_comp samples as KS (**Figure1**, **Table1**).

Data generation

DNA extraction

DNA was extracted from saliva with the DNA Genotek OG-250 kit for the Baka, Nzime, Bi.Aka_Mbati, Nsua, and Ba.Konjo samples. DNA was extracted from buffy coats with DNeasy Blood&Tissue spin-column Qiagen™ kits for the Ba.Kola, Ngumba and Ba.Kiga samples. Both extraction methods were used for the Ba.Twa samples. For the Nama, Ju|'hoansi, Karretjie People, !Xun, and Khutse_San samples, DNA was extracted from EDTA-blood using the salting-out method [111].

Library preparation and sequencing

Library preparation and sequencing was performed by the SciLifeLab SNP&SEQ Technology platform in Uppsala, Sweden. Libraries were prepared with TruSeq DNA preparation kits, and paired-end sequencing (150 bp read length), was performed on Illumina HiSeqX machines with v2 sequencing chemistry, to a coverage of 30X or more. For the Southern African samples, we included paired-end data (100 bp read length) sequenced on Illumina HiSeq2000 machines obtained previously for the same libraries [40].

Sequencing data quality-control processing and relatedness filtering

The processing pipeline used in the present study is adapted from the Genome Analyses Toolkit (GATK) "Germline short variant discovery (SNPs + Indels)" Best Practices workflow [112–114]. It is described and compared to the original GATK Best Practices workflow in [115]. **FigureS27.** gives an overview of the pipelines used for the processing; all template codes with accompanying detailed explanations for all processing steps are provided below and in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Reads were mapped to a decoy version of the human reference genome, GRCh38 (1000 Genomes Project version), with BWA-MEM [116] from BWAKIT v0.7.12 using a strategy appropriate for ALT contigs. The input was either FASTQ or BAM files that were reverted to unmapped BAM or to FASTQ prior to mapping. Mapped reads were selected with SAMtools version 1.1 and Picard version 1.126 (https://broadinstitute.github.io/picard/).

Note that, as mentioned in [117], the GRCh38 is a "mosaic haploid representation of the human genome". The dominant sequence contribution (70%) is from a donor of "likely African-European admixed ancestry". Therefore, this assembly of the reference genome is not equally genetically related to all extant individuals within and among human populations. It is important to keep in mind, thus, that while numbers of variants can be compared across studies referring to the same genome assembly, in any given study, some populations may have artificially high number of variants as compared to the reference.

Duplicate reads were marked (at the lane level) with Picard version 1.126 MarkDuplicates. Realignment around indels was performed (at the lane level) with GATK version 3.5.0 RealignerTargetCreator and IndelRealigner. We then performed a "triple mask base quality score recalibration (BQSR)", where the sample's variation is used together with dbSNP, as described in [40,115]. This step includes merging the reads from a given sample with SAMtools version 1.1; calling variants with GATK version 3.5.0 HaplotypeCaller with "--genotyping_mode DISCOVERY"; and (in a parallel track) standard BQSR with GATK version 3.5.0 BaseRecalibrator and PrintReads, and dbSNP version 144. Following triple mask BQSR (performed at the lane level), the recalibrated reads from a given sample are merged with SAMtools version 1.1, sorted and indexed with Picard version 1.126 SortSam. Duplicate marking and realignment around indels were performed again, at the sample level.

Variant calling was performed independently for the autosomes and the X chromosome, due to the difference in ploidy. SNP and short indels were first called with GATK version 3.7 HaplotypeCaller in each genome, with the "--emitRefConfidence BP_RESOLUTION" option. Multi-sample GVCFs were then generated with CombineGVCF, and variants were finally jointly called with GATK GenotypeGVCFs. Genotypes were emitted for each site (option "allSites").

Variants were filtered with GATK version 3.7 Variant Quality Score Recalibration (VQSR) using recommended resources from the GATK bundle: HapMap (version 3.3), 1000 Genomes Project Illumina Omni 2.5M SNP array, 1000 Genomes Project phase 1 high confidence SNPs, and dbSNP version 151 for the SNPs; and gold standard indels [118] and dbSNP version 151 for the indels. For the X chromosome, autosomal variants were included for training the VQSR model (VariantRecalibrator step).

We controlled the dataset for related individuals with KING [119] "--kinship" and plink version 1.90b4.9 [120] "--genome --ppc-gap 100". The dataset was pruned for linkage disequilibrium (LD) before relatedness estimation with plink, using plink "--indep-pairwise" with sliding windows of 50 SNPs, shifting by five SNPs, and a r^2 threshold of 0.5. Both methods identified two pairs of first-degree relatives. We excluded the sample with greatest missingness from each pair using GATK version 3.7 SelectVariants to produce the family unrelated working datasets used henceforth.

The callset was then further refined; a "FAIL" filter status was set on sites that are ambiguous in the reference genome (base "N") or had greater than 10% missingness, identified with VCFtools version 0.1.13 "--missing-site" [121]. Moreover, for the autosomes, variants heterozygous in all samples were marked as failed (Hardy Weinberg Equilibrium -HWE- filter, identified with VCFtools version 0.1.13 "--hardy").

Coverage was computed with QualiMap version 2.2 [122] including or not duplicates ("-sd" option for the latter). The average coverage (without duplicates) across individuals within populations is provided in **TableS3**.

Descriptive analyses

Variant counts

We obtained per-sample and aggregated metrics of variant counts (**Figure2**), with the tool *CollectVariantCallingMetrics* of the software package Picard v2.10.3 (https://broadinstitute.github.io/picard/), applied to the full 177 worldwide individuals' dataset after VQSR, relatedness, HWE and 10% site missingness filtering. We conducted variant counts procedures for chromosomes 1 to 22. We used dbSNP 156 as a reference; we downloaded "GCF_000001405.40.gz" and modified the contig names to match contig names in the VCFs. Separately, we applied the same variant counts pipeline as above to the Central and Southern African 73 individuals' original dataset extracted from the full dataset with BCFtools version 1.17 (https://github.com/samtools/bcftools), using view with the option "-S". All scripts are provided in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Genome-wide Heterozygosity

We calculated observed and expected heterozygosities, for the autosomes and for the X-chromosome separately using custom Python, Bash, and R scripts (full pipeline available at GitHub https://github.com/Gwennid/africa-wgs-descriptive), for the 177 worldwide individuals after variant-counts filtering described above. For autosomes, we considered the main contigs ("chr1" to "chr22"). For the X chromosome, we excluded the Pseudo-Autosomal

Regions with coordinates from GRCh38 (PAR1: 10,000 to 2,781,480, PAR2: 155,701,382 to 156,030,896).

For each population with more than one individual and for each autosome and X-chromosome separately, we counted the number of variable and non-variable sites and excluded multi-allelic sites, indels, and sites with missing genotypes in at least one individual in the population for simplicity and conservativeness. For each autosome and each individual, we then counted separately the numbers of observed homozygous and heterozygous sites of each configuration compared to the reference (0/0; 0/1; 1/0; 1/1). Based on these counts, for each population and each chromosome separately, we first calculated observed heterozygosities simply as the average proportion of heterozygous individuals per variable sites with no-missing data in the population sample. We also computed unbiased expected multi-locus heterozygosities for all variable loci with no missing genotypes in the population as in equations (2) and (3) in [47]. We averaged this value across all sites with no missing information in the population sample including both variable and non-variable sites, and finally corrected it for haploid population sample sizes (**Figure2**). We compared unbiased heterozygosities by chromosomes across groups of populations using Wilcoxon signed-rank tests. All scripts are provided in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Runs of homozygosity

We identified runs of homozygosity (ROH) using the *homozyg* tool from PLINK version 1.90b4.9 [120]. We selected autosomal biallelic SNPs from the variant-counts pipeline described above with GATK version 3.7 SelectVariants with options "-selectType SNP -restrictAllelesTo BIALLELIC -excludeFiltered". We converted the VCF to TPED and then binary plink fileset with VCFtools version 0.1.13 "-plink-tped" and plink. For the 177 individuals worldwide, we considered only ROHs measuring more than 200 Kb (*--homozyg-kb* 200), containing at least 200 SNPs (*--homozyg-snp* 200), containing at least one variable site per 20 Kb on average (*--homozyg-density* 20), and with possible gaps of up to 50 Kb (*--homozyg-gap* 50). Default values were considered for all other parameters: *--homozyg-window-het* 1 *--homozyg-window-snp* 50 *--homozyg-window-threshold* 0.05. Finally, the total mean ROH lengths were calculated for each population separately using the ".hom" output-file for each of four length categories: 0.2 to 0.5 Mb, 0.5 to 1 Mb, 1 to 2 Mb, and 2 to 4 Mb (**Figure2**). Corresponding pipelines are provided in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Individual pairwise genome-wide genetic differentiation

We explored genome-wide genetic differentiation between pairs of individuals with Neighbor-Joining Tree (NJT) [52,53], and Multi-Dimensional Scaling (MDS) approaches based on the pairwise matrix of Allele-Sharing Dissimilarities (ASD) [51] computed using 14,182,615 genome-wide autosomal SNPs pruned for low LD, with PLINK version 1.90b4.9 [120] *indep-pairwise* function (*--indep-pairwise 50 5 0.1*). We considered at first all 177 unrelated individuals in our dataset to calculate the ASD matrix, and then subset this ASD matrix for the 73 Central and Southern African unrelated individuals original to this study before computing NJT and MDS analyses anew (**Figure3, FigureS2**).

We computed the ASD matrix considering, for each pair of individuals, only those 14,182,615 SNPs without missing data, using the *asd* software (v1.1.0a; https://github.com/szpiech/asd; Szpiech, 2020). We computed the unrooted NJT using the *bionj* function of the R package *ape* and setting the branch-length option to "true". We computed the MDS using the *cmdscale* function in R.

We compared ASD values with populations' geographic locations by computing the standardized average ASD across all pairs of individuals from two different populations, for all pairs of African populations separately (**Figure1**, **Table1, TableS1**), and compared these values with the logarithm of geographic distances among pairs of populations (setting 0 km distances to "1"), using Spearman ρ and Pearson r correlations for different continental, sub-continental, and regional sets of populations. We tested these correlations using 1000 Mantel permutation tests (**TableS4**), with the *partial.mantel.test* function of the *ncf* package in R. We computed the shortest geographic distance between two populations' GPS coordinates on an Earth-like ellipsoid (with an earth radius of 6378.137 km by default in WGS84), using the *distm* function with *distGeo* method from the *geosphere* package in R.

We conducted permutational ANOVA analyses to test, separately, whether linguistic families, sub-linguistic families, or traditional food-producing strategies (**TableS1**), significantly explain average population-pairwise ASD patterns, using the *adonis2* function from the *vegan* package in R with 1000 permutations for each analysis conducted on subsets of the above ASD matrix (**Figure3, FigureS2**) at varying geographical scales (**TableS5**, **TableS6**, **TableS7**).

The clustering software ADMIXTURE [54] allows researchers to further explore inter-individual genome-wide levels of dissimilarity and resemblance. Indeed, while it is tedious and often cognitively difficult to explore multiple combinations of dimensions of genetic variation using an MDS or a NJT approach, ADMIXTURE instead allows exploring K higher such dimensions at once [54–58]. However, note that the visual representation of relative "distances" across pairs of individuals is lost in the classical bar-plot representation of ADMIXTURE, hence showing the complementarity of this descriptive analysis to the above MDS and NJT.

We considered here only 840,031 genome-wide autosomal SNPs pruned for LD (r^2 threshold 0.1) and with Minimum Allele Frequency above 0.1, for the 177 worldwide unrelated individuals, using PLINK *indep-pairwise* and *maf* functions. We computed unsupervised ADMIXTURE version 1.3.0 clustering with values of K ranging from 2 to 10, considering 20 independent runs for each value of K separately. We then calculated ADMIXTURE results symmetric-similarity-coefficient SSC for each value of K separately in order to find the groups of runs providing highly similar results (SSC>99.8%). Individual's genotype membership proportions to each K cluster were then averaged, per individual, across such highly resembling runs, and then plotted. SSC calculations, averaging results across similar runs, and producing barplots were conducted using the software PONG with the "greedy" algorithm [60]. Corresponding pipelines are provided in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive). For each value of K separately, results representing the majority of the 20 independent runs are presented in **Figure4**, and all other results are presented in supplementary **FigureS3**.

Machine-Learning Approximate Bayesian Computation scenario-choice and posterior parameter estimation

We reconstructed the complex demographic history of Central and Southern African populations using machine-learning Approximate Bayesian Computation (ABC) [41–45]. In principle, in ABC, researchers first simulate numerous genetic datasets under competing scenarios by drawing randomly a vector of parameter values for each simulation in distributions set a priori by the user. For each simulation separately, we then calculate a vector of summary statistics, thus corresponding to a vector of parameter values used for the simulation. The same set of summary statistics is computed on the observed data. ABC scenario-choice then allows the researcher to identify which one of the competing scenarios produces the simulations for which summary-statistics are closest to the observed ones. Under

1536 this best scenario, ABC posterior-parameter inference procedures allow the researcher to
1537 estimate the posterior distribution of parameter values most likely underlying the observed
1538 genetic patterns.

48 competing scenarios for the demographic history of Central and Southern African populations

We designed 48 competing demographic-history scenarios possibly underlying the genetic
patterns observed in five Central and Southern African populations of five individuals each
(**Table1**, **Figure5**); one Eastern Rainforest Hunter-Gatherer population (eRHG: Nsua, Ba.Twa,
or Mbuti), one Western Rainforest Hunter-Gatherer population (wRHG: Baka, Ba.Kola, or
Bi.Aka_Mbati), one Eastern or Western RHG neighboring population (eRHGn: Ba.Kiga or
Ba.Konjo; wRHGn: Nzime or Ngumba), one Northern Khoe-San population (nKS: Ju|'hoansi
or !Xun), and one Southern Khoe-San population (sKS: Karretjie People or Nama). Note that
we did not include the Khutse_San population in ABC inferences for simplicity, as this
population is located at intermediate geographic distances between the nKS and sKS groups of
populations.

The 48 competing scenarios differed in their ancient tree-topologies and relative timing of
the more recent divergence events, combined with different (duration and intensities) possibly
asymmetric gene-flow processes among ancient and recent lineages (**Figure5**). This design,
while complex, allowed us, for the first time to our knowledge, to consider explicitly the
expected confounding effects of gene-flow processes on otherwise different tree-topologies,
which may also possibly induce certain reticulations among ancient lineages, while jointly
estimating divergence times, gene-flow events' timing and intensities, and effective population
size changes over time.

Importantly, we relied on extensive previous findings having formally demonstrated the
common origin of Congo Basin RHG populations [4,75], and that of KS populations [6,40],
respectively, and thus did not consider all possible tree-topologies for five tree-leaves.
Moreover, we considered only a single RHGn sample in each combination, in turn from the
East or the West of Central Africa, in order to simplify already highly complex scenarios, as
we were not interested here in the detailed demographic history of RHGn which has been
extensively studied previously (e.g. [9,17]). This simplification was deemed reasonable as RHGn
populations throughout the Congo Basin and into Southern Africa have previously been shown
to be strongly more genetically resembling one another, compared to RHG and KS neighboring
populations, or compared to much more genetically dissimilar RHG and KS populations,
respectively (e.g. [4,5,8,9,17,77,91]); a result that we also obtained here (**Figure5**).
1570

Eight competing topologies

We investigated eight competing topologies starting with a single ancestral population and
resulting in five different sampled populations in the present (**Figure5**). The eight topologies
differ in which lineage ancestral to modern KS, RHG, or RHGn diverged first from the two
others, and in the relative order of divergence events internal to KS (nKS-sKS divergence) and
internal to RHG (wRHG-eRHG divergence), in order to consider the known variable
demographic histories of Sub-Saharan populations at a regional scale.

Note that divergence and introgression times (all *t* and *tad*, see **Figure5**, **TableS8**), were
each randomly drawn in uniform prior distributions between 10 and 15,000 generations, thus
effectively setting an upper limit for the most ancient divergence times among lineages at
roughly 450,000 years ago considering an upper-bound of 30 years for human generation
duration [123]. This was vastly anterior to the current estimates for the genetic or morphological
emergence of *Homo sapiens* [124,125], which allowed us to estimate *a posteriori* the most ancient
divergences among our populations, without constraining our assumptions based on previous

results obtained with different methods, data, and models. Furthermore, these prior-distribution
boundaries allowed for gene-flow events among ancient lineages (see next section), to
influence, potentially, the timing of the earliest divergence events among human lineages
previously estimated [2,22,26]. Note that we retained for simulations only those vectors of
randomly-drawn parameter-values that satisfied the chronological order of lineage divergences
set for each eight topologies, respectively (**Figure5**, **TableS8**)

In each eight topologies, we incorporated the possibility for changes in effective
population sizes, N (**Figure5**), during history along each lineage separately by defining
constant diploid effective population sizes parameters separately for each tree-branch, each
drawn randomly in U[10-100,000] (**Figure5**, **TableS8**).

1595

Asymmetric instantaneous or recurring gene-flows and their intensities
Migration of individuals between populations is ubiquitous in human history (for an overview
in Africa, see e.g. [2,3]). Here we aimed at disentangling the nature of migration processes that
may have occurred across pairs of lineages throughout the history of Central and Southern
African populations.

In particular, we first aimed at determining whether gene-flow events during history
occurred relatively instantaneously, leading to reticulations in tree-topologies, or, conversely,
occurred recurrently over longer periods of time leading to un- or weakly-differentiated
lineages throughout history [1,21,22]. To do so, for each eight topologies described above
(**Figure5**), we simulated gene-flow either as single-generation gene-flow pulses across pairs
of lineages, or as constant recurring gene-flow across pairs of lineages in-between each lineage-
divergence event. This design resulted in 16 competing scenarios encompassing eight different
possible topologies, contrasting instantaneous gene-flow events with recurring gene-flows.
Note that, for simplicity, we did not consider possible gene-flows between KS and RHG recent
lineages, as genetic signatures of such recent migrations have never been identified in previous
genetic studies nor in historical records to our knowledge.

Importantly, we aimed at determining whether gene-flow occurred symmetrically, or not,
across pairs of lineages, in particular in the past. Indeed, previous studies already identified
recent asymmetric admixture processes between RHG and RHGn [4,5,8,75], and such possible
asymmetries have not been explored among ancient lineages in previous studies [22,26], although
they may be influencing ancient tree-topologies or reticulations across lineages and the
subsequent past evolution of extant populations.

To do so, for each gene-flow event in each 16 competing scenarios (**Figure5**), we
parameterized separately the introgression of lineage A into lineage B, from that of lineage B
into lineage A, by drawing randomly the corresponding parameter values independently in the
same prior distribution (**TableS8**). ABC posterior-parameter estimations would thus reveal
possible asymmetries in gene-flows across pairs of lineages for each event separately, if
identifiable from genomic data.

Finally, for each one of the 16 competing combinations of topologies and gene-flow
processes, we considered three classes of gene-flow intensities by setting different boundaries
of gene-flow parameters' uniform prior-distributions (**TableS8**): no to low possible gene-flow
(U[0, 0.001] for recurring processes, and U[0, 0.01] for instantaneous processes), no to
moderate gene-flow (U[0,0.0125] for recurring processes, and U[0, 0.25] for instantaneous
processes), or no to intense gene-flow events (U[0,0.05] for recurring processes and U[0,1] for
instantaneous processes). Note that, for each three classes of gene-flow intensities, each gene-
flow parameter from one lineage to another at each point or period in time was drawn
independently in the above prior-distributions' boundaries, and thus may differ across events
within a scenario, and across scenarios (**Figure5**, **TableS8**).

Altogether, this design resulted in 16x3 = 48 competing scenarios for the complex
demographic history of Central and Southern African populations. Importantly, note that for
any given class of gene-flow process or intensity, the eight topologies are highly nested in
certain parts of the space of parameter values [92]. In particular, despite the fact that we did not
consider competing scenarios with explicitly trifurcating ancient tree-topologies, scenarios in
which the parameter values for the oldest and second oldest divergence times are similar are
expected to provide results highly resembling those obtained with ancient trifurcation
scenarios.

All scenario-parameters are represented schematically in **Figure5**, and their prior
distributions and constraints are indicated in **TableS8**.

1644

Simulating genomic datasets

We performed simulations under the coalescent using *fastsimcoal2* [126,127], for each one of the
48 scenarios described above, separately. For each simulation, we generated a vector of
parameter values randomly drawn in prior-distributions and satisfying the topological
constraints as described above and in **TableS8**, with custom-made Python and Bash scripts
available in the GitHub repository for this article (https://github.com/Gwennid/africa-wgs-
abc). The genetic mutation model is based on the model deployed in [95]. We simulated 100
independent loci (or "chromosomes") with the same structure. Each such "chromosome"
corresponds to a linkage block, with the following properties: the type of marker was "DNA",
the length of the loci 1 Mb, the recombination rate was 1×10^{-8} per base pair, the mutation rate
1.25×10^{-8} per base pair, and there was no transition bias (transition rate of 0.33).

We performed 5,000 such simulations for each 48 competing scenarios separately in order
to conduct Random Forest ABC scenario-choice, hence producing 240,000 separate simulated
dataset each corresponding to a single vector of parameter values randomly drawn in prior
distributions (**TableS8**). Then we performed an additional 95,000 separate simulations under
the best scenario obtained with RF-ABC for each 54 separate combinations of observed
population samples (see below). We thus reached 100,000 simulations under each best scenario
identified for each 54 observed datasets respectively, to be used for Neural Network ABC
posterior parameter inference.

1664

Building observed genome-wide data-sets for 54 sets of five Central and Southern African populations

We considered 54 separate combinations of four eRHG, wRHG, nKS, sKS, and one eRHGn
or wRHGn populations, each with five unrelated individuals (**Table1**). We prepared a call-set
of high-quality regions by applying the 1000 Genomes phase 3 accessibility mask and filtering
out indels on the variant callset comprising all individuals. This resulted in a "fragmented"
genome, with fragments of high-quality sites and fragments that are filtered out. We then
agglutinated high-quality fragments spanning 1 Mb in total agglutinated-length. We calculated
the mapped bp distance between the first and the last fragment of each 1Mb window and then
randomly chose 100 windows spanning less than 1.2 mapped Mb and located on chromosomes
1 to 10, among the 824 such windows identified on chromosomes 1 to 22. We extracted these
100 independent autosomal loci of 1 Mb each for the 54 sets of 25 individuals.

In order to test whether the 100 windows used in all ABC inferences were resembling
larger portions of the autosomal genome, we calculated the summary statistics used for ABC
inferences (see next section) for each one of the 54 combinations of five sampled-populations
each on the 100 windows and separately on the 724 other such windows identified genome-
wide. Some of the summary statistics, such as the total number of biallelic SNPs in the sample,
were scaled by the number of windows, to allow comparisons between the 100 and the 724

windows. We then performed Wilcoxon's two-sided rank test and Spearman's correlation test using the *wilcox.test* and *cor.test* functions in R.

Finally, we checked that these pruned datasets provided individual pairwise genomic differentiation patterns resembling those obtained with the entire SNP set, by computing standardized ASD matrices for each one of the 54 combinations of five populations separately. We compared these ASD matrices with the same individual subsets of the entire ASD matrix computed as above and used for MDS and NJT analyses in **Figure3,** using Spearman ρ and Pearson r correlations tested using 1000 Mantel permutation tests (**TableS10**), with the *partial.mantel.test* function of the *ncf* package in R.

Calculating 337 summary statistics

For each simulated genetic dataset, we computed 337 summary statistics. 41 summary statistics were computed within each five populations separately (hence 205 statistics for five populations), 132 summary statistics were computed across the five populations, for each one of the 54 combinations of five populations separately. In brief, all summary statistics were calculated with plink v1.90b4.9 [120], R v3.6.1 [128], Python v2.7.15, bash, awk, and scripts developed by [95], available at https://gitlab.inria.fr/ml_genetics/public/demoseq/-/tree/master, and using the software *asd* (https://github.com/szpiech/asd). The list of calculated summary statistics is provided in **TableS9**. The same statistics were computed for each one of the 54 observed data-sets separately, using the same computational tools and pipeline. Custom-made Python and Bash scripts for computations of all summary-statistics for each simulation and for the observed data are available in the GitHub repository for this article in the "fsc-simulations/code" and in the "observed-sum-stats" folders (https://github.com/Gwennid/africa-wgs-abc).

We used the complete set of 337 statistics to perform Random-Forest ABC scenario-choice, as this method is relatively fast and is unaffected by correlations among statistics [45]. We then considered a subset of 202 statistics among the 337 for Neural Network ABC posterior-parameter estimation, in order to diminish computation time (**TableS9**).

Concerning the Site Frequency Spectrum statistics, note that we inferred the ancestral state using an alignment of three non-human apes (gorilla – reference genome gorGor5, chimpanzee – reference genome panTro6 - and orangutan – reference genome ponAbe3). We considered that the ancestral state of a site could be reliably inferred if the site had the same allele (A, C, T or G) in the three apes. Only these sites were used to calculate the proportion of homozygous ancestral positions per population and the unfolded standardized SFS.

Prior-checking simulations' fit to the observed data

Before conducting ABC scenario-choice and posterior parameter estimation inferences, we checked that summary-statistics calculated on the observed data fell within the range of values of summary-statistics obtained from our simulations. First, we visually verified that each observed vector of summary-statistics computed separately from 54 combinations of five populations each, clustered with the 240,000 vectors of summary-statistics obtained from simulations under the 48 competing scenarios, using two-dimensional PCA calculated with the *prcomp* function R (**FigureS4**). Second, we used the *gfit* function in R to perform goodness-of-fit 100-permutations tests between the 240,000 vectors of summary-statistics obtained from simulations under the 48 competing scenarios and, in turn, each vector of summary-statistics obtained from the 54 observed datasets respectively.

Results both showed that our simulation design could successfully mimic summary-statistics observed in our five-population sample sets, for each 54 combinations of five populations separately. Hence, ABC inferences could be confidently conducted a priori based on the simulations and observed data here considered.

Random Forest ABC grouped scenario-choice

We conducted series of Random Forest ABC scenario-choice procedures for different groups of scenarios [45,63], elaborated specifically to address our different questions of interest regarding gene-flow processes, their intensities, and topological features of the history of Central and Southern African populations. RF-ABC scenario-choice has proven to be performing efficiently and satisfactorily with a significantly lower number of simulations compared to any other ABC scenario-choice procedure. Moreover, RF-ABC scenario choice is insensitive to correlations among summary-statistics [45,63].

Each RF-ABC scenario-choice procedure presented below was conducted using the *predict.abcrf* and the *abcrf* functions with the "group" option of the *abcrf* package in R [45], with 1000 decision trees to train the algorithm after checking that error rates were appropriately low with the *err.abcrf* function (**FigureS5**), separately for each 54 combinations of five sampled populations. Posterior probabilities that the selected scenario, for each analysis separately, is the true scenario, conditional on the classification error a priori is conducted with the abcrf function as explicated in Algorithm 3 in Section 2.4 of [64].

We conducted "out-of-bag" error cross-validation procedures [64], considering in turn each one of the simulations as pseudo-observed data and all remaining simulations to train the algorithm, for each RF-ABC analysis separately (**FigureS6, FigureS7, FigureS8, FigureS9**). While not necessarily predicting the outcome of the scenario-choice for the observed data, these out-of-bag procedures provide us with a sense of the discriminatory power, *a priori*, of RF-ABC for our set of competing-scenarios, as well as empirical levels of nestedness among scenarios or groups of scenarios [92,93]. Finally, we estimated the relative contribution of each one of the 337 summary-statistics to each one of the *a priori* RF-ABC scenario-choice procedure, i.e. the relative importance of each statistics to discriminate among competing scenarios for the entire simulation space using out-of-bag cross-validations on all simulations only without observed data, using the "n.var" option in the *abcrf* plotting function. For simplicity and readability, we provide results for the 50 most important such summary-statistics for each RF-ABC scenario-choice procedure performed in **FigureS10** to **FigureS19**.

Instantaneous or recurring gene-flows in the history of Central and Southern African populations?

The 5,000 simulations performed under each one of the 48 competing scenarios were first gathered into two groups of 24 scenarios each (**Figure5**), corresponding to either instantaneous asymmetric gene-flow processes or to recurring constant asymmetric gene-flow between each pair of lineages, respectively. Both groups thus contained all simulations from the eight competing topologies and all three sets of possible gene-flow intensities (low, moderate, or high, see above), and only differed in the process of gene-flow itself. For each 54 combinations of five sampled populations separately, we performed such RF-ABC scenario-choice to determine the best group of scenarios (**Figure6A**).

Low, moderate, or high gene-flow intensities?

For each 54 combinations of five sampled populations separately, we performed RF-ABC scenario-choice in order to determine which class of intensities of gene-flows best explained the data, everything else (topology and gene-flow process) being-equal. We thus considered three groups of scenarios in our RF-ABC scenario-choice, each corresponding to "low", "moderate", or possibly "high" intensities, and each grouping the 16 scenarios corresponding to eight topologies and instantaneous or recurring gene-flow processes indiscriminately (**Figure6B**).

Then, we conducted, for each 54 population combinations separately, the same RF-ABC
scenario-choice procedure to disentangle groups of gene-flow intensities "all topologies being
equal", considering here only the 24 scenarios from the best group among instantaneous or
recurring gene-flow processes obtained from the above scenario-choice procedure (**FigureS7**).
Finally, we performed, for each 54 combinations of five populations separately, RF-ABC
scenario choice for the 48 competing scenarios grouped in six different groups (encompassing
eight scenarios each), combining instantaneous or recurring processes with the three classes of
intensities respectively (**Figure6C**), thus "intersecting" the two corresponding group analyses
(see above).
1791
Which ancestral Central or Southern African lineage diverged first?
The eight topologies considered in the 48 competing scenarios can also be grouped according
to their ancient topologies, by considering which ancestral lineage diverged first from all others
at the oldest divergence event in the tree (**Figure5**). The ancient lineage which separated first
from the two others can either be the lineage ancestral to Northern and Southern Khoe-San
populations (AKS, scenario topologies 1a, 1b, 1c in **Figure5**), the lineage ancestral to Eastern
and Western Rainforest Hunter-Gatherer populations (ARHG, scenario topologies 2a, 2b, 2c
in **Figure5**), or the lineage leading to Rainforest Hunter-Gatherer Neighboring populations
(RHGn, scenario topologies 3a, 3b in **Figure5**).
For each 54 combinations of five populations separately, we conducted RF-ABC scenario-
choice across these three groups of scenarios, randomly drawing 2/3 of the 5,000 simulations
per scenario for the scenarios 1a, 1b, and 1c, and for the scenarios 2a, 2b, and 2c, respectively,
and kept all simulations for the scenarios 3a and 3b, in order to even the total number of
simulations in competition among the three groups of scenarios. We thus performed RF-ABC
scenario-choice across the three groups of topologies "all gene-flow processes and intensities
being equal" (**Figure6D**).
Then we restricted the RF-ABC scenario-choice of the three-competing groups of
topologies only for the 24 scenarios from the best instantaneous or recurring gene-flow
processes obtained above, all gene-flow intensities being equal (**Figure6E**).
1811
Northern and Southern Khoe-San populations diverged before or after the divergence
between Eastern and Western RHG?
The eight topologies considered in the 48 competing scenarios can alternatively be grouped
according to the relative order of divergence-time between the divergence event among
Northern and Southern KS populations, and that among Eastern and Western RHG
populations; an important question to understand the relative duration of separate evolution of
each group of populations.
To address this specific question we thus conducted, for the 54 combinations of five
populations separately, RF-ABC scenario-choice procedures by grouping the 48 competing
scenarios in two separate groups of 24 scenarios each. One group corresponded to KS
populations diverging from one-another before the RHG divergence (scenario topologies 1b,
2b, 3b, and 1c in **Figure5**), and the other group where RHG diverged from one another before
the KS divergence (scenario topologies 1a, 2a, 3a, and 2c in **Figure5**), instantaneous or
recurring gene-flow and all "low", "moderate", or "high" gene-flow intensities being equal
among the two groups (**Figure6F**). Last, we conducted the same RF-ABC scenario-choice
procedure between two groups of four topologies each, but restricted to the 24 best scenarios
among the instantaneous or recurring gene-flow processes as determined from above
procedures (**Figure6G**).

Finally, we conducted RF-ABC scenario choice procedures considering, respectively, all
48 competing scenarios separately (**FigureS8**), and all 24 best scenarios among the
instantaneous or recurring gene-flow processes as determined above (**FigureS9**).
1833

Neural Network ABC posterior-parameter estimation
We intersected all the RF-ABC scenario-choice results considering the different groups of
scenarios or all scenarios independently, as described above, in order to determine, which
single scenario was explaining data most accurately in the majority of the 54 combinations of
five Central and Southern African populations. We found that the Scenario i1-1b
conservatively produced simulations whose observed summary-statistics most resembled those
obtained in 25 out of the 54 possible population combinations here tested (see **Results**, the
detailed list of 25 population combinations is provided in **TableS12**). The second most often
best-scenario, Scenario i1-3b, only succeeded for four population combinations among the 54
tested.
Based on this *a posteriori* Scenario i1-1b, we performed an additional 95,000 simulations
considering the same prior distributions and constraints among parameters, in order to obtain
100,000 simulations for further ABC posterior parameter inferences, separately for each one
of the 25 population combinations for which it was identified confidently as the winner. As it
remains difficult to estimate jointly the posterior distribution of all parameters of the complex
scenarios here explored with RF-ABC [62], we instead conducted Neural Network ABC joint
posterior parameter inferences using the *abc* package in R [43,44], following best-practices
previously bench-marked for highly complex demographic scenarios [93,95].
There are no evident criteria to choose *a priori* the best tolerance level and numbers of
neurons in the neural network's hidden layer for parameterizing the NN-ABC posterior-
parameter estimation procedure [95,102]. As the total number of parameters in the best Scenario
i1-1b was large (43, **Figure5** and **TableS8**), and as the number of summary-statistics
considered was also large (202, **TableS9**), we chose to conduct the 25 NN-ABC posterior-
parameter inferences for the 25 combinations of sampled populations considering, in-turn, 7,
14, 21, 28, 35, 42, or 43 neurons in the hidden layer (43 being the assumed number of
dimensions equal to the number of parameters for this scenario [43,95]), and a tolerance level of
0.01, thus considering the 1,000 simulations closest to each observed dataset out of the 100,000
performed, respectively. We found *a posteriori* that, considering 43 neurons in the hidden layer
and a maximum allowable number of weights MaxNWts = 12100 provided overall posterior
parameter distributions departing from their priors, and in particular for divergence-times and
gene-flow-times parameters which we were highly interested in, and therefore decided to
provide all posterior parameter distributions' results using this parameterization (tolerance =
0.01, number of hidden neurons = 43, MaxNWts = 12100) for the training of each one of the
neural networks.
We thus performed 25 separate NN-ABC joint parameter posterior inferences, using the
"neuralnet" method option in the function *abc* of the *abc* package in R [44], with logit-
transformed ("logit" transformation option), an "epanechnikov" kernel, parameter values
within parameter-priors' boundaries, a tolerance level of 0.01, 43 neurons in the hidden layer,
and maximum allowable number of weights MaxNWts = 12100. Adjusted posterior parameter
distributions obtained with this method were then plotted for each parameter separately using
a gaussian kernel truncated at the boundaries of the parameter prior-distributions. We then
estimated the mode, median, mean, and 50% and 90% Credibility Intervals of these
distributions *a posteriori*. We provide all posterior-parameter distributions together with their
priors in figure-format (**Figure7**, **Figure8**, **FigureS20, FigureS21**), and in table-format
(**TableS15**, **TableS13**).
1879

Neural Network ABC posterior-parameter estimation cross-validation errors

For each one of the 25 combinations of sampled population separately, we randomly chose 50 simulations (5%) out of the 1000 simulations closest to each observed dataset as determined above. We then conducted NN-ABC joint posterior parameter inference with the same hyper-parameters as above and considering, for each chosen simulation considered as pseudo-observed, the other 99,999 simulations in the reference table, for each one of these 50 pseudo-observed simulations and for each 25 combinations of sampled populations separately (1250 separate NN-ABC joint posterior parameter inferences in total). We then calculated, for each 43 parameter and each 25 populations' combination, the cross-validation posterior parameter estimation error as 1. the mean-squared error of the median NN-ABC posterior estimate $(\hat{\theta}_i)$ scaled by the true-parameter's (θ_i) variance (**TableS14**), $\sum_1^{50}(\hat{\theta}_i - \theta_i)^2/(50 \times Variance(\theta_i))$ [44], and 2. simple average estimation bias of the median NN-ABC posterior estimate $\left(\sum_1^{50}((\hat{\theta}_i - \theta_i)/\theta_i)\right)/50$ (**TableS14**).

Neural Network ABC posterior-parameter estimation posterior checks

We performed 25 separate simulations of Scenario i1-1b each using the same mode point-estimate parameter-values obtained combining (concatenating) the results obtained separately for the 25 different observed combinations of five Central and Southern African populations each and presented in results (**Figure7**, **Figure 8**, **FigureS20**, **FigureS21**, **TableS15** and synthesized schematically in **Figure9**).

First, we aimed to determine *a posteriori* whether our synthetic scenario that best explains genomic patterns in 25 different observed combinations of five Central and Southern African populations each, provided summary-statistics reasonably resembling that of the 25 observed datasets. We performed PCA using the *prcomp* function in R, computed on the 202 summary-statistics used for our NN-ABC posterior-parameter inferences (**TableS9**), calculated on each one of the 25 pseudo-observed simulations and 25 observed datasets similarly as above (**FigureS22**).

Second, beyond the NN-ABC posterior parameter inferred CI and cross-validation error results here obtained, we aimed to empirically estimate whether the variance of posterior parameter estimates obtained among the 25 different combinations of five Central and Southern African populations originated from variation in performances across NN-ABC inferences rather than from different demographic evolutionary histories of the different population combinations considered.

To do so, we conducted, similarly as above for the 25 observed combinations of five populations, NN-ABC joint posterior-parameter inferences for each one of the 25 pseudo-observed simulated datasets under our proposed synthetic scenario (**Figure7**, **Figure 8**, **FigureS20**, **FigureS21**, **TableS15** and synthesized schematically in **Figure9**). We plotted obtained posterior-parameter distributions for each pseudo-observed simulations separately as well as for combined (concatenated) results similarly as for the observed datasets (**FigureS23**, **FigureS24**, **FigureS25**, **FigureS26**). For each parameter separately, we then performed Fligner and F tests of homogeneity of variance, using, respectively, the *fligner.test* and *var.test* functions in R between combined (concatenated) posterior-parameter distributions obtained with 25 pseudo-observed simulations and those obtained previously with 25 observed combinations of five populations (**TableS16**).

REFERENCES

1926

1.   Henn, B. M., Steele, T. E. & Weaver, T. D. Clarifying distinct models of modern human

origins in Africa. *Current Opinion in Genetics & Development* **53**, 148–156 (2018).

2.   Schlebusch, C. M. & Jakobsson, M. Tales of Human Migration, Admixture, and

Selection in Africa. *Annu. Rev. Genom. Hum. Genet.* **19**, 405–428 (2018).

3.   Pfennig, A., Petersen, L. N., Kachambwa, P. & Lachance, J. Evolutionary Genetics and

Admixture in African Populations. *Genome Biology and Evolution* **15**, evad054 (2023).

4.   Verdu, P. *et al.* Origins and Genetic Diversity of Pygmy Hunter-Gatherers from Western

Central Africa. *Current Biology* **19**, 312–318 (2009).

5.   Verdu, P. *et al.* Sociocultural Behavior, Sex-Biased Admixture, and Effective Population

Sizes in Central African Pygmies and Non-Pygmies. *Molecular Biology and Evolution*

30, 918–937 (2013).

6.   Schlebusch, C. M. *et al.* Genomic Variation in Seven Khoe-San Groups Reveals

Adaptation and Complex African History. *Science* **338**, 374–379 (2012).

7.   Breton, G. *et al.* Lactase Persistence Alleles Reveal Partial East African Ancestry of

Southern African Khoe Pastoralists. *Current Biology* **24**, 852–858 (2014).

8.   Patin, E. *et al.* The impact of agricultural emergence on the genetic history of African

rainforest hunter-gatherers and agriculturalists. *Nat Commun* **5**, 3163 (2014).

9.   Patin, E. *et al.* Dispersals and genetic adaptation of Bantu-speaking populations in

Africa and North America. *Science* **356**, 543–546 (2017).

10. Perry, G. H. *et al.* Adaptive, convergent origins of the pygmy phenotype in African

rainforest hunter-gatherers. *Proc. Natl. Acad. Sci. U.S.A.* **111**, (2014).

11. Busby, G. B. *et al.* Admixture into and within sub-Saharan Africa. *eLife* **5**, e15266

(2016).

12. Pierron, D. *et al.* Genomic landscape of human diversity across Madagascar. *Proc.*

Natl. Acad. Sci. U.S.A. **114**, (2017).

13. Semo, A. *et al.* Along the Indian Ocean Coast: Genomic Variation in Mozambique

Provides New Insights into the Bantu Expansion. *Molecular Biology and Evolution* **37**,

406–416 (2020).

14. Lucas-Sánchez, M., Serradell, J. M. & Comas, D. Population history of North Africa

based on modern and ancient genomes. *Human Molecular Genetics* **30**, R17–R23

(2021).

15. Sengupta, D. *et al.* Genetic substructure and complex demographic history of South

African Bantu speakers. *Nat Commun* **12**, 2080 (2021).

16. Fortes-Lima, C. *et al.* Demographic and Selection Histories of Populations Across the

Sahel/Savannah Belt. *Molecular Biology and Evolution* **39**, msac209 (2022).

17. Fortes-Lima, C. A. *et al.* The genetic legacy of the expansion of Bantu-speaking

peoples in Africa. *Nature* **625**, 540–547 (2024).

18. Laurent, R. *et al.* A genetic and linguistic analysis of the admixture histories of the

islands of Cabo Verde. *eLife* **12**, e79827 (2023).

19. Bergström, A., Stringer, C., Hajdinjak, M., Scerri, E. M. L. & Skoglund, P. Origins of

modern human ancestry. *Nature* **590**, 229–237 (2021).

20. Stringer, C. Modern human origins: progress and prospects. *Phil. Trans. R. Soc. Lond.*

B **357**, 563–579 (2002).

21. Hollfelder, N., Breton, G., Sjödin, P. & Jakobsson, M. The deep population history in

Africa. *Human Molecular Genetics* **30**, R2–R10 (2021).

22. Ragsdale, A. P. *et al.* A weakly structured stem for human origins in Africa. *Nature* **617**,

755–763 (2023).

23. Lachance, J. *et al.* Evolutionary History and Adaptation from High-Coverage Whole-

Genome Sequences of Diverse African Hunter-Gatherers. *Cell* **150**, 457–469 (2012).

24. Lorente-Galdos, B. *et al.* Whole-genome sequence analysis of a Pan African set of

samples reveals archaic gene flow from an extinct basal population of modern humans

into sub-Saharan populations. *Genome Biol* **20**, 77 (2019).

25. Lipson, M. *et al.* Ancient DNA and deep population structure in sub-Saharan African

foragers. *Nature* **603**, 290–296 (2022).

26. Fan, S. *et al.* Whole-genome sequencing reveals a complex African population

demographic history and signatures of local adaptation. *Cell* **186**, 923-939.e14 (2023).

27. Meyer, M. *et al.* A High-Coverage Genome Sequence from an Archaic Denisovan

Individual. *Science* **338**, 222–226 (2012).

28. Prüfer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai

Mountains. *Nature* **505**, 43–49 (2014).

29. Tishkoff, S. A. *et al.* The Genetic Structure and History of Africans and African

Americans. *Science* **324**, 1035–1044 (2009).

30. Skoglund, P. & Mathieson, I. Ancient Genomics of Modern Humans: The First Decade.

Annu. Rev. Genom. Hum. Genet. **19**, 381–404 (2018).

31. Gravel, S. Population Genetics Models of Local Ancestry. *Genetics* **191**, 607–619

(2012).

32. Ragsdale, A. P. & Gravel, S. Models of archaic admixture and recent history from two-

locus statistics. *PLoS Genet* **15**, e1008204 (2019).

33. Kamm, J., Terhorst, J., Durbin, R. & Song, Y. S. Efficiently Inferring the Demographic

History of Many Populations With Allele Count Data. *Journal of the American Statistical*

Association **115**, 1472–1487 (2020).

34. Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from 142 diverse

populations. *Nature* **538**, 201–206 (2016).

35. The 1000 Genomes Project Consortium *et al.* A global reference for human genetic

variation. *Nature* **526**, 68–74 (2015).

36. Rasmussen, M. *et al.* The genome of a Late Pleistocene human from a Clovis burial

site in western Montana. *Nature* **506**, 225–229 (2014).

37. Choudhury, A. *et al.* Whole-genome sequencing for an enhanced understanding of

genetic variation among South Africans. *Nat Commun* **8**, 2062 (2017).

38. *Hunter-Gatherers of the Congo Basin: Cultures, Histories and Biology of African Pygmies*. (Routledge, Taylor and Francis Group, London New York, 2017).

39. Li, J. Z. *et al.* Worldwide Human Relationships Inferred from Genome-Wide Patterns of Variation. *Science* **319**, 1100–1104 (2008).

40. Schlebusch, C. M. *et al.* Khoe-San Genomes Reveal Unique Variation and Confirm the Deepest Population Divergence in Homo sapiens. *Molecular Biology and Evolution* **37**, 2944–2954 (2020).

41. Tavaré, S., Balding, D. J., Griffiths, R. C. & Donnelly, P. Inferring coalescence times from DNA sequence data. *Genetics* **145**, 505–518 (1997).

42. Beaumont, M. A., Zhang, W. & Balding, D. J. Approximate Bayesian computation in population genetics. *Genetics* **162**, 2025–2035 (2002).

43. Blum, M. G. B. & François, O. Non-linear regression models for Approximate Bayesian Computation. *Stat Comput* **20**, 63–73 (2010).

44. Csilléry, K., François, O. & Blum, M. G. B. abc: an R package for approximate Bayesian computation (ABC): *R package: abc. Methods in Ecology and Evolution* **3**, 475–479 (2012).

45. Pudlo, P. *et al.* Reliable ABC model choice via random forests. *Bioinformatics* **32**, 859–866 (2016).

46. Breton, G., Fortes-Lima, C. & Schlebusch, C. M. Revisiting the demographic history of Central African populations from a genetic perspective. *Hum Popul Genet Genom* 1–29 (2021) doi:10.47248/hpgg2101010004.

47. Nei, M. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* **89**, 583–590 (1978).

48. Pemberton, T. J. *et al.* Genomic Patterns of Homozygosity in Worldwide Human Populations. *The American Journal of Human Genetics* **91**, 275–292 (2012).

49. Mooney, J. A. *et al.* Understanding the Hidden Complexity of Latin American Population Isolates. *The American Journal of Human Genetics* **103**, 707–726 (2018).

50. Szpiech, Z. A. *et al.* Ancestry-Dependent Enrichment of Deleterious Homozygotes in
Runs of Homozygosity. *The American Journal of Human Genetics* **105**, 747–762
(2019).

51. Bowcock, A. M. *et al.* Drift, admixture, and selection in human evolution: a study with
DNA polymorphisms. *Proceedings of the National Academy of Sciences* **88**, 839–843
(1991).

52. Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Molecular Biology and Evolution*
https://doi.org/10.1093/oxfordjournals.molbev.a040454 (1987)
doi:10.1093/oxfordjournals.molbev.a040454.

53. Gascuel, O. BIONJ: an improved version of the NJ algorithm based on a simple model
of sequence data. *Molecular Biology and Evolution* **14**, 685–695 (1997).

54. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research* **19**, 1655–1664 (2009).

55. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using
multilocus genotype data. *Genetics* **155**, 945–959 (2000).

56. Rosenberg, N. A. Genetic Structure of Human Populations. *Science* **298**, 2381–2385
(2002).

57. Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using
multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**,
1567–1587 (2003).

58. Lawson, D. J., van Dorp, L. & Falush, D. A tutorial on how not to over-interpret
STRUCTURE and ADMIXTURE bar plots. *Nat Commun* **9**, 3258 (2018).

59. Peter, B. M. A geometric relationship of F_2, F_3 and F_4-statistics with principal
component analysis. *Phil. Trans. R. Soc. B* **377**, 20200413 (2022).

60. Behr, A. A., Liu, K. Z., Liu-Fang, G., Nakka, P. & Ramachandran, S. pong: fast analysis
and visualization of latent clusters in population genetic data. *Bioinformatics* **32**, 2817–
2823 (2016).

61.   Pritchard, J. K., Seielstad, M. T., Perez-Lezaun, A. & Feldman, M. W. Population
growth of human Y chromosomes: a study of Y chromosome microsatellites. *Molecular*
Biology and Evolution **16**, 1791–1798 (1999).

62.   Raynal, L. *et al.* ABC random forests for Bayesian parameter inference. *Bioinformatics*
35, 1720–1728 (2019).

63.   Estoup, A., Raynal, L., Verdu, P. & Marin, J.-M. Model choice using Approximate
Bayesian Computation and Random Forests: analyses based on model grouping to
make inferences about the genetic history of Pygmy human populations. *Journal de la*
Société Française de Statistique 167–190 (2018).

64.   Pudlo, P. *et al.* Reliable ABC model choice via random forests. *Bioinformatics* **32**, 859–
866 (2016).

65.   Seidensticker, D. *et al.* Population collapse in Congo rainforest from 400 CE urges
reassessment of the Bantu Expansion. *Sci. Adv.* **7**, eabd8352 (2021).

66.   Verdu, P. & Rosenberg, N. A. A General Mechanistic Model for Admixture Histories of
Hybrid Populations. *Genetics* **189**, 1413–1426 (2011).

67.   Harris, K. & Nielsen, R. The Genetic Cost of Neanderthal Introgression. *Genetics* **203**,
881–891 (2016).

68.   Arredondo, A. *et al.* Inferring number of populations and changes in connectivity under
the n-island model. *Heredity* **126**, 896–912 (2021).

69.   Currat, M. & Excoffier, L. Modern Humans Did Not Admix with Neanderthals during
Their Range Expansion into Europe. *PLoS Biol* **2**, e421 (2004).

70.   Tournebize, R. & Chikhi, L. Ignoring population structure in hominin evolutionary
models can lead to the inference of spurious admixture events. *Nat Ecol Evol* **9**, 225–
236 (2024).

71.   Mazet, O., Rodríguez, W., Grusea, S., Boitard, S. & Chikhi, L. On the importance of
being structured: instantaneous coalescence rates and human evolution—lessons for
ancestral population size inference? *Heredity* **116**, 362–371 (2016).

72. Hamid, I., Korunes, K. L., Beleza, S. & Goldberg, A. Rapid adaptation to malaria

facilitated by admixture in the human population of Cabo Verde. *eLife* **10**, e63177

(2021).

73. Cuadros-Espinoza, S., Laval, G., Quintana-Murci, L. & Patin, E. The genomic

signatures of natural selection in admixed human populations. *The American Journal of*

Human Genetics **109**, 710–726 (2022).

74. Jakobsson, M. *et al.* Homo sapiens-specific evolution unveiled by ancient southern

African genomes. *Nature* https://doi.org/10.1038/s41586-025-09811-4 (2025)

doi:10.1038/s41586-025-09811-4.

75. Patin, E. *et al.* Inferring the Demographic History of African Farmers and Pygmy

Hunter–Gatherers Using a Multilocus Resequencing Data Set. *PLoS Genet* **5**,

e1000448 (2009).

76. Beyer, R. M., Krapp, M., Eriksson, A. & Manica, A. Climatic windows for human

migration out of Africa in the past 300,000 years. *Nat Commun* **12**, 4889 (2021).

77. Lopez, M. *et al.* The demographic history and mutational load of African hunter-

gatherers and farmers. *Nat Ecol Evol* **2**, 721–730 (2018).

78. Bartlein, P. J. *et al.* Pollen-based continental climate reconstructions at 6 and 21 ka: a

global synthesis. *Clim Dyn* **37**, 775–802 (2011).

79. Perry, G. H. & Verdu, P. Genomic perspectives on the history and evolutionary ecology

of tropical rainforest occupation by humans. *Quaternary International* **448**, 150–157

(2017).

80. Ziegler, M. *et al.* Development of Middle Stone Age innovation linked to rapid climate

change. *Nat Commun* **4**, 1905 (2013).

81. Scerri, E. M. L. *et al.* Did Our Species Evolve in Subdivided Populations across Africa,

and Why Does It Matter? *Trends in Ecology & Evolution* **33**, 582–594 (2018).

82. Gosling, W. D., Scerri, E. M. L. & Kaboth-Bahr, S. The climate and vegetation backdrop

to hominin evolution in Africa. *Phil. Trans. R. Soc. B* **377**, 20200483 (2022).

83. Thomas, D. S. G. *et al.* Lacustrine geoarchaeology in the central Kalahari: Implications for Middle Stone Age behaviour and adaptation in dryland conditions. *Quaternary Science Reviews* **297**, 107826 (2022).

84. Long, J. C. The genetic structure of admixed populations. *Genetics* **127**, 417–428 (1991).

85. Cornelissen, E. [No title found]. *Journal of World Prehistory* **16**, 197–235 (2002).

86. Mesfin, I., Oslisly, R. & Forestier, H. Technological analysis of the quartz industry of Maboué 5 – Layer 3 (Lopé National Park, Gabon): Implications for the Late Stone Age emergence in western Central Africa. *Journal of Archaeological Science: Reports* **39**, 103130 (2021).

87. Bader, G. D., Mabuza, A., Price Williams, D. & Will, M. Rethinking the Middle to Later Stone Age transition in southern Africa - A perspective from the highveld of Eswatini. *Quaternary Science Reviews* **286**, 107540 (2022).

88. Lézine, A.-M., Izumi, K., Kageyama, M. & Achoundong, G. A 90,000-year record of Afromontane forest responses to climate change. *Science* **363**, 177–181 (2019).

89. Phillipson, D. W. *African Archaeology*. (Cambridge University Press, 2005). doi:10.1017/CBO9780511800313.

90. Bostoen, K. *et al.* Middle to Late Holocene Paleoclimatic Change and the Early Bantu Expansion in the Rain Forests of Western Central Africa. *Current Anthropology* **56**, 354–384 (2015).

91. Lopez, M. *et al.* Genomic Evidence for Local Adaptation of Hunter-Gatherers to the African Rainforest. *Current Biology* **29**, 2926-2935.e4 (2019).

92. Robert, C. P., Mengersen, K. & Chen, C. Model choice versus model criticism. *Proceedings of the National Academy of Sciences* **107**, E5–E5 (2010).

93. Fortes-Lima, C. A., Laurent, R., Thouzeau, V., Toupance, B. & Verdu, P. Complex genetic admixture histories reconstructed with Approximate Bayesian Computation. *Mol Ecol Resour* **21**, 1098–1117 (2021).

94. Boitard, S., Rodríguez, W., Jay, F., Mona, S. & Austerlitz, F. Inferring Population Size

History from Large Samples of Genome-Wide Molecular Data - An Approximate

Bayesian Computation Approach. *PLoS Genet* **12**, e1005877 (2016).

95. Jay, F., Boitard, S. & Austerlitz, F. An ABC Method for Whole-Genome Sequence Data:

Inferring Paleolithic and Neolithic Human Expansions. *Molecular Biology and Evolution*

36, 1565–1579 (2019).

96. Verdu, P. *et al.* Sociocultural Behavior, Sex-Biased Admixture, and Effective Population

Sizes in Central African Pygmies and Non-Pygmies. *Molecular Biology and Evolution*

30, 918–937 (2013).

97. Patin, E. *et al.* The impact of agricultural emergence on the genetic history of African

rainforest hunter-gatherers and agriculturalists. *Nat Commun* **5**, 3163 (2014).

98. Patin, E. *et al.* Dispersals and genetic adaptation of Bantu-speaking populations in

Africa and North America. *Science* **356**, 543–546 (2017).

99. Batini, C. *et al.* Insights into the Demographic History of African Pygmies from

Complete Mitochondrial Genomes. *Molecular Biology and Evolution* **28**, 1099–1110

(2011).

100. Wang, H., Czerminski, R. & Jamieson, A. C. Neural Networks and Deep Learning. in

The Machine Age of Customer Insight (eds Einhorn, M., Löffler, M., De Bellis, E.,

Herrmann, A. & Burghartz, P.) 91–101 (Emerald Publishing Limited, 2021).

doi:10.1108/978-1-83909-694-520211010.

101. Leung, F. H. F., Lam, H. K., Ling, S. H. & Tam, P. K. S. Tuning of the structure and

parameters of a neural network using an improved genetic algorithm. *IEEE Trans.*

Neural Netw. **14**, 79–88 (2003).

102. Huang, X., Rymbekova, A., Dolgova, O., Lao, O. & Kuhlwilm, M. Harnessing deep

learning for population genetic inference. *Nat Rev Genet* **25**, 61–78 (2024).

103. Mooney, J. A., Agranat-Tamir, L., Pritchard, J. K. & Rosenberg, N. A. On the number of

genealogical ancestors tracing to the source groups of an admixed population.

GENETICS **224**, iyad079 (2023).

104. Agranat-Tamir, L., Mooney, J. A. & Rosenberg, N. A. Counting the genetic ancestors
from source populations in members of an admixed population. *GENETICS* **226**,
iyae011 (2024).

105. Murtagh, F. Multilayer perceptrons for classification and regression. *Neurocomputing* **2**,
183–197 (1991).

106. Chen, J., Zhao, F., Sun, Y. & Yin, Y. Improved XGBoost model based on genetic
algorithm. *IJCAT* **62**, 240 (2020).

107. Yelmen, B. & Jay, F. An Overview of Deep Generative Models in Functional and
Evolutionary Genomics. *Annu. Rev. Biomed. Data Sci.* **6**, 173–189 (2023).

108. Skoglund, P. *et al.* Reconstructing Prehistoric African Population Structure. *Cell* **171**,
59-71.e21 (2017).

109. Vicente, M. & Schlebusch, C. M. African population history: an ancient DNA
perspective. *Current Opinion in Genetics & Development* **62**, 8–15 (2020).

110. Schlebusch, C. Issues raised by use of ethnic-group names in genome study. *Nature*
464, 487–487 (2010).

111. Miller, S. A., Dykes, D. D. & Polesky, H. F. A simple salting out procedure for extracting
DNA from human nucleated cells. *Nucl Acids Res* **16**, 1215–1215 (1988).

112. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

113. DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat Genet* **43**, 491–498 (2011).

114. Van Der Auwera, G. A. *et al.* From FastQ Data to High-Confidence Variant Calls: The
Genome Analysis Toolkit Best Practices Pipeline. *CP in Bioinformatics* **43**, (2013).

115. Breton, G., Johansson, A. C. V., Sjödin, P., Schlebusch, C. M. & Jakobsson, M.
Comparison of sequencing data processing pipelines and application to
underrepresented African human populations. *BMC Bioinformatics* **22**, 488 (2021).

116. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

117. Schneider, V. A. *et al.* Evaluation of GRCh38 and de novo haploid genome assemblies

demonstrates the enduring quality of the reference assembly. *Genome Res.* **27**, 849–

864 (2017).

118. Mills, R. E. *et al.* Natural genetic variation caused by small insertions and deletions in

the human genome. *Genome Res.* **21**, 830–839 (2011).

119. Manichaikul, A. *et al.* Robust relationship inference in genome-wide association

studies. *Bioinformatics* **26**, 2867–2873 (2010).

120. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-

Based Linkage Analyses. *The American Journal of Human Genetics* **81**, 559–575

(2007).

121. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

122. Okonechnikov, K., Conesa, A. & García-Alcalde, F. Qualimap 2: advanced multi-

sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294

(2016).

123. Fenner, J. N. Cross-cultural estimation of the human generation interval for use in

genetics-based population divergence studies. *Am. J. Phys. Anthropol.* **128**, 415–423

(2005).

124. Hublin, J.-J. *et al.* New fossils from Jebel Irhoud, Morocco and the pan-African origin of

Homo sapiens. *Nature* **546**, 289–292 (2017).

125. Richter, D. *et al.* The age of the hominin fossils from Jebel Irhoud, Morocco, and the

origins of the Middle Stone Age. *Nature* **546**, 293–296 (2017).

126. Excoffier, L. & Foll, M. fastsimcoal: a continuous-time coalescent simulator of genomic

diversity under arbitrarily complex evolutionary scenarios. *Bioinformatics* **27**, 1332–

1334 (2011).

127. Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V. C. & Foll, M. Robust

Demographic Inference from Genomic and SNP Data. *PLoS Genet* **9**, e1003905

(2013).

128. R Core Team. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing (2015).

2228

2229

2230

Acknowledgments

We are grateful to all subjects who participated in this research. The computations were performed at the Swedish National Infrastructure for Computing (SNIC-UPPMAX). Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. We thank the platform Paléogénomique et Génétique Moléculaire (P2GM) of the MNHN at the Musée de l'Homme in Paris for helping to prepare, in part, the Central African samples for sequencing. We thank the Working Group of Indigenous Minorities in Southern Africa (WIMSA) and the South African San Council for their support and facilitating fieldwork. The project was approved by the South African San Council.

Funding

GB was funded in part by a grant from the Wilhelm och Martina Lundgrens Vetenskapsfond (2023-SA-4327). CS is funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 759933) and the Knut and Alice Wallenberg foundation. MJ is funded by the Knut and Alice Wallenberg foundation, and the Vetenskapsrådet/Swedish Research Council (no: 642-2013-8019 and 2018-05537). PV was funded in part by the UMR7206 Eco-anthropology and the French Agence Nationale de la Recherche ANR-METHIS (15-CE32-0009-1).

Authors' contributions

GB – Designed the study, conducted data curation and quality controls, conducted statistical descriptions, conducted simulations and ABC inferences, interpreted the results, wrote the original draft.

PS – Designed the study, conducted statistical descriptions, interpreted the results, participated in writing and editing.

PLZ – Conducted data curation and quality controls, conducted statistical descriptions, participated in editing.

RL –Conducted simulations and ABC inferences, participated in writing and editing.

AF – Conducted fieldwork and provided samples, participated in editing.

AES – Conducted fieldwork and provided samples, participated in editing.

BSH – Conducted fieldwork and provided samples, participated in editing.

LBB – Conducted fieldwork and provided samples, participated in editing.

GHP – Conducted fieldwork and provided samples, participated in editing.

HS – Conducted fieldwork and provided samples, participated in editing.

EH – Acquired funding, interpreted the results, participated in editing.

CMS – Supervised the project, designed the study, acquired funding, conducted fieldwork and provided samples, interpreted the results, participated in writing and editing.

MJ – Supervised the project, designed the study, acquired funding, interpreted the results, participated in writing and editing.

PV – Supervised the project, designed the study, acquired funding, conducted fieldwork and provided samples, conducted statistical descriptions, conducted simulations and ABC inferences, interpreted the results, wrote the original draft.

Competing interests

The authors declare no competing nor conflict of interests.

GitHub repositories for material availability

2278 https://github.com/Gwennid/africa-wgs-descriptive
2279 https://github.com/Gwennid/africa-wgs-abc
2280

Data availability statement

All raw whole-genome sequence FASTQ data original to this study are made available on the
European Genome-Phenome Archive (https://ega-archive.org/), provided that requests
conform to ethical requirements and informed consents provided by donors, as detailed in the
corresponding Data Access Policies associated with each data repository. Data access requests
are evaluated by the associated Data Access Committees. EGA accession numbers:
EGAD50000001559 and EGAD50000001560.
2288

Ethical statement

DNA samples from individuals were collected with the subjects' informed consent following
the Declaration of Helsinki guidelines and approved by the following Ethics committees:
Institut de Recherche pour le Développement (IRD, France); Cameroon Ministry of Scientific
Research and Technology; Central African Republic Ministère de l'Enseignement Supérieur
et de la Recherche Scientifique; France Ministère de l'Enseignement Supérieur, de la
Recherche et de l'Innovation (Comité de Protection des Personnes DC-2016-2740); French
Commission Nationale Informatique et Liberté CNIL (declaration n°1972648); Uganda
National Council for Science and Technology; Institutional Review Board of the University of
Chicago Biological Sciences Division and the Pritzker School of Medicine University of
Chicago Hospitals (protocol number 16986A); Committee for the Protection of Human
Subjects of the Trustees of Dartmouth College and the Dartmouth-Hitchcock Medical Center
(protocol number 22410); Human Research Ethics Committee (Medical) at the University of
the Witwatersrand, Johannesburg (Protocol Numbers: M980553, M050902, M090576,
M10270, M180654); Working Group of Indigenous Minorities in Southern Africa (WIMSA);
South African San Council (SASC); and the Swedish Ethical Review Authority (Dnr 2019-
05174).
2306
2307

Supplementary Material

27 Supplementary Figures and 16 Supplementary Tables

FigureS1: **Correlations between genome-wide autosomal number of biallelic SNPs, unbiased estimates of heterozygosities and total ROH lengths at the world-wide scale.**
Median values across individuals within populations reported in **Figure2** panels A-C are used here in panel **A**) Unbiased estimates of heterozygosity vs. total number of autosomal biallelic SNPs, and in panel **B**) Unbiased estimates of heterozygosity vs. total autosomal length found in ROH. As expected, we found a significant positive correlation in Panel **A** (Spearman rho=0.9490, p-value<2.2x10^{-16}), and a significant negative correlation in Panel **B** (Spearman rho=-0.59128, p-value=5.243x10^{-5}).

2324 **FigureS2: Individual pairwise genetic differentiation patterns at the world-wide scale.**
2325 Multi-Dimensional scaling representation of genome-wide Allele Sharing Dissimilarities [51] among pairs of individuals at
2326 worldwide scale. We considered the 14,182,615 genome-wide autosomal SNPs pruned for low LD (r^2 threshold 0.1) to
2327 calculate ASD between all pairs of individuals. This figure complements the NJ representation provided in **Figure3** panel A.
2328
2329
2330

FigureS3: Alternative ADMIXTURE modal results for K=4 to 10.
Modal results each containing the highest number of runs among the 20 independent runs performed for each value of K are
provided in main **Figure4**. All alternative modal results are provided here. See **Material and Methods** for details.

FigureS4: PCA for ABC prior-checking of simulations fit to the observed data.
We calculated 337 summary-statistics for each one of the 240 000 simulated data sets, 5000 under each one of the 48 competing
scenarios (**Figure5** and **Material and Methods**), which we projected on the first two axes of a principal component analysis.
Each vector (corresponding thus to a single simulation) is represented by a gray cross. We separately computed the same 337
summary-statistics, separately for each one of the 54 sets of five observed Central and Southern African populations included
in our analyses. A) Each observed vector is then projected, in turn, on the PCA obtained from simulations only. Each such
observed vector is represented as a single yellow dot. B) Alternatively, a PCA is performed on all simulated and observed
vectors of summary statistics together. Each observed vector is also represented as a single yellow dot in panel B.
We can see that, in both cases, the 54 observed sets of populations each fall well within the space of simulated dataset, thus
allowing us a priori to conduct machine-learning ABC scenario choice and posterior parameter estimation procedures.
2347
2348
2349

FigureS5: Cross-validation prior error of Random-Forest ABC scenario-choice procedures as a function of the number
of trees in the forest.
For panels A to G, out-of-bag cross-validation prior error results for each RF-ABC scenario choice analysis are shown in
FigureS6 panels A to G respectively. For panel H, the corresponding out-of-bag cross-validation prior error results are shown
in **FigureS7 panel B.** For panel I, the corresponding out-of-bag cross-validation prior error results are shown in **FigureS8**
panel B. For panel J, the corresponding out-of-bag cross-validation prior error results are shown in **FigureS9 panel B.**
2356
2357
2358

FigureS6: Out-of-bag cross-validation errors for Random-Forest ABC scenario-choices among 48 competing scenarios.
All RF-ABC cross-validation analyses are conducted *a priori*, without using any observed data, by considering, in-turn, each
simulation as observed data while all remaining simulations are used to train the Random Forest. Each panel provides the *a*
priori cross-validation errors for the corresponding RF-ABC scenario-choice analysis detailed in each panel in **Figure6**. While
these cross-validation errors do not predict the specific outcome of RF-ABC prediction for the observed data, they provide
levels of scenarios and groups of scenarios nestedness *a priori* for the entire space of parameters used for simulations. (**A**)
shows that intermediate and recurring gene-flow processes are, all topologies and gene-flow intensities "being equal", clearly
distinguishable by RF-ABC and little nested. (**B**) and (**C**) show that different classes of gene-flow intensities are also relatively
clearly identifiable, albeit more nested, as expected by design since the "high gene-flow" class comprises parameter values of
the moderate and the low intensity classes of gene-flows (**Material and Methods**, **TableS8**). (**D**) and (**E**) show that ancient
tree topologies are relatively well distinguishable *a priori* all gene-flow processes "being equal", despite some amounts of
errors due to the expected nestedness in the space of parameter values where ancient divergence times are resembling and thus
undistinguishable. (**F**) and (**G**) show that the chronological order between the divergence-time of Northern and Southern KS
lineages, and that of Western and Eastern RHG lineages, are also relatively well distinguishable a priori with RF-ABC
scenario-choice, albeit with an increased cross-validation error expected due to high levels of nestedness in the spaces of
parameter-values where both separate events may occur at similar times.
2375

<u>**FigureS7**</u>**: RF-ABC scenario choice among 24 competing scenarios with instantaneous asymmetric gene-flow processes,**
for three groups of gene-flow intensities, all topologies being equal.
(**A**) corresponds to the scenario-choice results for the same test as in **Figure6C**, restricted to the 24 competing scenarios
considering instantaneous gene-flow processes only, all scenario topologies "being equal". Description of the x and y axis
legends are provided in **Figure6**. Posterior probabilities of choosing the correct scenarios for each 54 sets of population
combinations ranged from 0.63935 to 0.87000, compared to a prior probability of 1/3 in this particular test. (**B**) corresponds
to the RF-ABC cross-validation prior error for the scenario-choice procedure conducted in (**A**). Errors were obtained without
considering observed data and using, instead, all simulations in turn as pseudo-observed data and the remaining simulations
to train the RF.
2385

FigureS8: RF-ABC scenario choice among 48 competing scenarios without groups.
(**A**) corresponds to the RF-ABC scenario-choice results obtained when considering all 48 scenarios (5,000 simulations per
scenario), as separate competitors without grouping them, for each 54 combinations of five sampled-populations separately.
Description of the x and y axis legends are provided in **Figure6**. Posterior probabilities of choosing the correct scenarios for
each 54 sets of population combinations ranged from 0.20553 to 0.28588, compared to a prior probability of 1/48 in this
particular test. (**B**) corresponds to the RF-ABC cross-validation prior error for the scenario-choice procedure conducted in (**A**).
Errors were obtained without considering observed data and using, instead, all simulations in turn as pseudo-observed data
and the remaining simulations to train the RF. Note that, in (**B**), whichever the intensity of the gene-flow, when considering
only recurring gene-flow processes, topologies are harder to distinguish from one another a priori. Conversely, such difficulty
to distinguish a priori among topologies remains overall limited for instantaneous gene-flow processes, and errors increase
only when considering the possibility of intense instantaneous gene-flows. However, as mentioned throughout the article, the
capacity to distinguish a priori among scenarios in the entire space of summary-statistics values does not predict the power to
predict a best scenario in the specific space occupied by observed data [45].

2399

2400

2401

FigureS8: RF-ABC scenario choice among 48 competing scenarios without groups.

FigureS9: RF-ABC scenario choice among 24 competing scenarios without groups.
(**A**) corresponds to the RF-ABC scenario-choice results obtained when considering all 24 instantaneous gene-flow scenarios
(5,000 simulations per scenario), as separate competitors without grouping them, for each 54 combinations of five sampled-
populations separately. Description of the x and y axis legends are provided in **Figure6**. Posterior probabilities of choosing
the correct scenarios for each 54 sets of population combinations ranged from 0.22380 to 0.30273, compared to a prior
probability of 1/24 in this particular test. (**B**) corresponds to the RF-ABC cross-validation prior error for the scenario-choice
procedure conducted in (**A**). Errors were obtained without considering observed data and using, instead, all simulations in turn
as pseudo-observed data and the remaining simulations to train the RF. Note that prediction and cross-validation error results
resemble those obtained for these 24 specific scenarios obtained when considering instead all the 48 scenarios in competition
(**FigureS8**).
2414

2415   **FigureS10: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based**
2416   **on simulations only without observed data – Instantaneous vs Recurring gene-flow processes all gene-flow intensities**
2417   **and all topologies "being equal".**
2418   Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel A.**
2419
2420
2421

RF-ABC scenario-choice
Instantaneous vs. *Recurring* gene-flow
48 competing scenarios - 2 groups
Out-of-bag cross-validation *a priori*
240 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

2422 **FigureS11**: **50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based**
2423 **on simulations only without observed data – Low vs Moderate vs High gene-flow intensities, all gene-flow processes**
2424 **(instantaneous and recurring) and all topologies "being equal".**
2425 Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel B.**

2431 **FigureS12**: **50 best summary statistics relative contribution to the Random Forest ABC scenario-choice** *a priori* **based**
2432 **on simulations only without observed data – Instantaneous vs Recurring & Low vs Moderate vs High gene-flow**
2433 **intensities, all topologies "being equal".**
2434 Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel C.**
2435
2436
2437

RF-ABC scenario-choice
Gene-flows processes & intensities
48 competing scenarios - 6 groups
Out-of-bag cross-validation *a priori*
240 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

2438 **FigureS13**: **50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based**
2439 **on simulations only without observed data – Ancient topologies, all gene-flow processes (instantaneous and recurring)**
2440 **and intensities (Low, moderate, and high), and all recent topologies "being equal".**
2441 Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel D.**
2442
2443
2444
2445

RF-ABC scenario-choice
Ancient topologies
48 competing scenarios - 3 groups
Out-of-bag cross-validation *a priori*
180 000 simulations: all sims per sce. 3a-b,
2/3 sims per all other scenarios
Cross-validation simulations only, no observed data

FigureS14: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based on simulations only without observed data – Ancient topologies for instantaneous gene-flow processes only, all intensities (Low, moderate, and high), and all recent topologies "being equal".
Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel E.**

RF-ABC scenario-choice
Ancient topologies
24 competing scenarios - 3 groups
Out-of-bag cross-validation *a priori*
90 000 simulations: all sims per sce. 3a-b,
2/3 sims per all other scenarios
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

FigureS15: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based on simulations only without observed data – Internal divergence order, all gene-flow processes (instantaneous and recurring) and intensities (Low, moderate, and high), and all ancient topologies "being equal".
Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel F.**

RF-ABC scenario-choice
Internal divergence order
48 competing scenarios - 2 groups
Out-of-bag cross-validation *a priori*
240 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

2462 **FigureS16: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based**
2463 **on simulations only without observed data – Internal divergence order for instantaneous gene-flow processes only, all**
2464 **intensities (Low, moderate, and high), and all ancient topologies "being equal".**
2465 Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS6 panel G.**
2466
2467
2468
2469

RF-ABC scenario-choice
Internal divergence order
24 competing scenarios - 2 groups
Out-of-bag cross-validation *a priori*
120 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

FigureS17: **50 best summary statistics relative contribution to the Random Forest ABC scenario-choice** *a priori* **based on simulations only without observed data – Low vs Moderate vs High gene-flow intensities for instantaneous gene-flow processes only, all topologies "being equal".**
Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS7 panel B.**

2478 **FigureS18: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based**
2479 **on simulations only without observed data – All models in competition.**
2480 Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS8 panel B.**
2481
2482
2483
2484

RF-ABC scenario-choice
Topologies and Low vs. ***Moderate*** vs ***High***
Instantaneous and Recurring gene-flows
48 competing scenarios - no groups
240 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

FigureS19: 50 best summary statistics relative contribution to the Random Forest ABC scenario-choice *a priori* based on simulations only without observed data – All models in competition for instantaneous gene-flow processes only. Out-of-bag cross-validation prior error results for this RF-ABC scenario choice analysis are shown in **FigureS9 panel B.**

RF-ABC scenario-choice
Topologies and Low vs. ***Moderate*** vs ***High***
Instantaneous gene-flows
24 competing scenarios - no groups
120 000 simulations: 5000 per scenario
Cross-validation simulations only, no observed data

Summary statistics Importance
in RF-ABC scenario choice *a priori* (50 best)

FigureS20: ABC posterior distribution of effective population size parameters.

Neural Network Approximate Bayesian Computation [43,44], posterior parameter joint estimations of Effective population sizes *Ne* (in generations before present) for 25 sets of five Central and Southern African populations for which the best scenario identified by RF-ABC was Scenario i1-1b (**Figure5**, **TableS12**). NN ABC posterior parameter estimation procedures were conducted using 100,000 simulations under Scenario i1-1b, each simulation corresponding to a single vector of parameter values drawn randomly from prior distributions provided in **TableS8**. We considered 43 neurons in the hidden layer of the NN and a tolerance level of 0.01, corresponding to the 1,000 simulations providing summary-statistics closest to the observed ones, for each 25 separate analyses. NN posterior estimates are based on the logit transformation of parameter values using an Epanechnikov kernel between the corresponding parameter's prior bounds (see **Material and Methods** and **TableS8**). Posterior parameter densities are represented with solid blue lines. 50% Credibility Intervals are represented as the light blue area under the density. The median and mode values are represented as a solid and dotted blue vertical line, respectively. Parameter prior distributions are represented as dotted grey lines. For all panels, the left plots represent the NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern African populations under Scenario i1-1b, separately (**TableS12** and **TableS13**). For all panels, the right plots represent a single parameter posterior distribution obtained from combining (concatenating) the 25 posterior distributions together. See **Figure5** and **TableS8** for all parameters' descriptions in each panel. Results are also provided in **TableS15**.

FigureS21: ABC posterior distribution of potentially asymmetric instantaneous gene-flow intensity parameters.
Neural Network Approximate Bayesian Computation [43,44], posterior parameter joint estimations of gene-flow intensity
parameters *m* (in generations before present) for 25 sets of five Central and Southern African populations for which the best
scenario identified by RF-ABC was Scenario i1-1b (**Figure5**, **TableS12**). NN ABC posterior parameter estimation procedures
were conducted using 100,000 simulations under Scenario i1-1b, each simulation corresponding to a single vector of parameter
values drawn randomly from prior distributions provided in **TableS8**. We considered 43 neurons in the hidden layer of the
NN and a tolerance level of 0.01, corresponding to the 1,000 simulations providing summary-statistics closest to the observed
ones, for each 25 separate analyses. NN posterior estimates are based on the logit transformation of parameter values using an
Epanechnikov kernel between the corresponding parameter's prior bounds (see **Material and Methods** and **TableS8**).
Posterior parameter densities are represented with solid blue lines. 50% Credibility Intervals are represented as the light blue
area under the density. Parameter prior distributions are represented as dotted grey lines. For all panels, the plots represent the
NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern African populations under Scenario
i1-1b, separately (**TableS12** and **TableS13**). See **Figure5** and **TableS8** for all parameters' descriptions in each panel. Note
that, overall, parameters are relatively little departing from their priors for numerous sets of population combinations among
the 25. Also, posterior parameter distributions that are substantially departing from their priors are often highly differing from
one another for each parameter. Therefore, we considered these parameters as unsatisfactorily estimated in our analyses and
discuss this limitation in the **Discussion** section of the main text.
2526

FigureS22: PCA posterior-check for the schematic NN-ABC inferred demographic and migration history of Central
and Southern African populations proposed in Figure9 and TableS15.
First two axes of a PCA computed on 202 summary-statistics used to conduct Neural Network ABC posterior-parameter
estimations under the winning Scenario i1-1b. Observed summary-statistics for the 25 combinations of five Central and
Southern African populations each (see **Material and Methods**), are indicated with red dots. Pseudo-observed summary-
statistics for 25 separate simulated datasets of five populations each conducted using mode point-estimate values for each
model-parameter provided in **TableS15** and schematized in **Figure9** (see **Material and Methods**), are indicated with blue
crosses.
Pseudo-observed simulated datasets were obtained under a single demographic model synthesizing the 25 models obtained
with NN-ABC inferences conducted on 25 different combinations of observed data separately. Nevertheless, pseudo-observed
simulations provide summary-statistics reasonably close to that calculated on the 25 observed datasets. Note that, as expected,
most variation in this PCA 2D plot is explained by differences among the 25 observed combinations of five populations each.
2539

2540
2541

2542 **FigureS23: NN-ABC parameter inference posterior check: posterior distribution of divergence time parameters.**
2543 Figure analog to **Figure7** obtained using 25 pseudo-observed simulations under the synthetic scenario proposed in **Figure9**
2544 and **TableS15** instead of the 25 observed datasets (see **Material and Methods**). Vertical red-doted lines indicate, for each
2545 scenario-parameter, the mode point-estimate values from **TableS15** used to perform the 25 simulations under Scenario i1-1b.
2546
2547
2548

2549　**FigureS24: NN-ABC parameter inference posterior check: posterior distribution of potentially asymmetric**
2550　**instantaneous gene-flow time parameters.**
2551　Figure analog to **Figure8** obtained using 25 pseudo-observed simulations under the synthetic scenario proposed in **Figure9**
2552　and **TableS15** instead of the 25 observed datasets (see **Material and Methods**). Vertical red-doted lines indicate, for each
2553　scenario-parameter, the mode point-estimate values from **TableS15** used to perform the 25 simulations under Scenario i1-1b.
2554
2555
2556

FigureS25: **NN-ABC parameter inference posterior check: posterior distribution of effective population size parameters.**
Figure analog to **FigureS20** obtained using 25 pseudo-observed simulations under the synthetic scenario proposed in **Figure9** and **TableS15** instead of the 25 observed datasets (see **Material and Methods**). Vertical red-doted lines indicate, for each scenario-parameter, the mode point-estimate values from TableS15 used to perform the 25 simulations under Scenario i1-1b.

FigureS26: **NN-ABC parameter inference posterior check: posterior distribution of potentially asymmetric instantaneous gene-flow intensity parameters.**
Figure analog to **FigureS21** obtained using 25 pseudo-observed simulations under the synthetic scenario proposed in **Figure9** and **TableS15** instead of the 25 observed datasets (see **Material and Methods**). Vertical red-doted lines indicate, for each scenario-parameter, the mode point-estimate values from **TableS15** used to perform the 25 simulations under Scenario i1-1b.

2573 **FigureS27: Schematic overview of the sequencing data processing pipeline.**
2574 All template codes with accompanying detailed explanations for all steps are provided in **Material and Methods** and in the
2575 corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

2576

TableS1: Complementary population information for genetic-geographic comparisons, and for linguistic and life-style
categories for ANOVA analyses.
Geographical coordinates correspond to locations in **Table1** and **Figure1**. "Traditional food-producing strategies" were
obtained via ethno-anthropological investigations on the field for all original population-data, and from previously published
material for comparison reference-datasets from the SGDP, HGDP, and KGP (see **Table1** and **Main Text** for references).
"*nd*" stands for no explicit data concerning the genetic dataset specifically. "AGR" stands for agriculturalist and/or
agropastoralist populations. "HG" stands for hunting, gathering, fishing, and horticulturalist populations. "Herder" stands for
more strictly pastoral populations.

2585

Population Name	N	Dataset	GeoCoord:X	GeoCoord:Y	Linguistic Family	Sub-Linguistic Family	Traditional Food-Producing Strategy
Baka	7	this study	13.86667	3.11667	Niger-Congo	Niger-Congo,Adamawa-Ubangian	HG
Nzime	5	this study	14.10000	2.98333	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Ba.Kola	5	this study	10.48010	3.12090	Niger-Congo	Niger-Congo, Narrow-Bantu	HG
Ngumba	5	this study	10.72630	3.23490	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Bi.Aka_Mbati	5	this study	17.83462	3.76069	Niger-Congo	Niger-Congo, Narrow-Bantu	HG
Ba.Kiga	5	this study	29.61434	-0.98659	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Ba.Twa	6	this study	29.61434	-0.98659	Niger-Congo	Niger-Congo, Narrow-Bantu	HG
Nsua	5	this study	30.25151	0.95051	Nilo-Saharan	Nilo-Saharan, Eastern-Sudanic	HG
Ba.Konjo	5	this study	30.25151	0.91938	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Karretjie People	5	this study	25.10101	-30.71264	Khoisan	Khoisan	Herder
Khutse-San	5	this study	24.66980	-23.65090	Khoisan	Khoisan	HG
Ju|'hoansi	5	this study	20.49500	-19.59740	Khoisan	Khoisan	HG
Nama	5	this study	17.07275	-22.55856	Khoisan	Khoisan	Herder
!Xun	5	this study	17.66602	-14.62894	Khoisan	Khoisan	Herder
Bantu_Herero	2	SGDP	19.00000	-22.00000	Niger-Congo	Niger-Congo, Narrow-Bantu	Herder
Bantu_Kenya	2	SGDP	37.00000	-3.00000	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Bantu_Tswana	2	SGDP	24.00000	-28.00000	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Biaka	2	SGDP	17.00000	4.00000	Niger-Congo	Niger-Congo, Narrow-Bantu	HG
Coloured	8	SAHGP	18.41740	-33.92899	Indo-European	Indo-European, Afrikaans	*nd*
Dinka	4	SGDP, HGDP	27.40000	8.78333	Nilo-Saharan	Nilo-Saharan, Eastern-Sudanic	AGR
Esan	3	SGDP, KGP	6.00000	6.50000	Niger-Congo	Niger-Congo, Volta-Niger	AGR
Gambian	2	SGDP	-16.70000	13.40000	Niger-Congo	Niger-Congo, Mande	AGR
Igbo	2	SGDP	7.00000	6.00000	Niger-Congo	Niger-Congo, Volta-Niger	AGR
Ju|'hoansi_comp	4	SGDP, HGDP	21.50000	-18.90000	Khoisan	Khoisan	HG
#Khomani	2	SGDP	20.80000	-27.00000	Khoisan	Khoisan	Herder
Kongo	1	SGDP	11.80000	2.50000	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Lemande	2	SGDP	11.10000	4.50000	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Luhya	3	SGDP, KGP	36.80000	1.30000	Niger-Congo	Niger-Congo, Narrow-Bantu	AGR
Luo	2	SGDP	34.30000	-0.10000	Nilo-Saharan	Nilo-Saharan, Eastern-Sudanic	AGR
Maasai	2	SGDP	35.20000	-1.50000	Nilo-Saharan	Nilo-Saharan, Eastern-Sudanic	Herder
Mandenka	4	SGDP, HGDP	-12.00000	12.00000	Niger-Congo	Niger-Congo, Mande	AGR
Mandinka	1	KGP	-16.70000	13.40000	Niger-Congo	Niger-Congo, Mande	AGR
Mbuti	5	SGDP, HGDP	29.00000	1.00000	Niger-Congo	Niger-Congo, Narrow-Bantu	HG
Mende	3	SGDP	-13.20000	8.20000	Niger-Congo	Niger-Congo, Mande	AGR
Mozabite	2	SGDP	3.00000	32.00000	Afro-Asiatic	Afro-Asiatic, Berber	AGR
Saharawi	2	SGDP	-8.90000	27.30000	Afro-Asiatic	Afro-Asiatic, Hassaniya	*nd*
Somali	1	SGDP	48.30000	5.60000	Afro-Asiatic	Afro-Asiatic, Cushitic	*nd*
Sotho	7	SAHGP	27.13588	-28.08954	Niger-Congo	Niger-Congo, Narrow-Bantu	*nd*
Xhosa	8	SAHGP	25.62075	-33.96171	Niger-Congo	Niger-Congo, Narrow-Bantu	*nd*
Yoruba	4	SGDP, HGDP	3.90000	7.40000	Niger-Congo	Niger-Congo, Volta-Niger	AGR
Zulu	1	SAHGP	28.04972	-26.20500	Niger-Congo	Niger-Congo, Narrow-Bantu	*nd*

2586
2587

TableS2: Whole genome total variant counts among 177 unrelated worldwide individuals.
Variant counts compared to the reference sequence for the human genome GRCh38 and previously reported variants in dbSNP 156. "SD" stands for "standard deviation". See **Material and Methods** for details about the quality control, relatedness filtering, and variant count procedures.

	Total variant count		Per individual variant count among 177 worldwide unrelated individuals			
	Original dataset: 73 Central and Southern African unrelated individuals	Entire dataset: 177 worldwide unrelated individuals	Total	SD	Min	Max
Number of biallelic SNPs	26,780,319	36,272,545	4,245,000	375,379	3,093,785	4,642,544
Number of biallelic SNPs not in dbSNP 156	854,114	1,055,245	8,052	4,563	1,152	27,406
Number of multiallelic SNPs	241,428	257,906	56,330	5,162	38,057	61,716
Number of simple indels	2,454,965	3,159,306	385,391	35,522	278,237	421,976
Number of novel simple indels not in dbSNP 156	114,362	189,124	1,651	514	771	627
Number of complex indels	969,025	977,542	414,037	71,029	289,993	566,951
Number of variants appearing in a single sample	8,404,499	12,638,316	71,403	15,094	22,444	104,211
Number of filtered SNPs	2,940,629	3,296,775	247,172	48,579	66,597	350,377

<u>TableS3</u>: Whole genome biallelic SNPs counts in 46 worldwide populations.

Mean number of bi-allelic SNPs across unrelated individuals from 46 worldwide populations, and associated standard deviations (when applicable), compared to the reference sequence for the human genome GRCh38 and compared to previously reported variants in dbSNP 156. See **Material and Methods** for details about the quality control, relatedness filtering, and variant count procedures. Populations are ordered in increasing mean number of SNPs per pop. The information original to this study is indicated in bold. "SD" stands for standard deviation. "*na*" stands for not applicable. Additional population information and geographical location of samples can be found in **Figure1**, **Table1**, and **TableS1**.

Population Name[1]	Sampling location	Dataset[2]	N[3]	Mean coverage X (SD)	Mean number of biallelic SNPs (SD)	Mean number of biallelic SNPs not in dbSNP 156 (SD)
Karitiana	Brazil	SGDP, HGDP	5	31.1 (7.8)	3,214,086 (69,241)	1,030 (547)
Papuan	Papua New Guinea	SGDP, HGDP	6	35.8 (7.8)	3,414,518 (74,893)	841 (290)
French	France	SGDP, HGDP	4	33.8 (8.7)	3,455,358 (77,954)	652 (225)
Dai	China	SGDP, KGP	6	37.7 (10.5)	3,480,744 (67,644)	695 (407)
CEU	United States of America	KGP	2	67.3 (0.2)	3,551,792 (7,819)	1,539 (57)
Saharawi	Western Sahara	SGDP	2	43.7 (2.2)	3,726,018 (27,793)	590 (21)
Mozabite	Algeria	SGDP	2	34.9 (1.6)	3,736,308 (69,393)	563 (11)
Somali	Kenya	SGDP	1	37.7 (*na*)	3,940,793 (*na*)	540 (*na*)
Coloured	South Africa	SAHGP	8	44.9 (2.2)	4,000,396 (97,447)	10,169 (2,319)
Dinka	Sudan	SGDP, HGDP	4	31.3 (5.8)	4,142,138 (97,001)	3,146 (4,811)
Maasai	Kenya	SGDP	2	38.6 (5.5)	4,169,732 (5,281)	654 (27)
Mandinka	Gambia	KGP	1	27.2 (*na*)	4,193,651 (*na*)	1,602 (*na*)
Mandenka	Senegal	SGDP, HGDP	4	30 (7.6)	4,217,398 (107,847)	914 (464)
Yoruba	Nigeria	SGDP, HGDP	4	31.1 (5)	4,233,500 (11,5371)	755 (195)
Ba.Kiga	**Mukono (Uganda)**	**this study**	**5**	**39.4 (4.6)**	**4,274,466 (25,201)**	**5,391 (191)**
Gambian	Gambia	SGDP	2	35.5 (0.8)	4,277,874 (12,976)	816 (107)
Esan	Nigeria	SGDP, KGP	3	39.7 (9.5)	4,284,041 (18,359)	704 (67)
Bantu_Kenya	Kenya	SGDP	2	38.7 (6.8)	4,288,533 (10,954)	611 (88)
Luo	Kenya	SGDP	2	34 (0.2)	4,293,632 (1,003)	722 (4)
Luhya	Kenya	SGDP, KGP	3	40.6 (12.1)	4,294,198 (28,096)	704 (37)
Igbo	Nigeria	SGDP	2	36.3 (8)	4,294,870 (1,867)	1,166 (9)
Lemande	Cameroon	SGDP	2	36.3 (0.5)	4,306,880 (5,284)	1,530 (37)
Bantu_Herero	Namibia	SGDP	2	38.4 (2.2)	4,318,604 (33,525)	690 (161)
Ba.Konjo	**Mulimassenge (Uganda)**	**this study**	**5**	**43 (6.4)**	**4,322,177 (6,336)**	**5,349 (773)**
Nzime	**Messea (Cameroon)**	**this study**	**5**	**35.5 (2.6)**	**4,322,734 (18,671)**	**5,305 (208)**
Ngumba	**Dispersed between Lolodorf and Kribi (Cameroon)**	**this study**	**5**	**40.2 (2.2)**	**4,332,324 (9,273)**	**4,810 (552)**
Kongo	Cameroon	SGDP	1	42.8 (*na*)	4,336,731 (*na*)	1233 (*na*)
Mende	Sierra Leone	SGDP, KGP	3	39.6 (8.3)	4,341,512 (13,486)	866 (180)
Zulu	South Africa	SAHGP	1	42.8 (*na*)	4,376,691 (*na*)	8,146 (*na*)
Xhosa	South Africa	SAHGP	8	42.4 (1.1)	4,385,032 (56,902)	10,085 (1,300)
Ba.Twa	**Kebiremu, Byumba, Kitariro, Mgungu, Nteko (Uganda)**	**this study**	**6**	**53.8 (17.7)**	**4,426,223 (32,552)**	**20,251 (1,022)**
Bantu_Tswana	South Africa	SGDP	2	36.7 (3.1)	4,430,870 (55,537)	720 (37)
Mbuti	Democratic Republic of Congo	SGDP, HGDP	5	32.1 (7.4)	4,434,425 (110,658)	1,227 (488)
Sotho	South Africa	SAHGP	7	42.9 (6.4)	4,446,796 (23,172)	12,040 (1,150)
Ba.Kola	**Dispersed between Lolodorf and Kribi (Cameroon)**	**this study**	**5**	**39.2 (1.9)**	**4,475,922 (50,261)**	**6,995 (1,948)**
Nsua	**Bundimassoli (Uganda)**	**this study**	**5**	**37.6 (4.9)**	**4,486,217 (22,774)**	**16,443 (984)**
Baka	**Bosquet (Cameroon)**	**this study**	**7**	**40.7 (3.5)**	**4,503,977 (23,946)**	**3,927 (144)**
Bi.Aka_Mbati	**Bombeketi section of Bagandou (Central African Republic)**	**this study**	**5**	**43.4 (13.6)**	**4,514,318 (17,376)**	**4,587 (198)**
Biaka	Central African Republic	SGDP	2	39.1 (0.3)	4,521,140 (8,155)	726 (52)
Nama	**Windhoek (Namibia)**	**this study**	**5**	**48.9 (4.3)**	**4,530,168 (18,807)**	**27,902 (462)**
Ju|'hoansi_comp	Namibia	SGDP, HGDP	4	33.9 (7)	4,554,593 (103,099)	1,382 (273)
#Khomani	South Africa	SGDP	2	44.7 (2.8)	4,559,846 (79,851)	1,364 (48)
Khutse_San	**Kutse Game reserve (Botswana)**	**this study**	**5**	**46.3 (6.7)**	**4,592,852 (27,794)**	**27,989 (1,353)**
Karretjie	**Colesberg (South Africa)**	**this study**	**5**	**46.6 (5.4)**	**4,601,724 (26,951)**	**27,720 (1,968)**
!Xun	**Omega camp (Namibia) and Schmidtsdrift (Sout Africa)[4]**	**this study**	**5**	**45.6 (1.9)**	**4,614,835 (10,047)**	**28,923 (1,371)**
Ju|'hoansi	**Tsumkwe (Namibia)**	**this study**	**5**	**49 (4.6)**	**4,628,957 (8,139)**	**26,665 (6,491))**

2605 **TableS4**: Correlations between genetic pairwise differentiation and geography.
2606 Standardized average Allele Sharing Dissimilarities across pairs of individuals from different populations are compared, for
2607 all population pairs, with the logarithm of geographic distances using Spearman and Pearson correlations, and tested using
2608 1000 Mantel permutations. See **Figure1**, **Table1, and TableS1** for population geographic location and information.
2609

Geographic Region	Population Samples	Nb pops.	Nb indivs.	Spearman ρ	Mantel Permutation p	Pearson r	Mantel Permutation p
Africa	All African population samples in **Figure1, Table1, and TableS1**	41	154	0.03573	0.32268	0.05562	0.16384
Sub-Sahara Africa	All African population samples in **Figure1, Table1, and TableS1,** except Northern African Sarahwi and Mozabites	38	150	0.07321	0.16883	0.09707	0.05294
West Western Africa	Gambian, Mandenka, Mandinka, Mende	4	10	0.50022	0.09291	0.38444	0.33467
Wider Central Africa	Igbo, Esan, Yoruba, Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG), Dinka	17	71	0.15587	0.07393	0.06298	0.21378
Central Africa - Congo Basin	Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG)	13	58	0.19952	0.03996	0.13451	0.05594
Eastern Africa	Bantu_Kenya, Luhya, Luo, Maasai, Somali	5	10	0.13939	0.25574	-0.04705	0.36863
Southern Africa	!Xun (nKS), Ju|'hoansi (nKS), Ju|'hoansi_comp (nKS), Bantu_Herero, Nama (sKS), Khutse San (Central KS), #Khomani, Zulu, Sotho, Bantu_Tswana, Karretjie_People (sKS), Coloured, Xhosa	13	59	0.07123	0.20979	0.10440	0.15485

2610
2611

2612 **TableS5: Permutation ANOVA of average population pairwise ASD explained by population linguistic families at**
2613 **different geographical scale in Africa.**
2614 See **Figure1**, **Table1** and **TableS1** for population geographical location and linguistic family categorization criteria.
2615 Significant ANOVA results are indicated in bold and non-significant results in italics. "*na*" indicates that the ANOVA test is
2616 not applicable due to insufficient numbers of levels in the linguistic family category.
2617

| | | | | ANOVA partial r^2: % Var(Av. ASD among pairs of populations) explained | | |
Geographic Region	Population Samples	Nb pops	Nb Linguistic families	among linguistic families	within linguistic families	1000 ANOVA permutation p
Africa	All African population samples in **Figure1**, **Table1**, and **TableS1**	41	5	**15.064%**	**84.936%**	**<0.001**
Sub-Sahara Africa	All African population samples in **Figure1**, **Table1**, and **TableS1**, except Northern African Sarahwi and Mozabites	39	5	**14.811%**	**85.189%**	**<0.001**
West Western Africa	Gambian, Mandenka, Mandinka, Mende	4	1	*Na*	*Na*	*na*
Wider Central Africa	Igbo, Esan, Yoruba, Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG), Dinka	17	2	*6.389%*	*93.611%*	*0.1698*
Central Africa - Congo Basin	Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG)	13	2	*9.172%*	*90.828%*	*0.1489*
Eastern Africa	Bantu_Kenya, Luhya, Luo, Maasai, Somali	5	3	*50.311%*	*49.689%*	*0.297*
Southern Africa	!Xun (nKS), Ju|'hoansi (nKS), Ju|'hoansi_comp (nKS), Bantu_Herero, Nama (sKS), Khutse San (Central KS), #Khomani, Zulu, Sotho, Bantu_Tswana, Karretjie_People (sKS), Coloured, Xhosa	13	3	**21.544%**	**78.456%**	**<0.001**

2618
2619

TableS6: Permutation ANOVA of average population pairwise ASD explained by population sub-linguistic families at different geographical scale in Africa.
See **Figure1**, **Table1** and **TableS1** for population geographical location and sub-linguistic family categorization criteria. Significant ANOVA results are indicated in bold and non-significant results in italics. "*na*" indicates that the ANOVA test is not applicable due to insufficient numbers of levels in the linguistic family category.

Geographic Region	Population Samples	Nb pops	Nb Sub-linguistic families	ANOVA partial r^2: % Var(Av. ASD among pairs of populations) explained		1000 ANOVA permutation p
				among Sub-linguistic families	within Sub-linguistic families	
Africa	All African population samples in **Figure1**, **Table1**, and **TableS1**	41	10	**26.109%**	**73.891%**	**<0.001**
Sub-Sahara Africa	All African population samples in **Figure1**, **Table1**, and **TableS1**, except Northern African Sarahwi and Mozabites	39	8	**22.759%**	**77.241%**	**<0.001**
West Western Africa	Gambian, Mandenka, Mandinka, Mende	4	1	*na*	*na*	*na*
Wider Central Africa	Igbo, Esan, Yoruba, Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG), Dinka	17	4	*19.441%*	*80.559%*	*0.1728*
Central Africa - Congo Basin	Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG)	13	3	*17.832%*	*82.168%*	*0.1538*
Eastern Africa	Bantu_Kenya, Luhya, Luo, Maasai, Somali	5	3	*50.311%*	*49.689%*	*0.297*
Southern Africa	!Xun (nKS), Ju|'hoansi (nKS), Ju|'hoansi_comp (nKS), Bantu_Herero, Nama (sKS), Khutse San (Central KS), #Khomani, Zulu, Sotho, Bantu_Tswana, Karretjie_People (sKS), Coloured, Xhosa	13	3	**21.544%**	**78.456%**	**<0.001**

2628 **TableS7: Permutation ANOVA of average population pairwise ASD explained by population traditional food-**
2629 **production strategies at different geographical scale in Africa.**
2630 See **Figure1**, **Table1** and **TableS1** for population geographical location and traditional food-production strategy categorization
2631 criteria. Significant ANOVA results are indicated in bold and non-significant results in italics. "*na*" indicates that the ANOVA
2632 test is not applicable due to insufficient numbers of levels in the linguistic family category.
2633

Geographic Region	Population Samples	Nb pops	Nb Traditional food-production strategies	ANOVA partial r^2: % Var(Av. ASD among pairs of populations) explained		
				among strategies	within strategies	1000 ANOVA permutation p
Africa	All African population samples in Figure1 and Table1, except Saharawi, Coloured, Somali, Sotho, Xhosa, and Zulu populations (TableS1)	35	3	**9.79%**	**90.21%**	**<0.001**
Sub-Sahara Africa	All African population samples in Figure1 and Table1, except Northern African Saharawi and Mozabite, and Southern African Coloured, Somali, Sotho, Xhosa, and Zulu populations (TableS1)	34	3	**10.01%**	**89.99%**	**<0.001**
West Western Africa	Gambian, Mandenka, Mandinka, Mende	4	1	*na*	*Na*	*na*
Wider Central Africa	Igbo, Esan, Yoruba, Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG), Dinka	17	2	**9.538%**	**90.462%**	**<0.001**
Central Africa - Congo Basin	Ba.Kola (wRHG), Ngumba (wRHGn), Baka (wRHG), Nzime(wRHGn), Kongo, Lemande, Bi.Aka_Mbati (wRHG), Bi.Aka (wRHG), Ba.Twa (eRHG), Ba.Kiga (eRHGn), Nsua (eRHG), Ba.Konjo (eRHGn), Mbuti (eRHG)	13	2	**11.153%**	**88.847%**	**0.003**
Eastern Africa	Bantu_Kenya, Luhya, Luo, Maasai	4	2	*34.198%*	*65.802%*	*0.25*
Southern Africa	!Xun (nKS), Ju|'hoansi (nKS), Ju|'hoansi_comp (nKS), Bantu_Herero, Nama (sKS), Khutse San (Central KS), #Khomani, Bantu_Tswana, Karretjie_People (sKS)	9	3	*25.979%*	*74.201%*	*0.1728*

2634
2635

2636 **TableS8: 48 competing scenarios parameters' description and prior distributions.**
2637 Codes for scenarios topologies and gene-flow processes and all scenario parameters are detailed in **Figure5** and in **Material**
2638 **and Methods**.
2639

TableS8: 48 competing scenarios parameters' description and prior distributions.

Parameter	Description	Prior	Scenario topology	Scenario gene-flow	Condition
t_{RHG}	Time of divergence between wRHG and eRHG	Uniform[1,15000]	1a-2a-2c-3a 1b-1c-2b-3b	i1-i2-i3 and r1-r2-r3	$t_{RHG}>t_{KS}$ $t_{KS}>t_{RHG}$
t_{KS}	Time of divergence between nKS and sKS	Uniform[1,15000]	1a-2a-2c-3a 1b-1c-2b-3b	i1-i2-i3 and r1-r2-r3	$t_{RHG}>t_{KS}$ $t_{KS}>t_{RHG}$
$t_{RHG-RHGn}$	Time of divergence between the RHGn lineage and the lineage ancestral to all RHG lineages	Uniform[1,15000]	1a-1b-1c	i1-i2-i3 and r1-r2-r3	$t_{RHG-RHGn}>t_{RHG}$
$t_{KS-RHGn}$	Time of divergence between the RHGn lineage and the lineage ancestral to all KS lineages	Uniform[1,15000]	2a-2b-2c	i1-i2-i3 and r1-r2-r3	$t_{KS-RHGn}>t_{KS}$
t_{KS-RHG}	Time of divergence between the KS and RHG ancestral lineages	Uniform[1,15000]	3a-3b	i1-i2-i3 and r1-r2-r3	$t_{KS-RHG}>t_{KS}$ and $t_{KS-RHG}>t_{RHG}$
$t_{KS-RHG-RHGn}$	Time of the divergence event ancestral to all lineages	Uniform[1,15000]	all 8 topol.	i1-i2-i3 and r1-r2-r3	$t_{KS-RHG-RHGn}>$ all other times
$tad_{nKS-sKS}$	Time for both the unidirectional instantaneous gene-flow events between nKS and sKS	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{nKS-sKS}$
m(nKS-sKS) or m(sKS-nKS)	Instantaneous gene-flow intensity from sKs into nKS, or from nKS into sKS independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(nKS-sKS) or m(sKS-nKS)	Recurring gene-flow intensity from sKS into nKS, or from nKS into sKs independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{nKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between nKS and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{nKS-RHGn}$ and $t_{RHG-RHGn}>tad_{nKS-RHGn}$
m(nKS-RHGn) or m(RHGn-nKS)	Instantaneous gene-flow intensity from RHGn into nKS, or from nKS into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(nKS-RHGn) or m(RHGn-nKS)	Recurring gene-flow intensity from RHGn into nKS, or from nKS into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{sKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between sKS and RHGn lineages	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{sKS-RHGn}$ and $t_{RHG-RHGn}>tad_{sKS-RHGn}$
m(sKS-RHGn) or m(RHGn-sKS)	Instantaneous gene-flow intensity from RHGn into sKS, or from sKS into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(sKS-RHGn) or m(RHGn-sKS)	Recurring gene-flow intensity from RHGn into sKS, or from sKS into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{wRHG-eRHG}$	Time for both the unidirectional instantaneous gene-flow events between wRHG and eRHG	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{wRHG-eRHG}$
m(wRHG-eRHG) or m(eRHG-wRHG)	Instantaneous gene-flow intensity from eRHG into wRHG, or from wRHG into eRHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(wRHG-eRHG) or m(eRHG-wRHG)	Recurring gene-flow intensity from eRHG into wRHG, or from wRHG into eRHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{wRHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between wRHG and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{wRHG-RHGn}$ and $t_{KS-RHGn}>tad_{nwRHG-RHGn}$
m(wRHG-RHGn) or m(RHGn-wRHG)	Instantaneous gene-flow intensity from RHGn into wRHG, or from wRHG into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(wRHG-RHGn) or m(RHGn-wRHG)	Recurring gene-flow intensity from RHGn into wRHG, or from wRHG into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{eRHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between eRHG and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{eRHG-RHGn}$ and $t_{KS-RHGn}>tad_{eRHG-RHGn}$
m(eRHG-RHGn) or m(RHGn-eRHG)	Instantaneous gene-flow intensity from RHGn into eRHG, or from eRHG into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(eRHG-RHGn) or m(RHGn-eRHG)	Recurring gene-flow intensity from RHGn into eRHG, or from eRHG into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{ARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the RHGn and the lineage ancestral to wRHG and eRHG lineages	Uniform[1,15000]	all 8 topol. except 2c	i1-i2-i3	$tad_{RHGn-ARHG}>t_{RHG}$
m(RHGn-ARHG) or m(ARHG-RHGn)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into RHGn, or from RHGn into ARHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol. except 2c	i1 i2 i3	x
m(RHGn-ARHG) or m(ARHG-RHGn)	Recurring gene-flow intensity from the ancestral lineage ARHG into RHGn, or from RHGn into ARHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol. except 2c	r1 r2 r3	x
$tad_{AKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to nKS and sKS, and the RHGn lineage	Uniform[1,15000]	all 8 topol. except 1c	i1-i2-i3	$tad_{AKS-RHGn}>t_{KS}$
m(RHGn-AKS) or m(AKS-RHGn)	Instantaneous gene-flow intensity from the ancestral lineage AKS into RHGn, or from RHGn into AKS independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol. except 1c	i1 i2 i3	x
m(RHGn-AKS) or m(AKS-RHGn)	Recurring gene-flow intensity from the ancestral lineage AKS into RHGn, or from RHGn into AKS independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol. except 1c	r1 r2 r3	x
$tad_{AKS-ARHG}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to nKS and sKS, and the lineage ancestral to wRHG and eRHG lineages	Uniform[1,15000]	all 8 topol. except 1c-2c	i1-i2-i3	$tad_{AKS-ARHG}>t_{KS}$ and $tad_{AKS-ARHG}>t_{RHG}$
m(AKS-ARHG) or		Uniform[0,1]		i1	x

Parameter	Description	Prior	Topology	Index		Constraints
m(ARHG-AKS)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into the ancestral lineage AKS, or from AKS into ARHG independantly	Uniform[0,0.125]	all 8 topol. except 1c-2c	i2		
		Uniform[0,0.01]		i3		
m(AKS-ARHG) or m(ARHG-AKS)	Recurring gene-flow intensity from the ancestral lineage ARHG into the ancestral lineage AKS, or from AKS into ARHG independantly	Uniform[0,0.05]	all 8 topol. except 1c-2c	r1		
		Uniform[0,0.0125]		r2	x	
		Uniform[0,0.001]		r3		
$tad_{nKS\text{-}ARHG\text{-}RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the nKS and the lineage ancestral to ARHG and RHGn lineages	Uniform[1,15000]	1c	i1-i2-i3		$t_{KS} > tad_{nKS\text{-}ARHG\text{-}RHGn}$ and $tad_{nKS\text{-}ARHG\text{-}RHGn} > t_{RHG\text{-}RHGn}$
m(nKS-ARHG-RHGn) or (ARHG-RHGn-nKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into nKS, or from nKS into ARHG-RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	1c	i2	x	
		Uniform[0,0.01]		i3		
m(nKS-ARHG-RHGn) or (ARHG-RHGn-nKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into nKS, or from nKS into ARHG-RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	1c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{sKS\text{-}ARHG\text{-}RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the sKS and the lineage ancestral to ARHG and RHGn lineages	Uniform[1,15000]	1c	i1-i2-i3		$t_{KS} > tad_{sKS\text{-}ARHG\text{-}RHGn}$ and $tad_{sKS\text{-}ARHG\text{-}RHGn} > t_{RHG\text{-}RHGn}$
m(sKS-ARHG-RHGn) or m(ARHG-RHGn-sKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into sKS, or from sKS into ARHG-RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	1c	i2	x	
		Uniform[0,0.01]		i3		
m(sKS-ARHG-RHGn) or m(ARHG-RHGn-sKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into sKS, or from sKS into ARHG-RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	1c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{AKS\text{-}RHGn\text{-}wRHG}$	Time for both the unidirectional instantaneous gene-flow events between the wRHG and the lineage ancestral to AKS and RHGn lineages	Uniform[1,15000]	2c	i1-i2-i3		$t_{RHG} > tad_{AKS\text{-}RHGn\text{-}wRHG}$ and $tad_{AKS\text{-}RHGn\text{-}wRHG} > t_{KS\text{-}RHGn}$
m(wRHG-AKS-RHGn) or m(AKS-RHGn--wRHG)	Instantaneous gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into wRHG, or from wRHG into AKS-RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	2c	i2	x	
		Uniform[0,0.01]		i3		
m(wRHG-AKS-RHGn) or m(AKS-RHGn--wRHG)	Recurring gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into wRHG, or from wRHG into AKS-RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	2c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{AKS\text{-}RHGn\text{-}eRHG}$	Time for both the unidirectional instantaneous gene-flow events between the eRHG and the lineage ancestral to AKS and RHGn lineages	Uniform[1,15000]	2c	i1-i2-i3		$t_{RHG} > tad_{AKS\text{-}RHGn\text{-}eRHG}$ and $tad_{AKS\text{-}RHGn\text{-}eRHG} > t_{KS\text{-}RHGn}$
m(eRHG-AKS-RHGn) or m(AKS-RHGn--eRHG)	Instantaneous gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into eRHG, or from eRHG into AKS-RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	2c	i2	x	
		Uniform[0,0.01]		i3		
m(eRHG-AKS-RHGn) or m(AKS-RHGn--eRHG)	Recurring gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into eRHG, or from eRHG into AKS-RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	2c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{AKS\text{-}ARHG\text{-}RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and the lineage ancestral to RHG and RHGn lineages	Uniform[1,15000]	1a-1b-1c	i1-i2-i3		$tad_{AKS\text{-}ARHG\text{-}RHGn} > t_{RHG\text{-}RHGn}$ and $tad_{AKS\text{-}ARHG\text{-}RHGn} > t_{KS}$ and $t_{KS\text{-}RHG\text{-}RHGn} > tad_{AKS\text{-}ARHG\text{-}RHGn}$
m(AKS-ARHG-RHGn) or (ARHG-RHGn-AKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages ARHG and RHGn into the ancestral lineage AKS, or from AKS into ARHG-RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	1a-1b-1c	i2	x	
		Uniform[0,0.01]		i3		
m(AKS-ARHG-RHGn) or (ARHG-RHGn-AKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages ARHG and RHGn into the ancestral lineage AKS, or from AKS into ARHG-RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	1a-1b-1c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{AKSRHGn\text{-}ARHG}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and RHGn, and the lineage ancestral to RHG lineages	Uniform[1,15000]	2a-2b-2c	i1-i2-i3		$tad_{AKSRHGn\text{-}ARHG} > t_{KS\text{-}RHGn}$ and $tad_{AKSRHGn\text{-}ARHG} > t_{RHG}$ and $t_{KS\text{-}RHG\text{-}RHGn} > tad_{AKSRHGn\text{-}ARHG}$
m(AKSRHGn-ARHG) or m(ARHG-AKSRHGn)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into the lineage AKSRHGn ancestral to AKS and RHGn, or from AKSRHGn into ARHG independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	2a-2b-2c	i2	x	
		Uniform[0,0.01]		i3		
m(AKSRHGn-ARHG) or m(ARHG-AKSRHGn)	Recurring gene-flow intensity from the ancestral lineage ARHG into the lineage AKSRHGn ancestral to AKS and RHGn, or from AKSRHGn into ARHG independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	2a-2b-2c	r2	x	
		Uniform[0,0.001]		r3		
$tad_{AKSARHG\text{-}RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and RHG, and the RHGn lineage	Uniform[1,15000]	3a-3b	i1-i2-i3		$tad_{AKSARHG\text{-}RHGn} > t_{KS\text{-}RHG}$ and $t_{KS\text{-}RHG\text{-}RHGn} > tad_{AKSARHG\text{-}RHGn}$
m(AKSARHG-RHGn) or m(RHGn-AKSARHG)	Instantaneous gene-flow intensity from the lineage RHGn into the lineage AKSRHG ancestral to AKS and ARHG, or from AKSRHG into RHGn independantly	Uniform[0,1]		i1		
		Uniform[0,0.125]	3a-3b	i2	x	
		Uniform[0,0.01]		i3		
m(AKSARHG-RHGn) or m(RHGn-AKSARHG)	Recurring gene-flow intensity from the lineage RHGn into the lineage AKSRHG ancestral to AKS and ARHG, or from AKSRHG into RHGn independantly	Uniform[0,0.05]		r1		
		Uniform[0,0.0125]	3a-3b	r2	x	
		Uniform[0,0.001]		r3		
N_{nKS}	Effective population size for the nKS lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{sKS}	Effective population size for the sKS lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{AKS}	Effective population size for the lineage ancestral to the nKS and the sKS		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{RHGn}	Effective population size for the RHGn lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{wRHG}	Effective population size for the wRHG lineage	Uniform[10,100000]	all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{eRHG}	Effective population size for the eRHG lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
N_{ARHG}	Effective population size for the lineage ancestral to the wRHG and the eRHG		all 8 topol.	i1-i2-i3 and r1-r2-r3	x	
$N_{ARHG\text{-}RHGn}$	Effective population size for the lineage ancestral to the RHGn and the ARHG		1a-1b-1c	i1-i2-i3 and r1-r2-r3	x	

$N_{AKS-RHGn}$	Effective population size for the lineage ancestral to the RHGn and the AKS		2a-2b-2c	i1-i2-i3 and r1-r2-r3 x
$N_{AKS-ARHG}$	Effective population size for the lineage ancestral to the AKS and the ARHG		3a-3b	i1-i2-i3 and r1-r2-r3 x
N_A	Effective population size for the lineage ancestral to all lineages		all 8 topol.	i1-i2-i3 and r1-r2-r3 x
Mutation rate	Fixed mutation rate	$1.25*10^{-8}$	all 8 topol.	i1-i2-i3 and r1-r2-r3 x
Recombination rate	Fixed recombination rate	$1*10^{-8}$	all 8 topol.	i1-i2-i3 and r1-r2-r3 x
Transition to transversion ratio	Fixed transition to transversion rate	0.33	all 8 topol.	i1-i2-i3 and r1-r2-r3 x

2640
2641

TableS9: Summary-statistics used in Random-Forest ABC scenario-choice and Neural-Network ABC posterior-parameter estimation.

Name of statistic (or group of statistics)	What was computed	Number of values	RF-ABC scenario-choice	NN-ABC posterior-parameter estimation
Total number of biallelic sites among the five populations	sum	1	yes	yes
Total number of multiallelic sites among the five populations	sum	1	yes	yes
ASD distance within population for each five population separately	average and variance	10	yes	no
First five dimensions of the projection of the ASD matrix within population for each five population separately	average and variance	50	yes	yes
Proportion of biallelic sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Proportion of homozygous sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Proportion of homozygous ancestral sites within each five population separately	sum in a population divided by total number of homozygous sites in the population (no missing data)	5	yes	no
Proportion of private biallelic sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Frequency of the minor allele of the private biallelic sites within each five population separately	average and variance	10	yes	no
Expected heterozygosity at a segregating site within each five population separately	average and variance	10	yes	no
Tajima's D within each five population separately	average and variance for chromosomes with at least one variant	10	yes	yes
1D Unfolded standardized site frequency spectrum (SFS) within each five population separately	proportion for each of 9 classes	45	yes	no
Watterson's theta within each five population separately	sum of biallelic sites divided by the harmonic number of the sample size (n=10)	5	yes	no
Total number of runs of homozygosity (ROH) within each five population separately	sum	5	yes	no
Total number of ROH by length class (<200 kbp, 200-500 kbp, >500 kbp) within each five population separately	sum	15	yes	yes
ROH length for each length class within each five population separately	average	15	yes	no
Total ROH length within each five population separately	average and variance	10	yes	no
ASD distance between each pair of five populations	average and variance	20	yes	no
First five dimensions of the projection of the ASD matrix between each pair of five populations	average and variance	100	yes	yes
Pairwise FST between each pair of five population	Pairwise FST (Weir and Cockerham 1984)	10	yes	yes
	TOTAL number of statistics used in ABC inference		337	202

TableS10: Correlations between standardized ASD matrices for each one of the 54 combinations of five population samples computed using 100 separate high-quality 1 Mb-long genomic windows, and corresponding individual-subsets of the standardized ASD matrix computed using the full genomic SNP set and used for Figure3.

Combinations of five populations used for ABC inferences					*Spearman_ρ*	1000 Mantel prem. p	*Pearson_r*	1000 Mantel prem. p
nKS	**sKS**	**RHGn**	**wRHG**	**eRHG**				
Ju'hoansi	Karretjie_People	Ba.Konjo	Ba.Kola	Mbuti	0.893	<0.002	0.962	<0.002
Ju'hoansi	Karretjie_People	Ba.Konjo	Ba.Kola	Nsua	0.896	<0.002	0.963	<0.002
Ju'hoansi	Karretjie_People	Ba.Konjo	Baka	Mbuti	0.898	<0.002	0.964	<0.002
Ju'hoansi	Karretjie_People	Ba.Konjo	Baka	Nsua	0.903	<0.002	0.967	<0.002
Ju'hoansi	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Mbuti	0.894	<0.002	0.963	<0.002
Ju'hoansi	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Nsua	0.895	<0.002	0.965	<0.002
Ju'hoansi	Karretjie_People	Ba.Kiga	Ba.Kola	Ba.Twa	0.901	<0.002	0.961	<0.002
Ju'hoansi	Karretjie_People	Ba.Kiga	Baka	Ba.Twa	0.908	<0.002	0.965	<0.002
Ju'hoansi	Karretjie_People	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	0.887	<0.002	0.961	<0.002
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Ba.Twa	0.885	<0.002	0.960	<0.002
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Mbuti	0.907	<0.002	0.963	<0.002
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Nsua	0.910	<0.002	0.964	<0.002
Ju'hoansi	Karretjie_People	Nzime	Baka	Ba.Twa	0.914	<0.002	0.968	<0.002
Ju'hoansi	Karretjie_People	Nzime	Baka	Mbuti	0.929	<0.002	0.971	<0.002
Ju'hoansi	Karretjie_People	Nzime	Baka	Nsua	0.935	<0.002	0.973	<0.002
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Ba.Twa	0.896	<0.002	0.965	<0.002
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Mbuti	0.925	<0.002	0.970	<0.002
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Nsua	0.927	<0.002	0.970	<0.002
Ju'hoansi	Nama	Ba.Konjo	Ba.Kola	Nsua	0.932	<0.002	0.966	<0.002
Ju'hoansi	Nama	Ba.Konjo	Baka	Nsua	0.948	<0.002	0.969	<0.002
Ju'hoansi	Nama	Ba.Konjo	Bi.Aka_Mbati	Nsua	0.944	<0.002	0.967	<0.002
Ju'hoansi	Nama	Ba.Kiga	Ba.Kola	Ba.Twa	0.948	<0.002	0.964	<0.002
Ju'hoansi	Nama	Ba.Kiga	Baka	Ba.Twa	0.958	<0.002	0.966	<0.002
Ju'hoansi	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	0.945	<0.002	0.962	<0.002
Ju'hoansi	Nama	Ngumba	Ba.Kola	Ba.Twa	0.935	<0.002	0.962	<0.002
Ju'hoansi	Nama	Ngumba	Ba.Kola	Nsua	0.945	<0.002	0.965	<0.002
Ju'hoansi	Nama	Nzime	Baka	Ba.Twa	0.957	<0.002	0.970	<0.002
Ju'hoansi	Nama	Nzime	Baka	Nsua	0.963	<0.002	0.974	<0.002
Ju'hoansi	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa	0.945	<0.002	0.965	<0.002
Ju'hoansi	Nama	Nzime	Bi.Aka_Mbati	Nsua	0.945	<0.002	0.971	<0.002
!Xun	Karretjie_People	Ba.Konjo	Ba.Kola	Nsua	0.927	<0.002	0.952	<0.002
!Xun	Karretjie_People	Ba.Konjo	Baka	Nsua	0.936	<0.002	0.957	<0.002
!Xun	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Nsua	0.934	<0.002	0.956	<0.002
!Xun	Karretjie_People	Ba.Kiga	Ba.Kola	Ba.Twa	0.923	<0.002	0.949	<0.002
!Xun	Karretjie_People	Ba.Kiga	Baka	Ba.Twa	0.936	<0.002	0.954	<0.002
!Xun	Karretjie_People	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	0.922	<0.002	0.950	<0.002
!Xun	Karretjie_People	Ngumba	Ba.Kola	Ba.Twa	0.913	<0.002	0.948	<0.002
!Xun	Karretjie_People	Ngumba	Ba.Kola	Nsua	0.930	<0.002	0.952	<0.002
!Xun	Karretjie_People	Nzime	Baka	Ba.Twa	0.945	<0.002	0.959	<0.002
!Xun	Karretjie_People	Nzime	Baka	Nsua	0.956	<0.002	0.963	<0.002
!Xun	Karretjie_People	Nzime	Bi.Aka_Mbati	Ba.Twa	0.931	<0.002	0.955	<0.002
!Xun	Karretjie_People	Nzime	Bi.Aka_Mbati	Nsua	0.935	<0.002	0.962	<0.002
!Xun	Nama	Ba.Konjo	Ba.Kola	Nsua	0.932	<0.002	0.954	<0.002
!Xun	Nama	Ba.Konjo	Baka	Nsua	0.945	<0.002	0.957	<0.002
!Xun	Nama	Ba.Konjo	Bi.Aka_Mbati	Nsua	0.937	<0.002	0.956	<0.002
!Xun	Nama	Ba.Kiga	Ba.Kola	Ba.Twa	0.933	<0.002	0.948	<0.002
!Xun	Nama	Ba.Kiga	Baka	Ba.Twa	0.942	<0.002	0.951	<0.002
!Xun	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	0.920	<0.002	0.947	<0.002
!Xun	Nama	Ngumba	Ba.Kola	Ba.Twa	0.910	<0.002	0.945	<0.002
!Xun	Nama	Ngumba	Ba.Kola	Nsua	0.930	<0.002	0.950	<0.002
!Xun	Nama	Nzime	Baka	Ba.Twa	0.940	<0.002	0.956	<0.002
!Xun	Nama	Nzime	Baka	Nsua	0.954	<0.002	0.962	<0.002
!Xun	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa	0.915	<0.002	0.951	<0.002
!Xun	Nama	Nzime	Bi.Aka_Mbati	Nsua	0.919	<0.002	0.959	<0.002

TableS11: Correlations between 332 summary-statistics computed on 100 genomic windows of 1Mb each used for ABC inferences and average values for 724 other such windows genome wide, for each one of the 54 combinations of five sampled-populations.

We performed separately Spearman correlation tests and Wilcoxon two-sided tests for each one of the 54 combinations of sampled populations considered here between 332 summary-statistics used for ABC inferences computed on 100 genomic windows of 1Mb each and the same summary statistics computed for 724 other such windows genome-wide. Note that we excluded 5 variance related summary-statistics from the comparisons due to scaling irrelevances across sets of windows. Values in italic were not-significant. Values in bold were significant.

Combinations of five populations used for ABC inferences					Spearman_ρ	p-value	Wilcoxon two-sided	p-value
nKS	sKS	RHGn	wRHG	eRHG				
Ju'hoansi	Karretjie_People	Ba.Konjo	Ba.Kola	Mbuti	**0.90025**	**<2.2x10-16**	*55505*	*0.8738*
Ju'hoansi	Karretjie_People	Ba.Konjo	Ba.Kola	Nsua	**0.94243**	**<2.2x10-16**	*55699*	*0.8124*
Ju'hoansi	Karretjie_People	Ba.Konjo	Baka	Mbuti	**0.88935**	**<2.2x10-16**	*54958*	*0.9505*
Ju'hoansi	Karretjie_People	Ba.Konjo	Baka	Nsua	**0.97795**	**<2.2x10-16**	*55561*	*0.8560*
Ju'hoansi	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Mbuti	**0.88791**	**<2.2x10-16**	*54975*	*0.9560*
Ju'hoansi	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Nsua	**0.98917**	**<2.2x10-16**	*55071*	*0.9869*
Ju'hoansi	Karretjie_People	Ba.Kiga	Ba.Kola	Ba.Twa	**0.99203**	**<2.2x10-16**	*55755*	*0.7949*
Ju'hoansi	Karretjie_People	Ba.Kiga	Baka	Ba.Twa	**0.88887**	**<2.2x10-16**	*55249*	*0.9560*
Ju'hoansi	Karretjie_People	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	**0.86963**	**<2.2x10-16**	*54818*	*0.9055*
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Ba.Twa	**0.93363**	**<2.2x10-16**	*56089*	*0.6928*
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Mbuti	**0.89903**	**<2.2x10-16**	*55667*	*0.8225*
Ju'hoansi	Karretjie_People	Ngumba	Ba.Kola	Nsua	**0.89407**	**<2.2x10-16**	*55683*	*0.8174*
Ju'hoansi	Karretjie_People	Nzime	Baka	Ba.Twa	**0.93279**	**<2.2x10-16**	*55110*	*0.9995*
Ju'hoansi	Karretjie_People	Nzime	Baka	Mbuti	**0.93047**	**<2.2x10-16**	*54903*	*0.9328*
Ju'hoansi	Karretjie_People	Nzime	Baka	Nsua	**0.97638**	**<2.2x10-16**	*55518*	*0.8697*
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Ba.Twa	**0.91203**	**<2.2x10-16**	*55193*	*0.9740*
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Mbuti	**0.88534**	**<2.2x10-16**	*54846*	*0.9145*
Ju'hoansi	Karretjie_People	Nzime	Bi.Aka_Mbati	Nsua	**0.92615**	**<2.2x10-16**	*55363*	*0.9193*
Ju'hoansi	Nama	Ba.Konjo	Ba.Kola	Nsua	**0.98709**	**<2.2x10-16**	*55784*	*0.7859*
Ju'hoansi	Nama	Ba.Konjo	Baka	Nsua	**0.95181**	**<2.2x10-16**	*55751*	*0.7961*
Ju'hoansi	Nama	Ba.Konjo	Bi.Aka_Mbati	Nsua	**0.99637**	**<2.2x10-16**	*55416*	*0.9023*
Ju'hoansi	Nama	Ba.Kiga	Ba.Kola	Ba.Twa	**0.94583**	**<2.2x10-16**	*55820*	*0.7747*
Ju'hoansi	Nama	Ba.Kiga	Baka	Ba.Twa	**0.97595**	**<2.2x10-16**	*55829*	*0.7719*
Ju'hoansi	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	**0.94859**	**<2.2x10-16**	*55672*	*0.8209*
Ju'hoansi	Nama	Ngumba	Ba.Kola	Ba.Twa	**0.89772**	**<2.2x10-16**	*56235*	*0.6497*
Ju'hoansi	Nama	Ngumba	Ba.Kola	Nsua	**0.97607**	**<2.2x10-16**	*56112*	*0.6859*
Ju'hoansi	Nama	Nzime	Baka	Ba.Twa	**0.89324**	**<2.2x10-16**	*55925*	*0.7423*
Ju'hoansi	Nama	Nzime	Baka	Nsua	**0.89274**	**<2.2x10-16**	*55425*	*0.8994*
Ju'hoansi	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa	**0.94542**	**<2.2x10-16**	*55174*	*0.9801*
Ju'hoansi	Nama	Nzime	Bi.Aka_Mbati	Nsua	**0.91175**	**<2.2x10-16**	*55064*	*0.9847*
!Xun	Karretjie_People	Ba.Konjo	Ba.Kola	Nsua	**0.93807**	**<2.2x10-16**	*55773*	*0.7893*
!Xun	Karretjie_People	Ba.Konjo	Baka	Nsua	**0.92892**	**<2.2x10-16**	*55492*	*0.8780*
!Xun	Karretjie_People	Ba.Konjo	Bi.Aka_Mbati	Nsua	**0.99421**	**<2.2x10-16**	*55025*	*0.9721*
!Xun	Karretjie_People	Ba.Kiga	Ba.Kola	Ba.Twa	**0.94753**	**<2.2x10-16**	*55946*	*0.7359*
!Xun	Karretjie_People	Ba.Kiga	Baka	Ba.Twa	**0.94283**	**<2.2x10-16**	*55320*	*0.9331*
!Xun	Karretjie_People	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	**0.88960**	**<2.2x10-16**	*55579*	*0.8503*
!Xun	Karretjie_People	Ngumba	Ba.Kola	Ba.Twa	**0.85660**	**<2.2x10-16**	*55508*	*0.8729*
!Xun	Karretjie_People	Ngumba	Ba.Kola	Nsua	**0.94814**	**<2.2x10-16**	*55514*	*0.8709*
!Xun	Karretjie_People	Nzime	Baka	Ba.Twa	**0.97947**	**<2.2x10-16**	*55807*	*0.7787*
!Xun	Karretjie_People	Nzime	Baka	Nsua	**0.89921**	**<2.2x10-16**	*55432*	*0.8971*
!Xun	Karretjie_People	Nzime	Bi.Aka_Mbati	Ba.Twa	**0.94073**	**<2.2x10-16**	*55083*	*0.9908*
!Xun	Karretjie_People	Nzime	Bi.Aka_Mbati	Nsua	**0.90798**	**<2.2x10-16**	*54884*	*0.9267*
!Xun	Nama	Ba.Konjo	Ba.Kola	Nsua	**0.92443**	**<2.2x10-16**	*55854*	*0.7642*
!Xun	Nama	Ba.Konjo	Baka	Nsua	**0.88787**	**<2.2x10-16**	*56213*	*0.6561*
!Xun	Nama	Ba.Konjo	Bi.Aka_Mbati	Nsua	**0.96611**	**<2.2x10-16**	*55590*	*0.8468*
!Xun	Nama	Ba.Kiga	Ba.Kola	Ba.Twa	**0.94429**	**<2.2x10-16**	*55924*	*0.7427*
!Xun	Nama	Ba.Kiga	Baka	Ba.Twa	**0.97282**	**<2.2x10-16**	*55637*	*0.8319*
!Xun	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa	**0.94463**	**<2.2x10-16**	*55365*	*0.9186*
!Xun	Nama	Ngumba	Ba.Kola	Ba.Twa	**0.85774**	**<2.2x10-16**	*55709*	*0.8093*
!Xun	Nama	Ngumba	Ba.Kola	Nsua	**0.93875**	**<2.2x10-16**	*56097*	*0.6904*
!Xun	Nama	Nzime	Baka	Ba.Twa	**0.94204**	**<2.2x10-16**	*55685*	*0.8168*
!Xun	Nama	Nzime	Baka	Nsua	**0.95260**	**<2.2x10-16**	*55684*	*0.8171*
!Xun	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa	**0.88873**	**<2.2x10-16**	*55332*	*0.9292*
!Xun	Nama	Nzime	Bi.Aka_Mbati	Nsua	**0.89805**	**<2.2x10-16**	*55200*	*0.9718*

2664 **TableS12: 25 combinations of five population samples out of 54 combinations tested for which Scenario i1-1b is best**
2665 **explaining observed data with Random-Forest ABC among the 48 competing scenarios.**
2666 Values in the table correspond to posterior densities plotted in the left panels of **Figure7**, **Figure8**, **FigureS20**, and **FigureS21**.
2667 Parameter definitions and priors are provided in **TableS8** and represented graphically in the corresponding scenario panel of
2668 **Figure5**.

2669

	nKS	sKS	RHGn	wRHG	eRHG
Combination 1	Ju|'hoansi	Karretjie People	Nzime	Baka	Nsua
Combination 2	Ju|'hoansi	Karretjie People	Ngumba	Ba.Kola	Nsua
Combination 3	Ju|'hoansi	Karretjie People	Ngumba	Ba.Kola	Ba.Twa
Combination 4	Ju|'hoansi	Karretjie People	Nzime	Bi.Aka_Mbati	Nsua
Combination 5	Ju|'hoansi	Karretjie People	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 6	Ju|'hoansi	Karretjie People	Ngumba	Ba.Kola	Mbuti
Combination 7	Ju|'hoansi	Karretjie People	Nzime	Bi.Aka_Mbati	Mbuti
Combination 8	Ju|'hoansi	Karretjie People	Ba.Konjo	Baka	Mbuti
Combination 9	Ju|'hoansi	Karretjie People	Ba.Konjo	Ba.Kola	Mbuti
Combination 10	Ju|'hoansi	Karretjie People	Ba.Konjo	Bi.Aka_Mbati	Mbuti
Combination 11	Ju|'hoansi	Nama	Ba.Konjo	Baka	Nsua
Combination 12	!Xun	Nama	Ba.Konjo	Baka	Nsua
Combination 13	Ju|'hoansi	Nama	Nzime	Baka	Nsua
Combination 14	Ju|'hoansi	Nama	Ngumba	Ba.Kola	Nsua
Combination 15	Ju|'hoansi	Nama	Nzime	Baka	Ba.Twa
Combination 16	Ju|'hoansi	Nama	Ngumba	Ba.Kola	Ba.Twa
Combination 17	Ju|'hoansi	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 18	Ju|'hoansi	Nama	Ba.Konjo	Ba.Kola	Nsua
Combination 19	Ju|'hoansi	Nama	Ba.Kiga	Ba.Kola	Ba.Twa
Combination 20	Ju|'hoansi	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa
Combination 21	!Xun	Nama	Nzime	Bi.Aka_Mbati	Nsua
Combination 22	!Xun	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 23	!Xun	Karretjie People	Nzime	Bi.Aka_Mbati	Nsua
Combination 24	!Xun	Karretjie People	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 25	!Xun	Karretjie People	Ba.Konjo	Bi.Aka_Mbati	Nsua

2670
2671

TableS13: NN-ABC posterior parameter estimation of all parameters in Scenario i1-1b for results from each 25 sets of
five populations each, separately.
Table summarizing posterior parameter distributions obtained with Neural Network ABC for each 25 combinations of
sampled-populations separately. Full posterior distributions are provided in plot formats in main **Figure7** (divergence times
t), **Figure8** (admixture times tad), supplementary **FigureS20** (Ne), and supplementary **FigureS21** (asymmetric admixture
rates). Column-headers indicate the categorization of sampled populations into Northern Khoe-San (nKS), Southern Khoe-
San (sKS), Rainforest Hunter-Gatherer neighbors (RHGn), Western Rainforest Hunter-Gatherer (wRHG), or Eastern
Rainforest Hunter-Gatherer (eRHG) groups, as per scenario topology and simulation design explicated in **Material and**
Methods and in **Figure5**. Note that each sampled population considered in the RF-ABC scenario choice analyses are
represented twice or more in the 25 combinations. "CI" stands for ABC Credibility Intervals. See main **Figure5**, supplementary
TableS8 and **Material and Methods** for detailed scenario-parameter descriptions.

2683

2684 See "**TableS13_new.xlsx**" file as the table is too large to reasonably fit in a A4 page format.

2685

2686

TableS14: **NN-ABC posterior parameter estimation cross-validation errors of all parameters in Scenario i1-1b for**
results from 25 sets of five populations each.
Cross-validation errors are based on 50 (5%) randomly chosen simulation out of the 1000 simulations closest to the observed
data for each 25 combinations of sampled populations, respectively, each considered in-turn as pseudo-observed data and for
which NN-ABC joint posterior parameter estimations were performed with the same hyper-parameters. For each parameter
and each 25 combinations of sampled populations, the first column provides variance-scaled mean-squared errors and the
second column provides simple average estimation bias (see **Material and Methods**). Relatively high variance-scaled mean-
squared errors (close-to or above 1) are highlighted in grey.
2695
2696 See "**TableS14_new2.xlsx**" file as the table is too large to reasonably fit in a A4 page format.
2697

TableS15: NN-ABC posterior parameter estimation of all parameters in Scenario i1-1b for results from 25 sets of five populations each, combined altogether.
Entire posterior distributions for each 25 analyses separately, for which summary statistics are presented in **TableS3**, are combined (concatenated) together before re-computing the Mode, Median, Mean, 50% CI and 90% CI presented in this table. Therefore, values in the table correspond to posterior densities plotted in the right panels of **Figure7**, **Figure8**, **FigureS20**, and **FigureS21**. Parameter definitions and priors are provided in **TableS8** and represented graphically in the corresponding scenario panel of **Figure5**. Values in this table are used for the synthetic schematic representation of the demographic and migration history of Central and Southern African populations presented in **Figure9**. Values in italic are not satisfactorily departing from the priors and are associated with high cross-validation posterior parameter estimation errors in the vicinity of the observed data (**TableS14**).

Parameter	Mode	Median	Mean	50% CI	90% CI
N_{nKS}	**13387**	**27960**	**33648**	**16400-46042**	**8648-78425**
N_{sKS}	**11496**	**22584**	**29465**	**12698-40535**	**6158-76677**
N_{wRHG}	**17083**	**23544**	**29128**	**15201-37922**	**7606-71133**
N_{eRHG}	**10702**	**17589**	**24112**	**10882-31019**	**5769-67086**
N_{RHGn}	**16730**	**31250**	**34896**	**18914-47062**	**9111-74177**
N_{AKS}	*62991*	*57534*	*55788*	*39887-72762*	*16051-90666*
N_{ARHG}	*53641*	*51921*	*51674*	*33228-69906*	*12751-90806*
$N_{ARHG-RHGn}$	*61127*	*55333*	*53839*	*35566-73204*	*11601-91048*
N_A	**17005**	**26518**	**31283**	**17496-40248**	**9890-69775**
t_{KS}	**2857**	**2881**	**2969**	**2304-3509**	**1624-4662**
t_{RHG}	**751**	**970**	**1089**	**670-1374**	**370-2222**
$t_{RHG-RHGn}$	**6347**	**6478**	**6617**	**5462-7566**	**4098-9734**
$t_{KS-RHG-RHGn}$	**13254**	**12412**	**11909**	**10715-13411**	**8044-14397**
$tad_{nKS-sKS}$	**550**	**611**	**718**	**434-847**	**227-1580**
$tad_{nKS-RHGn}$	**1241**	**1406**	**1524**	**969-1956**	**447-3003**
$tad_{sKS-RHGn}$	**1045**	**1243**	**1354**	**829-1739**	**406-2678**
$tad_{wRHG-RHGn}$	**239**	**310**	**373**	**197-478**	**86-874**
$tad_{eRHG-RHGn}$	**277**	**446**	**528**	**278-694**	**135-1190**
$tad_{wRHG-eRHG}$	**332**	**449**	**547**	**290-701**	**130-1263**
$tad_{RHGn-ARHG}$	**2905**	**3161**	**3369**	**2370-4114**	**1435-6007**
$tad_{AKS-RHGn}$	**4492**	**4483**	**4582**	**3710-5303**	**2728-6825**
$tad_{AKS-ARHG}$	**4228**	**4427**	**4579**	**3691-5261**	**2814-6954**
$tad_{AKS-ARHG-RHGn}$	**8881**	**9332**	**9307**	**7970-10762**	**5923-12505**
m(nKS-sKS)	*0.6058*	*0.6129*	*0.5888*	*0.4309-0.7707*	*0.1613-0.9241*
m(sKS-nKS)	*0.7823*	*0.6187*	*0.5925*	*0.4298-0.7824*	*0.1544-0.9309*
m(nKS-RHGn)	*0.3921*	*0.4800*	*0.4918*	*0.3218-0.6569*	*0.1356-0.8794*
m(RHGn-nKS)	*0.4897*	*0.5207*	*0.5203*	*0.3162-0.7340*	*0.0859-0.9382*
m(sKS-RHGn)	*0.5539*	*0.5241*	*0.5215*	*0.3463-0.6994*	*0.1343-0.8974*
m(RHGn-sKS)	*0.4941*	*0.4824*	*0.4816*	*0.2931-0.6633*	*0.0872-0.8868*
m(RHGn-wRHG)	*0.3666*	*0.5076*	*0.5055*	*0.2935-0.7189*	*0.0839-0.9216*
m(wRHG-RHGn)	*0.4899*	*0.5039*	*0.5038*	*0.3374-0.6693*	*0.1311-0.8837*
m(RHGn-eRHG)	*0.4183*	*0.4678*	*0.4749*	*0.2694-0.6774*	*0.0795-0.8886*
m(eRHG-RHGn)	*0.7038*	*0.6242*	*0.5957*	*0.4363-0.7746*	*0.1733-0.9270*
m(wRHG-eRHG)	*0.5720*	*0.5104*	*0.5087*	*0.3200-0.6976*	*0.1213-0.8929*
m(eRHG-wRHG)	*0.5011*	*0.4922*	*0.4936*	*0.3066-0.6788*	*0.0969-0.8911*
m(RHGn-ARHG)	*0.6155*	*0.5087*	*0.5011*	*0.3188-0.6820*	*0.1078-0.8798*
m(ARHG-RHGn)	*0.5104*	*0.4846*	*0.4893*	*0.2734-0.6991*	*0.0869-0.9104*
m(ARHG-AKS)	*0.6493*	*0.5414*	*0.5299*	*0.3316-0.7375*	*0.0935-0.9294*
m(AKS-ARHG)	*0.5627*	*0.4814*	*0.4807*	*0.2950-0.6624*	*0.0897-0.8766*
m(RHGn-AKS)	*0.5286*	*0.4940*	*0.4913*	*0.3026-0.6776*	*0.0998-0.8863*
m(AKS-RHGn)	*0.5057*	*0.4919*	*0.4888*	*0.3026-0.6751*	*0.0912-0.8781*
m(AKS-ARHG-RHGn)	*0.5948*	*0.5145*	*0.5096*	*0.3263-0.6970*	*0.1123-0.8887*
m(ARHG-RHGn-AKS)	*0.2813*	*0.4507*	*0.4642*	*0.2482-0.6707*	*0.0761-0.8966*

<u>**TableS16**</u>**: NN-ABC parameter inference posterior check: Fligner and F tests of homogeneity of variance between the**
NN-ABC posterior-parameter combined (concatenated) distributions obtained for 25 different observed combinations
of five sampled-populations and 25 pseudo-observed simulated datasets under the Scenario i1-1b proposed in Figure9
and TableS15.
Each line corresponds to a single scenario-parameter under Scenario i1-1b (**Figure5** and **TableS8**). For each parameter in
Scenario i1-1b, the first two columns provide the Mode and Variance of posterior distributions plotted in the right panels of
Figure7, **Figure8**, **FigureS20**, and **FigureS21**, and corresponding **TableS15** and synthetic **Figure9**. The third and fourth
columns provide the Mode and Variance of NN-ABC posterior-parameter estimated distributions obtained as for the 25
observed datasets but using instead 25 pseudo-observed simulated datasets, each obtained using the same point-value of
scenario-parameters provided in the first column (see **Material and Methods**). Therefore, values in the third and fourth
columns correspond to the Mode and Variance of the combined (concatenated) densities of **FigureS23**, **FigureS24**,
FigureS25, and in **FigureS26**. Fifth column corresponds to scaled difference in observed variance between NN-ABC results
obtained from observed data and NN-ABC results obtained from pseudo-observed simulations. Sixth and Seventh columns
present results from Fligner tests of homogeneity of variance performed between NN-ABC posterior parameter combined
distributions obtained between observed data and pseudo-observed simulations. The eighth and ninth columns present results
from simple F-tests of homogeneity of variance between the same distributions.
Values in italic correspond to scenario-parameters NN-ABC posterior estimates that are not satisfactorily departing from the
priors and are associated with large posterior CI and high cross-validation posterior parameter estimation errors in the vicinity
of the observed data (**TableS14**).
2730
2731
2732

2733 <u>**TableS16**</u>
2734

Parameter	25 Observed combinations of five sampled-populations		25 Pseudo-observed simulations		VarObs. / VarPseudoObs.	Fligner_test statistic	p-value	F_test statistic	p-value
	Mode	**Variance**	**Mode**	**Variance**					
N_{nKS}	13387	470652408	15022	244843368	1.9223	2834.9549	0	1.9223	0
N_{KS}	11496	480532055	10123	335167982	1.4337	757.9420	7.53e-167	1.4337	0
N_{wRHG}	17083	374510938	37740	405907201	0.9227	538.9419	3.20e-119	0.9227	1.97e-10
N_{eRHG}	10702	360795986	13586	396844095	0.9092	1008.5197	2.53e-221	0.9092	5.19e-14
N_{RHGn}	16730	399707441	37420	463160255	0.8630	242.5899	1.07e-54	0.8630	2.50e-31
N_{AKS}	*62991*	*505119215*	*54376*	*457143763*	*1.1049*	*60.2751*	*8.25e-15*	*1.1049*	*3.11e-15*
N_{ARHG}	*53641*	*563875758*	*69474*	*409652710*	*1.3765*	*801.0610*	*3.17e-176*	*1.3765*	*0*
$N_{ARHG-RHGn}$	*61127*	*590438189*	*48979*	*618052359*	*0.9553*	*12.1288*	*0.000496486*	*0.9553*	*0.00030234*
N_A	17005	345070021	8977	38413042	8.9831	12169.9104	0	8.9831	0
t_{KS}	2857	889610	2370	697809	1.2749	504.5351	9.80e-112	1.2749	0
t_{RHG}	751	351383	758	287469	1.2223	662.0090	5.46e-146	1.2223	0
$t_{RHG-RHGn}$	6347	2909824	6931	2181740	1.3337	301.5261	1.53e-67	1.3337	0
$t_{KS-RHG-RHGn}$	13254	4003176	13061	1573745	2.5437	3359.4043	0	2.5437	0
$tad_{nKS-sKS}$	550	232999	433	185207	1.2580	197.0105	9.38e-45	1.2580	0
$tad_{nKS-RHGn}$	1241	614188	1430	567071	*1.0831*	*16.8698*	*4.00e-05*	1.0831	2.81e-10
$tad_{sKS-RHGn}$	1045	534449	1283	458118	1.1666	83.2224	7.33e-20	1.1666	0
$tad_{wRHG-RHGn}$	239	70184	311	59430	*1.1810*	*11.0390*	*0.000892165*	1.1810	0
$tad_{eRHG-RHGn}$	277	119325	341	38320	3.1140	4191.0354	0	3.1140	0
$tad_{wRHG-eRHG}$	332	158937	389	105809	1.5021	350.3356	3.58e-78	1.5021	0
$tad_{RHGn-ARHG}$	2905	2067072	3276	2030903	1.0178	32.9386	9.51e-09	1.0178	0.162856127
$tad_{AKS-RHGn}$	4492	1599960	4421	1374801	1.1638	106.9052	4.67e-25	1.1638	0
$tad_{AKS-ARHG}$	4228	1605327	4357	1327935	1.2089	113.3627	1.80e-26	1.2089	0
$tad_{AKS-ARHG-RHGn}$	8881	4027523	9368	2160405	1.8642	2194.7651	0	1.8642	0
m(nKS-sKS)	*0.60584*	*0.05318*	*0.61484*	*0.05479*	*0.9706*	*18.7261*	*1.51e-05*	*0.9706*	*0.018451958*
m(sKS-nKS)	*0.78226*	*0.05541*	*0.66954*	*0.06263*	*0.8847*	*161.0983*	*6.51e-37*	*0.8847*	*3.48e-22*
m(nKS-RHGn)	*0.39207*	*0.05045*	*0.32679*	*0.05789*	*0.8715*	*221.6304*	*3.99e-50*	*0.8715*	*1.66e-27*
m(RHGn-nKS)	*0.48966*	*0.06839*	*0.39841*	*0.05102*	*1.3404*	*744.3383*	*6.83e-164*	*1.3404*	*0*
m(sKS-RHGn)	*0.55393*	*0.05338*	*0.37490*	*0.05968*	*0.8945*	*156.9520*	*5.24e-36*	*0.8945*	*1.24e-18*
m(RHGn-sKS)	*0.49410*	*0.05850*	*0.43655*	*0.06190*	*0.9451*	*55.1932*	*1.09e-13*	*0.9451*	*8.10e-06*
m(RHGn-wRHG)	*0.36658*	*0.06794*	*0.35286*	*0.06667*	*1.0191*	*1.6272*	*0.202083905*	*1.0191*	*0.134522467*
m(wRHG-RHGn)	*0.48987*	*0.05066*	*0.53936*	*0.04750*	*1.0667*	*37.2593*	*1.03e-09*	*1.0667*	*3.33e-07*
m(RHGn-eRHG)	*0.41833*	*0.06319*	*0.86078*	*0.06300*	*1.0030*	*0.0004*	*0.984365963*	*1.0030*	*0.812655593*
m(eRHG-RHGn)	*0.70376*	*0.05236*	*0.85783*	*0.05600*	*0.9351*	*37.1605*	*1.09e-09*	*0.9351*	*1.12e-07*
m(wRHG-eRHG)	*0.57199*	*0.05670*	*0.38485*	*0.05271*	*1.0758*	*56.7234*	*5.02e-14*	*1.0758*	*7.69e-09*
m(eRHG-wRHG)	*0.50114*	*0.05788*	*0.45735*	*0.06453*	*0.8970*	*129.8901*	*4.33e-30*	*0.8970*	*8.74e-18*
m(RHGn-ARHG)	*0.61550*	*0.05550*	*0.38558*	*0.06492*	*0.8549*	*293.0791*	*1.06e-65*	*0.8549*	*3.06e-35*
m(ARHG-RHGn)	*0.51038*	*0.06655*	*0.61734*	*0.05319*	*1.2511*	*572.2042*	*1.86e-126*	*1.2511*	*0*
m(ARHG-AKS)	*0.64930*	*0.06576*	*0.50831*	*0.06199*	*1.0608*	*58.4878*	*2.05e-14*	*1.0608*	*3.09e-06*
m(AKS-ARHG)	*0.56272*	*0.05714*	*0.52198*	*0.07406*	*0.7715*	*758.6786*	*5.20e-167*	*0.7715*	*3.11e-93*
m(RHGn-AKS)	*0.52861*	*0.05774*	*0.69502*	*0.06205*	*0.9304*	*49.7306*	*1.76e-12*	*0.9304*	*1.19e-08*
m(AKS-RHGn)	*0.50575*	*0.05785*	*0.61882*	*0.06447*	*0.8973*	*153.6100*	*2.82e-35*	*0.8973*	*1.10e-17*
m(AKS-ARHG-RHGn)	*0.59481*	*0.05661*	*0.27899*	*0.06865*	*0.8247*	*438.1366*	*2.75e-97*	*0.8247*	*2.41e-52*
m(ARHG-RHGn-AKS)	*0.28125*	*0.06601*	*0.10737*	*0.08326*	*0.7929*	*669.5820*	*1.23e-147*	*0.7929*	*4.86e-75*

2735
2736