Title:
Ancient tree-topologies and gene-flow processes among human lineages in Africa

Short Title:
Ancient demographic histories of African genetic lineages

Gwenna Breton 1,2*
Per Sjödin 1
Panagiotis I. Zervakis 1
Romain Laurent 3
Alain Froment 4
Agnès E. Sjöstrand 1
Barry S. Hewlett 5
Luis B. Barreiro 6
George H. Perry 7,8
Himla Soodyall 9,10
Evelyne Heyer 3
Carina M. Schlebusch 1,11,12,*
Mattias Jakobsson 1,11,12,*
Paul Verdu 2,*

Human Evolution, Department of Organismal Biology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18C, SE-752 36 Uppsala, Sweden
Centre for Medical Genomics, Department of Clinical Genetics and Genomics, Sahlgrenska University Hospital, Gothenburg, Sweden
UMR7206 Eco-anthropology, CNRS-MNHN-Université Paris Cité, Paris, France
UMR 208 Patrimoines locaux, IRD-MNHN, Paris, France
Department of Anthropology, Washington State University, Vancouver, Washington, USA
Department of Medicine, Section of Genetic Medicine, University of Chicago, Chicago, IL, USA
Department of Anthropology, Pennsylvania State University, University Park, PA 16801, USA
Department of Biology, Pennsylvania State University, University Park, PA 16801, USA
Division of Human Genetics, School of Pathology, Faculty of Health Sciences, University of the Witwatersrand and National Health Laboratory Service, Johannesburg, South Africa
Academy of Science of South Africa
Palaeo-Research Institute, University of Johannesburg, P.O. Box 524, Auckland Park, 2006, South Africa
SciLifeLab, Uppsala, Sweden

* Authors for correspondence:
gwenna.breton@gu.se,   paul.verdu@mnhn.fr,   mattias.jakobsson@ebc.uu.se, carina.schlebusch@ebc.uu.se

Keywords:
Africa; demographic history; admixture; migration; machine-learning; Approximate Bayesian Computations; Whole Genome Sequences; Rainforest Hunter-Gatherers; Agriculturalists; Khoe and San populations.

Authors' contributions
GB – Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software,
Supervision, Validation, Visualization, Writing – Original Draft Preparation.
PS – Conceptualization, Formal Analysis, Supervision, Validation, Writing – Review and Editing.
PLZ – Data Curation, Software, Writing – Review and Editing.
RL – Methodology, Software, Validation, Writing – Review and Editing.
AF – Investigation, Resources, Writing – Review and Editing.
AES – Investigation, Resources, Writing – Review and Editing.
BSH – Investigation, Resources, Writing – Review and Editing.
LBB – Investigation, Resources, Writing – Review and Editing.
GHP – Investigation, Resources, Writing – Review and Editing.
HS – Investigation, Resources, Writing – Review and Editing.
EH – Investigation, Resources, Writing – Review and Editing.
CMS – Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration,
Resources, Supervision, Validation, Writing – Review and Editing.
MJ – Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration,
Resources, Supervision, Validation, Writing – Review and Editing.
PV – Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project
Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft
Preparation.

74
75

<u>Abstract</u>

The deep history of humans in Africa and the complex divergences and migrations among ancient human genetic lineages remain poorly understood and are the subject of ongoing debate. We produced 73 high-quality whole genome sequences from 14 Central and Southern African populations with diverse, well-documented, languages, subsistence strategies, and socio-cultural practices, and jointly analyze this novel data with 104 African and non-African previously-released whole genomes. We find vast genome-wide diversity and individual pairwise differentiation within and among African populations at continental, regional, and even local geographical scales, often uncorrelated with linguistic affiliations and cultural practices. We combine populations in 54 different ways and, for each population combination separately, we conduct extensive machine-learning Approximate Bayesian Computation inferences relying on genome-wide simulations of 48 competing evolutionary scenarios. We thus reconstruct jointly the tree-topologies and migration processes among ancient and recent lineages best explaining the diversity of extant genomic patterns. Our results show the necessity to explicitly consider the genomic diversity of African populations at a local scale, without merging population samples indiscriminately into larger *a priori* categories based on geography, subsistence-strategy, and/or linguistics criteria, in order to reconstruct the diverse evolutionary histories of our species. We find that, for all different combinations of Central and Southern African populations, a tree-like evolution with long periods of drift between short periods of unidirectional gene-flow among pairs of ancient or recent lineages best explain observed genomic patterns compared to recurring gene-flow processes among lineages. Moreover, we find that, for 25 combinations of populations, the lineage ancestral to extant Southern African Khoe-San populations diverged around 300,000 years ago from a lineage ancestral to Rainforest Hunter-Gatherers and neighboring agriculturalist populations. We also find that short periods of ancient or recent asymmetrical gene-flow among lineages often coincided with epochs of major cultural and ecological changes previously identified by paleo-climatologists and archaeologists in Sub-Saharan Africa.

Introduction

Unraveling genetic structure and gene flow among human lineages in Africa is crucial to the
understanding of the biological evolution and diversity of *Homo sapiens* throughout the continent
(Henn, Steele and Weaver, 2018; Schlebusch and Jakobsson, 2018; Pfennig *et al.*, 2023).
Population geneticists largely agree today that *Homo sapiens* spent a large part of its genetic
evolution within Africa only, between its gradual emergence from anatomically archaic forms 600.000-
200,000 years ago and the beginning of the Out-of-Africa expansions to the rest of the world around
100,000-50,000 years ago (Schlebusch and Jakobsson, 2018). For the past 20 years, the detailed
demographic and evolutionary history that shaped the genetic diversity of extant populations since the
Out-of-Africa has been, and is still, the subject of numerous investigations in various regions of Africa
(Verdu *et al.*, 2009, 2013; Schlebusch *et al.*, 2012; Breton *et al.*, 2014; Patin *et al.*, 2014, 2017; Perry
et al., 2014; Busby *et al.*, 2016; Pierron *et al.*, 2017; Semo *et al.*, 2020; Lucas-Sánchez, Serradell and
Comas, 2021; Sengupta *et al.*, 2021; Fortes-Lima *et al.*, 2022, 2024; Laurent *et al.*, 2023; Pfennig *et al.*,
2023).
However, how ancient genetic divergences, and the dynamics of admixture and/or migration events
among ancient human lineages, influenced the genetic landscape observed today throughout Africa,
largely remains to be assessed (Bergström *et al.*, 2021). Classically tested demographic models
(Stringer, 2002; Henn, Steele and Weaver, 2018; Hollfelder *et al.*, 2021; Ragsdale *et al.*, 2023), range
from a unique ancestral *Homo sapiens* genetic population having recently and rapidly diverged into
extant African populations via series of founding events and/or multifurcations -often referred to as
tree-like models-, to models where extant populations diverged in a remote past and remained isolated
over long periods of time until relatively recently -often referred to as multiregional models-. Moreover,
each model may encompass possible gene exchanges among lineages via migration or admixture
processes. Importantly, the timing, duration, and/or intensity of such gene-flow events may
fundamentally change the most likely topology of bifurcating tree-like models compared to results
obtained without gene flows, and may also create reticulations among pairs of lineages throughout
history (Ragsdale *et al.*, 2023). Finally, these classical albeit highly complex models have recently been
enriched with possible introgression events from now extinct, unknown, or unsampled non-*Homo*
sapiens or ancient "ghost" *Homo-sapiens* lineages (Lachance *et al.*, 2012; Lorente-Galdos *et al.*, 2019;
Lipson *et al.*, 2022; Fan *et al.*, 2023; Pfennig *et al.*, 2023; Ragsdale *et al.*, 2023). Such latter models
became plausible in Africa in analogy to the likely events of introgressions from now extinct hominid
species into *Homo sapien*s lineages unveiled outside of Africa by the major advances of paleogenomics
(Meyer *et al.*, 2012; Prüfer *et al.*, 2014).
In this context, several recent investigations tested a variety of the above models with maximum-
likelihood approaches relying on whole genome sequences from several extant populations sampled
throughout the continent. They often reached highly contrasted results, with different ancient tree-
topologies between different numbers of ancient lineages (Schlebusch and Jakobsson, 2018; Lorente-
Galdos *et al.*, 2019; Lipson *et al.*, 2022; Fan *et al.*, 2023; Ragsdale *et al.*, 2023). Furthermore, they
found that reticulations among a limited number of ancient lineages explained observed genetic patterns
without the necessity of archaic introgressions (Ragsdale *et al.*, 2023); or, instead, that admixture with
other non-*Homo sapiens* or ancient *Homo sapiens* "ghost" populations most satisfactorily explained the
results (Lachance *et al.*, 2012; Lorente-Galdos *et al.*, 2019; Lipson *et al.*, 2022; Fan *et al.*, 2023).
A possible source of such vast differences among obtained results is likely the variety of population
sets considered in each study at a continental scale. Indeed, genetic diversity and differentiation among
populations is maximal in Africa (Tishkoff *et al.*, 2009; Skoglund and Mathieson, 2018; Pfennig *et al.*,
2023), and previously shown to be due to substantial differences in local populations' demographic

histories (e.g. (Verdu *et al.*, 2009; Schlebusch *et al.*, 2012; Busby *et al.*, 2016; Patin *et al.*, 2017)).
Therefore, representing vast regions of the continent by a single population (Lorente-Galdos *et al.*,
2019; Ragsdale *et al.*, 2023), or even merging data from differentiated groups (Fan *et al.*, 2023), likely
influenced results obtained across studies and their interpretations.

In addition, and most importantly, all above models are highly nested, i.e. overlapping and
genetically largely indistinguishable for vast spaces of model-parameter values. Indeed, depending on
values of divergence times, effective population sizes, timing, duration and intensity of migration or
admixture events among lineages, tree-like, reticulated, and even multiregional models, with or without
ancient admixture, may be highly mimetic of one another. This essential methodological difficulty is
well illustrated by the often relatively similar values of posterior likelihoods of the various competing
models, thus indeed proved empirically hard to discriminate with the maximum-likelihood approaches
deployed, even more so when competing models differ in the number and specifications of parameters
explored with different statistics. These fundamental statistical issues remain despite the use of high-
quality whole genomes, an increasing number of populations considered at once, and increasingly
powerful methods based on innovative statistics (Gravel, 2012; Lorente-Galdos *et al.*, 2019; Ragsdale
and Gravel, 2019; Kamm *et al.*, 2020; Fan *et al.*, 2023; Ragsdale *et al.*, 2023). Finally, the lack of
ancient DNA data older than a few thousand years in Africa, strongly hampers the empirical testing of
the possible occurrence of ancient *Homo sapiens* "ghost" or non-*Homo sapiens* introgression events in
Homo sapiens lineages in Africa (Lachance *et al.*, 2012; Skoglund and Mathieson, 2018; Lorente-
Galdos *et al.*, 2019; Lipson *et al.*, 2022; Ragsdale *et al.*, 2023).

In this context, we investigated 74 high coverage (>30x) whole genome sequences from an
anthropologically well-characterized sample of hunter-gatherer, herder, and agricultural neighboring
populations from Central and Southern Africa (**FigureF1x**, **TableT1x**), merged together with 105
previously published high-quality whole genomes (SGDP-Mallick et al. 2016, HGDP-Meyer et al.
2012, 1KGP-Auton et al. 2015, Rasmussen et al. 2014, SAHGP-Choudhury et al. 2017). In particular,
we investigated genomic diversity patterns and aimed at reconstructing ancient demographic histories
among lineages having led to different groups of Khoe-San hunter-gatherer or herder populations from
Southern Africa (Schlebusch et al. 2012), and to different groups of both Eastern or Western Congo
Basin hunter-gatherer populations (often historically designated "Pygmies" by Europeans) and their
agriculturalist ("non-Pygmy") neighbors with whom they share, nowadays, complex socio-economic
interactions (Verdu et al. 2013; Hewlett 2014). Investigating detailed ancient demography and gene-
flow among these three groups of populations, taking explicitly into account local differentiations and
possible gene-flow among and within groups, is of particular interest as they have previously been
identified to be among the most deeply diverged lineages in our species (Li *et al.*, 2008; Tishkoff *et al.*,
2009; Schlebusch and Jakobsson, 2018; Schlebusch *et al.*, 2020; Pfennig *et al.*, 2023).

We conducted formal statistical choice of competing scenarios and subsequent joint estimation of
parameters under the winning scenario using machine-learning Approximate Bayesian Computation
(ABC) (Tavaré *et al.*, 1997; Beaumont, Zhang and Balding, 2002; Blum and François, 2010; Csilléry,
François and Blum, 2012; Pudlo *et al.*, 2016), a methodological approach fundamentally differing from
all maximum-likelihood methods previously deployed, and only rarely explored in previous studies
reconstructing ancient demography in Africa (Lorente-Galdos *et al.*, 2019). ABC allows us to compare
a range of complex demographic scenarios presumed to have led to observed genetic patterns in
numerous population samples, and to estimate posterior parameter distributions best mimicking the data
under the winning scenario (e.g. (Verdu *et al.*, 2009; Lorente-Galdos *et al.*, 2019; Laurent *et al.*, 2023)).
Briefly, this is achieved based on informative summary-statistics computed on the observed data and
on numerous explicit genetic simulations for which scenario-parameters are drawn randomly in large
prior distributions set by the user. We thus aimed at inferring jointly competing tree-topologies,
divergence times, effective population size changes, and timing, duration and intensity of possible

asymmetric gene-flow events, to reconstruct the detailed evolutionary mechanisms underlying the genomic diversity of extant Central and Southern African populations.

FigureF1x: Sampling location and number of whole-genome sequenced individuals.
Individuals and populations originally sequenced in this work are indicated with filled circles. Individuals and populations from previously published data merged with the original data set for comparison purposes are indicated with open circles. Circles' diameters are proportional to the number of individuals as indicated in the top row of the legend in the bottom left corner.

FigureF1x

216 --

217 **TableT1x: Population table**

218 Geographical location of population samples are indicated in **FigureF1x**.

Population Name[1]	N[2]	Dataset[3]	Sampling Location	Language Family (language)	Close socio-economic interactions with	Samples included in the ROH-ASD-ADMIXTURE analyses (FigureF2x, FigureF3x, FigureF4x)	Samples included in the ABC analyses (FigureF5x, FigureF6x, FigureF7x, FigureF8x, FigureF9x)
Original dataset							
Baka	7	this study	Bosquet (Cameroon)	Niger-Congo non Bantu Adamawa Ubangian Gbanzili (Baka)	Nzime	Yes	wRHG, 5 individuals with highest coverage
Nzime	5	this study	Messea (Cameroon)	Bantu A842 (Nzime)	Baka	Yes	RHGn (West)
Ba.Kola	5	this study	Dispersed between Lolodorf and Kribi (Cameroon)	Bantu A80 (Kola)	Ngumba	Yes	wRHG
Ngumba	5	this study	Dispersed between Lolodorf and Kribi (Cameroon)	Bantu A80 (Ngumba)	Ba.Kola	Yes	RHGn (West)
Bi.Aka_Mbati	5	this study	Bombeketi section of Bagandou (Central African Republic)	Bantu C10 (Mbati)	Mbati farmers (not in the dataset)	Yes	wRHG
Ba.Kiga	5	this study	Mukono (Uganda)	Bantu J10 (Kiga)	Ba.Twa	Yes	RHGn (East)
Ba.Twa	6	this study	Kebiremu, Byumba, Kitariro, Mgungu, Nteko (Uganda)	Bantu J11 (Twa)	Ba.Kiga	Yes	eRHG, 5 individuals with highest coverage
Nsua	5	this study	Bundimassoli (Uganda)	Sudanic Mangbutu (Efe)	Ba.Konjo	Yes	eRHG
Ba.Konjo	5	this study	Mulimassenge (Uganda)	Bantu J40 (Konjo)	Nsua	Yes	RHGn (East)
!Xun	5	this study	Omega camp (Namibia) and Schmidtsdrift (Sout Africa)[4]	Khoisan (Ju)	Not applicable	Yes	nKS
Ju|'hoansi	5	this study	Tsumkwe (Namibia)	Khoisan (Ju)	Not applicable	Yes	nKS
Nama	5	this study	Windhoek (Namibia)	Khoisan (KhoeKhoe)	Not applicable	Yes	sKS
Karretjie People	5	this study	Colesberg (South Africa)	Khoisan (Tuu ancestral language) / Indo-European (Afrikaans current language)	Not applicable	Yes	sKS
Khutse-San	5	this study	Kutse Game reserve (Botswana)	Khoisan (Khoe Kalahari)	Not applicable	Yes	No
Comparative dataset							
Bantu_Herero	2	SGDP	Namibia	Bantu R30 (Herero)	Not applicable	Yes	No
Bantu_Kenya	2	SGDP	Kenya	No data	Not applicable	Yes	No
Bantu_Tswana	2	SGDP	South Africa	Bantu S31 (Tswana)	Not applicable	Yes	No
Biaka	2	SGDP	Central African Republic	Bantu C10 (Aka)	Not applicable	Yes	No
Coloured	8	SAHGP	South Africa	Indo-European (Afrikaans)	Not applicable	Yes	No
Dinka	4	SGDP, HGDP	Sudan	Nilo-Saharan (Dinka)	Not applicable	Yes	No
Esan	3	SGDP, KGP	Nigeria	Niger-Congo non Bantu (Esan)	Not applicable	Yes	No
Gambian	2	SGDP	Gambia	Niger-Congo non Bantu (Mandinka)	Not applicable	Yes	No
Igbo	2	SGDP	Nigeria	Niger-Congo non Bantu (Igbo)	Not applicable	Yes	No
Ju|'hoansi_comp	4	SGDP, HGDP	Namibia	Khoisan (Ju)	Not applicable	Yes	No
‡Khomani	2	SGDP	South Africa	Khoisan (Tuu)	Not applicable	Yes	No
Kongo	1	SGDP	Cameroon	No data	Not applicable	Yes	No
Lemande	2	SGDP	Cameroon	Bantu A46 (Lemande)	Not applicable	Yes	No
Luhya	3	SGDP, KGP	Kenya	Bantu JE32 (Luhya)	Not applicable	Yes	No
Luo	2	SGDP	Kenya	Nilo-Saharan (Dholuo)	Not applicable	Yes	No
Maasai	2	SGDP	Kenya	Nilo-Saharan (Maasai)	Not applicable	Yes	No

Mandenka	4	SGDP, HGDP	Senegal	Niger-Congo non Bantu (West Maninkakan)	*Not applicable*	Yes	No
Mandinka	1	KGP	Gambia	Niger-Congo non Bantu (Mandinka)	*Not applicable*	Yes	No
Mbuti	5	SGDP, HGDP	Democratic Republic of Congo	Bantu D30 (Bambuti)	*Not applicable*	Yes	eRHG
Mende	3	SGDP	Sierra Leone	Niger-Congo non Bantu (Mende)	*Not applicable*	Yes	No
Mozabite	2	SGDP	Algeria	Afro-Asiatic (Mozabite)	*Not applicable*	Yes	No
Saharawi	2	SGDP	Western Sahara	Afro-Asiatic (Saharawi)	*Not applicable*	Yes	No
Somali	1	SGDP	Kenya	Afro-Asiatic (Somali)	*Not applicable*	Yes	No
Sotho	7	SAHGP	South Africa	Bantu S30 (Sotho)	*Not applicable*	Yes	No
Xhosa	8	SAHGP	South Africa	Bantu S41 (Xhosa)	*Not applicable*	Yes	No
Yoruba	4	SGDP, HGDP	Nigeria	Niger-Congo non Bantu (Yoruba)	*Not applicable*	Yes	No
Zulu	1	SAHGP	South Africa	Bantu S42 (Zulu)	*Not applicable*	Yes	No
French	4	SGDP, HGDP	France	Indo-European (French)	*Not applicable*	Yes	No
CEU	2	KGP	United States of America	Indo-European (English)	*Not applicable*	Yes	No
Dai	6	SGDP, KGP	China	Tai (Dai)	*Not applicable*	Yes	No
Papuan	6	SGDP, HGDP	Papua New Guinea	Papuan (no data)	*Not applicable*	Yes	No
Karitiana	5	SGDP, HGDP	Brazil	Tupian (Karitiana)	*Not applicable*	Yes	No

[1]Population Names are self-reported for the original dataset presented in this study
[2]Number of unrelated individuals considered in all analyses in this study (see **Material and Methods**)
[3]SGDP (Mallick *et al.*, 2016); KGP (The 1000 Genomes Project Consortium *et al.*, 2015); HGDP (Meyer *et al.*, 2012; Rasmussen *et al.*,
2014); SAHGP (Choudhury *et al.*, 2017).
[4]The place of origin of the !Xun is around Menongue in Angola.

226
227

<u>Results</u>

We generated high-coverage whole genome sequences (>30X) in 74 individuals from 14 Central and Southern African populations for whom detailed ethno-anthropological information was gathered in the field jointly with DNA samples (**FigureF1x** and **TableT1x**). We merged this original dataset with a comparative dataset comprising 105 individuals from 27 African and five non-African populations for whom similar high-quality raw whole genome sequencing data were made available to the community (Meyer *et al.*, 2012; Rasmussen *et al.*, 2014; The 1000 Genomes Project Consortium *et al.*, 2015; Mallick *et al.*, 2016; Choudhury *et al.*, 2017). After quality-control, variant-calling, and relatedness filtering procedures conducted on all 179 individuals together, we retained 73 and 104 unrelated individuals in our original and comparative datasets, respectively, for all subsequent analyses (see **Material and Methods**).

Considering only the 73 Central and Southern African unrelated individuals newly sequenced here, we identify a total of 26,780,319 biallelic SNPs (241,428 multiallelic SNPs), and 2,454,965 simple insertions/deletions (969,025 complex indels), compared to the reference sequence of the human genome GRCh38, out of which 854,114 (3.1893%) and 114,362 (4.6584%), respectively, were not previously reported in dbSNP 156 (**SupplementaryTableST1x**). Among the 177 unrelated worldwide individuals including our original dataset, we identify 36,272,545 biallelic SNPs (257,906 multiallelic SNPs), and 3,159,306 simple insertions/deletions (977,542 complex indels), out of which 1,055,245 (2.9092%) and 189,124 (5.9863%), respectively, were not previously reported in dbSNP 156 (**SupplementaryTableST1x**).

Furthermore, we find a large variability in the mean number of biallelic SNPs across populations (**FigureF2x-PanelA, SupplementaryTableST2x**). In particular, we find substantially more biallelic SNPs within African populations and more variation of the mean number of SNPs across African populations (from mean=3,726,018; SD=27,793 across 2 individuals in the Saharawi from Western Sahara, to mean=4,628,957; SD=8,139 across 5 individuals in the Ju|'hoansi from Namibia), than within and among non-African populations (from mean=3,214,086; SD=69,241 across 5 individuals in the Karitiana from Brazil, to mean=3,551,792; SD=7,819 across 2 individuals from the USA CEU population). Overall, Southern African Khoe-San populations exhibit the highest mean number of biallelic SNPs as well as numbers of previously unreported SNPs, followed by Central African Rainforest Hunter-Gatherer populations from the Congo Basin, and then by all other African populations in our dataset. These results show that a substantial number of previously unknown variants can still be found when investigating high-quality whole genome sequences from relatively under-studied Sub-Saharan African populations, consistent with previous studies (Meyer *et al.*, 2012; Rasmussen *et al.*, 2014; The 1000 Genomes Project Consortium *et al.*, 2015; Mallick *et al.*, 2016; Choudhury *et al.*, 2017; Schlebusch *et al.*, 2020; Breton, Fortes-Lima and Schlebusch, 2021; Fan *et al.*, 2023; Ragsdale *et al.*, 2023).

FigureF2x: Whole autosomal genome bi-allelic SNPs counts, unbiased heterozygosities, and distributions of Runs of Homozygosity by bins of length, in 37 African and 5 non-African populations.

(**A**) Numbers of bi-allelic SNPs across individuals within populations of more than one individual, compared to the reference sequence of the human genome GCRh38 and to previously reported variants in dbSNP 156. Detailed variant-counts are provided in **SupplementaryTableST1x and SupplementaryTableST2x**. (**B**) Unbiased estimates of multi-locus heterozygosities (Nei, 1978), averaged across all variable (bi-allelic SNPs only) and non-variable autosomal sites with no-missing genotype within each population of more than one individual, corrected then for haploid population sample sizes. (**C**) Mean total ROH lengths in four bins of length categories for each population of more than one individual, separately. For each panel, we present a box zooming specifically on the results obtained for the original dataset of 73 unrelated individuals whole-genome sequenced from Central and Southern Africa.

See **Material and Methods** for filtering and calculation details and the software packages used. For (**A**) and (**B**), box-plots indicate the population median in between the first and third quartile of the box-limits, whiskers extending to data points no more than 1.5 times the interquartile range of the distribution, and empty circles for all more extreme points beyond this limit, if any. Population geographical location, categorization, and descriptions are provided in **FigureF1x** and **TableT1x**.

----

283   **FigureF2x**

1. Large genetic diversities and differentiations in Central and Southern Africa

1.A. Heterozygosities and Runs of Homozygosity

We find very large differences across populations at a local scale in Central and Southern Africa in unbiased heterozygosities (Nei, 1978), calculated on both varying and non-varying autosomal sites (**FigureF2x-PanelA-B** and **SupplementaryTableST2x**). Indeed, we find varying genetic variation across Northern Khoe-San populations (!Xun and Ju|'hoansi) and Southern KS populations (Nama, Karretjie People and #Khomani). Furthermore, we find overall more variation in Rainforest Hunter-Gatherer populations from the western part of the Congo Basin than in RHG populations from the east; and western RHG populations exhibited much larger genetic diversities than all their respective RHG neighbors.

In addition, we investigated the distributions of Runs of Homozygosity (ROH) across individuals within each of the 46 populations in our dataset. In general (**FigureF2x-PanelC**), individuals from Eastern RHG populations (Nsua, Mbuti, Ba.Twa), exhibit the longest total proportion of their autosomal genome in ROH across Sub-Saharan African populations, for all ROH length-classes, with the exception of #Khomani individuals from Southern Africa, and that of Coloured individuals for short ROH only. Interestingly, we find that Western RHG populations (Ba.Kola, Baka, Biaka, Bi.Aka_Mbati) have much less ROH of all classes than Eastern RHG. Furthermore, we find less short ROH, but more long ROH, in Western RHG than in all RHG neighboring populations across the Congo Basin except for the Ba.Kiga RHGn from Uganda. Finally, we find that all Northern and Southern Khoe-San populations have relatively similar total lengths of short ROH compared to that of RHG neighbors, with the notable exception of the !Xun, who have the smallest proportion of their genomes in short ROH, worldwide. Nevertheless, we find substantially more ROH of longer class in KS populations than in RHG neighbors, a relative pattern similar to what was observed for Western RHG. The distribution of ROH sizes has been shown to be highly informative about important demographic processes such as levels of inbreeding or endogamy within populations, and reproductive isolation or recent admixture among populations (Pemberton *et al.*, 2012; Mooney *et al.*, 2018; Szpiech *et al.*, 2019; Laurent *et al.*, 2023). Thus, our ROH results highlight possibly vast differences in recent demographic processes and levels of endogamy across Central and Southern African populations.

Altogether, our heterozygosity and ROH results show the vast genetic diversity of Sub-Saharan African populations, as well as substantially diverging genomic patterns at a local geographical scale among and within groups of Rainforest Hunter-Gatherers, RHG neighbors, and Khoe-San populations.

1.B. Individual pairwise genetic differentiation

We investigated individual pairwise genomic differentiation across the 177 worldwide unrelated individuals in our dataset, including the 73 unrelated novel Central and Southern African individuals, using Allele-Sharing Dissimilarities (ASD, (Bowcock *et al.*, 1991)). The Neighbor Joining Tree (NJT, (Saitou and Nei, 1987; Gascuel, 1997)) representation of the pairwise ASD matrix shows three main clusters among all African individuals, separated by a longer genomic distance from all non-African individuals (**FigureF3x-PanelA**). One cluster corresponds to all Southern African Khoe-San populations (represented by the pink and red symbols, see **FigureF3x-PanelB**), largely separated from a cluster corresponding to Rainforest Hunter-Gatherer populations from the Congo Basin (represented by the blue symbols); itself also largely separated from a third cluster grouping all other Western, Eastern, Southern and Central African populations in our data set, the latter group including Rainforest Hunter-Gatherer neighboring populations in green symbols (see, **FigureF3x-PanelB)**.

FigureF3x: Individual pairwise genetic differentiation patterns.

Neighbor-Joining Tree NJT (Saitou and Nei, 1987; Gascuel, 1997) and Multi-Dimensional scaling representation of genome-
wide Allele Sharing Dissimilarities (Bowcock *et al.*, 1991) among pairs of individuals at worldwide and regional African
scales. We considered the 14,182,615 genome-wide autosomal SNPs pruned for low LD (r^2 threshold 0.1) to calculate ASD
between all pairs of individuals. (**A**) NJT computed for all 177 worldwide individuals in the ASD matrix. For easing the
visualization of the internal branches of the NJT, all terminal edges are represented in dotted lines each measuring 1/10th of
their true size. (**B**) Individual symbols and colors identifying the 73 Central and Southern African individuals originally whole-
genome sequenced with filled symbols, and the 104 individuals from worldwide populations, including Africa, merged with
the original data with open symbols. (**C**) and (**D**) First three axes of ASD-MDS computed on a subset of the full ASD matrix
comprising only the individuals sequenced anew, whose symbols are provided in (**B**). (**E**) NJT computed on the same
individual subset of the ASD matrix used for (**C**) and (**D**). Terminal edges of this latter tree are represented with dotted lines
each measuring 1/30th of their true size.

FigureF3x

352
353
354

Note importantly that individual pairwise dissimilarities at the genome-wide scale do not reflect
geography (**FigureF1x**, **TableT1x**), highlighting the large genetic differentiation found across African
populations and the complexity of its distribution at both a local and continental scale. From West to
East of the Congo Basin throughout Central Africa, RHG populations cluster separately from RHG
geographic neighbors with whom they nevertheless share close socio-economic interactions.
Furthermore, Central African RHGn are more genetically resembling certain other Western, Eastern
and Southern Africans than their immediate RHG geographical neighbors. Analogously, Southern
African Khoe-San populations are also clustering together and much more distant from other Southern
African geographically neighboring populations.
The Multidimensional Scaling (MDS) projection of the subsampled ASD matrix for only the 73
novel individuals from Central and Southern Africa, further shows substantial differentiation among
groups of individuals within each one of the three clusters identified at the continental scale. Indeed,
FigureF3x-PanelC, **D** and **E** show substantial differentiation between Western and Eastern Central
African Rainforest Hunter-Gatherer populations, and even substantial differentiation between Ba.Twa
and Nsua RHG very locally in Uganda. Conversely, we find relatively much shorter genetic
differentiation among pairs of RHG neighbors from West to East of the entire Congo Basin. Finally,
results also show substantial differentiation across Southern African Khoe-San populations, albeit these
populations are relatively closer from one another than the different RHG populations.
NJT and MDS based on the ASD pairwise matrix provide an important view of the major axes of
genetic differentiation and variation across samples, but do not easily allow to visually describe higher-
order axes of genomic variations. To do so, we conducted an ADMIXTURE analysis (Alexander et al.
2009), on the same entire worldwide dataset for increasing values of K (**FigureF4x**). This descriptive
method is known to capture the same information as ASD-MDS and ASD-NJT, but allows to explore
multiple axes of variation at the same time (Pritchard, Stephens and Donnelly, 2000; Rosenberg, 2002;
Falush, Stephens and Pritchard, 2003; Alexander, Novembre and Lange, 2009; Lawson, van Dorp and
Falush, 2018; Peter, 2022; Laurent *et al.*, 2023).

FigureF4x:
Worldwide interindividual genetic diversity patterns with ADMIXTURE
Each individual is represented by a single vertical line divided in K colors each proportional to an individual's genotype
membership proportion assigned by ADMIXTURE (Alexander, Novembre and Lange, 2009) into the virtual cluster of that
color. Two individuals showing the same relative proportions of each color at a given value of K are genetically more
resembling one another than two individuals with different relative proportions of the same colors. Population and regional
geographic groupings are not considered for the calculations, individuals are *a posteriori* grouped by populations and ordered
by the user. Each population is separated by a thin vertical black line. Unsupervised clustering conducted with 840,031
genome-wide SNPs pruned for LD (r^2 threshold 0.1) and minor allele frequency (0.1) using ADMIXTURE for values of K
ranging from 2 to 10, considering 177 worldwide individuals. For each value of K separately, we performed 20 independent
runs and used PONG (Behr *et al.*, 2016), to identify groups of resembling results (called "modes") based on Symmetric
Similarity Coefficient measures of individual results. The number of runs among the 20 independent runs belonging to the
same mode result (i.e. that have pairwise SSC above 0.998), is indicated below the number of K on the left of each barplot
separately. Only those runs are averaged per individual to provide the barplot result for each value of K separately.

398
399

400   **FigureF4x**

At K=2 in **FigureF4x**, the green virtual genetic cluster is maximized in Northern KS populations
while the gray cluster is maximized in non-African populations, consistent with largest genetic distances
in the ASD-NJT in **FigureF3x**, with all other individuals in our dataset presenting intermediate, non-
resolved, genotype membership proportions to either cluster.
At K=3, a novel red cluster is maximized in all Southern KS populations separating them from the
rest of African individuals, also consistent with the ASD-NJT. At K=4, the novel blue cluster
differentiates Central African RHG populations, and in particular certain Eastern RHG populations who
maximize their membership proportions to this cluster.
The new dark and darker gray clusters respectively at K=5 and K=6 differentiate Oceanian Papuan
individuals and South American Karitiana individuals from one another and from all other non-African
individuals, without affecting substantially genotype membership proportions from African individuals.
At K=7, the novel light blue cluster differentiates Eastern and Western RHG populations, as
observed in **FigureF3x**.
At K=8, the novel orange cluster differentiates mainly among Northern and Southern Khoe-San
populations, with the Central Khutse San showing substantial membership proportions to both the red
and the orange clusters, respectively maximized in Northern and Southern KS.
At K=9, an alternative clustering solution is shown where East Asian Dai, Oceania Papuan, and
South American Karitiana individuals are all clustering separately in three fully resolved clusters,
affecting the African clustering patterns where the differentiation previously observed at K=8 among
Northern and Southern Khoe-San individuals disappears to the benefit of a novel clustering solution
where Ba.Twa cluster separately from all RHG and from Eastern RHG with whom they shared closer
genotype membership proportions to the medium blue clusters at previous values of K in particular.
Finally, at K=10, these alternative clustering solutions are resolved into seven different virtual
genetic clusters maximized in different groups of African individuals. The light blue cluster is
maximized in certain Western RHG populations. The dark blue cluster is maximized in the Ba.Twa
Eastern RHG individuals only, while the medium blue cluster is maximized in the two other Eastern
RHG populations mostly. The red cluster is mainly represented in the Northern KS individuals, while
the orange cluster is maximized in Southern KS. Finally, the green cluster is maximized in all other
African populations, albeit note that it is not fully resolved at this value of K since no individual exhibits
100% genotype membership to this cluster.
Most importantly, note that numerous Central and Southern African individuals retain, from K=2
to 10, intermediate genotype membership proportions in between certain clusters. This should be
interpreted primarily as these individuals being at intermediate genetic distance between individuals
presenting 100% membership to either cluster, which is consistent with ASD-MDS and ASD-NJT
projections (**FigureF3x**). In turn, such intermediate distances may be either due to yet-unresolved
ADMIXTURE clustering, which may appear at higher values of K, or can be due to admixture having
occurred between ancestors respective to each cluster, the latter interpretation being likely but not
formally tested by the ADMIXTURE method. The possible occurrence of gene-flow events across pairs
of lineages and their influence on tree-topologies are incorporated explicitly in our Approximate
Bayesian Computation inferences detailed below (Tavaré *et al.*, 1997; Pritchard *et al.*, 1999; Beaumont,
Zhang and Balding, 2002; Blum and François, 2010; Csilléry, François and Blum, 2012; Pudlo *et al.*,
2016; Raynal *et al.*, 2019).
447
Altogether, our descriptive results highlight the vast genetic diversity and differentiation of African
populations, maximized in certain groups of Central and Southern African populations, as previously
reported (e.g. (Meyer *et al.*, 2012; Rasmussen *et al.*, 2014; Mallick *et al.*, 2016; Choudhury *et al.*, 2017;
Fan *et al.*, 2023; Ragsdale *et al.*, 2023; Fortes-Lima *et al.*, 2024)). Finally, and most importantly, our
results highlight in particular the vast genetic diversity and differentiation of populations at a very local

scale in Africa, including among immediate neighbors, not trivially correlated with geographical
distances among populations, linguistic classification, nor mode of subsistence. In fact, these results
anticipate that historical and demographic inferences in Africa may substantially differ when
considering different population samples across the continent and even at a local scale, as well as
advocate for extreme caution when grouping individuals and populations samples into larger categories.
458

2. Demographic and migration histories in Central and Southern African

2.A. Summary of the ABC inference design

We aimed at reconstructing the demographic and migration history that produced extant genomic
patterns observed within and among Central and Southern African populations (**FigureF2x**, **FigureF3x**,
FigureF4x), with machine-learning ABC scenario-choices followed by posterior-parameter
estimations. To do so, we considered eight competing possible tree-topologies among five extant
Northern and Southern Khoe-San, Western and Eastern Rainforest Hunter-Gatherer, and Rainforest
Hunter-Gatherer neighboring populations (**FigureF5x**). Furthermore, for each topology, we considered
two possible gene-flow processes among recent and ancient genetic lineages: instantaneous asymmetric
gene-flow corresponding to instantaneous unidirectional introgression events between pairs of lineages;
and recurring asymmetric gene-flow corresponding to unidirectional recurring migrations among pairs
of lineages (scenarios "i" and "r" respectively, **FigureF5x**). Finally, we considered, for each topology
and each gene-flow process, three nested gene-flow intensities for each event among pairs of lineages
separately: no to very high gene-flow rates (scenarios "i1" and "r1"); no to moderate rates (scenarios
"i2" and "r2"); and no to limited rates (scenarios "i3" and "r3").
This led to 8x2x3=48 competing scenarios in total which can be grouped in different ways to
address formally two nested major questions:
476
477 i) which ancestral lineages to extant populations diverged first from the others and when did they do so,
478 when considering complex gene-flow events among recent and ancient lineages?
479
480 ii) did gene-flow events occur recurrently or more instantaneously among pairs of lineages during the
481 evolutionary history of Central and Southern African populations? When did these events occur? How
482 intense were they?
483
Importantly, we aimed at considering the vast genetic diversity and differentiation within and
among populations and groups of populations at a local and regional scale (**FigureF2x**, **FigureF3x**,
FigureF4x, and (Tishkoff *et al.*, 2009; Skoglund and Mathieson, 2018; Pfennig *et al.*, 2023)).
Therefore, we replicated the same ABC scenario-choice and posterior-parameter estimation procedures
considering, in turn, 54 different combinations of five sampled populations of five individuals' whole
autosomal genomes each (**FigureF5x**).
We thus conducted a total of 240,000 coalescent simulations under these 48 competing scenarios
(5000 simulations per scenario), by drawing parameter values from prior distributions set by the user
(**FigureF5x**, **TableT2x**). For each simulation, we then calculated a vector of 337 summary-statistics
(**TableT3x**), within and among the five simulated populations; each vector of summary-statistics thus
corresponds to a vector of parameter values drawn randomly from prior distributions and used for one
simulation. We then deployed Random Forest ABC procedures (Pudlo *et al.*, 2016; Estoup *et al.*, 2018),
to identify the winning scenario or group of scenarios for each 54 combinations of five sampled "real"
populations separately. Finally, under the winning scenario, we produced 100,000 simulations,
computed 202 summary-statistics for each simulation (**TableT3x**), and performed Neural Network

ABC posterior parameter joint-estimation (Blum and François, 2010; Csilléry, François and Blum,
2012), providing posterior-parameter distributions most likely to have produced observed genomic data,
for each scenario-parameter.
502

FigureF5x: 48 competing scenarios for the history of Central and Southern African populations.
(**A**) Eight competing topologies for the demographic history of five Central and Southern African extant lineages. The Northern
Khoe-San lineage (nKS) is represented by either the Ju|'hoansi or the !Xun population. The Southern Khoe-San lineage (sKS)
is represented by either the Karretjie People or the Nama population. The Rainforest Hunter-Gatherer Neighbors lineage
(RHGn) is represented either by the Western Congo Basin Nzime or Ngumba populations, or the Eastern Ba.Konjo or Ba.Kiga
populations. The Western Rainforest Hunter-Gatherer lineage (wRHG) is represented by either the Baka, the Ba.Kola, or the
Bi.Aka_Mbati population. The Eastern Rainforest Hunter-Gatherer lineage (eRHG) is represented by either the Nsua, the
Ba.Twa, or the Mbuti population. As indicated in legend in the bottom right corner of the panel, the eight topologies can be
grouped according to i) which ancestral lineage diverged first from the two others, or ii) whether the nKS and sKS lineages
split earlier or later than the split of the wRHG and eRHG lineages. For each topology, we consider possible changes in
lineages' constant effective population size, *Ne*, after each divergence event, *t*. For each topology, we considered possible
gene-flow events among ancestral lineages and among recent lineages, represented as horizontal uni-directional arrows. (**B**)
Example for a given topology (1a), of the fact that each one of the eight topologies are considered under two alternative
competing "instantaneous" (Scenario i-1a) or "recurring" (Scenario r-1a) gene-flow processes. Instantaneous gene-flow
processes consider that genes can be exchanged among pairs of ancestral or recent lineages at a single parameterized time, *tad*,
with independent parameters of gene-flow intensity, *m*, from lineage "A" to "B" and from lineage "B" to "A". Recurring gene-
flow processes consider instead that gene-flow occur at each generation between pairs of lineages with a constant independent
rate, *m*, from lineage "A" to "B" and from lineage "B" to "A". Note that for all eight topologies and for both instantaneous
and recurring gene-flow processes, we parameterized separately the gene-flow from lineage "A" into lineage "B", and the
gene-flow from lineage "B" into "A", in order to allow for possibly asymmetrical gene-flow events among pairs of lineages.
For both instantaneous and recurring gene-flow processes, we considered three nested intensities of gene-flow parameters
indicated at the bottom of (**B**). Scenarios i1 or r1 consider that the gene-flow parameters *m* in each topology are independently
drawn in Uniform distributions between 0 and 1 or 0 and 0.05, respectively, thus allowing for the possibility of no to high
gene-flows from one lineage to the other. Scenarios i2 or r2 consider that the gene-flow parameters *m* in each topology are
independently drawn in Uniform distributions between 0 and 0.125 or 0 and 0.0125, respectively, thus allowing only for the
possibility of no to intermediate gene-flow from one lineage to the other. Finally, scenarios i3 or r3 consider that the gene-
flow parameters *m* in each topology are independently drawn in Uniform distributions between 0 and 0.01 or 0 and 0.001,
respectively, thus only allowing for the possibility of no to reduced gene-flow from one lineage to the other. Altogether we
thus considered 8x2x3=48 competing scenarios for the demographic and migration history of Central and Southern African
populations inferred with machine-learning Approximate Bayesian Computation. We conducted Random-Forest ABC
scenario choice procedures among groups of these 48 scenarios in turn for 54 different sets of five observed populations each,
represented as grey lines between population names, each population comprising five whole-genome sequenced individuals.
Sampled-population names considered in the 54 combinations are given below each tree-leave, and combinations are limited
to those pairing one RHGn with the specific eastern or western RHG population with whom they share complex socio-
economic interactions, as indicated in **TableT1x**. See the detailed description of scenarios and their grouping, their parameters
and their respective prior distributions used for ABC inference in **Material and Methods** and **TableT2x**.

541
542

543 **FigureF5x**

545

A

Scenario-groups Legend

Ancient topologies

group 1: (AKS,(ARHG,RHGn))

group 2: (ARHG,(AKS,RHGn))

group 3: (RHGn,(AKS,ARHG))

Internal divergences order

group A: $t_{KS} < t_{RHG}$

group B: $t_{RHG} < t_{KS}$

B

547   ---

548   <u>**TableT2x**</u>**: 48 competing scenarios parameters' description and prior distributions**

549   Codes for scenarios topologies and gene-flow processes and all scenario parameters are detailed in **FigureF5x** and in **Material**

550   **and Methods**.

Parameter	Description	Prior	Scenario topology	Scenario gene-flow	Condition
t_{RHG}	Time of divergence between wRHG and eRHG	Uniform[1,15000]	1a-2a-2c-3a 1b-1c-2b-3b	i1-i2-i3 and r1-r2-r3	$t_{RHG}>t_{KS}$ $t_{KS}>t_{RHG}$
t_{KS}	Time of divergence between nKS and sKS	Uniform[1,15000]	1a-2a-2c-3a 1b-1c-2b-3b	i1-i2-i3 and r1-r2-r3	$t_{RHG}>t_{KS}$ $t_{KS}>t_{RHG}$
$t_{RHG-RHGn}$	Time of divergence between the RHGn lineage and the lineage ancestral to all RHG lineages	Uniform[1,15000]	1a-1b-1c	i1-i2-i3 and r1-r2-r3	$t_{RHG-RHGn}>t_{RHG}$
$t_{KS-RHGn}$	Time of divergence between the RHGn lineage and the lineage ancestral to all KS lineages	Uniform[1,15000]	2a-2b-2c	i1-i2-i3 and r1-r2-r3	$t_{KS-RHGn}>t_{KS}$
t_{KS-RHG}	Time of divergence between the KS and RHG ancestral lineages	Uniform[1,15000]	3a-3b	i1-i2-i3 and r1-r2-r3	$t_{KS-RHG}>t_{KS}$ and $t_{KS-RHG}>t_{RHG}$
$t_{KS-RHG-RHGn}$	Time of the divergence event ancestral to all lineages	Uniform[1,15000]	all 8 topol.	i1-i2-i3 and r1-r2-r3	$t_{KS-RHG-RHGn}>$ all other times
$tad_{nKS-sKS}$	Time for both the unidirectional instantaneous gene-flow events between nKS and sKS	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{nKS-sKS}$
m(nKS-sKS) or m(sKS-nKS)	Instantaneous gene-flow intensity from sKs into nKS, or from nKS into sKS independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(nKS-sKS) or m(sKS-nKS)	Recurring gene-flow intensity from sKS into nKS, or from nKS into sKs independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{nKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between nKS and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{nKS-RHGn}$ and $t_{RHG-RHGn}>tad_{nKS-RHGn}$
m(nKS-RHGn) or m(RHGn-nKS)	Instantaneous gene-flow intensity from RHGn into nKS, or from nKS into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(nKS-RHGn) or m(RHGn-nKS)	Recurring gene-flow intensity from RHGn into nKS, or from nKS into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{sKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between sKS and RHGn lineages	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{KS}>tad_{sKS-RHGn}$ and $t_{RHG-RHGn}>tad_{sKS-RHGn}$
m(sKS-RHGn) or m(RHGn-sKS)	Instantaneous gene-flow intensity from RHGn into sKS, or from sKS into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(sKS-RHGn) or m(RHGn-sKS)	Recurring gene-flow intensity from RHGn into sKS, or from sKS into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{wRHG-eRHG}$	Time for both the unidirectional instantaneous gene-flow events between wRHG and eRHG	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{wRHG-eRHG}$
m(wRHG-eRHG) or m(eRHG-wRHG)	Instantaneous gene-flow intensity from eRHG into wRHG, or from wRHG into eRHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(wRHG-eRHG) or m(eRHG-wRHG)	Recurring gene-flow intensity from eRHG into wRHG, or from wRHG into eRHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{wRHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between wRHG and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{wRHG-RHGn}$ and $t_{KS-RHG}>tad_{nwRHG-RHGn}$
m(wRHG-RHGn) or m(RHGn-wRHG)	Instantaneous gene-flow intensity from RHGn into wRHG, or from wRHG into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(wRHG-RHGn) or m(RHGn-wRHG)	Recurring gene-flow intensity from RHGn into wRHG, or from wRHG into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{eRHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between eRHG and RHGn	Uniform[1,15000]	all 8 topol.	i1-i2-i3	$t_{RHG}>tad_{eRHG-RHGn}$ and $t_{KS-RHG}>tad_{eRHG-RHGn}$
m(eRHG-RHGn) or m(RHGn-eRHG)	Instantaneous gene-flow intensity from RHGn into eRHG, or from eRHG into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol.	i1 i2 i3	x
m(eRHG-RHGn) or m(RHGn-eRHG)	Recurring gene-flow intensity from RHGn into eRHG, or from eRHG into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol.	r1 r2 r3	x
$tad_{ARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the RHGn and the lineage ancestral to wRHG and eRHG lineages	Uniform[1,15000]	all 8 topol. except 2c	i1-i2-i3	$tad_{RHGn-ARHG}>t_{RHG}$
m(RHGn-ARHG) or m(ARHG-RHGn)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into RHGn, or from RHGn into ARHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol. except 2c	i1 i2 i3	x
m(RHGn-ARHG) or m(ARHG-RHGn)	Recurring gene-flow intensity from the ancestral lineage ARHG into RHGn, or from RHGn into ARHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol. except 2c	r1 r2 r3	x
$tad_{AKS-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to nKS and sKS, and the RHGn lineage	Uniform[1,15000]	all 8 topol. except 1c	i1-i2-i3	$tad_{AKS-RHGn}>t_{KS}$
m(RHGn-AKS) or m(AKS-RHGn)	Instantaneous gene-flow intensity from the ancestral lineage AKS into RHGn, or from RHGn into AKS independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol. except 1c	i1 i2 i3	x
m(RHGn-AKS) or m(AKS-RHGn)	Recurring gene-flow intensity from the ancestral lineage AKS into RHGn, or from RHGn into AKS independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol. except 1c	r1 r2 r3	x
$tad_{AKS-ARHG}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to nKS and sKS, and the lineage ancestral to wRHG and eRHG lineages	Uniform[1,15000]	all 8 topol. except 1c-2c	i1-i2-i3	$tad_{AKS-ARHG}>t_{KS}$ and $tad_{AKS-ARHG}>t_{RHG}$
m(AKS-ARHG) or m(ARHG-AKS)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into the ancestral lineage AKS, or from AKS into ARHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	all 8 topol. except 1c-2c	i1 i2 i3	x

m(AKS-ARHG) or m(ARHG-AKS)	Recurring gene-flow intensity from the ancestral lineage ARHG into the ancestral lineage AKS, or from AKS into ARHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	all 8 topol. except 1c-2c	r1 r2 r3	x
tad$_{nKS-ARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the nKS and the lineage ancestral to ARHG and RHGn lineages	Uniform[1,15000]	1c	i1-i2-i3	t$_{KS}$>tad$_{nKS-ARHG-RHGn}$ and tad$_{nKS-ARHG-RHGn}$>t$_{RHG-RHGn}$
m(nKS-ARHG-RHGn) or m(ARHG-RHGn-nKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into nKS, or from nKS into ARHG-RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	1c	i1 i2 i3	x
m(nKS-ARHG-RHGn) or m(ARHG-RHGn-nKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into nKS, or from nKS into ARHG-RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	1c	r1 r2 r3	x
tad$_{sKS-ARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the sKS and the lineage ancestral to ARHG and RHGn lineages	Uniform[1,15000]	1c	i1-i2-i3	t$_{KS}$>tad$_{sKS-ARHG-RHGn}$ and tad$_{sKS-ARHG-RHGn}$>t$_{RHG-RHGn}$
m(sKS-ARHG-RHGn) or m(ARHG-RHGn-sKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into sKS, or from sKS into ARHG-RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	1c	i1 i2 i3	x
m(sKS-ARHG-RHGn) or m(ARHG-RHGn-sKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages RHGn and ARHG into sKS, or from sKS into ARHG-RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	1c	r1 r2 r3	x
tad$_{AKS-RHGn-wRHG}$	Time for both the unidirectional instantaneous gene-flow events between the wRHG and the lineage ancestral to AKS and RHGn lineages	Uniform[1,15000]	2c	i1-i2-i3	t$_{RHG}$>tad$_{AKS-RHGn-wRHG}$ and tad$_{AKS-RHGn-wRHG}$>t$_{KS-RHGn}$
m(wRHG-AKS-RHGn) or m(AKS-RHGn--wRHG)	Instantaneous gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into wRHG, or from wRHG into AKS-RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	2c	i1 i2 i3	x
m(wRHG-AKS-RHGn) or m(AKS-RHGn--wRHG)	Recurring gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into wRHG, or from wRHG into AKS-RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	2c	r1 r2 r3	x
tad$_{AKS-RHGn-eRHG}$	Time for both the unidirectional instantaneous gene-flow events between the eRHG and the lineage ancestral to AKS and RHGn lineages	Uniform[1,15000]	2c	i1-i2-i3	t$_{RHG}$>tad$_{AKS-RHGn-eRHG}$ and tad$_{AKS-RHGn-eRHG}$>t$_{KS-RHGn}$
m(eRHG-AKS-RHGn) or m(AKS-RHGn--eRHG)	Instantaneous gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into eRHG, or from eRHG into AKS-RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	2c	i1 i2 i3	x
m(eRHG-AKS-RHGn) or m(AKS-RHGn--eRHG)	Recurring gene-flow intensity from the lineage AKS-RHGn ancestral to lineages AKS and RHGn into eRHG, or from eRHG into AKS-RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	2c	r1 r2 r3	x
tad$_{AKS-ARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and the lineage ancestral to RHG and RHGn lineages	Uniform[1,15000]	1a-1b-1c	i1-i2-i3	tad$_{AKS-ARHG-RHGn}$>t$_{RHG-RHGn}$ and tad$_{AKS-ARHG-RHGn}$>t$_{KS}$ and t$_{KS-RHG-RHGn}$>tad$_{AKS-ARHG-RHGn}$
m(AKS-ARHG-RHGn) or m(ARHG-RHGn-AKS)	Instantaneous gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages ARHG and RHGn into the ancestral lineage AKS, or from AKS into ARHG-RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	1a-1b-1c	i1 i2 i3	x
m(AKS-ARHG-RHGn) or m(ARHG-RHGn-AKS)	Recurring gene-flow intensity from the lineage ARHG-RHGn ancestral to lineages ARHG and RHGn into the ancestral lineage AKS, or from AKS into ARHG-RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	1a-1b-1c	r1 r2 r3	x
tad$_{AKSRHGn-ARHG}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and RHGn, and the lineage ancestral to RHG lineages	Uniform[1,15000]	2a-2b-2c	i1-i2-i3	tad$_{AKSRHGn-ARHG}$>t$_{KS-RHGn}$ and tad$_{AKSRHGn-ARHG}$>t$_{RHG}$ and t$_{KS-RHG-RHGn}$>tad$_{AKSRHGn-ARHG}$
m(AKSRHGn-ARHG) or m(ARHG-AKSRHGn)	Instantaneous gene-flow intensity from the ancestral lineage ARHG into the lineage AKSRHGn ancestral to AKS and RHGn, or from AKSRHGn into ARHG independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	2a-2b-2c	i1 i2 i3	x
m(AKSRHGn-ARHG) or m(ARHG-AKSRHGn)	Recurring gene-flow intensity from the ancestral lineage ARHG into the lineage AKSRHGn ancestral to AKS and RHGn, or from AKSRHGn into ARHG independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	2a-2b-2c	r1 r2 r3	x
tad$_{AKSARHG-RHGn}$	Time for both the unidirectional instantaneous gene-flow events between the lineage ancestral to KS and RHG, and the RHGn lineage	Uniform[1,15000]	3a-3b	i1-i2-i3	tad$_{AKSARHG-RHGn}$>t$_{KS-RHG}$ and t$_{KS-RHG-RHGn}$>tad$_{AKSARHG-RHGn}$
m(AKSARHG-RHGn) or m(RHGn-AKSARHG)	Instantaneous gene-flow intensity from the lineage RHGn into the lineage AKSRHG ancestral to AKS and ARHG, or from AKSRHG into RHGn independantly	Uniform[0,1] Uniform[0,0.125] Uniform[0,0.01]	3a-3b	i1 i2 i3	x
m(AKSARHG-RHGn) or m(RHGn-AKSARHG)	Recurring gene-flow intensity from the lineage RHGn into the lineage AKSRHG ancestral to AKS and ARHG, or from AKSRHG into RHGn independantly	Uniform[0,0.05] Uniform[0,0.0125] Uniform[0,0.001]	3a-3b	r1 r2 r3	x
N$_{nKS}$	Effective population size for the nKS lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{sKS}$	Effective population size for the sKS lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{AKS}$	Effective population size for the lineage ancestral to the nKS and the sKS		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{RHGn}$	Effective population size for the RHGn lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{wRHG}$	Effective population size for the wRHG lineage	Uniform[10,100000]	all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{eRHG}$	Effective population size for the eRHG lineage		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{ARHG}$	Effective population size for the lineage ancestral to the wRHG and the eRHG		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
N$_{ARHG-RHGn}$	Effective population size for the lineage ancestral to the RHGn and the ARHG		1a-1b-1c	i1-i2-i3 and r1-r2-r3	x
N$_{AKS-RHGn}$	Effective population size for the lineage ancestral to the RHGn and the AKS		2a-2b-2c	i1-i2-i3 and r1-r2-r3	x

$N_{AKS-ARHG}$	Effective population size for the lineage ancestral to the AKS and the ARHG		3a-3b	i1-i2-i3 and r1-r2-r3	x
N_A	Effective population size for the lineage ancestral to all lineages		all 8 topol.	i1-i2-i3 and r1-r2-r3	x
Mutation rate	Fixed mutation rate	$1.25*10^{-8}$	all 8 topol.	i1-i2-i3 and r1-r2-r3	x
Recombination rate	Fixed recombination rate	$1*10^{-8}$	all 8 topol.	i1-i2-i3 and r1-r2-r3	x
Transition to transversion ratio	Fixed transition to transversion rate	0.33	all 8 topol.	i1-i2-i3 and r1-r2-r3	x

551 --

554 ---

TableT3x: Summary-statistics used in Random-Forest ABC scenario-choice and Neural-Network ABC posterior-parameter estimation.

Name of statistic (or group of statistics)	What was computed	Number of values	RF-ABC scenario-choice	NN-ABC posterior-parameter estimation
Total number of biallelic sites among the five populations	sum	1	yes	yes
Total number of multiallelic sites among the five populations	sum	1	yes	yes
ASD distance within population for each five population separately	average and variance	10	yes	no
First five dimensions of the projection of the ASD matrix within population for each five population separately	average and variance	50	yes	yes
Proportion of biallelic sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Proportion of homozygous sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Proportion of homozygous ancestral sites within each five population separately	sum in a population divided by total number of homozygous sites in the population (no missing data)	5	yes	no
Proportion of private biallelic sites within each five population separately	sum in a population divided by total number of biallelic sites in the dataset (no missing data)	5	yes	yes
Frequency of the minor allele of the private biallelic sites within each five population separately	average and variance	10	yes	no
Expected heterozygosity at a segregating site within each five population separately	average and variance	10	yes	no
Tajima's D within each five population separately	average and variance for chromosomes with at least one variant	10	yes	yes
Unfolded site frequency spectrum (SFS) within each five population separately	proportion for each of 9 classes	45	yes	no
Watterson's theta within each five population separately	sum of biallelic sites divided by the harmonic number of the sample size (n=10)	5	yes	no
Total number of runs of homozygosity (ROH) within each five population separately	sum	5	yes	no
Total number of ROH by length class (<200 kbp, 200-500 kbp, >500 kbp) within each five population separately	sum	15	yes	yes
ROH length for each length class within each five population separately	average	15	yes	no
Total ROH length within each five population separately	average and variance	10	yes	no
ASD distance between each pair of five populations	average and variance	20	yes	no
First five dimensions of the projection of the ASD matrix between each pair of five populations	average and variance	100	yes	yes
Pairwise FST between each pair of five population	Pairwise FST (Weir and Cockerham 1984)	10	yes	yes
	TOTAL number of statistics used in ABC inference		337	202

557 ---

2.B. Are we able to mimic observed data with simulations under the 48 competing scenarios?

In order to conduct ABC inferences, we first checked whether we were able to simulate data, under the 48 competing scenarios, for which vectors of summary statistics could mimic those obtained separately from the 54 different combinations of five sampled populations each.

We first conducted goodness-of-fit permutation tests, separately for the 54 combinations of five populations, and found that the observed vectors of 337 summary statistics were never significantly different (54 goodness-of-fit permutation p-values > 0.56), from the 240,000 vectors of 337 summary statistics computed from simulations under the 48 competing scenarios. Second, we computed a two-dimensional PCA on the vectors of summary-statistics obtained from simulations on which we projected, separately, the 54 vectors of these summary-statistics computed on observed data, and found that observations each fell well within the space of simulations (**SupplementaryFigureSF1x**).

Both results show that we empirically were able to simulate data for which summary-statistics were reasonably close to the observed ones, at least in parts of the parameter space used for simulations under the 48 competing scenarios, for each one of the 54 separate combinations of five sampled populations each. We could therefore reasonably proceed with Random Forest ABC scenario-choice and Neural Network ABC posterior-parameter estimation procedures.

2.C. Which historic scenario best explains genomic patterns in extant African populations?

2.C.1. Instantaneous or recurring gene-flows of which intensity among African lineages?

Panel A in **FigureF6x** shows that gene-flow processes among recent or ancient genetic lineages most likely occurred during very limited "instantaneous" periods of time, rather than recurrently, throughout the entire evolutionary history of Central and Southern African populations. Indeed, for each one of the 54 different combinations of five sampled populations, the group of 24 scenarios considering instantaneous gene-flow processes provides vectors of summary statistics systematically closest to the observed ones, whichever the tree-topology or intensity of the gene-flow considered.

Furthermore, we find that we imperatively need these gene-flow processes to be potentially high in order to best explain the observed data for each one of the 54 combinations of population samples, whichever the tree-topology and gene-flow process (**Panel B** in **FigureF6x**), whichever the tree-topology but considering gene-flow processes separately (**Panel C** in **FigureF6x**), or even considering 24 competing scenarios under instantaneous gene-flow processes only, whichever the tree-topology (**SupplementaryFigureSF6xPart1**). Finally, note that when considering all 48 competing-scenarios separately, a very challenging task *a priori* given the number of competing scenarios and the high level of nestedness among scenarios, the winning scenarios predicted by the RF-ABC procedure is, among the 54 tests conducted, systematically found for scenarios considering an instantaneous gene-flow process allowing for possibly highly intense gene-flow rates (**SupplementaryFigureSF6xPart2-3**).

Overall, for all the above analyses, the *a priori* discriminatory powers of the RF, estimated as cross-validation procedures based on simulations used as pseudo-observed data, were overall good across groups of scenarios, despite increased confusion among scenarios based on gene-flow intensity within groups of instantaneous or recurring processes, as expected due to increased scenario-nestedness among classes of no-to-low, no-to-moderate, or no-to-high intensities (**SupplementaryFigureSF6xPart0-PanelA-B-C**, **SupplementaryFigureSF6xPart1-2-3**).

FigureF6x: Random-Forest ABC scenario-choices among 48 competing scenarios.

Random Forest Approximate Bayesian Computation scenario-choice results (Pudlo *et al.*, 2016; Estoup *et al.*, 2018), are
conducted, for each analysis in each panel, separately for 54 different combinations of five Central and Southern African
sampled populations each. Posterior proportions of votes obtained for each 54 sampled-populations combinations, and for RF-
ABC analysis in each panel respectively, are provided as box-plots indicating the median in between the first and third quartile
of the box-limits, whiskers extending to data points no more than 1.5 times the interquartile range of the distribution, and
empty circles for all more extreme points beyond this limit, if any. In each RF-ABC analysis in each panel, proportion of votes
(indicated in the y-axis) are obtained using 1,000 decision trees in the random forest for each group of competing scenarios
(indicated in the x-axis table and labels), using 5,000 simulations per scenario and 337 summary-statistics (see **Material and**
Methods and **TableT3x** for details). Detailed competing-scenarios grouped in each group are indicated in the two bottom lines
of the x-axis table, with topology and gene-flow processes codes provided in **FigureF5x** and explained in detail in **Material**
and Methods. The top-line of the x-axis table indicates, for each group of competing scenarios in each analysis, the number
of times the corresponding group of scenarios was predicted as the winning one among the 54 different combinations of five
sampled-populations. (**A**) Scenario-choice results for the two groups of instantaneous or recurring gene-flow processes (24
scenarios and 120,000 simulations in each two competing groups), all gene-flow intensities and all topologies "being equal".
(**B**) Scenario-choice results for the three groups of intensities for gene-flow processes (16 scenarios and 80,000 simulations in
each three competing groups), all topologies and both instantaneous and recurring gene-flow processes "being equal". (**C**)
Scenario-choice results for the six groups of gene-flow processes and intensities separately (8 scenarios and 40,000 simulations
in each six competing groups), all topologies "being equal". (**D**) Scenario-choice results for the three groups of scenario
topologies differing in the order of ancient lineages divergence (differing number of scenarios in each three competing groups
are evened by randomly sampling the same number of simulations, equal to 60,000, in each group, see **Material and**
Methods), all gene-flow processes and intensities "being equal". (**E**) Scenario-choice results for the same test as (**D**) restricted
to the 24 competing scenarios considering instantaneous gene-flow processes only (differing number of scenarios in each three
competing groups are evened by randomly sampling the same number of simulations, equal to 30,000, in each group, see
Material and Methods), all gene-flow intensities "being equal". (**F**) Scenario-choice results for the two groups of scenario
topologies differing in the relative order of divergences between Northern and Southern KS, and Western and Eastern RHG
lineages, respectively (24 competing scenarios and 120,000 simulations in each two groups), all gene-flow processes and
intensities "being equal". (**G**) Scenario-choice results for the same test as (**F**) restricted to the 24 competing scenarios
considering instantaneous gene-flow processes only (12 competing scenarios and 60,000 simulations in each two groups), all
gene-flow intensities "being equal".

FigureF6x

2.C.2. Ancient tree-topology among African lineages?
FigureF6x shows that, overall, the RF-ABC scenario-choice procedures vote in favor of an ancient
tree-topology where the ancestral Khoe-San lineage diverged first followed by the divergence between
the ancient Rainforest Hunter-Gatherer lineage and that of their neighbors (Scenarios tree-topologies
1a-1b-1c, **FigureF5x**), in a large majority (45/54) of the 54 combinations of observed populations tested
here, whichever the gene-flow process and intensity (**FigureF6x-PanelD**), or when considering
instantaneous gene-flow processes, only, whichever their intensities (**FigureF6x-PanelE**). Note that
the tree-topologies where the RHGn lineage diverge first from the lineage ancestral to that of the RHG
and the KS (Scenarios tree-topologies 3a-3b, **FigureF5x**), is winning in a minority of the tests (9 out of
54 combinations of sampled populations), whichever the gene-flow process and/or intensity.
Interestingly, we found that tree-topologies where the ancient lineage for RHG populations diverges
first from the two others (Scenarios tree-topologies 2a-2b-2c, **FigureF5x**), is never favored in our
analyses, whichever the gene-flow process and/or intensity.
657
2.C.3. Recent tree-topology among African lineages?
Finally, **FigureF6x** shows that Northern and Southern Khoe-San extant lineages likely diverged from
one-another before the divergence between Eastern and Western Rainforest Hunter-Gatherer lineages
(Scenarios tree-topologies 1b-1c-2b-3b, **FigureF5x**), whichever the ancient tree-topology and gene-
flow process and intensity (**FigureF6x-PanelF**), and whichever the ancient tree-topology for
instantaneous gene-flow processes only, all classes of intensities "being equal" (**FigureF6x-PanelG**).
Interestingly, while the "b" group of tree-topologies wins for 38 out of 54 combinations of sampled
populations, this majority of RF-ABC votes is larger (47/54) when considering only the group of 24
competing scenarios for which gene-flows are instantaneous rather than recurring.
667
2.C.4. Conclusion: intersecting scenario-choice results for the history of Africa.
Altogether, when intersecting RF-ABC scenario-choice inferences in groups of gene-flow processes
and tree-topologies (**FigureF6x**), with inferences considering all competing-scenarios separately
(**SupplementaryFigureSF6x-Part2-3**), and among 54 combinations of five sampled populations, we
find that Scenario i1-1b (**FigureF5x**), systematically wins in all conformations of scenario-choice tests
for 25 combinations of populations (**SupplementaryTableST3x**), while the second best scenario
(Scenario i1-3b) wins consistently across tests for only four combinations of population out of 54.
Therefore, our results point to a scenario for the history of Central and Southern African
populations where the ancestral KS lineage diverged first, followed by the divergence between the
ancestral RHG and the RHGn lineages, with a divergence between the extant Northern and Southern
KS lineages occurring independently before that of the extant Eastern and Western RHG lineages. Most
importantly, the winning scenario necessarily involves gene-flow events among all pairs of lineages
throughout history which occurred during relatively short "instantaneous" periods of time, rather than
recurrently over larger periods of time. Furthermore, these instantaneous gene-flow events must each
have been possibly of high intensity, rather than all limited to moderate or low intensities, and possibly
asymmetrical between pairs of lineages, to best explain the observed data. Finally, our results also
highlight that considering different combinations of extant populations separately in the analyses may
provide contrasted results where alternative evolutionary scenarios may best explain observations.
For conservativeness and to further consider the genetic diversity of extant Central and Southern
African populations at a local or regional scale, we will henceforth conduct all Neural-Network ABC
posterior-parameter estimations separately for the 25 combinations of sampled populations
systematically providing the Scenario i1-1b as the winner.
690
691

2.D. Which scenario-parameters best explain genomic patterns in extant African populations?

2.D.1. Divergence times.

Divergence times are most often well estimated by our Neural Network ABC posterior parameter inferences (**FigureF7x** and **SupplementaryTableST4x**), with relatively narrow 90% Credibility Intervals (CI) and posterior distributions very often largely departing from the priors. Interestingly, we find relatively consistent posterior estimates across the 25 different combinations of five populations for, separately, the Eastern and Western RHG divergence between 383 generations before present (gbp) (mode point estimate, 90%CI=[220-934]) and 2174 gbp (90%CI=[1,226-4,373], **SupplementaryTableST4x**), and for the Northern and Southern KS divergence between 1,486 gbp (mode point estimate, 90%CI=[844-3,798]) and 6,201 gbp (90%CI=[4,465-8,126]). Combining the posterior distributions for these two parameters (**FigureF7x-PanelA-B**, **TableT4x**), respectively, provides, synthetically, a modal point-estimate divergence time between eRHG and wRHG of 892 gbp (90%CI =[422-2,531]), and one between nKS and sKS, largely more ancient, having occurred some 2,623 gbp (90%CI=[1,481-5,763]).

In a more remote past, the divergence time between the lineage ancestral to all RHG populations and the RHG neighbors' lineage is also most often well estimated, based on CI-width and departure from prior distributions, for each 25 combinations of populations respectively, albeit results are more variable across sets of population combinations than for more recent divergence times (**FigureF7x-PanelC** and **SupplementaryTableST4x**). Indeed, ancient RHG and RHGn lineages diverged between 3,904 gbp (mode point estimate, 90%CI=[2,828-6,636]) and 10,727 gbp (90%CI =[9,268-12,736], **SupplementaryTableST4x**) across population combinations, and synthetically combining results together 6,726 gbp (90%CI=[3,876-10,748], **FigureF7x-PanelC** and **TableT4x**).

Finally, we also find relatively variable posterior estimates for the most ancient divergence time in our tree-topology, between the ancestral lineages to all extant KS lineages and the ancestral lineages to all RHG and the RHG neighbors; with, again, posterior-parameter distributions across 25 combinations of sampled populations very often satisfactorily estimated, with relatively reduced 90%CI and substantial departure from the priors (**FigureF7x-PanelD**, **SupplementaryTableT4x**). Overall, we find that the original most ancient divergences in our tree-topology occurred some 12,117 gbp (90%CI=[8,875-14,538]), when all results are combined together synthetically (**FigureF7x-PanelD** and **TableT4x**).

Altogether, the relatively large variation in divergence times posterior estimates further back in time across combinations of sampled populations empirically highlights that demographic and historical reconstructions largely depend on the specific samples considered when investigating highly differentiated Sub-Saharan populations. Our results thus explicitly advocate for caution when summarizing results obtained across population sets and further likely explain the, sometimes, apparent discrepancies that arise across studies considering different sample sets and/or artificially merging several samples from relatively differentiated populations (e.g. (Fan *et al.*, 2023; Ragsdale *et al.*, 2023)).

<u>FigureF7x</u>: Neural Network ABC posterior parameter distributions of divergence times.

Neural Network Approximate Bayesian Computation posterior parameter joint estimations (Blum and François, 2010;
Csilléry, François and Blum, 2012), of topological divergence-times (in generations before present) for 25 sets of five Central
and Southern African populations for which the winning scenario identified by RF-ABC was Scenario i1-1b (**FigureF5x**). NN
ABC posterior parameter estimation procedures were conducted using 100,000 simulations under Scenario i1-1b, each
simulation corresponding to a single vector of parameter values drawn randomly from prior distributions provided in
TableT2x. We considered 42 neurons in the hidden layer of the NN and a tolerance level of 0.01, corresponding to the 1,000
simulations providing summary-statistics closest to the observed one. NN posterior estimates are based on the logit
transformation of parameter values using an Epanechnikov kernel between the corresponding parameter's prior bounds (see
Material and Methods and **TableT2x**). Posterior parameter densities are represented with solid blue lines. 50% Credibility
Intervals are represented as the light blue area under the density. The median and mode values are represented as a solid and
dotted blue vertical line, respectively. Parameter prior distributions are represented as dotted gray lines. For all panels, the left
plots represent the NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern African populations
winning under Scenario i1-1b, separately (**SupplementaryTableST3x** and **SupplementaryTableST4x**). For all panels, the
right plots represent a single parameter posterior distribution obtained from combining the 25 posterior distributions together.
(**A**) Results for parameter t_{RHG} corresponding to the split time between the Western and Eastern Rainforest Hunter-Gatherer
(RHG) lineages (**FigureF5x**). (**B**) Results for parameter t_{KS} corresponding to the split time between Northern Khoe-San (nKS)
and Southern Khoe-San (sKS) lineages (**FigureF5x**). (**C**) Results for parameter $t_{RHG-RHGn}$ corresponding to the split time
between the Rainforest Hunter-Gatherer neighboring population lineage (RHGn) and the lineage ancestral to Western and
Eastern RHG (**FigureF5x**). (**D**) Results for parameter $t_{KS-RHG-RHGn}$ corresponding to the split time between the lineage ancestral
to KS populations and the lineage ancestral to all RHG and RHGn lineages; this event thus corresponding to the split time of
all lineages in the history of Central and Southern African populations (**FigureF5x**). Results for all left panels are summarized
in **TableT4x**.

759
760
761
762

TableT4x: NN-ABC posterior parameter estimation of all parameters in Scenario i1-1b for results from 25 sets of five populations each, combined altogether.

Entire posterior distributions for each 25 analyses separately, for which summary statistics are presented in **SupplementaryTableT2x**, are combined together before re-computing the Mode, Median, Mean, 50%CI and 90%CI presented in this table. Therefore, values in the table correspond to posterior densities plotted in the right panels of **FigureF7x**, **FigureF8x**, and **SupplementaryFigureF7xPart1-Part2**. Parameter definitions and priors are provided in **TableT2x** and represented graphically in the corresponding scenario panel of **FigureF5x**. Values in this table are used for the synthetic schematic representation of the demographic and migration history of Central and Southern African populations presented in **FigureF9x**. Values in italic are not satisfactorily departing from the priors for the results from 25 sets of five populations each when combined together. In the case of the parameters N_{AKS}, N_{ARHG}, and $N_{ARHG-RHGn}$, this is due to parameter un-identifiability for all 25 analyses separately as almost none of these posterior distributions depart from their respective priors (**SupplementaryFigureF7xPart1**). In the case of all the migration rates parameters, **SupplementaryFigureF7xPart2** clearly shows both an overall lack of identifiability of parameters in the majority of the 25 analyses, separately, and a large variability of posterior estimates among those sets of analyses for which posterior distributions satisfactorily depart from the priors.

Parameter	Mode	Median	Mean	50% CI	90% CI
N_{nKS}	**14184**	**23881**	**29908**	**14721-39894**	**7825-73423**
N_{sKS}	**11042**	**19672**	**26050**	**11059-35737**	**4623-68767**
N_{wRHG}	**9461**	**20586**	**26551**	**11506-36934**	**5074-67317**
N_{eRHG}	**12592**	**15735**	**19968**	**10059-24368**	**3993-53065**
N_{RHGn}	**18022**	**28739**	**33965**	**18076-45474**	**9150-77157**
N_{AKS}	*86572*	*61359*	*59978*	*41254-80294*	*19523-94743*
N_{ARHG}	*57397*	*55256*	*54989*	*36771-73803*	*16922-91847*
$N_{ARHG-RHGn}$	*46968*	*48438*	*48585*	*30042-66886*	*9894-88177*
N_A	**18507**	**24995**	**28406**	**17135-35537**	**9399-60616**
t_{KS}	**2623**	**2960**	**3199**	**2287-3921**	**1481-5763**
t_{RHG}	**892**	**1111**	**1255**	**773-1574**	**422-2531**
$t_{RHG-RHGn}$	**6726**	**6729**	**6931**	**5393-8271**	**3876-10748**
$t_{KS-RHG-RHGn}$	**12117**	**12152**	**12001**	**10797-13422**	**8875-14538**
$tad_{nKS-sKS}$	**395**	**532**	**597**	**347-786**	**147-1232**
$tad_{nKS-RHGn}$	**1383**	**1659**	**1835**	**1140-2333**	**591-3710**
$tad_{sKS-RHGn}$	**1142**	**1378**	**1533**	**928-1970**	**418-3165**
$tad_{wRHG-RHGn}$	**302**	**348**	**418**	**227-550**	**97-944**
$tad_{eRHG-RHGn}$	**393**	**520**	**616**	**342-801**	**171-1358**
$tad_{wRHG-eRHG}$	**411**	**555**	**653**	**367-851**	**157-1429**
$tad_{RHGn-ARHG}$	**2756**	**3193**	**3523**	**2352-4347**	**1369-6828**
$tad_{AKS-RHGn}$	**4490**	**4588**	**4831**	**3731-5590**	**2828-7892**
$tad_{AKS-ARHG}$	**4408**	**4458**	**4730**	**3629-5498**	**2658-7963**
$tad_{AKS-ARHG-RHGn}$	**9679**	**9532**	**9524**	**7888-11066**	**6358-12919**
m(nKS-sKS)	*0.6727*	*0.6018*	*0.5685*	*0.3770-0.7830*	*0.0892-0.9385*
m(sKS-nKS)	*0.7330*	*0.6314*	*0.6072*	*0.4481-0.7911*	*0.1851-0.9346*
m(nKS-RHGn)	*0.4827*	*0.5336*	*0.5324*	*0.3595-0.7206*	*0.1277-0.9082*
m(RHGn-nKS)	*0.3668*	*0.4842*	*0.4966*	*0.2930-0.7092*	*0.0875-0.9153*
m(sKS-RHGn)	*0.4810*	*0.5093*	*0.5082*	*0.3190-0.6977*	*0.1062-0.9089*
m(RHGn-sKS)	*0.4920*	*0.4525*	*0.4587*	*0.2549-0.6432*	*0.0771-0.8880*
m(RHGn-wRHG)	*0.6461*	*0.5674*	*0.5482*	*0.3609-0.7438*	*0.1142-0.9258*
m(wRHG-RHGn)	*0.3925*	*0.4341*	*0.4447*	*0.2729-0.6058*	*0.0873-0.8413*
m(RHGn-eRHG)	*0.3305*	*0.4140*	*0.4365*	*0.2319-0.6291*	*0.0618-0.8750*
m(eRHG-RHGn)	*0.6072*	*0.5726*	*0.5507*	*0.3583-0.7577*	*0.1073-0.9243*
m(wRHG-eRHG)	*0.5034*	*0.4724*	*0.4766*	*0.2736-0.6758*	*0.0802-0.8891*
m(eRHG-wRHG)	*0.5453*	*0.5396*	*0.5340*	*0.3291-0.7465*	*0.1069-0.9388*
m(RHGn-ARHG)	*0.5802*	*0.4897*	*0.4851*	*0.2797-0.6870*	*0.0781-0.8904*
m(ARHG-RHGn)	*0.4081*	*0.5080*	*0.5110*	*0.2993-0.7299*	*0.0853-0.9345*
m(ARHG-AKS)	*0.4734*	*0.5082*	*0.5056*	*0.3033-0.7097*	*0.0850-0.9212*
m(AKS-ARHG)	*0.5824*	*0.5220*	*0.5121*	*0.3036-0.7210*	*0.0773-0.9255*
m(RHGn-AKS)	*0.7032*	*0.5235*	*0.5139*	*0.3053-0.7261*	*0.0887-0.9158*
m(AKS-RHGn)	*0.4066*	*0.5027*	*0.5066*	*0.3136-0.7197*	*0.0929-0.8935*
m(AKS-ARHG-RHGn)	*0.5357*	*0.5008*	*0.4916*	*0.3017-0.6756*	*0.0834-0.8904*
m(ARHG-RHGn-AKS)	*0.3359*	*0.4501*	*0.4680*	*0.2424-0.6856*	*0.0663-0.9148*

2.D.2. Effective population sizes.
Effective population sizes (Ne) for all recent Northern and Southern KS, Western and Eastern RHG,
and RHG neighbors' lineages are often reasonably well estimated with relatively reduced 90%CI and
substantial departure from their priors, for each 25 sets of sampled population combinations
(**SupplementaryFigureSF7xPart1**, **SupplementaryTableST4x**). Combining posterior distributions
among the 25 separate tests, we find Ne posterior estimates ranging from modal point-estimates of
9,461 diploid effective individuals (90%CI=[5,074-67,317]) in the wRHG to 18,022 (90%CI=[9,150-
77,157]) in the RHGn extant lineages (**SupplementaryFigureSF7xPart1**, **TableT4x**). Importantly, we
were unable to satisfactorily estimate, for almost all 25 combinations of sampled populations, the three
effective population sizes for, respectively, the ancient KS lineage, the ancient RHG lineage, and the
lineage ancestral to all RHG and RHG neighbors' extant lineages (**TableT4x**,
SupplementaryFigureSF7xPart1, **SupplementaryTableST4x**).
Conversely, our posterior estimates of the effective size of the lineage most ancestral to all our
populations was satisfactorily estimated, with relatively narrow 90%CI and substantial departure from
the prior distributions, for almost all 25 sets of population combinations. Combining posterior
distributions, despite a noteworthy variation across the 25 tests (**SupplementaryFigureSF7xPart1**,
SupplementaryTableST4x), we find an ancestral effective size for the lineage ancestral to all our
Central and Southern African extant populations of 18,507 diploid effective individuals
(90%CI=[9,399-60,616]), the largest posterior estimate across all recent and ancient lineages in our
analyses (**TableT4x**). Note that we considered large priors (Uniform[100-100,000] diploid individuals)
for constant effective population sizes in all lineages with possible changes at each divergence time, for
simplicity. Therefore, the posterior estimates here found may be different in future procedures
considering more complex effective demographic regimes likely to have occurred in certain lineages,
such as possible bottlenecks and/or population expansions (e.g. (Patin *et al.*, 2014; Schlebusch *et al.*,
2020; Seidensticker *et al.*, 2021)).
810
2.D.3. Instantaneous gene-flow times.
In between each lineage divergence times, we estimated the time of occurrence of instantaneous and
potentially asymmetric gene-flow exchanges across pairs of lineages (**FigureF5x**), for each 25
combinations of sampled populations separately. For four out of the six recent gene-flow times between
pairs of extant lineages ($tad_{nKS-sKS}$, $tad_{wRHG-eRHG}$, $tad_{wRHG-RHGn}$, $tad_{eRHG-RHGn}$, **FigureF5x**, **TableT2x**), we
find relatively consistent posterior estimates across the 25 tests, almost all with relatively reduced
90%CI and substantial departure from the priors (**FigureF8x-PanelA-D**, **SupplementaryTableST4x**,
TableT4x). Note that, while we also obtain similarly satisfactory posterior distributions among the 25
tests for gene-flows among Northern KS and RHGn lineages ($tad_{nKS-RHGn}$), and among Southern KS and
RHGn lineages ($tad_{sKS-RHGn}$), respectively, we found much larger variance across the 25 different
population sets (**FigureF8x-PanelE-F**, **SupplementaryTableST4x**).
Interestingly, we also obtain satisfactory posterior estimates of gene-flow timing among ancient
lineages for almost all 25 sets of population combinations ($tad_{ARHG-RHGn}$, $tad_{AKS-RHGn}$, $tad_{AKS-ARHG}$, tad_{AKS-}
$_{ARHG-RHGn}$, **FigureF5x**, **TableT2x**). Nevertheless, note that posterior estimates are increasingly variable
from one set of sampled populations to the other as the estimated parameter is further back in time in
the tree-topology (**FigureF8x-PanelG-J**, **TableT4x**).
These results further illustrate that different sample sets used in genetic inferences can provide
substantially differing demographic inferences, sometimes for even the most recent gene-flow events,
but most often for more ancient gene-flow events across ancestral lineages.
830
831

FigureF8x: ABC posterior parameter distribution of gene-flow times.

Neural Network Approximate Bayesian Computation (Blum and François, 2010; Csilléry, François and Blum, 2012), posterior
parameter joint estimations of gene-flow instantaneous times *tad* (in generations before present) for 25 sets of five Central and
Southern African populations for which the winning scenario identified by RF-ABC was Scenario i1-1b (**FigureF5x**,
SupplementaryTableST3x). Methodological details are provided in **FigureF7x** caption and detailed in **Material and**
Methods.

FigureF8x

2.D.4. Gene-flow intensities.

We obtain overall unsatisfactory posterior estimates of the 20 separate unidirectional gene-flow rates
for the 10 separate instantaneous gene-flow events, for a majority of the 25 sets of population
combinations (**SupplementaryFigureSF7xPart2, SupplementaryTableST4x**), with relatively large
90%CI and reduced posterior distributions' departure from priors. Considering only the several
satisfactory posterior estimates for each gene-flow parameter separately, we notice their very large
variability across the sets of population combinations for each parameter, making it unreasonable to
combine posterior distributions to obtain a synthetic value (**TableT4x**). The strong parameter-
estimation limitation encountered here is extensively discussed in the **Discussion** section below.

2.D.5. A synthesis of the demographic history of Central and Southern African populations.

We propose, in **FigureF9x**, a schematic synthesis of all the above inference results for the demographic
history of 25 out of 54 combinations of five Central and Southern African populations, for which all
our RF-ABC scenario-choice procedures provided systematically consistent results.

FigureF9x: Schematic inferred demographic and migration history of Central and Southern African populations.
Schematic representation of the winning Scenario i1-1b (**FigureF5x**), and Neural Network ABC posterior parameter mode
estimates summarizing results obtained separately for 25 sets of five Central and Southern African populations
(**SupplementaryTableST3x**), represented by the gray lines in between population names. For the time of each divergence and
gene-flow event, mode point estimates are provided in generations before present (gbp) in bold, and 90% credibility intervals
are provided between parentheses (**TableT4x**). We provide two estimates of the divergence times estimates in years before
present (ybp), one (upper) corresponding to 30 years per generation and the other (lower) to 20 years per generation. Mode
point estimates of effective population sizes *Ne* are provided in numbers of diploid effective individuals and width of lineages
are proportional to the estimated *Ne* (**TableT4x**, **SupplementaryFigureSF7xPart1**). Note that NN-ABC posterior
distributions for the effective population sizes of the ancestral Khoe-San lineage (AKS), for the ancestral RHG lineage
(ARHG), and for the lineage ancestral to RHGn and ARHG, were all three poorly distinguished from their respective prior
distributions; they were therefore considered them to be un-estimated (**TableT4x**, **SupplementaryFigureSF7xPart1**). Note
that posterior distributions for instantaneous asymmetric gene-flow rates were overall poorly departing from their priors, or
highly variable when substantially departing from their priors, among the 25 sets of population combinations
(**SupplementaryFigureSF7xPart2**, **SupplementaryTableST4x**). Instantaneous asymmetric gene-flow intensities are
therefore considered to be un-estimated. All posterior distributions are shown graphically in **FigureF7x**, **FigureF8x,**
SupplementaryFiguresSF7xPart1-2, and detailed in **TableT4x** and **SupplementaryTablesST4x**.

885
886
887

FigureF9x

Discussion

We aimed at inferring jointly the tree-topology, instantaneous or recurring gene-flow processes, and asymmetric gene-flow intensities across pairs of recent and ancient lineages that most likely produced genome-wide genetic patterns in 54 different combinations of five Central and Southern African populations. Therefore, we explicitly considered the large genetic differentiation observed across African populations at regional and local scales, and the possible confounding effects of complex gene-flow processes on tree-topologies' predictions among 48 complex scenarios in formal statistical competition.

First, our results altogether show that, in fact, considering different sets of populations at a regional and local scale in Africa highlights different aspects of the demographic histories of African populations, sometimes substantially divergent. This demonstrates in practice that apparently discrepant results obtained across previous studies may not necessarily be absolutely reconciled, but rather interpreted as separate illustrations of the large diversity of demographic histories experienced by our species in Africa since its emergence in a remote past. This further strongly advocates for avoiding artificially merging individual samples in un-discriminated *a priori* categories based on geography, subsistence-strategy, and/or linguistics criteria. Instead, we recommend developing novel approaches, probably at a large computational cost as experienced here, to explicitly take into account African genetic diversity at the continental, regional, and/or local scales, and *a posteriori* interpreting the diversity of results obtained separately.

Short periods of gene-flow rather than recurring migrations in the ancient history of Africa

As pointed out here and in numerous previous theoretical or empirical studies (e.g. (Verdu and Rosenberg, 2011; Gravel, 2012; Lachance *et al.*, 2012; Harris and Nielsen, 2016; Lorente-Galdos *et al.*, 2019; Fan *et al.*, 2023; Ragsdale *et al.*, 2023)), whether ancient gene-flow processes occurred recurrently or more instantaneously across lineages has fundamental consequences on the biologic and cultural evolution of our species. Indeed, recurrent gene-flow processes allow for continuous allelic exchanges across populations -thus never reproductively isolated throughout entire periods of evolution-, which may strongly influence the relative influence of drift and selection across populations and the notion of ancient tree-like evolution in human populations in Africa. Conversely, we found that instantaneous gene-flow processes systematically vastly outperformed recurring migration processes, whichever the tree-topology for ancient and recent Central and Southern African lineages divergences, and for all combinations of five observed populations. Therefore, our results unambiguously favor an evolutionary history of African lineages ancestral to a variety of Central and Southern African populations where *Homo sapiens* populations experienced long periods of isolation and drift, followed by short periods of possibly asymmetric gene-flow, which may in turn have induced some reticulations among lineages (Mazet *et al.*, 2016; Ragsdale *et al.*, 2023). Notably, this scenario further allows for differential selection and adaptation processes across lineages (e.g. (Lachance *et al.*, 2012; Schlebusch *et al.*, 2012; Perry *et al.*, 2014)), and for ancient admixture-related selection processes within Africa, analogous to previous findings of such processes having occurred more recently throughout the continent (Breton *et al.*, 2014; Patin *et al.*, 2017; Hamid *et al.*, 2021; Cuadros-Espinoza *et al.*, 2022)).

Nevertheless, it is certain that *Homo sapiens* evolutionary history of ancient and recent gene-flow in Africa has involved more complex processes among pairs of lineages than the ones here identified. Previous studies unquestionably considered arguably more realistic models to explain their data, with either a conjunction of recurring symmetric migrations among pairs of lineages with possible instantaneous introgressions among certain pre-specified recent and/or ancient lineages (Lorente-Galdos *et al.*, 2019; Ragsdale *et al.*, 2023), or only pre-specified instantaneous introgressions without

recurring gene-flows (Lachance *et al.*, 2012; Fan *et al.*, 2023), including specific events involving the
contributions from *Homo sapiens* or non-*Homo sapiens* unsampled lineages. In this context, our results
highlight that complex instantaneous introgression scenarios should probably be preferred as a starting-
point from where one could then increase complexity, for instance by considering more than a single
instantaneous gene-flow event among pairs of ancient lineages.
Moreover, our results advocate for the possibility of at least some such introgressions to be
relatively intense, as scenarios considering low or moderate gene-flows, whether instantaneous or
recurring, are systematically poorly mimetic of observed genomic patterns. Echoing this result, a
previous study identified such high levels of instantaneous introgressions across African lineages (Fan
et al., 2023), albeit the authors considered very different maximum-likelihood approaches based on
different statistics and scenario specifications, as well as different population samples, sometimes
artificially merged in larger categories at a continental scale.
Nevertheless, our approach failed to provide satisfactory posterior estimates of gene-flow
intensities, a serious limitation which we discuss extensively in the following methodological limits and
perspectives section below. Importantly, previous studies investigating conjunctions of recurring and
more instantaneous migrations overall estimated that symmetric recurring gene-flows were weak,
whether recent or ancient (e.g. (Ragsdale *et al.*, 2023)). Although our results are difficult to compare
with previous studies due to strong differences in model specifications, statistics used and
methodological approaches as well as population samples, this latter previous result may be reflected
in ours that disfavor recurring migration models as being poorly explicative of our data.
957
<u>Inferred demographic history in Central and Southern Africa</u>
Extensive previous work agrees that divergences among lineages ancestral respectively to extant Khoe-
San, Rainforest Hunter-Gatherers, and Rainforest Hunter-Gatherer neighbors were among the most
ancient in *Homo sapiens* evolution (e.g. reviewed in (Schlebusch and Jakobsson, 2018; Pfennig *et al.*,
2023)). However, the relative order of their divergences as well as their timing has been a matter of
extensive debate (Schlebusch *et al.*, 2020; Lipson *et al.*, 2022; Fan *et al.*, 2023). This is due in part to
split-time inferences conducted on tree-like topologies considering gene-flow or no gene-flow, using
ancient DNA data or not, and to differing methods, statistics, and population samples.
966
Most ancient divergences in Africa
Here, we formally tested, with Random-Forest ABC scenario-choice, which ancient tree-topology best
explained extant genomic patterns, whichever the gene-flow processes and intensities across pairs of
recent and ancient lineages, for 54 different combinations of five different Khoe-San, Rainforest
Hunter-Gatherer, and Rainforest Hunter-Gatherer neighboring populations. It is the first time to our
knowledge that such formal comparison of numerous competing-scenarios is conducted systematically
for a variety of sets of population samples at a regional and local scale, rather than comparing
maximum-likelihood values from vastly differing models, each obtained separately using differing
population samples (Lipson *et al.*, 2022; Fan *et al.*, 2023; Ragsdale *et al.*, 2023). Importantly, note that
(Lorente-Galdos *et al.*, 2019) also explored highly complex demographic scenarios in Africa with ABC,
focusing on varying unidirectional introgression processes among certain lineages for a fixed tree-shape
among, in particular, three extant African populations at the continental scale.
Our results show that whichever the gene-flow processes considered, the lineage ancestral to
extant KS populations diverged first from the lineage ancestral to RHG and RHGn, for almost all
combinations of sampled populations. Furthermore, we estimated with Neural Network ABC posterior-
parameter inferences that this original divergence likely occurred in a remote past ~300,000 years ago,
followed by the divergence between ancestral RHG and RHGn lineages ~165,000 years ago,
considering between 20 and 30 years per generations.

Both estimates fell within the upper bound of previously obtained results considering a variety of
similar population samples, genetic data, inference methods, and statistics, albeit scenarios may not
have always been specified in analogous ways (Patin *et al.*, 2009, 2014; Verdu *et al.*, 2009, 2013;
Schlebusch *et al.*, 2012, 2020; Lipson *et al.*, 2022; Fan *et al.*, 2023). Note however, that considering
different combinations of populations at the regional and local scales allowed us to identify a small
minority (4/54) of such combinations for which an alternative scenario (Scenario i1-3b), where RHGn
lineages diverged first from a lineage ancestral to all RHG and KS extant populations, somewhat
analogous to those proposed in (Fan *et al.*, 2023), would better explain the data consistently across our
analyses, again whichever the gene-flow processes and range of intensities considered. This latter result
may explain apparently discrepant results across some previous studies, which would then be due to
differing population samples used for demographic inferences and/or to artificial merging of
differentiated populations into larger groups (Lorente-Galdos *et al.*, 2019; Lipson *et al.*, 2022; Fan *et*
al., 2023; Ragsdale *et al.*, 2023), thus advocating for further accounting for the vast genetic diversity
among African populations even at a local scale in future studies.
Note that, in (Ragsdale *et al.*, 2023), the fundamental divergence between their Nama Khoe-San
sample and other Sub-Saharan populations was dated to ~110,000-135,000 years ago under their two
best-fitting models, thus strongly discrepant with our findings. However, these important results are
very difficult to compare with ours, since the authors did not consider any Central African Rainforest
Hunter-Gatherers nor their neighbors in their inferences, and since they investigated highly complex
models specified completely differently from those here envisioned. In particular, Ragsdale and
colleagues included in their models possible very ancient genetic structures, long before *Homo sapiens*
emergence, a feature that is unspecified in our scenarios which considered simply a single ancestral
population in which all extant lineages ultimately coalesce. Nevertheless, note that substructure and
reticulation within the ancestral population is not *per se* incompatible with our scenarios. In fact, it may
be compatible with our posterior estimates of a large effective population ancestral to all extant
populations here investigated, the largest among all inferred ancient and recent Central and Southern
African effective population sizes. Therefore, it will be reasonable in future work to complexify the
scenarios here proposed to evaluate whether very ancient substructures and reticulations within our
ancestral population, prior to the original divergence between Southern and Central African
populations, may improve the fit to the observed genomic data, as proposed by (Ragsdale *et al.*, 2023).
1015
More recent divergences in Central and Southern Africa
More recently during the evolutionary history of Sub-Saharan Africa, we found that the divergence
among Northern and Southern Khoe-San populations largely pre-dated the divergence of Western and
Eastern Congo Basin Rainforest Hunter-Gatherers, a question rarely addressed to our knowledge. First,
we found that KS divergence dated sometime between 50,000 and 80,000 years ago, thus substantially
more recently than estimates previously proposed (Schlebusch and Jakobsson, 2018). Beyond vast
differences in models, methods, and statistics used to provide either inferences, note that our results for
this divergence time were substantially variable across pairs of sampled populations used in each
analysis, some specific sets of populations providing posterior estimates consistent with previous result.
It thus further highlights that complex inferences in Africa imperatively need to explicitly consider
population variation at a local scale.
Interestingly, our synthesized estimates for the Northern and Southern Khoe-San population
divergence were relatively synchronic to the genetic onset of the Out-of-Africa (e.g. (Schlebusch and
Jakobsson, 2018)). Population genetics inferences only provide possible mechanisms to explain
observed genetic patterns and are, in essence, not addressing the possible causes underlying the inferred
mechanisms. In this context, we may hypothesize that the global climatic shifts inducing massive
ecological changes that have occurred in Africa at that time (e.g. (Beyer *et al.*, 2021)), sometimes

proposed to have triggered ancient *Homo sapiens* movements Out-of-Africa, may also have triggered, independently, the genetic isolation among ancestral Khoe-San populations. Nevertheless, where the ancestors of extant Khoe-San populations lived at that time remains unknown and is nevertheless crucial to further elaborate possible scenarios for the causes of the genetic divergence here inferred.

Long after this divergence, we found that Rainforest Hunter-Gatherer populations across the Congo Basin diverged roughly between 17,000 and 27,000 years ago, relatively consistently across pairs of sampled populations used for inferences; estimates highly consistent with previous studies (Patin *et al.*, 2009, 2014; Lopez *et al.*, 2018), despite major differences in gene-flow specifications across RHG groups between studies. Interestingly, this divergence time is relatively synchronic with absolute estimates for the Last-Glacial Maximum in Sub-Saharan Africa (e.g. (Bartlein *et al.*, 2011)). The fragmentation of the rainforest massif during this period in the Congo Basin may have induced isolation between Eastern and Western RHG extant populations, as plausibly previously proposed (Patin *et al.*, 2009). However, similarly as above for the Northern and Southern Khoe-San populations divergence, where the ancestors of extant Eastern and Western Rainforest Hunter-Gatherers lived remains unknown, which prevents us from formally testing this hypothesis (Perry and Verdu, 2017).

Altogether, these results show that Central African Rainforest Hunter-Gatherer and Southern African Khoe-San populations have had, respectively, extensive time for selection processes, including adaptive introgression processes, to have influenced independently both groups of populations as well as populations within each group separately (e.g. (Schlebusch *et al.*, 2012; Breton *et al.*, 2014; Perry *et al.*, 2014; Patin *et al.*, 2017)).

Ancient and recent instantaneous gene-flow times in Africa

We obtained reasonably well estimated instantaneous asymmetric gene-flow times among almost all pairs of ancient and recent lineages, albeit the variation of estimates across sets of Central and Southern African sampled populations increased substantially with most ancient times. In this context, we deem it hard to confidently try to interpret the most ancient event of instantaneous gene-flow between the two most ancestral lineages in our tree-topology. However, other instantaneous gene-flow time estimates throughout the topology were more consistently estimated overall, and showed relative synchronicity in some cases, which has never been reported before to our knowledge, even if they were specified independently in our models and drawn from large distribution *a priori*.

Interestingly, we found strong indications for almost synchronic events of introgressions having occurred during the Last Interglacial Maximum in Africa (Mazet *et al.*, 2016), between ~90,000 and ~135,000 years ago (when considering 20 or 30 years per generation). They involved gene-flow between lineages ancestral to Khoe-San populations and ancestors of Rainforest Hunter-Gatherer neighbors on the one hand and, on the other hand, between lineages ancestral to Khoe-San populations and the lineage ancestral to all Rainforest Hunter-Gatherers. An increase in material-based culture diversification and innovation, possibly linked to climatic and environmental changes locally, has previously been observed during this period of the Middle Stone Age in diverse regions of continental Africa; prompting a long-standing debate as to its causes if human populations were subdivided and isolated biologically and culturally at the time (Ziegler *et al.*, 2013; Scerri *et al.*, 2018; Gosling, Scerri and Kaboth-Bahr, 2022; Thomas *et al.*, 2022).

In this context, our results instead may suggest that population movements at that time among previously isolated populations may itself have triggered the observed increased cultural diversification locally, even if obvious signs of pan-African cultural spread at the time are difficult to assess (Gosling, Scerri and Kaboth-Bahr, 2022). In turn, it would interestingly echo the known effect of within-population genetic diversity increase induced by genetic admixture between previously isolated populations (e.g. (Long, 1991; Verdu and Rosenberg, 2011; Gravel, 2012; Laurent *et al.*, 2023)).

Then, note that we estimated that the instantaneous gene-flow event between the ancestral Rainforest Hunter-Gatherers lineage and that of their extant neighbors seemingly occurred synchronically to the genetic Out-of-Africa ((Beyer *et al.*, 2021); see above). This would imply that possible climatic and ecological shifts at that time may not have only induced population divergences and displacement, but may also have triggered population gene-flow.

Relatively more recently, around 30,000 years ago, we found two loosely synchronic gene-flow events between ancestors to extant Central African Rainforest Hunter-Gatherer neighbors' lineages and, separately, Northern and Southern Khoe-San lineages. This corresponds to the end of the Interglacial Maximum and a period of major cultural changes and innovations during the complex transition from Middle Stone Age to Late Stone Age in Central and Southern Africa (Cornelissen, 2002; Ziegler *et al.*, 2013; Mesfin, Oslisly and Forestier, 2021; Bader *et al.*, 2022; Thomas *et al.*, 2022). Nevertheless, connecting the two lines of genetic and archaeological evidence to conclude for increased population movements at the time and their possible causes should be considered with caution. Indeed, in addition to genetic-dating credibility-intervals being inherently much larger than archaeological dating, this period remains highly debated in paleoanthropology mainly due to the scarcity and complexity of the material-based culture records, and that of climatic and ecological changes locally, across vast regions going from the Congo Basin to the Cape of Good Hope (Cornelissen, 2002; Ziegler *et al.*, 2013; Mesfin, Oslisly and Forestier, 2021; Bader *et al.*, 2022; Thomas *et al.*, 2022).

Finally, we found strong signals for multiple instantaneous gene-flow events having occurred between almost all five recent Central and Southern African lineages between 6000 and 12,000 years ago, during the onset of the Holocene in that region, shortly before or during the beginning of the last Post Glacial Maximum climatic crisis in Western Central Africa (Lézine *et al.*, 2019), the emergence and spread of agricultural techniques (Phillipson, 2005), and the demic expansion of now-Bantu-speaking populations from West Central Africa into the rest of Central and Southern Africa (Bostoen *et al.*, 2015; Patin *et al.*, 2017; Fortes-Lima *et al.*, 2024). These results are consistent with previous investigations that demonstrated the determining influence of Rainforest Hunter-Gatherer neighboring populations' migrations through the Congo Basin in shaping complex socio-culturally determined admixture patterns (Patin *et al.*, 2009, 2014; Verdu *et al.*, 2009, 2013), including admixture-related natural selection processes (Perry *et al.*, 2014; Patin *et al.*, 2017; Lopez *et al.*, 2018, 2019). As our estimates for introgression events are in the upper bound of previous estimates for the onset of the so-called "Bantu expansion" throughout Central and Southern Africa, we may hypothesize here that major climatic and ecological changes that have occurred at that time may have triggered increased population mobility and gene-flow events between previously isolated populations, rather than consider that the Bantu-expansions themselves were the cause for all the gene-flow events here identified.

Finally, we did not find signals of more recent introgression events from Bantu-speaking agriculturalists populations into Northern or Southern Khoe-San populations, in particular among the !Xun, albeit such events have been identified in several previous studies (see (Schlebusch and Jakobsson, 2018)). This is likely due to the fact that we considered only a limited number of individual samples from each population, and therefore may lack power to detect these very recent events with our data and approach.

Conceptual, methodological, and empirical limitations and perspectives for inferring ancient histories from observed genomic data

All previous attempts at reconstructing the human evolutionary histories which led to genomic patterns observed today across African populations have faced major conceptual, methodological, and empirical challenges. Conceptually, large amounts of more-or-less nested scenarios can be envisioned *a priori* to explain extant genetic diversity, based on previous results from paleo-anthropology, population genetics, and paleogenomics. These scenarios may range from tree-like models without gene-flow

events among ancient or recent lineages to complex networks of weakly differentiated populations exchanging migrants over large periods of time, with or without the contribution of ancient *Homo* or non-*Homo* now extinct or unsampled lineages. Systematically exploring all possible models is often methodologically out of reach due to differing fundamental scenario-specifications or, when scenarios are specified and parameterized in analogous ways, due to scenarios' nestedness and un-identifiability in certain parts of their parameter spaces (e.g. (Robert, Mengersen and Chen, 2010)). Empirically, formal scenario comparisons are first hampered by necessarily limited amounts of genomic data representative, at continental, regional, and local scales, of the known diversity and differentiation of human populations in Africa; in addition to yet limited amounts of ancient DNA data throughout the continent. Finally, empirical limitations also emerge from the use of different statistics to explore genomic diversity patterns, which possibly each capture different facets of human evolutionary histories, thus providing discrepant results and interpretations only in appearances. While machine-learning ABC procedures provide significant advantages over other maximum-likelihood approaches, in particular concerning the formal exploration of competing scenarios' fit to observed data across numerous highly complex sometimes nested scenarios and using numerous summary-statistics (Blum and François, 2010; Robert, Mengersen and Chen, 2010; Pudlo *et al.*, 2016; Estoup *et al.*, 2018), the above challenges are also largely faced in this study.

     While most divergence and gene-flow times, as well as effective sizes, were inferred satisfactorily in our results, the lack of satisfactory posterior estimates for all 20 gene-flow rates parameters consistent among the 25 sets of population combinations likely stems from different limitations. First, considering only five individual genomes per population is inherently limiting when trying to estimate gene-flow rates from inter-population summary-statistics in ABC (Fortes-Lima *et al.*, 2021). In future studies, increasing sample sizes and adding statistics based on the inter and intra-individual distribution of admixture fractions (Verdu and Rosenberg, 2011; Gravel, 2012; Ragsdale and Gravel, 2019; Fortes-Lima *et al.*, 2021), will likely improve the posterior estimation of these parameters using machine-learning ABC approaches (Lorente-Galdos *et al.*, 2019). Note, however, that these statistics require non-trivial computation time in ABC frameworks comprising hundreds of thousands of simulations (Boitard *et al.*, 2016; Jay, Boitard and Austerlitz, 2019). Furthermore, in all cases, considering multiple sets of different and substantially genetically diverse populations as the ones here considered from sub-Saharan Africa may inevitably leads to large variation across gene-flow rates' posterior estimates depending on the specific population-combinations.

     Second, for simplicity, we considered a single instantaneous time for the two separate unidirectional gene-flow events, for each gene-flow event between pairs of lineages separately. Therefore, while our RF-ABC scenario-choice results strongly support instantaneous asymmetric gene-flow events rather than recurring asymmetric gene-flows to best explain extant genomic patterns, it is plausible that scenarios where each unidirectional gene-flow event may occur at a different time (Lorente-Galdos *et al.*, 2019), will more realistically explain the observed data. If this is the case, it is possible that the choice of a unique time for two separate gene-flow events rendered the corresponding gene-flow rates harder to identify with our joint NN-ABC posterior parameter estimation procedure. Furthermore, we explored two extreme gene-flow processes, establishing an open competition between only instantaneous gene-flow and only recurring gene-flow processes. While we demonstrated that only instantaneous gene-flow processes vastly outperformed only recurring gene-flow processes, whichever the range of intensities for each event, we only established here a starting point for the future complexification of evolutionary scenarios to be tested. For instance, it will be of natural interest to consider, next, more than a single "pulse" of gene-flow between any two ancient or recent lineage.

     Third, the neural-networks used here may have been unable to satisfactorily identify the 20 gene-flow parameters among the 43 jointly-estimated parameters, due to their lack of complexity when considering a unique layer of hidden neurons. For instance, and as a first step, considering Multilayer-

Perceptrons in the future, *i.e.* more complex neural-networks with additional layers of hidden neurons,
may help posterior estimations of these parameters (e.g. (Wang, Czerminski and Jamieson, 2021)), at
the known cost of non-trivial parameterization of the neural-networks themselves (e.g. (Leung *et al.*,
2003; Jay, Boitard and Austerlitz, 2019; Huang *et al.*, 2024)).
Altogether, joint posterior estimation of numerous gene-flow rates parameters and the timing of
their occurrence under highly complex demographic scenarios remains one of the most challenging
tasks in population genetics. It will unquestionably benefit from future analytical theoretical
developments (Mooney *et al.*, 2023; Agranat-Tamir, Mooney and Rosenberg, 2024), and the
improvement of machine-learning-based inference procedures (e.g. (Murtagh, 1991; Chen *et al.*, 2020;
Yelmen and Jay, 2023; Huang *et al.*, 2024)).
1186

Ancient admixture with *Homo sapiens* or non-*Homo sapiens* unsampled lineages?
We did not explore possible contributions from unsampled lineages, whether from non-*Homo sapiens*
or from ancient "ghost" human populations, and therefore cannot formally evaluate the likeliness of the
occurrence of such events to explain observed data. In all cases, our results demonstrate that explicitly
considering ancient admixture from unsampled populations is not a necessity to explain satisfactorily
large parts of the observed genomic diversity of extant Central and Southern African populations,
consistently with a previous study (Ragsdale *et al.*, 2023), and conversely to others (Lipson *et al.*, 2022;
Fan *et al.*, 2023; Pfennig *et al.*, 2023); at least when considering jointly the 337 relatively classical
population genetics summary-statistics used here for demographic inferences. As discussed above, our
results formally comparing competing-scenarios rather than comparing posterior likelihoods of highly
complex yet vastly differing models, provide a clear and reasonable starting point for future
complexification of scenarios comprising possible contributions from ancient or ghost unsampled
populations, which will unquestionably benefit from the explicit use of additional novel summary-
statistics ((Ragsdale and Gravel, 2019; Fan *et al.*, 2023; Ragsdale *et al.*, 2023); see also above).
In any case, the complexification of scenario-specifications to account for possible past "archaic"
or "ancient" introgressions will not fundamentally solve the issue of the current lack of reliable ancient
genomic data older than a few hundreds or thousands of years from Sub-Saharan Africa (Skoglund *et*
al., 2017; Vicente and Schlebusch, 2020; Pfennig *et al.*, 2023). Indeed, analogously to archaic
admixture signals that were unambiguously identified outside Africa only when ancient DNA data were
made available for Neanderthals and Denisovans (e.g. (Meyer *et al.*, 2012; Prüfer *et al.*, 2014)), we
imperatively need to overcome this lack of empirical ancient DNA data in Africa to formally test
whether, or not, ancient human or non-human now extinct lineages have contributed to shaping extant
African diversity.
1210
1211

1212

Material and Methods

Population samples

Central and Southern Africa dataset

We investigated high-coverage whole genomes newly generated for 74 individual samples (73 after relatedness filtering, see below), from 14 Central and Southern African populations (**FigureF1x**, **TableT1x**). Based on extensive ethno-anthropological data collected from semi-directed interviews with donors and their respective communities, we grouped *a posteriori* the 73 individual samples from 14 populations in three larger categories.

The Baka, Ba.Kola, Bi.Aka_Mbati, Ba.Twa, and Nsua individuals were categorized as Rainforest Hunter-Gatherers (RHG), based on several criteria including self-identification and relationships with other-than-self, socio-economic practices and ecology, mobility behavior, and musical practices (Verdu *et al.*, 2009; Hewlett, 2017).

The Nzime, Ngumba, Ba.Kiga, and Ba.Konjo individuals were categorized as Rainforest Hunter-Gatherer neighbors (RHGn), based on the same set of categorization-criteria as RHG. Indeed, RHG and RHGn populations in Central Africa are known to identify themselves separately and to live in different ways in the Central African rainforest, whilst often sharing languages locally as well as complex socio-economic relationships and interactions that may include intermarriages (Verdu *et al.*, 2013). Historically, RHG populations have been designated as "Pygmies" by European colonists, an exogenous term from ancient Greek that is sometimes used derogatorily by the RHGn.

Finally, the Nama, Ju|'hoansi, Karretjie People, !Xun, and Khutse_San individuals were categorized as Khoe-San populations (KS) based primarily on self-identification and relationships with other-than-self locally, lifestyle, and languages (Schlebusch, 2010). Note that research was approved by the South African San Council. Individuals from these populations were whole-genome sequenced anew with improved methods (see below), compared to previous work (Schlebusch *et al.*, 2020).

Note that the three groups can be further subdivided based on geography into Western and Eastern RHG and RHGn, and into Northern, Central, and Southern KS (**FigureF1x**, **TableT1x**).

Comparative dataset

The 74 genomes were analyzed together with 105 (104 after relatedness filtering, see below) high-coverage genomes from 32 worldwide populations (**FigureF1x**, **TableT1x**). 64 samples from the Simon's Genome Diversity Project (SGDP) (Mallick *et al.*, 2016); 9 HGDP samples (Meyer *et al.*, 2012); 7 samples from the 1000 Genomes project (KGP) (The 1000 Genomes Project Consortium *et al.*, 2015); one Karitiana sample (Rasmussen *et al.*, 2014); 24 samples from the South African Human Genome Project (SAHGP) (Choudhury *et al.*, 2017). Selection criteria were: Illumina paired-end reads, high-coverage (>30X), access to raw data and no (known) population substructure.

Note that, although samples from these datasets have not been collected by us for the purpose of this study, based on previous knowledge and publications, the Biaka and Mbuti HGDP samples can be categorized as RHG and the #Khomani and Ju|'hoansi_comp samples as KS (**FigureF1x**, **TableT1x**).

Data generation

DNA extraction

DNA was extracted from saliva with the DNA Genotek OG-250 kit for the Baka, Nzime, Bi.Aka Mbati, Nsua, and Ba.Konjo samples. DNA was extracted from buffy coats with DNeasy Blood&Tissue spin-column Qiagen™ kits for the Ba.Kola, Ngumba and Ba.Kiga samples. Both extraction methods were used for the Ba.Twa samples. For the Nama, Juǀʼhoansi, Karretjie People, !Xun, and Khutse_San samples, DNA was extracted from EDTA-blood using the salting-out method (Miller, Dykes and Polesky, 1988).

Library preparation and sequencing

Library preparation and sequencing was performed by the SciLifeLab SNP&SEQ Technology platform in Uppsala, Sweden. Libraries were prepared with TruSeq DNA preparation kits, and paired-end sequencing (150 bp read length), was performed on Illumina HiSeqX machines with v2 sequencing chemistry, to a coverage of 30X or more. For the Southern African samples, we included paired-end data (100 bp read length) sequenced on Illumina HiSeq2000 machines obtained previously for the same libraries (Schlebusch *et al.*, 2020).

Sequencing data quality-control processing and relatedness filtering

The processing pipeline used in the present study is adapted from the Genome Analyses Toolkit (GATK) "Germline short variant discovery (SNPs + Indels)" Best Practices workflow (McKenna *et al.*, 2010; DePristo *et al.*, 2011; Van Der Auwera *et al.*, 2013). It is described and compared to the original GATK Best Practices workflow in (Breton *et al.*, 2021). **SupplementaryFigureSF8x** gives an overview of the pipelines used for the processing; all template codes with accompanying detailed explanations for all processing steps are provided below and in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Reads were mapped to a decoy version of the human reference genome, GRCh38 (1000 Genomes Project version), with BWA-MEM (Li and Durbin, 2009) from BWAKIT v0.7.12 using a strategy appropriate for ALT contigs. The input was either FASTQ or BAM files that were reverted to unmapped BAM or to FASTQ prior to mapping. Mapped reads were selected with SAMtools version 1.1 and Picard version 1.126 (https://broadinstitute.github.io/picard/).

Duplicate reads were marked (at the lane level) with Picard version 1.126 MarkDuplicates. Realignment around indels was performed (at the lane level) with GATK version 3.5.0 RealignerTargetCreator and IndelRealigner. We then performed a "triple mask base quality score recalibration (BQSR)", where the sample's variation is used together with dbSNP, as described in (Schlebusch *et al.*, 2020; Breton *et al.*, 2021). This step includes merging the reads from a given sample with SAMtools version 1.1; calling variants with GATK version 3.5.0 HaplotypeCaller with "--genotyping_mode DISCOVERY"; and (in a parallel track) standard BQSR with GATK version 3.5.0 BaseRecalibrator and PrintReads, and dbSNP version 144. Following triple mask BQSR (performed at the lane level), the recalibrated reads from a given sample are merged with SAMtools version 1.1, sorted and indexed with Picard version 1.126 SortSam. Duplicate marking and realignment around indels were performed again, at the sample level.

Variant calling was performed independently for the autosomes and the X chromosomes, due to the difference in ploidy. SNP and short indels were first called with GATK version 3.7 HaplotypeCaller in each genome, with the "--emitRefConfidence BP_RESOLUTION" option. Multi-sample GVCFs

were then generated with CombineGVCF, and variants were finally jointly called with GATK
GenotypeGVCFs. Genotypes were emitted for each site (option "allSites").

Variants were filtered with GATK version 3.7 Variant Quality Score Recalibration (VQSR) using
recommended resources from the GATK bundle: HapMap (version 3.3), 1000 Genomes Project
Illumina Omni 2.5M SNP array, 1000 Genomes Project phase 1 high confidence SNPs, and dbSNP
version 151 for the SNPs; and (Mills *et al.*, 2011) gold standard indels and dbSNP version 151 for the
indels. For the X chromosome, autosomal variants were included for training the VQSR model
(VariantRecalibrator step).

We controlled the dataset for related individuals with KING (Manichaikul et al. 2010) "--kinship"
and plink version 1.90b4.9 (Purcell *et al.*, 2007) "--genome --ppc-gap 100". The dataset was pruned for
linkage disequilibrium (LD) before relatedness estimation with plink, using plink "--indep-pairwise"
with sliding windows of 50 SNPs, shifting by five SNPs, and a r^2 threshold of 0.5. Both methods
identified two pairs of first-degree relatives. We excluded the sample with greatest missingness from
each pair to produce the family unrelated working datasets henceforth used.

The callset was then further refined; two samples with a first degree relative in the dataset were
excluded with GATK version 3.7 SelectVariants; a "FAIL" filter status was set on sites that are
ambiguous in the reference genome (base "N") or had greater than 10% missingness, identified with
VCFtools version 0.1.13 "--missing-site" (Danecek *et al.*, 2011). Moreover, for the autosomes, variants
heterozygous in all samples were marked as failed (Hardy Weinberg Equilibrium -HWE- filter,
identified with VCFtools version 0.1.13 "--hardy").

Coverage was computed with QualiMap version 2.2 (Okonechnikov, Conesa and García-Alcalde,
2016) including or not duplicates ("-sd" option for the latter). The average coverage (without duplicates)
across individuals within populations is provided in **SupplementaryTableST2x**.

Descriptive analyses

Variant counts

We obtained per-sample and aggregated metrics of variant counts (**FigureF2x**), with the tool
CollectVariantCallingMetrics of the software package Picard v2.10.3
(https://broadinstitute.github.io/picard/), applied to the full 177 worldwide individuals dataset after
VQSR, relatedness, HWE and 10% site missingness filtering. We conducted variant counts procedures
for chromosomes 1 to 22. We used dbSNP 156 as a reference; we downloaded
"GCF_000001405.40.gz" and modified the contig names to match contig names in the VCFs.
Separately, we applied the same variant counts pipeline as above to the Central and Southern African
73 individuals' original dataset extracted from the full dataset with BCFtools version 1.17
(https://github.com/samtools/bcftools), using view with the option "-S". All scripts are provided in the
corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Genome-wide Heterozygosity

We calculated observed and expected heterozygosities, for the autosomes and for the X-chromosome
separately using custom Python, Bash, and R scripts (full pipeline available at GitHub
https://github.com/Gwennid/africa-wgs-descriptive), for the 177 worldwide individuals after variant-
counts filtering described above. For autosomes, we considered the main contigs ("chr1" to "chr22").
For the X chromosome, we excluded the Pseudo-Autosomal Regions with coordinates from GRCh38
(PAR1: 10,000 to 2,781,480, PAR2: 155,701,382 to 156,030,896).

For each population with more than one individual and for each autosome and X-chromosome
separately, we counted the number of variable and non-variable sites and excluded multi-allelic sites,
indels, and sites with missing genotypes in at least one individual in the population for simplicity and
conservativeness. For each autosome and each individual, we then counted separately the numbers of
observed homozygous and heterozygous sites of each configuration compared to the reference (0/0;
0/1; 1/0; 1/1). Based on these counts, for each population and each chromosome separately, we first
calculated observed heterozygosities simply as the average proportion of heterozygous individuals per
variable sites with no-missing data in the population sample. We also computed unbiased expected
multi-locus heterozygosities for all variable loci with no missing genotypes in the population as in
equations (2) and (3) in (Nei, 1978). We averaged this value across all sites with no missing information
in the population sample including both variable and non-variable sites, and finally corrected it for
haploid population sample sizes (**FigureF2x**). All scripts are provided in the corresponding GitHub
repository (https://github.com/Gwennid/africa-wgs-descriptive).

Runs of homozygosity

We identified runs of homozygosity (ROH) using the *homozyg* tool from PLINK version 1.90b4.9
(Purcell *et al.*, 2007). We selected autosomal biallelic SNPs from the variant-counts pipeline described
above with GATK version 3.7 SelectVariants with options "-selectType SNP -restrictAllelesTo
BIALLELIC -excludeFiltered". We converted the VCF to TPED and then binary plink fileset with
VCFtools version 0.1.13 "-plink-tped" and plink. For the 177 individuals worldwide, we considered
only ROHs measuring more than 200 Kb (*--homozyg-kb* 200), containing at least 200 SNPs (*--homozyg-*
snp 200), containing at least one variable site per 20 Kb on average (*--homozyg-density* 20), and with
possible gaps of up to 50 Kb (*--homozyg-gap* 50). Default values were considered for all other
parameters: *--homozyg-window-het* 1 *--homozyg-window-snp* 50 *--homozyg-window-threshold* 0.05.
Finally, the total mean ROH lengths were calculated for each population separately using the ".hom"
output-file for each of four length categories: 0.2 to 0.5 Mb, 0.5 to 1 Mb, 1 to 2 Mb, and 2 to 4 Mb
(**FigureF2x**). Corresponding pipelines are provided in the corresponding GitHub repository
(https://github.com/Gwennid/africa-wgs-descriptive).

Individual pairwise genome-wide genetic differentiation

We explored genome-wide genetic differentiation between pairs of individuals with Neighbor-Joining
Tree (NJT) (Saitou and Nei, 1987; Gascuel, 1997), and Multi-Dimensional Scaling (MDS) approaches
based on the pairwise matrix of Allele-Sharing Dissimilarities (ASD) (Bowcock *et al.*, 1991) computed
using 14,182,615 genome-wide autosomal SNPs pruned for low LD, with PLINK version 1.90b4.9
(Purcell *et al.*, 2007) *indep-pairwise* function (*--indep-pairwise 50 5 0.1*). We considered at first all 177
unrelated individuals in our dataset to calculate the ASD matrix, and then subset this ASD matrix for
the 73 Central and Southern African unrelated individuals original to this study before computing NJT
and MDS analyses anew (**FigureF3x**).
We computed the ASD matrix considering, for each pair of individuals, only those 14,182,615
SNPs without missing data, using the *asd* software (v1.1.0a; https://github.com/szpiech/asd; Szpiech,
2020). We computed the unrooted NJT using the *bionj* function of the R package *ape* and setting the
branch-length option to "true". We computed the MDS using the *cmdscale* function in R.
The clustering software ADMIXTURE (Alexander, Novembre and Lange, 2009) allows
researchers to further explore inter-individual genome-wide levels of dissimilarity and resemblance.
Indeed, while it is tedious and often cognitively difficult to explore multiple combinations of
dimensions of genetic variation using an MDS or a NJT approach, ADMIXTURE instead allows
exploring K higher such dimensions at once (Pritchard, Stephens and Donnelly, 2000; Rosenberg, 2002;

Falush, Stephens and Pritchard, 2003; Alexander, Novembre and Lange, 2009; Lawson, van Dorp and Falush, 2018). However, note that the visual representation of relative "distances" across pairs of individuals is lost in the classical bar-plot representation of ADMIXTURE, hence showing the complementarity of this descriptive analysis to the above MDS and NJT.

We considered here only 840,031 genome-wide autosomal SNPs pruned for LD (r^2 threshold 0.1) and with Minimum Allele Frequency above 0.1, for the 177 worldwide unrelated individuals, using PLINK *indep-pairwise* and *maf* functions. We computed unsupervised ADMIXTURE version 1.3.0 clustering with values of K ranging from 2 to 10, considering 20 independent runs for each value of K separately. We then calculated ADMIXTURE results symmetric-similarity-coefficient SSC for each value of K separately in order to find the groups of runs providing highly similar results (SSC>99.8%). Individual's genotype membership proportions to each K cluster were then averaged, per individual, across such highly resembling runs, and then plotted (**FigureF4x**). SSC calculations, averaging results across similar runs, and producing barplots were conducted using the software PONG with the "greedy" algorithm (Behr *et al.*, 2016). Corresponding pipelines are provided in the corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

Machine-Learning Approximate Bayesian Computation scenario-choice and posterior parameter estimation

We reconstructed the complex demographic history of Central and Southern African populations using machine-learning Approximate Bayesian Computation (Tavaré *et al.*, 1997; Beaumont, Zhang and Balding, 2002; Blum and François, 2010; Csilléry, François and Blum, 2012; Pudlo *et al.*, 2016). In principle, in ABC, researchers first simulate numerous genetic datasets under competing scenarios by drawing randomly a vector of parameter values for each simulation in distributions set a priori by the user. For each simulation separately, we then calculate a vector of summary statistics, thus corresponding to a vector of parameter values used for the simulation. The same set of summary statistics is then computed on the observed data. ABC scenario-choice then allows the researcher to identify which one of the competing scenarios produces the simulations for which summary-statistics are closest to the observed ones. Under this winning scenario, ABC posterior-parameter inference procedures allow the researcher to estimate the posterior distribution of parameter values most likely underlying the observed genetic patterns.

48 competing scenarios for the demographic history of Central and Southern African populations

We designed 48 competing demographic-history scenarios possibly underlying the genetic patterns observed in five Central and Southern African populations of five individuals each (**TableT1x**, **FigureF5x**); one Eastern Rainforest Hunter-Gatherer population (eRHG: Nsua, Ba.Twa, or Mbuti), one Western Rainforest Hunter-Gatherer population (wRHG: Baka, Ba.Kola, or Aka Mbati), one Eastern or Western RHG neighboring population (eRHGn: Ba.Kiga or Ba.Konjo; wRHGn: Nzimé or Ngumba), one Northern Khoe-San population (nKS: Ju|'hoansi or !Xun), and one Southern Khoe-San population (sKS: Karretjie People or Nama). Note that we did not include the Khutse-San population in ABC inferences for simplicity, as this population is located at intermediate geographic distances between the nKS and sKS groups of populations.

The 48 competing scenarios differed in their ancient tree-topologies and relative timing of the more recent divergence events, combined with different (duration and intensities) possibly asymmetric gene-flow processes among ancient and recent lineages (**FigureF5x**). This design, while complex, allowed

us, for the first time to our knowledge, to consider explicitly the expected confounding effects of gene-
flow processes on otherwise different tree-topologies, which may also possibly induce certain
reticulations among ancient lineages, while jointly estimating divergence times, gene-flow events'
timing and intensities, and effective population size changes over time.

Importantly, we relied on extensive previous findings having formally demonstrated the common
origin of Congo Basin RHG populations (Patin *et al.*, 2009; Verdu *et al.*, 2009), and that of KS
populations (Schlebusch *et al.*, 2012, 2020), respectively, and thus did not consider all possible tree-
topologies for five tree-leaves. Moreover, we considered only a single RHGn sample in each
combination, in turn from the East or the West of Central Africa, in order to simplify already highly
complex scenarios, as we were not interested here in the detailed demographic history of RHGn which
has been extensively studied previously (e.g. (Patin *et al.*, 2017; Fortes-Lima *et al.*, 2024)). This
simplification was deemed reasonable as RHGn populations throughout the Congo Basin and into
Southern Africa have previously been shown to be strongly more genetically resembling one another,
compared to RHG and KS neighboring populations, or compared to much more genetically dissimilar
RHG and KS populations, respectively (e.g. (Verdu *et al.*, 2009, 2013; Patin *et al.*, 2014, 2017; Lopez
et al., 2018, 2019; Fortes-Lima *et al.*, 2024)); a result that we also obtained here (**FigureF5x**).

1438

Eight competing topologies
We investigated eight competing topologies starting with a single ancestral population and resulting in
five different sampled populations in the present (**FigureF5x**). The eight topologies differ in which
lineage ancestral to modern KS, RHG, or RHGn diverged first from the two others, and in the relative
order of divergence events internal to KS (nKS-sKS divergence) and internal to RHG (wRHG-eRHG
divergence), in order to consider the known variable demographic histories of Sub-Saharan populations
at a regional scale.

Note that divergence and introgression times (all *t* and *tad*, see **FigureF5x**, **TableT2x**), were each
randomly drawn in uniform prior distributions between 10 and 15,000 generations, thus effectively
setting an upper limit for the most ancient divergence times among lineages at roughly 450,000 years
ago considering an upper-bound of 30 years for human generation duration (Fenner, 2005). This was
vastly anterior to the current estimates for the genetic or morphological emergence of *Homo sapiens*
(Hublin *et al.*, 2017; Richter *et al.*, 2017), which allowed us to estimate *a posteriori* the most ancient
divergences among our populations, without constraining our assumptions based on previous results
obtained with different methods, data, and models. Furthermore, these prior-distribution boundaries
allowed for gene-flow events among ancient lineages (see next section), to influence, potentially, the
timing of the earliest divergence events among human lineages previously estimated (Schlebusch and
Jakobsson, 2018; Fan *et al.*, 2023; Ragsdale *et al.*, 2023). Note that we retained for simulations only
those vectors of randomly-drawn parameter-values that satisfied the chronological order of lineage
divergences set for each eight topologies, respectively (**Figure5x**, **TableT2x**)

In each eight topologies, we incorporated the possibility for changes in effective population sizes,
N (**FigureF5x**), during history along each lineage separately by defining constant diploid effective
population sizes parameters separately for each tree-branch, each drawn randomly in U[10-100,000]
(**Figure5x**, **TableT2x**).

1463

1464   *Asymmetric instantaneous or recurring gene-flows and their intensities*
1465   Migration of individuals between populations is ubiquitous in human history (for an overview in Africa,
1466   see e.g. (Schlebusch and Jakobsson, 2018; Pfennig *et al.*, 2023)). Here we aimed at disentangling the
1467   nature of migration processes that may have occurred across pairs of lineages throughout the history of
1468   Central and Southern African populations.

In particular, we first aimed at determining whether gene-flow events during history occurred relatively instantaneously, leading to reticulations in tree-topologies, or, conversely, occurred recurrently over longer periods of time leading to un- or weakly-differentiated lineages throughout history (Henn, Steele and Weaver, 2018; Hollfelder *et al.*, 2021; Ragsdale *et al.*, 2023). To do so, for each eight topologies described above (**FigureF5x**), we simulated gene-flow either as single-generation gene-flow pulses across pairs of lineages, or as constant recurring gene-flow across pairs of lineages in-between each lineage-divergence event. This design resulted in 16 competing scenarios encompassing eight different possible topologies, contrasting instantaneous gene-flow events with recurring gene-flows. Note that, for simplicity, we did not consider possible gene-flows between KS and RHG recent lineages, as genetic signatures of such recent migrations have never been identified in previous genetic studies nor in historical records to our knowledge.

Importantly, we aimed at determining whether gene-flow occurred symmetrically, or not, across pairs of lineages, in particular in the past. Indeed, previous studies already identified recent asymmetric admixture processes between RHG and RHGn (Patin *et al.*, 2009, 2014; Verdu *et al.*, 2009, 2013), and such possible asymmetries have not been explored among ancient lineages in previous studies (Fan *et al.*, 2023; Ragsdale *et al.*, 2023), although they may be influencing ancient tree-topologies or reticulations across lineages and the subsequent past evolution of extant populations.

To do so, for each gene-flow event in each 16 competing scenarios (**FigureF5x**), we parameterized separately the introgression of lineage A into lineage B, from that of lineage B into lineage A, by drawing randomly the corresponding parameter values independently in the same prior distribution (**TableT2x**). ABC posterior-parameter estimations would thus reveal possible asymmetries in gene-flows across pairs of lineages for each event separately, if identifiable from genomic data.

Finally, for each one of the 16 competing combinations of topologies and gene-flow processes, we considered three classes of gene-flow intensities by setting different boundaries of gene-flow parameters' uniform prior-distributions (**TableT2x**): no to low possible gene-flow (U[0, 0.001] for recurring processes, and U[0, 0.01] for instantaneous processes), no to moderate gene-flow (U[0,0.0125] for recurring processes, and U[0, 0.25] for instantaneous processes), or no to intense gene-flow events (U[0,0.05] for recurring processes and U[0,1] for instantaneous processes). Note that, for each three classes of gene-flow intensities, each gene-flow parameter from one lineage to another at each point or period in time was drawn independently in the above prior-distributions' boundaries, and thus may differ across events within a scenario, and across scenarios (**FigureF5x**, **TableT2x**).

Altogether, this design resulted in 16x3 = 48 competing scenarios for the complex demographic history of Central and Southern African populations. Importantly, note that for any given class of gene-flow process or intensity, the eight topologies are highly nested in certain parts of the space of parameter values (Robert, Mengersen and Chen, 2010). In particular, despite the fact that we did not consider competing scenarios with explicitly trifurcating ancient tree-topologies, scenarios in which the parameter values for the oldest and second oldest divergence times are similar are expected to provide results highly resembling those obtained with ancient trifurcation scenarios.

All scenario-parameters are represented schematically in **FigureF5x**, and their prior distributions and constraints are indicated in **TableT2x**.

Simulating genomic datasets

We performed simulations under the coalescent using *fastsimcoal2* (Excoffier and Foll, 2011; Excoffier *et al.*, 2013), for each one of the 48 scenarios described above, separately. For each simulation, we generated a vector of parameter values randomly drawn in prior-distributions and satisfying the topological constraints as described above and in **TableT2x**, with custom-made Python and Bash scripts available in the GitHub repository for this article (https://github.com/Gwennid/africa-wgs-abc). The

genetic mutation model is based on the model deployed in (Jay, Boitard and Austerlitz, 2019). We simulated 100 independent loci (or "chromosomes") with the same structure. Each such "chromosome" corresponds to a linkage block, with the following properties: the type of marker was "DNA", the length of the loci 1 Mb, the recombination rate was 1×10^{-8} per base pair, the mutation rate 1.25×10^{-8} per base pair, and there was no transition bias (transition rate of 0.33).

We performed 5,000 such simulations for each 48 competing scenarios separately in order to conduct Random Forest ABC scenario-choice, hence producing 240,000 separate simulated dataset each corresponding to a single vector of parameter values randomly drawn in prior distributions (**TableT2x**). Then we performed an additional 95,000 separate simulations under the winning scenario obtained with RF-ABC for each 54 separate combinations of observed population samples (see below). We thus reached 100,000 simulations under each winning scenario identified for each 54 observed datasets respectively, to be used for Neural Network ABC posterior parameter inference.

Building observed genome-wide data-sets for 54 sets of five Central and Southern African populations

We considered 54 separate combinations of four eRHG, wRHG, nKS, sKS, and one eRHGn or wRHGn populations, each with five unrelated individuals (**TableT1x**). We prepared a callset of high-quality regions by applying the 1000 Genomes phase 3 accessibility mask and filtering out indels. We pieced high quality windows together to create 1 Mb-long windows. We selected 100 of these regions for which the total length from start to end is less than 1.2 Mb. We then extracted these 100 independent autosomal loci of 1 Mb each from the 25 individuals, hence mimetic of the simulated data.

Calculating 337 summary statistics

For each simulated genetic dataset, we computed 337 summary statistics. 41 summary statistics were computed within each five populations separately (hence 205 statistics for five populations), 132 summary statistics were computed across the five populations, for each one of the 54 combinations of five populations separately. In brief, all summary statistics were calculated with plink v1.90b4.9 (Purcell *et al.*, 2007), R v3.6.1 (R Core Team, 2015), Python v2.7.15, bash, awk[1], and scripts developed by (Jay, Boitard and Austerlitz, 2019), available at https://gitlab.inria.fr/ml_genetics/public/demoseq/-/tree/master, and using the software *asd* (https://github.com/szpiech/asd). The list of calculated summary statistics is provided in **TableT3x**. The same statistics were computed for each one of the 54 observed data-sets separately, using the same computational tools and pipeline. Custom-made Python and Bash scripts for computations of all summary-statistics for each simulation and for the observed data are available in the GitHub repository for this article in the "fsc-simulations/code" folder (https://github.com/Gwennid/africa-wgs-abc).

We used the complete set of 337 statistics to perform Random-Forest ABC scenario-choice, as this method is relatively fast and is unaffected by correlations among statistics (Pudlo *et al.*, 2016). We then considered a subset of 202 statistics among the 337 for Neural-Network ABC posterior-parameter estimation, in order to diminish computation time (**TableT3x**).

Prior-checking simulations' fit to the observed data

Before conducting ABC scenario-choice and posterior parameter estimation inferences, we checked that summary-statistics calculated on the observed data fell within the range of values of summary-statistics obtained from our simulations. First, we visually verified that each observed vector of summary-statistics computed separately from 54 combinations of five populations each, clustered with the 240,000 vectors of summary-statistics obtained from simulations under the 48 competing scenarios,

using two-dimensional PCA calculated with the *prcomp* function R (**SupplementaryFigureSF1x**).
Second, we used the *gfit* function in R to perform goodness-of-fit 100-permutations tests between the
240,000 vectors of summary-statistics obtained from simulations under the 48 competing scenarios and,
in turn, each vector of summary-statistics obtained from the 54 observed datasets respectively.

Results both showed that our simulation design could successfully mimic summary-statistics
observed in our five-population sample sets, for each 54 combinations of five populations separately.
Hence, ABC inferences could be confidently conducted a priori based on the simulations and observed
data here considered.

Random Forest ABC grouped scenario-choice

We conducted series of Random Forest ABC scenario-choice procedures for different groups of
scenarios (Pudlo *et al.*, 2016; Estoup *et al.*, 2018), elaborated specifically to address our different
questions of interest regarding gene-flow processes, their intensities, and topological features of the
history of Central and Southern African populations. Random Forest-ABC scenario-choice has proven
to be performing efficiently and satisfactorily with a significantly lower number of simulations
compared to any other ABC scenario-choice procedure. Moreover, RF-ABC scenario choice is
insensitive to correlations among summary-statistics (Pudlo *et al.*, 2016; Estoup *et al.*, 2018).

Each RF-ABC scenario-choice procedure presented below was conducted using the *predict.abcrf*
and the *abcrf* functions with the "group" option of the *abcrf* package in R (Pudlo *et al.*, 2016), with
1000 decision trees to train the algorithm after checking that error rates were appropriately low with the
err.abcrf function, separately for each 54 combinations of five sampled populations (**FigureF6x**).

We conducted cross-validation procedures considering in turn each one of the simulations as
pseudo-observed data and all remaining simulations to train the algorithm, for each RF-ABC analysis
separately (**SupplementaryFigureSF6xPart0**). While not necessarily predicting the outcome of the
scenario-choice for the observed data, these cross-validation procedures provide us with a sense of the
discriminatory power, *a priori*, of RF-ABC for our set of competing-scenarios, as well as empirical
levels of nestedness among scenarios or groups of scenarios (Robert, Mengersen and Chen, 2010;
Fortes-Lima *et al.*, 2021).

Instantaneous or recurring gene-flows in the history of Central and Southern African populations?
The 5,000 simulations performed under each one of the 48 competing scenarios were first gathered into
two groups of 24 scenarios each (**FigureF5x**), corresponding to either instantaneous asymmetric gene-
flow processes or to recurring constant asymmetric gene-flow between each pair of lineages,
respectively. Both groups thus contained all simulations from the eight competing topologies and all
three sets of possible gene-flow intensities (low, moderate, or high, see above), and only differed in the
process of gene-flow itself. For each 54 combinations of five sampled populations separately, we
performed such RF-ABC scenario-choice to determine the winning group of scenarios (**FigureF6x-panelA**).

Low, moderate, or high gene-flow intensities?
For each 54 combinations of five sampled populations separately, we performed RF-ABC scenario-
choice in order to determine which class of intensities of gene-flows best explained the data, everything
else (topology and gene-flow process) being-equal. We thus considered three groups of scenarios in our
RF-ABC scenario-choice, each corresponding to "low", "moderate", or possibly "high" intensities, and
each grouping the 16 scenarios corresponding to eight topologies and instantaneous or recurring gene-
flow processes indiscriminately (**FigureF6x-panelB**).

Then, we conducted, for each 54 population combinations separately, the same RF-ABC scenario-choice procedure to disentangle groups of gene-flow intensities "all topologies being equal", considering here only the 24 scenarios from the winning group among instantaneous or recurring gene-flow processes obtained from the above scenario-choice procedure (**SupplementaryFigureSF6xPart1**).

Finally, we performed, for each 54 combinations of five populations separately, RF-ABC scenario choice for the 48 competing scenarios grouped in six different groups (encompassing eight scenarios each), combining instantaneous or recurring processes with the three classes of intensities respectively (**FigureF6x-panelC**), thus "intersecting" the two corresponding group analyses (see above).

Which ancestral Central or Southern African lineage diverged first?

The eight topologies considered in the 48 competing scenarios can also be grouped according to their ancient topologies, by considering which ancestral lineage diverged first from all others at the oldest divergence event in the tree (**FigureF5x**). The ancient lineage which separated first from the two others can either be the lineage ancestral to Northern and Southern Khoe-San populations (AKS, scenario topologies 1a, 1b, 1c in **FigureF5x**), the lineage ancestral to eastern and western Rainforest Hunter-Gatherer populations (ARHG, scenario topologies 2a, 2b, 2c in **FigureF5x**), or the lineage leading to Rainforest Hunter-Gatherer Neighboring populations (RHGn, scenario topologies 3a, 3b in **FigureF5x**).

For each 54 combinations of five populations separately, we conducted RF-ABC scenario-choice across these three groups of scenarios, randomly drawing 2/3 of the 5,000 simulations per scenario for the scenarios 1a, 1b, and 1c, and for the scenarios 2a, 2b, and 2c, respectively, and kept all simulations for the scenarios 3a and 3b, in order to even the total number of simulations in competition among the three groups of scenarios. We thus performed RF-ABC scenario-choice across the three groups of topologies "all gene-flow processes and intensities being equal" (**FigureF6x-panelD**).

Then we restricted the RF-ABC scenario-choice of the three-competing groups of topologies only for the 24 scenarios from the winning instantaneous or recurring gene-flow processes obtained above, all gene-flow intensities being equal (**FigureF6x-panelE**).

Northern and Southern Khoe-San populations diverged before or after the divergence between Eastern and Western RHG?

The eight topologies considered in the 48 competing scenarios can alternatively be grouped according to the relative order of divergence-time between the divergence event among Northern and Southern KS populations, and that among Eastern and Western RHG populations; an important question to understand the relative duration of separate evolution of each groups of populations.

To address this specific question we thus conducted, for the 54 combinations of five populations separately, RF-ABC scenario-choice procedures by grouping the 48 competing scenarios in two separate groups of 24 scenarios each. One group corresponded to KS populations diverging from one-another before the RHG divergence (scenario topologies 1b, 2b, 3b, and 1c in **FigureF5x**), and the other group where RHG diverged from one another before the KS divergence (scenario topologies 1a, 2a, 3a, and 2c in **FigureF5x**), instantaneous or recurring gene-flow and all "low", "moderate", or "high" gene-flow intensities being equal among the two groups (**FigureF6x-panelF**). Last, we conducted the same RF-ABC scenario-choice procedure between two groups of four topologies each, but restricted to the 24 scenarios winning among the instantaneous or recurring gene-flow processes as determined from above procedures (**FigureF6x-panelG**).

Finally, we conducted RF-ABC scenario choice procedures considering, respectively, all 48 competing scenarios separately (**SupplementaryFigureSF6xPart2**), and all 24 scenarios winning

1655 among the instantaneous or recurring gene-flow processes as determined above
1656 (**SupplementaryFigureSF6xPart3**).
1657

Neural Network ABC posterior-parameter estimation

We intersected all the RF-ABC scenario-choice results considering the different groups of scenarios or
all scenarios independently, as described above, in order to determine, which single scenario was
winning in the majority of the 54 combinations of five Central and Southern African populations. We
found that the Scenario i1-1b conservatively produced simulations whose observed summary-statistics
most resembled those obtained in 25 out of the 54 possible population combinations here tested (see
Results, the detailed list of 25 population combinations is provided in **SupplementaryTableST3x**).
The second most often winning scenario, Scenario i1-3b, only succeeded for four population
combinations among the 54 tested.
Based on this *a posteriori* winning Scenario i1-1b, we performed an additional 95,000 simulations
considering the same prior distributions and constraints among parameters, in order to obtain 100,000
simulations for further ABC posterior parameter inferences, separately for each one of the 25 population
combinations for which it was identified confidently as the winner, separately. As it remains difficult
to estimate jointly the posterior distribution of all parameters of the complex scenarios here explored
with RF-ABC (Raynal *et al.*, 2019), we instead conducted Neural Network ABC joint posterior
parameter inferences using the *abc* package in R (Blum and François, 2010; Csilléry, François and
Blum, 2012), following best-practices previously bench-marked for highly complex demographic
scenarios (Jay, Boitard and Austerlitz, 2019; Fortes-Lima *et al.*, 2021). There are no rules of thumb in
order to determine a priori the best number of neurons and tolerance rate to be set for training the neural
network in NN-ABC (Jay, Boitard and Austerlitz, 2019).
There are no evident criteria to choose *a priori* the best tolerance level and numbers of neurons in
the neural network's hidden layer for parameterizing the NN-ABC posterior-parameter estimation
procedure (Jay, Boitard and Austerlitz, 2019; Huang *et al.*, 2024). As the total number of parameters in
the winning Scenario i1-1b was large (43, **FigureF5x** and **TableT2x**), and as the number of summary-
statistics considered was also large (202, **TableT3x**), we chose to conduct the 25 NN-ABC posterior-
parameter inferences for the 25 combinations of sampled populations considering, in-turn, 7, 14, 21,
28, 35, 42, or 43 neurons in the hidden layer (43 being the assumed number of dimensions equal to the
number of parameters for this scenario (Blum and François, 2010; Jay, Boitard and Austerlitz, 2019)),
and a tolerance level of 0.01, thus considering the 1,000 simulations closest to each observed dataset
out of the 100,000 performed, respectively. We found *a posteriori* that, considering 42 neurons in the
hidden layer provided overall posterior parameter distributions departing from their priors, and in
particular for divergence-times parameters which we were highly interested in, and therefore decided
to provide all posterior parameter distributions' results using this parameterization (tolerance = 0.01,
number of hidden neurons = 42) for the training of the neural network.
We thus performed 25 separate NN-ABC joint parameter posterior inferences, using the
"neuralnet" method option in the function *abc* of the *abc* package in R (Csilléry, François and Blum,
2012), with logit-transformed ("logit" transformation option), an "epanechnikov" kernel, parameter
values within parameter-priors' boundaries, a tolerance level of 0.01, and 42 neurons in the hidden
layer. Adjusted posterior parameter distributions obtained with this method were then plotted for each
parameter separately using a gaussian kernel truncated at the boundaries of the parameter prior-
distributions. We then estimated the mode, median, mean, and 50% and 95% Credibility Intervals of
these distributions *a posteriori*. We provide all posterior-parameter distributions together with their
priors in figure-format (**FigureF7x**, **FigureF8x**, **SupplementaryFigureSF7xPart1-2**), and in table-
format (**TableT4x**, **SupplementaryTableST4x**).

1702

GitHub repositories for this work

1703

1704

1705   https://github.com/Gwennid/africa-wgs-descriptive

1706

1707   https://github.com/Gwennid/africa-wgs-abc

1708

1709

Data availability statement

1710

1711   All raw whole-genome sequence FASTQ data original to this study are made available on the European
1712   Genome-Phenome Archive (https://ega-archive.org/), provided that requests conform to ethical
1713   requirements and informed consents provided by donors, as detailed in the corresponding Data Access
1714   Policies associated with each data repository. Data access requests are evaluated by the associated Data
1715   Access Committees. **EGA ACCESSION NUMBERS PENDING ACCEPTANCE**.

1716

1717

Supporting Information

8 Supplementary Figures and 4 Supplementary Tables

1724 **SupplementaryTableST1x: Whole genome total variant counts among 177 unrelated worldwide individuals**
1725 Variant counts compared to the reference sequence for the human genome GRCh38 and previously reported variants in dbSNP
1726 156. "SD" stands for "standard deviation". See **Material and Methods** for details about the quality control, relatedness
1727 filtering, and variant count procedures.
1728
1729

	Total variant count		Per individual variant count among 177 worldwide unrelated individuals			
	Original dataset: 73 Central and Southern African unrelated individuals	Entire dataset: 177 worldwide unrelated individuals	Total	SD	Min	Max
Number of biallelic SNPs	26,780,319	36,272,545	4,245,000	375,379	3,093,785	4,642,544
Number of biallelic SNPs not in dbSNP 156	854,114	1,055,245	8,052	4,563	1,152	27,406
Number of multiallelic SNPs	241,428	257,906	56,330	5,162	38,057	61,716
Number of simple indels	2,454,965	3,159,306	385,391	35,522	278,237	421,976
Number of novel simple indels not in dbSNP 156	114,362	189,124	1,651	514	771	627
Number of complex indels	969,025	977,542	414,037	71,029	289,993	566,951
Number of variants appearing in a single sample	8,404,499	12,638,316	71,403	15,094	22,444	104,211
Number of filtered SNPs	2,940,629	3,296,775	247,172	48,579	66,597	350,377

1730
1731

SupplementaryTableST2x: Whole genome biallelic SNPs counts in 46 worldwide populations
Mean number of bi-allelic SNPs across unrelated individuals from 46 worldwide populations, and associated standard deviations (when applicable), compared to the reference sequence for the human genome GRCh38 and compared to previously reported variants in dbSNP 156. See **Material and Methods** for details about the quality control, relatedness filtering, and variant count procedures. Populations are ordered in increasing mean number of SNPs per pop. The information original to this study is indicated in bold. "SD" stands for standard deviation. "*na*" stands for not applicable. Additional population information and geographical location of samples can be found in **FigureF1x** and **TableT1x**.

Population Name[1]	Sampling location	Dataset[2]	N[3]	Mean coverage X (SD)	Mean number of biallelic SNPs (SD)	Mean number of biallelic SNPs not in dbSNP 156 (SD)
Karitiana	Brazil	SGDP, HGDP	5	31.1 (7.8)	3,214,086 (69,241)	1,030 (547)
Papuan	Papua New Guinea	SGDP, HGDP	6	35.8 (7.8)	3,414,518 (74,893)	841 (290)
French	France	SGDP, HGDP	4	33.8 (8.7)	3,455,358 (77,954)	652 (225)
Dai	China	SGDP, KGP	6	37.7 (10.5)	3,480,744 (67,644)	695 (407)
CEU	United States of America	KGP	2	67.3 (0.2)	3,551,792 (7,819)	1,539 (57)
Saharawi	Western Sahara	SGDP	2	43.7 (2.2)	3,726,018 (27,793)	590 (21)
Mozabite	Algeria	SGDP	2	34.9 (1.6)	3,736,308 (69,393)	563 (11)
Somali	Kenya	SGDP	1	37.7 (*na*)	3,940,793 (*na*)	540 (*na*)
Coloured	South Africa	SAHGP	8	44.9 (2.2)	4,000,396 (97,447)	10,169 (2,319)
Dinka	Sudan	SGDP, HGDP	4	31.3 (5.8)	4,142,138 (97,001)	3,146 (4,811)
Maasai	Kenya	SGDP	2	38.6 (5.5)	4,169,732 (5,281)	654 (27)
Mandinka	Gambia	KGP	1	27.2 (*na*)	4,193,651 (*na*)	1,602 (*na*)
Mandenka	Senegal	SGDP, HGDP	4	30 (7.6)	4,217,398 (107,847)	914 (464)
Yoruba	Nigeria	SGDP, HGDP	4	31.1 (5)	4,233,500 (11,5371)	755 (195)
Ba.Kiga	**Mukono (Uganda)**	**this study**	**5**	**39.4 (4.6)**	**4,274,466 (25,201)**	**5,391 (191)**
Gambian	Gambia	SGDP	2	35.5 (0.8)	4,277,874 (12,976)	816 (107)
Esan	Nigeria	SGDP, KGP	3	39.7 (9.5)	4,284,041 (18,359)	704 (67)
Bantu_Kenya	Kenya	SGDP	2	38.7 (6.8)	4,288,533 (10,954)	611 (88)
Luo	Kenya	SGDP	2	34 (0.2)	4,293,632 (1,003)	722 (4)
Luhya	Kenya	SGDP, KGP	3	40.6 (12.1)	4,294,198 (28,096)	704 (37)
Igbo	Nigeria	SGDP	2	36.3 (8)	4,294,870 (1,867)	1,166 (9)
Lemande	Cameroon	SGDP	2	36.3 (0.5)	4,306,880 (5,284)	1,530 (37)
Bantu_Herero	Namibia	SGDP	2	38.4 (2.2)	4,318,604 (33,525)	690 (161)
Ba.Konjo	**Mulimassenge (Uganda)**	**this study**	**5**	**43 (6.4)**	**4,322,177 (6,336)**	**5,349 (773)**
Nzime	**Messea (Cameroon)**	**this study**	**5**	**35.5 (2.6)**	**4,322,734 (18,671)**	**5,305 (208)**
Ngumba	**Dispersed between Lolodorf and Kribi (Cameroon)**	**this study**	**5**	**40.2 (2.2)**	**4,332,324 (9,273)**	**4,810 (552)**
Kongo	Cameroon	SGDP	1	42.8 (*na*)	4,336,731 (*na*)	1233 (*na*)
Mende	Sierra Leone	SGDP, KGP	3	39.6 (8.3)	4,341,512 (13,486)	866 (180)
Zulu	South Africa	SAHGP	1	42.8 (*na*)	4,376,691 (*na*)	8,146 (*na*)
Xhosa	South Africa	SAHGP	8	42.4 (1.1)	4,385,032 (56,902)	10,085 (1,300)
Ba.Twa	**Kebiremu, Byumba, Kitariro, Mgungu, Nteko (Uganda)**	**this study**	**6**	**53.8 (17.7)**	**4,426,223 (32,552)**	**20,251 (1,022)**
Bantu_Tswana	South Africa	SGDP	2	36.7 (3.1)	4,430,870 (55,537)	720 (37)
Mbuti	Democratic Republic of Congo	SGDP, HGDP	5	32.1 (7.4)	4,434,425 (110,658)	1,227 (488)
Sotho	South Africa	SAHGP	7	42.9 (6.4)	4,446,796 (23,172)	12,040 (1,150)
Ba.Kola	**Dispersed between Lolodorf and Kribi (Cameroon)**	**this study**	**5**	**39.2 (1.9)**	**4,475,922 (50,261)**	**6,995 (1,948)**
Nsua	**Bundimassoli (Uganda)**	**this study**	**5**	**37.6 (4.9)**	**4,486,217 (22,774)**	**16,443 (984)**
Baka	**Bosquet (Cameroon)**	**this study**	**7**	**40.7 (3.5)**	**4,503,977 (23,946)**	**3,927 (144)**
Bi.Aka_Mbati	**Bombeketi section of Bagandou (Central African Republic)**	**this study**	**5**	**43.4 (13.6)**	**4,514,318 (17,376)**	**4,587 (198)**
Biaka	Central African Republic	SGDP	2	39.1 (0.3)	4,521,140 (8,155)	726 (52)
Nama	**Windhoek (Namibia)**	**this study**	**5**	**48.9 (4.3)**	**4,530,168 (18,807)**	**27,902 (462)**
Ju|'hoansi_comp	Namibia	SGDP, HGDP	4	33.9 (7)	4,554,593 (103,099)	1,382 (273)
#Khomani	South Africa	SGDP	2	44.7 (2.8)	4,559,846 (79,851)	1,364 (48)
Khutse_San	**Kutse Game reserve (Botswana)**	**this study**	**5**	**46.3 (6.7)**	**4,592,852 (27,794)**	**27,989 (1,353)**
Karretjie	**Colesberg (South Africa)**	**this study**	**5**	**46.6 (5.4)**	**4,601,724 (26,951)**	**27,720 (1,968)**
!Xun	**Omega camp (Namibia) and Schmidtsdrift (Sout Africa)[4]**	**this study**	**5**	**45.6 (1.9)**	**4,614,835 (10,047)**	**28,923 (1,371)**
Ju|'hoansi	**Tsumkwe (Namibia)**	**this study**	**5**	**49 (4.6)**	**4,628,957 (8,139)**	**26,665 (6,491))**

[1]Population Names are self reported for the original dataset presented in this study

[2]SGDP: Mallick et al 2016; KGP: Auton et al. 2015; HGDP: Meyer et al. 2012, Rasmussen et al. 2014; SAHGP: Choudhury et al. 2017

[3]Number of unrelated individuals considered in all analyses in this study (see Material and Methods)

[4]The place of origin of the !Xun is around Menongue in Angola.

SupplementaryTableST3x: 25 combinations of five population samples out of 54 combinations tested for which Scenario i1-1b is winning with Random-Forest ABC among the 48 competing scenarios.

Values in the table correspond to posterior densities plotted in the left panels of **FigureF7x**, **FigureF8x**, and **SupplementaryFigureSF7xPart1-Part2**. Parameter definitions and priors are provided in **TableT2x** and represented graphically in the corresponding scenario panel of **FigureF5x**. Values in italic are not satisfactorily departing from the priors.

1754

	nKS	sKS	RHGn	wRHG	eRHG
Combination 1	Ju|'hoansi	Karretjie	Nzime	Baka	Nsua
Combination 2	Ju|'hoansi	Karretjie	Ngumba	Ba.Kola	Nsua
Combination 3	Ju|'hoansi	Karretjie	Ngumba	Ba.Kola	Ba.Twa
Combination 4	Ju|'hoansi	Karretjie	Nzime	Bi.Aka_Mbati	Nsua
Combination 5	Ju|'hoansi	Karretjie	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 6	Ju|'hoansi	Karretjie	Ngumba	Ba.Kola	Mbuti
Combination 7	Ju|'hoansi	Karretjie	Nzime	Bi.Aka_Mbati	Mbuti
Combination 8	Ju|'hoansi	Karretjie	Ba.Konjo	Baka	Mbuti
Combination 9	Ju|'hoansi	Karretjie	Ba.Konjo	Ba.Kola	Mbuti
Combination 10	Ju|'hoansi	Karretjie	Ba.Konjo	Bi.Aka_Mbati	Mbuti
Combination 11	Ju|'hoansi	Nama	Ba.Konjo	Baka	Nsua
Combination 12	!Xun	Nama	Ba.Konjo	Baka	Nsua
Combination 13	Ju|'hoansi	Nama	Nzime	Baka	Nsua
Combination 14	Ju|'hoansi	Nama	Ngumba	Ba.Kola	Nsua
Combination 15	Ju|'hoansi	Nama	Nzime	Baka	Ba.Twa
Combination 16	Ju|'hoansi	Nama	Ngumba	Ba.Kola	Ba.Twa
Combination 17	Ju|'hoansi	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 18	Ju|'hoansi	Nama	Ba.Konjo	Ba.Kola	Nsua
Combination 19	Ju|'hoansi	Nama	Ba.Kiga	Ba.Kola	Ba.Twa
Combination 20	Ju|'hoansi	Nama	Ba.Kiga	Bi.Aka_Mbati	Ba.Twa
Combination 21	!Xun	Nama	Nzime	Bi.Aka_Mbati	Nsua
Combination 22	!Xun	Nama	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 23	!Xun	Karretjie	Nzime	Bi.Aka_Mbati	Nsua
Combination 24	!Xun	Karretjie	Nzime	Bi.Aka_Mbati	Ba.Twa
Combination 25	!Xun	Karretjie	Ba.Konjo	Bi.Aka_Mbati	Nsua

SupplementaryTableST4x: NN-ABC posterior parameter estimation of all parameters in Scenario i1-1b for results
from each 25 sets of five populations each, separately.
Each sampled-population combination is provided as a single line. Table-headers in bold indicate the categorization of sampled
populations into Northern Khoisan (nKS), Southern Khoisan (sKS), Rainforest Hunter-Gatherer neighbors (RHGn), Western
Rainforest Hunter-Gatherer (wRHG), or Eastern Rainsforest Hunter-Gatherer (eRHG) groups, as per scenario topology and
simulation design explicated in **Material and Methods** and in **FigureF5x**. Note that each sampled populations considered in
the RF-ABC scenario choice analyses are represented twice or more in the 25 combinations. See ".xlsx" file as the table is to
large to reasonably fit in a A4 page format.
1763
1764
1765

SupplementaryFigureSF1x: PCA for ABC prior-checking of simulations fit to the observed data.
We calculated 337 summary-statistics for each one of the 240 000 simulated data sets, 5000 under each one of the 48 competing
scenarios (**FigureF5x** and **Material and Methods**), which we projected on the first two axes of a principal component analysis.
Each vector (corresponding thus to a single simulation) is represented by a gray cross. We separately computed the same 337
summary-statistics, separately for each one of the 54 sets of five observed Central and Southern African populations included
in our analyses. Each observed vector is then projected, in turn, on the PCA obtained from simulations only. Each such
observed vector is represented as a single yellow dot. We can see that all the 54 observed sets of populations fall well within
the space of simulated dataset, thus allowing us a priori to conduct machine-learning ABC scenario choice and posterior
parameter estimation procedures.
1775
1776
1777

SupplementaryFigureSF1x

SupplementaryFigureSF6xPart0: Out-of-bag cross-validation errors for Random-Forest ABC scenario-choices among 48 competing scenarios.

All RF-ABC cross-validation analyses are conducted *a priori*, without using any observed data, by considering, in-turn, each simulation as observed data while all remaining simulations are used to train the Random Forest. Each panel provides the *a priori* cross-validation errors for the corresponding RF-ABC scenario-choice analysis detailed in each panel in **FigureF6x**. While these cross-validation errors do not predict the specific outcome of RF-ABC prediction for the observed data, they provide levels of scenarios and groups of scenarios nestedness *a priori* for the entire space of parameters used for simulations. (**A**) shows that intermediate and recurring gene-flow processes are, all topologies and gene-flow intensities "being equal", clearly differentiable by RF-ABC and little nested. (**B**) and (**C**) show that different classes of gene-flow intensities are also relatively clearly identifiable, albeit more nested, as expected by design since the "high gene-flow" class comprises parameter values of the moderate and the low intensity classes of gene-flows (**Material and Methods**, **TableT2x**). (**D**) and (**E**) show that ancient tree topologies are relatively well distinguishable *a priori* all gene-flow processes "being equal", despite some amounts of errors due to the expected nestedness in the space of parameter values where ancient divergence times are resembling and thus undistinguishable. (**F**) and (**G**) show that the chronological order between the divergence-time of Northern and Southern KS lineages, and that of Western and Eastern RHG lineages, are also relatively well distinguishable a priori with RF-ABC scenario-choice, albeit with an increased cross-validation error expected due to high levels of nestedness in the spaces of parameter-values where both separate events may occur at similar time.

SupplementaryFigureSF6xPart0

SupplementaryFigureSF6xPart1: RF-ABC scenario choice among 24 competing scenarios with instantaneous
asymmetric gene-flow processes, for three groups of gene-flow intensities, all topologies being equal.
(**A**) corresponds to the scenario-choice results for the same test as in **FigureF6x-PanelC**, restricted to the 24 competing
scenarios considering instantaneous gene-flow processes only, all scenarios topologies "being equal". Description of the x and
y axis legends are provided in **FigureF6x**. (**B**) corresponds to the RF-ABC cross-validation prior error for the scenario-choice
procedure conducted in (**A**). Errors were obtained without considering observed data and using, instead, all simulations in turn
as pseudo-observed data and the remaining simulations to train the RF.
1812
1813
1814

SupplementaryFigureSF6xPart2: RF-ABC scenario choice among 48 competing scenarios without groups.
(**A**) corresponds to the RF-ABC scenario-choice results obtained when considering all 48 scenarios (5,000 simulations per
scenario), as separate competitors without grouping them, for each 54 combinations of five sampled-populations separately.
Description of the x and y axis legends are provided in **FigureF6x**. (**B**) corresponds to the RF-ABC cross-validation prior error
for the scenario-choice procedure conducted in (**A**). Errors were obtained without considering observed data and using, instead,
all simulations in turn as pseudo-observed data and the remaining simulations to train the RF. Note that, in (**B**), whichever the
intensity of the gene-flow, when considering only recurring gene-flow processes, topologies are harder to distinguish from one
another a priori. Conversely, such difficulty to distinguish a priori among topologies remains overall limited for instantaneous
gene-flow processes, and errors increase only when considering the possibility of intense instantaneous gene-flows. However,
as mentioned throughout the article, the capacity to distinguish a priori among scenarios in the entire space of summary-
statistics values does not predict the power to predict a winning scenario in the specific space occupied by observed data (Pudlo
et al. 2016).
1831
1832
1833

A

RF-ABC scenario-choice
Topologies and Low vs. *Moderate* vs *High*
Instantaneous and Recurring gene-flows
48 competing scenarios - no groups
54 population sets

B

RF-ABC scenario-choice
48 competing scenarios - no groups
Out-of-bag cross-validation *a priori*
240 000 simulations: 5000 per scenario
no real data

RF−ABC predicted model

SupplementaryFigureSF6xPart3: RF-ABC scenario choice among 24 competing scenarios without groups.
(**A**) corresponds to the RF-ABC scenario-choice results obtained when considering all 24 instantaneous gene-flow scenarios
(5,000 simulations per scenario), as separate competitors without grouping them, for each 54 combinations of five sampled-
populations separately. Description of the x and y axis legends are provided in **FigureF6x**. (**B**) corresponds to the RF-ABC
cross-validation prior error for the scenario-choice procedure conducted in (**A**). Errors were obtained without considering
observed data and using, instead, all simulations in turn as pseudo-observed data and the remaining simulations to train the
RF. Note that prediction and cross-validation error results resemble those obtained for these 24 specific scenarios obtained
when considering instead all the 48 scenarios in competition (**SupplementaryFigureSF6xPart2**).
1846
1847
1848

SupplementaryFigureSF6xPart3

A

RF-ABC scenario-choice
Topologies and Low vs. *Moderate* vs *High*
Instantaneous gene-flows
24 competing scenarios - no groups
54 population sets

B

RF-ABC scenario-choice
24 competing scenarios - no groups
Out-of-bag cross-validation *a priori*
120 000 simulations: 5000 per scenario
no real data

SupplementaryFigureSF7xPart1: ABC posterior distribution of effective population size parameters.
Neural Network Approximate Bayesian Computation (Csilléry et al. 2012), posterior parameter joint estimations of Effective
population sizes *Ne* (in generations before present) for 25 sets of five Central and Southern African populations for which the
winning scenario identified by RF-ABC was Scenario i1-1b (**FigureF5x**, **SupplementaryTableST3x**). NN ABC posterior
parameter estimation procedures were conducted using 100,000 simulations under Scenario i1-1b, each simulation
corresponding to a single vector of parameter values drawn randomly from prior distributions provided in **TableT2x**. We
considered 42 neurons in the hidden layer of the NN and a tolerance level of 0.01, corresponding to the 1,000 simulations
providing summary-statistics closest to the observed ones, for each 25 separate analyses. NN posterior estimates are based on
the logit transformation of parameter values using an Epanechnikov kernel between the corresponding parameter's prior
bounds (see **Material and Methods** and **TableT2x**). Posterior parameter densities are represented with solid blue lines. 50%
Credibility Intervals are represented as the light blue area under the density. The median and mode values are represented as
a solid and dotted blue vertical line, respectively. Parameter prior distributions are represented as dotted grey lines. For all
panels, the left plots represent the NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern
African populations winning under Scenario i1-1b, separately (**SupplementaryTableST3x** and **SupplementaryTableST4x**).
See **FigureF5x** and **TableT2x** for all parameters' descriptions in each panel. Results are also provided in **TableT4x**.
1868
1869
1870

SupplementaryFigureSF7xPart1

SupplementaryFigureSF7xPart2: ABC posterior distribution of potentially asymmetric instantaneous gene-flow intensity parameters.

Neural Network Approximate Bayesian Computation (Csilléry et al. 2012), posterior parameter joint estimations of gene-flow intensity parameters m (in generations before present) for 25 sets of five Central and Southern African populations for which the winning scenario identified by RF-ABC was Scenario i1-1b (**FigureF5x**, **SupplementaryTableST3x**). NN ABC posterior parameter estimation procedures were conducted using 100,000 simulations under Scenario i1-1b, each simulation corresponding to a single vector of parameter values drawn randomly from prior distributions provided in **TableT2x**. We considered 42 neurons in the hidden layer of the NN and a tolerance level of 0.01, corresponding to the 1,000 simulations providing summary-statistics closest to the observed ones, for each 25 separate analyses. NN posterior estimates are based on the logit transformation of parameter values using an Epanechnikov kernel between the corresponding parameter's prior bounds (see **Material and Methods** and **TableT2x**). Posterior parameter densities are represented with solid blue lines. 50% Credibility Intervals are represented as the light blue area under the density. Parameter prior distributions are represented as dotted grey lines. For all panels, the plots represent the NN-ABC posterior parameter distributions for each 25 sets of five Central and Southern African populations winning under Scenario i1-1b, separately (**SupplementaryTableST3x** and **SupplementaryTableST4x**). See **FigureF5x** and **TableT2x** for all parameters' descriptions in each panel. Note that, overall, parameters are relatively little departing from their priors for numerous sets of population combinations among the 25. Also, posterior parameter distributions that are substantially departing from their priors are often highly differing from one another for each parameter. Therefore, we considered these parameters as unsatisfactorily estimated in our analyses and discuss this limitation in the **Discussion** section of the main text.

SupplementaryFigureSF7xPart2

1901 **SupplementaryFigureSF8x: Schematic overview of the sequencing data processing pipeline.**
1902 All template codes with accompanying detailed explanations for all steps are provided in **Material and Methods** and in the
1903 corresponding GitHub repository (https://github.com/Gwennid/africa-wgs-descriptive).

1904

1905

1906   **SupplementaryFigureSF8x**

1907

1908

1909

Acknowledgments

We are grateful to all subjects who participated in this research. The computations were performed at the Swedish National Infrastructure for Computing (SNIC-UPPMAX). Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. We thank the platform Paléogénomique et Génétique Moléculaire (P2GM) of the MNHN at the Musée de l'Homme in Paris for helping to prepare, in part, the Central African samples for sequencing. We thank the Working Group of Indigenous Minorities in Southern Africa (WIMSA) and the South African San Council for their support and facilitating fieldwork. The project was approved by the South African San Council.

Ethical statement

DNA samples from individuals were collected with the subjects' informed consent following the Declaration of Helsinki guidelines and approved by the following Ethics committees: Institut de Recherche pour le Développement (IRD, France); Cameroon Ministry of Scientific Research and Technology; Central African Republic Ministère de l'Enseignement Supérieur et de la Recherche Scientifique; France Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation (Comité de Protection des Personnes DC-2016-2740); French Commission Nationale Informatique et Liberté CNIL (declaration n°1972648); Uganda National Council for Science and Technology; Institutional Review Board of the University of Chicago Biological Sciences Division and the Pritzker School of Medicine University of Chicago Hospitals (protocol number 16986A); Committee for the Protection of Human Subjects of the Trustees of Dartmouth College and the Dartmouth-Hitchcock Medical Center (protocol number 22410); Human Research Ethics Committee (Medical) at the University of the Witwatersrand, Johannesburg (Protocol Numbers: M980553, M050902, M090576, M10270, M180654); Working Group of Indigenous Minorities in Southern Africa (WIMSA); South African San Council (SASC); and the Swedish Ethical Review Authority (Dnr 2019-05174).

Funding

CS is funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 759933) and the Knut and Alice Wallenberg foundation. MJ is funded by the Knut and Alice Wallenberg foundation, and the Vetenskapsrådet/Swedish Research Council (no: 642-2013-8019 and 2018-05537). PV was funded in part by the UMR7206 Eco-anthropology and the French Agence Nationale de la Recherche ANR-METHIS (15-CE32-0009-1).

REFERENCES

Agranat-Tamir, L., Mooney, J.A. and Rosenberg, N.A. (2024) 'Counting the genetic ancestors from source populations in members of an admixed population', *GENETICS*. Edited by J. Novembre, 226(4), p. iyae011. Available at: https://doi.org/10.1093/genetics/iyae011.

Alexander, D.H., Novembre, J. and Lange, K. (2009) 'Fast model-based estimation of ancestry in unrelated individuals', *Genome Research*, 19(9), pp. 1655–1664. Available at: https://doi.org/10.1101/gr.094052.109.

Bader, G.D. *et al.* (2022) 'Rethinking the Middle to Later Stone Age transition in southern Africa - A perspective from the highveld of Eswatini', *Quaternary Science Reviews*, 286, p. 107540. Available at: https://doi.org/10.1016/j.quascirev.2022.107540.

Bartlein, P.J. *et al.* (2011) 'Pollen-based continental climate reconstructions at 6 and 21 ka: a global synthesis', *Climate Dynamics*, 37(3–4), pp. 775–802. Available at: https://doi.org/10.1007/s00382-010-0904-1.

Beaumont, M.A., Zhang, W. and Balding, D.J. (2002) 'Approximate Bayesian computation in population genetics', *Genetics*, 162(4), pp. 2025–2035.

Behr, A.A. *et al.* (2016) 'pong: fast analysis and visualization of latent clusters in population genetic data', *Bioinformatics*, 32(18), pp. 2817–2823. Available at: https://doi.org/10.1093/bioinformatics/btw327.

Bergström, A. *et al.* (2021) 'Origins of modern human ancestry', *Nature*, 590(7845), pp. 229–237. Available at: https://doi.org/10.1038/s41586-021-03244-5.

Beyer, R.M. *et al.* (2021) 'Climatic windows for human migration out of Africa in the past 300,000 years', *Nature Communications*, 12(1), p. 4889. Available at: https://doi.org/10.1038/s41467-021-24779-1.

Blum, M.G.B. and François, O. (2010) 'Non-linear regression models for Approximate Bayesian Computation', *Statistics and Computing*, 20(1), pp. 63–73. Available at: https://doi.org/10.1007/s11222-009-9116-0.

Boitard, S. *et al.* (2016) 'Inferring Population Size History from Large Samples of Genome-Wide Molecular Data - An Approximate Bayesian Computation Approach', *PLOS Genetics*. Edited by M.A. Beaumont, 12(3), p. e1005877. Available at: https://doi.org/10.1371/journal.pgen.1005877.

Bostoen, K. *et al.* (2015) 'Middle to Late Holocene Paleoclimatic Change and the Early Bantu Expansion in the Rain Forests of Western Central Africa', *Current Anthropology*, 56(3), pp. 354–384. Available at: https://doi.org/10.1086/681436.

Bowcock, A.M. *et al.* (1991) 'Drift, admixture, and selection in human evolution: a study with DNA polymorphisms.', *Proceedings of the National Academy of Sciences*, 88(3), pp. 839–843. Available at: https://doi.org/10.1073/pnas.88.3.839.

Breton, G. *et al.* (2014) 'Lactase Persistence Alleles Reveal Partial East African Ancestry of Southern African Khoe Pastoralists', *Current Biology*, 24(8), pp. 852–858. Available at: https://doi.org/10.1016/j.cub.2014.02.041.

Breton, G. *et al.* (2021) 'Comparison of sequencing data processing pipelines and
application to underrepresented African human populations', *BMC Bioinformatics*, 22(1), p.
488. Available at: https://doi.org/10.1186/s12859-021-04407-x.

Breton, G., Fortes-Lima, C. and Schlebusch, C.M. (2021) 'Revisiting the demographic history
of Central African populations from a genetic perspective', *Human Population Genetics and*
Genomics, pp. 1–29. Available at: https://doi.org/10.47248/hpgg2101010004.

Busby, G.B. *et al.* (2016) 'Admixture into and within sub-Saharan Africa', *eLife*, 5, p. e15266.
Available at: https://doi.org/10.7554/eLife.15266.

Chen, J. *et al.* (2020) 'Improved XGBoost model based on genetic algorithm', *International*
Journal of Computer Applications in Technology, 62(3), p. 240. Available at:
https://doi.org/10.1504/IJCAT.2020.106571.

Choudhury, A. *et al.* (2017) 'Whole-genome sequencing for an enhanced understanding of
genetic variation among South Africans', *Nature Communications*, 8(1), p. 2062. Available
at: https://doi.org/10.1038/s41467-017-00663-9.

Cornelissen, E. (2002) '[No title found]', *Journal of World Prehistory*, 16(3), pp. 197–235.
Available at: https://doi.org/10.1023/A:1020949501304.

Csilléry, K., François, O. and Blum, M.G.B. (2012) 'abc: an R package for approximate
Bayesian computation (ABC): *R package: abc*', *Methods in Ecology and Evolution*, 3(3), pp.
475–479. Available at: https://doi.org/10.1111/j.2041-210X.2011.00179.x.

Cuadros-Espinoza, S. *et al.* (2022) 'The genomic signatures of natural selection in admixed
human populations', *The American Journal of Human Genetics*, 109(4), pp. 710–726.
Available at: https://doi.org/10.1016/j.ajhg.2022.02.011.

Danecek, P. *et al.* (2011) 'The variant call format and VCFtools', *Bioinformatics*, 27(15), pp.
2156–2158. Available at: https://doi.org/10.1093/bioinformatics/btr330.

DePristo, M.A. *et al.* (2011) 'A framework for variation discovery and genotyping using next-
generation DNA sequencing data', *Nature Genetics*, 43(5), pp. 491–498. Available at:
https://doi.org/10.1038/ng.806.

Estoup, A. *et al.* (2018) 'Model choice using Approximate Bayesian Computation and
Random Forests: analyses based on model grouping to make inferences about the genetic
history of Pygmy human populations.', *Journal de la Société Française de Statistique*. Tome
159, pp. 167–190. Available at: http://www.numdam.org/item/JSFS_2018__159_3_167_0/.

Excoffier, L. *et al.* (2013) 'Robust Demographic Inference from Genomic and SNP Data',
PLoS Genetics. Edited by J.M. Akey, 9(10), p. e1003905. Available at:
https://doi.org/10.1371/journal.pgen.1003905.

Excoffier, L. and Foll, M. (2011) 'fastsimcoal: a continuous-time coalescent simulator of
genomic diversity under arbitrarily complex evolutionary scenarios', *Bioinformatics*, 27(9),
pp. 1332–1334. Available at: https://doi.org/10.1093/bioinformatics/btr124.

Falush, D., Stephens, M. and Pritchard, J.K. (2003) 'Inference of population structure using
multilocus genotype data: linked loci and correlated allele frequencies', *Genetics*, 164(4), pp.
1567–1587.

Fan, S. *et al.* (2023) 'Whole-genome sequencing reveals a complex African population
demographic history and signatures of local adaptation', *Cell*, 186(5), pp. 923-939.e14.
Available at: https://doi.org/10.1016/j.cell.2023.01.042.

Fenner, J.N. (2005) 'Cross-cultural estimation of the human generation interval for use in
genetics-based population divergence studies', *American Journal of Physical Anthropology*,
128(2), pp. 415–423. Available at: https://doi.org/10.1002/ajpa.20188.

Fortes-Lima, C. *et al.* (2022) 'Demographic and Selection Histories of Populations Across
the Sahel/Savannah Belt', *Molecular Biology and Evolution*. Edited by E. Heyer, 39(10), p.
msac209. Available at: https://doi.org/10.1093/molbev/msac209.

Fortes-Lima, C.A. *et al.* (2021) 'Complex genetic admixture histories reconstructed with
Approximate Bayesian Computation', *Molecular Ecology Resources*, 21(4), pp. 1098–1117.
Available at: https://doi.org/10.1111/1755-0998.13325.

Fortes-Lima, C.A. *et al.* (2024) 'The genetic legacy of the expansion of Bantu-speaking
peoples in Africa', *Nature*, 625(7995), pp. 540–547. Available at:
https://doi.org/10.1038/s41586-023-06770-6.

Gascuel, O. (1997) 'BIONJ: an improved version of the NJ algorithm based on a simple
model of sequence data', *Molecular Biology and Evolution*, 14(7), pp. 685–695. Available at:
https://doi.org/10.1093/oxfordjournals.molbev.a025808.

Gosling, W.D., Scerri, E.M.L. and Kaboth-Bahr, S. (2022) 'The climate and vegetation
backdrop to hominin evolution in Africa', *Philosophical Transactions of the Royal Society B:
Biological Sciences*, 377(1849), p. 20200483. Available at:
https://doi.org/10.1098/rstb.2020.0483.

Gravel, S. (2012) 'Population Genetics Models of Local Ancestry', *Genetics*, 191(2), pp.
607–619. Available at: https://doi.org/10.1534/genetics.112.139808.

Hamid, I. *et al.* (2021) 'Rapid adaptation to malaria facilitated by admixture in the human
population of Cabo Verde', *eLife*, 10, p. e63177. Available at:
https://doi.org/10.7554/eLife.63177.

Harris, K. and Nielsen, R. (2016) 'The Genetic Cost of Neanderthal Introgression', *Genetics*,
203(2), pp. 881–891. Available at: https://doi.org/10.1534/genetics.116.186890.

Henn, B.M., Steele, T.E. and Weaver, T.D. (2018) 'Clarifying distinct models of modern
human origins in Africa', *Current Opinion in Genetics & Development*, 53, pp. 148–156.
Available at: https://doi.org/10.1016/j.gde.2018.10.003.

Hewlett, B.S. (ed.) (2017) *Hunter-gatherers of the Congo Basin: cultures, histories and
biology of African Pygmies*. First published by Transaction Publishers 2014. London New
York: Routledge, Taylor and Francis Group.

Hollfelder, N. *et al.* (2021) 'The deep population history in Africa', *Human Molecular
Genetics*, 30(R1), pp. R2–R10. Available at: https://doi.org/10.1093/hmg/ddab005.

Huang, X. *et al.* (2024) 'Harnessing deep learning for population genetic inference', *Nature
Reviews Genetics*, 25(1), pp. 61–78. Available at: https://doi.org/10.1038/s41576-023-
00636-3.

Hublin, J.-J. *et al.* (2017) 'New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens', *Nature*, 546(7657), pp. 289–292. Available at: https://doi.org/10.1038/nature22336.

Jay, F., Boitard, S. and Austerlitz, F. (2019) 'An ABC Method for Whole-Genome Sequence Data: Inferring Paleolithic and Neolithic Human Expansions', *Molecular Biology and Evolution*. Edited by R. Hernandez, 36(7), pp. 1565–1579. Available at: https://doi.org/10.1093/molbev/msz038.

Kamm, J. *et al.* (2020) 'Efficiently Inferring the Demographic History of Many Populations With Allele Count Data', *Journal of the American Statistical Association*, 115(531), pp. 1472–1487. Available at: https://doi.org/10.1080/01621459.2019.1635482.

Lachance, J. *et al.* (2012) 'Evolutionary History and Adaptation from High-Coverage Whole-Genome Sequences of Diverse African Hunter-Gatherers', *Cell*, 150(3), pp. 457–469. Available at: https://doi.org/10.1016/j.cell.2012.07.009.

Laurent, R. *et al.* (2023) 'A genetic and linguistic analysis of the admixture histories of the islands of Cabo Verde', *eLife*, 12, p. e79827. Available at: https://doi.org/10.7554/eLife.79827.

Lawson, D.J., van Dorp, L. and Falush, D. (2018) 'A tutorial on how not to over-interpret STRUCTURE and ADMIXTURE bar plots', *Nature Communications*, 9(1), p. 3258. Available at: https://doi.org/10.1038/s41467-018-05257-7.

Leung, F.H.F. *et al.* (2003) 'Tuning of the structure and parameters of a neural network using an improved genetic algorithm', *IEEE Transactions on Neural Networks*, 14(1), pp. 79–88. Available at: https://doi.org/10.1109/TNN.2002.804317.

Lézine, A.-M. *et al.* (2019) 'A 90,000-year record of Afromontane forest responses to climate change', *Science*, 363(6423), pp. 177–181. Available at: https://doi.org/10.1126/science.aav6821.

Li, H. and Durbin, R. (2009) 'Fast and accurate short read alignment with Burrows–Wheeler transform', *Bioinformatics*, 25(14), pp. 1754–1760. Available at: https://doi.org/10.1093/bioinformatics/btp324.

Li, J.Z. *et al.* (2008) 'Worldwide Human Relationships Inferred from Genome-Wide Patterns of Variation', *Science*, 319(5866), pp. 1100–1104. Available at: https://doi.org/10.1126/science.1153717.

Lipson, M. *et al.* (2022) 'Ancient DNA and deep population structure in sub-Saharan African foragers', *Nature*, 603(7900), pp. 290–296. Available at: https://doi.org/10.1038/s41586-022-04430-9.

Long, J.C. (1991) 'The genetic structure of admixed populations.', *Genetics*, 127(2), pp. 417–428. Available at: https://doi.org/10.1093/genetics/127.2.417.

Lopez, M. *et al.* (2018) 'The demographic history and mutational load of African hunter-gatherers and farmers', *Nature Ecology & Evolution*, 2(4), pp. 721–730. Available at: https://doi.org/10.1038/s41559-018-0496-4.

Lopez, M. *et al.* (2019) 'Genomic Evidence for Local Adaptation of Hunter-Gatherers to the African Rainforest', *Current Biology*, 29(17), pp. 2926-2935.e4. Available at: https://doi.org/10.1016/j.cub.2019.07.013.

Lorente-Galdos, B. *et al.* (2019) 'Whole-genome sequence analysis of a Pan African set of
samples reveals archaic gene flow from an extinct basal population of modern humans into
sub-Saharan populations', *Genome Biology*, 20(1), p. 77. Available at:
https://doi.org/10.1186/s13059-019-1684-5.

Lucas-Sánchez, M., Serradell, J.M. and Comas, D. (2021) 'Population history of North Africa
based on modern and ancient genomes', *Human Molecular Genetics*, 30(R1), pp. R17–R23.
Available at: https://doi.org/10.1093/hmg/ddaa261.

Mallick, S. *et al.* (2016) 'The Simons Genome Diversity Project: 300 genomes from 142
diverse populations', *Nature*, 538(7624), pp. 201–206. Available at:
https://doi.org/10.1038/nature18964.

Mazet, O. *et al.* (2016) 'On the importance of being structured: instantaneous coalescence
rates and human evolution—lessons for ancestral population size inference?', *Heredity*,
116(4), pp. 362–371. Available at: https://doi.org/10.1038/hdy.2015.104.

McKenna, A. *et al.* (2010) 'The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data', *Genome Research*, 20(9), pp. 1297–1303.
Available at: https://doi.org/10.1101/gr.107524.110.

Mesfin, I., Oslisly, R. and Forestier, H. (2021) 'Technological analysis of the quartz industry
of Maboué 5 – Layer 3 (Lopé National Park, Gabon): Implications for the Late Stone Age
emergence in western Central Africa', *Journal of Archaeological Science: Reports*, 39, p.
103130. Available at: https://doi.org/10.1016/j.jasrep.2021.103130.

Meyer, M. *et al.* (2012) 'A High-Coverage Genome Sequence from an Archaic Denisovan
Individual', *Science*, 338(6104), pp. 222–226. Available at:
https://doi.org/10.1126/science.1224344.

Miller, S.A., Dykes, D.D. and Polesky, H.F. (1988) 'A simple salting out procedure for
extracting DNA from human nucleated cells', *Nucleic Acids Research*, 16(3), pp. 1215–1215.
Available at: https://doi.org/10.1093/nar/16.3.1215.

Mills, R.E. *et al.* (2011) 'Natural genetic variation caused by small insertions and deletions in
the human genome', *Genome Research*, 21(6), pp. 830–839. Available at:
https://doi.org/10.1101/gr.115907.110.

Mooney, J.A. *et al.* (2018) 'Understanding the Hidden Complexity of Latin American
Population Isolates', *The American Journal of Human Genetics*, 103(5), pp. 707–726.
Available at: https://doi.org/10.1016/j.ajhg.2018.09.013.

Mooney, J.A. *et al.* (2023) 'On the number of genealogical ancestors tracing to the source
groups of an admixed population', *GENETICS*. Edited by J. Novembre, 224(3), p. iyad079.
Available at: https://doi.org/10.1093/genetics/iyad079.

Murtagh, F. (1991) 'Multilayer perceptrons for classification and regression',
Neurocomputing, 2(5–6), pp. 183–197. Available at: https://doi.org/10.1016/0925-
2312(91)90023-5.

Nei, M. (1978) 'Estimation of average heterozygosity and genetic distance from a small
number of individuals', *Genetics*, 89(3), pp. 583–590. Available at:
https://doi.org/10.1093/genetics/89.3.583.

Okonechnikov, K., Conesa, A. and García-Alcalde, F. (2016) 'Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data', *Bioinformatics*, 32(2), pp. 292–294. Available at: https://doi.org/10.1093/bioinformatics/btv566.

Patin, E. *et al.* (2009) 'Inferring the Demographic History of African Farmers and Pygmy Hunter–Gatherers Using a Multilocus Resequencing Data Set', *PLoS Genetics*. Edited by A. Di Rienzo, 5(4), p. e1000448. Available at: https://doi.org/10.1371/journal.pgen.1000448.

Patin, E. *et al.* (2014) 'The impact of agricultural emergence on the genetic history of African rainforest hunter-gatherers and agriculturalists', *Nature Communications*, 5(1), p. 3163. Available at: https://doi.org/10.1038/ncomms4163.

Patin, E. *et al.* (2017) 'Dispersals and genetic adaptation of Bantu-speaking populations in Africa and North America', *Science*, 356(6337), pp. 543–546. Available at: https://doi.org/10.1126/science.aal1988.

Pemberton, T.J. *et al.* (2012) 'Genomic Patterns of Homozygosity in Worldwide Human Populations', *The American Journal of Human Genetics*, 91(2), pp. 275–292. Available at: https://doi.org/10.1016/j.ajhg.2012.06.014.

Perry, G.H. *et al.* (2014) 'Adaptive, convergent origins of the pygmy phenotype in African rainforest hunter-gatherers', *Proceedings of the National Academy of Sciences*, 111(35). Available at: https://doi.org/10.1073/pnas.1402875111.

Perry, G.H. and Verdu, P. (2017) 'Genomic perspectives on the history and evolutionary ecology of tropical rainforest occupation by humans', *Quaternary International*, 448, pp. 150–157. Available at: https://doi.org/10.1016/j.quaint.2016.04.038.

Peter, B.M. (2022) 'A geometric relationship of F_2, F_3 and F_4 -statistics with principal component analysis', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 377(1852), p. 20200413. Available at: https://doi.org/10.1098/rstb.2020.0413.

Pfennig, A. *et al.* (2023) 'Evolutionary Genetics and Admixture in African Populations', *Genome Biology and Evolution*. Edited by A. Eyre-Walker, 15(4), p. evad054. Available at: https://doi.org/10.1093/gbe/evad054.

Phillipson, D.W. (2005) *African Archaeology*. 3rd edn. Cambridge University Press. Available at: https://doi.org/10.1017/CBO9780511800313.

Pierron, D. *et al.* (2017) 'Genomic landscape of human diversity across Madagascar', *Proceedings of the National Academy of Sciences*, 114(32). Available at: https://doi.org/10.1073/pnas.1704906114.

Pritchard, J.K. *et al.* (1999) 'Population growth of human Y chromosomes: a study of Y chromosome microsatellites', *Molecular Biology and Evolution*, 16(12), pp. 1791–1798. Available at: https://doi.org/10.1093/oxfordjournals.molbev.a026091.

Pritchard, J.K., Stephens, M. and Donnelly, P. (2000) 'Inference of population structure using multilocus genotype data', *Genetics*, 155(2), pp. 945–959.

Prüfer, K. *et al.* (2014) 'The complete genome sequence of a Neanderthal from the Altai Mountains', *Nature*, 505(7481), pp. 43–49. Available at: https://doi.org/10.1038/nature12886.

Pudlo, P. *et al.* (2016) 'Reliable ABC model choice via random forests', *Bioinformatics*, 32(6), pp. 859–866. Available at: https://doi.org/10.1093/bioinformatics/btv684.

Purcell, S. *et al.* (2007) 'PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses', *The American Journal of Human Genetics*, 81(3), pp. 559–575. Available at: https://doi.org/10.1086/519795.

R Core Team (2015) 'R: A Language and Environment for Statistical Computing'. Vienna, Austria: R Foundation for Statistical Computing. Available at: https://www.R-project.org.

Ragsdale, A.P. *et al.* (2023) 'A weakly structured stem for human origins in Africa', *Nature*, 617(7962), pp. 755–763. Available at: https://doi.org/10.1038/s41586-023-06055-y.

Ragsdale, A.P. and Gravel, S. (2019) 'Models of archaic admixture and recent history from two-locus statistics', *PLOS Genetics*. Edited by J.M. Akey, 15(6), p. e1008204. Available at: https://doi.org/10.1371/journal.pgen.1008204.

Rasmussen, M. *et al.* (2014) 'The genome of a Late Pleistocene human from a Clovis burial site in western Montana', *Nature*, 506(7487), pp. 225–229. Available at: https://doi.org/10.1038/nature13025.

Raynal, L. *et al.* (2019) 'ABC random forests for Bayesian parameter inference', *Bioinformatics*. Edited by O. Stegle, 35(10), pp. 1720–1728. Available at: https://doi.org/10.1093/bioinformatics/bty867.

Richter, D. *et al.* (2017) 'The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age', *Nature*, 546(7657), pp. 293–296. Available at: https://doi.org/10.1038/nature22335.

Robert, C.P., Mengersen, K. and Chen, C. (2010) 'Model choice versus model criticism', *Proceedings of the National Academy of Sciences*, 107(3), pp. E5–E5. Available at: https://doi.org/10.1073/pnas.0911260107.

Rosenberg, N.A. (2002) 'Genetic Structure of Human Populations', *Science*, 298(5602), pp. 2381–2385. Available at: https://doi.org/10.1126/science.1078311.

Saitou, N. and Nei, M. (1987) 'The neighbor-joining method: a new method for reconstructing phylogenetic trees.', *Molecular Biology and Evolution* [Preprint]. Available at: https://doi.org/10.1093/oxfordjournals.molbev.a040454.

Scerri, E.M.L. *et al.* (2018) 'Did Our Species Evolve in Subdivided Populations across Africa, and Why Does It Matter?', *Trends in Ecology & Evolution*, 33(8), pp. 582–594. Available at: https://doi.org/10.1016/j.tree.2018.05.005.

Schlebusch, C. (2010) 'Issues raised by use of ethnic-group names in genome study', *Nature*, 464(7288), pp. 487–487. Available at: https://doi.org/10.1038/464487a.

Schlebusch, C.M. *et al.* (2012) 'Genomic Variation in Seven Khoe-San Groups Reveals Adaptation and Complex African History', *Science*, 338(6105), pp. 374–379. Available at: https://doi.org/10.1126/science.1227721.

Schlebusch, C.M. *et al.* (2020) 'Khoe-San Genomes Reveal Unique Variation and Confirm the Deepest Population Divergence in Homo sapiens', *Molecular Biology and Evolution*. Edited by C. Mulligan, 37(10), pp. 2944–2954. Available at: https://doi.org/10.1093/molbev/msaa140.

Schlebusch, C.M. and Jakobsson, M. (2018) 'Tales of Human Migration, Admixture, and Selection in Africa', *Annual Review of Genomics and Human Genetics*, 19(1), pp. 405–428. Available at: https://doi.org/10.1146/annurev-genom-083117-021759.

Seidensticker, D. *et al.* (2021) 'Population collapse in Congo rainforest from 400 CE urges reassessment of the Bantu Expansion', *Science Advances*, 7(7), p. eabd8352. Available at: https://doi.org/10.1126/sciadv.abd8352.

Semo, A. *et al.* (2020) 'Along the Indian Ocean Coast: Genomic Variation in Mozambique Provides New Insights into the Bantu Expansion', *Molecular Biology and Evolution*. Edited by D. Falush, 37(2), pp. 406–416. Available at: https://doi.org/10.1093/molbev/msz224.

Sengupta, D. *et al.* (2021) 'Genetic substructure and complex demographic history of South African Bantu speakers', *Nature Communications*, 12(1), p. 2080. Available at: https://doi.org/10.1038/s41467-021-22207-y.

Skoglund, P. *et al.* (2017) 'Reconstructing Prehistoric African Population Structure', *Cell*, 171(1), pp. 59-71.e21. Available at: https://doi.org/10.1016/j.cell.2017.08.049.

Skoglund, P. and Mathieson, I. (2018) 'Ancient Genomics of Modern Humans: The First Decade', *Annual Review of Genomics and Human Genetics*, 19(1), pp. 381–404. Available at: https://doi.org/10.1146/annurev-genom-083117-021749.

Stringer, C. (2002) 'Modern human origins: progress and prospects', *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 357(1420), pp. 563–579. Available at: https://doi.org/10.1098/rstb.2001.1057.

Szpiech, Z.A. *et al.* (2019) 'Ancestry-Dependent Enrichment of Deleterious Homozygotes in Runs of Homozygosity', *The American Journal of Human Genetics*, 105(4), pp. 747–762. Available at: https://doi.org/10.1016/j.ajhg.2019.08.011.

Tavaré, S. *et al.* (1997) 'Inferring coalescence times from DNA sequence data', *Genetics*, 145(2), pp. 505–518.

The 1000 Genomes Project Consortium *et al.* (2015) 'A global reference for human genetic variation', *Nature*, 526(7571), pp. 68–74. Available at: https://doi.org/10.1038/nature15393.

Thomas, D.S.G. *et al.* (2022) 'Lacustrine geoarchaeology in the central Kalahari: Implications for Middle Stone Age behaviour and adaptation in dryland conditions', *Quaternary Science Reviews*, 297, p. 107826. Available at: https://doi.org/10.1016/j.quascirev.2022.107826.

Tishkoff, S.A. *et al.* (2009) 'The Genetic Structure and History of Africans and African Americans', *Science*, 324(5930), pp. 1035–1044. Available at: https://doi.org/10.1126/science.1172257.

Van Der Auwera, G.A. *et al.* (2013) 'From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline', *Current Protocols in Bioinformatics*, 43(1). Available at: https://doi.org/10.1002/0471250953.bi1110s43.

Verdu, P. *et al.* (2009) 'Origins and Genetic Diversity of Pygmy Hunter-Gatherers from Western Central Africa', *Current Biology*, 19(4), pp. 312–318. Available at: https://doi.org/10.1016/j.cub.2008.12.049.

Verdu, P. *et al.* (2013) 'Sociocultural Behavior, Sex-Biased Admixture, and Effective
Population Sizes in Central African Pygmies and Non-Pygmies', *Molecular Biology and*
Evolution, 30(4), pp. 918–937. Available at: https://doi.org/10.1093/molbev/mss328.

Verdu, P. and Rosenberg, N.A. (2011) 'A General Mechanistic Model for Admixture Histories
of Hybrid Populations', *Genetics*, 189(4), pp. 1413–1426. Available at:
https://doi.org/10.1534/genetics.111.132787.

Vicente, M. and Schlebusch, C.M. (2020) 'African population history: an ancient DNA
perspective', *Current Opinion in Genetics & Development*, 62, pp. 8–15. Available at:
https://doi.org/10.1016/j.gde.2020.05.008.

Wang, H., Czerminski, R. and Jamieson, A.C. (2021) 'Neural Networks and Deep Learning',
in M. Einhorn et al. (eds) *The Machine Age of Customer Insight*. Emerald Publishing Limited,
pp. 91–101. Available at: https://doi.org/10.1108/978-1-83909-694-520211010.

Yelmen, B. and Jay, F. (2023) 'An Overview of Deep Generative Models in Functional and
Evolutionary Genomics', *Annual Review of Biomedical Data Science*, 6(1), pp. 173–189.
Available at: https://doi.org/10.1146/annurev-biodatasci-020722-115651.

Ziegler, M. *et al.* (2013) 'Development of Middle Stone Age innovation linked to rapid climate
change', *Nature Communications*, 4(1), p. 1905. Available at:
https://doi.org/10.1038/ncomms2897.

2285
2286