Proteostasis modulates gene dosage evolution in antibiotic-resistant bacteria

Chinmaya Jena, Saillesh Chinnaraj, Soham Deolankar and Nishad Matange[#]

Department of Biology, Indian Institute of Science Education and Research, Pashan, Pune
411008 India

[#]nishad@iiserpune.ac.in

Abstract

Evolution of gene expression frequently drives antibiotic resistance in bacteria. We had
previously (Patel and Matange, *eLife*, 2021) shown that in *Escherichia coli*, mutations at the
mgrB locus were beneficial in trimethoprim and led to overexpression of dihydrofolate
reductase (DHFR), encoded by the *folA* gene. Here, we show that DHFR levels are further
enhanced by spontaneous duplication of a genomic segment encompassing *folA* and spanning
hundreds of kilobases. This duplication was rare in wild type *E. coli*. However, its frequency
was elevated in a *lon*-knockout strain, altering the mutational landscape early during
trimethoprim adaptation. We then exploit this system to investigate the relationship between
trimethoprim pressure and *folA* copy number. During long-term evolution, *folA* duplications
were frequently reversed. Reversal was slower under antibiotic pressure, first requiring the
acquisition of point mutations in DHFR or its promoter. Unexpectedly, despite resistance-
conferring point mutations, some populations under high trimethoprim pressure maintained
folA duplication to compensate for low abundance DHFR mutants. We find that evolution of
gene dosage depends on expression demand, which is generated by antibiotic and
exacerbated by proteolysis of drug-resistant mutants of DHFR. We propose a novel role for
proteostasis as a determinant of copy number evolution in antibiotic-resistant bacteria.

Introduction

The dosage of a gene, i.e., copy number relative to the genome, directly impacts its level of expression and can alter organismal phenotype and fitness [1-3]. Mutations that increase copy numbers, termed as gene duplications and amplifications (GDAs), are implicated in diverse biological phenomena including cancer progression [4-6], speciation [7-9], reproductive isolation [10-12] and antimicrobial resistance (AMR) [2, 13]. GDAs are important for rapid adaptation since their frequencies of occurrence are higher by several orders of magnitude (~1000-fold) than point mutations [2, 13-16]. Following the initial GDA mutation, however, amplified genes could be lost over time, pseudogenized, sub-functionalized or neo-functionalized [17-19]. Which of these possibilities is realized depends on the nature and strength of selection imposed by the environment, as well as the fitness cost associated with maintaining multiple gene copies [17-19].

In bacteria, GDA mutations are frequently encountered during adaptation to antibiotics [13] or heavy metals [20, 21]. They occur by RecA-dependent nonequal recombination as well as RecA-independent processes and often involve repeat sequences such as in IS-elements [22-25]. Amplified genes in drug-resistant bacteria code for effectors of resistance such as drug inactivating enzymes, drug target modifiers or "by-pass" mechanisms [26-28], but may also compensate for resistance-associated fitness costs [29]. More recently, the view that GDAs are less "hard-wired" than point mutations has emerged. For instance, transient alterations in gene dosage are a common mechanism of heteroresistance, i.e. resistant sub-populations within a drug-sensitive bacterial isolate [20, 30, 31]. There is also evidence to suggest that copy number variations are more common in bacteria than appreciated. It is estimated that as many as 20% of cells in a bacterial population may harbour a duplication of some genomic region [32]. These copy number polymorphisms create within-population heterogeneities in gene expression, serving as a bet-hedging strategy with adaptive value in fluctuating environments [33].

Dihydrofolate reductase (DHFR) is a ubiquitous enzyme crucial for growth and division. It catalyzes the formation of tetrahydrofolate, an important cofactor in nucleotide and amino acid biosynthesis. DHFR is coded by an essential gene in most organisms and its deletion or pharmacological inhibition results in cell death or growth stasis [34]. Not surprisingly, anti-folates are effective therapeutic agents for diverse pathologies [35, 36]. For instance,

methotrexate, used in the treatment of cancers and autoimmune conditions [37], pyrimethamine, used as an anti-malarial [38], and trimethoprim, used as an anti-bacterial [39], all of which are competitive inhibitors of DHFR. Interestingly, copy number amplification and consequent overexpression of DHFR confers resistance in all three systems [40-43]. Thus, the gene dosage of DHFR appears to be tightly regulated across evolutionarily divergent organisms, making it a useful system to understand the principles underlying selection and evolutionary fate of GDA mutations.

In *Escherichia coli* DHFR is coded by the *folA* gene. Several mutation hot spots for trimethoprim resistance have been identified in this enzyme [44-47]. Additionally, mechanisms that transcriptionally up-regulate *folA* expression are also known in trimethoprim resistant *E. coli*. These include cis-regulatory mutations in the *folA* promoter and trans-acting loss of function mutations in *mgrB*, which upregulate *folA* transcription in a PhoQP-dependent manner [44-46, 48]. Earlier work from our group identified Lon, a conserved quality control protease in bacteria, as a post-translational regulator of trimethoprim resistance. Lon degraded mutant DHFRs and reduced their half-lives *in vivo*, directly affecting genotype-phenotype relationships of resistance-conferring mutations [49, 50]. In this study, we first compare early mutational events in wild type and Lon-deficient *E. coli* challenged with trimethoprim. We show that a large genomic duplication encompassing *folA* confers trimethoprim resistance in both genetic backgrounds, though its frequency of occurrence is elevated in *lon*-knockout *E. coli*. We then use this system to ask what factors determine the fate of duplicated *folA* gene copies during evolution at different trimethoprim pressures.

Results

A large IS-flanked duplication encompassing *folA* mediates rapid adaptation to trimethoprim in *E. coli*

We had previously evolved 5 replicate populations of wild type *E. coli* K-12 MG1655 in sub-MIC trimethoprim (300 ng/mL) for 25 generations. We then isolated and sequenced the genome of a dominant trimethoprim-resistant clone from each population (designated as WTMPR1-5). These sequencing results have been reported earlier, in Patel and Matange (2021) [46]. Briefly, all isolates harboured loss-of-function mutations in the *mgrB* gene (Figure 1 A) that resulted in PhoQP-dependent up-regulation of DHFR expression [46]. Additionally, isolates WTMPR1, 3 and 5 had mutations at the active site of DHFR, while WTMPR4 harboured a large genomic duplication of ~0.6 Mb encompassing the *folA* gene (Figure 1 A, B). We performed similar evolution experiments starting with an isogenic *E. coli* *Δlon* strain to ask if early adaptation to trimethoprim was altered by Lon-deficiency. Among trimethoprim-resistant isolates derived from *E. coli* *Δlon*, designated as LTMPR1-5, four harboured mutations at the *mgrB* locus (Figure 1 A). However, none of them had mutations in DHFR or its promoter. Instead, LTMPR2-4 had evolved GDA mutations similar to WTMPR4, while LTMPR1 had a duplication in an overlapping but shorter genomic stretch (Figure 1 A, B). In all trimethoprim-resistant isolates, duplicated genomic regions were flanked by IS-elements (Figure 1 B). Isolates with and without genomic duplications showed comparable trimethoprim IC50 values (Figure 1 A), indicating that GDA mutations were an alternate genetic path to evolving resistance and occurred more frequently in Lon-deficient *E. coli*.

Seeking to explain the higher frequency of GDA mutations in the *lon*-knockout, we asked whether IS-transposases were sensitive to proteolysis by Lon. Indeed, heterologously expressed InsB and InsL had greater stability in the *lon*-knockout (Figure S1), suggesting that their accumulation may contribute to higher GDA frequency in *E. coli* Δlon. If true it would follow that a higher proportion of GDA mutants would be isolated from *E. coli* Δlon during adaptation to other environments as well. In agreement with this prediction, GDA mutations were detected after evolution of *E. coli* Δlon in 2 of the 5 additional growth conditions tested, namely antibiotic-free and spectinomycin-supplemented media (Figure 1 C, S2). In all cases

duplications were flanked by IS elements (Figure S2). Further, the regions of the genome that
were duplicated were similar among isolates derived from the same environment, but
different across environments (Figure 1 A, S2). On the other hand, wild type *E. coli* did not
show GDA mutations in any of the other tested conditions.

The duplicated genomic region in trimethoprim-resistant isolates included genes previously
associated with resistance to trimethoprim, namely, *folA* which codes for DHFR, and *acrA*
and *B*, which code for components of the AcrAB-TolC efflux pump (Figure 1 B). In one of
the isolates, i.e. LTMPR1, the amplified stretch encompassed *folA* but not *acrAB* without
affecting trimethoprim IC50 values (Figure 1 A, B). We therefore reasoned that duplication of
folA was the main effector of adaptation to trimethoprim. Immunoblotting showed that DHFR
protein was indeed overproduced in isolates LTMPR1 and WTMPR4 over and above the
increase mediated by inactivation of *mgrB* alone (Figure 2 A).

Though DHFR overproduction may contribute to the selection of genomic amplifications in
trimethoprim-resistant bacteria, ~173 (LTMPR1) and ~531 (WTMPR4) other genes were also
duplicated in these isolates. To understand the global transcriptional burden imposed by this
amplification, we analysed the gene expression profiles of WTMPR4 and LTMPR1 using
RNA-sequencing (Supplementary File 1). Since both isolates had loss-of-function mutations
in *mgrB*, which cause large changes in global gene expression [51], we also analysed the
transcriptomes of *E. coli ΔmgrB* and *E. coli ΔlonΔmgrB* strains for comparison
(Supplementary File 1). The level of *folA* mRNA was higher in LTMPR1 (2.3-fold) and
WTMPR4 (5.7-fold) over respective controls (Supplementary File 1), in agreement with
immunoblotting data. Other duplicated genes were mildly overexpressed by ~1.7-fold on an
average in LTMPR1 (over *E. coli ΔlonΔmgrB*) and ~1.3-fold in WTMPR4 (over *E. coli*
ΔmgrB) (Figure 2 B, Supplementary File 1). However, there was large variation in expression
level among duplicated genes in these isolates (Figure 2 B), indicating that increase in gene
dosage did not necessarily result in enhanced expression levels. Instead, global gene
expression profiles of WTMPR4 and LTMPR1 were correlated with *E. coli ΔmgrB* and *E.*
coli ΔlonΔmgrB respectively (Figure 2 C, D) and most of the top 50 up/down-regulated genes
in WTMPR4 and LTMPR1 could be attributed to loss of *mgrB* alone (Figure S3). In line with
this result, strains with and without genomic amplifications had comparable fitness costs (~20
%) in drug-free media, which could also be attributed to inactivation of *mgrB* (Figure 2 E).

Thus, we concluded that large GDA mutations encompassing *folA* led to DHFR
overexpression and represented an alternate mechanism for trimethoprim-resistance in *E. coli*
with undetectable additional cost.

Maintenance of *folA* duplication is linked to antibiotic-generated demand for DHFR
expression

We next investigated the evolutionary fate of duplicate *folA* gene copies by evolving the
LTMPR1 isolate, which harboured the shortest duplication, in the absence of antibiotic. We
also evolved this isolate at sustained drug pressure, i.e. trimethoprim at the same
concentration at which it had originally emerged (300 ng/mL). Using three replicate
populations in each condition, we asked how DHFR expression and trimethoprim IC50
values changed over ~252 generations of evolution and used genome sequencing to identify
the underlying mutational basis. As controls, we performed similar evolution experiments
with WTMPR4 (which harboured a GDA and a functional Lon protease) and WTMPR5
(which harboured a *folA* point mutation and a functional Lon protease).

Evolution in trimethoprim-free media

In drug-free conditions, trimethoprim IC50 of LTMPR1 rapidly declined and a concomitant
loss of colony formation on trimethoprim-supplemented plates occurred (Figure 3 A, B).
Evolved WTMPR4, but not WTMPR5, showed similar loss of colony formation on
trimethoprim indicating that resistance mediated by GDA mutations was far less stable than
point mutations regardless of the presence of Lon (Figure 3 B). For LTMPR1, these trends
were explained by a decline in DHFR expression levels over evolution and reversion to a
single *folA* copy (Figure 3 C, D, S4). Mutation in *mgrB* was, however, consistently
maintained till the end of the experiment (Supplementary File 2). In line with these data,
trimethoprim IC50s of evolving populations stabilized at a value similar to *E. coli*
ΔlonΔmgrB (Figure 3 A). Interestingly, a second GDA mutation ("GDA-2", ~2.2 Mb to 3.3
Mb) emerged in these populations (Figure 3 D, S4). The GDA-2 mutation was apparently lost
and regained multiple times during the experiment (Figure 3 D, S4), suggesting a dynamic
maintenance in the population rather than fixation. Further, the region of the genome
amplified in GDA-2 was similar to what we had observed when *E. coli Δlon* was evolved in

drug-free media (Figure S2, S4), indicating that GDA-2 was likely a fitness enhancing

mutation and unrelated to trimethoprim adaptation.

To further investigate the emergence of GDA-2 during evolution of LTMPR1 in drug-free

media we sequenced 6 randomly picked colonies from each of the lineages at the final time

point (Figure 3 E). All the sequenced isolates had a single copy of *folA* and a majority of

isolates showed the presence of GDA-2, corroborating population sequencing results (Figure

3 E, Supplementary File 2). Interestingly, some of the isolates that did not harbour GDA-2

had evolved mutations in the *rpoS* gene (Figure 3 E, Supplementary File 2). Loss-of-function

mutations in RpoS are fixed at late time points during adaptation to trimethoprim and

enhance the fitness of *E. coli* without conferring resistance [46, 51]. We also sequenced a few

randomly picked isolates from WTMPR4 and WTMPR5 evolved lines (2 from each of the 3

replicate lineages in each case) to check for the presence of GDA-2 and *rpoS* mutations. As

expected, all WTMPR4 isolates had reverted to a single copy of *folA*, while WTMPR5

isolates retained their ancestrally inherited *folA* mutation. None of them had evolved the

GDA-2 mutation, while a few of the isolates from WTMPR4 and WTMPR5 lines had

acquired *rpoS* mutations (Figure 3 F, G).

Consistent with earlier studies, and the role of mutations in *rpoS* in general fitness

enhancement in *E. coli*, we found that an LTMPR1-derived isolate with mutant *rpoS*

recovered fitness in drug-free media (Figure 4 A). Interestingly, an isolate with the GDA-2

mutation had also recovered fitness (Figure 4 A). Coupled with the fact that the GDA-2

mutation occurred only in a *lon*-deficient background, and GDA-2 and *rpoS* mutations were

mutually exclusive we wondered whether the fitness enhancing effects of GDA-2 were also

mediated by reduced RpoS activity. Lon is known to reduce the levels of RpoS during the

exponential growth phase of *E. coli* [52]. Therefore, in a *lon*-knockout we reasoned that at

least some targets of RpoS would be overexpressed. To test this prediction, we analysed the

transcriptome of *E. coli Δlon* and specifically looked at the expression levels of 168 known

targets of RpoS (Supplementary File 3). Of these, 41 genes were overexpressed in the *lon*-

knockout by at least 1.5-fold over wild type, including genes induced in the stationary phase

and by various stresses such as *poxB*, *aidB* and *cbpA* (Figure 4 B, C). Levels of *rpoS* mRNA

were however unaffected in this strain. Next, we compared the expression levels of these 41

RpoS-targets in LTMPR1 and *E. coli Δlon ΔmgrB* with *E. coli Δlon* as the reference and

found that most had similar expression levels in these three strains (Figure 4 C). Indeed a few of these genes showed higher expression levels in LTMPR1 and *E. coli Δlon ΔmgrB*, which was not unexpected since loss of *mgrB* is known stabilize RpoS [53]. Finally, we checked their expression in isolates derived from LTMPR1 which harboured either GDA-2 or *rpoS* mutations. Indeed, in both isolates RpoS target genes were down-regulated indicating a re-setting of RpoS activity (Figure 4 C). Further, in the isolate harbouring GDA-2, *rpoS* mRNA levels were themselves lowered to 0.49x of *E. coli Δlon* (Supplementary File 3). Finally, deletion of *rpoS* from *E. coli Δlon* enhanced the fitness of the strain (Figure 4 D). Thus, evolution of LTMPR1 in antibiotic-free media was characterized by rapid reversal of GDA-mediated resistance and acquisition of mutations that enhanced fitness by reducing the levels of RpoS-transcribed genes (Figure 4 D).

Evolution at constant trimethoprim pressure

In the presence of trimethoprim (300 ng/mL), antibiotic resistance level and DHFR expression remained high over the course of evolution of LTMPR1 (Figure 5 A, B, C). WTMPR4 and WTMPR5 too remained resistant, and in the case of WTMPR4 an increased colony forming efficiency was observed at a late time point (Figure 5 B). These data suggested that under drug pressure high *folA* copy number was likely to be maintained. Indeed, population sequencing of LTMPR1 lineages showed that the ancestral GDA mutation was maintained until 196 generations of evolution in all three replicates (Figure 5 D, S5). By 252 generations, however, a majority of bacteria in Lines A and C had reverted to a single *folA* copy (Figure 5 D, S5) and had acquired a point mutation either in the *folA* gene (Line C) or its promoter (Lines A) (Figure 5 D, Supplementary File 4). Line B, on the other hand, retained high *folA* copy number, but with a reduction in the length of the genomic duplication (Figure 5 D, S5). Line B had also acquired a mutation in the *folA* promoter, however its frequency in the population was low (Figure 5 D, Supplementary File 4).

In agreement with population sequencing, none of the resistant isolates derived from LTMPR1 Line A at the final time point showed the ancestral GDA. Instead, they all harboured a $C_{-35}T$ mutation in the *folA* promoter that is known to up-regulate DHFR expression [51] (Figure 5 E, Supplementary File 4). Among Line B isolates, many had retained the ancestral duplication, without acquiring any additional mutations in *folA*. One of

the isolates from this Line that had reverted to a single *folA* copy had gained the $A_{-64}T$ *folA*

promoter mutation (Figure 5 E, Supplementary File 4). In Line C too we found that isolates

that had lost the ancestral GDA had evolved a point mutation in the *folA* promoter ($+TG_{-30}$) or

a $Trp_{30}Arg$ mutation in DHFR (Figure 5 E, Supplementary File 4). We found similar results in

sequenced isolates from WTMPR4 lineages, i.e. loss of GDA and gain of a point mutation

either in *folA* or its promoter (Figure 5 F). WTMPR5-derived isolates did not accumulate

additional mutations in *folA* but did acquire mutations in *rpoS* (Figure 5 G), showing that at

this drug pressure additional mutations in *folA* were not required. Interestingly, two isolates

from LTMPR1 Lines B and C had evolved new GDA mutations encompassing *folA*. In one of

the Line B isolates (Isolate 2) the ancestral GDA was replaced by 5x amplification of a

shorter genomic region (~12 kb) encompassing *folA* (Figure S6). Both ends of this newly

amplified region were flanked by IS1A (Figure S6). On the other hand, one of the Line C

isolates (Isolate 5) had an expanded duplication beyond what was originally found in

LTMPR1 (Figure S6). Also, both copies of *folA* in this isolate coded for DHFR-$Trp_{30}Arg$

(Figure 5 E, S6), suggesting that the original GDA was lost and an extended GDA mutation

was gained after fixation of the mutant *folA* allele. Thus, even under drug pressure, in most

bacteria GDA mutations were reversed. Notably, reversion to a single *folA* copy was slower

than in drug-free media as it first required the fixation of point mutations at the *folA* locus.

However, in a smaller sub-population, more than one *folA* gene copy was maintained. This

maintenance was dynamic and likely involved repeated loss and gain of GDA mutations.

To further investigate the replacement of GDA by point mutants, we checked to see if under

drug pressure point mutants could invade a population harbouring GDA mutations. We

allowed LTMPR1-derived isolates with point mutations in *folA* (marked with Δ*lacZ::Kan*) to

compete with a GDA-harbouring isolate from the same time point in evolution (to nullify the

effects of other media adaptations) and asked how rapidly the former could invade a

population of the latter. Since for this assay we used point mutant strains genetically marked

with Δ*lacZ::Kan* we verified the neutrality of the Δ*lacZ::Kan* marker by checking if the

marked isolate could invade its unmarked parent. Indeed, in the presence of trimethoprim

(300 ng/mL) point mutants were enriched over the GDA mutant in co-culture by about 100-

fold (Figure 6 A) recreating the results of our evolution experiment. A possible explanation

for this finding may be higher resistance level conferred by *folA* point mutations than GDA.

However, most isolates with *folA* point mutations or harbouring GDAs had similar IC50

values as the ancestral LTMPR1 (Figure 6 B). These isolates also had comparable doubling

times in the presence of 300 ng/mL trimethoprim (Figure 6 B). Thus, though there were no

large observable differences in growth rate or drug IC50 values between strains, point

mutants could indeed invade and outcompete GDA mutations, reflecting either small

differences in fitness between them or their relative stabilities in culture.

Maintenance of high *folA* gene copy number under drug pressure is driven by unstable

mutants of DHFR

Though sustained drug pressure did extend the lifetime of *folA* gene duplication, acquisition

of trimethoprim-resistant point mutations facilitated reversion to a single gene copy. We

therefore asked whether *folA* duplication could be maintained under conditions of increasing

antibiotic pressure, i.e., trimethoprim concentrations starting at 1 μg/mL and ending at 10

μg/mL over 252 generations of evolution (Figure 7 A). Under these conditions, trimethoprim

IC50 values increased sharply over the course of evolution eventually reaching ~40 to ~160-

fold higher than ancestral LTMPR1 (Figure 7 A). Colony formation at high trimethoprim

concentrations also increased at later generations of evolution (Figure 7 B), reflecting the

high drug pressure. Similarly evolved WTMPR4 and WTMPR5 also showed enhanced

colony formation at high trimethoprim concentrations (Figure 7 B).

Curiously, we observed that the 3 replicate lineages of LTMPR1 showed different trends in

DHFR expression. Line A maintained a high level of DHFR protein for the duration of the

experiment. On the other hand, Lines B and C showed high DHFR expression until 196

generations, after which it sharply declined (Figure 7 C). Genome sequencing revealed that

all 3 lines convergently evolved the $Trp_{30}Arg$ mutation in DHFR early during evolution and

had fixed this mutation by 196 generations. All 3 lines also retained *folA* duplication until this

time point (Figure 7 D, S7). Subsequently, Line A evolved an additional $C_{-35}T$ *folA* promoter

mutation and had partially lost the ancestral GDA (Figure 7 D, S7, Supplementary File 5). In

Lines B and C, a second missense mutation in DHFR, i.e. $Tyr_{151}Asp$, appeared between 196

and 252 generations (Figure 7 D, Supplementary File 5). Line B also showed an $Ile_{94}Leu$

mutation in DHFR at low frequency at generation 252 (Figure 7 D, Supplementary File 5).

Unexpectedly, despite acquiring these additional mutational hits and lower DHFR expression

level, bacteria in LTMPR1 Lines B and C retained two copies of *folA* in contrast to Line A (Figure 7 D, S7, Supplementary File 5).

Individual resistant clones from each of the lineages provided greater insight into the genetic trajectories of evolution. In LTMPR1 Line A, 2 of the 6 sequenced isolates had lost gene duplication and acquired the $Trp_{30}Arg$ coding and $C_{-35}T$ promoter mutations in *folA* (Figure 7 E, Supplementary File 5). Interestingly, the other 4 sequenced isolates from Line A retained the ancestral GDA and harboured the same combination of *folA* point mutations, but only in 1 of the 2 gene copies (Figure 7 E, Supplementary File 5). Ten of the twelve isolates from LTMPR1 Lines B and C, on the other hand, retained *folA* duplication. In all ten isolates, both copies of *folA* had acquired $Trp_{30}Arg$ and $Tyr_{151}Asp$ mutations (Figure 7 E, Supplementary File 5). Unlike LTMPR1, all sequenced isolates of WTMPR4 at 252 generations of evolution had lost their GDA mutation and gained 2 *folA* mutations (Figure 7 F). WTMPR5 too had gained an additional *folA* mutation (Figure 7 G). Notably, none of the isolates from WTMPR4 and WTMPR5 lineages had the $Trp_{30}Arg$ and $Tyr_{151}Asp$ combination of mutations that occurred repeatedly in LTMPR1 lineages.

We next sought to understand why some LTMPR1 bacteria had retained high *folA* copy number despite acquiring multiple point mutations in the drug target. To address this question, we first asked what the advantage of *folA* duplication was in the two different mutational trajectories that were observed for LTMPR1. Isolates from Line A had similar IC50 values for trimethoprim regardless of whether they had one or two copies of *folA* (Figure 8 A). On the contrary, isolates from Line B that harboured only one copy of mutant *folA* (coding for DHFR-$Trp_{30}Arg$-$Tyr_{151}Asp$) displayed ~3.5-fold lower IC50 than those that harboured two copies (Figure 8 A). Immunoblotting showed that isolates from Line A had similar expression level of DHFR (Figure 8 B). However, DHFR protein level was significantly compromised by the $Trp_{30}Arg$-$Tyr_{151}Asp$ mutation combination and double the number of *folA* gene copies restored this defect (Figure 8 B). Thus, higher gene dosage of DHFR was dispensable once a promoter up-regulatory mutation was fixed. On the other hand, bacteria with alleles of DHFR harbouring multiple missense mutations continued to benefit from overexpression due to *folA* duplication.

Mutations at position Trp_{30} in DHFR are accompanied by lower *in vivo* proteolytic stability due to perturbation of the hydrophobic core of the protein [54]. This, in turn, affects the

expression level of DHFR [49, 54]. Interestingly, residue Tyr_{151} lies in the same hydrophobic

pocket as Trp_{30} (Figure 8 C). To test the effect of combination of mutations at these sites on

trimethoprim resistance and DHFR expression, we first heterologously expressed wild type

DHFR, $Trp_{30}Arg$ or $Tyr_{151}Asp$ single mutants and the $Trp_{30}Arg$-$Tyr_{151}Asp$ double mutant in

wild type *E. coli*. As expected, DHFR $Trp_{30}Arg$ expression increased the IC50 of

trimethoprim. Surprisingly, neither DHFR $Tyr_{151}Asp$, nor the $Trp_{30}Arg$-$Tyr_{151}Asp$ double

mutant conferred resistance to the antibiotic. Indeed, both mutants conferred lower IC50

values than wild type DHFR. This effect was due to loss of hydrophobicity at the Tyr_{151}

position, as DHFR-$Tyr_{151}Leu$ and DHFR-$Tyr_{151}Phe$ conferred similar IC50 values as wild

type DHFR (Figure 8 D). Since the $Trp_{30}Arg$-$Tyr_{151}Asp$ mutation combination had evolved in

a *lon*-deficient background, we next expressed these mutants in *E. coli Δlon*. Here too, DHFR

$Trp_{30}Arg$ conferred trimethoprim resistance, while the $Tyr_{151}Asp$ mutant did not. Strikingly,

the combination of $Trp_{30}Arg$ and $Tyr_{151}Asp$ conferred very high-level trimethoprim resistance

(Figure 8 D) demonstrating synergistic epistasis that was contingent on the absence of Lon

protease. The levels of expression of DHFR mutants in wild type and *lon*-deficient bacteria

mirrored trimethoprim resistance levels indicating that clearance by Lon altered their

genotype-phenotype relationship (Figure 8 E). Based on these data, we concluded that the

action of proteostatic machinery generated additional demand for expression by clearing

mutant-DHFR proteins. This additional demand drove the maintenance of *folA* duplications

in bacteria evolving at very high drug pressures.

Discussion

Ohno's hypothesis, proposed in 1970, provided a conceptual framework to think about the evolution and eventual fate of duplicated genes [55]. Several theoretical and experimental studies have built upon the original idea to delineate the evolutionary outcomes of gene duplication [18, 56, 57]. More recently, experimental evolution of microbes has been used to test some of the suggestions made by Ohno and others. This approach has the inherent advantage of greater control over the selection pressure applied and available genetic tools to interrogate molecular mechanisms. In most studies of this kind, an artificial system was constructed which was then evolved under defined selection pressures [33, 58-61]. Our study began with serendipitous isolation of spontaneous trimethoprim-resistant mutants of *E. coli* that overproduced DHFR because of a *folA* gene duplication. We exploited this system to understand the relationship between selection pressure and gene duplication in the context of drug-resistance evolution. The selection pressure for higher gene expression, i.e. "expression demand", can be understood as the amount of a gene product that is required to maintain fitness in a particular environment. Antibiotics directly increase expression demand by inhibiting the activity of their targets. Our study shows that increasing gene dosage is a rapid, but transient evolutionary response to ensure that expression demand is met. On the other hand, point mutations are relatively stable routes to mitigating expression demand. Over short time scales, changes in gene dosage can counter the effects of antibiotic. However, their inherent instability results in almost inevitable replacement with phenotypically equivalent point mutations. Thus, continued antibiotic exposure delayed but could not prevent reversion to low gene copy number in most of our experiments. Further, we discover a novel association between proteostasis and gene dosage evolution. In *E. coli* evolving at high trimethoprim pressure, bacterial protein quality control exacerbated expression demand by clearing mutant DHFR-s. As a result, drug-resistant mutants of DHFR with low proteolytic stability drove the maintenance of *folA* gene duplication (Figure 9). Based on these findings we propose that a combination of factors, which include environmental and physiological, alter the relationship between gene copy number and fitness, and drive gene dosage evolution.

A key finding from our study was the two-fold influence that proteostatic machinery had on gene copy number. Firstly, Lon protease-deficiency promoted GDA mutations, possibly by allowing accumulation of IS-tranposases in *E. coli*. This observation resonates with earlier

work by Nicoloff and co-workers who found that large IS-element flanked genomic

duplications in tetracycline-resistant *E. coli* were commonly preceded by spontaneous

mutations that inactivated *lon* [62, 63]. Further, the role of Lon protease in regulating IS-

element mobility by a similar mechanism as proposed by us has been previously

demonstrated for IS1 and IS903, strengthening our argument [64, 65]. We note however that

other mechanisms by which Lon may modulate the frequency of IS-mediated GDAs in *E.*

coli cannot be ruled out. For instance, Lon protease homologs from several organisms

directly bind DNA [66-70], though the role of this binding in DNA transactions is unclear.

Similarly, Lon is a regulator of the recombination protein, RecA, by its action on LexA [71]

and in *Salmonella* RecA stimulated the rate of gene duplications mediated by IS3 [23]. The

second impact of protein quality control on GDA was the action of proteases on

trimethoprim-resistant mutants of DHFR, which promoted gene duplication by generating

additional demand for expression (Figure 9). Importantly, which of these two effects

dominated was determined by a combination of environment, i.e. drug concentrations, and

other mutations acquired at the *folA* locus during evolution. In particular, mutations in the

folA promoter could alleviate expression demand and hence reduce selection for high gene

copy number.

The interplay between GDA and point mutations during evolution agrees in many respects

with a recent study by Tomanek and Guet [61]. They use an elegantly designed artificial

system [33, 61] in which expression of the *galK* gene in *E. coli* was driven by a very low

activity promoter. This strain was then allowed to evolve in different concentrations of

galactose. Like the present study, they found that the activity of IS-elements biased

evolutionary trajectories in favor of duplications. They also showed that co-occurrence of

duplications and point mutations was selected only at high expression demand, in agreement

with our findings. Finally, they argued that gene duplication events may constrain evolution

by point mutations. Once again, our study corroborates these results at the short term in that

the *lon*-knockout which showed a higher propensity for GDA mutations did not evolve point

mutations in *folA* at early time points of evolution. However, given the relative low stability

of large GDAs, these were inevitably lost over longer duration of evolution and replaced with

point mutations. Thus, it is likely that the different mutational profiles of IS-biased

populations may be relevant only at short durations of evolution. It is important to note that

though we have focused on the effects of mutations at the *folA* locus, which are directly under

trimethoprim pressure, many of the isolates from our evolution experiments also harboured

additional mutations at other loci. For instance, mutations in *rpoS* were seen throughout our
evolution lines and at least in drug-free media were significant contributors to bacterial
fitness. It is not yet clear what the role of other mutations are likely to be under antibiotic
selection, though it may be envisaged that these additional mutations play the role of media
adaptations. We did not take their effects into in this study account since we expected them to
be minor contributors to fitness especially during evolution under trimethoprim selection.
However, their role in modulating the stability and selection of GDA mutations in drug-
free/low drug environments may be more significant and remains to be investigated in the
future.
450
451 In summary, the present study uncovers the complex relationship between gene copy number,
452 point mutations and environment, and identifies proteostasis as a regulator of gene dosage
453 evolution in bacteria. These findings open the door to alternative explanations for why gene
454 duplications may be maintained over long periods of evolution and provides insight into how
455 gene expression demand is mitigated by multiple mechanisms in bacteria.
456
457
458

459   **Materials and Methods**
460

Strains, plasmids and culture conditions

Escherichia coli K-12 MG1655 was used as wild type for all experiments. *E. coli* DH10B

was used for plasmid construction and manipulation. *E. coli* strains were cultured in Luria-

Bertani (LB) broth with shaking at 180-200 rpm or on LB agar plates at 37 °C. Ampicillin

was used at a concentration of 100 μg/mL for selection of plasmid-transformed strains.

Trimethoprim was added to growth media at required concentrations as needed. The strains

and plasmids used in this study are listed in Table 1.

Table 1. List of strains and plasmids used in the study

Sr. No.	Strain	Description	Source
1.	*Escherichia coli* K-12 MG1655	Wild type *E. coli*; used as ancestor for evolution experiments and as reference strains for phenotypic and genotypic comparisons	Laboratory stocks
Gene knockouts generated in E. coli K-12 MG1655			
1.	*E. coli ΔmgrB*	Isogenic knockout of *mgrB* gene (*ΔmgrB::Kan*)	[46]
2.	*E. coli ΔlacZ*	Isogenic knockout of *lacZ* gene (*ΔlacZ::Cat*)	[46]
3.	*E. coli Δlon*	Isogenic knockout of *lon* gene (*Δlon::Kan*)	[49]
4.	*E. coli ΔlonΔlacZ*	Double knockout of *lon (Δlon::Kan)* and *lacZ (ΔlacZ::Cat)* genes	[46]
5.	*E. coli ΔmgrBΔlon*	Double knockout of *lon (Δlon::Cat)* and *mgrB (ΔmgrB::Kan)* genes	[46]
6.	*E. coli ΔlonΔrpoS*	Double knockout of *lon (Δlon::Cat)* and *rpoS (ΔrpoS::Kan)* genes	[46]
Strains/Isolates derived from short-term adaptive laboratory evolution			
1.	WTMPR1, WTMPR2, WTMPR3, WTMPR4 and	Trimethoprim-resistant derivatives of *E. coli* wild-type obtained by laboratory	[46] Note: These

	WTMPR5	evolution at 300 ng/ml of trimethoprim for ~25 generation, followed by selection on agar plates supplemented with 1 μg/mL of trimethoprim	strains are labelled as TMPR1-5 in the original publication. They have been renamed WTMPR1-5 to indicate that they were derived after evolution of wild type.
2.	LTMPR1, LTMPR2, LTMPR3, LTMPR4 and LTMPR5	Trimethoprim-resistant derivatives of *E. coli Δlon* obtained by laboratory evolution at 300 ng/ml of trimethoprim for ~25 generation, followed by selection on agar plates supplemented with 1 μg/mL of trimethoprim	This study
Plasmids			
1.	pPRO-*insB*	*InsB* gene cloned into the pPROEx-HtC expression plasmid in frame-with an N-terminal hexa-histidine tag.	This study
2.	pPRO-*insL*	*InsL* gene cloned into the pPROEx-HtA expression plasmid in frame-with an N-terminal hexa-histidine tag.	This study
3.	pPRO-*folA*	*FolA* gene cloned into the pPROEx-HtB expression plasmid in frame-with an N-terminal hexa-histidine tag.	[54]
4.	pPRO-*folA* $Trp_{30}Arg$	*FolA* gene cloned into the pPROEx-HtB expression plasmid, harbouring the $Trp_{30}Arg$ mutation in frame-with an N-terminal hexa-histidine tag.	[54]
5.	pPRO-*folA* $Tyr_{151}Asp$	*FolA* gene cloned into the pPROEx-HtB expression plasmid, harbouring the	This study

		Tyr$_{151}$Asp mutation in frame-with an N-terminal hexa-histidine tag.	
6.	pPRO-*folA* *Trp$_{30}$Arg Tyr$_{151}$Asp*	*FolA* gene cloned into the pPROEx-HtB expression plasmid, harbouring the Trp$_{30}$Arg and Tyr$_{151}$Asp mutations in frame-with an N-terminal hexa-histidine tag.	This study
7.	pPRO-*folA Tyr$_{151}$Phe*	*FolA* gene cloned into the pPROEx-HtB expression plasmid, harbouring the Tyr$_{151}$Phe mutation in frame-with an N-terminal hexa-histidine tag.	This study
8.	pPRO-*folA Tyr$_{151}$Leu*	*FolA* gene cloned into the pPROEx-HtB expression plasmid, harbouring the Tyr$_{151}$Leu mutation in frame-with an N-terminal hexa-histidine tag.	This study

Laboratory evolution of antibiotic resistance

Short-term evolution of antibiotic resistance

For short term laboratory evolution of antibiotic resistance, wild type and *E. coli Δlon* populations were cultured in 96-well plates in LB medium incorporating sub-inhibitory concentrations of different antibiotics. Following concentrations of antibiotics were used: trimethoprim (300 ng/mL; MIC/3); spectinomycin (5 µg/mL; MIC/10); nalidixic acid (4.25 µg/mL; MIC/3); erythromycin (10 µg/mL; MIC/10); rifampicin (4.25 µg/mL; MIC/3). Control populations were also set up in antibiotic-free LB. Five replicates were included for each strain in each antibiotic. Total volume of culture in each well was 150 µL. 10 % of the population was passaged to fresh growth media every 24 hours, for 7 days, resulting in ~25 generations of evolution. At the end of 7 days, the evolved populations were frozen as glycerol stocks.

To check for the evolution of resistance and isolation of resistant clones for further analyses, frozen populations were revived in drug-free LB broth and grown overnight (14-16 hours). Saturated cultures were serially diluted and spotted on LB agar plates supplemented with the appropriate antibiotic at MIC. Colonies were randomly picked, cultured overnight in drug-free LB and frozen in 50% glycerol at -80 °C until further use.

Long-term evolution at different trimethoprim pressures

For long-term evolution, trimethoprim-resistant isolates LTMPR1, WTMPR4 and WTMPR5

were first revived from frozen stocks in drug-free LB overnight. Triplicate evolving lineages

were then established by sub-culturing 1% of the saturated culture of each ancestor into 2 mL

of appropriate medium. For relaxed selection drug-free LB was used. For constant antibiotic

pressure, trimethoprim was added at a concentration of 300 ng/mL. For increasing antibiotic

pressure, trimethoprim was increased from 1 μg/mL to 10 μg/mL over course of the

experiment as shown in Figure 4 A. Cultures were grown for 10-14 hours (~7 generations)

and then passaged (1%) into fresh media. This was continued for 36 passages, i.e. ~252

generations. Aliquots of cultures were frozen periodically for further analyses.

For isolating individual clones, populations were revived from frozen stocks at 252

generations of evolution in drug-free LB, serially diluted and then spotted onto LB agar

plates supplemented with trimethoprim (1, 5 or 10 μg/mL). Plates were incubated for 18-20

hours at 37 °C and resistant colonies were picked randomly for further analyses.

Measuring trimethoprim resistance

Trimethoprim IC50

Trimethoprim Inhibitory Concentration - 50 (IC50), i.e. the antibiotic concentration needed

for 50 % growth inhibition, was calculated using a broth dilution assay. Frozen glycerol

stocks of appropriate bacterial strains were revived in 1-3 mL LB for 6-8 hours. In a 96-well

plate, 150 μL of LB was added to individual wells. Trimethoprim stocks were added, and the

antibiotic was serially diluted. No antibiotic was added to a few wells, which served as

controls. Next, 1.5 μL of the appropriate bacterial culture was added to each well. The

peripheral wells of the plate were filled with water to prevent dehydration. The 96-well plate

was then incubated at 37°C for 15-18 hrs. Optical density at 600 nm was measured for each

well using a plate reader (Ensight, Perkin Elmer). Obtained data were first normalised to

growth in drug-free wells (set to 1) and then fitted to a sigmoidal log(inhibitor)-response

curve using Graphpad Prism software. The equation for this model is as follows:

$Y = Min + (Max-Min)/(1+10^{(X-Log(IC50))})$

where,

Min: Minimum value of normalised OD and Max: maximum value of normalised OD

Values of IC50 were estimated from the above curve fit and averaged across 3 independent

measurements for every strain.

Colony forming efficiency

To calculate the colony forming efficiency of evolving populations at different trimethoprim

concentrations, frozen glycerol stocks were first revived in 1 mL LB for 14-15 hours.

Saturated cultures were serially diluted (10-fold). Diluted cultures were then spotted (10 mL)

on LB agar plates containing 0, 1, 5 and 10 µg/mL of trimethoprim. Spots were allowed to

dry completely and the plates were incubated at 37 °C for ~18 hours. Colonies were counted

from the least dense spot and multiplied by the appropriate dilution factor to determine

colony forming units (CFUs)/mL.

Growth curves

Appropriate frozen stocks were revived overnight in drug-free LB (1-3 mL) and 1µl of the

saturated culture was added to 200 µL of media supplemented with trimethoprim (300 ng/ml)

in the wells of a sterile uncoated 96-well plate. Optical density at 600 nm was measured

during growth at 37 °C at 15-minute intervals over a 24-hour period on a Perkin-Elmer

Ensight Multimode Plate Reader. Doubling time (DT) (in mins) was estimated from log

phase, using the following formula:

$$DT = \ln(2) \, / (\frac{\ln\left(\frac{OD2}{OD1}\right)}{T2 - T1})$$

Where OD1 and OD2 represent optical densities at the onset and end of log phase

respectively, and T1 and T2 represent time elapsed (in mins), corresponding to OD1 and

OD2.

Genome sequencing, identification of mutations and detection of GDA mutations

Genomic DNA was extracted from late log-phase cultures of appropriate strain/resistant

isolates/evolving populations using phenol:chloroform:isoamyl alcohol extraction as

previously described [72]. Extracted genomic DNA was then cleaned-up using HiGenoMB

DNA isolation and purification kit (Himedia, India) as per manufacturer's instructions and

quantitated spectrophotometrically. The integrity was checked by running purified DNA on
an agarose gel before further analyses.

For whole-genome resequencing, paired-end whole-genome next-generation sequencing
(NGS) was performed on a MiSeq system (Illumina, USA) with read lengths of 150-250 bps.
Library preparation and sequencing services were provided by Eurofins (Bangalore, India)
and Genotypic (Bangalore, India). Processed reads were aligned to the reference genome *E.*
coli K-12 MG1655 (U00096) using bowtie2. Variant calling and prediction of new junctions
in the genome to identify large structural mutations was done using Breseq [73] using default
settings in population or clonal modes as needed. All variants and new junctions that were
already present in the ancestor were excluded. Variants unique to the test sample were
classified as "mutations".

For detection of GDA mutations, 'coverage' vs 'coordinate in reference genome' plot
generated by the bam.cov command in Breseq was used [73]. Contiguous stretches greater
than 10 kb with coverage depth of 1.5x or more, compared to the rest of the genome, were
classified as GDA mutations. The presence of IS-elements or repeat sequences was manually
checked using the coordinates of the GDA.

Calculating relative fitness of E. coli strains

Relative fitness of gene knockouts or evolved clones was estimated using competition assays.
Saturated cultures of test strains and reference strains (*E. coli ΔlacZ* or *E. coli ΔlacZΔlon*)
were mixed 1:1 to a total of 1% in 3 mL LB. Strains were allowed to compete for ~7
generations (24 hours incubation). Mixed cultures were diluted and plated on LB agar
supplemented with IPTG (1 mM) and X-gal (50 μg/mL) at the start and end of the
competition experiment. Plates were incubated for 18-24 hrs, and blue and white colonies
were counted to determine the CFUs/mL at initial and final time points.

Relative fitness (w) was calculated using the following formula:

$$w = \ln\left(\frac{Tf}{Ti}\right) \Big/ \ln\left(\frac{Rf}{Ri}\right)$$

580 where T_f and T_i are the final and initial CFUs/mL of the test strain and R_f and R_i are
581 the final and initial CFUs/mL of the reference strain, respectively.

Estimation of DHFR expression by immunoblotting

Immunoblotting was used to determine the expression level of DHFR in various strains/mutants of *E. coli* as described in Patel and Matange (2021) [46]. Briefly, equal lysate protein (5 μg) was subjected to SDS-PAGE on a 15% gel and electroblotted onto PVDF membranes. The membrane was blocked with 5% BSA solution and cut into 2 halves at the 25 kDa molecular weight band. The lower part of the membrane was probed with anti-DHFR polyclonal IgG (100 ng/mL) and the upper part was probed with anti-FtsZ serum (1:50,000). HRP-linked anti-rabbit IgG (1:10,000) was used as the secondary antibody. Hybridised antibodies were detected using chemiluminescence. Band intensities were quantitated in ImageJ and normalised to appropriate control (set to 1). Fold change in DHFR was calculated as follows:

$$Fold\ change_{DHFR} = {Normalised\ intensity_{DHFR}} / {Normalised\ intensity_{FtsZ}}$$

Gene expression analysis using RNA-sequencing

Transcriptome analyses were performed using RNA sequencing. For RNA-seq, saturated overnight culture of appropriate strains/isolates was inoculated (1 %) into 3 mL LB in triplicate and incubated at 37 ºC for 3 hours with shaking at 180 rpm. The cultures were centrifuged and bacterial pellets were resuspended in 1 mL of RNAlater (Invitrogen) for storage. RNA extraction, sequencing and preliminary data analyses were performed by Eurofins (Bangalore, India). Total RNA was extracted using RNAeasy spin columns and quantitated using Qubit and Bioanalyzer. Bacterial rRNA was depleted using Ribozero kit. Bacterial mRNA was then reverse transcribed, library prepared and quality control performed using Tapestation platform. Paired end sequencing was performed on Illumina platform with read lengths of 150 bp read length. Processed reads were aligned to the reference genome (U00096.3) using HISTAT2(version 2.1.0). Abundance estimation was done using featureCounts(version1.34.0). Differential gene expression (DGE) analysis was done by comparing the expression of individual genes in test and reference samples. The output of the DGE analyses were $\log_2$(fold change) values for each gene and P-values for statistical significance.

Cloning and site-directed mutagenesis

Cloning of InsB and InsL transposases

InsB was amplified from *E. coli* K-12 MG1655 genomic DNA by PCR using primers

insB1_fwd_*Nco*I (5'-GGCGCCATGGATATGCCGGGCAACAGCCCG-3')

insB1_rev_*Eco*RI (5'- CCTTTGAATTCTTATTGATAGTGTTTTATG-3'). For amplification

of *insL*, primers insL_*Eco*RI_fwd (5'- ATGCCGAATTCATGAATTACTCTCACGATAAC-

3') and insL_*Hin*dIII_rev (5'- ATGCCAAGCTTTTAGTTCTTCTTTTCGGATCC-3') were

used. Amplified *insB* gene was digested with *Nco*I and *Eco*RI restriction endonuclease and

ligated with similarly digested pPROEx-HtC plasmid. Amplified *insL* was digested with

*Eco*RI and *Hin*dIII restriction endonucleases and ligated with similarly digested pPROEx-

HtA plasmid. Positive clones were screening using restriction digestion. Clones were

transformed into *E. coli* wild type or *Δlon* for further analyses.

Site-directed mutagenesis for DHFR

Site-directed mutagenesis was performed to introduce mutations into the coding sequence of

folA in pPRO-folA as described in [54]. Briefly, PCR was performed using pPRO-folA or

pPRO-folA *Trp$_{30}$Arg* as template and the appropriate forward and reverse mutagenic primers

given below:

folA_*Tyr$_{151}$Asp*_fwd_*Bgl*II:

5'-CTCTCACAGCGATTGCTTTGAGATCTTGGAGCGGCGGGGCC-3'

folA_*Tyr$_{151}$Asp*_rev_*Bgl*II:

5'-CGCCGCTCCAAGATCTCAAAGCAATCGCTGTGAGAGTTCTGCG-3'

folA_*Tyr$_{151}$Leu*_fwd_*Bgl*II:

5'-TCTCACAGCTTATGCTTTGAGATCTTGGAGCGGCGGGGCC-3'

folA_*Tyr$_{151}$Leu*_rev_*Bgl*II:

5'- CGCCGCTCCAAGATCTCAAAGCATAAGCTGTGAGAGTTCTGCG-3'

folA_*Tyr$_{151}$Phe*_fwd_*Bgl*II:

5'- CTCACAGCTTTTGCTTTGAGATCTTGGAGCGGCGGGGCC-3'

folA_*Tyr$_{151}$Phe*_rev_*Bgl*II:

5'- GCCGCTCCAAGATCTCAAAGCAAAAGCTGTGAGAGTTCTGCG-3'

PCR products were digested with *Dpn*I overnight, transformed into chemically competent *E.*

coli DH10B and selected on LB agar plates containing ampicillin. Plasmids habouring the

desired mutations were screened by restriction digestion and confirmed by sequencing.

Sequenced clones were transformed into *E. coli* wild type or *Δlon* for further analyses.

In-cell stability of InsB and InsL

A chloramphenicol-chase assay was used to assess the in-cell stability of heterologously

expressed InsB and InsL transposases as described in Matange (2020) [49] with a few

modifications. Overnight cultures of *E. coli* wild type or *Δlon* harbouring plasmids pPRO-

insB or pPRO-insL were diluted 1% in 5 mL LB supplemented with ampicillin and IPTG

(0.1mM). After 4 hours of growth at 37 °C, 1 mL of culture was pelleted. The cell pellet was

lysed in 100 μL of 1x Laemmli sample loading buffer by heating at 95°C for 10-20 mins.

Chloramphenicol was added to the remaining culture at a final concentration of 25 μg/mL to

stop protein synthesis and the culture was placed back in the incubator. After 15, 45, and 90

minutes 1 mL aliquots of the culture were pelleted and lysed. Lysates were subjected to SDS-

PAGE followed by immunoblotting to detect the levels of InsB and InsL. Anti-Hexahis

monoclonal antibody (Invitrogen) was used as the primary antibody for InsL. Anti-Hexahis

polyclonal antibody (Invitrogen) was used as the primary antibody for InsB. Band intensities

were calculated in ImageJ and normalised to control (set to 1) to calculate fraction of protein

remaining after chloramphenicol treatment.

Acknowledgments

We acknowledge Dr. Manjula Reddy for providing us with the anti-FtsZ polyclonal antibody.

We acknowledge Bruhad Dave for his valuable inputs during discussions at early stages of

this project and Prof. Satyajit Rath, Prof. Sandhya Visweswariah and Dr. Girish Deshpande

for their comments on the manuscript. Funding for this work was provided by

DBT/Wellcome Trust India Alliance. NM is a recipient of the India Alliance Intermediate

Fellowship. CJ is a recipient of a Senior Research Fellowship from the CSIR, Govt. of India.

Supplementary Files.

Supplementary File 1. Gene expression changes in LTMPR1 and WTMPR4 isolates

Supplementary File 2. Genomic changes associated with evolution of LTMPR1 in drug-free

media

681 **Supplementary File 3.** Gene expression changes in isolates derived from LTMPR1 evolution

682 in drug-free media containing mutation in *rpoS* or GDA-2

683 **Supplementary File 4.** Genomic changes associated with evolution of LTMPR1 in

684 trimethoprim (300 ng/mL)

685 **Supplementary File 5.** Genomic changes associated with evolution of LTMPR1 in

686 increasing trimethoprim pressure

687 **Supplementary File 6.** Genome sequencing of isolates derived from WTMPR4 evolution at

688 different trimethoprim pressures

689 **Supplementary File 7.** Genome sequencing of isolates derived from WTMPR5 evolution at

690 different trimethoprim pressures

Figures

Figure 1. Adaptation to antibiotics by large genomic duplications is favoured in *lon*-deficient *E. coli*. A. Coverage depth plots and mutations at the *folA* and *mgrB* loci of 5 independently evolved trimethoprim-resistant isolates derived from wild type (WTMPR1-5) or *Δlon E. coli* (LTMPR1-5). Resistant clones were isolated after ~25 generations of evolution at 300 ng/mL of trimethoprim. Coverage depth plots show the number of reads from Illumina short-read sequencing (y-axis) mapped to each genomic coordinate of the reference genome (x-axis). Coverage of 1x, 2x and 3x are marked by dotted lines for reference. Mutations in *mgrB* and *folA,* and trimethoprim IC50 values (mean ± S.D. from 3 independent measurements) are provided for each isolate below the appropriate coverage depth plot. **B.** Cartoon (not to scale) showing the duplicated genomic stretch and flanking IS-elements in trimethoprim-resistant isolates. Blue arrows represent IS1, while red arrows represent IS186. The positions of *folA* and *acrAB* genes, implicated in trimethoprim resistance in earlier studies, relative to the duplicated stretch are shown. Fold increase in the

number of reads corresponding to the *folA* gene are indicated for each isolate. **C.** Summary of the prevalence of GDA mutations detected in wild type or *E. coli Δlon* after ~25 generations of evolution in control media (no antibiotic) or in media supplemented with sub-MIC antibiotics. Each circle represents one of the 5 replicates in the evolution experiment. Light grey circles represent replicates in which no resistance was detected and hence were not sequenced. Dark grey circles indicate that the whole population or a resistant isolate was sequenced but no GDA mutation was detected. Red circles indicate that a GDA mutation was identified.

Figure 2. Overexpression of DHFR in trimethoprim-resistant isolates with genomic duplication. A. DHFR protein expression in trimethoprim-resistant isolates WTMPR4 and LTMPR1, compared with respective ancestors and *mgrB*-knockout strains. DHFR protein

was detected by immunoblotting using an anti-DHFR polyclonal antibody. FtsZ was used as loading control. A representative immunoblot from 3 biological replicates is shown. Quantitation was performed by calculating band intensities using image analysis. Fold change values of DHFR levels over controls (wild type or *E. coli Δlon*, set to 1) is shown as mean ± S.D. from triplicate measurements. **B.** Expression level of duplicated genes in LTMPR1 and WTMPR4, determined by RNA-sequencing, expressed as fold over *E. coli ΔlonΔmgrB* and *E. coli ΔmgrB* respectively. Each point represents fold change value for a single gene. Mean of the scatter is shown as a black line, and its value is provided along with standard deviation above the plot. **C, D.** Correlation between expression levels of genes in LTMPR1 (C) and WTMPR4 (D) with *E. coli ΔlonΔmgrB* and *E. coli ΔmgrB* respectively. Gene expression levels were estimated using RNA-sequencing and fold changes were calculated using wild type *E. coli* as reference for WTMPR4 and *E. coli ΔmgrB*, and *E. coli Δlon* as reference for LTMPR1 and *E. coli ΔlonΔmgrB*. Each point represents $\log_2$(fold change) value for a single gene. The resulting scatter was fit to a simple linear regression and the obtained R^2 values are provided. **E.** Relative fitness (w) of indicated *E. coli* strains calculated using competition experiments with *E. coli ΔlacZ* as reference. For strains derived from *E. coli Δlon, E. coli ΔlonΔlacZ* was used as reference. Neutrality of *ΔlacZ* was established by competition with the unmarked ancestral strains, i.e. wild type (WT) and *E. coli Δlon*. No change in relative fitness compared to the reference strain (w = 1) is indicated by a dotted line. Mean ± S.D. from three independent measurements is plotted.

Figure 3. Evolutionary fate of *folA*-duplication in the absence of drug pressure. A.
Trimethoprim IC50 values of 3 evolving lineages derived from LTMPR1 (A, B, C) over 252
generations of evolution in antibiotic-free medium. Mean ± S.D. from three measurements is
plotted at each time point. IC50 values of *E. coli Δlon* and *ΔlonΔmgrB* are shown as dotted
lines for reference. **B.** Colony formation of LTMPR1, WTMPR4 and WTMPR5 evolved in
antibiotic-free medium at different generations. Fraction of the population capable of forming

colonies at 1, 5 and 10 µg/mL trimethoprim was calculated across 3 replicate lines. Mean

value from the three lines is plotted. The results of similar experiments performed on

ancestors (0 generations) are also provided for reference. **C.** DHFR expression during

evolution in the absence of trimethoprim in LTMPR1 lineages A, B and C, measured by

immunoblotting using anti-DHFR polyclonal antibody. FtsZ was used as a loading control.

Quantitation was performed by calculating band intensities using image analysis. DHFR

expression at each time point was normalized to the ancestor (i.e. 0 generations, set to 1).

Mean of three independent measurements is shown below each lane. **D.** Copy number of the

ancestral GDA encompassing *folA* (GDA(*folA*)) and GDA-2 at different time points of

evolution in the absence of trimethoprim. For GDA(*folA*), copy number was determined by

dividing number of reads from an Illumina sequencing experiment corresponding to *folA* by

the average number of reads mapping to the rest of the genome. **E.** Point mutations in *folA*,

rpoS and *mgrB*, and the "GDA-2" genomic duplication in 6 randomly picked colonies from

each of the LTMPR1 lines at 252 generations of evolution are shown schematically using the

appropriate symbols. Asterix (*/**) marks the genotypes that were carried forward for further

analyses. **F, G.** Point mutations in *folA*, *mgrB* and *rpoS* in isolates derived from 252

generations of evolution of WTMPR4 (F) and WTMPR5 (G) in no antibiotic. From each of

the evolving lines 2 random isolates were picked for genome sequencing. Various point

mutants at the 3 gene loci of interest are represented by appropriate symbols as shown.

Figure 4. Fitness enhancement of LTMPR1 in drug-free media results from lower RpoS

activity. A. Relative fitness (w) of isolates with the indicated genotypes derived from

LTMPR1 evolution in antibiotic-free media (see also Figure 3) calculated using a competitive

growth assay using *E. coli ΔlonΔlacZ* as the reference strain. Mean ± S.D. from three

independent measurements is plotted. No change in relative fitness compared to the reference

(w = 1) is shown as a dotted line for reference. **B.** Expression level of 168 known targets of

RpoS in *E. coli Δlon* compared to wild type expressed as $\log_2$(fold change) values determined

by RNA sequencing. Dotted lines at 1.5-fold higher and lower than wild type represent cut-

offs used to identify overexpressed and under-expressed genes. The pie chart shows the

fraction of RpoS targets that were overexpressed in *E. coli Δlon* by at least 1.5-fold in green.

C. Expression levels of 41 RpoS targets in *E. coli Δlon* compared to wild type are shown as a

heat map in the first vertical. The expression levels of these genes in LTMPR1, *E. coli*

ΔlonΔmgrB or LTMPR1-derived isolates compared to *E. coli Δlon* are provided in the

subsequent verticals. **D.** Relative fitness of *E. coli ΔlonΔrpoS* in antibiotic-free media

calculated using a competitive growth assay using *E. coli ΔlonΔlacZ* as the reference strain.

Neutrality of the *ΔlacZ* genetic marker was verified by competition between *E. coli Δlon* and

E. coli ΔlonΔlacZ. Mean ± S.D. from three independent measurements is plotted. No change

in relative fitness compared to the reference (w = 1) is shown as a dotted line for reference. **E.**

Model for the effects of Lon deficiency on RpoS levels and bacterial fitness. The roles of

mutations within RpoS or the GDA-2 mutations in compensating for the defects of RpoS

overproduction are shown.

Figure 5. Evolutionary fate of *folA*-encompassing GDA in LTMPR1 at sustained trimethoprim pressure. A. Trimethoprim IC50 values of 3 evolving lineages starting from LTMPR1 (A, B, C) over 252 generations of evolution at 300 ng/mL trimethoprim. Mean ± S.D. from three measurements is plotted at each time point. IC50 values of *E. coli Δlon* and *ΔlonΔmgrB* are shown as dotted lines for reference. **B.** Colony formation of LTMPR1, WTMPR4 and WTMPR5 evolved in 300 ng/mL trimethoprim at indicated time points.

Fraction of the population capable of forming colonies at 1, 5 and 10 μg/mL trimethoprim was calculated across 3 replicate lines. Mean value from the three lines is plotted. The results of similar experiments performed on ancestors (0 generations) are also provided for reference. **C.** DHFR expression in LTMPR1 lineages A, B and C evolved in 300 ng/mL trimethoprim, measured by immunoblotting using anti-DHFR polyclonal antibody. FtsZ was used as a loading control. Quantitation was performed by calculating band intensities using image analysis. DHFR expression at each time point was normalized to the ancestor (i.e. 0 generations, set to 1). Mean of three independent measurements is shown below each lane. **D.** Copy number of *folA* (GDA(*folA*)) at different time points of evolution of LTMPR1 in trimethoprim is plotted on the left y-axis. Copy number was determined by dividing number of reads from an Illumina sequencing experiment corresponding to *folA* by the average number of reads mapping to the rest of the genome. Frequency of various *folA* alleles in the evolving populations at each of the time points is also plotted on the right y-axis. **E.** Point mutations in *folA*, *rpoS* and *mgrB* in 6 randomly picked trimethoprim-resistant colonies from each of the LTMPR1 lines at 252 generations of evolution are shown diagrammatically. Genotypes that were carried forward for further analysis are marked (* or #). **F, G.** Point mutations in *folA*, *mgrB* and *rpoS* in isolates derived from 252 generations of evolution of WTMPR4 (F) and WTMPR5 (G) in 300 ng/mL of trimethoprim. From each of the evolving lineages 2 random isolates were picked for genome sequencing. Various point mutants at the 3 gene loci are represented by appropriate symbols as shown.

Figure 6. Invasion of GDA-dominant populations by point mutations under
trimethoprim pressure. A. Upper panel. Competition between isolates with point mutations
in *folA* and an isolate that harboured a *folA* duplication derived from the same time point of
LTMPR1 evolution (upper panel). The genotypes of the isolates used are shown

diagrammatically (see also Figure 5, Supplementary File 4). The initial mixing ratio was

1000:1 in favour of the GDA mutant. Point mutants were marked genetically with

ΔlacZ::Cat. Neutrality of the *ΔlacZ::Cat* marker was verified by competing marked and

unmarked point mutants (lower panel). Percentage of the population constituted by the

marked point mutants over ~105 generations of serial transfer in media supplemented with

trimethoprim (300 ng/mL) is plotted. Traces for individual mutants are shown in grey.

Average of the 4 point mutants is shown in red. **B.** Trimethoprim IC50 values of the LTMPR1

ancestor or evolved isolates from 252 generations in trimethoprim with indicated genotypes

(see also Figure 5, Supplementary File 4) are plotted as bars. Mean ± S.D. from three

independent measurements are plotted. Doubling times relative to LTMPR1 (set to 1) in the

presence of 300 ng/mL of trimethoprim are provided above the graph as mean ± S.D. from

three independent measurements.

Figure 7. Evolutionary fate of *folA*-encompassing GDA in LTMPR1 at increasing trimethoprim pressure. A. Trimethoprim IC50 values of 3 evolving lineages starting from LTMPR1 (A, B, C) over 252 generations of evolution are plotted on the left Y-axis. Mean ± S.D. from three measurements is plotted at each time point. Trimethoprim concentrations used during evolution are plotted in the right Y-axis. **B.** Colony formation of LTMPR1,

WTMPR4 and WTMPR5 evolved in increasing trimethoprim at indicated time points.

Fraction of the population capable of forming colonies at 1, 5 and 10 μg/mL trimethoprim

was calculated across 3 replicate lines. Mean value from the three lines is plotted. The results

of similar experiments performed on ancestors (0 generations) are also provided for

reference. **C.** DHFR expression in LTMPR1 lineages A, B and C evolved in increasing

trimethoprim, measured by immunoblotting using anti-DHFR polyclonal antibody. FtsZ was

used as a loading control. Quantitation was performed by calculating band intensities using

image analysis. DHFR expression at each time point was normalized to the ancestor (i.e. 0

generations, set to 1). Mean of three independent measurements is shown below each lane. **D.**

Copy number of *folA* (GDA(*folA*)) at different time points of LTMPR1 evolution in

increasing trimethoprim is plotted on the left y-axis. Copy number was determined by

dividing number of reads from an Illumina sequencing experiment corresponding to *folA* by

the average number of reads mapping to the rest of the genome. Frequency of various *folA*

alleles in the evolving LTMPR1 populations at each of the time points is shown on the right

y-axis. **E.** Point mutations in *folA*, *rpoS* and *mgrB* in 6 randomly-picked trimethoprim-

resistant colonies from each of the LTMPR1 lines at 252 generations of evolution are shown

schematically. **F, G.** Point mutations in *folA*, *mgrB* and *rpoS* in isolates derived from 252

generations of evolution of WTMPR4 (F) and WTMPR5 (G) in increasing trimethoprim.

From each of the evolving lineages 2 random isolates were picked for genome sequencing.

Various point mutants at the 3 gene loci are represented by appropriate symbols as shown.

Figure 8. Proteostatic pressure facilitates maintenance of *folA* duplication A.
Trimethoprim-IC50 values of resistant isolates with the indicated genotypes (see also Figure
7) derived from LTMPR1 evolution in increasing antibiotic pressure. Mean ± S.D. from three
independent measurements is plotted. **B.** DHFR expression level in LTMPR1-derived
trimethoprim resistant isolates with indicated genotypes assessed by immunoblotting using
anti-DHFR polyclonal antibody. FtsZ was used as a loading control. Quantitation was
performed by calculating band intensities using image analysis. DHFR expression was
normalized to the ancestor (LTMPR1). Mean of three independent measurements is provided.
C. Cartoon representation of the structure of *E. coli* DHFR (PDB: 7DFR) bound to folate.
Residues Trp_{30}, Tyr_{151}, Phe_{153} and Ile_{155} which form hydrophobic interactions and are
required for proteolytic stability are shown as sticks and coloured by element (C: green,
O:red, N:blue). Distances of less than or equal to 4 Å, which indicates possible interactions,
are shown as dotted yellow traces. Folate bound in the active site of DHFR is shown as
sticks. **D.** Trimethoprim IC50 values of *E. coli* wild type or *E. coli* *Δlon* heterologously
expressing DHFR (wt) or its mutants. Mean ± S.D. from three independent measurements is
plotted. **E.** Expression level of plasmid-borne DHFR (wt) or its mutants in wild type or *Δlon*

E. coli assessed by immunoblotting using anti-DHFR polyclonal antibody. FtsZ was used as a

loading control. Quantitation was performed by calculating band intensities using image

analysis. Expression of mutants was normalized to wild type DHFR (set to 1). Mean of three

independent measurements is shown.

Figure 9. Model of expression demand induced selection of *folA* gene duplication,

followed by replacement by phenotypically equivalent point mutants. Expression demand

for DHFR is generated by trimethoprim pressure, which results in the selection of *folA* gene

duplications. Gene duplication is unstable and reverses to a single copy of wild type *folA*

when drug-pressure is withdrawn. Under drug pressure, *folA* gene duplication is maintained

until a point mutation that confers resistance arises in one of the copies of *folA* (promoter or

coding region). Upon acquisition of a point mutation, reversal of gene duplication ensues

unless additional expression demand is generated by the action of proteostatic machinery on

unstable drug-resistant DHFR mutants.

Supplementary Figures

 Figure S1. Stability of heterologously expressed InsB and InsL transposases in wild type or

 Δlon E. coli measured using a chloramphenicol (CMP) chase assay. Levels of the plasmid-

 expressed transposase were measured using immunoblotting with an anti-His

 monoclonal/polyclonal antibody in lysates prepared from cells at indicated time points

 following treatment with 50 μg/mL of CMP. A representative image from three independent

 experiments is shown. Band intensities were quantitated by image analyses and normalized to

 0' (set to 1). The fraction of protein remaining is plotted in blue for wild type and red for *E.*

 coli Δlon. Values from 3 independent experiments are plotted as mean ± S.D.

Representative coverage plots of *E. coli Δlon* evolved in no antibiotic or spectinomycin

910   **Figure S2.** Coverage plots and flanking IS elements for *E. coli Δlon* populations evolving in

911   no antibiotic or spectinomycin that showed GDA mutations. For no-antibiotic, only one of the

912   5 replicates is shown as similar coverage plots were obtained for all 5 replicates.

Figure S3. Gene expression changes associated with *folA*-encompassing duplication.

Heat map comparing the top 50 up/downregulated genes in LTMPR1 and WTMPR4 with *E.*

coli ΔlonΔmgrB and *E. coli ΔmgrB* respectively. Log$_2$(Fold change) values were calculated

based on RNA-sequencing with respect to wild type or *Δlon E. coli* and represented on a

green/red scale as indicated. Gene names are shown adjacent to the appropriate cell. Genes

are ranked according to their fold change in LTMPR1 and WTMPR4.

Figure S4. Coverage depth plots for population sequencing at 105, 196 and 252 generations

of the 3 lineages of LTMPR1 (A, B, C) evolving in trimethoprim-free media. Coverage depth

plots show the number of reads from Illumina sequencing (y-axis) mapped to each genomic

coordinate of the reference genome (x-axis). Coverage of 1x, 2x and 3x are marked by dotted

lines for reference.

Figure S5. Coverage depth plots for population sequencing at 105, 196 and 252 generations of the 3 lineages of LTMPR1 (A, B, C) evolving in trimethoprim-supplemented (300 ng/mL) media. Coverage depth plots show the number of reads from Illumina sequencing (y-axis mapped) to each genomic coordinate of the reference genome (x-axis). Coverage of 1x, 2x and 3x are marked by dotted lines for reference.

Figure S6. Coverage plots for Line C Isolate 5 and Line B Isolate 2. Line B Isolate 3 showed 5x amplification of a shorter genomic stretch encompassing *folA* which is shown diagrammatically next to the coverage plot. Line C Isolate 5 showed an expanded GDA encompassing *folA* compared to the LTMPR1 ancestor, the coordinates for which are below the coverage plot.

944 **Figure S7.** Coverage depth plots for population sequencing at 105, 196 and 252 generations

945 of the 3 lineages of LTMPR1 (A, B, C) evolving in increasing trimethoprim concentrations.

946 Coverage depth plots show the number of reads from Illumina sequencing (y-axis) mapped to

947 each genomic coordinate of the reference genome (x-axis). Coverage of 1x, 2x and 3x are

948 marked by dotted lines for reference.

References

1.   Andersson DI, Hughes D. Gene Amplification and Adaptive Evolution in Bacteria. Annu Rev Genet. 2009;43:167-95. doi: 10.1146/annurev-genet-102108-134805. PubMed PMID: WOS:000273580300008.

2.   Elliott KT, Cuff LE, Neidle EL. Copy number change: evolving views on gene amplification. Future Microbiol. 2013;8(7):887-99. doi: 10.2217/fmb.13.53. PubMed PMID: WOS:000325822800015.

3.   Kondrashov FA. Gene duplication as a mechanism of genomic adaptation to a changing environment. P Roy Soc B-Biol Sci. 2012;279(1749):5048-57. doi: 10.1098/rspb.2012.1108. PubMed PMID: WOS:000310999000025.

4.   Dürrbaum M, Storchová Z. Effects of aneuploidy on gene expression: implications for cancer. Febs J. 2016;283(5):791-802. doi: 10.1111/febs.13591. PubMed PMID: WOS:000372005600002.

5.   Jiang Z, Li CX, Li FY, Wang XS. EGFR Gene Copy Number as a Prognostic Marker in Colorectal Cancer Patients Treated with Cetuximab or Panitumumab: A Systematic Review and Meta Analysis. Plos One. 2013;8(2). doi: ARTN e56205
10.1371/journal.pone.0056205. PubMed PMID: WOS:000315159200024.

6.   Spielmann M, Lupiáñez DG, Mundlos S. Structural variation in the 3D genome. Nat Rev Genet. 2018;19(7):453-67. doi: 10.1038/s41576-018-0007-0. PubMed PMID: WOS:000435393700009.

7.   Pascual-Anaya J, D'Aniello S, Kuratani S, Garcia-Fernàndez J. Evolution of gene clusters in deuterostomes. Bmc Dev Biol. 2013;13. doi: Artn 26
10.1186/1471-213x-13-26. PubMed PMID: WOS:000321576400001.

8.   Taylor JS, Van de Peer Y, Meyer A. Genome duplication, divergent resolution and speciation. Trends Genet. 2001;17(6):299-301. doi: Doi 10.1016/S0168-9525(01)02318-6. PubMed PMID: WOS:000170644200002.

9.   Ting CT, Tsaur SC, Sun S, Browne WE, Chen YC, Patel NH, et al. Gene duplication and speciation in *Drosophila*: Evidence from the *Odysseus* locus. P Natl Acad Sci USA. 2004;101(33):12232-5. doi: 10.1073/pnas.0401975101. PubMed PMID: WOS:000223410100059.

10.   Jiao WB, Patel V, Klasen J, Liu F, Pecinkova P, Ferrand M, et al. The Evolutionary Dynamics of Genetic Incompatibilities Introduced by Duplicated Genes in *Arabidopsis thaliana*. Mol Biol Evol. 2021;38(4):1225-40. doi: 10.1093/molbev/msaa306. PubMed PMID: 33247726; PubMed Central PMCID: PMCPMC8042742.

11. Maclean CJ, Greig D. Reciprocal Gene Loss Following Experimental Whole-Genome
Duplication Causes Reproductive Isolation in Yeast. Evolution. 2011;65(4):932-45. doi:
10.1111/j.1558-5646.2010.01171.x. PubMed PMID: WOS:000289158300003.

12. Zuellig MP, Sweigart AL. Gene duplicates cause hybrid lethality between sympatric
species of *Mimulus*. Plos Genet. 2018;14(4). doi: ARTN e1007130
10.1371/journal.pgen.1007130. PubMed PMID: WOS:000431115700001.

13. Sandegren L, Andersson DI. Bacterial gene amplification: implications for the
evolution of antibiotic resistance. Nat Rev Microbiol. 2009;7(8):578-88. doi:
10.1038/nrmicro2174. PubMed PMID: 19609259.

14. Anderson P, Roth J. Spontaneous tandem genetic duplications in Salmonella
typhimurium arise by unequal recombination between rRNA (rrn) cistrons. Proc Natl Acad
Sci U S A. 1981;78(5):3113-7. doi: 10.1073/pnas.78.5.3113. PubMed PMID: 6789329;
PubMed Central PMCID: PMCPMC319510.

15. Seaton SC, Elliott KT, Cuff LE, Laniohan NS, Patel PR, Neidle EL. Genome-wide
selection for increased copy number in *Acinetobacter baylyi* ADP1: locus and context-
dependent variation in gene amplification. Mol Microbiol. 2012;83(3):520-35. Epub
20111229. doi: 10.1111/j.1365-2958.2011.07945.x. PubMed PMID: 22211470.

16. Sonti RV, Roth JR. Role of gene duplications in the adaptation of *Salmonella*
typhimurium to growth on limiting carbon sources. Genetics. 1989;123(1):19-28. doi:
10.1093/genetics/123.1.19. PubMed PMID: 2680755; PubMed Central PMCID:
PMCPMC1203782.

17. Hahn MW. Distinguishing Among Evolutionary Models for the Maintenance of Gene
Duplicates. J Hered. 2009;100(5):605-17. doi: 10.1093/jhered/esp047. PubMed PMID:
WOS:000269206800012.

18. Innan H, Kondrashov F. The evolution of gene duplications: classifying and
distinguishing between models. Nat Rev Genet. 2010;11(2):97-108. doi: 10.1038/nrg2689.
PubMed PMID: WOS:000273688200004.

19. Kuzmin E, Taylor JS, Boone C. Retention of duplicated genes in evolution. Trends
Genet. 2022;38(1):59-72. doi: 10.1016/j.tig.2021.06.016. PubMed PMID:
WOS:000732466400007.

20. Hufnagel DA, Choby JE, Hao S, Johnson AF, Burd EM, Langelier C, et al. Antibiotic-
Selected Gene Amplification Heightens Metal Resistance. mBio. 2021;12(1). Epub
20210119. doi: 10.1128/mBio.02994-20. PubMed PMID: 33468696; PubMed Central
PMCID: PMCPMC8545094.

21.   von Rozycki T, Nies DH. *Cupriavidus metallidurans*: evolution of a metal-resistant

bacterium. Antonie Van Leeuwenhoek. 2009;96(2):115-39. Epub 20081001. doi:

10.1007/s10482-008-9284-5. PubMed PMID: 18830684.

22.   Bi X, Liu LF. RecA-independent and recA-dependent intramolecular plasmid

recombination: Differential homology requirement and distance effect. J Mol Biol.

1994;235(2):414-23. doi: 10.1006/jmbi.1994.1002. PubMed PMID: 8289271.

23.   Reams AB, Kofoid E, Kugelberg E, Roth JR. Multiple Pathways of Duplication

Formation with and Without Recombination (RecA) in *Salmonella enterica*. Genetics.

2012;192(2):397-415.   doi: 10.1534/genetics.112.142570.   PubMed   PMID:

WOS:000309547400006.

24.   Reams AB, Kofoid E, Savageau M, Roth JR. Duplication frequency in a population of

Salmonella enterica rapidly approaches steady state with or without recombination. Genetics.

2010;184(4):1077-94. Epub 20100118. doi: 10.1534/genetics.109.111963. PubMed PMID:

20083614; PubMed Central PMCID: PMCPMC2865909.

25.   Reams AB, Neidle EL. Genome plasticity in *Acinetobacter*: new degradative

capabilities acquired by the spontaneous amplification of large chromosomal segments.

Molecular Microbiology. 2003;47(5):1291-304. doi: DOI 10.1046/j.1365-2958.2003.03342.x.

PubMed PMID: WOS:000181056400009.

26.   Min YH, Kwon AR, Yoon JM, Yoon EJ, Shim MJ, Choi EC. Molecular analysis of

constitutive mutations in ermB and ermA selected in vitro from inducibly MLSB-resistant

enterococci. Arch Pharm Res. 2008;31(3):377-80. Epub 20080413. doi: 10.1007/s12272-001-

1167-8. PubMed PMID: 18409053.

27.   Sun S, Berg OG, Roth JR, Andersson DI. Contribution of gene amplification to

evolution of increased antibiotic resistance in *Salmonella typhimurium*. Genetics.

2009;182(4):1183-95. Epub 20090527. doi: 10.1534/genetics.109.103028. PubMed PMID:

19474201; PubMed Central PMCID: PMCPMC2728858.

28.   Vinella D, Cashel M, D'Ari R. Selected amplification of the cell division genes ftsQ-

ftsA-ftsZ   in   *Escherichia   coli*.   Genetics.   2000;156(4):1483-92.   doi:

10.1093/genetics/156.4.1483.   PubMed   PMID: 11102351;   PubMed   Central   PMCID:

PMCPMC1461353.

29.   Nilsson AI, Zorzet A, Kanth A, Dahlstrom S, Berg OG, Andersson DI. Reducing the

fitness cost of antibiotic resistance by amplification of initiator tRNA genes. Proc Natl Acad

Sci U S A. 2006;103(18):6976-81. Epub 20060424. doi: 10.1073/pnas.0602171103. PubMed

PMID: 16636273; PubMed Central PMCID: PMCPMC1459004.

30.   Andersson DI, Nicoloff H, Hjort K. Mechanisms and clinical relevance of bacterial
heteroresistance. Nature Reviews Microbiology. 2019;17(8):479-96. doi: 10.1038/s41579-
019-0218-1. PubMed PMID: WOS:000475479100007.

31.   Nicoloff H, Hjort K, Levin BR, Andersson DI. The high prevalence of antibiotic
heteroresistance in pathogenic bacteria is mainly caused by gene amplification. Nat
Microbiol.   2019;4(3):504-14.   doi: 10.1038/s41564-018-0342-0.   PubMed   PMID:
WOS:000459201400019.

32.   Sun S, Ke R, Hughes D, Nilsson M, Andersson DI. Genome-wide detection of
spontaneous chromosomal rearrangements in bacteria. Plos One. 2012;7(8):e42639. Epub
20120803. doi: 10.1371/journal.pone.0042639. PubMed PMID: 22880062; PubMed Central
PMCID: PMCPMC3411829.

33.   Tomanek I, Grah R, Lagator M, Andersson AMC, Bollback JP, Tkacik G, et al. Gene
amplification as a form of population-level gene expression regulation. Nat Ecol Evol.
2020;4(4):612-25.   doi: 10.1038/s41559-020-1132-7.   PubMed   PMID:
WOS:000519008300005.

34.   Ahmad SI, Kirk SH, Eisenstark A. Thymine metabolism and thymineless death in
prokaryotes   and   eukaryotes.   Annu   Rev   Microbiol.   1998;52:591-625.   doi:
10.1146/annurev.micro.52.1.591. PubMed PMID: 9891809.

35.   Gangjee A, Jain HD, Kurup S. Recent advances in classical and non-classical
Antifolates as antitumor and antiopportunistic infection agents: part I. Anti-Cancer Agent Me.
2007;7(5):524-42.   doi: Doi   10.2174/187152007781668724.   PubMed   PMID:
WOS:000249469700005.

36.   Gangjee A, Jain HD, Kurup S. Recent advances in classical and non-classical
antifolates as antitumor and antiopportunistic infection agents: Part II. Anti-Cancer Agent
Me.   2008;8(2):205-31.   doi: Doi   10.2174/187152008783497064.   PubMed   PMID:
WOS:000255810200007.

37.   Huennekens FM. The methotrexate story: a paradigm for development of cancer
chemotherapeutic agents. Adv Enzyme Regul. 1994;34:397-419. doi: 10.1016/0065-
2571(94)90025-6. PubMed PMID: 7942284.

38.   Plowe CV. Malaria chemoprevention and drug resistance: a review of the literature
and policy implications. Malar J. 2022;21(1):104. Epub 20220324. doi: 10.1186/s12936-022-
04115-8. PubMed PMID: 35331231; PubMed Central PMCID: PMCPMC8943514.

39.   Gleckman R, Blagg N, Joubert DW. Trimethoprim: mechanisms of action,
antimicrobial   activity,   bacterial   resistance,   pharmacokinetics,   adverse   reactions,   and

therapeutic indications. Pharmacotherapy. 1981;1(1):14-20. doi: 10.1002/j.1875-
9114.1981.tb03548.x. PubMed PMID: 6985448.

40.   Brochet M, Couve E, Zouine M, Poyart C, Glaser P. A naturally occurring gene
amplification leading to sulfonamide and trimethoprim resistance in Streptococcus agalactiae.
J Bacteriol. 2008;190(2):672-80. Epub 20071116. doi: 10.1128/JB.01357-07. PubMed PMID:
18024520; PubMed Central PMCID: PMCPMC2223700.

41.   Goker E, Waltham M, Kheradpour A, Trippett T, Mazumdar M, Elisseyeff Y, et al.
Amplification of the Dihydrofolate-Reductase Gene Is a Mechanism of Acquired-Resistance
to Methotrexate in Patients with Acute Lymphoblastic-Leukemia and Is Correlated with P53
Gene-Mutations.   Blood.   1995;86(2):677-84.   doi: DOI
10.1182/blood.V86.2.677.bloodjournal862677. PubMed PMID: WOS:A1995RH80100030.

42.   Inselburg J, Bzik DJ, Horii T. Pyrimethamine resistant Plasmodium falciparum:
overproduction of dihydrofolate reductase by a gene duplication. Mol Biochem Parasitol.
1987;26(1-2):121-34. doi: 10.1016/0166-6851(87)90136-8. PubMed PMID: 3323903.

43.   Morales C, Garcia MJ, Ribas M, Miro R, Munoz M, Caldas C, et al. Dihydrofolate
reductase amplification and sensitization to methotrexate of methotrexate-resistant colon
cancer cells. Mol Cancer Ther. 2009;8(2):424-32. Epub 20090203. doi: 10.1158/1535-
7163.MCT-08-0759. PubMed PMID: 19190117.

44.   Baym M, Lieberman TD, Kelsic ED, Chait R, Gross R, Yelin I, et al. Spatiotemporal
microbial evolution on antibiotic landscapes. Science. 2016;353(6304):1147-51. doi:
10.1126/science.aag0822. PubMed PMID: 27609891; PubMed Central PMCID:
PMCPMC5534434.

45.   Palmer AC, Toprak E, Baym M, Kim S, Veres A, Bershtein S, et al. Delayed
commitment to evolutionary fate in antibiotic resistance fitness landscapes. Nat Commun.
2015;6:7385. Epub 20150610. doi: 10.1038/ncomms8385. PubMed PMID: 26060115;
PubMed Central PMCID: PMCPMC4548896.

46.   Patel V, Matange N. Adaptation and compensation in a bacterial gene regulatory
network evolving under antibiotic selection. Elife. 2021;10. Epub 20210930. doi:
10.7554/eLife.70931. PubMed PMID: 34591012; PubMed Central PMCID:
PMCPMC8483737.

47.   Watson M, Liu JW, Ollis D. Directed evolution of trimethoprim resistance in
Escherichia coli. Febs J. 2007;274(10):2661-71. Epub 20070419. doi: 10.1111/j.1742-
4658.2007.05801.x. PubMed PMID: 17451440.

48.   Toprak E, Veres A, Michel JB, Chait R, Hartl DL, Kishony R. Evolutionary paths to antibiotic resistance under dynamically sustained drug selection. Nat Genet. 2011;44(1):101-5. Epub 20111218. doi: 10.1038/ng.1034. PubMed PMID: 22179135; PubMed Central PMCID: PMCPMC3534735.

49.   Matange N. Highly Contingent Phenotypes of Lon Protease Deficiency in Escherichia coli upon Antibiotic Challenge. J Bacteriol. 2020;202(3). Epub 20200115. doi: 10.1128/JB.00561-19. PubMed PMID: 31740490; PubMed Central PMCID: PMCPMC6964741.

50.   Rodrigues JV, Bershtein S, Li A, Lozovsky ER, Hartl DL, Shakhnovich EI. Biophysical principles predict fitness landscapes of drug resistance. Proc Natl Acad Sci U S A. 2016;113(11):E1470-8. Epub 20160229. doi: 10.1073/pnas.1601441113. PubMed PMID: 26929328; PubMed Central PMCID: PMCPMC4801265.

51.   Vinchhi R, Yelpure C, Balachandran M, Matange N. Pervasive gene deregulation underlies adaptation and maladaptation in trimethoprim-resistant *E. coli*. mBio. 2023:e0211923. Epub 20231130. doi: 10.1128/mbio.02119-23. PubMed PMID: 38032208.

52.   Ranquet C, Gottesman S. Translational regulation of the Escherichia coli stress factor RpoS: a role for SsrA and Lon. J Bacteriol. 2007;189(13):4872-9. Epub 20070420. doi: 10.1128/JB.01838-06. PubMed PMID: 17449615; PubMed Central PMCID: PMCPMC1913435.

53.   Xu J, Li T, Gao Y, Deng J, Gu J. MgrB affects the acid stress response of Escherichia coli by modulating the expression of iraM. FEMS Microbiol Lett. 2019;366(11). doi: 10.1093/femsle/fnz123. PubMed PMID: 31158277.

54.   Matange N, Bodkhe S, Patel M, Shah P. Trade-offs with stability modulate innate and mutationally acquired drug resistance in bacterial dihydrofolate reductase enzymes. Biochem J. 2018;475(12):2107-25. Epub 20180629. doi: 10.1042/BCJ20180249. PubMed PMID: 29871875.

55.   Ohno S. Evolution by gene duplication. London, New York: Allen & Unwin; Springer-Verlag; 1970. xv, 160 p. p.

56.   Bergthorsson U, Andersson DI, Roth JR. Ohno's dilemma: Evolution of new genes under continuous selection. P Natl Acad Sci USA. 2007;104(43):17004-9. doi: 10.1073/pnas.0707158104. PubMed PMID: WOS:000250487600051.

57.   Long MY, VanKuren NW, Chen SD, Vibranovski MD. New Gene Evolution: Little Did We Know. Annual Review of Genetics, Vol 47. 2013;47:307-33. doi: 10.1146/annurev-genet-111212-133301. PubMed PMID: WOS:000329816000014.

58.   Holloway AK, Palzkill T, Bull JJ. Experimental evolution of gene duplicates in a bacterial plasmid model. J Mol Evol. 2007;64(2):215-22. Epub 20070108. doi: 10.1007/s00239-006-0087-x. PubMed PMID: 17211548.

59.   Kugelberg E, Kofoid E, Reams AB, Andersson DI, Roth JR. Multiple pathways of selected gene amplification during adaptive mutation. P Natl Acad Sci USA. 2006;103(46):17319-24.   doi: 10.1073/pnas.0608309103.   PubMed   PMID: WOS:000242249400047.

60.   Naseeb S, Ames RM, Delneri D, Lovell SC. Rapid functional and evolutionary changes follow gene duplication in yeast. Proc Biol Sci. 2017;284(1861). doi: 10.1098/rspb.2017.1393.   PubMed   PMID: 28835561;   PubMed   Central   PMCID: PMCPMC5577496.

61.   Tomanek I, Guet CC. Adaptation dynamics between copy-number and point mutations. Elife. 2022;11. doi: ARTN e82240

10.7554/eLife.82240. PubMed PMID: WOS:000912674700001.

62.   Nicoloff H, Perreten V, Levy SB. Increased genome instability in *Escherichia coli* lon mutants: relation to emergence of multiple-antibiotic-resistant (Mar) mutants caused by insertion sequence elements and large tandem genomic amplifications. Antimicrob Agents Chemother. 2007;51(4):1293-303. Epub 20070112. doi: 10.1128/AAC.01128-06. PubMed PMID: 17220404; PubMed Central PMCID: PMCPMC1855481.

63.   Nicoloff H, Perreten V, McMurry LM, Levy SB. Role for tandem duplication and lon protease in AcrAB-TolC-dependent multiple antibiotic resistance (Mar) in an *Escherichia coli* mutant   without   mutations   in   marRAB   or   acrRAB.   Journal   of   Bacteriology. 2006;188(12):4413-23. doi: 10.1128/Jb.01502-05. PubMed PMID: WOS:000238173900027.

64.   Derbyshire KM, Kramer M, Grindley ND. Role of instability in the cis action of the insertion sequence IS903 transposase. Proc Natl Acad Sci U S A. 1990;87(11):4048-52. doi: 10.1073/pnas.87.11.4048.   PubMed   PMID: 2161528;   PubMed   Central   PMCID: PMCPMC54044.

65.   Rouquette C, Serre MC, Lane D. Protective role for H-NS protein in IS1 transposition. J Bacteriol. 2004;186(7):2091-8. doi: 10.1128/JB.186.7.2091-2098.2004. PubMed PMID: 15028694; PubMed Central PMCID: PMCPMC374409.

66.   Hua CF, Wang TT, Shao XL, Xie YP, Huang H, Liu JG, et al. *Pseudomonas psyringae* dual-function protein Lon switches between virulence and metabolism by acting as both DNA-binding transcriptional regulator and protease in different environments. Environ

Microbiol. 2020;22(7):2968-88. doi: 10.1111/1462-2920.15067. PubMed PMID:
WOS:000537012700001.

67. Karlowicz A, Wegrzyn K, Gross M, Kaczynska D, Ropelewska M, Siemiatkowska M,
et al. Defining the crucial domain and amino acid residues in bacterial Lon protease for DNA
binding and processing of DNA-interacting substrates. J Biol Chem. 2017;292(18):7507-18.
doi: 10.1074/jbc.M116.766709. PubMed PMID: WOS:000400761300019.

68. Liu T, Lu B, Lee I, Ondrovicová G, Kutejová E, Suzuki CK. DNA and RNA binding
by the mitochondrial Lon protease is regulated by nucleotide and protein substrate. J Biol
Chem. 2004;279(14):13902-10. doi: 10.1074/jbc.M309642200. PubMed PMID:
WOS:000220478500077.

69. Lu B, Yadav S, Shah PG, Liu T, Tian B, Pukszta S, et al. Roles for the human ATP-
dependent Lon protease in mitochondrial DNA maintenance. J Biol Chem.
2007;282(24):17363-74. Epub 20070409. doi: 10.1074/jbc.M611540200. PubMed PMID:
17420247.

70. Sonezaki S, Okita K, Oba T, Ishii Y, Kondo A, Kato Y. Protein substrates and heat
shock reduce the DNA binding ability of *Escherichia coli* Lon protease. Appl Microbiol Biot.
1995;44(3-4):484-8. PubMed PMID: WOS:A1995TM49800033.

71. Breidenstein EB, Bains M, Hancock RE. Involvement of the lon protease in the SOS
response triggered by ciprofloxacin in Pseudomonas aeruginosa PAO1. Antimicrob Agents
Chemother. 2012;56(6):2879-87. Epub 20120326. doi: 10.1128/AAC.06014-11. PubMed
PMID: 22450976; PubMed Central PMCID: PMCPMC3370746.

72. Matange N, Hegde S, Bodkhe S. Adaptation Through Lifestyle Switching Sculpts the
Fitness Landscape of Evolving Populations: Implications for the Selection of Drug-Resistant
Bacteria at Low Drug Pressures. Genetics. 2019;211(3):1029-44. Epub 20190122. doi:
10.1534/genetics.119.301834. PubMed PMID: 30670539; PubMed Central PMCID:
PMCPMC6404241.

73. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol.
2014;1151:165-88. doi: 10.1007/978-1-4939-0554-6_12. PubMed PMID: 24838886; PubMed
Central PMCID: PMCPMC4239701.

1218
1219

1220

1221
1222