**Variation in a microbial mutualist has transcriptional and phenotypic consequences for host-parasite interactions**

Addison D. Buxton-Martin[1*], Emile Gluck-Thaler[2,3], Eunnuri Yi[1], John R. Stinchcombe[4], Corlett W. Wood[1*]

    (1) Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

    (2) Department of Plant Pathology, University of Wisconsin-Madison, Madison, WI, USA

    (3) Wisconsin Institute for Discovery, Madison, WI, USA

    (4) Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

   *Corresponding authors:

        ADBM: admartin@sas.upenn.edu

        CWW: corlett@sas.upenn.edu

**Abstract**

Strains of microbial symbionts often vary in their effect on their host. However, little is known about how the genetic variation in microbial symbiont populations impacts host interactions with other co-colonizing microbes. Here we investigated how different strains of nitrogen-fixing rhizobial bacteria affect their host plant's response to parasitic root-knot nematodes in the legume *Medicago truncatula*. Using dual RNAseq of the root organs harboring rhizobia or nematodes, we identified genes from plant host, rhizobia, and nematode whose expression differed between parasite-infected and -uninfected plants, and between plants inoculated with different rhizobial strains. At the site of host-parasite interactions (in nematode galls), hundreds of host genes and few nematode genes differed in expression between host plants inoculated with different rhizobia strains. At the site of host-mutualist interactions (in rhizobia nodules), hundreds of host genes and few rhizobial genes responded to parasite infection. The vast majority of parasite-induced changes in host gene expression depended on the resident rhizobia strain. We additionally observed phenotypic differences in the gall and nodule counts on host root systems and in some root architecture traits, showing that genetic variation in a mutualist symbiont impacted parasite colonization rates. The transcriptomic and phenotypic differences we observed suggest that microbial indirect genetic effects play an underappreciated role in structuring their host's interactions with other co-colonizing microorganisms, which may be mediated by changes in host resource budgets, defense response, or developmental trajectories.

**Keywords:** dual RNAseq, host-microbe interactions, indirect genetic effects, intraspecific genetic variation, mutualism, parasitism

**Introduction**

An individual plant hosts thousands of microorganisms, ranging from bacteria and viruses to eukaryotes like fungi and protozoa. These microbial species are composed of strains that vary – often dramatically – in their effects on their host (Heath, 2010; Heath and Stinchcombe, 2014; Lange et al., 2023; Lively and Dybdahl, 2000; Morand et al., 1997; Osborne

et al., 2009; Parker et al., 2017). Here we investigate how strain-level genetic variation in microbial symbiont populations cascades to affect a plant's interactions with other microbes.

Co-colonizing symbionts – parasites, commensals, or mutualists – often compete with or facilitate one another (Coyte et al., 2015; Stein et al., 2013) directly or indirectly via their effects on their shared host (Foster et al., 2017; Pedersen and Fenton, 2007). This framework is implicitly or explicitly employed by researchers studying interactions between hosts and microbial communities in a variety of disciplines: in plant pathology, as the priming of host defenses (Ahmad et al., 2010; Benjamin et al., 2022; Guo and Ge, 2017; Zamioudis and Pieterse, 2012); in parasitology within the context of co-infections or mixed infections (Alizon et al., 2013; Devi et al., 2021; Venter et al., 2022); and in community and microbial ecology, as priority effects, indirect effects, or higher-order interactions (Debray et al., 2022; Miller and Travis, 1996; Strauss, 1991). Across these disparate subfields, consensus has emerged that interactions between co-colonizing microbes of different species have far-reaching ramifications for the assembly and function of host-associated communities (Alizon et al., 2013; Banerjee et al., 2018; Dastogeer et al., 2020; Shetty et al., 2017; terHorst et al., 2018). Co-colonizing microbes of different species can influence the relationship their co-colonizers have with their joint host at all stages of their respective lifecycles and in different ways: from impacting each other's colonization success (Telfer et al., 2010), transmission dynamics (Bousema et al., 2008; Pollitt et al., 2015; Schneider et al., 2018), or the benefits they confer and costs they inflict on their shared host (Barrett et al., 2015; Larimer et al., 2014; Magnoli and Bever, 2023; Weldon et al., 2020).

However, within symbiont species there is extensive genetic and functional variation. For example, pathogen genotypes differ dramatically in virulence and transmissibility (Lively and Dybdahl, 2000; Morand et al., 1997), strains of bacterial mutualists vary in the costs and benefits they bring to their hosts (Heath, 2010; Heath and Stinchcombe, 2014; Heath and Tiffin, 2007; Calvert *et al.*, 2023), and genotypes of defensive mutualists differ in their protective effect (Lange et al., 2023; Osborne et al., 2009; Parker et al., 2017). These examples share a unifying feature: the genes in one species (the microbe) impact trait expression in another (the host). Such "interspecific indirect genetic effects" (De Lisle et al., 2022; Shuster et al., 2006) are thought to be pervasive in traits mediating symbiotic interactions (Fronk and Sachs, 2022) and alter the tempo and mode of trait evolution, because the genetic basis of trait expression and evolution in

one species (the host) resides in another species (the microbe) (De Lisle et al., 2022; O'Brien et al., 2021b; Queller, 2014; Shuster et al., 2006).

The extensive variation of symbiosis traits within microbial species raises a crucial question: do different microbial strains of the same species vary in their effect on the host's interaction with other microbes? Put another way: can genetic variation for traits mediating species interactions reside in species not directly involved in the interaction itself? The answer to this question has broad implications for the assembly, function, and evolution of host-associated communities. If strain-level (i.e., intraspecific) variation in a microbial symbiont influences its host's interaction with other symbionts, then strain-level changes in microbial symbiont populations such as changes in allele frequencies or strain abundance could have cascading impacts on the structure or function of the entire host-associated community. The few studies that have addressed this possibility have found that microbial genotypes do vary in their effect on the broader host-associated community, especially in defensive mutualisms (Lange et al., 2023; Osborne et al., 2009; Parker et al., 2017). However, when two microbes inhabit the same niche within the host (e.g., the animal gut) or colonize systemically, disentangling the host's responses to two co-colonizing symbionts and the indirect effects between symbionts can be challenging.

Plant galls provide a unique means of circumventing these challenges and investigating how genetic variation in one microbial symbiont species impacts its host's interactions with other co-colonizing symbionts. Galls are specialized structures that are made of plant tissue but are induced and colonized by another organism such as an insect or microbe. Many plant parasites, pathogens, and mutualists form galls (Harris and Pitzschke, 2020). Common examples include legume root nodules induced by nitrogen-fixing rhizobia, goldenrod galls induced by the goldenrod gall fly, and root knot galls induced by root-knot nematodes (Harris and Pitzschke, 2020). These galls are the locus of interaction between the gall inducers and their host plant (Fronk and Sachs, 2022). Because each gall is specific to a single pairwise species interaction, galls provide a unique opportunity to sample the symbiotic interface between the host and single symbiont even in hosts interacting with many other symbionts. The abundance and discrete nature of these structures is also helpful in quantifying and phenotyping aspects of host-symbiont interactions such as parasite or mutualist colonization rates or host susceptibility.

Transcriptomic methods when applied to plant galls are particularly useful in elucidating the otherwise difficult to observe interaction involving hosts and microbes, as they provide

genome-wide proxies of the underlying biology mediating the interaction between host and symbiont. In particular, dual RNAseq, where RNA from two symbiotic organisms are sequenced in the same sample, has become a useful tool to investigate in vivo interactions between a host and symbiont (Westermann et al., 2017; Westermann and Vogel, 2021; Wolf et al., 2018). Thus comparing the gene expression profiles of galls can be used to examine changes in the interactions between host plants and their microbial gall-forming symbionts under different conditions. Whatismore, interactions between plants and symbiotic microbes may drive systemic reactions outside of just the organ that mediates the host-symbiont interaction (Pieterse et al., 2014; Mauch-Mani et al., 2017) and plant-microbe interactions can often be regulated by signals produced in parts of the host plant not directly involved in the interaction (Li *et al.*, 2022). Comparing gene expression profiles of galls to that of other organs can thus be used to assess the systemic or local nature of an interaction between host and symbiont (Oates *et al.,* 2021).

We leveraged these unique features of root-knot nematode galls and rhizobia nodules  as well as dual RNAseq to investigate how strain-level variation in a microbial mutualist influences its host plant's transcriptomic response to parasite infection. We inoculated a single genotype of the model legume *Medicago truncatula* with one of two strains of mutualistic nitrogen-fixing rhizobia bacteria *Ensifer meliloti* and infected half of these plants with the parasitic root-knot nematode *Meloidogyne hapla*. We then profiled plant, mutualist, and parasite gene expression in rhizobia nodules and nematode galls (Figure **1**) and quantified host fitness, root architecture, and parasite and mutualist colonization rates. We asked: (1) how do mutualistic rhizobia strains influence host traits, parasite colonization rates, and gene expression in nematode galls?; (2) how do parasitic nematodes affect host traits, mutualist colonization rates, and gene expression in rhizobial nodules?; and (3) is the effect of parasite infection on host traits, mutualist colonization rates, and nodule gene expression influenced by the resident rhizobia strain?

On a phenotypic level, we found that nematode infection status impacted root architecture and rhizobia colonization rates, while rhizobia strain impacted parasite colonization rate. Neither nematode infection status, nematode infection severity, or rhizobia strain impacted a host fitness proxy, whereas rhizobia colonization rates increased host fitness. On a transcriptomic level, we found that strain-level variation in the rhizobial mutualist significantly alters the host plant's transcriptomic response to parasite infection. Plants inoculated with different rhizobial strains had differential gene expression in nematode galls, the site of parasite infection.

Furthermore, parasite infection altered the expression of hundreds of genes in rhizobia nodules, and this effect depended strongly on the resident rhizobia strain. Our results show that intraspecific genetic variation in a keystone microbial mutualist has far reaching consequences on host-parasite interactions with mutualist strain having transcriptional consequences in mature host-parasite structures and consequences for parasite colonization rates. Whatismore, parasite infection carried consequences for host-mutualist interactions with transcriptional consequences in mature host-mutualist structures and altered mutualist colonization rates.

**Materials and Methods**

| Scale of inference | The scale at which the factor of interest is applied | Number of replicates at the appropriate scale |
|---|---|---|
| Population of plants (transcriptomic response) | Manipulated parasitic nematode infection and rhizobia strain inoculation on individual plants; pooled organs (gall, nodule, or root) from 6 individual plants into a single biological replicate | 3 replicates of each organ per treatment (nematode infection and rhizobia strain combination) |
| Population of plants (phenotypic response) | Manipulated parasitic nematode infection and rhizobia strain inoculation on individual plants | 19 plants per combination of nematode treatment and rhizobia strain |

*Study system*

Medicago truncatula Gaertn. is an annual legume native to the Mediterranean (Cook, 1999). Upon the transfer of compatible chemical signals, *M. truncatula* plants form associations with nitrogen-fixing rhizobial bacteria in the soil, such as *Ensifer meliloti*. Rhizobia are housed in nodules that develop from the roots of their host plant. Nodules provide the requisite low-oxygen environment and resources for the rhizobia to differentiate into bacteroids within intracellular compartments called symbiosomes. Within these symbiosomes, rhizobia fix atmospheric nitrogen into biologically available forms of nitrogen and provide some to their host plant and in exchange hosts provide rhizobia with carbon-containing compounds derived from photosynthate. The *Medicago-Ensifer* system has become a well characterized model of resource-exchange mutualisms (Long, 1989).

The pathogenic root-knot nematode *Meloidogyne hapla* also associates with *M. truncatula* roots. Female *M. hapla* burrow through the intercellular spaces of host roots in search of a suitable infection site. There *M. hapla* uses its stylet mouthpiece to inject effector proteins into parenchymal host cells and root tissues that suppress immune responses and induce cell differentiation into giant cells from which the pathogenic nematode siphons nutrients. Nematode galls form around this site of infection (Jones and Goto, 2011).

We chose the in-bred line of *M. truncatula* accession HM145 from the French National Institute for Agriculture Research (INRAE) germplasm library for its high nodulation rate and

nematode susceptibility (Wood et al., 2018). We obtained inoculum containing the northern root-knot nematode *M. hapla* line LM originally collected in France and supplied by the Department of Entomology and Plant Pathology at North Carolina State University. We used *E. meliloti* strains USDA1021 and WSM1022 from lab stocks (JRS). WSM1022 is known as a high-efficiency nitrogen fixer in contrast to other strains (Terpolilli *et al*., 2013).

*Sample Preparation*

We grew *M. truncatula* HM145 in Cone-tainers and maintained them in growth chambers at the University of Pennsylvania following the instructions of Per Garcia *et al*. (2006). Plants were arranged in a randomized complete block design of rhizobia strain (USDA1021 or WSM1022) and nematode status (infected or uninfected) and plants were inoculated with the appropriate nematode and rhizobia treatment on day 14 after planting. Each treatment had three replicates, which were each composed of six individual plants. See Methods S1 and Table S3 for greater detail regarding sample preparation.

*Organ sample harvesting*

Organ samples were collected from all plants 52-53 days after planting. All six plants that constituted a replicate were harvested on the same day. Nodule samples were collected from all plants and root and nematode gall samples were collected from plants inoculated with both rhizobia and nematodes (Figure **1**). To harvest, the root system was washed in water, then organs were manually excised using a pair of sterilized scissors and immediately flash frozen in liquid nitrogen. Samples were taken from the top ⅓ of the root system so as to collect nematode galls and nodules at a consistent level of maturity. All samples were stored at -80 °C until RNA extraction which was performed within 30 days of harvesting.

Figure 1: Conceptual diagram of the experimental design, treatments, and samples used in this experiment. A) The host plant, *Medicago truncatula*, with rhizobia nodules and nematode galls in the host plant root system as magnified insets. B) Each of the four treatments of symbionts consisted of three biological replicates with six plants constituting each biological replicate, totaling 18 plants per treatment. C) Rhizobia nodules, roots, and nematode galls were harvested from hosts as indicated. Organs from the six plants constituting one biological replicate were pooled. D) The recoverable RNA reads from each organ.

*Dual RNA sequencing of nodules and nematode galls and RNA sequencing of roots*

To isolate species interactions in co-colonized hosts, we extracted RNA from nodules and nematode galls for Dual RNAseq. To assess the localized or systemic nature of host changes in gene expression, we also extracted RNA from samples of roots from co-colonized hosts. We used a Trizol and QIAGEN RNeasy Kit based RNA extract and a Nanodrop ND-1000 (Thermo Fisher Scientific) to confirm RNA quality. RNA extracts were shipped to Genewiz Azenta Life Sciences (South Plainfield, NJ) for sequencing. We had rRNA depletion performed during library preparation so as to preserve non-polyadenylated prokaryotic reads, enabling quantification of gene expression for both eukaryotes and prokaryotes. We distributed RNA

samples from the same treatment across different lanes to preclude sequencing lanes as a batch effect. See Methods S2 for details regarding RNA extraction and sequencing methods. Raw read data is available through the Sequence Read Archive under BioProjectID PRJNA1090526.

*Dual RNA-seq data analysis and gene expression quantification*

After sequencing, we used the default settings of Trimmomatic (Bolger et al., 2014) to remove sequencing adapters from reads and discard poor quality reads. We used Sortmerna (Kopylova et al., 2012) and the rRNA sequences of all three species in the study system in the SILVA database (Glöckner et al., 2017; Pruesse et al., 2007; Quast et al., 2013; Yilmaz et al., 2014, p. 2) to remove rRNA reads that were not removed by rRNA depletion. Concatenated genomes were created by concatenating the reference genomes (Methods S3) for the one or two organisms present in each sample. The remaining non-rRNA and high quality reads were aligned with default settings to the appropriate concatenated genomes for each sample (Espindula *et al.*, 2020) using HISAT2 (Kim et al., 2019). Reads with alignment quality scores <30 and whose paired read aligned to a separate chromosome or scaffold were discarded. The remaining alignments were used with the union method from HTSeq-count (Putri et al., 2022) to quantify gene expression of all genomic features not annotated as repeat regions, rRNA, or tRNA in concatenated reference genome annotations.

We found our gene expression quantification data to be suitable for downstream analyses (Chung et al., 2021; Love et al., 2014): host gene expression quantification data segregated on a principal component analysis by sample type (Figure S**1**) and the percentage of genes with a coefficient of variation within at least one treatment that was greater than across all samples never exceeded 20% (Table S**5**), indicating biological variation was greater than technical variation for most genes. See Methods S3 for greater details regarding reference genomes and software.

*Differential expression analyses*

We considered genes with an aligned read count of >1 RPKM within its genome as expressed (Hebenstreit et al., 2011) and applied this categorization across all samples. To identify differentially expressed genes (DEGs), we used DESeq2 (Love et al., 2014) with a likelihood ratio test and Benjamini–Hochberg multiple test correction on all genes expressed in a given

organ. We used several DESeq2 models to test for differential expression across different sample types. See Methods S4 for details about the models used with DESeq2.

*Gene ontology overrepresentation analyses*

We used Gene Ontology overrepresentation analyses (GOORA) of gene lists generated from our differential expression analyses to formulate hypotheses about the biological processes that may be different across our treatments. For each *M. truncatula* organ in our analysis, we created a custom database of the genes that were expressed in that organ and their corresponding Gene Ontology (GO) annotations from the *M. truncatula* genome annotation. We used the default settings (p value 0.05, q value 0.2, Benjamini–Hochberg correction) of the enrichGO function from the clusterProfile package in R (Wu et al., 2021) against the aforementioned databases to identify over represented GO terms in the gene lists generated by our differential expression analyses.

*Phenotypic effects of rhizobia strains on host fitness and nematode colonization*

We recapitulated the experiment as described above with an additional 76 plants (19 per treatment group). Upon harvesting, we recorded and removed a number of nodules from a subset of 7-8 plants from each treatment for a separate experiment. We separated root and shoot systems and dried the shoot systems for >48 hours before weighing them. Root systems were frozen for later processing. We scanned root systems at 1200 dpi following York (2023). After scanning all roots, root systems from plants infected with nematodes were stained with red dye following the procedure outlined by Thies *et al.* (2002) to make nematode galls more apparent and then scanned them. We manually inspected dyed and undyed images and manually counted nodules and nematode galls. RootPainter (Smith *et al.*, 2022), a deep learning tool designed to use convolutional neural networks to segment root images, was used to segment the undyed scanned root images and RhizoVision Explorer (Seethepalli and York, 2020; Seethepalli *et al.*, 2021) was then used to quantify root architecture from segmented images. Specifically, we examined above ground biomass as a proxy for host fitness (Younginger *et al.*, 2017), root biomass, total number of branch points and root tips, network area, root system perimeter and average, maximum, and median root diameter as indications of host nutrient acquisition strategy (van der Bom *et al.*, 2020).

We used linear mixed-effects models and analysis of variance in R to examine relationships between traits and rhizobia strain and nematode effects. We fit our models to the data using the lme4 (Bates *et al.,* 2015) and used the Satterthwaite's degrees of freedom method executed by the lmerTest package (Kuznetsova *et al*., 2017) to test for significance. We used the DHARMα package (Hartig, 2022) to check all of our models for outliers, normality, and homoscedasticity after fitting them to our data.

## Results

*Mutualist strain impacts the expression of host genes related to defense, development, and pathogenesis at the site of host-parasite interactions.*

Plant host gene expression in nematode galls was different in hosts inoculated with different rhizobia strains. Out of the 51,653 examined genes in the *M. truncatula* genome, 23,069 genes (44.661%) were expressed. Of these genes, 285 genes (1.24%) were differentially expressed (DE) across hosts with different rhizobia strains. Of these DEGs, the majority (208 genes; 73.0%) had higher expression in hosts inoculated with WSM1022 than hosts inoculated with USDA1021 (Figure **2a**,**c**).

Figure 2: Rhizobia strain influences gene expression at the site of host-parasite interactions with largely similar responses in gene expression occurring in nematode galls and nodules and in nematode galls and roots. Log fold change (LFC) and standard error values were calculated

using DESeq2. LFC values are the logarithm base 2 of the ratio of normalized mean expression in hosts inoculated with USDA1021 to that in hosts inoculated with WSM1022. Genes with significant log fold changes determined by likelihood ratio tests performed by DESeq2 are indicated with color and also with standard error bars in D and E. a) 261 host genes were differentially expressed (DE) in nematode galls across hosts with different rhizobia strains. b) Six nematode genes were DE across hosts with different rhizobia strains. c) Similar numbers of genes were DE in nematode galls and nodules across hosts with different rhizobia strains, but the magnitude of differential expression was higher in nodules. Differences in magnitude were tested using the optimal pooled t-test (Guo and Yuan, 2017) ('n.s.' indicates no significance, * indicates p-value < 0.05). d) Rhizobial differences produced largely similar responses in host gene expression in nodules and nematode galls. 712 genes showed a similar response to rhizobia strain in nodules and nematode galls (main effect) while 136 showed different responses in the two organs (interaction effect). e) Rhizobial differences produced largely similar responses in host gene expression in roots and nematode galls. 191 genes showed a similar response to rhizobia strain in roots and nematode galls (main effect) while 31 showed different responses in the two organs (interaction effect).

GOORA showed that annotations associated with diverse biological processes are overrepresented in all 285 host DEGs that responded to rhizobial strain differences in nematode galls (Dataset S6). Notably, genes annotated as related to carbohydrate metabolism were overrepresented. Functions related to the plant hormone abscisic acid, which coordinates plant responses to biotic and abiotic stress (Cao et al., 2011), were also overrepresented. Functions related to metabolism of cell wall components such as xylan, hemicellulose, arabinan, and pectin, and cell wall organization and biogenesis were also overrepresented. Changes in plant cell walls are both a symptom of and defense response against root knot nematode infection (Caillaud et al., 2008; Przybylska & Obrępalska-Stęplowska, 2020). Phosphoprotein regulation was also overrepresented in the host gene expression changes (Table S**2a**; see Dataset S6 for full details of GOORA results).

GOORA produced similar results when performed with only the DEGs in nematode galls that were upregulated in hosts inoculated with USDA1021. In contrast, within the DEGs in nematode galls that were upregulated in hosts inoculated with WSM1022, galacturan

1,4-alpha-galacturonidase activity (GO:0047911) and carbohydrate metabolic process (GO:0005975) were the only overrepresented annotations (Dataset S6). The activity of galacturan 1,4-alpha-galacturonidase degrades pectin in plant cell walls (Hasegawa and Nagel, 1968). Galacturan 1,4-alpha-galacturonidase enzymes are also secreted by root knot nematodes into cells at the infection site and pectin de-esterification and degradation is critical for successful root knot nematode infection (Jagdale et al., 2021; Wieczorek et al., 2014). Pectin metabolism and homeostasis affects many developmental pathways in plants (Guo et al., 2022; Wolf et al., 2009). Together these ORA results suggest the possibility that host carbohydrate metabolism, developmental processes, defense responses, or host pathogenesis are potentially altered in nematode galls when hosts are colonized by these two different rhizobia strains.

*Mutualist strain has little impact on global parasite gene expression but does alter a putative parasitism gene*

Nematode gene expression showed little response to differences in the rhizobia strains resident in host nodules. Out of the 14,419 genes in the *M. hapla* genome, 12,880 genes (89.326%) were expressed under the aforementioned expression criteria. Of these genes, only five (0.04%) were DEGs (Figure **2b**, Table S**1a**).

Many parasitic nematode effector proteins localize to the nucleus where they modulate host gene expression, so we investigated the predicted protein products of the five differentially expressed genes for secretory signal sequences using SignalP 6.0 (Teufel et al., 2022) and for nuclear localization signals using NLStradamus (Nguyen Ba et al., 2009). Four of the five DEGs (MhA1_Contig1073.frz3.gene2, MhA1_Contig1386.frz3.gene15, MhA1_Contig253.frz3.gene16, and MhA1_Contig953.frz3.gene1) were predicted to code for secreted proteins by Signal 6.0. One of these four genes (MhA1_Contig1073.frz3.gene2) is also predicted to contain a nuclear localization signal sequence by NLStradamus. The protein product of the predicted cylicin-2 gene (MhA1_Contig253.frz3.gene16) also shows significant sequence similarity (coverage of 95% and percent identity 81.13%) to Minc00331 in the closely related *Meloidogyne incognita* which is presumed to be a parasitism gene as it is upregulated during parasitic life stages (Nguyen et al., 2018).

*Mutualist strain variation elicits similar responses in nodules and nematode galls*

Out of the 26,875 *M. truncatula* genes expressed in all three organ types of nematode-infected hosts (52.030% of the genome), 9 genes showed a similar response to rhizobia strain across all organ types (Table S1c) and 705 genes (2.62% of expressed genes) showed a rhizobia strain by organ interaction effect. Notably, one gene that responded to rhizobia strain across all organ types (MtrunA17_Chr8g0364961/MtTPS9) and showed higher expression in hosts colonized by USDA1021 has been shown to be upregulated in response to abiotic stress and may mediate jasmonic acid response (Song *et al*., 2021). This indicates that the response to different rhizobia strains varies across organs and that very few host genes are responding similarly in all organs to intraspecific genetic variation in rhizobia.

Few genes are responding to differences in mutualist genotypes in similar ways across roots, nodules, and nematode galls, but similar responses can be observed across pairs of organs. Out of the 25,732 *M. truncatula* genes expressed in both symbiotic structures – nodules and nematode galls – 877 genes showed a similar response and 77 genes showed a different response to mutualist strain (Figure **2d**), indicating that the majority of genes that respond to different mutualist strains in nematode galls and nodules (81%) are responding in similar ways. Despite the lack of systemic response to different rhizobia strains across all organs in this study, symbiotic organs are responding similarly to differences in rhizobial strain. GOORA of the genes responding similarly in nematode galls and nodules suggests that processes associated with carbohydrate metabolism, cell walls, signal transduction, phosphorylation, and rhythmic/circadian processes may be responding in similar ways in the two symbiotic organs (Dataset S6). When changes in nematode galls were compared to changes in roots, we found that out of the 25,086 *M. truncatula* genes expressed in nematode galls and roots, 231 genes showed a similar response and 46 genes showed a different response to rhizobial strain across the two organ types (Figure **2e**), indicating that the majority of genes in nematode galls and roots that respond to rhizobial differences (83%) are responding to mutualist strain in similar ways in the two organs.

*Parasite infection impacts gene expression at the site of host-mutualist interactions in both strains of rhizobia*

Parasite infection caused little change in gene expression in rhizobia. WSM1022 showed no DEGs between infected and uninfected hosts and USDA1021 had only 13 DEGs (Figure **3b**,

Table S**1b**). In contrast, the host showed a large transcriptomic response in nodules to parasite infection. In nodules with rhizobia strain USDA1021, 22,993 *M. truncatula* genes (46.514% of the genome) were expressed. Of the expressed genes, 1,463 host genes (6.363%) were differentially expressed and similar numbers were up- and down-regulated in response to parasite infection (806 and 657 genes respectively) (Figure **3a**). Of these responding genes, 136 of them (9.30%) were conditionally expressed with 52 expressed only when parasites are present and 84 expressed only when parasites were absent. GOORA show that diverse processes are overrepresented in all 1,463 responsive host genes, but notably, terms related to host-mutualist interactions were overrepresented such as carbohydrate metabolic processes including oligosaccharide and glycogen metabolisms, energy reserve metabolism, defense response, transmembrane transport, structural constituents of cell walls, iron ion binding, and tripeptidyl-peptidase activity (Tables S**2b**,**c**,**d**; Dataset S**6**).

In nodules with WSM1022, 22,729 *M. truncatula* genes (44.003% of the genome) were expressed. Of the expressed genes, 475 genes (2.09%) were differentially expressed and the majority (433 genes; 91.2%) were downregulated in response to parasite infection (Figure **3a**). Of the responding genes, 25 of them (5.8%) were conditionally expressed with 6 of them expressed only in the presence of parasites and 19 of them expressed only in the absence of parasites. GOORA suggests that parasite infection may impact the expression of host defenses as processes related to defense were overrepresented in all 475 responsive host genes as annotations related to defense response, regulation of salicylic acid biosynthesis, response to chitin, stress response, signal transduction, and cell wall related terms were overrepresented (Table S**2e**; Dataset S**6**).

Figure 3: Nematode infection influences gene expression at the site of host-mutualist interactions in a largely strain specific manner. Log fold change (LFC) and standard error values were calculated using DESeq2. LFC values are the logarithm base 2 of the ratio of normalized mean expression in infected hosts to that in uninfected hosts. Genes with significant log fold changes determined by likelihood ratio tests performed by DESeq2 are indicated with color and also with standard error bars in d. All genes that were differentially expressed (DE) and functionally verified as critical to the functioning of the mutualism by Roy et al. (2019) have been circled in gold. a) 386 host genes were DE in response to nematode infection in nodules with WSM1022 and 1,428 were DE in nodules with USDA1021. b) 13 genes from USDA1021 and no genes from WSM1022 were DE in response to host nematode infection. c) The magnitude of host differential expression was higher in nodules with WSM1022. Differences in magnitude were tested using the optimal pooled t-test (B. Guo & Yuan, 2017) ('n.s.' indicates no significance, * indicates p-value < 0.05, ** indicates p-value < 0.01, *** indicates p-value < 0.001, **** indicates p-value < 0.0001). d) 1,632 host genes responded to parasite infection differently in nodules with different rhizobia strains (interaction effect) whereas only 193 genes responded similarly to parasite infection regardless of rhizobia strain (main effect).

*Parasite infection impacts host gene expression at the site of host-mutualist interactions in a largely mutualist strain specific manner*

Out of the 23,565 host genes expressed across all nodule samples, 230 genes (0.976%) responded similarly to parasite infection in nodules regardless of mutualist strain. On the other hand, 1,847 of the expressed genes (7.838%) showed a strain specific response to parasite infection. This indicates that the majority of genes responding to parasite infection in nodules (88.93%) respond in different ways to mutualist genotype (Figure **3d**). Notably many of these genes responded in opposite ways when different rhizobia were resident in the nodule. Of these genes showing a strain-specific response, 645 genes (34.9%) showed conditional expression in which the gene was not expressed in at least one treatment.

GOORA showed that processes potentially involved in host-mutualist interactions are overrepresented in the host genes showing a strain specific response to parasite infection. Notably, terms related to stress and defense responses, response to other organisms, cell communication and signal transduction, protein phosphorylation, autophosphorylation, and

kinase activity, carbohydrate and energy reserve metabolism of various types as well as the transferase or hydrolyzing activity of glycosyls or sugars, and various processes related to components or organization of the cell wall were overrepresented. Additionally, GO terms related to cellular context showed that the genes showing a strain specific response were overrepresented for anchored components of the plasma membrane, plasmodesma, and cell walls (Table **S2f**; see Dataset S6). As many of these functions are important for plant-rhizobia interactions, especially carbohydrate metabolism (Udvardi and Poole, 2013) and defense responses (Cao et al., 2017; Gourion et al., 2015), this suggests that the strain-specific effect of parasites on host gene expression in nodules may influence host-mutualist interactions.

The examination of individual genes of interest can underscore the potential functional import of the strain-specific response in the transcriptomic data. The leghemoglobin gene MtrunA17_Chr5g0427351, a critical component of creating an environment within nodules that is conducive to nitrogen fixation (Larrainzar et al., 2020), responded to parasite infection in a strain-specific manner (Dataset S3). Roy et al. (2019) compiled a list of 207 genes who have been functionally verified as integral to the proper functioning of the *M. truncatula-E. meliloti* mutualism. Out of these 207 genes, 25 genes were differentially expressed in nodules in a strain specific manner (interaction effect) while none of these genes showed a consistent response to nematode infection (main effect) across hosts regardless of rhizobia strain (Figure 3d).

*Rhizobia strain had no effect on root architecture while nematode infected plants had thicker and shorter roots with fewer root tips*

A number of the root architecture traits were highly correlated with one another (Figure S4; S5). Linear mixed models showed that nematode infection increased median root diameter ($\chi^2_{df=1} = 4.26$, p = 0.039) and average root diameter ($\chi^2_{df=1} = 7.14$, p = 0.008) and decreased root perimeter ($\chi^2_{df=1} = 5.61$, p = 0.018), total root length ($\chi^2_{df=1} = 4.66$, p = 0.031) and number of root tips ($\chi^2_{df=1} = 6.28$, p = 0.012) (Figure S6). However, host plants colonized by different rhizobia strains showed no difference in root architecture traits. The number of branch points, maximum root diameter, and network area were all unaffected by rhizobia strain and nematode infection and our models showed no interaction effect of rhizobia strain and nematode infection for any root architecture traits (Figure S6).

*Neither nematode infection nor rhizobia strain altered aboveground biomass, a proxy for host fitness*

Plants that formed more nodules had higher aboveground biomass (r = 0.46; Figure S4), confirming both strains of rhizobia used in this experiment confer benefits to their host plant. However, rhizobia strain was not a good predictor of above ground biomass (Figure 4d), indicating that the two strains did not differ in the benefits they provide their host. Neither nematode infection status nor parasite load (nematode gall count) affected aboveground biomass (Figure 4e), even when correcting for nodule count or root biomass. Together this suggests that, in concordance with other published data (Wood *et al.,* 2018), nematode infection status and nematode infection severity did not affect above ground biomass but nodulate rate did.

Figure 4: Rhizobia strain and nematode infection status could not explain variation in above ground biomass, but nodule count could. a) The above ground biomass of host plants did not

differ across rhizobia strain or nematode infection status. b) Plants differed in their nodules counts across both rhizobial strain and nematode infection status. Infected plants had fewer nodules than uninfected plants, and plants inoculated with USDA1021 had fewer nodules than plants inoculated with WSM1022. c) Nematode-infected plants had different number of nematode galls depending on the rhizobia strain they were inoculated with. Plants inoculated with WSM1022 had fewer nematode galls than plants inoculated with USDA1021. d) Nodule count was a good predictor of above ground biomass but neither rhizobia strain nor nematode infection status changed the relationship between nodule count and host plant above ground biomass. e) Nematode gall count was not a good predictor of above ground biomass.

*Nematode infection status impacted nodule count and rhizobia strain impacted nematode gall count*

As expected, plants with larger root systems had more rhizobia nodules and nematode nematode galls (Figure S4; S5). Nodule and nematode gall counts were positively correlated ($r = 0.42$), and a linear mixed model verified rhizobia colonization rates is a good predictor of nematode colonization rates ($\chi^2_{df=1} = 10.44$, $p = 0.001$; Figure S6). This held true even when correcting for root volume ($\chi^2_{df=1} = 3.86$, $p = 0.049$), indicating that the relationship between mutualist and parasite colonization is not merely a byproduct of overall plant size. Plants inoculated with WSM1022 had a lower mean nematode gall count than plants inoculated with USDA1022 ($\chi^2_{df=1} = 5.11$, $p = 0.024$; Figure 4c), even after correcting for root biomass ($\chi^2_{df=1} = 5.88$, $p = 0.015$). Nematode decreased the number of nodules ($\chi^2_{df=1} = 9.56$, $p = 0.002$; Figure 4b) even when corrected for by root biomass ($\chi^2_{df=1} = 12.96$, $p < 0.001$) and WSM1022 produced higher nodule counts than USDA1021 ($\chi^2_{df=1} = 8.14$, $p = 0.004$; Figure 4b) even when corrected for by root biomass ($\chi^2_{df=1} = 11.14$, $p < 0.001$). In summary, we saw that rhizobia strain had an effect on gall counts and nematode infection decreased nodule counts.

**Discussion**

Here we showed that intraspecific genetic variation in a major microbial mutualist impacts host-parasite interactions at the phenotypic and transcriptomic levels. Rhizobia strains affected their host's parasite load (Figure 4c). At the mature site of parasite infection (nematode galls), hundreds of genes were differentially expressed across plant hosts that were inoculated with one of two different strains of mutualistic rhizobia. At the mature site of mutualist colonization (rhizobial nodules), the effect of parasite infection on host gene expression depended on the resident mutualist strain. Our results suggest that microbial indirect genetic effects play an underappreciated role in structuring their host's interactions with other co-colonizing microorganisms at multiple stages of host-symbiont interaction.

*Microbial mutualists influence their host's parasite load and gene expression at the site of parasite infection*

Host plants inoculated with different strains of mutualistic rhizobia showed different numbers of nematode galls (Figure 4c) indicating that microbial mutualist genetic variation can affect the severity of parasite infection. This finding is in line with a variety of other studies that highlight the importance of microbial mutualists on traits that mediate other species interactions with hosts (Lange et al., 2023; Osborne et al., 2009; Parker et al., 2017). The difference in parasite infection severity that we observed was based on a count of mature parasitic structures and therefore could arise from differences in the speed at which parasite infection develops or differences in colonization rates due to varying host defenses or susceptibility.

Strain-level variation in a microbial mutualist was a major determinant of gene expression in parasite-infected hosts. This effect manifested in two ways. First, rhizobia had a comparable effect on host gene expression in nematode galls as in rhizobia nodules. The number of rhizobia-responsive plant genes in nematode galls was greater than in nodules, despite the magnitude of differential expression being slightly larger in nodules (Figure **2c**). Second, rhizobia largely determined the effect that parasite infection had on host gene expression. Hundreds of genes that were upregulated in response to parasite infection when the host was inoculated with one rhizobial strain were downregulated in response to parasite infection when inoculated with another strain (Figure **3c**).

Our results show that microbial effects on host gene expression extend to other species interactions their host engaged in. Furthermore, microbial effects on a host's third-party

interactions can rival their direct effect on hosts at their site of colonization. These results are consistent with and build on the growing body of evidence that microorganisms can strongly influence gene expression in their hosts: microbial modulation of host gene expression has been demonstrated in terrestrial and marine mutualisms (Burghardt et al., 2017; Heath et al., 2012; Palakurty et al., 2018; Rodriguez-Lanetty et al., 2006), in animal and plant pathogens (Apidianakis et al., 2005; Stringlis et al., 2018), and in the primate microbiome (Grieneisen et al., 2020).

*The consequences of cascading effects of mutualist strain variation*

Simple models of selection on mutualisms predict that selection should remove intraspecific variation in mutualism-related traits when, in reality, extensive intraspecific variation is a hallmark of many mutualism-related traits (Heath and Stinchcombe, 2014). This is true in the legume-rhizobia mutualism as rhizobia strains vary dramatically in the fitness benefit they provide their host (Heath, 2010). Similar patterns have been documented in mutualisms across the tree of life, from mycorrhizal fungi (Hart et al., 2013), to obligate pollinators (Jandér et al., 2012), to the bacterial symbionts of marine mussels (Ansorge et al., 2019). This disparity has motivated searches for mechanisms that maintain variation in mutualisms (Heath and Stinchcombe, 2014). Our experiment suggests that the mechanisms maintaining variation in mutualisms may have unanticipated and underappreciated consequences for the ecological communities in which mutualisms are embedded. However, predicting when and how intraspecific variation in mutualisms will cascade to impact other species interactions in host-associated communities requires understanding the mechanisms mediating crosstalk between microbial mutualists and other co-colonizing microorganisms.

*Potential mechanisms underlying mutualist strain effects on host-parasite interactions*

Why might strain-level variation in a microbial mutualist have such a large effect on the host's response to a parasite? We elaborate on three non-exclusive explanations below: a defense-centric, resource-centric, and development-centric model.

First, mutualist strains may elicit different defense responses in their host that may cascade to impact the expression of defense responses at the site of parasite infection. The establishment and maintenance of many mutualisms requires tightly coordinated control of the host's immune response (Zipfel and Oldroyd, 2017). In the legume-rhizobia mutualism,

suppression of the plant's immune response via nod factors and other effectors orchestrate the initiation and progression of the mutualism (Cao et al., 2017; Gourion et al., 2015). Mature nodules have a suppressed immune response, and functional genetic experiments have established that maintaining a functional symbiosis necessitates a controlled and tightly coordinated immune response (Zhang et al., 2023).

Consistent with this defense-centric model, our GOORA analysis of nematode gall samples uncovered enrichment of defense-related functions in DEGs. Notably, functions related to the plant hormone abscisic acid were overrepresented. Abscisic acid coordinates biotic and abiotic stress responses (Cao et al., 2011), although its role in plant responses to nematode infection remains poorly understood (Gheysen and Mitchum, 2019). changes in plant cell walls are a defense response against root knot nematode infection (Caillaud et al., 2008; Przybylska & Obrępalska-Stęplowska, 2020) and functions related to the metabolism of cell wall constituents were also overrepresented. Thus our findings about parasite severity may also be explained by rhizobia strains prompting varying defense responses in host plants that result in different degrees of host susceptibility.

Second, mutualist strains may differ in the resources they provide or costs they impose on their host. If strains differ in their impact on the host's resource budget, these differences could cascade to impact resource flux at the site of parasite infection or produce differences in the way mutualism resource fluxes respond to parasite infection. Intraspecific variation in the costs and benefits a mutualist provides is well documented in the legume-rhizobia mutualism (Heath, 2010; Calvert *et al.*, 2023) and rhizobia strains vary in the nitrogen and benefits they provide their plant host (Batstone et al., 2017; Terpolilli et al., 2008). If the strains in our experiment negotiated different carbon-for-nitrogen exchange rates with their host, the resulting impact on the host's resource budget could have driven differences in resource flux in nodules, nematode galls, or both, driving the transcriptomic differences we observed.

In support of this resource-centric model, our GOORA analyses suggest that rhizobia strain affected carbohydrate metabolism in nematode galls and nematode infection affected carbohydrate metabolism in nodules (Dataset S6). Carbohydrate metabolism plays a central function in the legume-rhizobia mutualism (Udvardi and Poole, 2013), and root knot nematodes siphon carbohydrates from the phloem, facilitated by active sugar transport into nematode galls (Zhou et al., 2023). Furthermore, the expression of leghemoglobin, a key regulator of the

nitrogen-fixation reaction in rhizobia nodules (Larrainzar et al., 2020), exhibited a strain-specific response to parasite infection. This observation raises the intriguing hypothesis that rhizobia strains control the sensitivity of nitrogen fixation—the central function of the legume-rhizobia mutualism—to parasite infection. Resource budget differences may also account for our phenotypic data, as parasite access to host resources can influence parasite infection severity or progress (Schultz *et al.*, 2013; Compson *et al.*, 2011). Differences in resource exchange across different rhizobia strains may alter host resource budgets in ways that impact the nematode's ability to siphon nutrients from its host, further impacting nematode infection severity, host defenses, or parasite pathogenesis.

Third, mutualist strains may differ in their impact on developmental processes. Symbiotic microbes are known to have profound effects on host development in both plants and animals on a systemic and local level (Fronk and Sachs, 2022; O'Brien et al., 2021a; Sommer and Bäckhed, 2013). Furthermore, resistance to infectious disease changes across ontogeny: juveniles are typically more susceptible to infection than adults (Ashby and Bruns, 2018; Izhar et al., 2020). If mutualist strains have different impacts on host systemic ontogeny and host ontogeny influences parasite resistance or infection trajectory, then hosts inoculated with different mutualists could be in different stages of infection when sampled at the same chronological age. On a local scale, mutualist strain differences may alter nematode gall or nodule development. Regardless of scale, in bulk RNAseq, differences in ontogeny can produce signals of differential expression if the ratio of different cell types in sequenced bulk organs differs across treatments of interest (Hunnicutt et al., 2022; Montgomery and Mank, 2016). Therefore, the differential expression that we observed may result from rhizobia strains influencing nematode gall, nodule, or systemic development such that cell composition in nematode galls or nodules differed across treatments. Developmental differences in hosts may also impact nematode infection severity as host ontogeny can influence parasite susceptibility and infection severity (Burr *et al*., 2022).

*Outstanding questions and future directions*

Our results suggest several promising avenues of future research. First, what are the relative contributions of defense-, resource-, and development-centric models to crosstalk in host-associated communities? Our GOORA implicates both the defense- and resource-centric models in mediating transcriptional crosstalk between the rhizobia mutualism and nematode parasitism but is unable to implicate development-centric models. Similarly, our phenotypic data

can be explained by all three models. Future experiments should independently manipulate host defense and resource flux to disentangle their independent contributions to crosstalk between co-infecting mutualists and parasites. Similarly, future experiments should be designed to examine different developmental stages or the contribution of different tissue types within symbiotic organs of the host-parasite interaction (Martinson *et al.*, 2022) to examine how rhizobial strain effects occur in different developmental stages. Experiments that implement sequencing technologies that de-confound the gene expression profiles of different cells or tissues, such as spatial transcriptomics (Rao et al., 2021) or scRNAseq (Saliba et al., 2014), could also shed light on the contribution of development to differential gene expression and phenotypic differences. Beyond illuminating the organismal processes that underlie crosstalk in host-associated communities, discriminating between defense-, resource-, and development-centric models is important to understand how these communities may evolve, as different underlying mechanisms implicate different regimes of pleiotropy and crosstalk that may constrain or alter evolutionary potential and shape organismal traits and species interactions. For example, if crosstalk is primarily mediated by the host's defense response, co-colonization may impact the evolution of genes and processes involved in host immunity. Alternatively, if crosstalk is primarily resource mediated, co-colonizers may influence the evolution of resource allocation and acquisition traits in their hosts.

Second, why was the transcriptomic response to co-colonization attenuated in symbionts relative to hosts? And why were changes in host gene expression in symbiotic structures not mirrored by the resident symbiont? In contrast to host gene expression, mutualist gene expression was largely unaffected by parasite infection, and parasite gene expression was largely unaffected by mutualist strain differences. Only 0-0.2% of rhizobia and nematode genes responded to their co-colonizer, while 1-6% of plant genes responded to variation in one or both symbionts. This observation suggests that the effect of mutualist strain on host-parasite interactions is primarily driven by the host. One straightforward explanation is that direct contact between species drives a larger transcriptomic response than indirect interactions. Another possibility is that generalist symbionts — like root-knot nematodes (Castagnone-Sereno et al., 2013; Catella et al., 2022), and, to a lesser extent, *Ensifer* rhizobia (Harrison et al., 2018) — have evolved transcriptional robustness across a wide variety of host environments.

Third, when does crosstalk in host-associated communities arise from systemic versus symbiosis-specific host responses? A significant insight from our experiment is that mutualist effects on gene expression at the site of parasite infection were specific to symbiotic organs and not a byproduct of systemic changes in host gene expression. Rhizobia elicited a different transcriptomic response in nodules and nematode galls than in uncolonized root tissue (Figure **2**). One possible explanation is that symbiosis-specific responses dominate whenever there is substantial overlap in the function of mature symbiotic structures induced by different colonizing species. Despite rhizobia being a mutualistic prokaryote and nematodes being a parasitic eukaryote, there is substantial overlap in their functions (Fronk and Sachs, 2022): both involve nutrient exchange, manipulation of the host defense response, and modification of host cell development and differentiation (Koltai et al., 2001). Regardless of the mechanism, our results underscore the importance of tissue specificity when studying the transcriptomic basis of host-symbiont interactions (Koltai et al., 2001; Przybylska and Obrępalska-Stęplowska, 2020).

Finally, our study raises a final fundamental question: How much genetic variation for traits mediating species interactions (e.g., parasite resistance and mutualism function) resides in species not directly involved in the interaction itself? Our results indicate that indirect genetic effects are important and overlooked coordinators of species interactions within multispecies contexts. The concept of indirect genetic effects—when one organism's genes influence another's phenotype—was originally formalized in the context of intraspecific interactions like aggression or maternal effects (Moore et al., 1997; Wolf et al., 1998). Recently, this framework was formally extended to interactions between species (De Lisle et al., 2022). However, this field primarily remains focused on pairwise species interactions: for example, how host genes influence parasite virulence or the benefit derived from a mutualistic symbiont. A significant insight from our study is that indirect genetic effects in species interactions extend to third parties. Given the potential that microbial communities have for impacting selection and overall health of their hosts (Lau and Lennon, 2011), future work should explore the role of intraspecific variation in microbial symbionts in shaping the structure and function of host-associated communities and its consequences for host health and fitness.

**References**

Ahmad, S., Gordon-Weeks, R., Pickett, J., Ton, J., 2010. Natural variation in priming of basal resistance: from evolutionary origin to agricultural exploitation. Mol. Plant Pathol. 11, 817–827. https://doi.org/10.1111/j.1364-3703.2010.00645.x

Alizon, S., de Roode, J.C., Michalakis, Y., 2013. Multiple infections and the evolution of virulence. Ecol. Lett. 16, 556–567. https://doi.org/10.1111/ele.12076

Ansorge, R., Romano, S., Sayavedra, L., Porras, M.Á.G., Kupczok, A., Tegetmeyer, H.E., Dubilier, N., Petersen, J., 2019. Functional diversity enables multiple symbiont strains to coexist in deep-sea mussels. Nat. Microbiol. 4, 2487–2497. https://doi.org/10.1038/s41564-019-0572-9

Apidianakis, Y., Mindrinos, M.N., Xiao, W., Lau, G.W., Baldini, R.L., Davis, R.W., Rahme, L.G., 2005. Profiling early infection responses: Pseudomonas aeruginosa eludes host defenses by suppressing antimicrobial peptide gene expression. Proc. Natl. Acad. Sci. 102, 2573–2578. https://doi.org/10.1073/pnas.0409588102

Ashby, B., Bruns, E., 2018. The evolution of juvenile susceptibility to infectious disease. Proc. R. Soc. B Biol. Sci. 285, 20180844. https://doi.org/10.1098/rspb.2018.0844

Banerjee, S., Schlaeppi, K., van der Heijden, M.G.A., 2018. Keystone taxa as drivers of microbiome structure and functioning. Nat. Rev. Microbiol. 16, 567–576. https://doi.org/10.1038/s41579-018-0024-1

Barker, D.G., Pfaff, T., Moreau, D., Groves, E., Ruffel, S., Lepetit, M., Whitehand, S., Maillet, F., Nair, R.M., Journet, E.-P., 2006. Growing M. truncatula: choice of substrates and growth conditions, in: Medicago Truncatula Handbook. Noble Institute, p. 26.

Barrett, L.G., Bever, J.D., Bissett, A., Thrall, P.H., 2015. Partner diversity and identity impacts on plant productivity in Acacia–rhizobial interactions. J. Ecol. 103, 130–142. https://doi.org/10.1111/1365-2745.12336

Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects Models Using lme4." Journal of Statistical Software, 67(1), 1–48. doi:10.18637/jss.v067.i01.

Batstone, R.T., Dutton, E.M., Wang, D., Yang, M., Frederickson, M.E., 2017. The evolution of symbiont preference traits in the model legume Medicago truncatula. New Phytol. 213, 1850–1861. https://doi.org/10.1111/nph.14308

Benjamin, G., Pandharikar, G., Frendo, P., 2022. Salicylic Acid in Plant Symbioses: Beyond

Plant Pathogen Interactions. Biology 11, 861. https://doi.org/10.3390/biology11060861

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

van der Bom, F. J., Williams, A., & Bell, M. J. (2020). Root architecture for improved resource capture: trade-offs in complex environments. Journal of Experimental Botany, 71(19), 5752-5763.

Bousema, J.T., Drakeley, C.J., Mens, P.F., Arens, T., Houben, R., Omar, S.A., Gouagna, L.C., Schallig, H., Sauerwein, R.W., 2008. Increased Plasmodium falciparum gametocyte production in mixed infections with P. malariae. Am. J. Trop. Med. Hyg. 78, 442–448.

Brown, C. Titus, Irber, Luiz., 2016. sourmash: a library for MinHash sketching of DNA. Journal of open source software 1.5, 27.

Burghardt, L.T., 2020. Evolving together, evolving apart: measuring the fitness of rhizobial bacteria in and out of symbiosis with leguminous plants. New Phytol. 228, 28–34. https://doi.org/10.1111/nph.16045

Burghardt, L.T., Epstein, B., Guhlin, J., Nelson, M.S., Taylor, M.R., Young, N.D., Sadowsky, M.J., Tiffin, P., 2018. Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. Proc. Natl. Acad. Sci. 115, 2425–2430. https://doi.org/10.1073/pnas.1714246115

Burghardt, L.T., Guhlin, J., Chun, C.L., Liu, J., Sadowsky, M.J., Stupar, R.M., Young, N.D., Tiffin, P., 2017. Transcriptomic basis of genome by genome variation in a legume-rhizobia mutualism. Mol. Ecol. 26, 6122–6135. https://doi.org/10.1111/mec.14285

Burr, A. A., Woods, K. D., Cassidy, S. T., & Wood, C. W. (2022). Priority effects alter the colonization success of a host‑associated parasite and mutualist. Ecology, 103(8), e3720.

Calvert, M. B., Hoque, M., & Wood, C. W. (2023). Toward a trait-based framework for studying mutualism evolution: genotypic variation in benefit and cost traits in the legume-rhizobia mutualism. bioRxiv, 2023-08.

Cao, F.Y., Yoshioka, K., Desveaux, D., 2011. The roles of ABA in plant–pathogen interactions. J. Plant Res. 124, 489–499. https://doi.org/10.1007/s10265-011-0409-y

Cao, Y., Halane, M.K., Gassmann, W., Stacey, G., 2017. The Role of Plant Innate Immunity in

the Legume-Rhizobium Symbiosis. Annu. Rev. Plant Biol. 68, 535–561. https://doi.org/10.1146/annurev-arplant-042916-041030

Carrï Re, S.B., Verdenaud, M., Gough, C., Gouzy, J.R.M., Gamas, P., 2020. LeGOO: An Expertized Knowledge Database for the Model Legume Medicago truncatula. Plant Cell Physiol. 61, 203–211. https://doi.org/10.1093/pcp/pcz177

Castagnone-Sereno, P., Danchin, E.G.J., Perfus-Barbeoch, L., Abad, P., 2013. Diversity and evolution of root-knot nematodes, genus Meloidogyne: new insights from the genomic era. Annu. Rev. Phytopathol. 51, 203–220. https://doi.org/10.1146/annurev-phyto-082712-102300

Catella, S.A., Olmsted, C.F., Markalanda, S.H., McFadden, C.J., Wood, C.W., Kuebbing, S.E., 2022. A generalist nematode destabilises plant competition: no evidence for direct effects, but strong evidence for indirect effects on rhizobium abundance. New Phytol. 233, 2561–2572. https://doi.org/10.1111/nph.17943

Chung, M., Bruno, V.M., Rasko, D.A., Cuomo, C.A., Muñoz, J.F., Livny, J., Shetty, A.C., Mahurkar, A., Dunning Hotopp, J.C., 2021. Best practices on the differential expression analysis of multi-species RNA-seq. Genome Biol. 22, 1–23. https://doi.org/10.1186/s13059-021-02337-8

Compson, Z. G., Larson, K. C., Zinkgraf, M. S., & Whitham, T. G. (2011). A genetic basis for the manipulation of sink–source relationships by the galling aphid Pemphigus batae. Oecologia, 167, 711-721.

Cook, D.R., 1999. Medicago truncatula-a model in the making! Curr. Opin. Plant Biol. 2, 301–304.

Coyte, K.Z., Schluter, J., Foster, K.R., 2015. The ecology of the microbiome: Networks, competition, and stability. Science 350, 663–666. https://doi.org/10.1126/science.aad2602

Dastogeer, K.M.G., Tumpa, F.H., Sultana, A., Akter, M.A., Chakraborty, A., 2020. Plant microbiome–an account of the factors that shape community composition and diversity. Curr. Plant Biol., Plant Interactions with Microbes and Environment 23, 100161. https://doi.org/10.1016/j.cpb.2020.100161

De Lisle, S.P., Bolnick, D.I., Brodie, E.D., Moore, A.J., McGlothlin, J.W., 2022. Interacting phenotypes and the coevolutionary process: Interspecific indirect genetic effects alter coevolutionary dynamics. Evol. Int. J. Org. Evol. 76, 429–444.

https://doi.org/10.1111/evo.14427

Debray, R., Herbert, R.A., Jaffe, A.L., Crits-Christoph, A., Power, M.E., Koskella, B., 2022. Priority effects in microbiome assembly. Nat. Rev. Microbiol. 20, 109–121. https://doi.org/10.1038/s41579-021-00604-w

Devi, P., Khan, A., Chattopadhyay, P., Mehta, P., Sahni, S., Sharma, S., Pandey, R., 2021. Co-infections as Modulators of Disease Outcome: Minor Players or Major Players? Front. Microbiol. 12.

Downie, J.A., Kondorosi, E., 2021. Why Should Nodule Cysteine-Rich (NCR) Peptides Be Absent From Nodules of Some Groups of Legumes but Essential for Symbiotic N-Fixation in Others? Front. Agron. 3.

Durgo, H., Klement, E., Hunyadi-Gulyas, E., Szucs, A., Kereszt, A., Medzihradszky, K.F., Kondorosi, E., 2015. Identification of nodule-specific cysteine-rich plant peptides in endosymbiotic bacteria. PROTEOMICS 15, 2291–2295. https://doi.org/10.1002/pmic.201400385

Espindula, E., Sperb, E. R., Bach, E., & Passaglia, L. M. P. (2020). The combined analysis as the best strategy for Dual RNA-Seq mapping. Genetics and Molecular Biology, 42, e20190215.

Foster, K.R., Schluter, J., Coyte, K.Z., Rakoff-Nahoum, S., 2017. The evolution of the host microbiome as an ecosystem on a leash. Nature 548, 43–51. https://doi.org/10.1038/nature23292

Fronk, D.C., Sachs, J.L., 2022. Symbiotic organs: the nexus of host–microbe evolution. Trends Ecol. Evol., Special issue: Symbiosis 37, 599–610. https://doi.org/10.1016/j.tree.2022.02.014

Gheysen, G., Mitchum, M.G., 2019. Phytoparasitic Nematode Control of Plant Hormone Pathways. Plant Physiol. 179, 1212–1226. https://doi.org/10.1104/pp.18.01067

Glöckner, F.O., Yilmaz, P., Quast, C., Gerken, J., Beccati, A., Ciuprina, A., Bruns, G., Yarza, P., Peplies, J., Westram, R., Ludwig, W., 2017. 25 years of serving the community with ribosomal RNA gene reference databases and tools. J. Biotechnol., Bioinformatics Solutions for Big Data Analysis in Life Sciences presented by the German Network for Bioinformatics Infrastructure 261, 169–176. https://doi.org/10.1016/j.jbiotec.2017.06.1198

Gourion, B., Berrabah, F., Ratet, P., Stacey, G., 2015. Rhizobium–legume symbioses: the crucial role of plant immunity. Trends Plant Sci. 20, 186–194. https://doi.org/10.1016/j.tplants.2014.11.008

Grieneisen, L., Muehlbauer, A.L., Blekhman, R., 2020. Microbial control of host gene regulation and the evolution of host–microbiome interactions in primates. Philos. Trans. R. Soc. B Biol. Sci. 375, 20190598. https://doi.org/10.1098/rstb.2019.0598

Guefrachi, I., Nagymihaly, M., Pislariu, C.I., Van de Velde, W., Ratet, P., Mars, M., Udvardi, M.K., Kondorosi, E., Mergaert, P., Alunni, B., 2014. Extreme specificity of NCR gene expression in Medicago truncatula. BMC Genomics 15, 712. https://doi.org/10.1186/1471-2164-15-712

Guo, B., Yuan, Y., 2017. A comparative review of methods for comparing means using partially paired data. Stat. Methods Med. Res. 26, 1323–1340. https://doi.org/10.1177/0962280215577111

Guo, H., Ge, F., 2017. Root nematode infection enhances leaf defense against whitefly in tomato. Arthropod-Plant Interact. 11, 23–33. https://doi.org/10.1007/s11829-016-9462-8

Guo, S., Wang, M., Song, X., Zhou, G., Kong, Y., 2022. The evolving views of the simplest pectic polysaccharides: homogalacturonan. Plant Cell Rep. 41, 2111–2123. https://doi.org/10.1007/s00299-022-02909-3

Harris, M.O., Pitzschke, A., 2020. Plants make galls to accommodate foreigners: some are friends, most are foes. New Phytol. 225, 1852–1872. https://doi.org/10.1111/nph.16340

Harrison, T.L., Simonsen, A.K., Stinchcombe, J.R., Frederickson, M.E., 2018. More partners, more ranges: generalist legumes spread more easily around the globe. Biol. Lett. 14, 20180616. https://doi.org/10.1098/rsbl.2018.0616

Hart, M.M., Forsythe, J., Oshowski, B., Bücking, H., Jansa, J., Kiers, E.T., 2013. Hiding in a crowd—does diversity facilitate persistence of a low-quality fungal partner in the mycorrhizal symbiosis? Symbiosis 59, 47–56. https://doi.org/10.1007/s13199-012-0197-8

Hartig, F. (2018). DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models.

Hasegawa, S., Nagel, C.W., 1968. Isolation of an oligogalacturonate hydrolase from a Bacillus specie. Arch. Biochem. Biophys. 124, 513–520.

https://doi.org/10.1016/0003-9861(68)90360-3

Heath, K.D., 2010. Intergenomic epistasis and coevolutionary constrain in plants and rhizobia. Evolution 64, 1446–1458. https://doi.org/10.1111/j.1558-5646.2009.00913.x

Heath, K.D., Burke, P.V., Stinchcombe, J.R., 2012. Coevolutionary genetic variation in the legume-rhizobium transcriptome. Mol. Ecol. 21, 4735–4747. https://doi.org/10.1111/j.1365-294X.2012.05629.x

Heath, K.D., Stinchcombe, J.R., 2014. Explaining mutualism variation: a new evolutionary paradox? Evolution 68, 309–317. https://doi.org/10.1111/evo.12292

Heath, K.D., Tiffin, P., 2007. Context dependence in the coevolution of plant and rhizobial mutualists. Proc. R. Soc. B Biol. Sci. 274, 1905–1912. https://doi.org/10.1098/rspb.2007.0495

Hebenstreit, D., Fang, M., Gu, M., Charoensawan, V., van Oudenaarden, A., Teichmann, S.A., 2011. RNA sequencing reveals two major classes of gene expression levels in metazoan cells. Mol. Syst. Biol. 7, 497. https://doi.org/10.1038/msb.2011.28

Hunnicutt, K.E., Good, J.M., Larson, E.L., 2022. Unraveling patterns of disrupted gene expression across a complex tissue. Evolution 76, 275–291. https://doi.org/10.1111/evo.14420

Izhar, R., Gilboa, C., Ben-Ami, F., 2020. Disentangling the steps of the infection process responsible for juvenile disease susceptibility. Funct. Ecol. 34, 1551–1563. https://doi.org/10.1111/1365-2435.13580

Jagdale, S., Rao, U., Giri, A.P., 2021. Effectors of Root-Knot Nematodes: An Arsenal for Successful Parasitism. Front. Plant Sci. 12.

Jain, Chirag, Rodriguez-R, Luis M., Phillippy, Adam M., Konstantinidis, Konstantinos T., Aluru, Srinivas, 2018. High Throughput ANI Analysis of 90K Prokaryotic Genomes Reveals Clear Species Boundaries. Nature Communications. 9 (1), 5114.

Jandér, K.C., Herre, E.A., Simms, E.L., 2012. Precision of host sanctions in the fig tree–fig wasp mutualism: consequences for uncooperative symbionts. Ecol. Lett. 15, 1362–1369. https://doi.org/10.1111/j.1461-0248.2012.01857.x

Jones, M.G.K., Goto, D.B., 2011. Root-knot Nematodes and Giant Cells, in: Jones, J., Gheysen, G., Fenoll, C. (Eds.), Genomics and Molecular Genetics of Plant-Nematode Interactions. Springer Netherlands, Dordrecht, pp. 83–100.

https://doi.org/10.1007/978-94-007-0434-3_5

Journet, E.-P., de Carvalho-Niebel, F., Andriankaja, A., Huguet, T., Barker, D.G., 2006. Rhizobial inoculation and nodulation of Medicago truncatula, in: Medicago Truncatula Handbook. Noble Institute, p. 6.

Kim, D., Paggi, J.M., Park, C., Bennett, C., Salzberg, S.L., 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37, 907–915. https://doi.org/10.1038/s41587-019-0201-4

Koltai, H., Dhandaydham, M., Opperman, C., Thomas, J., Bird, D., 2001. Overlapping Plant Signal Transduction Pathways Induced by a Parasitic Nematode and a Rhizobial Endosymbiont. Mol. Plant-Microbe Interactions® 14, 1168–1177. https://doi.org/10.1094/MPMI.2001.14.10.1168

Kopylova, E., Noé, L., Touzet, H., 2012. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. Bioinformatics 28, 3211–3217. https://doi.org/10.1093/bioinformatics/bts611

Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests in Linear Mixed Effects Models." Journal of Statistical Software, 82(13), 1–26. doi:10.18637/jss.v082.i13.

Lange, C., Boyer, S., Bezemer, T.M., Lefort, M.-C., Dhami, M.K., Biggs, E., Groenteman, R., Fowler, S.V., Paynter, Q., Verdecia Mogena, A.M., Kaltenpoth, M., 2023. Impact of intraspecific variation in insect microbiomes on host phenotype and evolution. ISME J. 1–10. https://doi.org/10.1038/s41396-023-01500-2

Larimer, A.L., Clay, K., Bever, J.D., 2014. Synergism and context dependency of interactions between arbuscular mycorrhizal fungi and rhizobia with a prairie legume. Ecology 95, 1045–1054. https://doi.org/10.1890/13-0025.1

Larrainzar, E., Villar, I., Rubio, M.C., Pérez-Rontomé, C., Huertas, R., Sato, S., Mun, J.-H., Becana, M., 2020. Hemoglobins in the legume–Rhizobium symbiosis. New Phytol. 228, 472–484. https://doi.org/10.1111/nph.16673

Lau, J.A., Lennon, J.T., 2011. Evolutionary ecology of plant–microbe interactions: soil microbial structure alters selection on plant traits. New Phytol. 192, 215–224. https://doi.org/10.1111/j.1469-8137.2011.03790.x

Li, Y., Pei, Y., Shen, Y., Zhang, R., Kang, M., Ma, Y., ... & Chen, Y. (2022). Progress in the

self-regulation system in legume nodule development-AON (autoregulation of nodulation). International Journal of Molecular Sciences, 23(12), 6676.

Lively, C.M., Dybdahl, M.F., 2000. Parasite adaptation to locally common host genotypes. Nature 405, 679–681. https://doi.org/10.1038/35015069

Long, S.R., 1989. Rhizobium-legume nodulation: Life together in the underground. Cell 56, 203–214. https://doi.org/10.1016/0092-8674(89)90893-3

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 1–21. https://doi.org/10.1186/s13059-014-0550-8

Magnoli, S.M., Bever, J.D., 2023. Plant productivity response to inter- and intra-symbiont diversity: Mechanisms, manifestations and meta-analyses. Ecol. Lett. 26, 1614–1628. https://doi.org/10.1111/ele.14274

Martinson, E. O., Werren, J. H., & Egan, S. P. (2022). Tissue‑specific gene expression shows a cynipid wasp repurposes oak host gene networks to create a complex and novel parasite‑specific organ. Molecular Ecology, 31(11), 3228-3240.

Mauch-Mani, B., Baccelli, I., Luna, E., & Flors, V. (2017). Defense priming: an adaptive part of induced resistance. Annual review of plant biology, 68(1), 485-512.

Miller, T.E., Travis, J., 1996. The evolutionary role of indirect effects in communities. Ecology 77, 1329.

Montgomery, S.H., Mank, J.E., 2016. Inferring regulatory change from gene expression: the confounding effects of tissue scaling. Mol. Ecol. 25, 5114–5128. https://doi.org/10.1111/mec.13824

Moore, A.J., Brodie, E.D., Wolf, J.B., 1997. Interacting phenotypes and the evoultionary process: I. Direct and indriect genetic effects of social interactions. Evol. Int. J. Org. Evol. 51, 1352–1362. https://doi.org/10.1111/j.1558-5646.1997.tb01458.x

Morand, S., Manning, S.D., Woolhouse, M.E.J., 1997. Parasite—host coevolution and geographic patterns of parasite infectivity and host susceptibility. Proc. R. Soc. Lond. B Biol. Sci. 263, 119–128. https://doi.org/10.1098/rspb.1996.0019

Moreau, D., Voisin, A.-S., Salon, C., Munier-Jolain, N., 2008. The model symbiotic association between Medicago truncatula cv. Jemalong and Rhizobium meliloti strain 2011 leads to N-stressed plants when symbiotic N2 fixation is the main N source for plant growth. J.

Exp. Bot. 59, 3509–3522. https://doi.org/10.1093/jxb/ern203

Nguyen Ba, A.N., Pogoutse, A., Provart, N., Moses, A.M., 2009. NLStradamus: a simple Hidden Markov Model for nuclear localization signal prediction. BMC Bioinformatics 10, 202. https://doi.org/10.1186/1471-2105-10-202

Nguyen, C.-N., Perfus-Barbeoch, L., Quentin, M., Zhao, J., Magliano, M., Marteu, N., Da Rocha, M., Nottet, N., Abad, P., Favery, B., 2018. A root-knot nematode small glycine and cysteine-rich secreted effector, MiSGCR1, is involved in plant parasitism. New Phytol. 217, 687–699. https://doi.org/10.1111/nph.14837

O'Brien, A.M., Ginnan, N.A., Rebolleda-Gómez, M., Wagner, M.R., 2021a. Microbial effects on plant phenology and fitness. Am. J. Bot. 108, 1824–1837. https://doi.org/10.1002/ajb2.1743

O'Brien, A.M., Jack, C.N., Friesen, M.L., Frederickson, M.E., 2021b. Whose trait is it anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. Proc. R. Soc. B Biol. Sci. 288, 20202483. https://doi.org/10.1098/rspb.2020.2483

Oates, C. N., Denby, K. J., Myburg, A. A., Slippers, B., & Naidoo, S. (2021). Insect egg‐induced physiological changes and transcriptional reprogramming leading to gall formation. Plant, Cell & Environment, 44(2), 535-547.

Opperman, C.H., Bird, D.M., Williamson, V.M., Rokhsar, D.S., Burke, M., Cohn, J., Cromer, J., Diener, S., Gajan, J., Graham, S., Houfek, T.D., Liu, Q., Mitros, T., Schaff, J., Schaffer, R., Scholl, E., Sosinski, B.R., Thomas, V.P., Windham, E., 2008. Sequence and genetic map of Meloidogyne hapla: A compact nematode genome for plant parasitism. Proc. Natl. Acad. Sci. U. S. A. 105, 14802–14807. https://doi.org/10.1073/pnas.0805946105

Osborne, S.E., Leong, Y.S., O'Neill, S.L., Johnson, K.N., 2009. Variation in Antiviral Protection Mediated by Different Wolbachia Strains in Drosophila simulans. PLOS Pathog. 5, e1000656. https://doi.org/10.1371/journal.ppat.1000656

Palakurty, S.X., Stinchcombe, J.R., Afkhami, M.E., 2018. Cooperation and coexpression: How coexpression networks shift in response to multiple mutualists. Mol. Ecol. 27, 1860–1873. https://doi.org/10.1111/mec.14550

Pan, H., Wang, D., 2017. Nodule cysteine-rich peptides maintain a working balance during nitrogen-fixing symbiosis. Nat. Plants 3, 1–6. https://doi.org/10.1038/nplants.2017.48

Parker, B.J., Hrček, J., McLean, A.H.C., Godfray, H.C.J., 2017. Genotype specificity among

hosts, pathogens, and beneficial microbes influences the strength of symbiont-mediated protection. Evolution 71, 1222–1231. https://doi.org/10.1111/evo.13216

Pecrix, Y., Staton, S.E., Sallet, E., Lelandais-Brière, C., Moreau, S., Carrère, S., Blein, T., Jardinaud, M.-F., Latrasse, D., Zouine, M., Zahm, M., Kreplak, J., Mayjonade, B., Satgé, C., Perez, M., Cauet, S., Marande, W., Chantry-Darmon, C., Lopez-Roques, C., Bouchez, O., Bérard, A., Debellé, F., Muños, S., Bendahmane, A., Bergès, H., Niebel, A., Buitink, J., Frugier, F., Benhamed, M., Crespi, M., Gouzy, J., Gamas, P., 2018. Whole-genome landscape of Medicago truncatula symbiotic genes. Nat. Plants 4, 1017–1025. https://doi.org/10.1038/s41477-018-0286-7

Pedersen, A.B., Fenton, A., 2007. Emphasizing the ecology in parasite community ecology. Trends Ecol. Evol. 22, 133–139. https://doi.org/10.1016/j.tree.2006.11.005

Pieterse, C. M., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C., & Bakker, P. A. (2014). Induced systemic resistance by beneficial microbes. Annual review of phytopathology, 52(1), 347-375.

Pollitt, L.C., Bram, J.T., Blanford, S., Jones, M.J., Read, A.F., 2015. Existing Infection Facilitates Establishment and Density of Malaria Parasites in Their Mosquito Vector. PLOS Pathog. 11, e1005003. https://doi.org/10.1371/journal.ppat.1005003

Pruesse, E., Quast, C., Knittel, K., Fuchs, B.M., Ludwig, W., Peplies, J., Glöckner, F.O., 2007. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. Nucleic Acids Res. 35, 7188–7196. https://doi.org/10.1093/nar/gkm864

Przybylska, A., Obrępalska-Stęplowska, A., 2020. Plant defense responses in monocotyledonous and dicotyledonous host plants during root-knot nematode infection. Plant Soil 451, 239–260. https://doi.org/10.1007/s11104-020-04533-0

Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E., Zanini, F., 2022. Analysing high-throughput sequencing data in Python with HTSeq 2.0. Bioinformatics 38, 2943–2945. https://doi.org/10.1093/bioinformatics/btac166

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res. 41, D590–D596. https://doi.org/10.1093/nar/gks1219

Queller, D.C., 2014. Joint phenotypes, evolutionary conflict and the fundamental theorem of natural selection. Philos. Trans. R. Soc. B Biol. Sci. 369, 20130423. https://doi.org/10.1098/rstb.2013.0423

Rao, A., Barkley, D., França, G.S., Yanai, I., 2021. Exploring tissue architecture using spatial transcriptomics. Nature 596, 211–220. https://doi.org/10.1038/s41586-021-03634-9

Rodriguez-Lanetty, M., Phillips, W.S., Weis, V.M., 2006. Transcriptome analysis of a cnidarian – dinoflagellate mutualism reveals complex modulation of host gene expression. BMC Genomics 7, 23. https://doi.org/10.1186/1471-2164-7-23

Roy, S., Liu, W., Nandety, R.S., Crook, A., Mysore, K.S., Pislariu, C.I., Frugoli, J., Dickstein, R., Udvardi, M.K., 2020. Celebrating 20 Years of Genetic Discoveries in Legume Nodulation and Symbiotic Nitrogen Fixation. Plant Cell 32, 15–41. https://doi.org/10.1105/tpc.19.00279

Saliba, A.-E., Westermann, A.J., Gorski, S.A., Vogel, J., 2014. Single-cell RNA-seq: advances and future challenges. Nucleic Acids Res. 42, 8845–8860. https://doi.org/10.1093/nar/gku555

Schneider, P., Greischar, M.A., Birget, P.L.G., Repton, C., Mideo, N., Reece, S.E., 2018. Adaptive plasticity in the gametocyte conversion rate of malaria parasites. PLOS Pathog. 14, e1007371. https://doi.org/10.1371/journal.ppat.1007371

Schultz, J. C., Appel, H. M., Ferrieri, A. P., & Arnold, T. M. (2013). Flexible resource allocation during plant defense responses. Frontiers in plant science, 4, 324.

Seemann, Torsten, 2014. Prokka: Rapid Prokaryotic Genome Annotation. Bioinformatics 30, 14, 2068–69.

Seethepalli, A., Dhakal, K., Griffiths, M., Guo, H., Freschet, G. T., York, L. M. (2021). RhizoVision Explorer: Open-source software for root image analysis and measurement standardization. AoB PLANTS, plab056, https://doi.org/10.1093/aobpla/plab056

Seethepalli, A. and York, L. M. (2020). RhizoVision Explorer - Interactive software for generalized root image analysis designed for everyone (Version 2.0.3). Zenodo. http://doi.org/10.5281/zenodo.4095629

Shen, Y., Ma, Y., Li, D., Kang, M., Pei, Y., Zhang, R., Tao, W., Huang, S., Song, W., Li, Y., Huang, W., Wang, D., Chen, Y., 2023. Biological and genomic analysis of a symbiotic nitrogen fixation defective mutant in Medicago truncatula. Front. Plant Sci. 14, 1209664.

https://doi.org/10.3389/fpls.2023.1209664

Shetty, S.A., Hugenholtz, F., Lahti, L., Smidt, H., de Vos, W.M., 2017. Intestinal microbiome landscaping: insight in community assemblage and implications for microbial modulation strategies. FEMS Microbiol. Rev. 41, 182–199. https://doi.org/10.1093/femsre/fuw045

Shuster, S.M., Lonsdorf, E.V., Wimp, G.M., Bailey, J.K., Whitham, T.G., 2006. Community heritability measures the evolutionary consequences of indirect effects on community structure. Evolution 60, 991–1003.

Smith, A. G., Han, E., Petersen, J., Olsen, N. A. F., Giese, C., Athmann, M., ... & Thorup‑Kristensen, K. (2022). RootPainter: deep learning segmentation of biological images with corrective annotation. New Phytologist, 236(2), 774-791.

Sommer, F., Bäckhed, F., 2013. The gut microbiota — masters of host development and physiology. Nat. Rev. Microbiol. 11, 227–238. https://doi.org/10.1038/nrmicro2974

Song, J., Mao, H., Cheng, J., Zhou, Y., Chen, R., Zeng, L., ... & Wang, Y. (2021). Identification of the trehalose-6-phosphate synthase gene family in Medicago truncatula and expression analysis under abiotic stresses. Gene, 787, 145641.

Stein, R.R., Bucci, V., Toussaint, N.C., Buffie, C.G., Rätsch, G., Pamer, E.G., Sander, C., Xavier, J.B., 2013. Ecological Modeling from Time-Series Inference: Insight into Dynamics and Stability of Intestinal Microbiota. PLOS Comput. Biol. 9, e1003388. https://doi.org/10.1371/journal.pcbi.1003388

Strauss, S.Y., 1991. Indirect Effects in Community Ecology:Their Definition, Study and Importance 6.

Stringlis, I.A., Proietti, S., Hickman, R., Van Verk, M.C., Zamioudis, C., Pieterse, C.M.J., 2018. Root transcriptional dynamics induced by beneficial rhizobacteria and microbial immune elicitors reveal signatures of adaptation to mutualists. Plant J. 93, 166–180. https://doi.org/10.1111/tpj.13741

Tan, S., Debellé, F., Gamas, P., Frugier, F., Brault, M., 2019. Diversification of cytokinin phosphotransfer signaling genes in Medicago truncatula and other legume genomes. BMC Genomics 20, 373. https://doi.org/10.1186/s12864-019-5724-z

Telfer, S., Lambin, X., Birtles, R., Beldomenico, P., Burthe, S., Paterson, S., Begon, M., 2010. Species Interactions in a Parasite Community Drive Infection Risk in a Wildlife Population. Science 330, 243–246. https://doi.org/10.1126/science.1190333

terHorst, C.P., Zee, P.C., Heath, K.D., Miller, T.E., Pastore, A.I., Patel, S., Schreiber, S.J., Wade, M.J., Walsh, M.R., 2018. Evolution in a Community Context: Trait Responses to Multiple Species Interactions. Am. Nat. 191, 368–380. https://doi.org/10.1086/695835

Terpolilli, J., Hill, Y., Tian, R., Howieson, J., Bräu, L., Goodwin, L., ... & Reeve, W. (2013). Genome sequence of Ensifer meliloti strain WSM1022; a highly effective microsymbiont of the model legume Medicago truncatula A17. Standards in Genomic Sciences, 9, 315-324.

Terpolilli, J.J., O'Hara, G.W., Tiwari, R.P., Dilworth, M.J., Howieson, J.G., 2008. The model legume Medicago truncatula A17 is poorly matched for N2 fixation with the sequenced microsymbiont Sinorhizobium meliloti 1021. New Phytol. 179, 62–66. https://doi.org/10.1111/j.1469-8137.2008.02464.x

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gíslason, M.H., Pihl, S.I., Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., Nielsen, H., 2022. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat. Biotechnol. 40, 1023–1025. https://doi.org/10.1038/s41587-021-01156-3

Thies, J. A., Merrill, S. B., & Corley, E. L. (2002). Red food coloring stain: New, safer procedures for staining nematodes in roots and egg masses on root surfaces. Journal of Nematology, 34(2), 179.

Tonkin-Hill, Gerry, MacAlasdair, Neil, Ruis, Christopher, Weimann, Aaron, Horesh, Gal, Lees, John A., Gladstone, Rebecca A., 2020. Producing Polished Prokaryotic Pangenomes with the Panaroo Pipeline. Genome Biology. 21, 1, 180.

Udvardi, M., Poole, P.S., 2013. Transport and Metabolism in Legume-Rhizobia Symbioses. Annu. Rev. Plant Biol. 64, 781–805. https://doi.org/10.1146/annurev-arplant-050312-120235

Venter, F., Matthews, K.R., Silvester, E., 2022. Parasite co-infection: an ecological, molecular and experimental perspective. Proc. R. Soc. B Biol. Sci. 289, 20212155. https://doi.org/10.1098/rspb.2021.2155

Weldon, S.R., Russell, J.A., Oliver, K.M., 2020. More Is Not Always Better: Coinfections with Defensive Symbionts Generate Highly Variable Outcomes. Appl. Environ. Microbiol. 86, e02537-19. https://doi.org/10.1128/AEM.02537-19

Westermann, A.J., Barquist, L., Vogel, J., 2017. Resolving host–pathogen interactions by dual

RNA-seq. PLOS Pathog. 13, e1006033. https://doi.org/10.1371/journal.ppat.1006033

Westermann, A.J., Vogel, J., 2021. Cross-species RNA-seq for deciphering host–microbe interactions. Nat. Rev. Genet. 22, 361–378. https://doi.org/10.1038/s41576-021-00326-y

Wieczorek, K., Elashry, A., Quentin, M., Grundler, F.M.W., Favery, B., Seifert, G.J., Bohlmann, H., 2014. A Distinct Role of Pectate Lyases in the Formation of Feeding Structures Induced by Cyst and Root-Knot Nematodes. Mol. Plant-Microbe Interactions® 27, 901–912. https://doi.org/10.1094/MPMI-01-14-0005-R

Wolf, J.B., Brodie III, E.D., Cheverud, J.M., Moore, A.J., Wade, M.J., 1998. Evolutionary consequences of indirect genetic effects. Trends Ecol. Evol. 13, 64–69. https://doi.org/10.1016/S0169-5347(97)01233-0

Wolf, S., Mouille, G., Pelloux, J., 2009. Homogalacturonan Methyl-Esterification and Plant Development. Mol. Plant 2, 851–860. https://doi.org/10.1093/mp/ssp066

Wolf, T., Kämmer, P., Brunke, S., Linde, J., 2018. Two's company: studying interspecies relationships with dual RNA-seq. Curr. Opin. Microbiol., Cell Regulation 42, 7–12. https://doi.org/10.1016/j.mib.2017.09.001

Wood, C.W., Pilkington, B.L., Vaidya, P., Biel, C., Stinchcombe, J.R., 2018. Genetic conflict with a parasitic nematode disrupts the legume–rhizobia mutualism. Evol. Lett. 2, 233–245. https://doi.org/10.1002/evl3.51

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X., Yu, G., 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation 2, 100141. https://doi.org/10.1016/j.xinn.2021.100141

Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., Glöckner, F.O., 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res. 42, D643–D648. https://doi.org/10.1093/nar/gkt1209

York, L. M. (2023) Root scanning using a flatbed scanner with transparency unit. Zenodo. doi.org/10.5281/zenodo.7825893

Younginger, B. S., Sirová, D., Cruzan, M. B., & Ballhorn, D. J. (2017). Is biomass a reliable estimate of plant fitness?. Applications in plant sciences, 5(2), 1600094.

Zamioudis, C., Pieterse, C.M.J., 2012. Modulation of Host Immunity by Beneficial Microbes.

Mol. Plant-Microbe Interactions® 25, 139–150.
https://doi.org/10.1094/MPMI-06-11-0179

Zhang, D., Wu, Q., Zhao, Y., Yan, Z., Xiao, A., Yu, H., Cao, Y., 2023. Dual RNA-Seq Analysis
Pinpoints a Balanced Regulation between Symbiosis and Immunity in Medicago
truncatula-Sinorhizobium meliloti Symbiotic Nodules. Int. J. Mol. Sci. 24, 16178.
https://doi.org/10.3390/ijms242216178

Zhou, Y., Zhao, D., Duan, Y., Chen, L., Fan, H., Wang, Y., Liu, X., Chen, L.-Q., Xuan, Y., Zhu,
X., 2023. AtSWEET1 negatively regulates plant susceptibility to root-knot nematode
disease. Front. Plant Sci. 14.

Zipfel, C., Oldroyd, G.E.D., 2017. Plant signaling in symbiosis and immunity. Nature 543,
328–336. https://doi.org/10.1038/nature22009

**Supporting Information**

The following Supporting Information is available for this article:

**Dataset S1**: A comma-separated values file containing gene expression quantification data output from HTSeq-union for all genes in all organisms.

**Dataset S2**: A comma-separated values file containing gene expression status for all genes in all samples.

**Dataset S3**: A comma-separated values file containing the results of all DESeq2 differential expression analyses for all *Medicago truncatula* genes.

**Dataset S4**: A comma-separated values file containing the results of all DESeq2 differential expression analyses for all *Ensifer meliloti* genes.

**Dataset S5**: A comma-separated values file containing the results of all DESeq2 differential expression analyses for all *Meloidogyne hapla* genes.

**Dataset S6**: A comma-separated values file containing all Gene Ontology overrepresentation analysis results.

**Figure S1**: Primary component analysis of *M. truncatula* gene expression quantification data showing segregation of datapoints by organ type.

**Figure S2**: Upset plot showing the intersection of *M. truncatula* gene lists produced by differential expression analyses comparing similar and dissimilar responses to rhizobial strain differences across different groups of organs.

**Figure S3**: Diagrams showing the expression and differential expression status of *M. truncatula* genes in different organ types.

**Figure S4:** Plot showing the correlation between root architecture traits and nodules counts across all samples.

**Figure S5:** Plot showing the correlation between root architecture traits and nodule and gall counts across all samples that were infected with nematodes.

**Figure S6:** Plot showing root architecture traits separated by nematode infection status and rhizobia strain.

**Table S1:** Gene lists of interest.

**Table S2**: Select Gene Ontology terms that were enriched in various gene lists produced by differential expression analyses.

**Table S3**: Constituents of the various fertilizer solutions used in this experiment.

**Table S4**: Symbiont treatment (nematode status and rhizobial strain) and host organ of origin for each sample.

**Table S5**: Parameters for the DESeq2 likelihood ratio tests performed for each model.

**Table S6**: Table showing the percentage of genes in each organ and treatment that had a coefficient of variation within the treatment that was greater than the coefficient of variation across all samples.

**Table S7**: Table showing total reads and percentage of reads that are filtered or aligned at various bioinformatic steps.

**Methods S1**: Sample preparation

**Methods S2**: RNA extraction and sequencing

**Methods S3**: Software and reference genomes

**Methods S4**: Differential expression analysis

**Results S1:** Nodule cysteine-rich protein gene expression is impacted by parasite infection