Local nuclear to cytoplasmic ratio regulates H3.3 incorporation via cell

cycle state during zygotic genome activation

Anusha D. Bhatt[1], Madeleine G. Brown[1], Aurora B. Wackford[1], Yuki Shindo[1], Amanda A.

Amodeo[1,2,*]

[1]Department of Biological sciences, Dartmouth College, Hanover, NH 03755, USA

[2] Lead contact

*Correspondence: amanda.amodeo@dartmouth.edu

Abstract

Early embryos often have unique chromatin states prior to zygotic genome activation (ZGA). In *Drosophila*, ZGA occurs after 13 reductive nuclear divisions during which the nuclear to cytoplasmic (N/C) ratio grows exponentially. Previous work found that histone H3 chromatin incorporation decreases while its variant H3.3 increases leading up to ZGA. In other cell types, H3.3 is associated with sites of active transcription and heterochromatin, suggesting a link between H3.3 and ZGA. Here, we test what factors regulate H3.3 incorporation at ZGA. We find that H3 nuclear availability falls more rapidly than H3.3 leading up to ZGA. We generate H3/H3.3 chimeric proteins at the endogenous H3.3A locus and observe that chaperone binding, but not gene structure, regulates H3.3 behavior. We identify the N/C ratio as a major determinant of H3.3 incorporation. To isolate how the N/C ratio regulates H3.3 incorporation we test the roles of genomic content, zygotic transcription, and cell cycle state. We determine that cell cycle regulation, but not H3 availability or transcription, controls H3.3 incorporation. Overall, we propose that local N/C ratios control histone variant usage via cell cycle state during ZGA.

Keywords: Zygotic genome activation, nuclear to cytoplasmic ratio, histones, chromatin, transcription, cell cycle

Introduction

Genome accessibility can be dynamically regulated through controlled incorporation of variant histones[1–3]. In most tissues, replication-coupled (RC) histones, produced during S-phase, generate the majority of nucleosomes[2,4,5]. RC histones have unusually high copy number, lack introns, and contain specialized UTRs to facilitate their rapid production during S-phase[3,6–9]. Conversely, replication-independent (RI), "variant" histones are made throughout the cell cycle and incorporated into specific genomic regions[4,10]. The exchange of RC and RI histones on chromatin is a common feature of early embryonic development especially during zygotic genome activation (ZGA)[11–17]. During ZGA, chromatin undergoes extensive remodeling to facilitate bulk transcription and establish heterochromatin[18–25].

The pre-ZGA cell cycles in many organisms depend on maternally supplied components, including histones[26–29]. These cycles are unusual since they oscillate between S and M without growth phases, leading to an exponential increase in the nuclear to cytoplasmic (N/C) ratio[29–34]. The N/C ratio, in turn, controls the timing of cell cycle slowing and ZGA[31–33,35–37]. Titration of maternal histones against the increasing amount of DNA has been proposed to contribute to N/C ratio sensing in the early embryo[38–45]. Another hallmark of ZGA is histone variant exchange on chromatin. In many organisms, maternally supplied, embryonic-specific linker histone variants are replaced by RC H1s during ZGA[12–14,16,17]. Concurrently, the RC nucleosomal H2A is also replaced by RI H2Av as a consequence of the lengthened interphase in cycles leading up to ZGA in *Drosophila*[15,46]. Similarly, we

have previously shown that RC H3 is replaced by RI H3.3 during these same cycles,

though the cause remains unclear[29].

H3.3 is essential for proper embryonic development in mice, *Xenopus*, and zebrafish[47–

51]. In *Xenopus,* the H3.3-specific S31 residue is required for gastrulation while its

chaperone binding site is dispensable[50]. In *Drosophila*, H3.3 nulls survive until adulthood

using maternal H3.3 but are sterile[52]. Flies expressing H3 from the H3.3 enhancer

generated conflicting results as to whether H3.3 protein or simply a source of replication-

independent H3 is required for fertility[52,53]. The H3/H3.3 pair is particularly interesting

during ZGA because H3.3 is enriched at sites of active transcription and in

heterochromatin, which are both established during ZGA[2,4,54].

Here, we examine the factors that contribute to H3.3 incorporation at ZGA in *Drosophila*.

We identify a more rapid decrease in the nuclear availability of H3 than H3.3 over the final

pre-ZGA cycles. We find that chaperone binding, not gene expression, controls

incorporation patterns using H3/H3.3 chimeric proteins at the endogenous H3.3A locus.

The increase in H3.3 incorporation depends on the N/C ratio. Since the N/C ratio affects

many parameters of embryogenesis, we further test the contributions of genomic content,

zygotic transcription, and cell cycle states. We identify cell cycle regulation, but not H3

availability or transcription, as a major determinant of H3.3 incorporation. Overall, we

propose a model in which local N/C ratios regulate chromatin composition via cell cycle

state during ZGA.

Results

The interphase nuclear availability of H3 decreases more rapidly than H3.3 over the pre-ZGA cycles

To understand in vivo dynamics of the H3/H3.3 pair during ZGA in *Drosophila*, we previously tagged H3 and H3.3 with a photo-convertible Dendra2 protein, (H3-Dendra2 and H3.3-Dendra2) at a pseudo-endogenous H3 locus and the endogenous H3.3A locus respectively (Figures S1A-B)[29]. *Drosophila* ZGA occurs after 13 rapid syncytial nuclear cycles (NCs) and is accompanied by cell cycle slowing and cellularization. We have previously shown that with each NC, the pool of free H3 is depleted and its levels on chromatin decrease (Figure S1C-D)[29]. In contrast, H3.3 chromatin levels increase during the same cycles (Figure S1C-D)[29]. To test if changes in the relative nuclear availability of H3 and H3.3 mirror the observed chromatin incorporation trends, we measured the nuclear intensities of H3-Dendra2 and H3.3-Dendra2 in each interphase. We observed that H3 nuclear intensities decreased by ~40% between NC10 and NC13 as previously shown (Figure 1A-B)[29]. However, when we measured H3.3-Dendra2 nuclear intensities we found that they decreased by only ~20% between NC10 and NC13 (Figure 1A-B). To further assess how nuclear uptake dynamics changed during these cycles, we tracked total nuclear H3 and H3.3 in each cycle and found that H3.3 accumulation reduced more slowly than H3 (Figure 1C-D).

The reduction in nuclear accumulation could be due to a decrease in nuclear import, an increase in nuclear export, or both. To test these possibilities, we quantified the rate of nuclear export by photo-converting Dendra2 during interphase and measuring red

Dendra2 signal over time. Using this method, we have previously shown that nuclear

export of H3 is negligible[29]. Here, we find that export of H3.3 is also negligible (Figure

S1E). These data suggest that the distinct dynamics of H3 and H3.3 nuclear availability

are due to their import dynamics. Though the change in initial import rates between NC10

and NC13 are similar between the two histones (Figure S1F), we observed a notable

difference in their behavior in NC13. H3 nuclear accumulation plateaus ~5 minutes into

NC13, whereas H3.3 nuclear accumulation merely slows (Figure 1C-D). These changes

in nuclear import and incorporation result in a less dramatic loss of the free nuclear H3.3

pool than previously seen for H3 (Figure 1E)[29].

Chaperone binding sites regulate the differences in H3 and H3.3 chromatin

incorporation

We next investigated what differences between H3 and H3.3 caused the observed trends

in chromatin incorporation. There are two major differences between H3 and H3.3: protein

sequence and expression pattern. H3 differs from H3.3 by four amino acids which create

an additional phosphosite in H3.3 and generate differing affinities for specific H3-family

histone chaperones[55]. H3 is also generally expressed at much higher levels and in a

replication-dependent manner. To determine which factor controls nuclear availability and

chromatin incorporation, we genetically engineered flies to express Dendra2-tagged

H3/H3.3 chimeras at the endogenous H3.3A locus. These chimeras include (i) H3.3's

phosphosite replaced with Alanine from H3 ($H3.3^{S31A}$) (ii) H3.3's chaperone binding

domain replaced with H3's ($H3.3^{SVM}$), and (iii) all four H3.3-specific amino acids replaced

with those of H3 ($H3.3^{ASVM}$), (Figure 2A). In all cases, the gene structure, including the

promoter, intron, and UTRs of H3.3, remained intact and no other codons were changed

to maximize similarity to the endogenous H3.3A locus.

To study how chromatin incorporation differed in these chimeras, we measured their total

intensities on mitotic chromatin during each nuclear cycle. We observed that, though

H3.3^{S31A} chromatin incorporation was significantly reduced compared to H3.3 by NC13,

its levels increased on chromatin over the nuclear cycles, resembling H3.3 more than H3

(Figure 2B, S2A). Conversely, the total amount of H3.3^{SVM} and H3.3^{ASVM} on mitotic

chromatin fell over the nuclear cycles, similar to H3 (Figure 2B, S2B-C). This suggests

that chromatin incorporation is mainly determined by the chaperone binding site. These

results are broadly consistent with the final interphase nuclear concentrations and import

dynamics where H3.3^{S31A} was intermediate between H3 and H3.3 while H3.3^{SVM} and

H3.3^{ASVM} were more similar to H3 (Figure 2C, S2A-G). However, both nuclear H3.3^{S31A}

and H3.3^{SVM} fell more quickly than H3.3 and H3 respectively, suggesting that chimeric

histones may not be as stable and/or efficiently imported as their canonical counterparts.

Together, these data indicate that the specific amino acid sequence of the chaperone

binding site is the primary factor in differentiating the two histones for chromatin

incorporation and nuclear import dynamics.

137
H3 chaperone binding site conveys independence from Hira for chromatin

incorporation

Since the chromatin incorporation of the H3/H3.3 chimeras appears to depend on their

chaperone binding sites, we asked whether they still required the canonical H3.3

chaperone, Hira. We used Hira$^{ssm-185b}$ (hereafter Hira^{ssm}) flies which have a point mutation

in the Hira locus[56]. This mutant Hira protein can bind but not incorporate H3.3 into

chromatin (Figure 2D-E, S2H), resulting in sperm chromatin decondensation defects.

These embryos develop as haploids and undergo one additional syncytial division before

ZGA (NC14)[56]. The fall in nuclear concentration of H3 is slightly more gradual in the

haploid Hira^{ssm} embryos than in wildtype, though H3 chromatin incorporation is not

disrupted (Figure 1B, 2F-G, S2I). To test if H3-like chimeras expressed from the H3.3A

locus use the canonical Hira pathway, we measured import and chromatin incorporation

of H3.3^{ASVM} in Hira^{ssm}. We found that H3.3^{ASVM} interphase nuclear concentration was

more stable than H3 or H3.3 in Hira^{ssm} embryos (Figure 2G, S2J). This stability is reflected

in H3.3^{ASVM} chromatin incorporation where it only drops by ~20% between NC10 and

NC14 compared to the observed ~40% drop in H3 (Figure 2H-I). These data indicate that

H3.3^{ASVM} chromatin incorporation is Hira independent, even when expressed from the

H3.3A locus.

Local N/C ratios determine H3 and H3.3 chromatin incorporation

Since the N/C ratio controls many aspects of pre-ZGA development we asked whether

the local N/C ratio determines histone chromatin incorporation within a nuclear cycle. To

test this, we employed mutants in the gene Shackleton (Shkl) whose embryos have non-

uniform nuclear densities across the anterior/posterior axis (Figure 3A-B, Movie 1-2)[57]. In

these embryos, impaired cortical migration of early nuclei increases the N/C ratio in the

center and decreases it in the posterior, which results in frequent partial extra divisions at

the posterior pole (Figure 3B, G)[57]. For our analyses, we manually defined low and high

nuclear density regions, with the low-density region always undergoing an extra division

(Figure 3B, see methods). To control for potential positional effects, we measured

chromatin incorporation at the middle and pole regions of control embryos for comparison

(Figure 3A). In control embryos, the drop in the total amount of H3 and rise in total H3.3

on chromatin are comparable between the middle and pole over the pre-ZGA cycles

(Figure 3C-D). In contrast, in Shkl embryos, we observe decreased incorporation of H3

on chromatin at high nuclear densities compared to low nuclear densities (Figure 3E,

S3A). This trend is reversed for H3.3, where chromatin from high density regions has

more total H3.3 than chromatin from low density regions (Figure 3F, S3B).

This observation indicates that incorporation of H3 and H3.3 are reciprocal and depend

on the local N/C ratio leading to several possible models (Figure 3H). First, the H3 pool

available for chromatin incorporation may become limiting at high N/C ratios leading to

increased H3.3 incorporation. Second, since H3.3 is known to be associated with sites of

active transcription[50,52,58–62], the increased H3.3 incorporation might be downstream of

N/C ratio dependent ZGA. Finally, since H3 is usually incorporated only during S-phase

the changing H3 to H3.3 incorporation rates may be the result of N/C ratio-dependent cell

cycle changes. Note, that all these processes feedback onto one another such as cell

cycle slowing allowing time for ZGA[37,63].

H3 nuclear availability depends on the local N/C ratio

To ask whether nuclear availability can explain the N/C ratio-dependent differences in H3

and H3.3 incorporation, we measured their interphase accumulation in Shkl embryos

(Figure 4A). Since H3 and H3.3 both have negligible nuclear export, their nuclear

availabilities are determined by their import rates (Figure 1B, S1E)[29]. To assess the

impact of the N/C ratio on nuclear import in individual nuclei, we calculated the number

of neighbors within a 20 µm radius for each nucleus at its minimum volume (Figure 3G,

Figure S3C). We then binned the nuclei by their number of neighbors and determined

their nuclear import curves for both H3 and H3.3. In control NC13 embryos, there is little

variation in the number of neighbors and all nuclei import H3 and H3.3 similarly (Figure

4B-E). In NC13 Shkl embryos, H3 import is anticorrelated with the local N/C ratio (Figure

4F, H, S3D). We observed slower H3 nuclear uptake at high N/C ratios resulting in lower

total interphase H3 accumulation (Figure 4F). This was also reflected in the initial H3

import rates where the nuclei with fewer neighbors had higher slopes (Figure 4H). H3.3

uptake was less affected by the local N/C ratio (Figure 4G, I, S3E). A similar trend was

also observed in NC12 for both histones, where more neighbors correspond to slower

import. However, the range of behaviors was not as large as seen in NC13 (Figure S3F-

G). These observations support a model where H3 pools are exhausted by the increasing

N/C ratio, increasing the relative availability of H3.3 to H3 over the pre-ZGA cycles.

H3.3 incorporation is not caused by exhaustion of H3 pools

Given that the available H3 seems to be depleted by the increasing N/C ratio we sought

to test if H3.3 chromatin incorporation depends on the size of the H3 pool (Figure 5A).

We hypothesized that as the embryo exhausted the supply of RC H3 it might increase the

use of RI H3.3 to compensate. We knocked down Stem-loop binding protein (Slbp), which

specifically binds and stabilizes the mRNAs of RC histones, including H3, but does not

interact with H3.3 mRNAs[7,64,65]. Slbp RNAi dramatically decreases the size of the

available H3 pool and results in frequent chromosomal segregation defects (Figure

S4A)[39]. For this reason, we only analyzed embryos that appeared reasonably healthy

until the final cell cycle under consideration. In embryos that survives through at least

NC12, we found that H3.3 incorporation is largely unaffected by the reduction in RC H3

(Figure 5B). To further validate that the lack of effect on H3.3 incorporation was not due

to inefficient Slbp-knockdown, we also tested H3.3 incorporation in embryos that already

display severe bridging in NC11. In these embryos, we detected no difference in the H3.3

incorporation in NC10 mitosis (Figure S4B). These results strongly indicate that simply

running out of H3 is not the cause of the observed increase in H3.3 on chromatin.

H3.3 incorporation does not depend on zelda-dependent ZGA

Since H3.3 is associated with sites of active transcription in other systems[50,52,58–62], we

next sought to test if H3.3 incorporation during ZGA depends on transcription. To do this,

we knocked down the critical pioneer transcription factor zelda (Figure 5C). zelda controls

the transcription of the majority of Pol II genes during ZGA and when zelda is disrupted,

Pol II relocates to the histone locus body[66–68]. We found that H3.3 chromatin incorporation

did not change in zelda RNAi embryos despite their inability to cellularize and longer

NC13s (Figure 5D and S4C). This suggests that the large increase in H3.3 incorporation

that we detect by microscopy in the final nuclear cycles does not depend on bulk ZGA.

H3.3 incorporation depends on cell cycle state, but not cell cycle duration

Finally, to test the contribution of the cell cycle on the N/C ratio dependent accumulation

of H3.3 on chromatin we used mutants in Chk1 (grapes in *Drosophila*) that are less

efficient in cell cycle slowing (Figure 5E, S4C). These mutants have an unusually rapid

NC13 and attempt to enter mitosis before their DNA is fully replicated resulting in mitotic

catastrophe[69]. We found that H3.3 accumulation is disrupted as early as NC12 (P-

value=10^{-8}) in Chk1 mutants (Figure 5F). Importantly, the Chk1 mutants have relatively

normal NC12 durations[69,70]. In our experiments, Chk1 NC12 was only ~1 minute faster

than wildtype and Chk1 embryos with comparable cell cycle durations still displayed

reduced H3.3 incorporation (Figure 5F and S4C). To further isolate the effect of cell cycle

length on H3.3 incorporation we used the natural variation in NC13 duration in control

embryos. When we plotted H3.3 chromatin signal against the total NC13 duration for

control embryos we found no correlation (Figure S4D). This result suggests that cell cycle

duration as such does not directly regulate H3.3 chromatin incorporation. Instead, Chk1

appears to regulate H3.3 incorporation in a manner that is not mediated solely by

lengthening the cell cycle.

Discussion

We demonstrate that H3.3 replaces H3 on chromatin leading up to ZGA in *Drosophila*.

This process depends on the specific H3.3 chaperone binding site and is controlled by

the N/C ratio. We tested which aspects of the N/C ratio control the dynamic incorporation

of H3.3 and found that cell cycle state, but not H3 availability or bulk transcription, is the

major regulator of H3.3 behavior. Chk1 mutants decrease H3.3 incorporation even before

the cell cycle is significantly slowed. Cell cycle slowing has been previously reported to

regulate the incorporation of other histone variants in *Drosophila*[15]. However, our results

indicate that cell cycle state and not duration per se, regulates H3.3 incorporation. We

speculate that this may be due to changes in chromatin state as a result of Chk1 activity.

Late replicating regions and heterochromatin first emerge during ZGA, and Chk1 can

control origin firing in many contexts[21,23,24,71,72]. Since H3.3 is often associated with

heterochromatin, the decreased H3.3 incorporation in Chk1 mutants may be an indirect

result of increased origin firing and decreased heterochromatin formation[71,72]. Another

possibility is that the additional Chk1 phosphosite that is found in H3.3-S31 may be

important for promoting H3.3 incorporation during ZGA[50].

The interaction between H3-type histones and Chk1 has additional significance since H3

nuclear concentration has been proposed to directly regulate cell cycle length through H3

interactions with Chk1[40]. In Hira^{ssm} embryos that undergo one extra division before cell

cycle slowing, the fall in nuclear H3 concentration between NC10 and the final fast cell

cycle is strikingly similar to that seen in wildtype. Moreover, H3 nuclear concentrations

appear to be strongly sensitive to the local N/C ratio in Shkl embryos. Together these

data are consistent with a model in which H3 nuclear concentrations regulate cell cycle

slowing. However, H3.3 nuclear concentrations are less sensitive to the local N/C ratio

than H3. Since H3.3 has an additional Chk1 phosphorylation site compared to H3 it may

have different regulatory interactions with Chk1[50,73]. The relative contributions of both H3

and H3.3 nuclear availability to cell cycle slowing will require further exploration.

Finally, how the changing histone landscape contributes to ZGA remains an important

open question. We have shown that bulk H3.3 incorporation does not depend on

transcription from zelda-dependent genes. However, the reciprocal relationship remains

untested. H3.3 incorporation may increase transcription factor accessibility at specific

genomic loci to mark them for activation. It is also possible that H3.3 incorporation occurs

as a response to transcription initiated by other transcription factors but does not

specifically respond to the pioneer factor zelda. We have shown that disruption of major

ZGA does not impair bulk H3.3 incorporation, but the role of H3.3 containing nucleosomes

in ZGA remains to be tested.

Materials and methods

***Drosophila* stocks and genetic crosses**

A) y,w; 1xHisC.H3-Dendra2;

B) y,w; H3.3A-Dendra2/CyO;

C) y,w; IX HisC.H3-Dendra2; *shkl*^{GM163}/TM3

D) y,w; IX HisC.H3-Dendra2; *shkl*^{GM130}/TM3

E) y,w; H3.3A-Dendra2/CyO; *shkl*^{GM163}/TM6B

F)   y,w; H3.3A-Dendra2/CyO; *shkl*^{GM130}/TM6B

G)   y,w; H3.3A-Dendra2^{S31A}/CyO;

H)   y,w; H3.3A-Dendra2^{SVM}/CyO;

I)   y,w; H3.3A-Dendra2^{ASVM}/CyO;

J) *ssm*^{185b},w/FM7c,*w*^{a}; IX HisC.H3-Dendra2;

K) *ssm*^{185b},w/FM7c,*w*^{a}; H3.3A-Dendra2/CyO;

L) *ssm*^{185b},w/FM7c,*w*^{a}; H3.3A-Dendra2^{S31A}/CyO;

M) *ssm*^{185b},w/FM7c,*w*^{a}; H3.3A-Dendra2^{SVM}/CyO;

N) *ssm*^{185b},w/FM7c,*w*^{a}; H3.3A-Dendra2^{ASVM}/CyO;

O) y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ21114}attP40 (Slbp-RNAi)

P) ;;UAS-Zld-shRNA

Q) yw; Mat-α-tub67-gal4, H3.3A-Dendra2 / CyO; Mat-α-tub15

R) y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00094}attP2 (white-RNAi)

S) y,w; H3.3A-Dendra2, grp^{1};

Fly stocks A, B, J, and K were generated previously in our lab and are described in Shindo

and Amodeo (2019). The Shkl lines y,w; Sp/CyO; *shkl*^{GM163}/TM3 and y,w; Sp/CyO;

shkl[GM130]/TM3 were a generous gift from Stefano Di Talia, Duke University. Stocks C, D, E, and F were generated in the lab by the genetic crossing of these Shkl lines with lines A and B. Stocks L, M, and N were generated by crossing the stocks G, H and I with *ssm*[185b],w/FM7c,*w*[a];; (a generous gift from Eric Weischaus). Stocks O and R were obtained from the Bloomington *Drosophila* Stock Center with IDs #51171 and #35573[74]. Stock P was a generous gift from Christine Rushlow, NYU. Stock Q was generated by performing recombination crosses of B with y,w; Mat-α-tub67-gal4; Mat-α-tub15 (a generous gift from Eric Weischaus)[75]. Stock S was generated by performing recombination crosses of B with y,w; grp1/CyO; flies (a generous gift from Eric Weischaus)[70].

Drosophila husbandry

All fly stocks were maintained at room temperature, on standard molasses media. The egg lay cages were set up to collect embryos at 25°C (except for the Slbp RNAi flies). Slbp RNAi egg lay cages and associated control w-RNAi cages were set up at 18°C. Embryos from these cages were collected on apple juice agar plates with yeast paste, dechorionated with 50% bleach for up to 2 minutes, and washed twice with dH$_2$O. The *ssm*[185b] embryos were collected from *ssm*[185b]/*ssm*[185b] homozygous females. For shkl embryos, the 2 shkl lines were crossed to obtain *shkl*[GM130e]/*shkl*[GM163e] transheterozygous females and their embryos were imaged. For all the RNAi crosses, males from the gal4 driver line Q, were crossed with virgins from UAS-RNAi lines (O, P or R) to obtain progeny expressing both UAS and Gal4. Embryos from these progeny flies were used for imaging.

333 Embryos from w-RNAi flies were used as controls for all RNAi experiments. Chk1[-/-]

334 embryos were collected from grp[1]/grp[1] homozygous females.

Plasmids and transgenesis

To generate stocks G, H, and I, CRISPR-Cas9 editing was performed at the endogenous

H3.3A locus. To this end, pScarlessHD-H3.3A-Dendra2-DsRed plasmid, reported in

Shindo and Amodeo (2019)[29] was modified through site-directed mutagenesis to express

H3.3 with H3-specific amino acids, generating pScarlessHD-H3.3A[S31A]-Dendra2-DsRed

(S31A mutation), pScarlessHD-H3.3A[SVM]-Dendra2-DsRed (A87S, I89V, G90M

mutations) and pScarlessHD-H3.3A[ASVM]-Dendra2-DsRed (S31A, A87S, I89V, G90M

mutations) plasmids (Genscript). Two CRISPR target sites were identified using Target

Finder[76], one near the stop codon and one near S31, and the corresponding gRNAs were

cloned into pU6-BbsI-chiRNA vector (a gift from Melissa Harrison & Kate O'Connor-Giles

& Jill Wildonger, Addgene plasmid #45946). Each mutant plasmid was co-injected with

both the gRNA plasmids into nos-Cas9 embryos (TH00787.N) and DsRed+ progeny were

selected (BestGene). These progeny were then crossed with nos-PBac flies (a generous

gift from Robert Marmion and Stas Shvartsman) to remove the DsRed marker. DsRed

negative single males were then crossed with y,w;Sp/CyO; to establish stocks G, H, and

I. Insertion of Dendra2 tagged mutants was verified by PCR and Sanger sequencing.

Microscopy

For live imaging, dechorionated embryos were mounted on glass-bottom MatTek dishes

in deionized water and imaged with the 20x, 0.8 NA, objective of Zeiss LSM980 confocal

microscope with Airyscan-2 at 45 s intervals for 2 h at room temperature (19-22°C). All

H3-Dendra2 tagged embryos were imaged using a 488 nm laser at 2% power and all

lines expressing Dendra2 tagged proteins from the endogenous H3.3A locus (H3.3 and

the chimeras) were imaged with a 488 nm with 0.5% power in Airyscan multiplex CO-8Y

mode. All but shkl embryos and their controls were imaged at a 700 x 700 pixels

resolution, with 1 μm Z-steps over a 15 μm range, with a frame time of 26.06ms. All Shkl

embryos and their controls (Figure 3, 4, S3) were imaged at a 2836 x 2836 pixels

resolution, with 1.2 μm Z-steps over a 14.4 μm range, with a frame time of 328.29ms. All

images were acquired with a pixel size of 0.149 μm x 0.149 μm.

Nuclear export and unbound H3.3 measurement through Dendra2 photoconversion

For measuring the nuclear export and amount of free histone H3.3 (Figure 1E and S1E),

we used the photoconvertible Dendra2 tag and the interactive bleaching panel in Zen

software. We used a 4μm diameter circular stencil to interactively photo-convert the

nuclei. H3.3-Dendra2 within a single nucleus was photoconverted from green to red using

a 405 nm laser at 3% power with 60 iterations of laser exposure at a speed of 1.37μs/pixel.

The nucleus was converted in the middle of each nuclear cycle for NC11-13 and then

imaged with 561 nm at 1% laser power and 488 nm with 0.5% laser power at 15-second

intervals until the end of the nuclear cycle. Images were captured at 576 x 576 pixels

resolution with 1 μm Z-steps over a 15 μm range, with a frame time of 66.55ms for each

channel. The images were acquired with a 40x oil immersion objective, 1.3 NA with a

pixel size of 0.092 μm x 0.092 μm.

Photobleaching corrections

To assess the potential effects of fluorophore photobleaching during our image capture, we performed parallel embryo experiments. In these experiments, we identified 2 embryos of the same age and imaged the interphase nucleus and the metaphase chromatin for both in NC10. Following this, we image only a sub-region of one of the two embryos continuously with our experimental settings described for H3-Dendra2 embryos above until NC13, while keeping the other embryo to develop parallelly without imaging. Once the imaged embryo reached NC13, both the imaged and unimaged parallel embryo were imaged again. We quantified the total nuclear signal from both embryos to evaluate the photobleaching effects. We then compared the continuously imaged section of the embryo, with the area outside the sub-region imaged as well as the unimaged parallel embryo. Using these comparisons, we determined that the effect of photobleaching was minimal and therefore did not apply a numeric photobleaching correction to our data (Figure S1G-H, Table S4-5).

Nuclear segmentation and intensity analysis

All raw CZI output files from ZEN 3.3 (blue edition) live imaging were first 3D Airyscan Processed at a strength of 3.7 and then converted into individual TIFF files.

For mitotic chromatin quantification, the time points corresponding to metaphase chromatin from each nuclear cycle were extracted and the z-stacks were sum projected in FIJI (2.14.0/1.54f) These files were segmented using the 'pixel classification + object classification' applet in the ilastik-1.4.0 software[77] into chromatin and cytoplasm. The

individually segmented mitotic chromatin objects were then exported as a single CSV file

containing object properties such as total intensity, mean intensity, and size. The total

intensity within each chromatin mass was calculated and normalized to the average NC10

chromatin values (or NC11 for shkl embryos and their controls) for that genotype.

In Shkl embryos and their controls (Figure 3C-F, S3A-B), chromatin was segmented from

different regions within an embryo (middle and pole regions for control, and from low and

high-density regions for Shkl). In control embryos, middle regions were defined by

outlining a box (250x250 pixels) in NC10 around the line separating the embryo into 2

halves. A similar-sized box was outlined with one edge at the tip of the embryo to define

the pole region. In shkl embryos, the regions with the highest apparent nuclear density

within the center was defined as the high-density region and the region that underwent

the partial extra division in NC14 was defined as the low-density region. To account for

the asynchronous nature of the divisions in the shkl embryo, within each region, 5-6 nuclei

that divided synchronously along the mitotic wave were quantified. For both control and

shkl embryos, at least 5 nuclei per embryo were quantified in each cycle for each region.

For analyzing the interphase nuclear concentrations, nuclei from 45 seconds before the

nuclear envelope breakdown were segmented in 3D using the 'pixel classification + object

classification' applet on ilastik software. The CSV file with the mean intensities of each

nucleus was exported and normalized to the average NC10 nuclear concentration values

for each genotype.

For obtaining the nuclear import curves, individual nuclear cycles were run through the

pixel classification + object classification applet in the ilastik software. The results were

exported as CSV files and processed with a custom R script. For Shkl embryos (Figure

4B-C,F-G), the pixel prediction maps were used with the 'tracking with learning' applet

(ilastik) to segment the nuclei as well as track them over time. The tracking result with

object properties was exported as a CSV file and processed with a custom R script.

Intensities were normalized by the average total intensity of the nuclei at their maximum

size in each cycle. For each case, the volume was calculated by multiplying the voxel size

with the 'size in pixels' of an object.

Neighborhood analysis

Nuclei within each embryo were tracked over a single nuclear cycle using the 'tracking +

learning' applet on ilastik. The tracking result with the coordinates of each nucleus over

time was obtained as a CSV file, along with other parameters including the total intensity,

mean intensity, and nuclear size. The CSV file was analyzed to calculate the number of

nuclear neighbors for each nucleus within a 20 μm radius using a custom R script. The

script calculates the number of neighbors each nucleus has at its minimum volume since

the maximum nuclear import occurs at this time point. To overcome the noise from the

incomplete edge nuclei, which are centered lower in the embryo, we utilized the

differences in their Z-coordinates to filter them out, after using them for the number of

neighbor calculations. For Shkl embryos, as the nuclear cycles are asynchronous, the

total intensity traces were aligned to match their minimum volumes to T0. Nuclei with the

same number of neighbors were binned together and weighted to reflect the number of

nuclei being averaged. The total intensity curves were then normalized such that the

average total intensity of the nuclei at their maximum size was equal to 1.

Cell cycle time measurements

Cell cycle durations were measured from metaphase to metaphase. To account for day-

to-day temperature variability, we normalized the mean NC11 durations in control

embryos to 10 minutes and scaled for other cell cycles in all embryos acquired on the

same day accordingly as done previously[70].

Western blot analysis

For western blotting, embryos were collected from white-RNAi flies or Slbp-RNA flies, for

a period of 1 hr. Following this the embryos were dechorionated with 50% bleach for 2

minutes followed by 2 washes with deionized water. They were then collected in a

microcentrifuge tube and lysed with forceps in ice-cold embryo lysis buffer (50 mM Tris

pH 8.0, 150 mM NaCl, 0.5% Triton-X, 1 mM $MgCl_2$, 0.1 mM EDTA, 1X protease inhibitor

cocktail (Sigma: P2714)). 25 embryos were collected per genotype to quantify pan-H3

levels. Lamelli buffer was added in 1:1 volume and the samples were boiled at 95°C for

5 minutes. The protein lysates were run on a TGX 12% acrylamide gel (Bio-Rad

Laboratories), stain-free activated for 45 secs under UV, and transferred onto a LF-PVDF

membrane. Membranes were incubated in rabbit anti-H3 antibody (1:1000, Abcam:

ab1791) overnight at 4°C. They were then washed and incubated 2 hrs in Alexa Fluor

647-conjugated donkey anti-rabbit IgG antibody (1:2000, Invitrogen: A31573). The

membranes were then imaged to detect for fluorescence using a gel imager (Bio-Rad

ChemiDoc MP).

Statistical analysis

Two-way ANOVA tests were conducted to assess the statistical significance between the

dataset means of different genotypes over the nuclear cycles. All studies were performed

with nuclei from at least 3 embryos. For Shkl embryos, a two-way ANOVA test was used

to determine the statistical significance of nuclei within different regions of the same

embryo over the different nuclear cycles, with each nucleus as a replicate. For all other

embryos, the average chromatin/nuclear values for each NC from each embryo were

considered as a replicate. Results from these tests are reported in supplementary tables

S1-9.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

We thank Shruthi Balachandra, Eric Alpert, Grace Carey and Kiera Schwarz for

constructive discussion of this manuscript. We further thank the members of Bickel,

Kasper, Lacefield, Moseley and Landino labs at Dartmouth for their helpful suggestions.

We thank Patrick Robison from the Dartmouth bioMT Core, Ann Lavanway and Britton

Johnson for technical support. We thank Stefano Di Talia for shkl stock flies, Eric

Weischaus for ssm, grp and gal4-driver stock flies; Robert Marmion for the nos-PBac

transposase flies, Christine Rushlow for the zelda-RNAi flies and Robert Duronio for the

1xHisC plasmid. We thank FlyBase, funded by NHGRI and NIGMS, and Bloomington

Drosophila Stock Center (NIHP40OD018537) for providing essential resources. This

work was funded by NIH/NIGMS (P20-GM113132 and R35GM150853 to AAA). ADB is

supported by Sondra and Charles Gilman Graduate Research Fellowship.

Author contributions

Conceptualization, ADB, MGB, YS and AAA.; Investigation and Analysis, ADB, MGB and

ABW; Writing – Original Draft, ADB, MGB and AAA; Writing – Review & Editing, ADB,

MGB, YS and AAA.

References

1. Khorasanizadeh, S. (2004). The Nucleosome: From Genomic Organization to Genomic
Regulation. Cell *116*, 259–272. https://doi.org/10.1016/S0092-8674(04)00044-3.

2. Talbert, P.B., and Henikoff, S. (2017). Histone variants on the move: substrates for
chromatin   dynamics.   Nat   Rev   Mol   Cell   Biol   *18*,   115–126.
https://doi.org/10.1038/nrm.2016.148.

3. Talbert, P.B., and Henikoff, S. (2021). Histone variants at a glance. Journal of Cell
Science *134*. https://doi.org/10.1242/jcs.244749.

4. Weber, C.M., and Henikoff, S. (2014). Histone variants: dynamic punctuation in
transcription. Genes Dev. *28*, 672–682. https://doi.org/10.1101/gad.238873.114.

5. Loppin, B., and Berger, F. (2020). Histone Variants: The Nexus of Developmental
Decisions and Epigenetic Memory. Annual Review of Genetics *54*, 121–149.
https://doi.org/10.1146/annurev-genet-022620-100039.

6. Lifton, R.P., Goldberg, M.L., Karp, R.W., and Hogness, D.S. (1978). The Organization
of the Histone Genes in Drosophila melanogaster: Functional and Evolutionary
Implications.   Cold   Spring   Harb   Symp   Quant   Biol   *42*,   1047–1051.
https://doi.org/10.1101/SQB.1978.042.01.105.

7. Marzluff, W.F., Wagner, E.J., and Duronio, R.J. (2008). Metabolism and regulation of
canonical histone mRNAs: life without a poly(A) tail. Nat Rev Genet *9*, 843–854.
https://doi.org/10.1038/nrg2438.

8. McKay, D.J., Klusza, S., Penke, T.J.R., Meers, M.P., Curry, K.P., McDaniel, S.L., Malek,
P.Y., Cooper, S.W., Tatomer, D.C., Lieb, J.D., et al. (2015). Interrogating the Function
of Metazoan Histones using Engineered Gene Clusters. Developmental Cell *32*, 373–
386. https://doi.org/10.1016/j.devcel.2014.12.025.

9. Dominski, Z., and Tong, L. (2021). U7 deciphered: the mechanism that forms the
unusual 3' end of metazoan replication-dependent histone mRNAs. Biochem Soc
Trans *49*, 2229–2240. https://doi.org/10.1042/BST20210323.

10.   Talbert, P.B., and Henikoff, S. (2010). Histone variants — ancient wrap artists of
the epigenome. Nat Rev Mol Cell Biol *11*, 264–275. https://doi.org/10.1038/nrm2861.

11.   Shindo, Y., Brown, M.G., and Amodeo, A.A. (2022). Versatile roles for histones in
early development. Current Opinion in Cell Biology *75*, 102069.
https://doi.org/10.1016/j.ceb.2022.02.003.

12.   Dimitrov, S., Almouzni, G., Dasso, M., and Wolffe, A.P. (1993). Chromatin
Transitions during Early *Xenopus* Embryogenesis: Changes in Histone H4 Acetylation
and in Linker Histone Type. Developmental Biology *160*, 214–227.
https://doi.org/10.1006/dbio.1993.1299.

13.   Pérez-Montero, S., Carbonell, A., Morán, T., Vaquero, A., and Azorín, F. (2013).
The Embryonic Linker Histone H1 Variant of *Drosophila*, dBigH1, Regulates Zygotic
Genome Activation. Developmental Cell *26*, 578–590.
https://doi.org/10.1016/j.devcel.2013.08.011.

14.   Müller, K., Thisse, C., Thisse, B., and Raz, E. (2002). Expression of a linker
histone-like gene in the primordial germ cells in zebrafish. Mechanisms of Development
117, 253–257. https://doi.org/10.1016/S0925-4773(02)00174-0.

15.   Johnson, M.R., Stephenson, R.A., Ghaemmaghami, S., and Welte, M.A.
Developmentally regulated H2Av buffering via dynamic sequestration to lipid droplets
in Drosophila embryos. eLife *7*, e36021. https://doi.org/10.7554/eLife.36021.

16.   Smith, R.C., Dworkin-Rastl, E., and Dworkin, M.B. (1988). Expression of a histone
H1-like protein is restricted to early Xenopus development. Genes Dev *2*, 1284–1295.
https://doi.org/10.1101/gad.2.10.1284.

17.   Wibrand, K., and Olsen, L.C. (2002). Linker histone *H1M* transcripts mark the
developing germ line in zebrafish. Mechanisms of Development *117*, 249–252.
https://doi.org/10.1016/S0925-4773(02)00173-9.

18.   Tadros, W., and Lipshitz, H.D. (2009). The maternal-to-zygotic transition: a play in
two acts. Development *136*, 3033–3042. https://doi.org/10.1242/dev.033183.

19.   Vastenhouw, N.L., Cao, W.X., and Lipshitz, H.D. (2019). The maternal-to-zygotic
transition revisited. Development *146*, dev161471. https://doi.org/10.1242/dev.161471.

20.   Blythe, S.A., and Wieschaus, E.F. (2016). Establishment and maintenance of
heritable chromatin structure during early Drosophila embryogenesis. eLife *5*, e20148.
https://doi.org/10.7554/eLife.20148.

21.   Seller, C.A., Cho, C.-Y., and O'Farrell, P.H. (2019). Rapid embryonic cell cycles
defer the establishment of heterochromatin by Eggless/SetDB1 in Drosophila. Genes
Dev. https://doi.org/10.1101/gad.321646.118.

22.   Hug, C.B., Grimaldi, A.G., Kruse, K., and Vaquerizas, J.M. (2017). Chromatin
Architecture Emerges during Zygotic Genome Activation Independent of Transcription.
Cell *169*, 216-228.e19. https://doi.org/10.1016/j.cell.2017.03.024.

23.   Shermoen, A.W., McCleland, M.L., and O'Farrell, P.H. (2010). Developmental
Control of Late Replication and S phase Length. Curr Biol *20*, 2067–2077.
https://doi.org/10.1016/j.cub.2010.10.021.

24.   McKnight, S.L., and Miller, O.L. (1977). Electron microscopic analysis of chromatin
replication in the cellular blastoderm drosophila melanogaster embryo. Cell *12*, 795–
804. https://doi.org/10.1016/0092-8674(77)90278-1.

25.   Zhang, Y., Vastenhouw, N.L., Feng, J., Fu, K., Wang, C., Ge, Y., Pauli, A., van
Hummelen, P., Schier, A.F., and Liu, X.S. (2014). Canonical nucleosome organization
at promoters forms during genome activation. Genome Res *24*, 260–266.
https://doi.org/10.1101/gr.157750.113.

26.   Horard, B., and Loppin, B. (2015). Histone storage and deposition in the early
Drosophila embryo. Chromosoma *124*, 163–175. https://doi.org/10.1007/s00412-014-
0504-7.

27.   Adamson, E.D., and Woodland, H.R. (1974). Histone synthesis in early amphibian
development: Histone and DNA syntheses are not co-ordinated. Journal of Molecular
Biology *88*, 263–285. https://doi.org/10.1016/0022-2836(74)90481-1.

28.   Woodland, H.R., and Adamson, E.D. (1977). The synthesis and storage of
histones during the oogenesis of Xenopus laevis. Dev Biol *57*, 118–135.
https://doi.org/10.1016/0012-1606(77)90359-1.

29.   Shindo, Y., and Amodeo, A.A. (2019). Dynamics of Free and Chromatin-Bound
Histone H3 during Early Embryogenesis. Current Biology *29*, 359-366.e4.
https://doi.org/10.1016/j.cub.2018.12.020.

30.   Farrell, J.A., and O'Farrell, P.H. (2014). From Egg to Gastrula: How the Cell Cycle
Is Remodeled During the Drosophila Mid-Blastula Transition. Annual Review of
Genetics *48*, 269–294. https://doi.org/10.1146/annurev-genet-111212-133531.

31.   Newport, J., and Kirschner, M. (1982). A major developmental transition in early
Xenopus embryos: I. characterization and timing of cellular changes at the midblastula
stage. Cell *30*, 675–686. https://doi.org/10.1016/0092-8674(82)90272-0.

32.   Newport, J., and Kirschner, M. (1982). A major developmental transition in early
Xenopus embryos: II. Control of the onset of transcription. Cell *30*, 687–696.
https://doi.org/10.1016/0092-8674(82)90273-2.

33.   Kane, D.A., and Kimmel, C.B. (1993). The zebrafish midblastula transition.
Development *119*, 447–456. https://doi.org/10.1242/dev.119.2.447.

34.   Blythe, S.A., and Wieschaus, E.F. (2015). Chapter Four - Coordinating Cell Cycle
Remodeling with Transcriptional Activation at the Drosophila MBT. In Current Topics in
Developmental Biology The Maternal-to-Zygotic Transition., H. D. Lipshitz, ed.
(Academic Press), pp. 113–148. https://doi.org/10.1016/bs.ctdb.2015.06.002.

35.   Edgar, B.A., Kiehle, C.P., and Schubiger, G. (1986). Cell cycle control by the
nucleo-cytoplasmic ratio in early Drosophila development. Cell *44*, 365–372.
https://doi.org/10.1016/0092-8674(86)90771-3.

36.   Edgar, B.A., and Schubiger, G. (1986). Parameters controlling transcriptional
activation during early drosophila development. Cell *44*, 871–877.
https://doi.org/10.1016/0092-8674(86)90009-7.

37.   Syed, S., Wilky, H., Raimundo, J., Lim, B., and Amodeo, A.A. (2021). The nuclear
to cytoplasmic ratio directly regulates zygotic transcription in *Drosophila* through
multiple modalities. Proceedings of the National Academy of Sciences *118*,
e2010210118. https://doi.org/10.1073/pnas.2010210118.

38.   Amodeo, A.A., Jukam, D., Straight, A.F., and Skotheim, J.M. (2015). Histone
titration against the genome sets the DNA-to-cytoplasm threshold for the Xenopus
midblastula transition. Proceedings of the National Academy of Sciences *112*, E1086–
E1095. https://doi.org/10.1073/pnas.1413990112.

39.   Chari, S., Wilky, H., Govindan, J., and Amodeo, A.A. (2019). Histone concentration
regulates the cell cycle and transcription in early development. Development *146*,
dev177402. https://doi.org/10.1242/dev.177402.

40.   Shindo, Y., and Amodeo, A.A. (2021). Excess histone H3 is a competitive Chk1
inhibitor that controls cell-cycle remodeling in the early Drosophila embryo. Current
Biology *31*, 2633-2642.e6. https://doi.org/10.1016/j.cub.2021.03.035.

41.   Almouzni, G., Méchali, M., and Wolffe, A.P. (1990). Competition between
transcription complex assembly and chromatin assembly on replicating DNA. The
EMBO Journal *9*, 573–582. https://doi.org/10.1002/j.1460-2075.1990.tb08145.x.

42.   Almouzni, G., Méchali, M., and Wolffe, A.P. (1991). Transcription Complex
Disruption Caused by a Transition in Chromatin Structure. Molecular and Cellular
Biology *11*, 655–665. https://doi.org/10.1128/mcb.11.2.655-665.1991.

43.   Prioleau, M.N., Huet, J., Sentenac, A., and Méchali, M. (1994). Competition
between chromatin and transcription complex assembly regulates gene expression
during early development. Cell *77*, 439–449. https://doi.org/10.1016/0092-
8674(94)90158-9.

44.   Almouzni, G., and Wolffe, A.P. (1995). Constraints on transcriptional activator
function contribute to transcriptional quiescence during early Xenopus embryogenesis.
EMBO J *14*, 1752–1765.

45.   Joseph, S.R., Pálfy, M., Hilbert, L., Kumar, M., Karschau, J., Zaburdaev, V.,
Shevchenko, A., and Vastenhouw, N.L. (2017). Competition between histone and
transcription factor binding regulates the onset of transcription in zebrafish embryos.
eLife *6*, e23326. https://doi.org/10.7554/eLife.23326.

46.   Li, Z., Johnson, M.R., Ke, Z., Chen, L., and Welte, M.A. (2014). *Drosophila* Lipid
Droplets Buffer the H2Av Supply to Protect Early Embryonic Development. Current
Biology *24*, 1485–1491. https://doi.org/10.1016/j.cub.2014.05.022.

47.   Santenard, A., Ziegler-Birling, C., Koch, M., Tora, L., Bannister, A.J., and Torres-
Padilla, M.-E. (2010). Heterochromatin formation in the mouse embryo requires critical
residues of the histone variant H3.3. Nat Cell Biol *12*, 853–862.
https://doi.org/10.1038/ncb2089.

48.   Jang, C.-W., Shibata, Y., Starmer, J., Yee, D., and Magnuson, T. (2015). Histone
H3.3 maintains genome integrity during mammalian development. Genes Dev *29*,
1377–1392. https://doi.org/10.1101/gad.264150.115.

49.   Klein, R.H., and Knoepfler, P.S. (2023). Knockout tales: the versatile roles of
histone H3.3 in development and disease. Epigenetics & Chromatin *16*, 38.
https://doi.org/10.1186/s13072-023-00512-8.

50.   Sitbon, D., Boyarchuk, E., Dingli, F., Loew, D., and Almouzni, G. (2020). Histone
variant H3.3 residue S31 is essential for Xenopus gastrulation regardless of the
deposition pathway. Nat Commun *11*, 1256. https://doi.org/10.1038/s41467-020-
15084-4.

51.   Delaney, K., Weiss, N., and Almouzni, G. (2023). The cell-cycle choreography of
H3 variants shapes the genome. Molecular Cell *83*, 3773–3786.
https://doi.org/10.1016/j.molcel.2023.08.030.

52.   Sakai, A., Schwartz, B.E., Goldstein, S., and Ahmad, K. (2009). Transcriptional and
developmental functions of the H3. 3 histone variant in Drosophila. Current Biology *19*,
1816–1820.

53.   Hödl, M., and Basler, K. (2009). Transcription in the absence of histone H3.3. Curr
Biol *19*, 1221–1226. https://doi.org/10.1016/j.cub.2009.05.048.

54.   Szenker, E., Ray-Gallet, D., and Almouzni, G. (2011). The double face of the
histone variant H3. 3. Cell research *21*, 421–434.

55.   Tagami, H., Ray-Gallet, D., Almouzni, G., and Nakatani, Y. (2004). Histone H3. 1
and H3. 3 complexes mediate nucleosome assembly pathways dependent or
independent of DNA synthesis. Cell *116*, 51–61.

56.   Loppin, B., Docquier, M., Bonneton, F., and Couble, P. (2000). The Maternal Effect
Mutation sésame Affects the Formation of the Male Pronucleus in Drosophila
melanogaster.   Developmental   Biology   *222*,   392–404.
https://doi.org/10.1006/dbio.2000.9718.

57.   Hayden, L., Chao, A., Deneke, V.E., Vergassola, M., Puliafito, A., and Di Talia, S.
(2022). Cullin-5 mutants reveal collective sensing of the nucleocytoplasmic ratio in
Drosophila   embryogenesis.   Curr   Biol   *32*,   2084-2092.e4.
https://doi.org/10.1016/j.cub.2022.03.007.

58.   Ahmad, K., and Henikoff, S. (2002). The histone variant H3. 3 marks active
chromatin by replication-independent nucleosome assembly. Molecular cell *9*, 1191–
1200.

59.   Chow, C.-M., Georgiou, A., Szutorisz, H., Silva, A.M. e, Pombo, A., Barahona, I.,
Dargelos, E., Canzonetta, C., and Dillon, N. (2005). Variant histone H3.3 marks
promoters of transcriptionally active genes during mammalian cell division. EMBO
reports. https://doi.org/10.1038/sj.embor.7400366.

60.   Jullien, J., Astrand, C., Szenker, E., Garrett, N., Almouzni, G., and Gurdon, J.B.
(2012). HIRA dependent H3.3 deposition is required for transcriptional reprogramming
following nuclear transfer to Xenopus oocytes. Epigenetics & Chromatin *5*, 17.
https://doi.org/10.1186/1756-8935-5-17.

61.   Chen, P., Zhao, J., Wang, Y., Wang, M., Long, H., Liang, D., Huang, L., Wen, Z.,
Li, W., Li, X., et al. (2013). H3.3 actively marks enhancers and primes gene
transcription via opening higher-ordered chromatin. Genes Dev *27*, 2109–2124.
https://doi.org/10.1101/gad.222174.113.

62.   Ng, R.K., and Gurdon, J.B. (2008). Epigenetic memory of an active gene state
depends on histone H3.3 incorporation into chromatin in the absence of transcription.
Nat Cell Biol *10*, 102–109. https://doi.org/10.1038/ncb1674.

63.   Strong, I.J.T., Lei, X., Chen, F., Yuan, K., and O'Farrell, P.H. (2020). Interphase-
arrested Drosophila embryos activate zygotic gene expression and initiate mid-blastula
transition events at a low nuclear-cytoplasmic ratio. PLOS Biology *18*, e3000891.
https://doi.org/10.1371/journal.pbio.3000891.

64.   Lanzotti, D.J., Kaygun, H., Yang, X., Duronio, R.J., and Marzluff, W.F. (2002).
Developmental Control of Histone mRNA and dSLBP Synthesis during Drosophila
Embryogenesis and the Role of dSLBP in Histone mRNA 3′ End Processing In Vivo.
Mol Cell Biol *22*, 2267–2282. https://doi.org/10.1128/MCB.22.7.2267-2282.2002.

65.   Wagner, E.J., Berkow, A., and Marzluff, W.F. (2005). Expression of an RNAi-
resistant SLBP restores proper S-phase progression. Biochem Soc Trans *33*, 471–473.
https://doi.org/10.1042/BST0330471.

66.   Harrison, M.M., Li, X.-Y., Kaplan, T., Botchan, M.R., and Eisen, M.B. (2011). Zelda
Binding in the Early Drosophila melanogaster Embryo Marks Regions Subsequently
Activated at the Maternal-to-Zygotic Transition. PLOS Genetics *7*, e1002266.
https://doi.org/10.1371/journal.pgen.1002266.

67.   Liang, H.-L., Nien, C.-Y., Liu, H.-Y., Metzstein, M.M., Kirov, N., and Rushlow, C.
(2008). The zinc-finger protein Zelda is a key activator of the early zygotic genome in
Drosophila. Nature *456*, 400–403. https://doi.org/10.1038/nature07388.

68.   Huang, S.-K., Whitney, P.H., Dutta, S., Shvartsman, S.Y., and Rushlow, C.A.
(2021). Spatial organization of transcribing loci during early genome activation in
Drosophila. Current Biology *31*, 5102-5110.e5.
https://doi.org/10.1016/j.cub.2021.09.027.

69.   Fogarty, P., Kalpin, R.F., and Sullivan, W. (1994). The Drosophila maternal-effect
mutation grapes causes a metaphase arrest at nuclear cycle 13. Development *120*,
2131–2142. https://doi.org/10.1242/dev.120.8.2131.

70.   Blythe, S.A., and Wieschaus, E.F. (2015). Zygotic Genome Activation Triggers the
DNA Replication Checkpoint at the Midblastula Transition. Cell *160*, 1169–1181.
https://doi.org/10.1016/j.cell.2015.01.050.

71.   Feijoo, C., Hall-Jackson, C., Wu, R., Jenkins, D., Leitch, J., Gilbert, D.M., and
Smythe, C. (2001). Activation of mammalian Chk1 during DNA replication arrest : a role
for Chk1 in the intra-S phase checkpoint monitoring replication origin firing. Journal of
Cell Biology *154*, 913–924. https://doi.org/10.1083/jcb.200104099.

72.   Moiseeva, T.N., Yin, Y., Calderon, M.J., Qian, C., Schamus-Haynes, S., Sugitani,
N., Osmanbeyoglu, H.U., Rothenberg, E., Watkins, S.C., and Bakkenist, C.J. (2019).
An ATR and CHK1 kinase signaling mechanism that limits origin firing during
unperturbed DNA replication. Proceedings of the National Academy of Sciences *116*,
13374–13383. https://doi.org/10.1073/pnas.1903418116.

73.   Chang, F.T.M., Chan, F.L., R. McGhie, J.D., Udugama, M., Mayne, L., Collas, P.,
Mann, J.R., and Wong, L.H. (2015). CHK1-driven histone H3.3 serine 31
phosphorylation is important for chromatin maintenance and cell survival in human ALT
cancer cells. Nucleic Acids Res *43*, 2603–2614. https://doi.org/10.1093/nar/gkv104.

74.　Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou,
D., Flockhart, I., Binari, R., Shim, H.-S., et al. (2015). The Transgenic RNAi Project at
Harvard Medical School: Resources and Validation. Genetics *201*, 843–852.
https://doi.org/10.1534/genetics.115.180208.

75.　Hunter, C., and Wieschaus, E. (2000). Regulated expression of nullo is required
for the formation of distinct apical and basal adherens junctions in the Drosophila
blastoderm. J Cell Biol *150*, 391–401. https://doi.org/10.1083/jcb.150.2.391.

76.　Gratz, S.J., Ukken, F.P., Rubinstein, C.D., Thiede, G., Donohue, L.K., Cummings,
A.M., and O'Connor-Giles, K.M. (2014). Highly specific and efficient CRISPR/Cas9-
catalyzed homology-directed repair in Drosophila. Genetics *196*, 961–971.
https://doi.org/10.1534/genetics.113.160713.

77.　Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C.,
Schiegg, M., Ales, J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine
learning for (bio)image analysis. Nat Methods *16*, 1226–1232.
https://doi.org/10.1038/s41592-019-0582-9.

Figure 1: Interphase Nuclear availability of H3 decreases more rapidly than H3.3 over the pre-ZGA cycles

(A) Maximum intensity projections of H3-Dendra2 (top) and H3.3-Dendra2 (bottom) interphase nuclei 45 seconds before nuclear envelope breakdown (NEB) from NC10-13. Images are pseudo-colored with non-linear look-up tables where purple indicates low and yellow indicates high intensities. **(B)** Average interphase nuclear pixel intensities for H3-Dendra2 and H3.3-Dendra2 45 seconds before NEB in NC10-13, normalized to the average individual NC10 values. H3 and H3.3 concentrations decrease over time, but H3 loss is relatively more rapid. **(C-D)** Summed (total) pixel intensities for each nucleus over time for NC11-13 normalized to the maximum NC11 values for H3-Dendra2 **(C)** and H3.3-Dendra2 **(D)**. Nuclear import plateaus after the first 5 mins for H3, but merely slows and does not plateau for H3.3 in NC13. **(E)** The fraction of photoconverted unbound H3.3-Dendra2 after NEB in NC11-13 (see materials and methods for details). The "free" pool of H3.3 falls with each cycle. (n=3 H3 and 5 H3.3 embryos in B-D and >= 5 embryos in F; Statistical comparisons for B can be found in Supplemental table 2).

Figure 2: The chaperone binding site determines H3 variant chromatin incorporation.

Figure 2: The chaperone binding site determines H3 variant chromatin incorporation

(A) Schematic of the Dendra2 tagged H3/H3.3 replacement chimeras at the endogenous H3.3A locus. S31A: H3.3 phosphosite (S) replaced with that of H3 (A), SVM: H3.3 chaperone binding site (AIG) replaced with that of H3 (SVM), and ASVM: all H3.3-specific amino acids replaced with those from H3. (B) Total intensities on mitotic chromatin of chimeras during NC10-13 normalized to their NC10 values. The same data for H3-Dendra2 and H3.3-Dendra2 are shown in Figure S1C. H3.3^{S31A} increases similarly to H3.3 while the constructs containing the H3 chaperone binding site decrease similarly to H3. (C) Interphase nuclear concentrations of chimeras 45 seconds before NEB during NC10-13 normalized to their NC10 values. H3-Dendra2 and H3.3-Dendra2 from Figure 1B included for reference. As seen for chromatin, nuclear accumulation generally follows the behavior of the chaperone binding site. (D) Schematic of H3.3 incorporation in control embryos and Hira^{ssm} mutants. H3.3 is imported to the nucleus, but the mutant Hira chaperone fails to incorporate H3.3. Hira mutants develop as haploids and undergo one additional fast nuclear division. (E, G, I) Representative maximum intensity projections during interphase and mitosis over NC10-14: interphase nuclei (top) and mitotic chromatin (bottom) for H3.3-Dendra2 (E), H3-Dendra2 (G), and H3.3^{ASVM}-Dendra2 (I). Images are pseudo-colored with non-linear look-up tables such that purple indicates low intensities and yellow indicates high intensities. H3 and H3.3^{ASVM} continue to accumulate on chromatin in the absence of Hira. (F, H) Total intensities of H3-Dendra2 (F) and H3.3^{ASVM}-Dendra2 (H) on mitotic chromatin in Hira^{ssm} embryos between NC10-14 normalized to their average NC10 values. Though H3.3^{ASVM} is successfully incorporated without active Hira, the chromatin amounts decrease more slowly than H3. (n=5 all chimeras, 3 H3 ssm, 4 H3.3 ssm, and 5 H3.3^{ASVM} ssm embryos. Statistical comparisons for B and C can be found in Supplemental tables 6-7).

Figure 3: Local N/C ratios determine H3 and H3.3 chromatin incorporation

Figure 3: Local N/C ratios determine H3 and H3.3 chromatin incorporation.

(A) Example NC11 control embryo with middle (blue) and pole (red) regions labeled as used in C and D. **(B)** Example NC11 shkl embryo with high (blue) and low (red) density regions labeled as used in E and F. **(C-D)** Total intensities on mitotic chromatin of H3-Dendra2 **(C)** and H3.3-Dendra2 **(D)** during NC11-13 in a representative control embryo where each point indicates a single nucleus. Total H3-Dendra2 intensities fall and H3.3-Dendra2 intensities rise uniformly between middle and pole regions within each cycle. **(E-F)** Total intensities on mitotic chromatin of H3-Dendra2 **(E)** and H3.3-Dendra2 **(F)** during NC11-13 in a representative shkl embryo where each point indicates a single nucleus. NC14 represents a partial extra division in the low-density region. Chromatin in the low-density region retains more H3 and incorporates less H3.3 within the same cell cycle compared to the high-density region. Similar results were observed in replicate embryos

(Figure S3A-B). **(G)** Gradient in the number of neighbors for each nucleus at its minimum

volume within a 20µm radius for the shkl embryo shown in B. **(H)** Direct and indirect

mechanisms of H3.3 incorporation in response to the N/C ratio. H3.3 incorporation could

be a direct result of reduced nuclear H3 availability. Here, the increasing demand for

nucleosomes with the increasing numbers of genomes would be met by H3.3. The N/C

ratio also controls transcription and cell cycle duration. H3.3 incorporation could be

downstream of either process. (Statistical significance was determined by 2-way ANOVA,

ns= $p > .05$, *** = $p < 0.001$).

Figure 4: Local N/C ratios differentially affect H3 and H3.3 nuclear availabilities

824 **Figure 4: Local N/C ratios differentially affect H3 and H3.3 nuclear availabilities**

825 **(A)** Schematic of how the N/C ratio might affect H3 and H3.3 chromatin incorporation

826 through loss of available H3. Bolded portion is the hypothesis under consideration. **(B-C)**

827 Total intensities over time for nuclei in representative NC13 control embryos binned by

828 the number of neighbors as in 3G for H3-Dendra2 **(B)** and H3.3-Dendra2 **(C)**. Total

intensity was normalized to the average maximum intensities achieved in NC13 and line

color represents the number of neighbors. In controls there is little variation in the number

of neighbors or the import of H3 and H3.3 across the length of the embryo. **(D-E)** Initial

slopes of nuclear import curves from representative NC13 control embryos from B and C

for H3-Dendra2 **(D)** and H3.3-Dendra2 **(E)** plotted by the number of nuclear neighbors.

Note the uniformity in the number of neighbors and similarity in nuclear import behaviors

in control embryos. **(F-G)** Total intensities over time for nuclei in representative NC13 shkl

embryos binned by the number of neighbors as in 3G for H3-Dendra2 **(F)** and H3.3-

Dendra2 **(G)**. Nuclear import and accumulation of H3 inversely correlate with the number

of neighbors, suggesting H3 nuclear import is N/C ratio sensitive. H3.3 nuclear import is

less N/C ratio sensitive than H3. Similar results were observed in replicate embryos

(Figure S3D-G). **(H-I)** Initial slopes of nuclear import curves from representative NC13

shkl embryos from F and G for H3-Dendra2 **(H)** and H3.3-Dendra2 **(I)** plotted by the

number of nuclear neighbors. The slopes reflect a faster H3 uptake in nuclei with fewer

neighbors and a slower H3 uptake in nuclei with more neighbors. Slopes in some nuclei

with more neighbors are near zero indicating that very little additional H3 is imported after

nuclear envelope formation. Though the slopes reduce with the number of neighbors for

H3.3, there is a non-negligible H3.3 import in the nuclei with the largest number of

neighbors.

Figure 5: H3.3 incorporation depends on cell cycle state but not H3 availability or zelda-dependent transcription

(A,C,E) Schematics of different parameters that may regulate H3.3 chromatin incorporation. Bolded portion is the hypothesis under consideration in B, D and F respectively. **(A)** Slbp-RNAi decreases the size of the available H3 pool. **(B)** Total intensities of H3.3-Dendra2 on mitotic chromatin in white-RNAi (control) and Slbp-RNAi backgrounds during NC10-13. H3.3 incorporation does not increase upon lowering H3 availability. Note that most Slbp-RNAi embryos are arrested in NC13 without dividing and therefore do not contribute to the mitotic NC13 data. **(C)** zelda-RNAi inhibits the majority of zygotic transcription allowing us to test if H3.3 incorporation depends on transcription.

(D) Total intensities of H3.3-Dendra2 on mitotic chromatin in white-RNAi (control) and
zelda-RNAi backgrounds during NC10-13 normalized to their NC10 values. H3.3
incorporation does not change upon inhibiting zelda-dependent transcription. **(E)** Chk1
(grp[1]) mutation prevents cell cycle slowing allowing us to test if H3.3 incorporation is
dependent on cell cycle state. **(F)** Total intensities of H3.3-Dendra2 on mitotic chromatin
in control and embryos from chk1[-/-] mothers during NC10-13. H3.3 incorporation is
reduced in both NC12 and NC13 indicating that cell cycle state though not cell cycle
duration regulated H3.3 incorporation. Note that these embryos are homozygous for
H3.3-Dendra2 and have double the fluorescent intensity compared to all previous
embryos. (n≥5 embryos, Statistical significance was determined by 2-way ANOVA, ns=
p>.05, *=p<.05, **=p<0.01, *** = p<0.001)