Systematically investigating and identifying bacteriocins in the human gut
microbiome

Dengwei Zhang[1], Yinai Zou[2], Yuqi Shi[1], Junliang Zhang[1], Jing Liu[1], Gengfan Wu[1], Jian Zhang[1],
Ying Gao[1], Muxuan Chen[2, *], Yong-Xin Li[1, 3, *]

[1] Department of Chemistry and The Swire Institute of Marine Science, The University of Hong
Kong, Pokfulam Road, Hong Kong, China

[2] Microbiome Medicine Center, Department of Laboratory Medicine, ZhuJiang Hospital,
Southern Medical University, Guangzhou 510280, China

[3] Lead contact

[*] Correspondence: muxuanchen@126.com (M.C.), yxpli@hku.hk (Y.L.)

Summary

Human gut microbiota produces unmodified bacteriocins, natural antimicrobial peptides that protect against pathogens and regulate host physiology. However, current bioinformatic tools limit the comprehensive investigation of bacteriocins' biosynthesis, obstructing research into their biological functions. Here, we introduce IIBacFinder, a superior analysis pipeline for identifying unmodified class II bacteriocins. Through large-scale bioinformatic analysis and experimental validation, we demonstrate their widespread distribution across the bacterial kingdom, with being niche-specific and enriched in the human microbiome. Analyzing over 280,000 bacterial genomes, we reveal the diverse potential of human gut bacteria to produce these bacteriocins. Guided by meta-omics analysis, we synthesized 26 hypothetical bacteriocins from gut commensal species, with 16 showing antibacterial activities. Further *ex vivo* tests show minimal impact of narrow-spectrum bacteriocins on human fecal microbiota. Our study highlights the huge biosynthetic potential of unmodified bacteriocins in the human gut, paving the way for understanding their biological functions and health implications.

Keywords

Bacteriocin, Antimicrobial peptides, Natural products, Antimicrobial activity, Human microbiome, Genome mining, Metagenomics, Omics analysis

Introduction

Bacteria are ubiquitously present on Earth, often found in a communal fashion where each species will compete for limited nutrients and space[1]. These common antagonistic interactions are crucial not only to influence the composition of their members but also to maintain the community stability[2,3]. The interference competition can be governed by various types of molecules, such as polyketides[4], protein toxins[5], non-ribosomally synthesized peptides (NRPs)[6], and ribosomally synthesized antimicrobial peptides that are referred to as bacteriocins[7,8]. Despite exhibiting incredible sequence and structural diversity, bacteriocins are usually categorized into two major classes: those that undergo posttranslational modifications by enzymatic tailoring (class I bacteriocins) and unmodified peptides (class II bacteriocins)[9]. On the basis of posttranslational modifications, class I bacteriocins can be further grouped into more than 40 subgroups such as lanthipeptide and lasso peptide, whereas class II bacteriocins are generally divided into four subgroups (class IIa, YGNG-motif containing bacteriocins; class IIb, two-peptide bacteriocins; class IIc, leaderless bacteriocins; class IId, other linear bacteriocins)[10].

Due to its antimicrobial nature, bacteriocins have been regarded as promising antibiotic alternatives or adjuncts to contain the antimicrobial resistance threats of current antibiotics[7]. Compared to class I bacteriocins, unmodified class II bacteriocins are simpler for their production in heterologous systems and chemical synthesis, thereby rendering it more practical for antimicrobial discovery at a large scale. Although a few class II bacteriocins have been identified generally via traditional isolation techniques (Table S1), they exhibit a variety of biological activities, such as antifungal and antiviral activities[11,12]. This makes class II bacteriocins promising candidates for the discovery of antimicrobials. Additionally, recent studies, albeit limited, have broadened the functions of bacteriocins beyond their antimicrobial nature, suggesting a role in interacting with the host[13–15]. Our recent study of antimicrobial peptides from human microbiota further suggests the largely unexplored potential of bacteriocins in the gastrointestinal tract (GIT) and their potential physiological roles in human health[16,17]. This further emphasizes the significance of studying host-associated class II bacteriocins, particularly those from the GIT where bacteriocins are crucial to human health[18,19].

In the GIT, bacteriocins act as essential protectors impacting human health. For example, plantaricin EF, a class IIb bacteriocin, can maintain the integrity of the epithelial barrier, reducing the effects of obesity-inducing diets in mice on a high-fat diet[15]. Another instance demonstrated the immunomodulatory activity of a lactococcin 972-like class IId bacteriocin from *Bifidobacterium longum* subsp. *infantis*[20], a bacterium found in the infant gut. Given that the human microbiome contains 150 times more genes than the human genome[21], class II bacteriocins within the human microbiome represent an untapped treasure for antimicrobial discovery and play a crucial role in mediating human health. Although previous investigations on hundreds of human GIT reference genomes or human metagenomic data revealed limited biosynthetic potential of class II bacteriocin in human gut microbiome[22,23], the overall understanding is significantly incomplete, considering over 280,000 reference genomes reconstructed from the human gut microbiome[24]. It is imperative to conduct a comprehensive investigation into class II bacteriocins in the human gut microbiome to better understand their biosynthetic potential, chemical diversity, distribution, and possible biological functions.

In the meta-omics era, with tremendous genomic data sets available, bacteriocin discovery has evolved from a traditional activity screening to a more prominent emphasis on genomics-guided discovery via predicting biosynthetic gene clusters (BGCs)[9]. The BGCs of class I or class II bacteriocins typically include the precursor genes, and other associated genes such

as transporter and peptidase (hereinafter referred to as context genes). Advanced by the rapid
development of computational techniques, a range of dedicated bioinformatic tools have been
in place for bacteriocin mining using distinct algorithms (Table 1). However, none of these
tools are specifically designed for high-precision identification of class II bacteriocins. Even
though class II bacteriocins belong to the group of antimicrobial peptides (AMPs) that do not
undergo post-translational modifications, they need a unique mining approach compared to
the general AMP prediction[25–27]. This is because while AMPs encompass entire small open
reading frames, the maturation of most class II bacteriocins involves the removal of the leader
peptide through peptidase, followed by secretion via transporter. Moreover, most tools predict
class II bacteriocins solely by identifying precursor genes. Tools like RMSCNN[28], BaPreS[29],
and BPAGS[30] identify bacteriocin precursor sequences, but they do not differentiate between
modified and unmodified bacteriocins. On the other hand, some tools, such as BAGEL4[31] and
PRISM 4[32], concentrate only on precursor sequences similar to those of known bacteriocins.
The most commonly used tool, antiSMASH[33], has limited detection rules due to its focus on
overall secondary metabolites, constraining a comprehensive identification of class II
bacteriocins. Therefore, developing a more dedicated tool for class II bacteriocin mining is a
prerequisite for interrogating them in the human gut microbiome.

Here, we commenced with the development of a specialized tool called IIBacFinder (Class II
Bacteriocin Finder), which integrates the detection of class II bacteriocins through both
precursor genes and context genes. Demonstrated by recovering known class II bacteriocins
from the genomes of bacteriocin producers, IIBacFinder outdoes the two wildly used tools,
BAGEL4 and antiSMASH, in detecting class II bacteriocins. Alongside experimental validation,
the systematic mining of ~260,000 bacterial genomes from RefSeq database revealed the
underappreciated biosynthetic potential of class II bacteriocins in the bacterial kingdom,
particularly in Gram-negative bacteria. Those class II bacteriocins exhibited high niche-
specificity and were enriched in the human microbiome. Leveraging the comprehensive
analysis of ~280,000 reference genomes reconstructed from human gut metagenomes, we
gained deep insights into the biosynthetic capacity and chemical diversity of class II
bacteriocins. Analyzing metagenomics and metatranscriptomics data of healthy human gut,
we prioritized and chemically synthesized 26 predicted class II bacteriocins that were
prevalent and abundant in healthy human gut microbiomes. They showed diverse narrow-
spectrum antibacterial activities but had minimal impacts on human fecal microbiota in an *ex
vivo* assay. Our work demonstrates the previously underappreciated biosynthetic potential of
unmodified bacteriocins in the human gut microbiome, which might contribute to maintaining
microbiome homeostasis and impacting human health.

Results

The development of IIBacFinder pipeline specializing in detecting unmodified bacteriocins

Several tools have been developed for bacteriocin identification with distinct mining strategies (Table 1), but they underperform in identifying class II bacteriocins primarily due to solely relying on precursor genes. To circumvent these limitations, we set out to develop an improved tool dedicated to identifying class II bacteriocins by integrating the detection of precursor genes and context genes. Given that profile hidden Markov models (pHMMs) offer a straightforward and rapid detection method to identify homologs within a specific protein family[34,35], we constructed the pHMM library of precursor genes and context genes by curating known sequences and detecting their homologs from the UniRef100 database (Figure S1A, Tables S1, 2). Leveraging the library of 1,153 pHMMs of precursor genes and 37 pHMMs of context genes (Tables S3 - 5), we could comprehensively pinpoint class II bacteriocins by either directly detecting precursor genes or indirectly detecting novel bacteriocins genes through anchoring context genes (Figure 1A, Figure S1B). We termed this pipeline IIBacFinder (Class II Bacteriocin Finder).

Following the detection of precursor genes, we sought to integrate the prediction of leader, which is a marked difference between class II bacteriocins and generic AMPs. Two types of leaders, the double-glycine leader and the sec-dependent leader, are generally present in class II bacteriocins. To streamline the prediction of double-glycine leader, we adopted and tuned a RiPP leader detection tool, NLPPrecursor[36], for the cleavage site prediction. By employing 10-fold cross-validation, we determined that the modified algorithm could attain an accuracy of 98.0 % when factoring in cleavage points that are ±3 amino acids (AAs) away from the actual prediction site (Figure S1C, D). On the other hand, we applied SignalP 6.0[37] to predict the cleavage site for the sec-dependent leader. Finally, the predicted core peptides were compared to known AMP sequences from four AMP databases[38–41], to gain an insight into their sequence novelty (Figure 1A).

Upon the development of IIBacFinder, we next sought to evaluate its performance in identifying class II bacteriocins. We curated 137 bacterial genomes containing 205 complete precursor sequences of known class II bacteriocins, spanning different leader types and subclasses (Table S6). Among them, 191 sequences were unutilized when constructing the pHMM library of precursor genes, while 12 sequences were not because they were unintentionally excluded when building pHMMs. We then applied three tools (antiSMASH 6, BAGEL4, and IIBacFinder) to these genomes for detecting class II bacteriocins. Reassuringly, IIBacFinder could recover all bacteriocin precursor sequences (205/205, 100%), including 12 sequences that were new to IIBacFinder, while BAGEL4 and antiSMASH 6 detected 134 (65.4%) precursor sequences and 112 (54.6%) corresponding bacteriocin gene clusters, respectively (Figure 1B). Notably, their performance in identifying leaderless class IIc bacteriocins was significantly lower. The performance benchmarking demonstrated that IIBacFinder has a superior detection capability for unmodified class II bacteriocins compared to the other two commonly used tools.

Underappreciated biosynthetic capacity of class II bacteriocins in the bacterial kingdom

Having attested to IIBacFinder's remarkable ability to identify class II bacteriocins, we then applied it to 257,997 bacterial genomes from the NCBI RefSeq database[42], in order to assess

the global biosynthetic capacity of unmodified bacteriocins in the bacterial kingdom. These
genomes spanned 56 phyla and 4,259 genera, corresponding to 96,325 Gram-positive and
161,472 Gram-negative bacteria. Using IIBacFinder, a total of 540,587 precursor sequences
were identified from 133,806 genomes. Among them, 387,888 (71.75%) and 152,598 (28.23%)
precursor sequences were detected from Gram-positive and Gram-negative bacteria,
respectively. For comparison, antiSMASH 6 could detect 206,288 RiPP-like BGCs
(encompassing BGCs encoding class II bacteriocins and other unclassified metabolites) in
119,697 genomes, consisting of 108,962 (52.84%) and 97,265 (47.16%) BGCs from Gram-
positive and Gram-negative bacteria; meanwhile, BAGEL4 detected 137,942 precursor
sequences with 93.36% being from Gram-positive genomes, indicating BAGEL4 biased the
prediction in Gram-positive bacteria in which most known bacteriocins were discovered
(Figure S2A, B).

To reduce the probability of false prediction, we only included 481,654 high-confidence
sequences for downstream analysis (Figure S2C). The length of core peptides was
approximately 40 amino acids for double-glycine and leaderless bacteriocins, but around 70
amino acids for sec-dependent bacteriocins (Figure S2D). Compared to known AMPs
deposited in four databases (i.e., APD3, DRAMP, DBAASP, and dbAMP2), they showed less
homology, with most being less than 40% identity (Figure S2E). Meanwhile, class II
bacteriocins had higher median length than AMPs. By analyzing two physicochemical features,
which play a pivotal role in elucidating the activity of AMPs, we found that double-glycine or
leaderless bacteriocins exhibited a lower number of electronic charges than AMPs (Mann–
Whitney U test, $P < 0.001$) but had higher GRAVY index scores, suggesting they are likely to
be hydrophobic. In contrast, significantly lower GRAVY scores than AMPs implied that sec-
dependent bacteriocins were more likely to be hydrophilic (Figure S2F).

We then conducted a rarefaction analysis on 8,610 precursor clusters that were categorized
from 481,654 sequences using a 50% identity threshold, aiming to assess the genetically
encoded biochemical diversity. The curve astonishedly showed that Gram-negative bacteria
had a higher biosynthetic potential than Gram-positive bacteria (Figure 2A). We then
decorated the GTDB bacterial tree with predicted precursor sequences to reveal the
biosynthetic capacity at the phylum level. Through this examination, we found that the
bacteriocin biosynthetic capacity stood out in several most sequenced phyla, such as
Firmicutes (13,401 unique sequences), Proteobacteria (13,104), Bacteroidota (4,494), and
Actinobacteriota (3,801) (Figure 2B). Among them, the bacteria from Firmicutes were the most
prolific producers, with the average number of precursors per genome being 4.97, in line with
the fact that most known bacteriocins have been discovered in this phylum. When down to the
genus level, *Streptococcus*, as expected, topped the list of the bacterial tree of life. Other
Gram-negative genera, such as the soil bacteria *Rhizobium* and *Bradyrhizobium*, also
displayed promising biosynthetic capacity (Figure 2C). The overall survey indicated the
previously underappreciated bacteriocin biosynthetic capacity in the bacterial kingdom,
particularly in Gram-negative bacteria.

To further demonstrate the prediction reliability, we arbitrarily selected and chemically
synthesized 28 hypothetical bacteriocins (HBs) with the length of deduced core peptides < 50
amino acids for experimental validation, but 7 failed (Table S7). The remaining 21 sequences
consisted of 13 double-glycine sequences, 5 sec-dependent sequences, and 3 leaderless
sequences from seven phyla (i.e., Firmicutes, Firmicutes_A, Firmicutes_C, Cyanobacteria,
Actinobacteria, Bacteroidota, Proteobacteria). Moreover, only HB17 could be detected by
BAGEL4, while six HBs with double-glycine precursors could be captured by antiSMASH 6 as
RiPP-like BGCs (Figure 2D). We examined their antibacterial potential against 19 Gram-
positive and 9 Gram-negative indicator strains from four phyla at 100 µg/mL concentration

(Figure 2D). The assay showed that 17 of 21 HBs could inhibit the growth of at least one
indicator strain (inhibition ratio > 25%, Dunnett's test *P* < 0.05). The active HBs primarily
targeted the Gram-positive bacteria from phyla Firmicutes and Actinobacteria and barely
inhibited Gram-negative bacteria. They displayed either narrow-spectrum activity (e.g., HB1
and HB2) or relatively broad-spectrum activity (e.g., HB7). Among them, 9 HBs (three from
Gram-negative bacteria and six from Gram-positive bacteria) exhibited a robust activity with
over 75% inhibition ratio, with minimal inhibitory concentrations (MICs) ranging from 0.78 to
100 µg/mL (Figure 2D). This suggests that IIBacFinder has the potential to identify genuine
new unmodified bacteriocins, a capability that the other two tools lacked.

Class II bacteriocins exhibit niche-specific enrichment in host-associated bacteria

The global inspection of class II bacteriocins revealed their wide distribution among diverse
bacterial phyla, which might account for the previous concept that bacteriocin producers are
widespread in natural environments[43]. To understand their biosynthetic potential in diverse
ecosystems, we applied IIBacFinder to ~50,000 metagenome-assembled genomes (MAGs)
recovered from metagenomes of diverse habitats, including host-associated environments
(primarily from human), aquatic environments, engineered environments, and terrestrial
environments[44]. We identified 4,956 unique high-confidence precursor sequences from the
genomic catalog of earth's microbiomes, half of which (2,482 sequences) were from host-
associated MAGs, followed by the engineered, aquatic, and terrestrial environments (Figure
3A). Notably, the bacteriocins identified from human contributed most to the host-associated
bacteriocin (Figure 3B). Interestingly, we observed that class II bacteriocins exhibited high
niche-specify, with >90 percent of unmodified bacteriocin sequences found in one specific
ecosystem (Figure 3C).

Although the fact that the largest number of bacteriocins were detected in host-associated
MAGs is informative, this might be skewed by the uneven number of reconstructed MAGs.
Therefore, we then adopted rarefaction analyses to address this bias (Figure 3D). The
rarefaction curve still indicated host-associated MAGs presented the greatest biosynthetic
potential than other ecosystems, with being around three times over aquatic and engineered
environments. The analysis of environmentally diverse bacteriocins implied class II
bacteriocins' potential to interact closely with community members in the host. They may also
be utilized to interact with the host itself, particularly in the case of human hosts[20].

Human gut microbiota has the potential to produce diverse health-beneficial unmodified bacteriocins

Recognizing the prevalence of class II bacteriocins in the human microbiome, we focused on
unmodified bacteriocin production in the human gut, where numerous microbes impact health
by producing various metabolites[45]. We, therefore, applied IIBacFinder to ~280,000 bacterial
genomes recovered from the human gut microbiome[24], resulting in 82,806 high-confidence
precursor sequences, with 6,238 being unique (100% identity). More than half of them were
double-glycine precursors (4,186, 67.10%), followed by sec-dependent leader precursors (765,
12.26%) and leaderless precursors (610, 9.78%). At the class level, Clostridia, Bacilli, and
Bacteroidia were the most abundant producers. Despite not being predominant in the human
gut microbiota, *Streptococcus* could produce the most diverse bacteriocins, aligning with its
global bacterial biosynthetic potential. Other key human gut bacteria like *Prevotella*,
Faecalibacterium, and *Bacteroides* also exhibited high biosynthetic potential (Figure 4A).

Although this analysis indicated the high biosynthetic potential of human gut bacteria, only a
few class II bacteriocins have been identified from gut microbiota yet[18]. We, thus, wondered
whether these predicted unmodified bacteriocins were chemically distinct from known ones.
We thus compared the 6,238 unique sequences with 298 known bacteriocin sequences at
varying identity thresholds of 50% - 90% (Figure 4B, Figure S3). To our surprise, we found
379 (6.07%) and 143 (2.29%) homologous sequences at the identity of 50% and 90%,
respectively, indicating that gut microbiota could produce similar bacteriocin products, but the
majority remained uncharacterized. Among the most diverse were UviB homologies produced
by *Clostridium perfringen*[46]. Some bacteriocins have established functions in mediating host
health, for example, a broad-spectrum class II bacteriocin, ABP118, can protect mice against
infection with pathogen *L. monocytogenes*[47]. These findings suggest that unmodified
bacteriocins with protective functions may have a wide range of health-promoting effects.

Recent advancements in functional metagenomics have facilitated the discovery of bioactive
RiPPs in the largely uncharted human microbiota, and have established links to their potential
physiological roles[48]. We next looked at evidence of their transcription in 724
metatranscriptomes of healthy individuals from three cohorts to look for insight into their
possible physiological roles. We found 1,966 (31.52%) sequences were actively transcribed
in at least one sample, but more sequences could be detected with an increase in sample size
as indicated by rarefaction analysis (Figure 4C). Most of actively transcribed bacteriocins are
predicted novel (Figure 4D), might involving uncharted roles in the human gut. Furthermore,
multiple distinct bacteriocin sequences were found to be transcribed in each individual, with a
median number of 139, 88, and 122 in three cohorts (Figure 4E), implying that hundreds of
transcribed bacteriocins might create a protective barrier for pathogen invasion in a synergistic
manner[49].

Omics analysis-guided prioritization and identification of unmodified
bacteriocins from the healthy gut microbiome

Inspecting the class II bacteriocins from human gut MAGs allowed us to appreciate their
significant biosynthetic potential in the human gut. Next, we set out to experimentally
characterize the bioactive class II bacteriocins that might play a crucial role in maintaining
healthy human gut homeostasis. As the species abundance and sequence depth would skew
the genome assembly from metagenomics data, we prioritized putatively bioactive bacteriocin
using mapping-based analysis of metagenomic and metatranscriptomic data instead of those
that were the most abundant in gut MAGs. To do so, we first created a reference database of
class II bacteriocin by applying IIBacFinder to ~700,000 bacterial genomes from multiple
sources, including RefSeq[42], human bodies[24,50,51], earth environment[44], ocean[52], mouse and
ruminant gastrointestinal tract[53,54]. Following the construction of this database encompassing
32,735 non-redundant high-confidence class II bacteriocin sequences, we examined their
profiles by re-visiting 1,901 metagenomic and 724 metatranscriptomic datasets from healthy
individuals (Figure 5A). To remove the redundancy, we computed their abundance at the
cluster level (80% identity and 80% coverage) rather than at the individual level, and 3,581
bacteriocin clusters were detected in this omics analysis. Given the fact that bioactive
bacteriocin could inhibit the growth of other bacteria and thus might regulate the overall
bacterial community, we ranked them by considering (1) their prevalence and (2) abundance,
(3) negative association with bacterial species, and (4) association with bacterial community
diversity assessed by Shannon index (Figure 5A). This analysis allowed us to prioritize
bioactive bacteriocins in the human gut microbiome (Table S8).

Finally, on the basis of the ranking analysis, we scrutinized and selected 40 clusters of human
gut microbiota-derived hypothetical bacteriocins (HGM-HBs) with a length of core peptide <
50 amino acids for experimental validation (see STAR Methods). We attempted to chemically
synthesize the core peptides from their representative sequences; however, 14 of these
attempts were unsuccessful. Consequently, we had 26 HGM-HBs available for examination
(Table S9). All of them were identified from human gut-derived bacterial genomes. None was
detected by BAGEL 4, while eight were identified by antiSMASH 6 (seven within RiPP-like
BGCs and one within NRPS BGC). Meanwhile, HGM-HP11 was identical to bacteroidetocin
A[55], but we did not include the intrinsic disulfide bonds for chemical synthesis as the putative
disulfide bonds in other HGM-HPs were not predictable. Likewise, we initially examined their
antibacterial activity against 21 indicator strains encompassing 13 clinical pathogens and 8
gut commensal bacteria at a concentration of 100 µg/mL. Under a cutoff of inhibition ratio >
25% and P < 0.05, 16 peptides (70%) were found to be active against, generally, Gram-
positive bacteria. Notably, 8 HGM-HBs, including HGM-HP11, effectively inhibited at least one
indicator strain with a > 75% inhibition ratio. Further activity examination for three HGM-HBs
(HGM-HP1, HGM-HP7, and HGM-HP17) showed their MICs ranging from 3.125 to 50 µg/mL
(Figure 5B). The *in vitro* assay indicated that the human gut microbiota-derived class II
bacteriocin could inhibit the growth of clinical pathogens and gut commensals.

Narrow-spectrum bacteriocins tweak human fecal microbiota

Although *in vitro* activity screening against a limited panel of indicator strains demonstrated
active unmodified bacteriocins from the human gut microbiome, whether and how they affect
the entire gut microbial community remains unclear. To address this, we utilized a human
fecal-derived *ex vivo* microbial community to model the effects of bacteriocins on gut
microbiota (Figure 6A). We tested three active bacteriocins (HGM-HP1, HGM-HP7, and HGM-
HP17) and an inactive one (HGM-HP14) as determined by *in vitro* assay (Figure 5B).
Vancomycin was included as a positive control. We investigated the effects of two
concentrations, 1 µg/mL and 100 µg/mL, to understand how these bacteriocins modulate the
microbial community at a low concentration, and to determine if they exhibit a broader activity
spectrum than previously observed *in vitro* which could be indicated at a high concentration.
Following a 48-hour treatment with bacteriocin peptides or vancomycin, no pronounced
change in fecal microbiota composition was observed, except with 100 µg/mL of vancomycin
(Figure 6B). Additionally, only the 100 µg/mL concentration of vancomycin, and not the other
treatments, significantly decreased microbial diversity as indicated by the Shannon diversity
index (Figure 6C), and markedly altered the microbial community structure as shown by
principal coordinate analysis based on Bray-Curtis distances; however, 100 µg/mL of HGM-
HP14 and HGM-HP17 slightly changed the overall microbial community (Figure 6D, Figure
S4A, B). These results suggest that putative bacteriocins have only slight effects on human
fecal microbiota, even at high concentrations.

We then used MaAsLin2[56] to examine the abundance difference in bacterial species between
treatment and control groups, identifying a variable number of species that could be affected
by bacteriocins or vancomycin (FDR-adjusted P value < 0.05) (Figure 6E, Figure S4C).
Results indicated that a low concentration (1 µg/mL) of bacteriocins had minimal impact on
fecal bacterial species. However, at a high concentration (100 µg/mL), two bacteriocins, HGM-
HP14 and HGM-HP17, significantly affected the abundance of 17 and 15 bacterial species,
respectively (Figure 6E). Notably, HGM-HP14, which showed no activity *in vitro*, decreased
the abundance of 10 bacterial species in the larger microbial community, suggesting that *in*
vitro screening might overlook some authentic bacteriocins. In contrast, the other two active
bacteriocins (HGM-HP1 and HGM-HP7) had negligible effects on fecal microbiota, even at

100 µg/mL. As expected, the broad-spectrum antibiotic vancomycin significantly affected
numerous bacterial species, particularly at the 100 µg/mL concentration, where it decreased
the abundance of 53 species (Figures 6E, F). Vancomycin primarily reduced species from the
classes Bacilli and Bacteroidia. Similarly, the narrow-spectrum bacteriocins HGM-HP14 and
HGM-HP17 also reduced several bacterial species from these classes. Taken together, the
ex vivo assay demonstrated slight effects of narrow-spectrum bacteriocins on human fecal
microbiota.

Discussion

The human gut microbiota is a rich source of bacteriocins that might combat pathogenic bacteria invasion and participate in human physiology[18,19]. Although a few efforts have been made to decipher their distribution in human gut microbiome[22,23], the unmodified bacteriocins remain incompletely characterized. In this study, we conducted the first large-scale survey, to our best knowledge, on unmodified class II bacteriocins in the bacterial kingdom and human gut microbiome by applying an in-house bioinformatic tool. Through systematic investigation in combination with experimental validation, we demonstrated novel class II bacteriocins throughout the tree of bacterial life as well as in the human gut microbiome. These investigations will not only disclose that the class II bacteriocins are largely untapped resources for antimicrobial discovery but also act as a crucial pathway for unraveling bacteriocins-mediated interactions between the microbiome and human host.

Antimicrobial resistance (AMR) is a major global health concern that requires immediate, effective responses. Bacteriocins, due to their diversity, high efficacy, and ease of bioengineering, are emerging as potential alternatives to traditional antibiotics[7]. They often specifically target pathogens, thus limiting the damage caused by broad-spectrum antibiotics to existing microbial communities and potentially reducing adverse health effects[57]. For instance, Thuricin CD targets *Clostridium difficile* with minimal impact on the microbial community, yet with efficacy comparable to broad-spectrum antibiotics[58]. Our study found that class II bacteriocins exhibit narrow-spectrum activity, suggesting their potential to target pathogens specifically. Unmodified class II bacteriocins are preferred over modified counterparts for heterologous production and chemical synthesis, making them more practical for large-scale antimicrobial discovery efforts. In this regard, big data omics mining offers a faster and more high-throughput approach than traditional screening methods. To enhance the detection of class II bacteriocins, we developed a specialized software pipeline that identifies them by scanning pHMMs of signature genes. This advanced tool has unveiled the untapped biosynthetic potential for class II bacteriocins within the bacterial kingdom and the human gut microbiome, facilitating their large-scale discovery. Their genomics-guided discovery could be achieved by cell-free protein synthesis systems or robotic automation systems; two techniques have been successfully applied to AMP identification and modified bacteriocin characterization[59–61]. Hence, we believe that future advances in techniques can accelerate the discovery of class II bacteriocins, providing a promising strategy to address the AMR issue.

Recent studies have uncovered and experimentally confirmed a multitude of AMPs originating from the global microbiome[25] and the human gut microbiome[26,27], highlighting the overlooked potential of antagonistic AMPs within the bacterial realm, particularly within the human gut microbiota. While there are still questions about whether these AMPs are genuinely produced, secreted, and undergo modifications similar to bacteriocin biosynthesis, they hold potential as new bacteriocins. Unlike previous studies, our research primarily focuses on unmodified bacteriocins that are likely naturally produced and secreted. We pay particular attention to mature peptides, as opposed to the full coding sequences typically examined in AMP prediction. We have determined that synthetic bacteriocins, originating from the human gut microbiome, demonstrate antibacterial properties. However, their *in vivo* functions might extend beyond this, as they also display additional antimicrobial activities such as combating biofilms, fungi, and viruses[2,64]. For instance, enterocin A inhibits biofilm formation in three known pathogens, which can aid in reducing harmful pathogens in the human gut[58]. Furthermore, plantaricins EF and JK[11], known for their antifungal properties, could influence human health indirectly by regulating the vital fungal community[62]. On top of antimicrobial activities, unmodified bacteriocins can also directly interplay with the human host, exhibiting

various activities such as anticancer[58] and immunomodulation[63–65]. Notably, plantaricin EF
increases the production of proteins that fortify the intestinal lining, preserving its integrity[15].
Our comprehensive review reveals that human gut bacteria can produce a variety of class II
bacteriocins, suggesting their potential roles in promoting human health. Consequently, we
advocate for considering the role of class II bacteriocins in studies exploring the relationship
between gut microbiota and human health.

In conclusion, we have developed an advanced tool, IIBacFinder, to provide a comprehensive
overview of unmodified bacteriocins within the bacterial kingdom, with a particular focus on
the human gut microbiome. Our study reveals the previously underestimated chemical
repertoire of unmodified bacteriocins, revealing an untapped treasure for antimicrobial
discovery. Diverse class II bacteriocins encoded by the human gut microbiome imply their
under-perceived roles that cannot be ignored in future studies. Thus, our study paves the way
for antimicrobial discovery from the untapped human microbiome and for deepening our
understanding of its role in human health and disease in future studies.

Limitations of the study

Identifying bacteriocins produced by human microbiota is crucial for comprehending the
communication mechanisms within gut microbiota and between the microbiota and the
human host. Although our tool, IIBacFinder, significantly improves the identification of
unmodified class II bacteriocins, it may still yield false predictions, especially when detecting
new bacteriocins based on context genes. Due to the limited number of characterized class
II bacteriocins, our current understanding makes it difficult to differentiate real bacteriocins
from false predictions without experimental validation. Moreover, while we can identify new
bacteriocins through anchoring context genes, we may miss other unknowns that don't fit
our current understanding of class II bacteriocin biosynthetic logic, leading to an incomplete
exploration of their biosynthetic potential. Future advancements in high-confidence
techniques are needed to identify authentic bacteriocins accurately.

In this study, we chemically synthesized predicted mature peptides of unmodified
bacteriocins, which may not accurately represent their natural mature form. Despite not
undergoing post-translational modifications, unmodified bacteriocins might have other
intramolecular features, like disulfide bonds and a reduced state of cysteine[10,66]. Additionally,
bacteriocin synergies might be common in the human gut[49], considering that hundreds of
bacteriocins are actively transcribed in individuals and many bacteriocin gene clusters often
contain more than one precursor gene. Thus, the *in vivo* functions of synthetic bacteriocins
might differ from our *in vitro* characterization. Other potential functions of these bacteriocins
remain unknown and require further investigation. Despite these limitations, our study
emphasizes that the gut microbiota could produce a variety of class II bacteriocins that might
interact with the microbial community or human host.

Resource availability

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Yong-Xin Li (yxpli@hku.hk).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- The bacterial genomes were collected from eight publicly available datasets, including the NCBI Assembly RefSeq database (https://ftp.ncbi.nlm.nih.gov/genomes/refseq/), human microbiome (http://segatalab.cibio.unitn.it/data/Pasolli_et_al.html), UHGG database (, https://www.ebi.ac.uk/ena/browser/view/PRJEB33885), HRGM database (https://www.mbiomenet.org/HRGM/), Earth's microbiome (https://portal.nersc.gov/GEM/genomes/), ocean microbiome (https://www.ebi.ac.uk/ena/browser/view/PRJEB45951), mouse (https://zenodo.org/records/4840600) and ruminant gut microbiome (https://www.ebi.ac.uk/ena/browser/view/PRJNA657473).
- The raw data for metagenomes and metatranscriptomes are publicly available in NCBI-SRA under the BioProject IDs: PRJNA422434, PRJNA319574, PRJEB4336, PRJEB1220, PRJDB4176, PRJNA354235, PRJEB51894, and PRJNA398089.
- Publicly available AMP data can be accessed from four databases – APD3: https://aps.unmc.edu/downloads; dbAMP v2.0: https://awi.cuhk.edu.cn/dbAMP/download.php; DRAMP v3.0: http://dramp.cpu-bioinfor.org/downloads/; DBAASP v3.0: https://dbaasp.org/.
- Protein family models can be found in the Pfam database (version 34.0, http://ftp.ebi.ac.uk/pub/databases/Pfam/releases/Pfam34.0) and NCBI database (version 11.0, https://ftp.ncbi.nlm.nih.gov/hmm/11.0/).
- Other source data required to reanalyze the data reported in this paper can be found in supplementary materials or the Zenodo repository (https://zenodo.org/records/13588376).
- Raw read sequences of the shotgun metagenomic sequences generated in this study were deposited at Sequence Read Archive (SRA, NCBI) under the accession of PRJNA1201247.
- The IIBacFinder codes can be found at https://github.com/ZhangDengwei/IIBacFinder and https://zenodo.org/records/14292149.

Acknowledgments

The author would like to thank Prof. Wang (University of Nebraska Medical Center) for graciously providing the AMP sequences from the latest APD3 database in response to an email request. The authors also thank Vecteezy.com for distributing free vector art at https://www.vecteezy.com, which is partially adopted and modified in the graphical abstract.

Author contributions

Conceptualization: Y.L. and D.Z. Methodology: D.Z., Y.Z., Y.S., J.Z., J.L., G.W., J.Z., Y.G. Investigation: Y.L. and M.C. Visualization: Y.L. and D.Z. Supervision: Y.L. and M.C. Writing— original draft: D.Z. Writing—review & editing: Y.L. and D.Z.

Declaration of interests

The authors declare no competing interests.

Funding

This work is partially funded by three Hong Kong Research Grants Council General research grants (HKU17101324, HKU17115322, and HKU17102123) and National Natural Science Foundation of China (NSFC82341218).

STAR Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
Bacillus timonensis	Leibniz Institute DSMZ	DSM 25372
Enterococcus caccae	Leibniz Institute DSMZ	DSM 19114
Enterococcus ureasiticus	Leibniz Institute DSMZ	DSM 23328
Enterococcus termitis	Leibniz Institute DSMZ	DSM 22803
Enterococcus rivorum	Leibniz Institute DSMZ	DSM 104544
Enterococcus faecium	American Type Culture Collection	ATCC 35667
Enterococcus faecalis	American Type Culture Collection	ATCC 29212
Lactobacillus delbrueckii	Leibniz Institute DSMZ	DSM 20081
Lactococcus lactis subsp. *cremoris*	N/A	MG1363
Paenibacillus ihumii	Leibniz Institute DSMZ	DSM 100664
Staphylococcus aureus	American Type Culture Collection	ATCC 43300
Streptococcus pneumoniae	N/A	CMCC 31002
Listeria monocytogenes	American Type Culture Collection	ATCC 19115
Clostridium polynesiense	Leibniz Institute DSMZ	DSM 27072
Staphylococcus aureus	American Type Culture Collection	ATCC 25923
Staphylococcus epidermidis	American Type Culture Collection	ATCC 12228
Virgibacillus massiliensis	Leibniz Institute DSMZ	DSM 28587
Enterocloster bolteae	Leibniz Institute DSMZ	DSM 15670
Escherichia coli	American Type Culture Collection	ATCC 25922
Escherichia coli	American Type Culture Collection	ATCC 35401
Pseudomonas aeruginosa	American Type Culture Collection	ATCC 27853
Acinetobacter baumannii	American Type Culture Collection	ATCC 19606
Klebsiella pneumoniae	American Type Culture Collection	ATCC 13883
Salmonella enteritidis	N/A	CMCC (B) 50335
Enterobacter cloacae	N/A	NRRL-B-413
Sphingomonas parapaucimobilis	Japan Collection of Microorganisms	JCM 7512
Micrococcus luteus	Leibniz Institute DSMZ	DSM 1790
Bifidobacterium longum subsp. *infantis*	Leibniz Institute DSMZ	DSM 20088
Bifidobacterium adolescentis	Leibniz Institute DSMZ	DSM 108263
Brevibacterium senegalense	Leibniz Institute DSMZ	DSM 25783
Rothia sp.	Leibniz Institute DSMZ	DSM 111809
Chryseobacterium gleum	Japan Collection of Microorganisms	JCM 2410
Mucilaginibacter defluvii	Japan Collection of Microorganisms	JCM 18283
Chemicals, peptides, and recombinant proteins		
Brain heart infusion medium (BHI)	hopebio	HB8297-1
Tryptic Soy Broth (TSB)	hopebio	HB4114
De Man–Rogosa–Sharpe medium (MRS)	BEIJING LAND BRIDGE	CM187
Peptone Yeast Glucose medium (modified) (mPYG)	TUOPU BIOTECHNOLOGY	MD090
Reinforced Clostridium Medium (RCM)	hopebio	HB0316

Gifu Anaerobic Broth (GAM)	TUOPU BIOTECHNOLOGY	M5650
Bifidobacterium Medium (BM)	hopebio	HB8527
M17 broth	Solarbio	LA0610
Deposited data		
Code for IIBacFinder	This study	https://github.com/ZhangDengwei/IIBacFinder
Metagenomic sequencing data of human feces	This study	PRJNA1201247
Software and algorithms		
Python 3.7.12	Python Software Foundation	https://www.python.org/
R version 4.2.2	The R Foundation	https://www.r-project.org/
Prodigal-short	Santos-Aberturas et al.[67]	https://github.com/streptomyces/ripper/tree/master
CD-HIT 4.8.1	Fu et al.[68]	https://github.com/weizhongli/cdhit
numpy		
pandas		
HMMer 3.1b2	Sean R. Eddy[69]	http://hmmer.org/
Ampir		
amPEPpy 1.0		
SAMtools 1.18		
Diamond v2.1.8.162	Buchfink et al.[70]	https://github.com/bbuchfink/diamond
MAFFT v7.508	Katoh et al.[71]	https://mafft.cbrc.jp/alignment/software/
trimAl v1.4.rev15	Capella-Gutiérrez et al.[72]	https://github.com/inab/trimal
amPEP v1.0	Lawrence et al.[73]	https://github.com/tlawrence3/amPEPpy
ampir 1.1.0	Fingerhut et al.[74]	https://github.com/Legana/ampir
NLPPrecursor	Merwin et al.[36]	https://github.com/magarveylab/NLPPrecursor
SignalP 6.0	Teufel et al.[37]	https://services.healthtech.dtu.dk/services/SignalP-6.0/
peptides.py	N/A	https://peptides.readthedocs.io/en/stable/index.html
BAGEL4	Teufel et al.[37]	https://github.com/annejong/BAGEL4
antiSMASH 6.1.1	Blin et al.[33]	https://github.com/antismash/antismash

MMseqs2 13.45111	Steinegger et al.[75]	https://github.com/soedinglab/MMseqs2
IQ-TREE version 2.2.0.3	Minh et al.[76]	http://www.iqtree.org/
GTDB-Tk v2.1.1	Chaumeil et al.[77]	https://github.com/Ecogenomics/GTDBTk
vegan 2.6.8	Jari Oksanen[78]	https://github.com/vegandevs/vegan
iNEXT 3.0.1	Hsieh et al.[79]	https://github.com/JohnsonHsieh/iNEXT
fastp v0.21.1	Chen et al.[80]	https://github.com/OpenGene/fastp
kneaddata	N/A	https://github.com/biobakery/kneaddata
PALADIN 1.4.6	Westbrook et al.[81]	https://github.com/ToniWestbrook/paladin
SAMtools 1.16.1	Li et al.[82]	https://www.htslib.org/
MetaPhlAn version 4.0.6	Blanco-Míguez et al.[83]	https://github.com/biobakery/MetaPhlAn
WGCNA 1.72.1	Langfelder et al.[84]	https://cran.r-project.org/web/packages/WGCNA/index.html
MaAsLin2 1.12.0	Mallick et al.[56]	https://github.com/biobakery/Maaslin2
ggplot2 v3.4.4	Villanueva et al.[85]	https://ggplot2.tidyverse.org/
Other		
UniRef100 release-2022_05	The UniProt Consortium[86]	https://www.uniprot.org/help/downloads
PFAM version 34.0	Mistry et al.[87]	http://ftp.ebi.ac.uk/pub/databases/Pfam/releases/Pfam34.0
NCBI version 11.0	NCBI Resource Coordinators[88]	https://ftp.ncbi.nlm.nih.gov/hmm/11.0/
GTDB repository release 207	Parks et al.[89]	https://gtdb.ecogenomic.org/downloads
GENCODE release 19 (GRCh37.p13)	Frankish et al.[90]	https://www.gencodegenes.org/human/

METHOD DETAILS

Development of IIBacFinder

Profile hidden Markov models (pHMMs) construction

A total of 286 non-redundant class II bacteriocin sequences (Table S1) were curated from previous studies. To comprehensively identify their homologous sequences, we used jackhammer[69] to iteratively align them against the UniProt UniRef100 database[86] (release-2022_05, comprising 323,519,324 sequences) with the parameters "-N 5 -T 20 --domT 20 --incT 20 --incdomT 20". Putative homologous sequences longer than 150 amino acids were considered false positives and discarded. We manually collated the homologous sequences, removing those with specific functions unrelated to antimicrobial activity or those with unknown functions, resulting in 297,986 remaining sequences. Combined with the 286 known bacteriocin sequences, these class II bacteriocin sequences were deduplicated to avoid significant similarity when constructing pHMMs using CD-HIT with the parameters "-n 2 -p 1 -c 0.8 -d 200 -M 5000 -aL 0.6 -g 1"[68]. The deduplicated sequences were then clustered at a 50% identity threshold using CD-HIT, resulting in 2,884 clusters with low sequence similarity among them. One cluster was removed due to containing a sequence with ambiguous amino acids. We used MAFFT[71] for multiple sequence alignment within each cluster. Following this, we used trimAl[72] to automatically remove poorly aligned regions with the model "-automated1". All aligned sequences were then used to build the pHMM for each cluster using hmmbuild[69]. This process resulted in 2,883 class II bacteriocin-related pHMMs. Additionally, we collected pHMMs from the PFAM[87] and NCBI databases[88], obtaining 16 and 14 pHMMs, respectively. To avoid redundancy between the 30 available pHMMs and the 2,883 self-constructed pHMMs, we used hmmscan[69] to annotate the seed sequences of the 2,883 clusters against the PFAM and NCBI databases. Clusters whose seed sequences were entirely annotated by either PFAM or NCBI domains were discarded, retaining a total of 1,123 clusters. Consequently, we obtained 1,153 precursor-related pHMMs (Table S3).

Subsequently, we curated 29 immunity protein sequences (Table S2) associated with class II bacteriocins from previous reports. To build the pHMMs for these immunity proteins, we first aligned the sequences against the UniRef100 database using Diamond[70] with the parameter "--id 40 --masking 0", retaining homologous sequences with identities greater than 40%. These homologous sequences were processed similarly to the precursor sequences, resulting in 29 pHMMs for class II bacteriocin immunity proteins. Additionally, we included eight publicly available pHMMs related to context genes, including protease genes, inducer genes, and accessory genes. Finally, we obtained 37 context gene-related pHMMs (Table S4).

Precursor gene identification

IIBacFinder takes a DNA sequence in FASTA format as input, which is subjected to open reading frame (ORF) annotation using Prodigal-short[67]. All ORFs are queried against a library of 1,153 precursor-related pHMMs and 37 context gene-related pHMMs using hmmsearch[69]. Putative precursor genes are detected using both precursor-based and context gene-based rules in parallel. For the precursor-based rule, significant hits identified by the 1,153 precursor-related pHMMs are initially considered putative precursor genes if the hit sequence is less than 150 amino acids in length. The region encompassing ±8 genes around the identified precursor genes is defined as the biosynthetic gene cluster (BGC). For the context gene-based rule, significant hits identified by the 37 context gene-related pHMMs are considered associated with class II bacteriocin biosynthesis. The eight genes upstream and downstream

of the identified context genes are included as the putative BGCs. BGC regions lacking a
transporter are discarded. Subsequently, the BGC region is examined for the presence of
proteins containing peptidase_C39 (PF03412) and LANC_like (PF05147) domains. BGCs
containing both domains are discarded as they are likely to be modified lantibiotics. All small
ORFs within the BGC, with lengths less than 150 AA, are retrieved and subjected to
antimicrobial peptide annotation using both amPEP[73] and ampir[74]. Small ORFs predicted as
antimicrobial peptides (AMPs) with a probability threshold of 0.8 by either tool are considered
putative precursor genes. The putative precursor genes detected by either the precursor-
based or context gene-based rules are then queried against the Pfam and NCBI databases
using hmmscan[69]. Sequences containing domains unrelated to class II bacteriocins are
removed (Table S5). The remaining sequences are regarded as putative precursors of class
II bacteriocins. Finally, BGC regions detected by both the precursor-based and context gene-
based rules are combined.

Leader type prediction

The leader type of precursor genes detected by the precursor-based rule is considered
identical to that of known homologous bacteriocins (Table S3). For precursor genes detected
by the context gene-based rule, we annotate the leader types based on the context genes
within the BGC region (Table S4). For example, if the BGC contains an ORF with a
peptidase_C39 domain (PF03412), the leader type of the precursor genes is considered to be
double-glycine.

Leader peptide prediction

To predict the cleavage site of the double-glycine leader, we tuned a deep learning model,
NLPPrecursor, specifically designed for the prediction of RiPP precursor cleavage sites[36]. The
protein sequences were tokenized using a vocabulary of 24 tokens, including a start and end
tag, twenty proteinogenic amino acids, an ambiguous amino acid, and a padding token. Unlike
RiPP precursors, which may contain leader and follower sequences at the N-terminal and C-
terminal, respectively, double-glycine bacteriocin precursors only have a leader sequence.
Therefore, we modified the sequence labels in the model. The allowed transitions of protein
sequences within the model were: (start-before), (before-before), (before-propeptide),
(propeptide-propeptide), (propeptide-stop), (stop-pad), and (pad-pad). All sequences were
unified to a final length of 150 by adding pad tokens at the end.

To train the deep learning model, we created a dataset of double-glycine class II bacteriocins.
Since only 142 known class II bacteriocin precursors contain a double-glycine leader, we
collected their homologous sequences from the UniProt UniRef100 database to expand the
training dataset. DIAMOND[91] was used to query the 142 sequences against the UniRef100
database with the parameters "--id 50 --query-cover 80 --masking 0", resulting in 1,224
homologs. To reduce sequence bias in training the prediction model, the 1,336 double-glycine
sequences were deduplicated at a sequence identity threshold of 90% using CD-HIT[68]. This
deduplication process yielded 439 non-redundant sequences. Subsequently, we used
MAFFT[71] with the parameters "--maxiterate 1000 --genafpair" to conduct multiple sequence
alignments and manually check whether the homologous sequences contained a conserved
double-glycine cleavage site. Finally, 412 class II bacteriocin precursor sequences with a
double-glycine leader were retained for model training with 100 epochs (Figure S1C).

For predicting the Sec-dependent leader, we adopted the well-established tool SignalP 6.0[37],
which predicts the signal peptides of proteins secreted by the secretory (Sec) pathway, using
default parameters.

Physicochemical property calculation

Following the prediction of the leader peptide, we compute common descriptors for the predicted core peptide using a pure-Python package, peptides.py (https://peptides.readthedocs.io/en/stable/index.html). The physicochemical properties include the aliphatic index, Boman index, isoelectric point, molecular weight, instability index, and net charge at a pH of 7.

Comparison with antimicrobial peptides (AMPs)

To gain insight into sequence novelty, predicted core peptides of class II bacteriocins were compared with known AMPs. AMP sequences were retrieved from four publicly available databases in April 2024: APD3[38], dbAMP[39], DRAMP[40], and DBAASP[41]. Sequences less than 10 amino acids or containing ambiguous amino acids were discarded, resulting in 51,692 AMP sequences, of which 21,403 were unique (100% identity). The predicted core peptides were queried against the AMP dataset using DIAMOND [41] with the parameters "--ultra-sensitive --max-target-seqs 1 --id 20 --query-cover 20 --masking 0". The best alignment hit was retained.

Prediction confidence assignment

Given that bioinformatic predictions can inevitably yield false positives, we added a confidence label to indicate whether the predicted class II bacteriocins are likely authentic or potentially false (Figure S2C). We considered several factors in determining this confidence level: the presence of a ribosome binding site (RBS), the completeness of the predicted ORF of the precursor gene, the detection rules (i.e., bacteriocin detection based on one rule or both rules), the leader peptide prediction, and length of the predicted core peptide. The decision tree in Figure S4 illustrates the classification into high-confidence or low-confidence categories. It is important to note that the inferred confidence level does not confirm whether the predicted bacteriocins are authentic or false without experimental validation; it only suggests a likelihood of the prediction's accuracy.

Performance benchmarking on identifying class II bacteriocins

To evaluate the performance of IIBacFinder, BAGEL4[41], and antiSMASH 6[33] in detecting class II bacteriocins, we applied these tools to the genomes of known bacteriocin producers. Since most producers' genomes have not been documented, we first attempted to locate known class II bacteriocins in sequenced bacterial genomes. The ORFs of 257,997 bacterial genomes from the RefSeq database were identified using Prodigal-short[67] with default parameters. These ORFs were then searched against 286 non-redundant known precursor sequences using MMseqs2 with the parameter "easy-search --max-seqs 1 --min-seq-id 0.95"[75]. A best hit with an identity > 95% and coverage > 95% was considered a known precursor sequence. Consequently, we obtained 137 genomes containing 205 complete and non-redundant precursor sequences of known class II bacteriocins.

We then applied the three prediction tools to these genomes using default parameters to assess their ability to identify class II bacteriocins. Notably, both IIBacFinder and BAGEL4 can pinpoint the precursor sequences, while antiSMASH 6 preferentially detects the biosynthetic region rather than the precursor sequences. Accordingly, if the precursor sequence identified by either IIBacFinder or BAGEL4 was within the BGC region detected by antiSMASH 6, we considered it detected by antiSMASH.

Phylogenetic tree construction

The multiple sequence alignment of 286 non-redundant known precursor sequences was achieved by MAFFT v7.508[71] with the parameter of "--maxiterate 1000 --localpair", which was further used to infer the phylogenetic tree. IQ-TREE version 2.2.0.3[76] was adapted for constructing maximum likelihood (ML) phylogenetic trees, with 1,000 ultrafast bootstrap replicates. The best-fit model was chosen as VT+F+I+I+R4 identified by the in-built ModelFinder[92]. The inferred phylogeny was visualized and annotated using iTOL[93].

Analysis of class II bacteriocins in bacterial genomes from multiple datasets

We used IIBacFinder to predict the biosynthetic potential of class II bacteriocins in bacterial genomes from eight datasets, consisting of the RefSeq database (RefSeq, n = 257,997), human microbiome (SGB, n = 126,702), the Unified Human Gastrointestinal Genome (UHGG, n = 285,835), the Human Reference Gut Microbiome (HRGM, n = 29,035), global earth microbiome (GEM, n = 49,466), ocean microbiome (Ocean, n = 21,592), mouse (MGBC, n = 26,640) and ruminant gut microbiome (RGM, n = 10,211). To confirm and unify the taxonomical annotation, GTDB-Tk v2.1.1[77] was adopted for taxonomical annotation against GTDB-Tk reference data version r207[89]. Genomes not belonging to bacteria were discarded.

Rarefaction analysis

To assess the bacteriocin biosynthetic potential of bacterial genomes from the RefSeq database, R package vegan[78] was used. CD-HIT[68] was used to group 481,654 high-confidence sequences from RefSeq genomes into 8,610 clusters at a 50% identity threshold The accumulations of bacteriocin clusters detected in Gram-positive and Gram-negative bacteria were computed using *specaccum* function with "random" method.

For the biosynthetic potential in Earth's microbiome, rarefaction analyses were conducted using the iNEXT R package[79]. We adopted CD-HIT[68] for deduplicating 10,581 high-confidence sequences from Earth's microbiome to 4,956 unique sequences at a 100% identity threshold. A unique sequence presence/absence table (sequence versus strain matrix) was constructed for each group considered. The type of input data was set as 'incidence-raw' data in the iNEXT main function. Besides, knots of 2,000 and an endpoint of 50,000 were adopted. The number of bootstrap replications is 50 by default.

Metagenomics and metatranscriptomics analysis

The raw metagenomic sequencing reads from five cohorts and the raw metatranscriptomic data from three cohorts were acquired from NCBI SRA (Sequence Read Archive) under respective project accession numbers. Only the sequencing data of healthy individuals was retained for analysis. Fastp v0.21.1[80] with default parameters was used to detect and remove low-quality sequencing reads. The sequencing reads belonging to the human host were detected and discarded by kneaddata (https://github.com/biobakery/kneaddata), through searching against the human reference genome (GRCh38.p13) from GENCODE[90]. The high-quality sequencing reads retained were subjected to downstream analysis.

Bacteriocin precursor profiling

Before profiling bacteriocin precursors in metagenomic and metatranscriptomic data, we constructed a reference dataset of class II bacteriocin precursor sequences. IIBacFinder identified 645,318 high-confidence precursor sequences in bacterial genomes from eight

databases. To remove duplicates and reduce computational load, we deduplicated these
sequences at a threshold of 95% identity and 95% coverage using CD-HIT[68], resulting in a
reference precursor dataset of 32,735 sequences. This reference dataset was then queried
by high-quality sequencing reads from metagenomic and metatranscriptomic data using
PALADIN software[81] to profile the abundance of bacteriocin precursors in each sample. We
used SAMtools[82] to calculate their abundance, normalizing the results using transcripts per
kilobase million (TPM). To reduce inaccuracies in abundance calculation caused by highly
similar sequences in the reference dataset, we grouped the sequences into clusters with a
threshold of 80% identity and 80% coverage using CD-HIT[68], and computed their abundance
at the cluster level rather than the individual sequence level.

Taxonomic profiling, community diversity, and correlation analysis

Metagenomic taxonomic profiling was conducted using MetaPhlAn version 4.0.6[83]. The
relative abundance of species was used to represent the abundance of bacterial species in
the community. The alpha diversity (Shannon index) of each sample was calculated with the
R package vegan v2.6.4[78]. Given that bioactive bacteriocins could inhibit the growth of other
bacteria and thus regulate the overall bacterial community, we computed the Spearman
correlation between taxonomic composition (at the species level) or Shannon diversity and
precursor clusters using the corAndPvalue function from the R package WGCNA[84]. The p-
values were then adjusted using the "BH" method to control the false discovery rate in multiple
comparisons[94]. Considering that bioactive bacteriocins would inhibit the growth of specific
bacterial species, we retained only the significantly negative associations (*rho* < -0.3 and
adjusted *P* < 0.05).

Bacteriocin precursor prioritization in metagenomic and metatranscriptomic data

700 To infer the importance of class II bacteriocins in metagenomic and metatranscriptomic data,
701 we contrived a scheme to assign ranking scores for each profiled precursor clusters, through
702 considering the ecological importance (prevalence and abundance of bacteriocins) and
703 functional importance (association between bacterial species and bacteriocins and between
704 Shannon diversity and bacteriocins). Mathematically, the ecological importance is scored as:

705
$$RS\_eco = \frac{1}{\frac{1}{R_p} \times 0.5 + \frac{1}{R_a} \times 0.5}$$

706 Where RS_eco indicates the ranking score (ecological importance) of a particular bacteriocin
707 within a cohort. R_p and R_a represent the ranking percentiles based on prevalence and mean
708 abundance (largest to smallest), respectively. A bacteriocin with high prevalence and high
709 abundance would give rise to a larger RS_eco.

710 The functional importance is mathematically scored as:

711
$$RS\_fun = \frac{RS_{cor} + RS_{reg}}{2}$$

Where RS_fun is the ranking score (functional importance) of a particular bacteriocin within a
cohort. RS_{cor} indicates the ranking percentiles based on the negative correlation between
bacteriocins and bacterial species, while RS_{reg} is the ranking percentiles based on the
correlation between bacteriocins and the Shannon diversity index.

The overall ranking score of a bacteriocin in metagenomic and metatranscriptomic data is the
average ranking score of ecological and functional importance in all cohorts, mathematically
represented as:

719
$$RS = \frac{\sum_{i=1}^{n} RS\_eco + \sum_{i=1}^{n} RS\_fun}{n}$$

Where n represents the number of metagenomic and metatranscriptomic datasets. A high
score of RS implies the great importance of the bacteriocin in the gut microbiome.

After prioritizing the bacteriocins in the human gut microbiome, we selected bacteriocin
clusters to validate their antibacterial activities based on the ranking scores (RS) and the
following criteria: (1) The hypothetical bacteriocin cluster should include ≥ 5 precursor
sequences, as sporadically identified bacteriocins are likely to be false predictions. (2) The
cleavage site for double-glycine bacteriocins should be a typical "GG" rather than alternative
sites such as "GA" or "GS" to reduce false positives. (3) The length of the deduced core
peptide should be less than 50 amino acids due to cost considerations and the feasibility of
chemical synthesis. (4) The bacteriocin sequences should not be annotated as peptides with
functions other than antimicrobials when queried against the NCBI database. Based on these
criteria, we chose to synthesize and test the representative sequences, assigned by CD-HIT[68],
from 40 bacteriocin clusters.

Three-dimensional peptide structure prediction

734 ColabFold was used to predict the three-dimensional structure of peptides, with a parameter
735 of "--amber --templates --num-recycle 3 --use-gpu-relax --model-type auto"[95]. The model with
736 the highest relaxed rank was retained and visualized using PyMol [96].

Peptide synthesis

738 The deducted core peptides of predicted bacteriocins were chemically synthesized by solid-
739 phase peptide synthesis by Sangon Biotech (Shanghai, China). Their molecular weights were
740 confirmed by mass spectrometry, and their required purity was greater than 90% determined
741 by high-performance liquid chromatography. The synthesized peptide powder was stored
742 at − 80 °C and dissolved in Dimethyl sulfoxide (DMSO) to 50 mg/mL upon use.

Microbial strains and culturing

A total of 33 bacterial strains are included in this study and are detailed in Table S10 regarding
the strains and growth conditions. The following media were used: Brain heart infusion
medium (BHI); Tryptic Soy Broth with supplement of 0.3% yeast extract (TSB-YE); De Man–
Rogosa–Sharpe medium (MRS); Peptone Yeast Glucose medium (PYG); Reinforced
Clostridium Medium (RCM); Gifu Anaerobic Broth (GAM); Bifidobacterium Medium (BM); M17
broth. Bacterial strains were grown and streaked on the agar plate, and incubated under
respective optimal growth conditions. Individual colonies were then inoculated into 5 mL
corresponding culture broth. Once the bacterial culture reached sufficient density, 500 μL
culture bacterial suspension was supplemented with 500 μL 50% glycerol to generate liquid
glycerol stocks and stored at −80 °C.

Antibacterial activity screening

The bacterial inhibition experiment was adopted from a previous study[26]. All bacterial
indicators were revived on agar medium under respective growth conditions (Table S10). The

individual colonies were picked into broth medium and incubated for one or two days. The
bacterial suspension was diluted to a particular concentration [optical density at 600 nm (OD$_{600}$)
= 0.1] and then diluted 100 or 1,000 times for the antibacterial activity screening (Table S10).
Three experimental groups were set: (1) blank control group, 100 µL of broth culture; (2)
bacterial control group, 50 µL of broth culture and 50 µL of bacterial solution with DMSO
(equivalent amount to the solvent of peptides); (3) experimental group, 50 µL of broth culture
and 50 µL of bacterial solution with 100 µg/mL peptide. Experiments were conducted on flat
bottom 96-well plates. The bacterial growth was measured with OD$_{600}$ value after statically
incubating for 24h or 48h. The inhibition ratio was calculated as:

766
$$inhibition\ ratio\ = \frac{OD_{experiment} - OD_{blank}}{OD_{control} - OD_{blank}}$$

Where OD$_{blank}$, OD$_{control}$, and OD$_{experiment}$ represent the OD$_{600}$ of the blank control group,
bacterial control group, and experimental group, respectively.

All experiments were conducted with three independent replicates. We performed Dunnett's
test for computing the statistical difference between the control and experiment groups. The
peptide with a criterion of inhibition ratio ≥ 25% and P value < 0.05 was considered effective
against the tested indicator strain. We categorized the activity results into four groups: (1) no
activity, inhibition ratio < 25% or P value ≥ 0.05; (2) low activity, 25% ≤ inhibition ratio < 50%
and P value < 0.05; (3) medium activity, 50% ≤ inhibition ratio < 75% and P value < 0.05; (4)
high activity, inhibition ratio ≥ 75% and P value < 0.05. Peptides with an inhibition ratio > 90%
were further examined the MIC against this indicator strain.

Determination of minimum inhibitory concentrations (MICs)

The minimum inhibitory concentrations (MICs) of synthetic peptides were performed by broth
microdilution in 96-well flat bottom plates as previously reported[97]. Vancomycin was used as
a positive control. Peptides were added to the plate containing 100 µL aliquots of bacterial
suspensions and were two-fold serially diluted (ranging from 100 µg/mL to 0.195 µg/mL). After
incubating for 24h or 48h, bacterial growth was assessed by determining OD$_{600}$. The MIC
value was defined as the lowest concentration of the peptides where no bacterial or fungal
growth was detected. The assays were conducted in triplicate.

Human feces collection

Human studies were performed in accordance with all relevant ethical regulations and were
approved by the Medical Ethics Committee of ZhuJiang Hospital, Southern Medical University
(2022-KY-119). Fecal samples were collected from five healthy Chinese adults aged 20 to 42
who had not taken antibiotics or probiotics within one month before sample collection.

Ex vivo assay was conducted as previously described[17]. Briefly, the collected feces were
combined and homogenized in 65 mL of a rich medium mGAM. Following resuspension, the
tubes were gently centrifuged at 1,000 rpm for 2 minutes, and the supernatants were then
retained for further assay. Four selected bacteriocin peptides were pre-dissolved in DMSO to
final concentrations of 0.1 mg/mL, and 10 mg/mL. Subsequently, 10 µL of peptides and 40 µL
of fecal supernatant were inoculated into 950 µL of mGAM media, resulting in a final volume
of 1 mL. The mixtures were then anaerobically incubated at 37°C for 48 hours. Following the
incubation period, the bacterial growth media were centrifuged at 12,000 rpm for 10 minutes.
The supernatants were carefully removed, and the resulting pellets were washed twice with 1
mL of PBS before being collected for DNA extraction. The pelleted samples were extracted

using the QIAamp® DNA Micro Kit (Qiagen) according to the manufacturer's instructions.
Finally, all DNA samples were prepared for shotgun metagenomics sequencing.

Data analysis of human fecal metagenome

All DNA samples were submitted to Novogene for shotgun metagenomics sequencing using
a 150bp paired-end protocol as per the manufacturer's instructions. The raw sequencing data
process was described above to obtain the taxonomic profiles. The Vegan package[78] in R was
used to compute alpha diversity and beta diversity at the species level, indicated by Shannon
diversity and Bray-Curtis dissimilarity respectively. Differential abundance analysis between
groups was conducted using MaAsLin2 (microbiome multivariable associations with linear
models) [56]. Species with a false discovery rate (FDR)-adjusted p-value of less than 0.05 were
deemed significantly different between the two groups.

Visualization

The intersection of identified bacteriocins from three tools was visualized using the R package
UpSetR v1.4.0[98]. The sankey plot showing the biosynthetic potential of bacteriocins in the
human gut microbiome was achieved by the package ggsankey v0.0.99999[99]. The heat maps
were plotted using the package ComplexHeatmap v2.14.0[100]. The similarity network was
visualized using Cytoscape 3.9.0[101]. The organization of gene clusters was visualized by R
package gggenes v0.5.0[102]. Without a specific statement, other figures were generated using
ggplot2 v3.4.4[85].

QUANTIFICATION AND STATISTICAL ANALYSIS

820 All statistical analyses were finished in R v4.2.3. Dunnett's test (two sided) was done by the
821 function *DunnettTest* in R package DescTools v0.99.49[103].

References

1. Ghoul, M., and Mitri, S. (2016). The Ecology and Evolution of Microbial Competition. Trends Microbiol. *24*, 833–845. https://doi.org/10.1016/j.tim.2016.06.011.

2. García-Bayona, L., and Comstock, L.E. (2018). Bacterial antagonism in host-associated microbial communities. Science (80-.). *361*. https://doi.org/10.1126/science.aat2456.

3. Palmer, J.D., and Foster, K.R. (2022). Bacterial species rarely work together. Science (80-.). *376*, 581–582. https://doi.org/10.1126/science.abn5093.

4. Özçam, M., Oh, J.H., Tocmo, R., Acharya, D., Zhang, S., Astmann, T.J., Heggen, M., Ruiz-Ramírez, S., Li, F., Cheng, C.C., et al. (2022). A secondary metabolite drives intraspecies antagonism in a gut symbiont that is inhibited by cell-wall acetylation. Cell Host Microbe *30*, 824-835.e6. https://doi.org/10.1016/j.chom.2022.03.033.

5. Roelofs, K.G., Coyne, M.J., Gentyala, R.R., Chatzidaki-Livanis, M., and Comstock, L.E. (2016). Bacteroidales Secreted Antimicrobial Proteins Target Surface Molecules Necessary for Gut Colonization and Mediate Competition In Vivo. MBio *7*, 1–10. https://doi.org/10.1128/mBio.01055-16.

6. Piewngam, P., Zheng, Y., Nguyen, T.H., Dickey, S.W., Joo, H.S., Villaruz, A.E., Glose, K.A., Fisher, E.L., Hunt, R.L., Li, B., et al. (2018). Pathogen elimination by probiotic Bacillus via signalling interference. Nature *562*, 532–537. https://doi.org/10.1038/s41586-018-0616-y.

7. Cotter, P.D., Ross, R.P., and Hill, C. (2013). Bacteriocins-a viable alternative to antibiotics? Nat. Rev. Microbiol. *11*, 95–105. https://doi.org/10.1038/nrmicro2937.

8. Heilbronner, S., Krismer, B., Brötz-Oesterhelt, H., and Peschel, A. (2021). The microbiome-shaping roles of bacteriocins at Nature Publishing Group, https://doi.org/10.1038/s41579-021-00569-w https://doi.org/10.1038/s41579-021-00569-w.

9. Sugrue, I., Ross, R.P., and Hill, C. (2024). Bacteriocin diversity, function, discovery and application as antimicrobials. Nat. Rev. Microbiol. https://doi.org/10.1038/s41579-024-01045-x.

10. Acedo, J.Z., Chiorean, S., Vederas, J.C., and van Belkum, M.J. (2018). The expanding structural variety among bacteriocins from Gram-positive bacteria at Oxford University Press, https://doi.org/10.1093/femsre/fuy033 https://doi.org/10.1093/femsre/fuy033.

11. Sharma, A., and Srivastava, S. (2014). Anti-Candida activity of two-peptide bacteriocins, plantaricins (Pln E/F and J/K) and their mode of action. Fungal Biol. *118*, 264–275. https://doi.org/10.1016/j.funbio.2013.12.006.

12. Teiar, R., Sane, F., Erol, I., Nekoua, M.P., Lecouturier, D., Boukherroub, R., Durdağı, S., Hober, D., and Drider, D. (2024). Enterocin DD14 can inhibit the infection of eukaryotic cells with enveloped viruses. Arch. Microbiol. *206*, 269. https://doi.org/10.1007/s00203-024-04002-7.

13. Nazari, M., Yaghoubian, I., and Smith, D.L. (2022). The stimulatory effect of Thuricin 17, a PGPR-produced bacteriocin, on canola (Brassica, napus L.) germination and vegetative growth under stressful temperatures. Front. Plant Sci. *13*, 1–17. https://doi.org/10.3389/fpls.2022.1079180.

14. Hu, J., Ma, L., Nie, Y., Chen, J., Zheng, W., Wang, X., Xie, C., Zheng, Z., Wang, Z., Yang, T., et al. (2018). A Microbiota-Derived Bacteriocin Targets the Host to Confer Diarrhea Resistance in Early-Weaned Piglets. Cell Host Microbe *24*, 817-832.e8.

https://doi.org/10.1016/j.chom.2018.11.006.

15. Heeney, D.D., Zhai, Z., Bendiks, Z., Barouei, J., Martinic, A., Slupsky, C., and Marco,
M.L. (2019). Lactobacillus plantarum bacteriocin is associated with intestinal and
systemic improvements in diet-induced obese mice and maintains epithelial barrier
integrity in vitro. Gut Microbes *10*, 382–397.
https://doi.org/10.1080/19490976.2018.1534513.

16. Zhang, D., Zhang, J., Kalimuthu, S., Liu, J., Song, Z., He, B., Cai, P., Zhong, Z., Feng,
C., Neelakantan, P., et al. (2023). A systematically biosynthetic investigation of lactic
acid bacteria reveals diverse antagonistic bacteriocins that potentially shape the human
microbiome. Microbiome *11*, 91. https://doi.org/10.1186/s40168-023-01540-y.

17. Zhang, J., Zhang, D., Xu, Y., Zhang, J., Liu, R., Gao, Y., Shi, Y., Cai, P., Zhong, Z., He,
B., et al. (2024). Elucidating the Diversity and Potential Function of Ribosomally
Synthesized and Post-translationally Modified Peptides in the Human Microbiome.
bioRxiv, 107–125. https://doi.org/10.1101/2024.07.16.603675.

18. Garcia-Gutierrez, E., Mayer, M.J., Cotter, P.D., and Narbad, A. (2019). Gut microbiota
as a source of novel antimicrobials. Gut Microbes *10*, 1–21.
https://doi.org/10.1080/19490976.2018.1455790.

19. Gu, Q., Yan, J., Lou, Y., Zhang, Z., Li, Y., Zhu, Z., Liu, M., Wu, D., Liang, Y., Pu, J., et
al. (2024). Bacteriocins: Curial guardians of gastrointestinal tract. Compr. Rev. Food
Sci. Food Saf. *23*, 1–27. https://doi.org/10.1111/1541-4337.13292.

20. Javvadi, S.G., Kujawska, M., Papp, D., Jordan, A., Lawson, M.A.E., Clarke, P., Beraza,
N., Hall, L.J., Microbes, G., Microbiome, I., et al. (2022). A novel bacteriocin produced
by Bifidobacterium longum subsp. infantis has dual antimicrobial and
immunomodulatory activity. bioRxiv. https://doi.org/10.1101/2022.01.27.477972.

21. Zhu, B., Wang, X., and Li, L. (2010). Human gut microbiome: The second genome of
human body. Protein Cell *1*, 718–725. https://doi.org/10.1007/s13238-010-0093-z.

22. Walsh, C.J., Guinane, C.M., Hill, C., Ross, R.P., O'Toole, P.W., and Cotter, P.D. (2015).
In silico identification of bacteriocin gene clusters in the gastrointestinal tract, based on
the Human Microbiome Project's reference genome database. BMC Microbiol. *15*.
https://doi.org/10.1186/s12866-015-0515-4.

23. Zheng, J., Gänzle, M.G., Lin, X.B., Ruan, L., and Sun, M. (2015). Diversity and
dynamics of bacteriocins from human microbiome. Environ. Microbiol. *17*, 2133–2143.
https://doi.org/10.1111/1462-2920.12662.

24. Almeida, A., Nayfach, S., Boland, M., Strozzi, F., Beracochea, M., Shi, Z.J., Pollard,
K.S., Sakharova, E., Parks, D.H., Hugenholtz, P., et al. (2021). A unified catalog of
204,938 reference genomes from the human gut microbiome. Nat. Biotechnol. *39*, 105–
114. https://doi.org/10.1038/s41587-020-0603-3.

25. Santos-Júnior, C.D., Torres, M.D.T., Duan, Y., Rodríguez del Río, Á., Schmidt, T.S.B.,
Chong, H., Fullam, A., Kuhn, M., Zhu, C., Houseman, A., et al. (2024). Discovery of
antimicrobial peptides in the global microbiome with machine learning. Cell, 1–18.
https://doi.org/10.1016/j.cell.2024.05.013.

26. Ma, Y., Guo, Z., Xia, B., Zhang, Y., Liu, X., Yu, Y., Tang, N., Tong, X., Wang, M., Ye,
X., et al. (2022). Identification of antimicrobial peptides from the human gut microbiome
using deep learning. Nat. Biotechnol. *40*, 921–931. https://doi.org/10.1038/s41587-022-
01226-0.

27. Torres, M.D.T., Brooks, E.F., Cesaro, A., Sberro, H., Gill, M.O., Nicolaou, C., Bhatt,

A.S., and de la Fuente-Nunez, C. (2024). Mining human microbiomes reveals an
untapped source of peptide antibiotics. Cell *187*, 5453-5467.e15.
https://doi.org/10.1016/j.cell.2024.07.027.

28.   Cui, Z., Chen, Z.H., Zhang, Q.H., Gribova, V., Filaretov, V.F., and Huang, D.S. (2022).
919   RMSCNN: A Random Multi-Scale Convolutional Neural Network for Marine Microbial
920   Bacteriocins Identification. IEEE/ACM Trans. Comput. Biol. Bioinforma. *19*, 3663–3672.
921   https://doi.org/10.1109/TCBB.2021.3122183.

922   29.   Akhter, S., and Miller, J.H. (2023). BaPreS: a software tool for predicting bacteriocins
923   using an optimal set of features. BMC Bioinformatics *24*, 313.
924   https://doi.org/10.1186/s12859-023-05330-z.

925   30.   Akhter, S., and Miller, J.H. (2024). BPAGS: a web application for bacteriocin prediction
926   via feature evaluation using alternating decision tree, genetic algorithm, and linear
927   support vector classifier. Front. Bioinforma. *3*, 1–11.
928   https://doi.org/10.3389/fbinf.2023.1284705.

929   31.   van Heel, A.J., de Jong, A., Song, C., Viel, J.H., Kok, J., and Kuipers, O.P. (2018).
930   BAGEL4: a user-friendly web server to thoroughly mine RiPPs and bacteriocins.
931   Nucleic Acids Res. *46*, W278–W281. https://doi.org/10.1093/nar/gky383.

932   32.   Skinnider, M.A., Johnston, C.W., Gunabalasingam, M., Merwin, N.J., Kieliszek, A.M.,
933   MacLellan, R.J., Li, H., Ranieri, M.R.M., Webster, A.L.H., Cao, M.P.T., et al. (2020).
934   Comprehensive prediction of secondary metabolite structure and biological activity from
935   microbial genome sequences. Nat. Commun. *11*, 6058. https://doi.org/10.1038/s41467-
936   020-19986-1.

937   33.   Blin, K., Shaw, S., Kloosterman, A.M., Charlop-Powers, Z., van Wezel, G.P., Medema,
938   M.H., and Weber, T. (2021). antiSMASH 6.0: improving cluster detection and
939   comparison capabilities. Nucleic Acids Res. *49*, W29–W35.
940   https://doi.org/10.1093/nar/gkab335.

941   34.   Baldi, P., Chauvin, Y., Hunkapiller, T., and Mcclure, M.A. (1994). Hidden Markov
942   models of biological primary sequence information. Proc. Natl. Acad. Sci. U. S. A. *91*,
943   1059–1063. https://doi.org/10.1073/pnas.91.3.1059.

944   35.   Eddy, S.R. (1998). Profile hidden Markov models. Bioinformatics *14*, 755–763.
945   https://doi.org/10.1093/bioinformatics/14.9.755.

946   36.   Merwin, N.J., Mousa, W.K., Dejong, C.A., Skinnider, M.A., Cannon, M.J., Li, H., Dial,
947   K., Gunabalasingam, M., Johnston, C., and Magarvey, N.A. (2020). DeepRiPP
948   integrates multiomics data to automate discovery of novel ribosomally synthesized
949   natural products. Proc. Natl. Acad. Sci. *117*, 371–380.
950   https://doi.org/10.1073/pnas.1901493116.

951   37.   Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gíslason, M.H., Pihl, S.I.,
952   Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022). SignalP
953   6.0 predicts all five types of signal peptides using protein language models. Nat.
954   Biotechnol. *40*, 1023–1025. https://doi.org/10.1038/s41587-021-01156-3.

955   38.   Wang, G., Li, X., and Wang, Z. (2016). APD3: The antimicrobial peptide database as a
956   tool for research and education. Nucleic Acids Res. *44*, D1087–D1093.
957   https://doi.org/10.1093/nar/gkv1278.

958   39.   Jhong, J.H., Yao, L., Pang, Y., Li, Z., Chung, C.R., Wang, R., Li, S., Li, W., Luo, M., Ma,
959   R., et al. (2022). dbAMP 2.0: Updated resource for antimicrobial peptides with an
960   enhanced scanning method for genomic and proteomic data. Nucleic Acids Res. *50*,
961   D460–D470. https://doi.org/10.1093/nar/gkab1080.

40.   Shi, G., Kang, X., Dong, F., Liu, Y., Zhu, N., Hu, Y., Xu, H., Lao, X., and Zheng, H.
(2022). DRAMP 3.0: An enhanced comprehensive data repository of antimicrobial
peptides. Nucleic Acids Res. *50*, D488–D496. https://doi.org/10.1093/nar/gkab651.

41.   Pirtskhalava, M., Amstrong, A.A., Grigolava, M., Chubinidze, M., Alimbarashvili, E.,
Vishnepolsky, B., Gabrielian, A., Rosenthal, A., Hurt, D.E., and Tartakovsky, M. (2021).
DBAASP v3: Database of antimicrobial/cytotoxic activity and structure of peptides as a
resource for development of new therapeutics. Nucleic Acids Res. *49*, D288–D297.
https://doi.org/10.1093/nar/gkaa991.

42.   Haft, D.H., DiCuccio, M., Badretdin, A., Brover, V., Chetvernin, V., O'Neill, K., Li, W.,
Chitsaz, F., Derbyshire, M.K., Gonzales, N.R., et al. (2018). RefSeq: An update on
prokaryotic genome annotation and curation. Nucleic Acids Res. *46*, D851–D860.
https://doi.org/10.1093/nar/gkx1068.

43.   Arbulu, S., and Kjos, M. (2024). Revisiting the Multifaceted Roles of Bacteriocins: The
Multifaceted Roles of Bacteriocins. Microb. Ecol. *87*. https://doi.org/10.1007/s00248-
024-02357-4.

44.   Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu, D.,
Paez-Espino, D., Chen, I.-M., Huntemann, M., et al. (2021). A genomic catalog of
Earth's microbiomes. Nat. Biotechnol. *39*, 499–509. https://doi.org/10.1038/s41587-
020-0718-6.

45.   Fan, Y., and Pedersen, O. (2021). Gut microbiota in human metabolic health and
disease. Nat. Rev. Microbiol. *19*, 55–71. https://doi.org/10.1038/s41579-020-0433-9.

46.   Gamier, T., and Cole, S.T. (1988). Studies of UV-inducible promoters from Clostridium
perfringens in vivo and in vitro. Mol. Microbiol. *2*, 607–614.
https://doi.org/10.1111/j.1365-2958.1988.tb00069.x.

47.   Corr, S.C., Li, Y., Riedel, C.U., O'Toole, P.W., Hill, C., and Gahan, C.G.M. (2007).
Bacteriocin production as a mechanism for the antiinfective activity of Lactobacillus
salivarius UCC118. Proc. Natl. Acad. Sci. U. S. A. *104*, 7617–7621.
https://doi.org/10.1073/pnas.0700440104.

48.   Li, J., Jin, J., Li, S., Zhong, Y., Jin, Y., Zhang, X., Xia, B., Zhu, Y., Guo, R., Sun, X., et
al. (2022). Tonsillar Microbiome-Derived Lantibiotics Induce Structural Changes of IL-6
and IL-21 Receptors and Modulate Host Immunity. Adv. Sci. *9*, 2202706.
https://doi.org/10.1002/advs.202202706.

49.   Lazzaro, B.P., Zasloff, M., and Rolff, J. (2020). Antimicrobial peptides: Application
informed by evolution at American Association for the Advancement of Science,
https://doi.org/10.1126/science.aau5480 https://doi.org/10.1126/science.aau5480.

50.   Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F.,
Manghi, P., Tett, A., Ghensi, P., et al. (2019). Extensive Unexplored Human Microbiome
Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning Age,
Geography, and Lifestyle. Cell *176*, 649-662.e20.
https://doi.org/10.1016/j.cell.2019.01.001.

51.   Kim, C.Y., Lee, M., Yang, S., Kim, K., Yong, D., Kim, H.R., and Lee, I. (2021). Human
reference gut microbiome catalog including newly assembled genomes from under-
represented Asian metagenomes. Genome Med. *13*. https://doi.org/10.1186/s13073-
021-00950-7.

52.   Paoli, L., Ruscheweyh, H.-J., Forneris, C.C., Hubrich, F., Kautsar, S., Bhushan, A., Lotti,
A., Clayssen, Q., Salazar, G., Milanese, A., et al. (2022). Biosynthetic potential of the
global ocean microbiome. Nature *607*, 111–118. https://doi.org/10.1038/s41586-022-

04862-3.

53.   Beresford-Jones, B.S., Forster, S.C., Stares, M.D., Notley, G., Viciani, E., Browne, H.P., Boehmler, D.J., Soderholm, A.T., Kumar, N., Vervier, K., et al. (2022). The Mouse Gastrointestinal Bacteria Catalogue enables translation between the mouse and human gut microbiotas via functional mapping. Cell Host Microbe *30*, 124-138.e8. https://doi.org/10.1016/j.chom.2021.12.003.

54.   Xie, F., Jin, W., Si, H., Yuan, Y., Tao, Y., Liu, J., Wang, X., Yang, C., Li, Q., Yan, X., et al. (2021). An integrated gene catalog and over 10,000 metagenome-assembled genomes from the gastrointestinal microbiome of ruminants. Microbiome *9*, 1–20. https://doi.org/10.1186/s40168-021-01078-x.

55.   Coyne, M.J., Béchon, N., Matano, L.M., McEneany, V.L., Chatzidaki-Livanis, M., and Comstock, L.E. (2019). A family of anti-Bacteroidales peptide toxins wide-spread in the human gut microbiota. Nat. Commun. *10*. https://doi.org/10.1038/s41467-019-11494-1.

56.   Mallick, H., Rahnavard, A., McIver, L.J., Ma, S., Zhang, Y., Nguyen, L.H., Tickle, T.L., Weingart, G., Ren, B., Schwager, E.H., et al. (2021). Multivariable association discovery in population-scale meta-omics studies. PLoS Comput. Biol. *17*. https://doi.org/10.1371/journal.pcbi.1009442.

57.   de Nies, L., Kobras, C.M., and Stracy, M. (2023). Antibiotic-induced collateral damage to the microbiota and associated infections. Nat. Rev. Microbiol. https://doi.org/10.1038/s41579-023-00936-9.

58.   Ankaiah, D., Esakkiraj, P., Perumal, V., Ayyanna, R., and Venkatesan, A. (2017). Probiotic characterization of Enterococcus faecium por1: Cloning, over expression of Enterocin-A and evaluation of antibacterial, anti-cancer properties. J. Funct. Foods *38*, 280–292. https://doi.org/10.1016/j.jff.2017.09.034.

59.   Pandi, A., Adam, D., Zare, A., Trinh, V.T., Schaefer, S.L., Burt, M., Klabunde, B., Bobkova, E., Kushwaha, M., Foroughijabbari, Y., et al. (2023). Cell-free biosynthesis combined with deep learning accelerates de novo-development of antimicrobial peptides. Nat. Commun. *14*, 7197. https://doi.org/10.1038/s41467-023-42434-9.

60.   Ayikpoe, R.S., Shi, C., Battiste, A.J., Eslami, S.M., Ramesh, S., Simon, M.A., Bothwell, I.R., Lee, H., Rice, A.J., Ren, H., et al. (2022). A scalable platform to discover antimicrobials of ribosomal origin. Nat. Commun. *13*, 1–15. https://doi.org/10.1038/s41467-022-33890-w.

61.   Liu, W.-Q., Ji, X., Ba, F., Zhang, Y., Xu, H., Huang, S., Zheng, X., Liu, Y., Ling, S., Jewett, M.C., et al. (2024). Cell-free biosynthesis and engineering of ribosomally synthesized lanthipeptides. Nat. Commun. *15*, 4336. https://doi.org/10.1038/s41467-024-48726-y.

62.   Yan, Q., Li, S., Yan, Q., Huo, X., Wang, C., Wang, X., Sun, Y., Zhao, W., Yu, Z., Zhang, Y., et al. (2024). A genomic compendium of cultivated human gut fungi characterizes the gut mycobiome and its relevance to common diseases. Cell, 1–21. https://doi.org/10.1016/j.cell.2024.04.043.

63.   Teiar, R., Pérez-Ramos, A., Zgheib, H., Cudennec, B., Belguesmia, Y., and Drider, D. (2022). Anti-adhesion and Anti-inflammatory Potential of the Leaderless Class IIb Bacteriocin Enterocin DD14. Probiotics Antimicrob. Proteins *14*, 613–619. https://doi.org/10.1007/s12602-022-09954-0.

64.   Śmiałek, J., Mak, P., Bzowska, M., Hinz, A., Mężyk-Kopeć, R., and Sołtys, K. (2022). Bacteriocin BacSp222 and Its Succinylated Forms Exhibit Proinflammatory Activities Toward Innate Immune Cells. J. Inflamm. Res. *15*, 4601–4621.

1056   https://doi.org/10.2147/JIR.S362066.

1057   65.   Antoshina, D. V., Balandin, S. V., Bogdanov, I. V., Vershinina, M.A., Sheremeteva, E.
1058   V., Toropygin, I.Y., Finkina, E.I., and Ovchinnikova, T. V. (2022). Antimicrobial Activity
1059   and Immunomodulatory Properties of Acidocin A, the Pediocin-like Bacteriocin with the
1060   Non-Canonical   Structure.   Membranes   (Basel).   *12*.
1061   https://doi.org/10.3390/membranes12121253.

1062   66.   Venema, K., Abee, T., Haandrikman, A.J., Leenhouts, K.J., Kok, J., Konings, W.N., and
1063   Venema, G. (1993). Mode of Action of Lactococcin B, a Thiol-Activated Bacteriocin from
1064   Lactococcus   lactis.   Appl.   Environ.   Microbiol.   *59,*   1041–1048.
1065   https://doi.org/10.1128/aem.59.4.1041-1048.1993.

1066   67.   Santos-Aberturas, J., Chandra, G., Frattaruolo, L., Lacret, R., Pham, T.H., Vior, N.M.,
1067   Eyles, T.H., and Truman, A.W. (2019). Uncovering the unexplored diversity of
1068   thioamidated ribosomal peptides in Actinobacteria using the RiPPER genome mining
1069   tool. Nucleic Acids Res. *47,* 4624–4637. https://doi.org/10.1093/nar/gkz192.

1070   68.   Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: Accelerated for clustering
1071   the   next-generation   sequencing   data.   Bioinformatics   *28,*   3150–3152.
1072   https://doi.org/10.1093/bioinformatics/bts565.

1073   69.   Eddy, S.R. (2011). Accelerated profile HMM searches. PLoS Comput. Biol. *7,*
1074   e1002195. https://doi.org/10.1371/journal.pcbi.1002195.

1075   70.   Buchfink, B., Xie, C., and Huson, D.H. (2014). Fast and sensitive protein alignment
1076   using DIAMOND. Nat. Methods *12,* 59–60. https://doi.org/10.1038/nmeth.3176.

1077   71.   Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software
1078   Version 7: Improvements in Performance and Usability. Mol. Biol. Evol. *30,* 772–780.
1079   https://doi.org/10.1093/molbev/mst010.

1080   72.   Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: A tool for
1081   automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics *25,*
1082   1972–1973. https://doi.org/10.1093/bioinformatics/btp348.

1083   73.   Lawrence, T.J., Carper, D.L., Spangler, M.K., Carrell, A.A., Rush, T.A., Minter, S.J.,
1084   Weston, D.J., and Labbé, J.L. (2021). amPEPpy 1.0: a portable and accurate
1085   antimicrobial   peptide   prediction   tool.   Bioinformatics   *37,*   2058–2060.
1086   https://doi.org/10.1093/bioinformatics/btaa917.

1087   74.   Fingerhut, L.C.H.W., Miller, D.J., Strugnell, J.M., Daly, N.L., and Cooke, I.R. (2021).
1088   ampir: an R package for fast genome-wide prediction of antimicrobial peptides.
1089   Bioinformatics *36,* 5262–5263. https://doi.org/10.1093/bioinformatics/btaa653.

1090   75.   Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein sequence
1091   searching for the analysis of massive data sets at Nature Publishing Group,
1092   https://doi.org/10.1038/nbt.3988 https://doi.org/10.1038/nbt.3988.

1093   76.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., Von
1094   Haeseler, A., Lanfear, R., and Teeling, E. (2020). IQ-TREE 2: New Models and Efficient
1095   Methods for Phylogenetic Inference in the Genomic Era. Mol. Biol. Evol. *37,* 1530–1534.
1096   https://doi.org/10.1093/molbev/msaa015.

1097   77.   Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2020). GTDB-Tk: a
1098   toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics *36,*
1099   1925–1927. https://doi.org/10.1093/bioinformatics/btz848.

1100   78.   Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R.,
1101   O'Hara, R.B., Solymos, P., Stevens, M.H.H., Szoecs, E., et al. (2001). vegan:

Community Ecology Package. CRAN Contrib. Packag. *368*, 11.
https://doi.org/10.32614/CRAN.package.vegan.

79. Hsieh, T.C., Ma, K.H., and Chao, A. (2016). iNEXT: an R package for rarefaction and
extrapolation of species diversity (Hill numbers). Methods Ecol. Evol. *7*, 1451–1456.
https://doi.org/10.1111/2041-210X.12613.

80. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics *34*, i884–i890.
https://doi.org/10.1093/bioinformatics/bty560.

81. Westbrook, A., Ramsdell, J., Schuelke, T., Normington, L., Bergeron, R.D., Thomas,
W.K., and MacManes, M.D. (2017). PALADIN: Protein alignment for functional profiling
whole metagenome shotgun data. Bioinformatics *33*, 1473–1478.
https://doi.org/10.1093/bioinformatics/btx021.

82. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and
SAMtools. Bioinformatics *25*, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

83. Blanco-Míguez, A., Beghini, F., Cumbo, F., McIver, L.J., Thompson, K.N., Zolfo, M.,
Manghi, P., Dubois, L., Huang, K.D., Thomas, A.M., et al. (2023). Extending and
improving metagenomic taxonomic profiling with uncharacterized species using
MetaPhlAn 4. Nat. Biotechnol. *41*, 1633–1644. https://doi.org/10.1038/s41587-023-
01688-w.

84. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 559. https://doi.org/10.1186/1471-2105-9-559.

85. Villanueva, R.A.M., and Chen, Z.J. (2019). ggplot2: Elegant Graphics for Data Analysis
(2nd ed.). Meas. Interdiscip. Res. Perspect. *17*, 160–167.
https://doi.org/10.1080/15366367.2019.1565254.

86. Bateman, A., Martin, M.J., Orchard, S., Magrane, M., Agivetova, R., Ahmad, S., Alpi,
E., Bowler-Barnett, E.H., Britto, R., Bursteinas, B., et al. (2021). UniProt: the universal
protein knowledgebase in 2021. Nucleic Acids Res. *49*, D480–D489.
https://doi.org/10.1093/nar/gkaa1100.

87. Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer,
E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The
protein families database in 2021. Nucleic Acids Res. *49*, D412–D419.
https://doi.org/10.1093/nar/gkaa913.

88. Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor,
R., Funk, K., Kelly, C., Kim, S., et al. (2022). Database resources of the national center
for biotechnology information. Nucleic Acids Res. *50*, D20–D26.
https://doi.org/10.1093/nar/gkab1112.

89. Parks, D.H., Chuvochina, M., Rinke, C., Mussig, A.J., Chaumeil, P.-A., and Hugenholtz,
P. (2022). GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy.
Nucleic Acids Res. *50*, D785–D794. https://doi.org/10.1093/nar/gkab776.

90. Frankish, A., Diekhans, M., Ferreira, A.-M., Johnson, R., Jungreis, I., Loveland, J.,
Mudge, J.M., Sisu, C., Wright, J., Armstrong, J., et al. (2019). GENCODE reference
annotation for the human and mouse genomes. Nucleic Acids Res. *47*, D766–D773.
https://doi.org/10.1093/nar/gky955.

91. Buchfink, B., Reuter, K., and Drost, H.-G. (2021). Sensitive protein alignments at tree-

of-life scale using DIAMOND. Nat. Methods *18*, 366–368.
https://doi.org/10.1038/s41592-021-01101-x.

92.   Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and Jermiin, L.S.
(2017). ModelFinder: fast model selection for accurate phylogenetic estimates. Nat.
Methods *14*, 587–589. https://doi.org/10.1038/nmeth.4285.

93.   Letunic, I., and Bork, P. (2019). Interactive Tree of Life (iTOL) v4: Recent updates and
new developments. Nucleic Acids Res. *47*, 256–259.
https://doi.org/10.1093/nar/gkz239.

94.   Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A
Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Stat.
Methodol. *57*, 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x.

95.   Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M.
(2022). ColabFold: making protein folding accessible to all. Nat. Methods *19*, 679–682.
https://doi.org/10.1038/s41592-022-01488-1.

96.   DeLano, W.L., and others (2002). Pymol: An open-source molecular graphics tool.
CCP4 Newsl. Protein Crystallogr *40*, 82–92.

97.   Cheng, Z., He, B.B., Lei, K., Gao, Y., Shi, Y., Zhong, Z., Liu, H., Liu, R., Zhang, H., Wu,
S., et al. (2024). Rule-based omics mining reveals antimicrobial macrocyclic peptides
against drug-resistant clinical isolates. Nat. Commun. *15*.
https://doi.org/10.1038/s41467-024-49215-y.

98.   Conway, J.R., Lex, A., and Gehlenborg, N. (2017). UpSetR: An R package for the
visualization of intersecting sets and their properties. Bioinformatics *33*, 2938–2940.
https://doi.org/10.1093/bioinformatics/btx364.

99.   Sjoberg, D. (2021). Ggsankey: Sankey, alluvial and Sankey bump plots. R Packag.
version 0.0 *99999*.

100.   Gu, Z. (2022). Complex heatmap visualization. iMeta *1*, 1–15.
https://doi.org/10.1002/imt2.43.

101.   Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: A Software Environment for
Integrated Models of Biomolecular Interaction Networks. Genome Res. *13*, 2498–2504.
https://doi.org/10.1101/gr.1239303.

102.   Wilkins, D., and Kurtz, Z. (2019). gggenes: draw gene arrow maps in 'ggplot2.' R
Packag. version 0.4. 0 *342*.

103.   Signorell, A., Aho, K., Alfons, A., Anderegg, N., Aragon, T., Arppe, A., Baddeley, A.,
Barton, K., Bolker, B., Borchers, H.W., et al. (2019). DescTools: Tools for descriptive
statistics. R Packag. version 0.99 *28*, 17.

1184

1185

Figure legends

Figure 1. Overview of IIBacFinder specializing in detecting unmodified bacteriocins

(A) Schematic overview of the genome mining procedure in IIBacFinder. IIBacFinder detects potential unmodified class II bacteriocins from nucleotide sequences in FASTA format, by leveraging the library of 1,153 pHMMs of precursor genes and 37 pHMMs of context genes. The putative core peptides of hypothetical bacteriocins are deduced and compared against known AMPs from four databases (i.e., APD3, dbAMP v2.0, DRAMP v3.0, and DBAASP v3.0). (B) Performance of three tools in recovering 205 known class II bacteriocins from 137 bacterial genomes. The maximum likelihood phylogenetic tree was inferred based on full bacteriocin sequences. Symbols at the leaf tips represent the subclass of class II bacteriocins, and the black triangles indicate the 12 bacteriocin sequences which are not included when building pHMM models in IIBacFinder. Numbers in brackets indicate the number of bacteriocins detected by each prediction tool.

Figure 2. Biosynthetic potential of class II bacteriocins in the bacterial kingdom

(A) Rarefaction curves illustrating how the number of detected bacteriocins increases with the inclusion of more bacterial genomes. (B) The GTDB bacterial tree, decorated with the number of genomes, detected bacteriocin sequences, unique bacteriocin sequences, and bacteriocin sequences per genome, demonstrates the bacteriocin biosynthetic potential across different bacterial phyla. (C) The top 20 bacterial genera containing the most abundant class II bacteriocins. (D) Antibacterial activity of 21 synthetic hypothetical bacteriocins (HBs) against both Gram-positive (+) and Gram-negative (-) bacteria. A red cross signifies that HBs cannot be identified by the prediction tool, while a green tick indicates successful detection. The activity screening was conducted at a concentration of 100 µg/mL, with activity being categorized into four groups (no activity, < 25% inhibitory ratio or P value > 0.05; low activity, 25% ≤ inhibitory ratio < 50% & P value < 0.05; medium activity, 50% ≤ inhibitory ratio < 75% & P value < 0.05; high activity, inhibitory ratio > 75% & P value < 0.05). HBs with high activity are highlighted in red. Asterisks in the activity screening heatmap denote an inhibitory ratio of ≥ 90%, which is subject to further MIC examination (rightmost panel). Exact MIC values are indicated in the heatmap, with antibiotic vancomycin serving as the positive control. All assays are performed in three independent replicates.

Figure 3. Class II bacteriocins exhibit niche-specific enrichment in host-associated bacteria

(A) Bar plots illustrating the number of genomes analyzed for class II bacteriocin identification and the corresponding number of non-redundant bacteriocin sequences detected. (B) Bar plot showing the number of non-redundant bacteriocin sequences detected from bacterial genomes across different habitats. (C) Venn diagram depicting the overlap of bacteriocin sequences detected from four distinct habitats, with pie charts indicating the proportion of niche-specific sequences, labeled with percentages. (D) Accumulation curves of bacteriocin sequences detected from the four habitats, with solid lines representing interpolated and actual data, and dashed lines representing extrapolated data.

Figure 4. Human gut bacteria have significant potential to produce a diverse array of unmodified bacteriocins

(A) Sankey plot showing the bacteriocin biosynthetic potential of 285,835 human gut bacteria across different taxonomic levels. Numbers in brackets indicate the number of non-redundant class II bacteriocin sequences detected. (B) Heatmap illustrating the number of gut microbiota-derived class II bacteriocins homologous to specific (left panel) and all (right panel) known

class II bacteriocins at varying identity thresholds (50%-90%). Only the top 20 most abundant
known class II bacteriocins are shown. (C) Rarefaction curves showing the increase in
detected bacteriocins with the inclusion of metatranscriptomes. The pie chart indicates the
percentage of class II bacteriocins detected (red) and not detected (grey) in 724
metatranscriptomes. (D) Scatter plot illustrating the prevalence of 6,238 putative bacteriocins
in metatranscriptomes. (E) Violin plot depicting the distribution of the number of class II
bacteriocins actively transcribed in individuals from three cohorts.

Figure 5. Prioritization and experimental validation of class II bacteriocins from human
gut microbiota

(A) Schematic overview of prioritizing class II bacteriocins using metagenomics and
metatranscriptomics analysis. Class II bacteriocins are detected from >700,000 bacterial
genomes, which are subsequently surveyed in human gut metagenomes (MG) and
metatranscriptomes (MT). A total of 3,581 bacteriocin clusters detected gut MG/MTare ranked
based on a meta-ranking scheme consisting of four criteria (i.e., prevalence, abundance,
negative association with bacterial species, and correlation with community diversity). (B)
Antibacterial activity of 26 human gut microbiota-derived hypothetical bacteriocins (HGM-HBs).
A red cross signifies that HGM-HBs cannot be identified by the prediction tool, while a green
tick indicates successful detection. The heatmap in the middle panel depicts the prevalence
and abundance of class II bacteriocins in different cohorts. The ones in the right panel show
the activity screening for 26 HGM-HBs at a concentration of 100 μg/mL. HGM-HBs with high
activity are highlighted in red. Asterisks in the activity screening heatmap denote an inhibitory
ratio of ≥ 90%, which is subject to further MIC examination (rightmost panel). Exact MIC values
are indicated in the heatmap, with antibiotic vancomycin serving as the positive control. All
assays are performed in three independent replicates.

Figure 6. Narrow-spectrum bacteriocins tweak human fecal microbiota in *ex vivo* assay

(A) Schematic overview of the *ex vivo* assay to explore the impact of bacteriocins on human
fecal microbiota. (B) Stacked bar plot displaying the average relative abundance of bacterial
families, with a top dendrogram illustrating hierarchical clustering based on the Bray–Curtis
dissimilarity of microbial communities at the family level. (C) Bar plot depicting the alpha
diversity of the microbial community at the species level, measured by the Shannon diversity
index. Data are presented as mean ± standard deviation. Statistical significance between
treatment groups and the control group (DMSO) was determined using a two-sided Welch's
t-test. (D) Principal Coordinate Analysis (PCoA) of the microbial community with the
treatment of 100 μg/mL of vancomycin or bacteriocins, based on Bray-Curtis dissimilarity at
the species level. (E) Stacked bar plot showing the number of significantly differential
species identified by MaAsLin2. (F) Heatmap representing the 71 significantly differentially
abundant species between three treatment groups (100 μg/mL of vancomycin, HGM_HP14,
and HGM_HP17) vs. the DMSO group. Each treatment group is symbolized by the
bacteriocin and the concentration tested (e.g., HGM_HP1-1 represents 1 μg/mL of
HGM_HP1, whereas HGM_HP1-100 represents 100 μg/mL of HGM_HP1). Statistical
significance is denoted as follows: $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; ns, not significant.

Tables

Table 1: Comparison of current bioinformatic tools for bacteriocin identification

Tools	Availability	Detection rules	Detection approach	Input file	Detection bacteriocin class	Class classification	Limitations
BPAGS	Web server	Precursor based	Machine learning algorithm [a]	Protein sequences	Mixed	No	1. Not detect bacteriocin based on context genes
BaPreS	Standalone tool	Precursor based	Machine learning algorithm [a]	Protein sequences	Mixed	No	2. Cannot infer the bacteriocin class of predicted sequences which might undergo posttranslational modification
RMSCNN	NA	Precursor based	Deep learning algorithm [b]	NA	Mixed	No	
NeuBI	Standalone tool	Precursor based	Deep learning algorithm [c]	Protein sequences	Mixed	No	
antiSMASH 6	Web server and standalone tool	Precursor and genomic context based	To scan signature gene pHMMs [d]	Genome sequences	RiPPs and RiPP-like [e]	No	1. Precursor-derived and context genes-derived HMMs are incomplete 2. RiPP-like BGCs encompass other unspecified RiPPs, not only limited to class II bacteriocins
PRISM 4	Web server	Precursor based	To scan signature gene pHMMs [f]	Genome sequences	Bacteriocin (Mixture of class II and III)	No	1. Precursor-derived pHMMs are incomplete 2. Not detect bacteriocin based on context genes
BAGEL 4	Web server and standalone tool	Precursor based	To blast core peptide database	Genome sequences	Class I, Class II, and Class III	Yes	1. Blast algorithm can only detect bacteriocin sequences similar to known class II bacteriocin 2. Not detect bacteriocin based on context genes

a, When constructing models using machine learning algorithm, BPAGS and BaPreS adopt BACTIBASE and BAGEL databases
as the training data set, which contains modified and unmodified bacteriocins.

b, When constructing models using deep learning algorithm, RMSCNN adopts APD vr3 database and MMBs dataset as the
training data set, which contains modified and unmodified bacteriocins, and other AMPs.

c, When constructing models using deep learning algorithm, NeuBI adopts BAGEL databases as the training data set, which
contain modified and unmodified bacteriocins.

d, In total, antiSMASH implements 20 pHMMs for categorizing RiPP-like BGCs. Among them, 8 pHMMs are derived from
precursors of class II bacteriocins, and one pHMM (TIGR01193) is a genomic context domain for bacteriocin cleavage/export
ABC   transporter.   Detection   details   can   be   found   at
https://github.com/antismash/antismash/blob/master/antismash/detection/hmm_detection/cluster_rules/relaxed.txt.

e, In antiSMASH, bacteriocin BGCs were removed and renamed to RiPP-like BGCs after version 6.0
(https://docs.antismash.secondarymetabolites.org/glossary/).

f, PRISM 4 adopts 133 HMMs of class II and class III bacteriocin precursors for bacteriocin identification.

g, BAGEL4 can detect class I bacteriocins based on genomic context, while it detects class II bacteriocins only by blasting against
the core peptide database.

NA, Not available.